Contextual effects in motor adaptation adhere to associative learning rules

Guy Avraham*[1,2], Jordan A. Taylor[3], Assaf Breska[1,2], Richard B. Ivry[1,2], Samuel D. McDougle[4]

1. Department of Psychology, University of California, Berkeley, Berkeley, CA 94720, USA

2. Helen Wills Neuroscience Institute, University of California, Berkeley, Berkeley, CA 94720, USA

3. Department of Psychology, Princeton University, Princeton, NJ 08544, USA

4. Department of Psychology, Yale University, New Haven, CT 06520, USA

* Corresponding author

E-mail: guyavraham@berkeley.edu

Keywords: Sensorimotor adaptation; associative learning; cerebellum; forward model; Rescorla-Wagner model

ABSTRACT

Two influential paradigms, eyeblink conditioning and motor adaptation, have deepened our understanding of the theoretical and neural foundations of sensorimotor learning. Although both forms of error-based learning are dependent on the cerebellum, the two literatures typically operate within distinct theoretical frameworks. For eyeblink conditioning, the focus is on the formation of associations between an error signal and arbitrary stimuli. For adaptation, the error signal is used to modulate an internal model of a sensorimotor map for motor planning. Here we take a step towards an integrative account of these two forms of learning, examining the relevance of core concepts from associative learning for motor adaptation. Using a task that drives implicit adaptation of reaching movements, we paired movement-related feedback with neutral auditory or visual cues that served as conditioning stimuli (CSs). Trial-by trial changes in feedforward movement kinematics exhibited two key signatures of associative learning: Differential conditioning and compound conditioning. Moreover, after compound conditioning, a robust negative correlation was observed between responses to the two elemental CSs of the compound (i.e., overshadowing), consistent with the additivity principle posited by models of associative learning. Computational modeling demonstrated that these results could not be captured by conventional, context-insensitive algorithms used to describe motor adaptation. Associative learning effects in motor adaptation provide a proof-of-concept for linking cerebellar-dependent learning paradigms within a common theoretical framework.

SIGNIFICANCE STATEMENT

Motor adaptation is a cerebellar-dependent process, describing how the motor system remains calibrated in response to environmental and bodily changes. Another cerebellar-dependent learning phenomenon, eyeblink conditioning, is viewed as an associative learning process. Here we sought to bringing together these two approaches for studying sensorimotor learning. We

demonstrate that core associative learning phenomena are manifest during motor adaptation,

pointing to a common framework for these distinct cerebellar-dependent motor learning

processes.

INTRODUCTION

A hallmark of many motor learning tasks is the use of error information to improve performance.

Sensorimotor errors not only signal that an action has not been properly executed but can provide

information on how the output of the system should be changed to increase the likelihood of future

success in a similar context. Two paradigmatic tasks for studying error-based learning are

sensorimotor adaptation and eyeblink conditioning. Sensorimotor adaptation refers to the gradual

adjustment of movements in response to changes in the environment or body. The key principle

in this process is that on each trial, sensorimotor prediction errors are used to update an internal

model of a sensorimotor mapping to help keep the sensorimotor system precisely calibrated

(Shadmehr and Krakauer, 2008; Wolpert et al., 1995; Wolpert and Ghahramani, 2000). In eyeblink

conditioning, a form of classical conditioning (Pavlov, 1927), learning centers on the formation of

associations between an aversive stimulus (i.e., an error signal) and arbitrary cues. In a standard

variant of this task, an aversive air puff to the cornea (unconditioned stimulus, or US) elicits a

reflexive blink (unconditioned response, or UR). The US can be repeatedly paired with a predictive

conditioning stimulus (CS, such as a tone or light flash) to gradually induce a preemptive

conditioned eyeblink response (CR). The CR is adaptative, mitigating the anticipated aversive

sensory consequences of the US.

In addition to serving as paradigmatic examples of implicit, error-based motor learning

(Clark et al., 2002; Clark and Squire, 1998; Mazzoni and Krakauer, 2006; Morehead et al., 2017),

sensorimotor adaptation and eyeblink conditioning share core properties. First, although one task

involves active planning (motor adaptation) and the other passive conditioning of a reflex (eyeblink

conditioning), both involve a learning signal based on a sensorimotor error that must occur close

in time to an initial prediction to drive learning (Brudner et al., 2016; Kitazawa et al., 1995;

Rasmussen et al., 2008; Schneiderman and Gormezano, 1964; Schween and Hegele, 2017;

Shadmehr et al., 2010). Second, both are strongly associated with the cerebellum (Donchin et al.,

2011; Garcia et al., 1999; Gerwig et al., 2007; Izawa et al., 2012; Kim and Thompson, 1997; Popa

et al., 2016; Schlerf et al., 2012).

While these parallels suggest a link between sensorimotor adaptation and eyeblink

conditioning, it is surprising that the two literatures have generally operated within distinct

theoretical frameworks. For adaptation, the focus has been on how an error signal is used to

directly update a motor command and the anticipated sensory consequences of that command.

For eyeblink conditioning, the focus has been on how an error signal is used to build an

association between an arbitrary stimulus (the CS) and an aversive event (the US) to produce a

predictive, adaptive response (the CR). This last point highlights one aspect of apparent

divergence between adaptation and conditioning: A prominent feature of sensorimotor adaptation

has been the finding that, at least when putatively restricted to implicit learning, arbitrary cues are

not effective for this form of learning. For example, people do not differentially adapt their reaching

movements to two opposing perturbations when the perturbation is signaled by a distinct color

cue in the environment (Howard et al., 2012, 2013; Gandolfo et al., 1996).

Subsequent work has shown that contextual cues can be highly effective when the cue is

directly relevant to the movement. For example, people can simultaneously adapt to opposing

perturbations if the context is established by movements prior to, or following, the perturbed

segment of a reach (Howard et al., 2015, 2012; Sheahan et al., 2016). In these situations, the

contextual cues are thought to affect learning because the cues are incorporated into the motor

plan itself (Howard et al., 2013, 2012). This notion contrasts with classical conditioning, where

predictive, arbitrary sensory cues need not have a direct relationship to the events they predict,

accruing associative value as long as they comply with strict timing requirements (Schneiderman

and Gormezano, 1964; Smith et al., 1969).

Here we take a first step toward bridging these two worlds by applying some key

theoretical concepts and design features of classical conditioning studies to a visuomotor rotation

task. First, we test the idea that arbitrary sensory cues can indeed shape motor adaptation, even

though they have no direct link to motor commands. Second, we test the hypothesis that the effect

of these cues on adaptation will adhere to established principles of associative learning. By

demonstrating these parallels between conditioning and motor adaptation, we offer a more

parsimonious framework for understanding these disparate cerebellar-dependent learning

processes.

RESULTS

We modified a visuomotor rotation task to ask if implicit motor adaptation exhibits associative

behavior when a perturbation is paired with an arbitrary sensory cue. Participants were asked to

reach from a start location to a target with movement feedback provided by a cursor (Fig. 1A).

Drawing on classic work in animal learning (Pavlov, 1927; Rescorla and Wagner, 1972), we

presented sensory events (a tone or light stimulus) as imperative cues for movement initiation.

The use of a neutral stimulus to cue movement initiation provides a means to establish a tight

temporal link between these arbitrary cues – the CSs in the task – and sensorimotor feedback,

the US. This temporal linkage of the CS and US, where the CS onset precedes the US and the

two events occur close in time, are prerequisites for cerebellar-dependent delay conditioning

(Schneiderman and Gormezano, 1964). To promote this timing constraint, a warning was given if

the participant did not move within 400 ms of the imperative, reminding them to initiate the

movement faster. Participants complies with this requirement, exhibiting rapid reaction times

(Mean ± STD: 287±45.2 ms).

Rather than impose a typical visuomotor perturbation on the feedback (e.g., rotate the

cursor relative to the true hand position), we opted to use "clamped" feedback. In this technique,

which is designed to isolate implicit motor adaptation, the cursor path is not tied to the direction

of the reach but rather follows an invariant ("clamped") path, with the radial position of the cursor

corresponding to the radial position of the hand (Morehead et al., 2017; Shmuelof et al., 2012).

As in past studies with clamped feedback, the participant was fully informed of the clamp

manipulation and instructed to ignore the task-irrelevant cursor and reach straight to the target on

all trials. As such, this method eliminates the potentially confounding effects of strategic processes

(McDougle et al., 2016; Kim et al., 2020). Nonetheless, participants' reach angles gradually shift

in the direction opposite to the clamp and shows the cardinal signatures of implicit motor

adaptation without awareness (Morehead et al., 2017; Kim et al., 2018, 2019; Parvin et al., 2018;

Tsay et al., 2020a, 2020b, 2021a, 2021b; Avraham et al., 2021; Poh et al., 2021; Vandevoorde

and Orban de Xivry, 2019; Yin and Wei, 2020).

Figure 1. Experiment 1: Differential Conditioning.

(A) Reaching task. Participants reach from a start location (white circle) to a target (blue dot). Online feedback is provided by a cursor (white dot). The target is displayed in a fixed location for the entire experiment (location varied across participants), and the direction of the cursor trajectory is fixed ("clamped") on all trials. After a random delay, a tone or a light (white frame) is presented, serving as the movement imperative. The cue persists for the duration of the movement. (B) Experimental protocol. During acquisition (white background), a 15° clamp (CW/CCW, counterbalanced across participants) was associated with CS+ (e.g., a tone) and a 0° clamp with CS- (e.g., a light; counterbalancing the associations with the tone and light across participants). During the probe phase (gray background), CS+ and CS- were presented with no feedback. Throughout the entire experiment, CS+ and CS- trials were randomly interleaved. (C, D) Experimental results (N=16) for trial-by-trial change (Δ) in hand angle during the acquisition (*C*) and probe (*D*) phases. Left panels present analysis results for an adaptation effect (main effect of trial *n-1*, dark vs light blue) and a Pavlovian effect (main effect of the presented CS on the current trial *n*, filled vs empty bars). The black outlined bar (right panel) presents the Pavlovian effect, i.e., the subtraction of hand angle changes between CS+ and CS- trials. (E, F) Rescorla-Wagner model simulation results during the acquisition (*E*) and probe (*F*) phases are consistent with the experimental results. Error bars represent SEM. Dots represent individual participants.

 Motor adaptation is modulated by arbitrary sensory cues

In differential conditioning, one CS is paired with the US (CS+) and another CS is presented

without the US (CS-). Thus, only the CS+ should become associated with the US and result in a

CR. To implement this in Experiment 1, we used two arbitrary cues for the CS's – a tone or a light

cue – and clamped cursor feedback for the US's (Morehead et al., 2017). Rather than vary the

presence or absence of feedback, we manipulated the trajectory of the cursor feedback to signal

the presence or absence of an error: For the CS+ condition, the feedback cursor followed a

clamped path that was rotated from the target by 15° (i.e., error/US present); for the CS- condition,

the feedback cursor always moved directly to the target (i.e., error/US absent). The CR in this

scenario is the angular deviation of the movement in the opposite direction of the 15° clamp.

During a 600-trial acquisition phase, CS+ and CS- trials were randomly interleaved.

Participants exhibited a marked change in movement direction during this phase, reaching an

asymptote of ~15° (Fig. S1A). This rapid adaptation is consistent with previous adaptation studies,

particularly those in which the target appears at a fixed location (Bond and Taylor, 2015; Day et

al., 2016; McDougle et al., 2015, 2017; Poh et al., 2021).

The main analysis centered on trial-by-trial changes in hand angle. The change in hand

angle from trial *n-1* to trial *n* is normally dictated by the feedback experienced in trial *n-1*. Thus,

following experience with an error on CS+ trials, participants should show increased adaptation

(a positive change in hand angle), and following no error on CS- trials, decreased adaptation. We

refer to these trial-by-trial changes as the "adaptation effect", the standard measure of learning in

sensorimotor adaptation studies. However, the conditioning framework makes a critical additional

prediction: The CS+ and CS- should differentially modulate the hand angle on trial *n* itself. That

is, the presentation of the light or tone should produce a CR associated with that cue, leading to

a difference in hand angle between the two cues that is independent of the feedback during the

previous trial. We refer to this as the "Pavlovian effect." Importantly, if implicit adaptation is not

sensitive to arbitrary contextual cues, we expect to only observe an adaptation effect.

A robust adaptation effect was observed (Fig. 1C, left panel): Trial-by-trial changes in

reaching direction (Δ hand angle) were significantly affected by the CS presented on the previous

trial [$F(1,15)=13.3$, $p=0.002$, $\eta_p^2=0.47$] such that the change in hand angle was larger after CS+

trials compared to after CS- trials. That is, the error occurring on CS+ trials (US+) resulted in

learning that carried over to the next trial, whereas the absence of an error on a previous CS- trial

(US-) resulted in a relative reversion to baseline (extinction). This is the canonical signature of

incremental motor adaptation.

The critical test of our hypothesis centers on the Pavlovian effect. As shown in Figure 1C

(both panels), the results revealed clear Pavlovian effects: The hand angle increased in the

direction of adaptation on CS+ trials (i.e., a positive change in hand angle relative to previous

trials) and decreased on CS- trials [$F(1,15)=5.83$, $p=0.029$, $\eta_p^2=0.28$]. This effect provides a novel

demonstration that arbitrary sensory cues can lawfully influence implicit motor adaptation.

We also observed an interaction between CS identity on trials *n-1* and *n* [$F(1,15)=7.11$,

$p=0.018$, $\eta_p^2=0.32$]. That is, the difference between CS+ and CS- was larger on trials following a

CS- (Mean±SE, 1.09°±0.39°) compared to trials following a CS+ (0.65°±0.35°). This interaction

effect likely reflects an asymmetry between the rate of the acquisition and extinction processes

once CS-US associations are established. That is, the state following a CS+ trial may be closer

to its asymptotic limit than after a CS- trial and is thus more limited in its potential for further

change.

We note that the visual feedback was different on CS+ and CS- trials, with the cursor

deviating from the target in the former and moving in a straight line to the target in the latter. This

raises the possibility that the hand angle differences on CS+ and CS- trials could be affected by

rapid online feedback responses. This explanation is unlikely given that the movements were

quite rapid ([Mean±STD], 103±31.2 ms). To directly assess the online correction hypothesis, we

calculated the difference in hand angle 50 ms after movement initiation and at the radial distance

of the target. There was no overall change in hand angle between the time points (-0.32°±2.92°),

and no significant difference between CS+ and CS- trials on this metric [$t(15)=-0.47$, $p=0.640$, $d=$-

0.12]. Thus, our results appear to pertain exclusively to feedforward learning.

Following the acquisition phase, participants completed a probe phase in which CS+ and

CS- trials were randomly presented in the absence of any visual feedback (no US). This phase

provides a "clean" test for associative learning effects since it removes trial-by-trial effects that

arise from the differential feedback given during CS+ and CS- acquisition trials. Here too we

observed a significant Pavlovian effect: Although there was an overall decrease in hand angle

across the probe phase (i.e., a partial washout of adaptation, Fig. S1A), there was a significant

main effect of the CS presented on trial n [$F(1,15)=7.98$, $p=0.013$, $\eta_p^2=0.35$] (Fig. 1D), with a

relative increase in hand angle on CS+ trials and a decrease on CS- trials. Moreover, there was

neither an effect of the trial n-1 CS [$F(1,15)=1.74$, $p=0.206$, $\eta_p^2=0.10$] nor a trial n-1 x n interaction

[$F(1,15)=0.25$, $p=0.621$, $\eta_p^2=0.02$], consistent with trial-by-trial adaptation being dependent on

error feedback.

Importantly, the Pavlovian effect did not appear to be driven by explicit awareness of the

CS-US contingency. A post-experiment survey was used to classify participants as either aware

or unaware of the CS-US associations (see *Methods*). Participants who reported being aware of

the contingencies (N=7) did not show a different Pavlovian effect compared to those who reported

being unaware (N=9) in either the acquisition ($t(14)=-0.333$, $p=0.744$, $BF_{10}=0.446$, $d=-0.17$) or

probe ($t(14)=-0.081$, $p=0.937$, $BF_{10}=0.431$, $d=-0.04$) phases. A similar null effect of awareness on

the strength of Pavlovian conditioning has also been reported in studies of human delay eyeblink

conditioning (Clark and Squire, 1998).

In summary, the observed effects of context on implicit motor adaptation in both the

acquisition and probe phases in Experiment 1 are consistent with differential conditioning effects

observed in classical conditioning. Feedforward implicit motor adaptation – here operationalized

as a type of CR – was differentially modulated by an arbitrary sensory CS+ versus CS-, with a

greater response to the CS+, the cue that was paired with a visuomotor error.

The standard approach to describe motor adaptation is based on a "state-space" model

in which the motor state is updated according to the error observed on the previous trial. The

parameters of this model determine both the trial-by-trial retention of the previous state and the

rate of learning from the error (see Eq. 3 in the *Methods*). By definition, a model of this form only

predicts a behavioral change based on trial *n*-1 (i.e., an adaptation effect), where no change in

hand angle is produced by the trial *n* CS; thus, this standard model cannot capture any context

effects (e.g., our Pavlovian effect).

In contrast, the Rescorla-Wagner model, a classic model of associative learning, provides

a straightforward account of context effects. It formalizes changes in conditioned responses via

the modulation of (arbitrary) learned associations. Here, the associative strengths, *V*, of the

conditioning stimuli are updated according to the learning rule (Eq. 1):

223   (1) $V^{[n]} = V^{[n-1]} + \alpha \cdot \beta \cdot SPE^{[n-1]}$,

where *V* represents the associative strengths between the US and the CS. It is updated based on

the sensory prediction error (*SPE*) presented on trial *n*-1. The *SPE* is defined as λ-$V^{[n-1]}$, where λ

is the maximum conditioning (asymptotic) level of the US. β is the learning rate parameter of the

US and α represents the salience of the CS. We note that the Rescorla-Wagner model does not

provide a mechanistic account for error-correction itself (e.g., the fact that the motor system

"knows" to update movements in the direction opposite of the error). For simplicity, we assume

that the sign of the change in movement direction is determined by a specialized neural circuit

attempting to reduce directional motor errors (Hadjiosif et al., 2021; Herzfeld et al., 2018; Wolpert

et al., 1998).

To illustrate that the Rescorla-Wagner model can capture differential conditioning behavior

similar to the results observed in Experiment 1, we performed model simulations. The simulation

results demonstrate both the adaptation effect and Pavlovian effect for the acquisition phase (Fig.

1E), as well as a clear Pavlovian effect for the Probe phase (Fig. 1F). Importantly, the behavioral

signature of differential conditioning, the larger changes in hand angle on CS+ trials compared to

CS- trials, holds for essentially all combinations of parameters in the Rescorla-Wagner model

(Fig. S2, left side). In contrast, there are no parameter combinations for the state-space model

that can produce this behavior (Fig. S2, right side). As noted above, the failure of the state-space

model should be expected given that it has no mechanism for handling contextual effects (we

return to this point in the *Discussion*).

To formally compare the Rescorla-Wagner and state-space models in terms of the

observed results, we conducted a model comparison by fitting each participant's data with the

two models. The Rescorla-Wagner model provided a better fit to the data than a standard state-

space model (Fig. S3; *t*-test comparing sum of squared residuals, t(15)=-3.62, p=0.003, *d*=-0.90;

t-test comparing Akaike Information Criterion (AIC) values, t(15)=-6.44, p<0.001, *d*=-1.61).

An additional analysis provided further support for an associative learning account of the

results of Experiment 1. The Rescorla-Wagner model not only provides a framework for

understanding how associations can be formed with arbitrary stimuli, but can also capture how

the strength of these associations is constrained by the relevance of the cues; for instance,

gustatory cues are much more likely to be associated with an internal state (e.g., nausea) than a

visual cue (Garcia and Koelling, 1966). In the current study, the clamped feedback (the US) is a

highly relevant stimulus for reaching; as such, we should expect it to have an immediate strong

influence on motor behavior. In contrast, the imperative cues, the tone and light (the CSs) have

no natural relevance for reaching; as such, their contribution to the CR should initially be quite

modest, gradually increasing over time (Fig. 2A). To test this prediction, we examined the time-

course of the adaptation and Pavlovian effects during acquisition using a linear mixed model

analysis. As expected, at early stages, the adaptation effect emerged quickly whereas the

contribution of the Pavlovian effect was small. Over experience, the relative contribution of the

two effects reversed (Type and Bin interaction effect: $F_{(1,365)}=16.7$, $p<0.001$): The Pavlovian

effect gradually grew (mean slope, [95% CI]; 0.14, [0.04 0.24]) at the expense of the adaptation

effect, which eventually exhibited a reduced contribution (-0.15, [-0.25 -0.05]) (Fig. 2B). We note

that both the decrease in the adaptation effect and the increase in the Pavlovian effect are not

captured by a typical state-space model (Fig. S4).

Figure 2. Dynamics of adaptation and Pavlovian effects.

Time course of the mean weights (regression β) of the adaptation (trial *n-1*, grey) and Pavlovian (trial *n*,

blue) effects in Experiment 1 as predicted by the Rescorla-Wagner model (A) and derived from fits of the

experimental results (B). Solid lines in *B* represent least squares regression lines. Shaded region represent

SEM.

Additivity principle in response to compound stimuli is observed in motor adaptation

The differential conditioning results of Experiment 1 show that implicit motor adaptation

manifests a prominent feature of classical conditioning, the gradual associability of an error signal

with arbitrary sensory cues. In Experiment 2, we tested a second core phenomenon of associative

learning, the principle of additivity (Mackintosh, 1976; Pavlov, 1927; Rescorla and Wagner, 1972).

This principle is based on the idea that there is an associative capacity for a given US – the *V*

term in the Rescorla-Wagner equation. Multiple CSs can become associated with a given US, but

the combined associative strength is bounded by *V*. As a result of this capacity constraint, CSs

effectively compete with one another, with the associative strength split among multiple cues.

The classic method to test for additivity is compound conditioning, where two or more

stimuli are presented simultaneously to form a "compound" CS (Eq. 2). When paired with a US,

this compound CS will come to elicit CRs. Importantly, the associative strength of the Compound

CS (V_{comp}) is the sum of the associative strengths of the elemental CSs (V_i), where *nS* in Eq. 2

represents the number of elements forming the Compound CS. Consequently, each element of

the compound, when presented alone, elicits a proportionally weaker CR, with the degree of

attenuation being a function of the associative strength of that CS.

$$(2)\ V_i^{[n]} = V_i^{[n-1]} + \alpha_i \cdot \beta \cdot (\lambda - V_{comp}^{[n-1]});\ V_{comp}^{[n-1]} = \textstyle\sum_{i=1}^{nS} V_i^{[n-1]}$$

The additivity principle has received ample support in behavioral and neural studies of

associative learning (Giurfa, 2007; Kehoe et al., 1994; Kehoe and Schreurs, 1986; Rescorla and

Wagner, 1972; Weiss, 1972), but has not, to our knowledge, been tested in motor adaptation. In

Experiment 2 we used a compound conditioning design pairing a 15° error clamp stimulus with a

compound CS (simultaneous presentation of the tone and light; Fig. 3A) on all trials during the

acquisition phase. As in Experiment 1, we again observed robust adaptation in the acquisition

phase, manifest as a change in hand angle in the direction opposite to the clamp (Fig. S1).

Figure 3. Experiment 2: Compound Conditioning.

(A) During acquisition, a tone and light were presented simultaneously (Compound CS), serving as the imperative signal for the reaching movement. They were paired on all trials with a 15° clamp. During the probe phase, no feedback was provided, and the CSs were presented either together (Compound CS) or alone (Single CS; tone or light). (B) Experimental results (N=22) for the Δ hand angle during the probe phase, showing a positive relative change for the compound CS on the current trial *n* (filled bar) and a negative relative change for each of its elements (empty bars). (C) Predictions from the Rescorla-Wagner model for trial-by-trial Δ hand angle during the probe phase. Note that the two elements were assumed to have equal weight in the simulation. (D) Scatter plot showing the between-participant trade-off in terms of the associative strength of the two CSs (the dotted black line represents the unity line). (E) The Δ hand angle after pooling the two single-CS conditions to measure the effects of the previous CS type (compound versus singleton, dark and light pink, respectively) and current CS (filled and empty bars). Error bars represent SEM.

The critical test in this experiment comes from the probe phase. Here, the clamped feedback was eliminated, and the imperative was either the original compound CS, or just the

tone or light alone. We observed a significant Pavlovian effect of CS type on these no-feedback

trials [$F_{(2,42)}=4.78$, $p=0.014$, $\eta_p^2=0.19$], with larger hand angles observed on compound CS trials

relative to the tone-alone or light-alone trials, conforming to the first prediction of the additivity

principle (Fig. 3B, 3C). Crucially, the additivity principle posits that there should be a negative

correlation between the associative strengths of competing CSs (Rescorla and Wagner, 1972).

That is, if a strong associative bond is formed between one CS and the US, this will come at the

expense of the associative strength accrued by any competing CSs (Eq. 2) given that there is a

capacity limit of associability (*V*). This prediction was strikingly confirmed in an analysis of the

hand angle changes on tone-alone and light-alone trials: Participants who were more sensitive to

the tone stimulus were less sensitive to the light stimulus, and *vice versa* (Fig. 3D, Pearson

correlation: $r=-0.72$, $p<0.001$).

As in Experiment 1, since the probe phase of Experiment 2 consisted of different types of

CSs presented randomly across trials, the behavior for a given trial should reflect not only the CS

on trial *n* but also the motor state on trial *n-1* (which is also influenced by the CS on that trial). As

a further test of compound conditioning, we pooled the two single-CS conditions to measure the

effects of the previous and current CS type (singleton versus compound) on the observed

changes in hand angle. We again observed the Pavlovian effect [$F_{(1,21)}=21.2$, $p<0.001$,

$\eta_p^2=0.50$]: There was a relative increase in hand angle for the compound CS (Mean ± SE,

0.24°±0.06°) and a relative decrease for the single CSs (-0.18°±0.06°) (Fig. 3E). The main effect

of the CS on trial *n-1* was not significant [$F_{(1,21)}=0.012$, $p=0.915$, $\eta_p^2=0.00$] nor was the trial *n-1*

× trial *n* interaction [$F_{(1,21)}=0.276$, $p=0.605$, $\eta_p^2=0.01$], presumably due to the elimination of trial-

by-trial adaptation given the absence of visual feedback in the probe phase.

DISCUSSION

Sensorimotor adaptation and eyeblink conditioning have provided foundational paradigms for the study of error-based sensorimotor learning. They have yielded a rich empirical foundation in the development of theoretical and neural models of learning and memory, particularly with respect to cerebellar function (Albus, 1971; Ito, 1984; Marr, 1969; Wolpert et al., 1998). Adaptation has typically been modeled from a control engineering perspective, centered on the idea that changes in behavior are driven by sensory prediction errors arising when the experienced feedback deviates from that predicted by a forward model operating on an efference copy of the motor command (Krakauer et al., 2019; Shadmehr and Krakauer, 2008; Wolpert and Flanagan, 2001). In contrast, eyeblink conditioning is treated as an associative learning process that can be driven by arbitrary sensory cues.

Our goal in the present study was to take a first step towards establishing explicit links between these two frameworks. To that end, we asked if core phenomena identified in the study of classical conditioning are operative in sensorimotor adaptation. We showed that visuomotor adaptation exhibited the hallmarks of both differential conditioning (Experiment 1) and compound conditioning (Experiment 2). These results provide the first evidence, to our knowledge, that pairing neutral stimuli (the tone and light) with distinct visuomotor outcomes can differentially influence implicit feedforward motor adaptation and do so in a manner consistent with the principles of associative learning rules.

The role of context in sensorimotor adaptation has been the subject of considerable debate. A number of labs have employed sensory cues, asking if these are sufficient to negate interference effects observed in response to interleaved opposing perturbations. The results from this work have generally shown that arbitrary visual cues such as differently colored cursors are ineffective, resulting in considerable interference between the perturbations (Gandolfo et al., 1996; Karniel and Mussa-Ivaldi, 2002; Howard et al., 2012, 2013; but see Krouchev and Kalaska,

2003; Osu et al., 2004; Addou et al., 2011). In contrast, postural and movement-related variables,

such as different lead-in and follow-through movements, have been shown to act as reliable

contextual cues for overcoming interference effects (Howard et al., 2012, 2013, 2015; Sheahan

et al., 2016). One interpretation of these results has focused on the difference between static and

dynamic cues, with the idea that the latter are incorporated into the motor state, allowing for the

separation of distinct internal models (Howard et al., 2012, 2013).

In the current study, we found compelling contextual effects using static, arbitrary sensory

cues. There are several key features of our design that differ from prior work with static cues.

First, drawing on the eyeblink literature (Schneiderman and Gormezano, 1964; Smith et al., 1969),

we imposed a strong temporal constraint on the interval between the CS (the contextual cue) and

the US by having the cues serve as imperative signals and enforcing rapid response times. This

strict temporal constraint was absent in prior studies; For example, in Howard et al. (Howard et

al., 2013), the color cue was presented 1,000 ms before the imperative. Second, the salience of

the arbitrary cues was arguably increased in the present study by having them serve as movement

imperatives.

More broadly, we propose that an associative learning framework may provide a useful

lens for understanding various aspects of sensorimotor adaptation. Consider a standard

adaptation study in which there are no contextual cues: Adaptation is observed when feedback is

perturbed while the participant reaches to a visual target. In such tasks, the target itself is not only

a salient stimulus that defines the task goal and movement plan, but its onset usually serves as

the imperative for movement initiation. Under these conditions, the target can be viewed as one

(highly effective) CS. Given that RTs in these tasks are typically below 500 ms (Kim et al., 2019,

2018; Avraham et al., 2021), such designs might establish a tight temporal link between target

appearance and movement, echoing (perhaps inadvertently) the CS-US temporal constraints

essential for eyeblink conditioning (Schneiderman and Gormezano, 1964).

More precisely, we propose that it is the movement plan itself, rather than the target cue,

that constitutes the primary CS in standard adaptation tasks. While the plan and target usually

coincide, this is not always the case. For example, with a contingent visuomotor rotation,

participants often deliberately aim away from the target, especially when the perturbation is large

(Hegele and Heuer, 2010; Taylor et al., 2014). Recognizing this "re-aimed" plan as a CS provides

a parsimonious account of a number of phenomena. First, generalization from adaptation is

centered around the direction of the movement plan but not the target (Day et al., 2016; McDougle

et al., 2017). This phenomenon echoes generalization effects seen in eyeblink conditioning,

where variation of the CS (e.g., tone frequency) leads to parametric changes in CR probability

(Siegel et al., 1968). Second, even in the absence of distinct contextual movements (e.g. follow-

through), the activation of different motor plans can serve as efficient contextual cues, negating

interference effects from opposing perturbations (Sheahan et al., 2016). Third, an emphasis on

the plan is in accord with general models of cerebellar-dependent adaptation, where the prediction

that constitutes the basis for sensory prediction error is computed using an efference copy of the

motor intention (Blakemore et al., 2001; Gao et al., 2016; Kawato and Gomi, 1992; Wolpert et al.,

1998).

To formally relate the eyeblink and adaptation worlds, we implemented the Rescorla-

Wagner model, a classic associative learning model that has been widely employed in the

classical and operant conditioning literature (Rescorla and Wagner, 1972). The success of this

model to capture the general features of the current data sets should not be surprising since the

model was developed to account for phenomena such as differential conditioning and compound

conditioning. Thus, it was obvious to us that the Rescorla-Wagner model would provide better fits

than the standard state-space model, given that the latter cannot capture contextual effects. In

theory, the standard state-space model can be modified such that different sensory cues become

associated with different states (Heald et al., 2018), allowing for context-dependent learning (e.g.,

differential conditioning). However, we emphasize that to account for compound conditioning, and

in particular, the additivity principle, a state-space model would essentially have to become a

variant of a Rescorla-Wagner model.

Although the Rescorla-Wagner model provides a ready account of how arbitrary sensory

cues, visual targets, and even movement plans may provide a CS to drive adaptation, there are

other phenomena in the sensorimotor adaptation literature that would not be accounted for by the

model, at least in its simplest form. One such effect is spontaneous recovery, the re-manifestation

of a previously adapted state in the absence of error feedback. A variant of the state-space model,

one that allows for multiple states with different learning and forgetting rates, can capture

spontaneous recovery (Smith et al., 2006). Second is the relationship between learning rate and

environmental consistency; learning is faster in response to a consistent versus inconsistent

perturbation (Albert et al., 2021; Avraham et al., 2020; Gonzalez Castro et al., 2014; Herzfeld et

al., 2014; Hutter and Taylor, 2018). Here, the state-space model has been modified to allow the

learning rate to vary with experience (Herzfeld et al., 2014). In its simplest form, the Rescorla-

Wagner model cannot account for these effects. However, recent efforts to model associative

learning have focused on how this framework may require more sophisticated, holistic

computational approaches that complement simple associative mechanisms with inference

processes (Collins and Frank, 2013; Gershman, 2015). A similar approach has been incorporated

in a recent work looking at contextual effects in sensorimotor adaptation, and could successfully

capture the aforementioned effects (Heald et al., 2020).

While our results highlight principles that address context-dependent motor adaptation,

they do not speak to the error-correcting adaptation algorithm itself. Conventional models of this

algorithm focus on the role of forward models in predicting future sensory states and updating

motor commands to reduce sensory prediction errors, which are signed error signals (Wolpert

and Ghahramani, 2000; but see Hadjiosif et al., 2021). Associative learning does not provide a

mechanism for the error correction process (e.g., a directional change in reaching movement

given rotated feedback, or a well-timed eyeblink response to an airpuff); rather, it describes the

association between a context and an error signal. Speculatively, motor adaptation perhaps

operates as a lookup table of a multitude of context-error associations, built up from a lifetime of

experience with subtle movement errors, with behavioral adjustments determined by a built-in,

direction-sensitive error correcting mechanism (Herzfeld et al., 2018). A model of this form could

bring sensorimotor adaptation closer to more classic models of cerebellar learning and plasticity

(Albus, 1971; Ito, 1984; Marr, 1969). More generally, exploring the computational links between

different sensorimotor learning tasks will be critical for understanding general principles of motor

learning and their neural mechanisms.

METHODS

Participants

Thirty-eight healthy volunteers (aged 18-31 years; 31 females) participated in either Experiment

1 (N=16) or Experiment 2 (N=22). All participants were right-handed, as self-reported and verified

with the Edinburgh Handedness Inventory. The protocol was approved by the Institutional Review

Board at the University of California, Berkeley.

Experimental setup and task

The participant sat at a custom-made table that housed a horizontally mounted LCD screen (53.2

cm by 30 cm, ASUS), positioned 27 cm above a digitizing tablet (49.3 cm by 32.7 cm, Intuos 4XL;

Wacom, Vancouver, WA). The participant held in their right hand a hockey paddle that contained

an embedded digitizing stylus. The monitor occluded direct vision of the hand, and the room lights

were extinguished to minimize peripheral vision of the arm. Reaching movements were performed

by sliding the paddle across the tablet. The sampling rate of the tablet was 200 Hz.

At the beginning of each trial, a white circle (0.5 cm diameter) appeared at the center of

the black screen, indicating the start location (Fig. 1A). The participant moved the stylus to the

start location. Feedback of hand position (i.e., the stylus position) was indicated by a white cursor

(0.3 cm diameter), provided only when the hand was within 1 cm of the start location. A single

blue target (0.5 cm diameter) was positioned 8 cm from the start location. In most studies of

adaptation, the appearance of the target specifies both the movement goal (where to reach) and

serves as the imperative (when to reach). From a classical conditioning perspective, the target

should constitute a very salient CS given that its onset is temporally contingent with the US, the

visual feedback associated with the movement (see below). To eliminate this temporal

contingency, the target remained visible at the same location during the entire experiment. For

each participant, the target was placed at one of four locations, either 45°, 135°, 225°, and 315°,

and this location was counterbalanced across participants.

Movement initiation was cued by the presentation of the neutral (non-spatial) CS(s). We

used two different CSs, a tone and a light, both of which have no inherent association with the

US. The tone CS was a pure sine wave tone with a frequency of 440 Hz. The light CS was a white

rectangular frame [39.4 cm X 26.2 cm] that spanned the perimeter of the visual workspace. The

large frame was selected to provide a salient visual stimulus, but one that would not be confused

with the target. The onset of the CS occurred following a pseudo-random and predetermined

delay after the hand was positioned at the start location. This was done to mitigate predictions

regarding the timing of the CS onset, and thus to enhance its salient role as an imperative. The

delay ranged between 800-1,200 ms (in steps of 100 ms), and was drawn from a uniform

distribution.

Depending on the specific experimental protocol and task phase, the CSs could appear

alone or together on a trial. The onset of the CS served as the imperative signal, with the

participant instructed to rapidly reach directly towards the target, slicing through the target. The

CS was terminated when the hand reached 8 cm, the radial distance to the target (Fig. 1A). To

minimize the delay between the onset of the CS and the US (the error clamp), the auditory

message "start faster" was played whenever a reaction time (RT) exceeded 400 ms. RT was

operationalized as the interval between CS onset and the time required for the radial distance of

the hand to exceed 1 cm. Given our objective to test the link between feedforward adaptation and

classical conditioning, we sought to eliminate online feedback corrections. Participants were

instructed to make rapid movements and the auditory message "move faster" was played

whenever movement time exceeded 300 ms. The end of the movement was operationalized as

the point where the radial distance of the hand reached 8 cm.

For the unconditioned stimulus (US), we used task-irrelevant clamped feedback

(Morehead et al., 2017). With clamped feedback, the radial position of the visual cursor is matched

to the radial position of the hand. However, the angular position of the cursor is fixed. The

participant thus controlled the speed and radial distance of the cursor, but not its direction. When

designed to produce a prediction error and elicit implicit sensorimotor adaptation, the clamp

followed a path that deviated from the target by 15°, with the direction, i.e., clockwise (CW) or

counterclockwise (CCW), counterbalanced across participants. We also included no-error trials

(Experiment 1) by presenting a clamped feedback that followed a path directly to the target (0°

clamp; Fig. 1B). The nature of the clamp manipulation was described in detail to the participant,

and they were explicitly instructed strictly to ignore the feedback, aiming their reach directly toward

the target on every trial. These instructions were designed to emphasize that the participant did

not control the cursor position, and that they should always attempt to reach directly to the target.

The instructions of the task were reinforced by the presentation of short video animations to

demonstrate how the CSs would serve as imperative signals, as well as to show the invariant

direction of the clamped feedback.

The experimental software was custom written in Matlab (The MathWorks, Natick, MA),

using the Psychtoolbox package (Brainard, 1997).

Experimental protocol

Both Experiments 1 and 2 included an acquisition phase and a probe phase. During the

acquisition phase, clamped feedback was presented on each trial, serving as the US. During the

probe phase, the clamped feedback was not presented. In both phases, the participants were

instructed to reach straight to the target as soon as the imperative CS appeared. Note that we

opted to not include baseline reaching blocks prior to the start of the acquisition phases to avoid

introducing any incidental associations between the baseline feedback and the target, movement

plan, and any other contextual variables. A break of approximately 1-minute was provided in the

middle of the experiment.

<u>Experiment 1: Differential conditioning</u>

Experiment 1 (N=16) was designed to test differential conditioning in the context of a sensorimotor

adaptation task. The session consisted of 800 trials: 600 acquisition trials followed by 200 probe

trials (Fig. 1B). One of two CSs (tone or light) was presented on each trial, serving as the

imperative for the reaching response. During the acquisition phase, one CS was paired with a 15°

clamped error feedback (CS+ condition) and the other CS was paired with a 0° clamped feedback

(CS- condition). Each CS was presented on 50% of the trials, and the assignment of the tone and

light to the CS+ and CS- was counterbalanced across participants. During the probe phase, each

CS was presented alone on half of the trials, and there was no visual feedback.

For both the acquisition and probe phases, the CS+ and CS- trials were interleaved in a

pseudo-random order that was unique to each participant. To ensure that the participant would

not be able to predict the CS type, the generated trial sequence for each participant was assessed

to verify that there was no significant lag-1 autocorrelation in the time series. At the end of the

experimental session, we assessed whether participants were aware of the contingency between

each CS and its associated feedback, asking, "Did you identify any pattern in the experiment?" in

a free-response survey.

Experiment 2: Compound conditioning

The adaptation task was modified in Experiment 2 (N=22) to provide a test of compound

conditioning. The procedure was similar to that used in Experiment 1 with the following changes.

First, the session consisted of 600 acquisition trials and 300 probe trials (Fig. 3A). Second, a

compound CS, consisting of both the tone and light, served as the imperative throughout the

acquisition phase, and was always paired with a 15° clamped feedback. Third, the probe phase

(no visual feedback) consisted of 100 trials for each of the compound CS, tone alone CS, and

light alone CS.

Data analysis

The recorded position of the digitizing stylus was analyzed using custom-written MATLAB scripts.

Our main analyses focused on the reach direction (hand angle) and the trial-by-trial changes in

hand angle (Δ hand angle). Hand angle was defined by two imaginary lines, one from the start

position to the target and the other from the start position to the hand position at maximum

movement velocity.

Trials in which the hand angle was larger than 100° off from the target, or in which the

trial-to-trial change in hand angle was larger than 25°, were considered outliers and not included

in the analyses. These outliers constituted 0.03% and 0.16% of all trials in Experiments 1 and 2,

respectively. For the change in hand angle analysis, but not for the presentation of hand angle

time courses (Fig. S1), we excluded trials in which the reaction time exceeded 400 ms and/or

movement time exceeded 300 ms (Experiment 1: 11% of acquisition trials, 4.3% of probe trials;

Experiment 2: 3.4% of acquisition trials, 4.9% of probe trials).

For all analyses and to visualize the results, the sign of the hand angle was flipped for

participants who experienced a CCW clamp, such that a positive hand angle is in the direction of

expected adaptation (i.e., opposite the direction of the perturbed feedback). Moreover, the hand

angle on the first acquisition trial was treated as the baseline reaching angle and subtracted from

the hand angle on all subsequent trials. (We note that the results remain unchanged in terms of

statistical comparisons if this baseline subtraction step is omitted.)

In Experiment 1, the primary analyses examined how the Δ hand angle was influenced by

the CS type (CS+ vs CS-), either in terms of the previous trial (*n-1*, adaptation effect) or current

trial (*n*, Pavlovian effect). For each participant and phase, we calculated the average Δ hand angle

for four types of trials: CS+ trials that follow CS+ trials, CS- after CS+, CS+ after CS-, and CS-

after CS-. For each phase, a two-way repeated-measures ANOVA was conducted with two within-

participant independent factors, the CS on trial *n-1* and the CS on trial *n*, each with two levels,

CS+ and CS-. The Δ hand angle was the dependent variable in the two ANOVAs. To examine

the dynamics of the adaptation and Pavlovian effects (Fig. 2B), we binned the hand angle data

into epochs of 50 trials. Within each bin, we performed multiple regression analysis to test whether

trial-to-trial change in hand angle can be predicted from the previous trial CS, the current trial CS

and their interaction. Figure 2B presents the mean ± SEM regression β weights of all simulated

time courses for the previous (adaptation effect) and current (Pavlovian effect) predictors. To

evaluate statistically the changes in β weights for each type of effect across bins, we used a linear

mixed model (R statistical package: lmerTest), with Type (adaptation and Pavlovian) and Bin as

fixed effects and participants as random effects.

To assess whether the Pavlovian effect is influenced by the awareness about the

contingency between each CS and its respective feedback, we divided the participants based on

their responses to the post experiment questionnaire (see Experimental protocol): 7 out of the 16

participants stated the correct contingency between the CS and the visual feedback and were

thus considered the "aware sub-group." The rest of the participants (N=9) reported that they did

not identify any pattern related to the CS-US contingency and were considered the "unaware sub-

group." Independent two-sample *t* tests were used to compare the Pavlovian effects between

these groups during the acquisition and probe phases of the experiment. For these tests we also

report Bayes factor BF_{10}, the ratio of the likelihood of the alternative hypothesis (H_1) over the null

hypothesis (H_0) (Kass and Raftery, 1995).

Feedback correction was operationalized as the difference between the hand angle

measured at the radial distance to the target and at 50 ms after movement initiation. We estimated

the mean and standard deviation of feedback correction across all acquisition trials in all of the

participants in Experiment 1. In addition, we calculated, for each participant, the mean feedback

correction for each of the CS+ and CS- trials, and used a paired-sample *t* test to examine within-

participant changes in feedback correction between the two trial types.

In Experiment 2, the analysis focused on the probe phase in which there was no visual

feedback. We compared the Δ hand angle in response to the three CSs on trial *n* (compound CS,

light alone, tone alone) regardless of the CS presented on trial *n-1*. We fit a one-way repeated-

measures ANOVA, with the Δ hand angle as the dependent variable, and the CS type as the

within-participant independent variable.

The additivity principle of the Rescorla-Wagner model states that the association strengths

of each element of the compound CS will compete for associative strength with respect to the US

(Eq. 2). This principle was tested in our data by examining the correlation between the Δ hand

angle associated with the tone and light CSs in Experiment 2.

We report effect size using Cohen's d for all t tests and partial eta-squared (η_p^2) for the

ANOVA. All of the measurements met the assumption of normality based on the Lilliefors test

(Lilliefors, 1967).

Model simulations

Trial-by-trial reach angles were simulated using the Rescorla-Wagner model as a representative

model for associative learning. In this model, the motor state is updated based on the associative

strength (V) between the US (error signal) and all the CSs present on a given trial (Eqs. 1-2). The

extent of learning is determined by the maximum conditioning level (λ), and the rate of behavioral

change are determined by the learning rate of the US (β) and the salience of each presented CS

(a). To illustrate the predictions of the model in Figures 1E, 1F and 3C, we chose parameters that

result in qualitatively similar effects to the experimental results. For both Experiments 1 and 2, we

set λ to 15, similar to the observed implicit adaptation asymptotes these experiments, as well as

in other studies (Bond and Taylor, 2015; Morehead et al., 2017). The value of β was set to 0.12

(Exp. 1) or 0.02 (Exp. 2), the salience parameter (a) for the movement plan CS to 0.99, and both

the tone CS and light CS to 0.002 (Exp. 1) or 0.1 (Exp. 2). These divergent salience values are

consistent with the assumption that most of the associative strength of the US would be absorbed

by the movement plan CS given its central relevance to the task of reaching to a target. The

remainder of potential associative strength to the US is thus available for the tone and light CSs.

To demonstrate how the particular choice of parameters in the Rescorla-Wagner model

influences the predicted Pavlovian effects, we simulated behavior using various combinations of

the free parameters. We chose several combinations of values for β and λ, and, for each

combination, a wide range of values for α_{goal}, α_{tone} and α_{light}. (For simplicity, we constrained α_{tone}

and α_{light} to the same value.) Figure S2 displays heatmaps resulting from these simulations, with

the color of each cell corresponding to the simulated difference in the trial-by-trial change in hand

angle between CS+ and CS- trials. In addition, we illustrated that a single-process state-space

model, a standard model of motor adaptation, cannot capture these Pavlovian effects. In this

model of motor adaptation, the motor state (x) is updated according to the following learning rule

(Eq. 3):

639　　　　　　　　　　　　　　　(3)　　$x^{[n]} = A \cdot x^{[n-1]} + B \cdot SPE^{[n-1]}$

where SPE is the sensory prediction error – the difference between the predicted and the actual

sensory feedback – experienced on trial n-1, A is the retention factor, and B is the learning rate.

For the simulation presented in Figure S2, the error was fixed to one of two values during the

acquisition phase, either -15° on CS+ trials or 0° on CS- trials, and in the probe phase, it was set

to 0° on all trials.

While the two models share similar features and parameters (e.g., learning rate of an error

signal), the additional parameters in the Rescorla-Wagner model allow it to capture the effects of

differential and compound conditioning, should these processes be operative in our experiments.

For example, in the differential conditioning case, modifying the parameters of the Rescorla-

Wagner model could produce changes in the magnitude of both the within-trial "Pavlovian" effects

(i.e., the modulation of behavior in response to the current CS, or "trial n" effects) and across-trial

adaptation effects (i.e., the modulation of behavior in response to the current state of learning, or

"trial n-1" effects). However, the Rescorla-Wagner model will always yield unique responses to

the CS+ and CS- given non-zero salience parameters. In contrast, no combination of the A and

B parameters in the state-space model will produce differential responses to the tone and light

CSs.

To simulate the dynamics of the adaptation and Pavlovian effects in differential

conditioning (Fig. 2A, S4), and to compare it to the dynamics observed in the actual data, we

simulated the trial-by-trial change in hand angle during acquisition (600 trials) according to the

Rescorla-Wagner model (Eqs. 1 and 2) and the state-space model (Eq. 3), based on the

schedules of CS+ and CS- trials that were presented to the participants (16 simulated time

courses). For the Rescorla-Wagner model, we used the same parameters values that generated

the simulation results presented in Figures 1E and 1F. For the state-space model, we set the *A*

parameter to 0.9 and *B* to 0.12. We then did the same trial-by-trial regression analysis described

above for the actual hand angle data (see *Data analysis*) with the exception that the dependent

variable was now the simulated change in hand angle.

Model fitting and comparison

We conducted a *post-hoc* model comparison analysis (Fig. S2). In this analysis, we fit the

Rescorla-Wagner model (Eqs. 1 and 2) and the standard state-space model (Eq. 3) to

participants' hand angle time course data of Experiment 1. The two models were fit by minimizing

the sum of squared residuals between the measured and modeled movement data, using the

MATLAB function *fmincon*. To avoid local minima, 200 randomized sets of initial parameter values

were used during fitting and the best fit of each model was selected for model comparison. Models

were compared using both the sum of squared residuals and the Akaike Information Criterion

approximated on the residuals (AIC; Akaike, 1974). All free parameters were bound at [0, 1], with

the exception of λ, which was bounded at [-30, 60]. We did not perform model fitting for

Experiment 2 due to the fact that each trial was identical in the acquisition phase. As such, the

salience parameters in the Rescorla-Wagner model for the goal, tone and light are unidentifiable,

and both models make indistinguishable behavioral predictions during acquisition.

ACKNOWLEDGEMENTS

We thank the Alkis Hadjiosif, John Krakauer, Adrian Haith and other members of the BLAM lab

(Johns Hopkins Medicine) for helpful discussions. We thank Marina Iranmanesh, Janet Hwang,

and Sarvenaz Pakzad for their assistance with data collection. RBI is funded by the NIH

(NS116883, NS105839 and DC077091).

REFERENCES

Addou T, Krouchev N, Kalaska JF. 2011. Colored context cues can facilitate the ability to learn
and to switch between multiple dynamical force fields. *Journal of Neurophysiology*
106:163–183. doi:10.1152/jn.00869.2010

Akaike H. 1974. A new look at the statistical model identification. *IEEE Transactions on
Automatic Control* **19**:716–723. doi:10.1109/TAC.1974.1100705

Albert ST, Jang J, Sheahan HR, Teunissen L, Vandevoorde K, Herzfeld DJ, Shadmehr R. 2021.
An implicit memory of errors limits human sensorimotor adaptation. *Nature Human
Behaviour* 1–15. doi:10.1038/s41562-020-01036-x

Albus JS. 1971. A theory of cerebellar function. *Mathematical Biosciences* **10**:25–61.
doi:10.1016/0025-5564(71)90051-4

Avraham G, Keizman M, Shmuelof L. 2020. Environmental consistency modulation of error
sensitivity during motor adaptation is explicitly controlled. *Journal of Neurophysiology*
123:57–69. doi:10.1152/jn.00080.2019

Avraham G, Morehead JR, Kim HE, Ivry RB. 2021. Reexposure to a sensorimotor perturbation
produces opposite effects on explicit and implicit learning processes. *PLOS Biology*
19:e3001147. doi:10.1371/journal.pbio.3001147

Blakemore S-J, Frith CD, Wolpert DM. 2001. The cerebellum is involved in predicting the
sensory consequences of action. *NeuroReport* **12**:1879–1884.

Bond KM, Taylor JA. 2015. Flexible explicit but rigid implicit learning in a visuomotor adaptation
task. *Journal of Neurophysiology* **113**:3836–3849. doi:10.1152/jn.00009.2015

Brainard DH. 1997. The Psychophysics Toolbox. *Spatial Vision* **10**:433–436.
doi:10.1163/156856897X00357

Brudner SN, Kethidi N, Graeupner D, Ivry RB, Taylor JA. 2016. Delayed feedback during

sensorimotor learning selectively disrupts adaptation but not strategy use. *Journal of*

Neurophysiology **115**:1499–1511. doi:10.1152/jn.00066.2015

Clark RE, Manns JR, Squire LR. 2002. Classical conditioning, awareness, and brain systems.

Trends in Cognitive Sciences **6**:524–531. doi:10.1016/S1364-6613(02)02041-7

Clark RE, Squire LR. 1998. Classical Conditioning and Brain Systems: The Role of Awareness.

Science **280**:77–81. doi:10.1126/science.280.5360.77

Collins AGE, Frank MJ. 2013. Cognitive control over learning: Creating, clustering, and

generalizing task-set structure. *Psychological Review* **120**:190–229.

doi:10.1037/a0030852

Day KA, Roemmich RT, Taylor JA, Bastian AJ. 2016. Visuomotor Learning Generalizes Around

the Intended Movement. *eNeuro* **3**. doi:10.1523/ENEURO.0005-16.2016

Donchin O, Rabe K, Diedrichsen J, Lally N, Schoch B, Gizewski ER, Timmann D. 2011.

Cerebellar regions involved in adaptation to force field and visuomotor perturbation.

Journal of Neurophysiology **107**:134–147. doi:10.1152/jn.00007.2011

Gandolfo F, Mussa-Ivaldi FA, Bizzi E. 1996. Motor learning by field approximation. *PNAS*

93:3843–3846. doi:10.1073/pnas.93.9.3843

Gao Z, Proietti-Onori M, Lin Z, ten Brinke MM, Boele H-J, Potters J-W, Ruigrok TJH, Hoebeek

FE, De Zeeuw CI. 2016. Excitatory Cerebellar Nucleocortical Circuit Provides Internal

Amplification during Associative Conditioning. *Neuron* **89**:645–657.

doi:10.1016/j.neuron.2016.01.008

Garcia J, Koelling RA. 1966. Relation of cue to consequence in avoidance learning. *Psychon*

Sci **4**:123–124. doi:10.3758/BF03342209

Garcia KS, Steele PM, Mauk MD. 1999. Cerebellar Cortex Lesions Prevent Acquisition of

Conditioned Eyelid Responses. *J Neurosci* **19**:10940–10947.

doi:10.1523/JNEUROSCI.19-24-10940.1999

Gershman SJ. 2015. A Unifying Probabilistic View of Associative Learning. *PLoS Comput Biol*

11. doi:10.1371/journal.pcbi.1004567

Gerwig M, Kolb FP, Timmann D. 2007. The involvement of the human cerebellum in eyeblink

conditioning. *Cerebellum* **6**:38. doi:10.1080/14734220701225904

Giurfa M. 2007. Behavioral and neural analysis of associative learning in the honeybee: a taste

from the magic well. *J Comp Physiol A* **193**:801–824. doi:10.1007/s00359-007-0235-9

Gonzalez Castro LN, Hadjiosif AM, Hemphill MA, Smith MA. 2014. Environmental Consistency

Determines the Rate of Motor Adaptation. *Current Biology* **24**:1050–1061.

doi:10.1016/j.cub.2014.03.049

Hadjiosif AM, Krakauer JW, Haith AM. 2021. Did we get sensorimotor adaptation wrong?

Implicit adaptation as direct policy updating rather than forward-model-based learning. *J*

Neurosci. doi:10.1523/JNEUROSCI.2125-20.2021

Heald JB, Ingram JN, Flanagan JR, Wolpert DM. 2018. Multiple motor memories are learned to

control different points on a tool. *Nature Human Behaviour* **2**:300–311.

doi:10.1038/s41562-018-0324-5

Heald JB, Lengyel M, Wolpert DM. 2020. Contextual inference underlies the learning of

sensorimotor repertoires. *bioRxiv* 2020.11.23.394320. doi:10.1101/2020.11.23.394320

Hegele M, Heuer H. 2010. Implicit and explicit components of dual adaptation to visuomotor

rotations. *Consciousness and Cognition* **19**:906–917. doi:10.1016/j.concog.2010.05.005

Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R. 2018. Encoding of error and learning to

correct that error by the Purkinje cells of the cerebellum. *Nature Neuroscience* **21**:736–

743. doi:10.1038/s41593-018-0136-y

Herzfeld DJ, Vaswani PA, Marko M, Shadmehr R. 2014. A memory of errors in sensorimotor

learning. *Science* 1253138. doi:10.1126/science.1253138

Howard IS, Ingram JN, Franklin DW, Wolpert DM. 2012. Gone in 0.6 Seconds: The Encoding of

Motor Memories Depends on Recent Sensorimotor States. *J Neurosci* **32**:12756–12768.

doi:10.1523/JNEUROSCI.5909-11.2012

Howard IS, Wolpert DM, Franklin DW. 2015. The Value of the Follow-Through Derives from

Motor Learning Depending on Future Actions. *Current Biology* **25**:397–401.

doi:10.1016/j.cub.2014.12.037

Howard IS, Wolpert DM, Franklin DW. 2013. The effect of contextual cues on the encoding of

motor memories. *Journal of Neurophysiology* **109**:2632–2644.

doi:10.1152/jn.00773.2012

Hutter SA, Taylor JA. 2018. Relative sensitivity of explicit reaiming and implicit motor

adaptation. *Journal of Neurophysiology* **120**:2640–2648. doi:10.1152/jn.00283.2018

Ito M. 1984. The Cerebellum and Neural Control. Raven Press.

Izawa J, Criscimagna-Hemminger SE, Shadmehr R. 2012. Cerebellar Contributions to Reach

Adaptation and Learning Sensory Consequences of Action. *J Neurosci* **32**:4230–4239.

doi:10.1523/JNEUROSCI.6353-11.2012

Karniel A, Mussa-Ivaldi FA. 2002. Does the motor control system use multiple models and

context switching to cope with a variable environment? *Exp Brain Res* **143**:520–524.

doi:10.1007/s00221-002-1054-4

Kass RE, Raftery AE. 1995. Bayes Factors. *Journal of the American Statistical Association*

90:773–795. doi:10.1080/01621459.1995.10476572

Kawato M, Gomi H. 1992. A computational model of four regions of the cerebellum based on

feedback-error learning. *Biol Cybern* **68**:95–103. doi:10.1007/BF00201431

Kehoe EJ, Horne AJ, Horne PS, Macrae M. 1994. Summation and configuration between and

within sensory modalities in classical conditioning of the rabbit. *Animal Learning &*

Behavior **22**:19–26. doi:10.3758/BF03199952

Kehoe EJ, Schreurs BG. 1986. Compound conditioning of the rabbit's nictitating membrane

response: Test trial manipulations. *Bull Psychon Soc* **24**:79–81.

doi:10.3758/BF03330509

Kim HE, Avraham G, Ivry RB. 2020. The Psychology of Reaching: Action Selection, Movement

Implementation, and Sensorimotor Learning. *Annu Rev Psychol.* doi:10.1146/annurev-

psych-010419-051053

Kim HE, Morehead JR, Parvin DE, Moazzezi R, Ivry RB. 2018. Invariant errors reveal limitations

in motor correction rather than constraints on error sensitivity. *Communications Biology*

1:19. doi:10.1038/s42003-018-0021-y

Kim HE, Parvin DE, Ivry RB. 2019. The influence of task outcome on implicit motor learning.

eLife **8**:e39882. doi:10.7554/eLife.39882

Kim JJ, Thompson RE. 1997. Cerebellar circuits and synaptic mechanisms involved in classical

eyeblink conditioning. *Trends in Neurosciences* **20**:177–181. doi:10.1016/S0166-

2236(96)10081-3

Kitazawa S, Kohno T, Uka T. 1995. Effects of delayed visual information on the rate and amount

of prism adaptation in the human. *J Neurosci* **15**:7644–7652.

doi:10.1523/JNEUROSCI.15-11-07644.1995

Krakauer JW, Hadjiosif AM, Xu J, Wong AL, Haith AM. 2019. Motor LearningComprehensive

Physiology. American Cancer Society. pp. 613–663. doi:10.1002/cphy.c170043

Krouchev NI, Kalaska JF. 2003. Context-Dependent Anticipation of Different Task Dynamics:

Rapid Recall of Appropriate Motor Skills Using Visual Cues. *Journal of Neurophysiology*

89:1165–1175. doi:10.1152/jn.00779.2002

Lilliefors HW. 1967. On the Kolmogorov-Smirnov Test for Normality with Mean and Variance

Unknown. *Journal of the American Statistical Association* **62**:399–402.

doi:10.1080/01621459.1967.10482916

809 Mackintosh NJ. 1976. Overshadowing and stimulus intensity. *Animal Learning & Behavior*

810   **4**:186–192. doi:10.3758/BF03214033

811 Marr D. 1969. A theory of cerebellar cortex. *The Journal of Physiology* **202**:437–470.

812   doi:10.1113/jphysiol.1969.sp008820

813 Mazzoni P, Krakauer JW. 2006. An Implicit Plan Overrides an Explicit Strategy during

814   Visuomotor Adaptation. *J Neurosci* **26**:3642–3645. doi:10.1523/JNEUROSCI.5317-

815   05.2006

816 McDougle SD, Bond KM, Taylor JA. 2017. Implications of plan-based generalization in

817   sensorimotor adaptation. *Journal of Neurophysiology* **118**:383–393.

818   doi:10.1152/jn.00974.2016

819 McDougle SD, Bond KM, Taylor JA. 2015. Explicit and Implicit Processes Constitute the Fast

820   and Slow Processes of Sensorimotor Learning. *J Neurosci* **35**:9568–9579.

821   doi:10.1523/JNEUROSCI.5061-14.2015

822 McDougle SD, Ivry RB, Taylor JA. 2016. Taking Aim at the Cognitive Side of Learning in

823   Sensorimotor Adaptation Tasks. *Trends in Cognitive Sciences* **20**:535–544.

824   doi:10.1016/j.tics.2016.05.002

825 Morehead JR, Taylor JA, Parvin DE, Ivry RB. 2017. Characteristics of Implicit Sensorimotor

826   Adaptation Revealed by Task-irrelevant Clamped Feedback. *Journal of Cognitive*

827   *Neuroscience* **29**:1061–1074. doi:10.1162/jocn_a_01108

828 Osu R, Hirai S, Yoshioka T, Kawato M. 2004. Random presentation enables subjects to adapt

829   to two opposing forces on the hand. *Nature Neuroscience* **7**:111–112.

830   doi:10.1038/nn1184

831 Parvin DE, McDougle SD, Taylor JA, Ivry RB. 2018. Credit assignment in a motor decision

832   making task is influenced by agency and not sensorimotor prediction errors. *J Neurosci*

833   3601–17. doi:10.1523/JNEUROSCI.3601-17.2018

834 Pavlov IP. 1927. Conditioned reflexes, translated by GV Anrep. *London: Oxford.*

Poh E, Al-Fawakari N, Tam R, Taylor JA, McDougle SD. 2021. Generalization of motor learning

in psychological space. *bioRxiv* 2021.02.09.430542. doi:10.1101/2021.02.09.430542

Popa LS, Streng ML, Hewitt AL, Ebner TJ. 2016. The Errors of Our Ways: Understanding Error

Representations in Cerebellar-Dependent Motor Learning. *Cerebellum* **15**:93–103.

doi:10.1007/s12311-015-0685-5

Rasmussen A, Jirenhed D-A, Hesslow G. 2008. Simple and Complex Spike Firing Patterns in

Purkinje Cells During Classical Conditioning. *Cerebellum* **7**:563. doi:10.1007/s12311-

008-0068-2

Rescorla R, Wagner A. 1972. A theory of Pavlovian conditioning: Variations in the effectiveness

of reinforcement and nonreinforcement. *Classical conditioning II: Current research and*

theory **2**:64–69.

Schlerf JE, Xu J, Klemfuss NM, Griffiths TL, Ivry RB. 2012. Individuals with cerebellar

degeneration show similar adaptation deficits with large and small visuomotor errors.

Journal of Neurophysiology **109**:1164–1173. doi:10.1152/jn.00654.2011

Schneiderman N, Gormezano I. 1964. Conditioning of the nictitating membrane of the rabbit as

a function of CS-US interval. *Journal of Comparative and Physiological Psychology*

57:188–195. doi:10.1037/h0043419

Schween R, Hegele M. 2017. Feedback delay attenuates implicit but facilitates explicit

adjustments to a visuomotor rotation. *Neurobiology of Learning and Memory* **140**:124–

133. doi:10.1016/j.nlm.2017.02.015

Shadmehr R, Krakauer JW. 2008. A computational neuroanatomy for motor control. *Exp Brain*

Res **185**:359–381. doi:10.1007/s00221-008-1280-5

Shadmehr R, Smith MA, Krakauer JW. 2010. Error Correction, Sensory Prediction, and

Adaptation in Motor Control. *Annual Review of Neuroscience* **33**:89–108.

doi:10.1146/annurev-neuro-060909-153135

Sheahan HR, Franklin DW, Wolpert DM. 2016. Motor Planning, Not Execution, Separates Motor

Memories. *Neuron* **92**:773–779. doi:10.1016/j.neuron.2016.10.017

Sheahan HR, Ingram JN, Žalalytė GM, Wolpert DM. 2018. Imagery of movements immediately

following performance allows learning of motor skills that interfere. *Scientific Reports*

8:14330. doi:10.1038/s41598-018-32606-9

Shmuelof L, Huang VS, Haith AM, Delnicki RJ, Mazzoni P, Krakauer JW. 2012. Overcoming

Motor "Forgetting" Through Reinforcement Of Learned Actions. *J Neurosci* **32**:14617–

14621a. doi:10.1523/JNEUROSCI.2184-12.2012

Siegel S, Hearst E, George N. 1968. Generalization gradients obtained from individual subjects

following classical conditioning. *Journal of Experimental Psychology* **78**:171–174.

doi:10.1037/h0026178

Smith MA, Ghazizadeh A, Shadmehr R. 2006. Interacting Adaptive Processes with Different

Timescales Underlie Short-Term Motor Learning. *PLOS Biology* **4**:e179.

doi:10.1371/journal.pbio.0040179

Smith MC, Coleman SR, Gormezano I. 1969. Classical conditioning of the rabbit's nictitating

membrane response at backward, simultaneous, and forward CS-US intervals. *Journal*

of Comparative and Physiological Psychology **69**:226–231. doi:10.1037/h0028212

Taylor JA, Krakauer JW, Ivry RB. 2014. Explicit and Implicit Contributions to Learning in a

Sensorimotor Adaptation Task. *J Neurosci* **34**:3023–3032.

doi:10.1523/JNEUROSCI.3619-13.2014

Tsay JS, Avraham G, Kim HE, Parvin DE, Wang Z, Ivry RB. 2020a. The effect of visual

uncertainty on implicit motor adaptation. *Journal of Neurophysiology* **125**:12–22.

doi:10.1152/jn.00493.2020

Tsay JS, Kim HE, Parvin DE, Stover AR, Ivry RB. 2021a. Individual differences in proprioception

predict the extent of implicit sensorimotor adaptation. *Journal of Neurophysiology*

125:1307–1321. doi:10.1152/jn.00585.2020

Tsay JS, Lee AS, Ivry RB, Avraham G. 2021b. Moving outside the lab: The viability of

conducting sensorimotor learning studies online. *bioRxiv* 2021.01.30.181370.

doi:10.1101/2021.01.30.181370

Tsay JS, Parvin DE, Ivry RB. 2020b. Continuous reports of sensed hand position during

sensorimotor adaptation. *Journal of Neurophysiology* **124**:1122–1130.

doi:10.1152/jn.00242.2020

Vandevoorde K, Orban de Xivry J-J. 2019. Internal model recalibration does not deteriorate with

age while motor adaptation does. *Neurobiology of Aging* **80**:138–153.

doi:10.1016/j.neurobiolaging.2019.03.020

Weiss SJ. 1972. Stimulus compounding in free-operant and classical conditioning: A review and

analysis. *Psychological Bulletin* **78**:189–208. doi:10.1037/h0032956

Wolpert DM, Flanagan JR. 2001. Motor prediction. *Current Biology* **11**:R729–R732.

doi:10.1016/S0960-9822(01)00432-8

Wolpert DM, Ghahramani Z. 2000. Computational principles of movement neuroscience. *Nature*

Neuroscience **3**:1212–1217. doi:10.1038/81497

Wolpert DM, Ghahramani Z, Jordan MI. 1995. An internal model for sensorimotor integration.

Science **269**:1880–1882. doi:10.1126/science.7569931

Wolpert DM, Miall RC, Kawato M. 1998. Internal models in the cerebellum. *Trends in Cognitive*

Sciences **2**:338–347. doi:10.1016/S1364-6613(98)01221-2

Yin C, Wei K. 2020. Savings in sensorimotor adaptation without an explicit strategy. *Journal of*

Neurophysiology **123**:1180–1192. doi:10.1152/jn.00524.2019

 FIGURE CAPTIONS

Figure 1. Experiment 1: Differential Conditioning.

(A) Reaching task. Participants reach from a start location (white circle) to a target (blue dot).

Online feedback is provided by a cursor (white dot). The target is displayed in a fixed location for

the entire experiment (location varied across participants), and the direction of the cursor

trajectory is fixed ("clamped") on all trials. After a random delay, a tone or a light (white frame) is

presented, serving as the movement imperative. The cue persists for the duration of the

movement. (B) Experimental protocol. During acquisition (white background), a 15° clamp

(CW/CCW, counterbalanced across participants) was associated with CS+ (e.g., a tone) and a 0°

clamp with CS- (e.g., a light; counterbalancing the associations with the tone and light across

participants). During the probe phase (gray background), CS+ and CS- were presented with no

feedback. Throughout the entire experiment, CS+ and CS- trials were randomly interleaved. (C,

D) Experimental results (N=16) for trial-by-trial change (Δ) in hand angle during the acquisition

(*C*) and probe (*D*) phases. Left panels present analysis results for an adaptation effect (main

effect of trial *n-1*, dark vs light blue) and a Pavlovian effect (main effect of the presented CS on

the current trial *n*, filled vs empty bars). The black outlined bar (right panel) presents the Pavlovian

effect, i.e., the subtraction of hand angle changes between CS+ and CS- trials. (E, F) Rescorla-

Wagner model simulation results during the acquisition (*E*) and probe (*F*) phases are consistent

with the experimental results. Error bars represent SEM. Dots represent individual participants.

Figure 2. Dynamics of adaptation and Pavlovian effects.

Time course of the mean weights (regression *β*) of the adaptation (trial *n-1*, grey) and Pavlovian

(trial *n*, blue) effects in Experiment 1 as predicted by the Rescorla-Wagner model (A) and derived

from fits of the experimental results (B). Solid lines in *B* represent least squares regression lines.

Shaded region represent SEM.

Figure 3. Experiment 2: Compound Conditioning.

(A) During acquisition, a tone and light were presented simultaneously (Compound CS), serving

as the imperative signal for the reaching movement. They were paired on all trials with a 15°

clamp. During the probe phase, no feedback was provided, and the CSs were presented either

together (Compound CS) or alone (Single CS; tone or light). (B) Experimental results (N=22) for

the Δ hand angle during the probe phase, showing a positive relative change for the compound

CS on the current trial *n* (filled bar) and a negative relative change for each of its elements (empty

bars). (C) Predictions from the Rescorla-Wagner model for trial-by-trial Δ hand angle during the

probe phase. Note that the two elements were assumed to have equal weight in the simulation.

(D) Scatter plot showing the between-participant trade-off in terms of the associative strength of

the two CSs (the dotted black line represents the unity line). (E) The Δ hand angle after pooling

the two single-CS conditions to measure the effects of the previous CS type (compound versus

singleton, dark and light pink, respectively) and current CS (filled and empty bars). Error bars

represent SEM.

Figure 1

Figure 2

Figure 3

A

Acquisition

Compound CS
🔊&💡
15° Clamp

Probe

Compound CS
🔊&💡
No feedback

Single CS
💡/🔊
No feedback

×60   ●●●   600   610   ×30

1   10   900

Trial #

B

p=0.016

p=0.012

ΔHand Angle (deg)

0.5

-0.5

Trial n: Comp Tone Light

C

Rescorla-Wagner
predictions

ΔHand Angle (deg)

0.5

-0.5

Trial n: Comp Tone Light

D

ΔHand Angle (deg)
Light Alone

-1

-1   0   1

ΔHand Angle (deg)
Tone Alone

E

p=0.915

p<0.001

ΔHand Angle (deg)

0.8

0.4

-0.4

Trial n: Comp Single Comp Single
Trial n-1:   Comp   Single