An associative learning account of sensorimotor adaptation

Guy Avraham[1,2], Jordan A. Taylor[3], Richard B. Ivry[1,2], Samuel D. McDougle[4]

1. Department of Psychology, University of California, Berkeley, Berkeley, CA 94720, USA

2. Helen Wills Neuroscience Institute, University of California, Berkeley, Berkeley, CA 94720, USA

3. Department of Psychology, Princeton University, Princeton, NJ 08544, USA

4. Department of Psychology, Yale University, New Haven, CT 06520, USA

* Corresponding author

E-mail: guyavraham@berkeley.edu

1 ABSTRACT

Two influential paradigms, sensorimotor adaptation and eyeblink conditioning, have deepened

our understanding of the theoretical and neural foundations of motor learning, and in particular,

the role of the cerebellum. Although there has been some cross-pollination between these two

lines of research, they typically operate within distinct theoretical frameworks, with the incremental

updating of an internal forward model explaining adaptation, and associative learning processes

explaining eyeblink conditioning. Here we ask if a unified framework might be parsimonious,

directly linking sensorimotor adaptation to associative learning. Using a task that isolates implicit

sensorimotor adaptation, we paired movement-related feedback with neutral auditory or visual

cues that served as conditioning stimuli (CSs) to test two key signatures of associative learning-

differential conditioning and compound conditioning. We observed clear Pavlovian effects in both

cases: Implicit trial-by-trial changes in movement kinematics were reliably modulated by the CSs

in the predicted directions. Moreover, after compound conditioning, we observed a robust

negative correlation between the responses of individuals to the two elemental CSs of the

compound, consistent with the additivity principle posited by the Rescorla-Wagner model of

classical conditioning. Computational modelling demonstrates that these results cannot be

captured by the conventional algorithm used to explain the operation of a forward model. We

believe that associative learning effects in implicit sensorimotor adaptation provide a proof-of-

concept for linking multiple motor learning paradigms within a similar theoretical framework.

INTRODUCTION

Sensorimotor adaptation refers to the gradual adjustment of movements in response to changes in the environment or body. Studies of human sensorimotor adaptation typically involve imposing a perturbation during reaching (Shadmehr and Mussa-Ivaldi, 1994), walking (Reisman et al., 2005), or speech (Houde and Jordan, 1998) to generate errors that drive recalibration of an existing sensorimotor map. Over numerous iterations, the adaptation process works to reduce the errors brought on by these perturbations, helping to optimize motor performance.

The primary theoretical framework used to describe adaptation is built around the engineering concept of an internal forward model (Shadmehr and Krakauer, 2008; Wolpert et al., 1995; Wolpert and Ghahramani, 2000). Here, a simulation process operating on the motor intent generates a prediction of the expected sensory outcome. This representation is compared to the actual sensory outcome and, if they are different, a sensory prediction error is generated to incrementally tune the forward model to reduce future errors.

Adaptation via sensory prediction error is considered to be an implicit form of learning. When properly isolated from other processes that contribute to learning in typical adaptation tasks, such as explicit cognitive strategies (McDougle et al., 2016), implicit adaptation appears to be curiously limited: It does not exhibit savings (Avraham et al., 2020; Haith et al., 2015; Morehead et al., 2015), is generally insensitive to both the magnitude and consistency of errors (Bond and Taylor, 2015; Hutter and Taylor, 2018; Avraham et al., 2019), and appears to be impervious to information relevant to task performance (Mazzoni and Krakauer, 2006; Morehead et al., 2017; but see Leow et al., 2018 and Kim et al., 2019). Moreover, as shown in several recent studies, implicit adaptation fails, at least initially, in producing appropriate changes in behavior in response to other simple perturbations like mirror reversals (i.e., when visual feedback is mirrored across an unseen axis with respect to the veridical limb position), a task that should be natural for a

canonical forward model to solve (Hadjiosif et al., 2020; Telgen et al., 2014; Wilterson and Taylor,

2020).

These constraints suggest that it may be appropriate to consider a simpler, model-free

computational framework for understanding implicit adaptation. As a starting point, we note

intriguing parallels between adaptation and another important paradigm in the study of motor

learning – Pavlovian delay eyeblink conditioning. In eyeblink conditioning, an aversive air puff

(unconditioned stimulus, or US) to the cornea elicits a reflexive blink (unconditioned response, or

UR). The US can be repeatedly paired with a predictive conditioning stimulus (CS, such as a tone

or light flash) to gradually induce a preemptive conditioned eyeblink response (CR) that, over

time, will come to be elicited even when the US is withheld. Both adaptation and eyeblink

conditioning are strongly associated with learning mechanisms in the cerebellum (Donchin et al.,

2011; Garcia et al., 1999; Gerwig et al., 2007; Izawa et al., 2012; Kim and Thompson, 1997; J. E.

Schlerf et al., 2012). In addition, although one task involves active planning (motor adaptation)

and the other passive conditioning of a reflex (eyeblink conditioning), both are thought to rely on

some form of prediction error (Ohmae and Medina, 2015; Rasmussen et al., 2008; J. Schlerf et

al., 2012; Shadmehr et al., 2010), which needs to occur close in time to the initial prediction to

produce learning (Brudner et al., 2016; Kitazawa et al., 1995; Schneiderman and Gormezano,

1964; Schween and Hegele, 2017). Lastly, both adaptation and eyeblink conditioning can proceed

outside of conscious awareness (Clark et al., 2002; Clark and Squire, 1998; Mazzoni and

Krakauer, 2006; Morehead et al., 2017).

Motivated by these similarities, we ask here how one might forge more direct links between

motor adaptation and eyeblink conditioning. Consider first the unconditioned events: The air puff

to the cornea and the imposed perturbation in adaptation studies can both be viewed as salient

USs, aversive sensory events that should be avoided or attenuated. Similarly, the eyeblink and

corrective feedback responses to a volitional action can be viewed as URs, automatic actions

made in response to their respective USs. When the US becomes predictable (i.e., when paired with a CS), the output of the system is modified to produce a response that marches forward in time, eventually resulting in an adaptive, feedforward response. In this light, sensorimotor adaptation can be seen as echoing the UR-to-CR transition in eyeblink conditioning (Albert and Shadmehr, 2016), with CRs defined as feedforward adjustments of movements in response to CSs. As such, sensorimotor adaptation could in theory come about via a simple associative learning rule rather than a multi-step process that requires internal simulations to generate a sensory prediction, compare that prediction to observed sensory feedback, and then use the difference to update the forward model.

In eyeblink conditioning, a neutral sensory stimulus is introduced as the CS, and learning requires that the CS and US be repeatedly paired in a relatively fixed temporal relationship. Continuing our analogy, what are putative CSs in adaptation? One candidate CS is the visual target that cues an action. For example, in reaching studies, the CS could simply be the appearance of a visual target (Welch and Abel, 1970), particularly when the target onset not only provides information required to plan the reach (i.e., direction and extent), but also serves as an imperative for movement initiation. Typical experimental designs thus create a tight temporal link between a neutral sensory cue (the target) and movement, echoing the CS-US temporal constraints essential for eyeblink conditioning (Schneiderman and Gormezano, 1964).

Some adaptation studies have revealed so-called context-dependent effects, which can be viewed as conceptually similar to classical conditioning. For instance, when participants are presented with two opposing perturbations applied to the same movement path across interleaved trials, they exhibit marked interference, failing to compensate for either perturbation (Karniel and Mussa-Ivaldi, 2002; Wainscott et al., 2005). However, this interference can be attenuated if the opposing perturbations are associated with distinct contextual cues (Addou et al., 2011; Gandolfo et al., 1996; Hirashima and Nozaki, 2012; Hwang et al., 2006; Osu et al., 2004). Dynamic cues,

especially those that are task-relevant, such as varying the path of the early and late phases of

movement, are especially effective in reducing this interference (Howard et al., 2013, 2015;

Sheahan et al., 2016). Moreover, the temporal relationship between the cue and subsequent

movement is critical: When a contextual cue precedes the movement by more than 600 ms, the

interference between conflicting perturbations persists (Howard et al., 2012). Less clear is the

efficacy of arbitrary, static sensory stimuli, such as color cues (Howard et al., 2012), although

studies attempting to modulate adaptation with static cues have generally not imposed the strict

temporal constraints required for context-dependent adaptation. The importance of temporal

specificity is clearly reminiscent of eyeblink conditioning, where static cues (e.g., tones and light

flashes) are extremely effective CSs insofar as they conform to strict temporal requirements

(Smith et al., 1969).

While contextual cueing studies can be viewed as providing a foundation for an

associative learning model of implicit adaptation, these handful of studies leave open three key

issues, two practical and one theoretical. First, contextual cues have the potential to introduce

confounding effects from explicit strategies (Howard et al., 2013; McDougle et al., 2015; Schween

et al., 2020), and thus their influence might not reflect a pure modulation of implicit adaptation.

Second, as mentioned above, the effect of static (versus dynamic) contextual cues remains

controversial, perhaps due to variations in temporal requirements of the task (Howard et al.,

2012). This issue must be addressed to convincingly link classical conditioning and implicit

adaptation, given that the former historically relies on static CSs.

Third, and most important, discussion of potential similarities between classical

conditioning and sensorimotor adaptation have been largely descriptive, focused mostly on their

shared dependency on the cerebellum (Popa et al., 2016). The contextual cueing studies

described above relate to one exemplary conditioning paradigm – differential conditioning – where

distinct CSs predict distinct outcomes. However, building a theoretical bridge between

sensorimotor adaptation and classical conditioning requires drawing on more subtle concepts

from the associative learning literature. In particular, from a conditioning perspective, the

associative strengths of different CSs are additive (Rescorla and Wagner, 1972). For example, in

compound conditioning, if two different CSs are simultaneously paired with a US, the associative

strength of the compound stimulus is equal to the sum of the associative strength of each

individual CS, which compete with one another. Fundamental conditioning principles like CS

additivity (or competition) have ample support in behavioral and neural studies of associative

learning, but have not, to our knowledge, been tested in the context of sensorimotor adaptation.

Here, we address these issues and examine the plausibility of an associative learning

framework for understanding implicit sensorimotor adaptation. We measured and computationally

modeled the effect of predictive CSs on trial-by-trial modulation of movements during adaptation.

We employed a method that strictly isolates implicit adaptation and is readily amenable to

presenting static CSs with precise temporal constraints. In Experiment 1, we verified differential

conditioning effects under these conditions. In Experiment 2, we used a compound conditioning

design to reveal that the conditioning principle of CS additivity applies to implicit sensorimotor

adaptation. Together, our behavioral and modeling results provide evidence for a unifying

framework for cerebellar-dependent learning, and may offer a parsimonious account of implicit

sensorimotor adaptation.

RESULTS

In two experiments, we modified a visuomotor adaptation task to ask if learning would show two foundational classical conditioning phenomena, differential conditioning and compound conditioning. We opted to use clamped visual feedback, a technique that excludes strategic contributions and thus restricts learning to implicit motor adaptation (Morehead et al., 2017). With this method, the participant makes a rapid reaching movement to a visual target and is presented with a feedback cursor that follows a fixed ("clamped") path. On error trials, the cursor trajectory is rotated by an invariant angle from the target, regardless of the movement direction of the hand. On error-free trials, the cursor trajectory goes directly to the target. Participants are fully aware of the clamp manipulation and instructed to reach straight to the target on all trials attempting to ignore the task-irrelevant cursor. Critically, when presented with the deviated trajectory repeatedly over trials, the participant's reach angle gradually shifts in the direction opposite the clamp (Fig. 1A), and their behavior shows the cardinal signatures of implicit motor adaptation (Morehead et al., 2017; Kim et al., 2018, 2019; Parvin et al., 2018; Avraham et al., 2020; Tsay et al., 2020).

Figure 1. Reaching Task. (A) Clamped-feedback induces implicit adaptation. (B) Adaptation described as a conditioning process. (C) Participants reach to a single target (blue dot), moving a cursor (white dot) from a start location (white circle). The target is displayed in a fixed location for the entire experiment (location varied across participants), and the direction of the cursor trajectory is fixed ("clamped") on all trials. After a random delay, a tone and/or a light (white frame) is presented, serving as the movement imperative. This cue persists for the duration of the movement.

In the present experiments, we modified the clamp procedure by introducing

manipulations borrowed from classical conditioning studies (Fig. 1B). We consider the clamped

feedback as an unconditioned stimulus (US), timed instantaneously with the reach. The

movement plan can be viewed as one conditioning stimulus (CS; Hirashima and Nozaki, 2012;

Sheahan et al., 2016). Repeated pairing of this plan-based CS with the clamped feedback (US)

forms an association that causes an adaptive conditioned response (CR) whenever the

movement plan is elicited, even when visual feedback is absent. The movement plan is present

on all reaching trials as it is fundamental to volitional movement. In most adaptation studies, the

onset of the target cues the movement plan and serves as the imperative, making the target

appearance a salient task-relevant stimulus. To eliminate this confound, a single target location

was used in the current study and the target was visible throughout the entire task (Fig. 1C).

Critically, we tested predictions of classical conditioning using arbitrary sensory CSs:

Drawing on classic work in animal learning (Pavlov, 1927; Rescorla and Wagner, 1972), we

accomplished this by presenting sensory events (i.e., a tone or light) as the imperative cues for

movement initiation (Fig. 1C). We thus created a situation where there were two putative CSs,

one consisting of the movement plan directed at the target location (presumably with a strong

preconditioned association with the task goal, i.e., visual feedback contacting the target), and one

serving as the movement imperative (tone/light, which are neutral with respect to the task goal).

The use of a neutral sensory stimulus to cue movement initiation provides a means to establish

a tight temporal link between the light or tone and the US (clamp feedback), a prerequisite for

cerebellar-dependent delay conditioning (Schneiderman and Gormezano, 1964). If the participant

did not move within 400 ms of the imperative, a warning was given, reminding them to initiate the

movement faster (Howard et al., 2012). Participants were able to comply with this requirement,

exhibiting rapid reaction times (Mean±STD: 287±45.2 ms).

In terms of conditioning, feedback corrections elicited by a deviated clamp could constitute

the unconditioned responses (UR). However, our primary goal here was to test the link between

feedforward adaptation and classical conditioning. As such, we opted to focus on trial-to-trial

changes resulting from adaptation. To this end, we required the participant to make rapid

movements, "shooting" through the target. If movement time was longer than 300 ms, the

participant was requested to move faster via an onscreen message. The conditioned response

(CR) was operationalized as changes across trials in feedforward movement commands.

In differential conditioning, one CS is paired with the US (CS+) and another CS is

presented without the US (CS-). Differential conditioning refers to the fact that only the CS+ should

become associated with the US and result in a CR. We implemented this in Experiment 1 (Fig.

2A) by pairing one imperative cue (e.g., the tone) with a 15° error clamp to create a CS+ (i.e.,

prediction error present), and the other imperative cue (e.g., the light) with a 0° clamp to create a

CS- (no prediction error). During the acquisition phase, CS+ and CS- trials, paired with their

associated US+/US-, were randomly interleaved (Fig. 2B). This phase was followed by a probe

phase in which CS+ and CS- trials were randomly presented in the absence of any visual

feedback (no US).

Participants exhibited a marked change in movement direction during the acquisition

phase, reaching an asymptote of ~15° (Fig. 2C). This rapid adaptation is consistent with previous

adaptation studies, particularly those in which a single target always appears at the same location

(Bond and Taylor, 2015; Day et al., 2016; McDougle et al., 2015, 2017).

The main analysis centered on the trial-by-trial changes in hand angle, with these data

used to compare two canonical models of motor learning. The internal model framework was

represented by the standard state-space model, a linear dynamical system model that captures

the time course of learning in motor adaptation tasks (Eq. 1, see Methods). The classical

conditioning framework was represented by the Rescorla-Wagner model, a model that formalizes

changes in conditioned responses via modulations in the associative strength of conditioned (and

unconditioned) stimuli (Eqs. 2-4). For each model, we generated predictions using a fixed set of

parameters (Fig. 2D). The state-space model predicts that the hand angle would be dictated by

the state of the internal model that is updated based on the error experienced in the previous trial

(trial *n-1*); we refer to this as the "adaptation effect." Critically, in this model, the hand angle would

not be influenced by the current CS (trial *n*). In contrast, the Rescorla-Wagner model predicts that

the two CSs would have different associative strengths in relation to the US, and as such, elicit

different CRs on trial *n* ("Pavlovian effect") in addition to the adaptation effect of the CS

experienced on trial *n-1*.

Consistent with the predictions of both models, a robust adaptation effect was observed

(Fig. 2E, left panel): Trial-by-trial changes in reaching direction (Δ hand angle) were significantly

affected by the CS presented on the previous trial [$F(1,15)=13.3$, $p=0.002$], such that the change

in hand angle was larger after CS+ trials compared to after CS- trials. That is, the error occurring

on CS+ trials (US+) resulted in learning that carried over to the next trial, whereas the absence of

an error on a previous CS- trial (US-) resulted in a relative reversion to baseline (extinction).

The critical test of our hypothesis centers on within-trial, or "Pavlovian" effects. That is, will

the reach angle on trial *n* be affected by whether the imperative is the CS+ or the CS-? The results

revealed clear Pavlovian effects (Fig, 2E, both panels): The hand angle increased in the direction

of adaptation on CS+ trials (i.e., a positive change in hand angle relative to previous trials) and

decreased on CS- trials [$F(1,15)=5.83$, $p=0.029$]. This effect is consistent with classical

conditioning, and accords with previous studies of contextual cueing (Hirashima and Nozaki,

2012; Howard et al., 2012, 2015; Sheahan et al., 2016). We also observed an interaction between

CS identity on trials *n-1* and *n* [$F(1,15)=7.11$, $p=0.018$]. That is, the difference between CS+ and

CS- was larger on trials following a CS- (Mean±SE, 1.09°±0.39°) compared to trials following a

CS+ (0.65°±0.35°). This interaction effect may reflect an asymmetry between the rate of the

acquisition and extinction processes once the CS-US associations are established. That is, the

state (or associative strength) following a CS+ trial may be closer to its asymptotic limit than after

a CS- trial, and is thus more limited in its potential for further change.

Figure 2. Experiment 1: Differential Conditioning. (A) Trial types. A 15° clamp (CW/CCW, counterbalanced across participants) was associated with CS+ and a 0° clamp with CS- (counterbalancing

the associations with the tone and light across participants). (B) Experimental protocol. CS+ (e.g., tone)

and CS- (e.g. light) trials were randomly interleaved. Clamped feedback (±15°/0°) was presented during

the acquisition phase (white background), and no feedback was provided during the probe phase (gray).

(C) Mean hand angle time course (N=16). (D) Simulation results for trial-by-trial change (Δ) in hand angle

during acquisition. While the state-space model predicts only an adaptation effect (main effect of trial *n-1*,

dark vs light blue), the Rescorla-Wagner model also predicts a Pavlovian effect (main effect of the

presented CS on the current trial *n*, filled vs empty bars). (E) Experimental results for trial-by-trial Δ hand

angle during acquisition comport with the Rescorla-Wagner model simulations but not the state-space

model simulations (left panel). The black outlined bar (right panel) presents the Pavlovian effect, i.e. the

subtraction of hand angle changes between CS+ and CS- trials. (F, G) Similar to (D) and (E), respectively,

but for the probe phase, showing a significant Pavlovian effect that is consistent with the Rescorla-Wagner

model. Shaded regions and error bars represent SEM. Dots represent individual participants.

We note that the feedback was different on CS+ and CS- trials, with the cursor deviating

from the target in the former and moving in a straight line to the target in the latter. This raises the

possibility that the hand angle differences on CS+ and CS- trials could be affected by rapid

feedback responses to the error. This explanation is unlikely given that the movements were quite

rapid (Mean±STD, 103±31.2 ms). Moreover, when we directly computed feedback corrections

(operationalized as the difference between the hand angle 50 ms after movement initiation and

at the radial distance of the target), we observed no overall change in hand angle between the

time points (Mean±STD, -0.32°±2.92°), and no significant difference between CS+ and CS- trials

[$t(15)$=-0.47, p=0.640].

The probe phase, where the visual feedback is eliminated, provides a more stringent test

for associative learning (Fig. 2F). Critically, here too we observed significant Pavlovian effects:

Although there was an overall decrease in hand angle across the probe phase (i.e., a partial

washout of adaptation, Fig. 2C), there was a significant main effect of the CS presented on trial *n*

[$F(1,15)$=7.98, p=0.013] (Fig. 2G), with a relative trial-by-trial increase in hand angle on CS+ trials,

and a relative trial-by-trial decrease on CS- trials. Moreover, there was neither effect of the *n-1*

CS [F(1,15)=1.74, p=0.206] nor an *n-1* x *n* interaction [F(1,15)=0.25, p=0.621], consistent with the

expectation that trial-by-trial adaptation effects are abolished in the absence of error feedback.

The simulations shown in Figures 2D and 2F illustrate the basic predictions of each model,

with parameter values chosen to approximate the mean experimental results. Crucially, the

construction of the standard state-space model offers no potential parameter combination that

can account for the observed behavior (Fig. S1). In contrast, differential conditioning effects of

the CS+ versus CS- hold for virtually any combination of Rescorla-Wagner model parameters

(Fig. S1).

Two additional analyses provide further support for the conditioning account over the

internal model account. First, a direct model comparison favored the Rescorla-Wagner model

over the state-space model (Fig. S2; *t*-test comparing sum of squared residuals, t(15)=-3.36,

p=0.004; *t*-test comparing Akaike Information Criterion (AIC) values, t(15)=-5.60, p<0.001).

Second, an *a posteriori* analysis tested a subtle but important aspect of our conditioning

hypothesis: Given the putative low associative salience, and, consequently, lower learning rates

(Eq. 4) for the neutral sensory imperative cues in our model, the contribution of these cues to the

CR should initially be rather modest and gradually increase over time. Because these cues

deterministically appear on different trial types, we can isolate their effects from the other putative

CS, the movement plan (which appears on all trials, and is thus mainly realized in cumulative

learning effects). That is, we can isolate the Pavlovian effect from the adaptation effect. We

investigated the time-course of these dissociable effects by performing a trial-by-trial multiple

regression analysis on the hand angle data (divided into 50-trial epochs), using the identity of the

previous (*n*-1) and current (*n*) CS as two separable predictor variables competing for variance.

Results of this analysis supported the conditioning model: As shown in Figure S3, the Pavlovian

effect related to the (trial *n*) imperative cues emerged gradually during acquisition, whereas the

adaptation effect (i.e., trial *n-1*) was much larger, emerged quickly, and then stabilized. Rescorla-

Wagner model simulations echoed this result.

Importantly, the observed Pavlovian effects did not appear to be affected by awareness

of the CS-US contingency. Participants that correctly identified the pattern of associations

between the imperatives and the type of feedback in a post-experiment survey ("aware" sub-

group) did not exhibit a different Pavlovian effect than an "unaware" sub-group, in either the

acquisition ($t(14)=-0.333$, $p=0.744$, $BF_{10}=0.446$) or probe ($t(14)=-0.081$, $p=0.937$, $BF_{10}=0.431$)

phases. This finding is consistent with work on human delay eyeblink conditioning, which was

also found to proceed outside of awareness (Clark and Squire, 1998).

In summary, the results from both the acquisition and probe phases in Experiment 1 are

consistent with predictions derived from an associative learning. Feedforward implicit motor

adaptation – here operationalized as the CR – was differentially modulated by the CS+ and CS-,

with a greater response to the CS+, the cue paired with a visuomotor error.

In Experiment 2, we put the associative learning account through a more stringent test.

We used a compound conditioning design to investigate a fundamental tenet of the Rescorla-

Wagner model, where the strength of the CR reflects the sum of the associative effects of all

presented CSs (Eq. 2; Rescorla and Wagner, 1972). In compound conditioning, two or more

sensory stimuli form a compound CS by being presented simultaneously. When paired with a US,

this compound CS comes to elicit CRs. Critically, each element of the compound, when presented

alone, elicits a proportionally weaker CR, with the degree of attenuation being a function of the

associative strength of that CS.

During the acquisition phase of Experiment 2 we presented a 15° error clamp stimulus on

all trials, and paired it with a compound CS (i.e., the simultaneous presentation of the tone and

light; Fig. 3A, 3B). As in Experiment 1, we again observed robust adaptation in the acquisition

phase, manifest as a rapid change in hand angle in the direction opposite to the clamp (Fig. 3C).

Figure 3. Experiment 2: Compound Conditioning. (A, B) During acquisition, a 15° clamp was associated
with a compound CS: Tone and light were presented simultaneously as movement imperatives. During the
probe phase, no feedback was provided, and the CSs were presented either together (Compound CS) or
alone (Single CS; tone or light). (C) Mean hand angle time course (N=22). (D) Simulation results for trial-
by-trial Δ hand angle during the probe phase for the compound CS (filled bar) and each of its elements
(empty bars). Unlike the state-space model, the Rescorla-Wagner model predicts a positive relative change
for the compound CS on the current trial *n* and a negative relative change for each element. (E)
Experimental results for the Δ hand angle during the probe are in line with the Rescorla-Wagner model. (F)
Scatter plot showing the relationship between individuals' responses to the elemental CSs (the dotted black
line represents the unity line). Shaded regions and error bars represent SEM. Dots represent individual
participants.

Acquisition was followed by a probe phase in which the clamp feedback was eliminated

(Fig. 3B). The imperative in the probe phase could now be either the same compound CS used

during the acquisition phase, or a singleton cue presented alone (i.e., either the tone or light). We

observed a significant Pavlovian effect of CS type in the participants' movements on these trials

[$F(2,42)=4.78$, $p=0.014$], with larger hand angles on compound CS trials versus both the tone-

alone or light-alone trials (Fig. 3E). These results are again consistent with the Rescorla-Wagner

model but cannot be explained by the standard state-space model (Fig. 3D).

The additivity (or cue competition) principle in classical conditioning also states that there

should be a negative correlation between the associative strengths of competing CSs (Rescorla

and Wagner, 1972). That is, if a strong associative bond is formed between one CS and the US,

this will come at the expense of the associative strength accrued by any competing CSs (Eq. 2).

This pattern was indeed observed when comparing responses to the tone-alone and light-alone

trials: Participants showing a large trial-by-trial increase in hand angle for the tone tended to

exhibit a relative decrease in hand angle for the light, and vice versa (Fig. 3F, Pearson correlation:

$r=-0.72$, $p<0.001$). This striking correlation reflects competition between the elements of the

compound CS, a key prediction of the Rescorla-Wagner model.

To this point, our analyses of the compound conditioning results have focused on the

responses to the CS on trial *n*, regardless of the CS identity on the previous trial. However, as

observed in the results of Experiment 1 with differential conditioning, even in the probe phase

where the visual feedback is eliminated (Fig. 2G), the trial-by-trial change in hand angle was

influenced not only by the CS on trial *n*, but also by the CS on trial *n-1*. Thus, to examine the

distinct response of the compound CS with respect to its elements, while also accounting for the

behavior on the previous trial, we pooled the two elemental CSs of the compound (the singletons

conditions) into a single CS-alone condition. Then we examined the effects of the previous and

current CS type (single versus compound) on the observed change in hand angle. We again

observed the predicted Pavlovian effect [$F_{(1,21)}=21.2$, $p<0.001$]: There was a marked relative

increase in hand angle for the compound CS (Mean±SE, 0.24°±0.06°) and a relative decrease for

the single CS (-0.18°±0.06°). There was neither significant main effect of the CS on trial *n-1*

[$F_{(1,21)}=0.012$, $p=0.915$] nor any trial *n-1* × trial *n* interaction [$F_{(1,21)}=0.276$, $p=0.605$], likely due

to the absence of visual feedback having eliminated trial-by-trial adaptation.

DISCUSSION

Sensorimotor adaptation and eyeblink conditioning have provided foundational paradigms for the study of motor learning, yielding key insights for theoretical and neural models of learning and memory, particularly with respect to cerebellar function. A control engineering perspective has generally been applied to modeling sensorimotor adaptation (Krakauer et al., 2019; Shadmehr and Krakauer, 2008; Wolpert and Flanagan, 2001), advancing the notion that changes in behavior arise from simulations of internal models of motor commands to anticipate their consequences. In contrast, eyeblink conditioning has been described as a model-free associative learning process. Our motivation in this work was the desire to bring together these two diverging literatures, and to highlight similar behavioral effects and overlapping neural mechanisms. Inspired by parsimonious models of classical conditioning (Rescorla and Wagner, 1972), we sought to reconsider adaptation as a manifestation of associative learning.

To this end, we tested if, and how, behavioral changes observed in sensorimotor adaptation can be captured by fundamental computational principles of classical conditioning, specifically, associability (Experiment 1) and additivity (Experiment 2). Our results provide the first evidence that, when properly timed, neutral and static sensory stimuli (CSs) paired with distinct visuomotor outcomes can differentially influence feedforward adaptation of reaching movements. In addition, we discovered that associating a visuomotor error with a compound CS resulted in a higher adaptive response to the compound than to each of its two elements, and observed a striking between-participant anti-correlation between the adaptive responses to these elements, comporting with the additivity principle put forth by Rescorla and Wagner (1972). Our results cannot be captured by the typical state-space model of adaptation, regardless of how it is parameterized (Fig. S1); however, they are clearly consistent with the Rescorla-Wagner model (Figs. 2 and 3). We believe these findings go beyond superficial links between adaptation and eyeblink conditioning. Rather, we believe they constitute a first step in the development of a more

cohesive algorithmic framework for understanding cerebellar learning across widely different

tasks.

The predominant model of motor adaptation states that adaptation is driven by the

updating of an internal forward model. A core function of the forward model is to simulate predicted

sensory consequences of descending motor commands, providing a representation that can be

compared with the observed outcomes, with any discrepancy between these quantities being

used to update the internal model (Wolpert et al., 1995). This framework is well-suited for a control

system hampered by the transmission delays inherent in an embodied agent. However, a number

of lines of evidence have highlighted how implicit adaptation, when isolated from other learning

processes, is a curiously stereotyped learning system that can even, given certain experimental

constraints, produce outputs that are counterproductive to performance (Mazzoni and Krakauer,

2006; Morehead et al., 2017). For example, participant's initial response to a mirror reversal visual

perturbation is in the opposite direction to the optimal solution, as if the visual feedback is rotated

(Wilterson and Taylor, 2020). These observations have motivated recent efforts to reframe

sensorimotor adaptation as the result of more rigid "direct policy updating" rather than the

updating of a flexible forward model (Hadjiosif et al., 2020). Here, we offer a novel account that is

compatible with a putative direct policy update, one that does not require any model-based

internal simulation – associative learning, with a strong prior to adapt movements in the direction

opposite to the observed errors.

An attempt to find consilience between associative learning processes like classical

conditioning of eyeblinks and other cerebellar-dependent learning tasks is not without

precedence. For instance, Raymond et al. (1996) made the conjecture that adaptation of the

vestibulo-ocular reflex (VOR) could be reframed as a conditioning process, with the error signal

given by experimentally-induced image motion on the retina acting as a US, and the vestibular

stimulus, driven by a head turn, acting as a conditioning stimulus (CS) (Raymond et al., 1996).

Moreover, several studies of human motor adaptation to force-field perturbations have

demonstrated that certain movement-related contextual cues can help form separate motor

memories during adaptation in a manner consistent with classical conditioning (Addou et al.,

2011; Hirashima and Nozaki, 2012; Howard et al., 2012, 2013, 2015; Osu et al., 2004; Sheahan

et al., 2016). However, to our knowledge, no study to date has directly tested fundamental

computational principles of classical conditioning in the context of visuomotor adaptation, while

simultaneously isolating implicit learning.

In our effort to frame adaptation as a form of conditioning, it is important to note that the

former has various features that are absent in the latter. First, adaptation is sensitive to the sign

of the error; in visuomotor adaptation, this sensitivity is reflected in the gradual deviation of the

movement direction opposite to the direction of the perceived error. At first blush, it is not clear

that the error in most conditioning tasks, like eyeblink conditioning, is similarly signed. However,

it is important to consider that the update of the CR is driven by a prediction error, which could be

positive or negative with respect to the expected feedback (Albus, 1971). Moreover, both the

cerebellar-dependent VOR and saccade adaptation processes are determined by the directional

sign of a prediction error (Boyden et al., 2004; Kojima et al., 2004). Indeed, such directional tuning

to errors has been observed in the cerebellum (Herzfeld et al., 2018; Kojima et al., 2010; Soetedjo

et al., 2008), and likely extends to reaching errors.

Second, by definition, classical conditioning paradigms like eyeblink conditioning do not

involve an instrumental component, while visuomotor adaptation tasks require volitional

movement. In that sense, motor adaptation is perhaps more akin to operant than classical

conditioning. While the elements of models for operant and classical conditioning are indeed

different, and likely engage non-identical processes (Brembs and Plendl, 2008; Lorenzetti et al.,

2006; Ostlund and Balleine, 2007), it is also clear that both can be viewed as outcomes of a CS-

US association (Brembs and Heisenberg, 2000). Fundamentally, this distinction does not change

our account; concordantly, the underlying computational model we employed (Rescorla and

Wagner, 1972) is applicable to both cases. Furthermore, the volitional movement plan, driven by

the presentation of the visual target (or imperative cue), could be considered as a "pre-

conditioned" CS, one that has a strong baseline association with the non-perturbed sensory

feedback experienced in everyday movement.

The idea of a movement plan acting as a CS is supported by multiple lines of evidence.

First, "follow-through" movements can serve as sufficient cues to allow seemingly identical pre-

movements be associated with distinct perturbative errors (Sheahan et al., 2016). Even though

the implementations of these follow-through movements are deferred until after errors have been

experienced, it is presumably planned before the movement is made. This can be seen as a form

of differential conditioning, with different motor plans serving as unique CSs. Furthermore, these

high-level motor plans appear to maintain this associative quality even when they are simply

imagined by the participant without being executed (Sheahan et al., 2018). Second, recent

evidence suggests that the generalization function for implicit adaptation is yoked to participants'

explicit movement goals (Day et al., 2016; McDougle et al., 2017), reinforcing the notion that

implicit adaptation is tied to the movement plan. Third, there is ample neurophysiological support

for a top-down motor command behaving like a CS: Descending motor commands (efference

copies) constitute a major source of input to the cerebellar cortex, and are associated with motor

errors during learning (Blakemore et al., 2001; Gao et al., 2016; Kawato and Gomi, 1992;

Khilkevich et al., 2018; Wolpert et al., 1998). Thus, our focus on associative learning fits with the

conventional notion of the cerebellum as a high-throughput pattern recognition device (Marr,

1969), which is, arguably, an optimal substrate for representing a diverse set of associations.

We also note that considering the movement plan as akin to a CS implies that an internal

representation links motor commands to predicted sensory consequences. As such, our account

is broadly consistent with a conventional internal forward model. However, under the conditioning

framework, this representation is fully described as the outcome of a straightforward associative

learning process (i.e., essentially a look-up table) without requiring any form of model-based

forward simulation. Thus, while functionally related, the forward model framework and the

associative learning framework are algorithmically and mechanistically distinct.

On a more speculative note, a parsimonious account of sensorimotor adaptation as

associative learning could avoid one additional problem in the forward model account; namely,

the computational cost one would expect to accompany a forward simulation process. Simulation-

based accounts for movement planning and motor learning are perhaps better suited for

describing top-down, cognitive behaviors, such as the strategic aiming of movements (Haith et

al., 2015; McDougle and Taylor, 2019; Schween et al., 2020). Indeed, forward simulations are

typically linked to high-level cognitive tasks that require planning, such as spatial navigation and

multi-step decision-making (Doll et al., 2012). Moreover, the framing of implicit sensorimotor

adaptation as an outcome of a model-based simulation process rather than a model-free

associative learning process also echoes certain broader distinctions between control

engineering and machine learning approaches. In the former, feedforward adaption could be

assumed to reflect changes in a complex internal model, whereas in the latter, it can be described

as the result of robust associations established through repeated training (e.g., via deep

reinforcement learning; Heess et al., 2017).

Further work is certainly required to provide a more systematic investigation of the utility

of associative learning models for understanding sensorimotor adaptation. First, as currently

framed, we do not provide an analog of the unconditioned response (UR) in reach adaptation.

Online feedback corrections could perhaps fill this gap: Rapid feedback corrections are

comparable to reflexive eyeblinks in that they entail automatic responses to errors, evolving into

feedforward (conditioned) responses over time (Albert and Shadmehr, 2016). Because we opted

to use rapid shooting movements to isolate feedforward adaptation, no reliable feedback

corrections were observed. Importantly, specifying a UR is not a necessary precondition for

characterizing a conditioning process; indeed, at the level of the UR, the line between classical

and instrumental conditioning is blurred. For example, in Pavlovian-to-Instrumental transfer, a

behavior learned through classical conditioning influences subsequent volitional behavior (Cartoni

et al., 2016).

Another curiosity is the incomplete unlearning we observed in the no-feedback probe

phase of both experiments, with reach angles remaining deviated in the direction of adaptation

(Figs. 2C and 3C). Given the rapid pace of learning in the acquisition phase, these persistent

aftereffects are not predicted by either the basic state-space model or the Rescorla-Wagner

model. However, persistent aftereffects are not uncommon in adaptation tasks, and are especially

robust when error feedback is simply removed rather than replaced by veridical feedback

(Brennan and Smith, 2015), perhaps reflecting a use-dependent Hebbian form of learning

(Verstynen and Sabes, 2011). The prolonged aftereffects we observed in the probe phase could,

in theory, be captured by state-space models that contain two parallel learning components, such

as separate fast and slow learning states (Smith et al., 2006). While multi-rate state-space model

may be able to capture the overall shape of the acquisition and extinction functions, they would

still fail to explain the differential and compound Pavlovian effects we observed. Future studies

using more complex tasks designed to dissociate multiple learning components could be used to

expand the conditioning account to accommodate a two-state learning model and additional use-

dependent effects.

Lastly, we caution that our findings do not directly refute nor falsify conventional

explanations of adaptation, such as the forward model, nor do they speak to the utility of such

models in explaining other types of motor behaviors. The results do suggest that, at the very least,

associative learning processes may be superimposed onto model-based adaptation. However, if

an associative learning account can successfully explain various forms of cerebellar motor

learning, we believe parsimony should favor it over a forward model-based account. Moreover,

the associative learning framework is supported by decades of conditioning literature; thus, the

proposal is rooted in well-known psychological and neurophysiological mechanisms.

METHODS

Participants

Thirty-eight healthy volunteers (aged 18-31 years; 31 females) participated in one of two

experiments, 16 participants in Experiment 1 and 22 in Experiment 2. All participants were right-

handed, as verified with the Edinburgh Handedness Inventory. The protocol was approved by the

Institutional Review Board at the University of California, Berkeley.

Experimental setup and task

The participant sat at a custom-made table that housed a horizontally mounted LCD screen (53.2

cm by 30 cm, ASUS), positioned 27 cm above a digitizing tablet (49.3 cm by 32.7 cm, Intuos 4XL;

Wacom, Vancouver, WA). The participant held in their right hand a hockey paddle that contained

an embedded digitizing stylus. The monitor occluded direct vision of the hand, and the room lights

were extinguished to minimize peripheral vision of the arm. Reaching movements were performed

by sliding the paddle across the tablet. The sampling rate of the tablet was 200 Hz. The

experimental software was custom written in Matlab (The MathWorks, Natick, MA), using the

Psychtoolbox package (Brainard, 1997).

At the beginning of each trial, a white circle (0.5 cm diameter) appeared at the center of

the black screen, indicating the start location (Fig. 1C). The participant moved the stylus to the

start location. Feedback of hand position (i.e., the stylus position) was indicated by a white cursor

(0.3 cm diameter), only provided when the hand was within 1 cm of the start location. A single

blue target (0.5 cm diameter) was positioned 8 cm from the start location. In the classical

conditioning framework, the target, which also drives the movement plan, should constitute a very

salient CS if it is presented in temporal contingency with the US, the visual feedback associated

with the movement (see below). To eliminate this temporal contingency, the target remained

visible at the same location during the entire experiment. For each participant, the target was

placed at one of four locations, either 45°, 135°, 225°, and 315°, and this location was

counterbalanced across participants.

Movement initiation was cued by the presentation of the neutral (non-spatial) CS(s). We

used two different CSs, a tone and a light, which are both assumed to have no inherent

association with the US. The tone CS was a pure sine wave tone with a frequency of 440 Hz. The

light CS was a white rectangular frame [39.4 cm X 26.2 cm] that spanned the perimeter of the

visual workspace. The large frame was selected to provide a salient visual stimulus, but one that

would not be confused with the target. The CSs appeared following a pseudo-random and

predetermined delay from the time the cursor was in the start location. The delay ranged between

800-1,200 ms (in steps of 100 ms), and was drawn from a uniform distribution.

Depending on the specific experimental protocol and task phase, the CSs could appear

alone or together on a trial. The onset of the CS served as the imperative signal, with the

participant instructed to rapidly reach directly towards the target upon perceiving the imperative,

slicing through the target. The CS was terminated when the hand reached 8 cm, the radial

distance to the target (Fig. 1C). To minimize the delay between the onset of the CS and the US

(the error clamp), the auditory message "start faster" was played whenever the reaction time (RT)

exceeded 400 ms. RT was operationalized as the interval between CS onset and the time required

for the radial distance of the hand to exceed 1 cm. To encourage rapid movements, the auditory

message "move faster" was played when the movement time exceeded 300 ms, with the end of

the movement operationalized as the point where the radial distance of the hand reached 8 cm.

For the unconditioned stimulus (US), we used task-irrelevant clamped feedback

(Morehead et al., 2017). With clamped feedback, the radial position of the visual cursor is matched

to the radial position of the hand. However, the angular position of the cursor is fixed. The

participant thus controlled the speed and radial distance of the cursor, but not its direction. When

designed to produce a prediction error and elicit implicit sensorimotor adaptation, the clamp

followed a path that deviated from the target by 15°, with the direction, i.e., clockwise (CW) or

counterclockwise (CCW), counterbalanced across participants. We also included no-error trials

(Experiment 1) by presenting a clamped feedback that followed a path directly to the target (0°

clamp; Fig. 2A). The nature of the clamp manipulation was described in detail to the participant,

and they were explicitly instructed strictly to ignore the feedback, aiming their reach directly toward

the target on every trial. These instructions were designed to emphasize that the participant did

not control the cursor position, and that they should always attempt to reach directly to the target.

The instructions of the task were reinforced by the presentation of short video animations to

demonstrate how the CSs would serve as imperative signals, as well as to show the invariant

direction of the clamped feedback.

Experimental protocol

Both Experiments 1 and 2 included two phases: acquisition and probe. During the acquisition

phase, clamped feedback was presented on each trial, serving as the US. During the probe

phase, the clamped feedback was not presented. In both phases, the participants were instructed

to reach straight to the target as soon as the imperative CS appeared. Note that we opted to not

include baseline reaching blocks prior to the start of the acquisition phases to avoid introducing

any incidental associations between the target, movement plan, and any other contextual

variables. A ~1 minute break was provided in the middle of the experiment.

Experiment 1: Differential conditioning

Experiment 1 (N=16) was designed to test differential conditioning in the context of a sensorimotor

adaptation task. The session consisted of 800 trials: 600 acquisition trials followed by 200 probe

trials (Fig. 2A, 2B). One of two CSs (tone or light) was presented on each trial, serving as the

imperative for the reaching response. During the acquisition phase, one CS was paired with a 15°

clamped error feedback (CS+ condition) and the other CS was paired with a 0° clamped feedback

(CS- condition). Each CS was presented on 50% of the trials, and the assignment of the tone and

light to the CS+ and CS- was counterbalanced across participants. During the probe phase, each

CS was presented alone on half of the trials, and there was no visual feedback. For both the

acquisition and probe phases, the CS+ and CS- trials were interleaved in a pseudo-random order

that was unique to each participant. To ensure that the participant would not be able to predict

the CS type, the generated trial sequence for each participant was assessed to verify that there

was no significant lag-1 autocorrelation in the time series. At the end of the experimental session,

we assessed whether participants were aware of the contingency between each CS and its

associated feedback, asking, "did you identify any pattern in the experiment?" in a free-response

survey.

<u>Experiment 2: Compound conditioning</u>

The adaptation task was modified in Experiment 2 (N=22) to provide a test of compound

conditioning. The procedure was mostly the same as that used in Experiment 1, with the following

changes. First, the session consisted of 600 acquisition trials and 300 probe trials (Fig. 3A,3B).

Second, a compound CS, consisting of both the tone and light, served as the imperative

throughout the acquisition phase. Moreover, the compound CS was always paired with 15°

clamped feedback. Lastly, the probe phase, in which the visual feedback was removed, consisted

of 100 trials each of the compound CS, tone alone CS, and light alone CS.

Model simulations

Trial-by-trial reach angles were simulated using two canonical learning models: the state-space

model and the Rescorla-Wagner model. The former was employed to represent the internal model

framework, and the latter to represent the classical conditioning framework. To illustrate the

predictions of the models, we used a single set of fixed parameters for each model in all

simulations (Figs. 2D, 2F, 3D). For the Rescorla-Wagner model, we chose parameters that result

in qualitatively similar effects to the experimental results. No such parameters exist for the state-

space model. To demonstrate that the conditioning effects predicted by the Rescorla-Wagner

model are qualitatively independent of the particular parameters used in the simulations, and that

they cannot be captured by the state-space model, we also simulated behavior over a wide range

of parameter values to generate various predicted effect sizes (Fig. S1, see details below).

In the standard state-space model of motor adaptation, the state of the internal model (x)

is updated according to the learning rule:

$$(1) \quad x_{n+1} = Ax_n + Be_n$$

where e is the sensory prediction error experienced on trial n, A is the retention factor, and B is

the learning rate. With clamped feedback, the sensory prediction error is fixed to one of two values

during the acquisition phase, either -15° on CS+ trials or 0° on CS- trials. The sensory prediction

error is set to 0° on all trials in the probe phase. For the simulations presented in Figures 2D, 2F

and 3D, we set the A to a value of 0.9 and B to 0.15.

As an alternative to the state-space model, the learning function during adaptation can be

described with a variant of the Rescorla-Wagner model. Here, the associative strengths, V, of the

conditioning stimuli are updated according to the following equations:

$$(2) \quad V(S)_n = \sum_{i=1}^{nS} V(s)_i$$

$$(3) \quad \partial(s)_n = \alpha(s) \cdot (\lambda - V(S)_n)$$

$$(4) \quad V(s)_{n+1} = V(s)_n + \beta \partial(s)_n$$

V(*S*) in Equation 2 represents the sum of associative strengths for each individual conditioning

stimulus *V*(*s*) over the number of stimuli *nS* presented on trial *n*. The value of ∂ reflects the

prediction error (Eq. 3), which is dependent on the salience of stimulus s, *α*(*s*), the asymptotic

learning level for the US, *λ*, and the current total associative strength, *V*(*S*). Equation 4 describes

the update process for the associative strength of each stimulus *s*, where *β* represents the

learning rate. Note that the summation in Equation 2 is used in the computation of the prediction

error (Eq. 3), reflecting the additivity (or competition) principle.

In typical simulations of classical conditioning behaviors such as eyeblink conditioning, *λ*

and *V* are normalized and the dependent variable is expressed as a percentage of trials in which

a conditioned response occurs. However, the Rescorla-Wagner model can, in principle, compute

any dependent variable (e.g., reaching angle). For all simulations, we set *λ* to 20° (similar to the

observed implicit adaptation asymptotes in other studies, Bond and Taylor, 2015; Morehead et

al., 2017; Kim et al., 2018, 2019), *β* to 0.2, and the three salience parameters (*α*'s) for the

movement plan CS, tone CS, and light CS, to, respectively, 0.6, 0.0009, and 0.0009. These

divergent salience values reflect our assumption that most of the associative strength of the US

would be absorbed by the movement plan CS, given its central relevance to the task goal of

reaching to a target. The remainder of potential associative strength to the US is thus available

for the tone and light CSs.

Although the two models yield many similar features of learning (e.g., sharing the Markov

property, producing exponential-family learning curves, etc.), the additional parameters in the

Rescorla-Wagner model allow it to capture the effects of differential and compound conditioning,

should these processes be operative in our experiments. For example, in the differential

conditioning case, modifying the parameters of the Rescorla-Wagner model could produce

changes in the magnitude of both the within-trial "Pavlovian" effects (i.e., the modulation of

behavior in response to the current CS presented, or "trial *n*" effects) and across-trial adaptation

effects (i.e., the modulation of behavior in response to the current state of learning, or "trial *n-1*"

effects). However, the Rescorla-Wagner model will always yield unique responses to the CS+

and CS- given non-zero salience parameters. In contrast, no combination of the *A* and *B*

parameters in the state-space model will produce differential responses to the tone and light CSs.

To illustrate how the particular choice of parameters in the Rescorla-Wagner model influences

the predicted conditioning effects, we simulated behavior using various combinations of the free

parameters. We chose several combinations of values for β and λ, and, for each combination, a

wide range of values for α_{Plan}, α_{tone} and α_{light}. (For simplicity, we constrained α_{tone} and α_{light} to the

same value.) Figure S1 displays heatmaps resulting from these simulations, with the color of each

cell corresponding to the simulated difference in the trial-by-trial change in hand angle between

CS+ and CS- trials.

To simulate the dynamics of the adaptation and Pavlovian effects in differential

conditioning (Fig. S3), and to compare it to the dynamics observed in the actual data, we

simulated the hand angle during acquisition (600 trials) according to the Rescorla-Wagner model

($V(s)$ in Eqs. 2-4), based on the schedules of CS+ and CS- trials that were presented to the

participants (16 simulated time courses). We then binned the simulated hand angle into epochs

of 50 trials. For each bin, we performed the same trial-by-trial multiple regression analysis we

performed on the actual data (Fig. S3, see *Data analysis* below). The simulated hand angle was

the dependent variable, and the identity of the previous (*n*-1) and current (*n*) CS were separable

predictor variables (z-scored) that, including an interaction term, competed for variance. Figure

S3 presents the mean±SEM regression β weights of all simulated time courses for the previous

(adaptation effect) and current (Pavlovian effect) predictors.

Model fitting and comparison

We conducted a *post-hoc* model comparison analysis (Fig. S2). In this analysis, we fit the

standard state-space model (Eq. 1) and the Rescorla-Wagner model (Eqs. 2-4) to participants'

hand angle time course data from Experiment 1. The two models were fit by minimizing the sum

of squared residuals between the measured and modeled movement data, using the MATLAB

function *fmincon*. To avoid local minima, 200 randomized sets of initial parameter values were

used during fitting and the best fit of each model was selected for model comparison. Models

were compared using both the sum of squared residuals and the Akaike Information Criterion

approximated on the residuals (AIC; Akaike, 1974). All free parameters were bound at [0, 1], with

the exception of λ, which was bounded at [0, 60]. We did not perform model fitting for Experiment

2 due to the fact that each trial was identical in the acquisition phase. As such, the salience

parameters in the Rescorla-Wagner model for the light, tone, and plan are unidentifiable, and both

models make indistinguishable behavioral predictions during acquisition.

Data analysis

The recorded position of the digitizing stylus was analyzed using custom-written MATLAB scripts.

Our main analyses focused on the reach direction (hand angle) and, for learning, the trial-by-trial

changes in hand angle (Δ hand angle). Hand angle was defined by two imaginary lines, one from

the start position to the target and the other from the start position to the hand position at maximum

movement velocity.

Trials in which the hand angle was larger than 100° off from the target, or in which the

trial-to-trial change in hand angle was larger than 25°, were considered outliers and not included

in the analyses. These outliers constituted 0.03% and 0.16% of all trials in Experiments 1 and 2,

respectively. For the change in hand angle analysis, but not for the presentation of hand angle

time courses, we excluded trials in which the reaction time exceeded 400 ms and/or movement

time exceeded 300 ms (Experiment 1: 11% of acquisition trials, 4.3% of probe trials; Experiment

2: 3.4% of acquisition trials, 4.9% of probe trials).

For all analyses and to visualize the results, the sign of the hand angle was flipped for

participants who experienced a CCW clamp, such that a positive hand angle is in the direction of

expected adaptation (i.e., opposite the direction of the perturbed feedback). Moreover, the hand

angle on the first acquisition trial was treated as the baseline reaching angle and subtracted from

the hand angle on all subsequent trials. (We note that the results remain unchanged in terms of

statistical comparisons if this baseline subtraction step is omitted.)

In Experiment 1, the primary analyses examined how the Δ hand angle was influenced by

the CS type (CS+ vs CS-), either in terms of the previous trial (n-1, adaptation effect) or current

trial (n, Pavlovian effect). For each participant and phase, we calculated the average Δ hand angle

for four types of trials: CS+ trials that follow CS+ trials, CS- after CS+, CS+ after CS-, and CS-

after CS-. For each phase, a two-way repeated-measures ANOVA was conducted with two within-

participant independent factors, the CS on trial n-1 and the CS on trial n, each with two levels,

CS+ and CS-. The Δ hand angle was the dependent variable in the two ANOVAs. To examine

the dynamics of the adaptation and Pavlovian effects (Fig. S3), we did the same trial-by-trial

regression analysis described above for the simulated hand angle (see Model simulations) with

the exception that the dependent variable was now the actual hand angle data of each participant.

To assess whether the Pavlovian effect is influenced by the awareness about the

contingency between each CS and its respective feedback, we divided the participants based on

their responses to the post experiment questionnaire (see Experimental protocol): 7 out of the 16

participants stated the correct contingency between the CS and the visual feedback, and were

thus considered the "aware sub-group." The rest of the participants (N=9) reported that they did

not identify any pattern related to the CS-US contingency and were considered the "unaware sub-

group." Independent two-sample t tests were used to compare the Pavlovian effects between

these groups during the acquisition and probe phases of the experiment. For these tests we also

report Bayes factor BF_{10}, the ratio of the likelihood of the alternative hypothesis (H_1) over the null

hypothesis (H_0) (Kass and Raftery, 1995).

Feedback correction was operationalized as the difference between the hand angle

measured at the radial distance to the target and at 50 ms after movement initiation. We estimated

the mean and standard deviation of feedback correction across all acquisition trials in all of the

participants in Experiment 1. In addition, we calculated, for each participant, the mean feedback

correction for each of the CS+ and CS- trials, and used a paired-sample *t* test to examine within-

participant changes in feedback correction between the two trial types.

In Experiment 2, the analysis focused on the probe phase in which there was no visual

feedback, comparing hand angles in response to the three imperatives (compound CS, light

alone, tone alone). We compared the Δ hand angle to the three CSs on trial *n* regardless of the

CS presented on trial *n-1*. We fit a one-way repeated-measures ANOVA, with the Δ hand angle

as the dependent variable, and the CS type as the within-participant independent variable.

The additivity principle of the Rescorla-Wagner model states that the association strengths

of each element of the compound CS will compete for associative strength with respect to the US

(Eqs. 2-4). This principle was tested in our data by examining the correlation between the Δ hand

angle associated with the tone and light CSs in Experiment 2.

Figure S1. Simulations of Pavlovian effects in differential conditioning. Simulations of current trial CS effect, i.e. the difference between the change in hand angle on CS+ and CS- trials, for the acquisition (top panel) and probe (bottom panel) phases of Experiment 1, as predicted by the state-space (left) and Rescorla-Wagner (right) models. Contours represent mean±SEM of the Pavlovian effect from Experiment 1.

Figure S2. Model comparison. The sum of squares residuals (SSR) and Akaike Information Criterion
(AIC) difference between Rescorla-Wagner (RW) and the state-space (SS) models for each participant of
Experiment 1.

Figure S3. Dynamics of adaptation and Pavlovian effects. Time courses of the mean weights
(regression *β*) of the adaptation (trial *n-1*, purple) and Pavlovian (trial *n*, cyan) effects in Experiment 1 as
predicted by the Rescorla-Wagner model (left), and the actual experimental results (right). Shaded regions
represent SEM.

ACKNOWLEDGEMENTS

We would like to thank the CognAc Lab (UC Berkeley) for helpful discussions. RBI is funded by

the NIH (NS116883 and NS105839). JAT is funded by the NIH (grant NS084948).

REFERENCES

Addou T, Krouchev N, Kalaska JF. 2011. Colored context cues can facilitate the ability to learn

and to switch between multiple dynamical force fields. *Journal of Neurophysiology*

106:163–183. doi:10.1152/jn.00869.2010

Akaike H. 1974. A new look at the statistical model identification. *IEEE Transactions on*

Automatic Control **19**:716–723. doi:10.1109/TAC.1974.1100705

Albert ST, Shadmehr R. 2016. The Neural Feedback Response to Error As a Teaching Signal

for the Motor Learning System. *J Neurosci* **36**:4832–4845.

doi:10.1523/JNEUROSCI.0159-16.2016

Albus JS. 1971. A theory of cerebellar function. *Mathematical Biosciences* **10**:25–61.

doi:10.1016/0025-5564(71)90051-4

Avraham G, Keizman M, Shmuelof L. 2019. Environmental consistency modulation of error

sensitivity during motor adaptation is explicitly controlled. *Journal of Neurophysiology*

123:57–69. doi:10.1152/jn.00080.2019

Avraham G, Morehead JR, Kim HE, Ivry RB. 2020. Re-exposure to a sensorimotor perturbation

produces opposite effects on explicit and implicit learning processes. *bioRxiv*

2020.07.16.205609. doi:10.1101/2020.07.16.205609

Blakemore S-J, Frith CD, Wolpert DM. 2001. The cerebellum is involved in predicting the

sensory consequences of action. *NeuroReport* **12**:1879–1884.

Bond KM, Taylor JA. 2015. Flexible explicit but rigid implicit learning in a visuomotor adaptation

792   task. *Journal of Neurophysiology* **113**:3836–3849. doi:10.1152/jn.00009.2015

793   Boyden ES, Katoh A, Raymond JL. 2004. CEREBELLUM-DEPENDENT LEARNING: The Role

794   of Multiple Plasticity Mechanisms. *Annual Review of Neuroscience* **27**:581–609.

795   doi:10.1146/annurev.neuro.27.070203.144238

796   Brainard DH. 1997. The Psychophysics Toolbox. *Spatial Vision* **10**:433–436.

797   doi:10.1163/156856897X00357

798   Brembs B, Heisenberg M. 2000. The Operant and the Classical in Conditioned Orientation of

799   Drosophila melanogaster at the Flight Simulator. *Learn Mem* **7**:104–115.

800   doi:10.1101/lm.7.2.104

801   Brembs B, Plendl W. 2008. Double Dissociation of PKC and AC Manipulations on Operant and

802   Classical Learning in Drosophila. *Current Biology* **18**:1168–1171.

803   doi:10.1016/j.cub.2008.07.041

804   Brennan AE, Smith MA. 2015. The Decay of Motor Memories Is Independent of Context

805   Change Detection. *PLOS Computational Biology* **11**:e1004278.

806   doi:10.1371/journal.pcbi.1004278

807   Brudner SN, Kethidi N, Graeupner D, Ivry RB, Taylor JA. 2016. Delayed feedback during

808   sensorimotor learning selectively disrupts adaptation but not strategy use. *Journal of*

809   *Neurophysiology* **115**:1499–1511. doi:10.1152/jn.00066.2015

810   Cartoni E, Balleine B, Baldassarre G. 2016. Appetitive Pavlovian-instrumental Transfer: A

811   review. *Neuroscience & Biobehavioral Reviews* **71**:829–848.

812   doi:10.1016/j.neubiorev.2016.09.020

813   Clark RE, Manns JR, Squire LR. 2002. Classical conditioning, awareness, and brain systems.

814   *Trends in Cognitive Sciences* **6**:524–531. doi:10.1016/S1364-6613(02)02041-7

815   Clark RE, Squire LR. 1998. Classical Conditioning and Brain Systems: The Role of Awareness.

816   *Science* **280**:77–81. doi:10.1126/science.280.5360.77

Day KA, Roemmich RT, Taylor JA, Bastian AJ. 2016. Visuomotor Learning Generalizes Around

the Intended Movement. *eNeuro* **3**. doi:10.1523/ENEURO.0005-16.2016

Doll BB, Simon DA, Daw ND. 2012. The ubiquity of model-based reinforcement learning.

Current Opinion in Neurobiology, Decision making **22**:1075–1081.

doi:10.1016/j.conb.2012.08.003

Donchin O, Rabe K, Diedrichsen J, Lally N, Schoch B, Gizewski ER, Timmann D. 2011.

Cerebellar regions involved in adaptation to force field and visuomotor perturbation.

Journal of Neurophysiology **107**:134–147. doi:10.1152/jn.00007.2011

Gandolfo F, Mussa-Ivaldi FA, Bizzi E. 1996. Motor learning by field approximation. *PNAS*

93:3843–3846. doi:10.1073/pnas.93.9.3843

Gao Z, Proietti-Onori M, Lin Z, ten Brinke MM, Boele H-J, Potters J-W, Ruigrok TJH, Hoebeek

FE, De Zeeuw CI. 2016. Excitatory Cerebellar Nucleocortical Circuit Provides Internal

Amplification during Associative Conditioning. *Neuron* **89**:645–657.

doi:10.1016/j.neuron.2016.01.008

Garcia KS, Steele PM, Mauk MD. 1999. Cerebellar Cortex Lesions Prevent Acquisition of

Conditioned Eyelid Responses. *J Neurosci* **19**:10940–10947.

doi:10.1523/JNEUROSCI.19-24-10940.1999

Gerwig M, Kolb FP, Timmann D. 2007. The involvement of the human cerebellum in eyeblink

conditioning. *Cerebellum* **6**:38. doi:10.1080/14734220701225904

Hadjiosif AM, Krakauer JW, Haith AM. 2020. Did we get sensorimotor adaptation wrong?

Implicit adaptation as direct policy updating rather than forward-model-based learning.

bioRxiv 2020.01.22.914473. doi:10.1101/2020.01.22.914473

Haith AM, Huberdeau DM, Krakauer JW. 2015. The Influence of Movement Preparation Time

on the Expression of Visuomotor Learning and Savings. *J Neurosci* **35**:5109–5117.

doi:10.1523/JNEUROSCI.3869-14.2015

Heess N, TB D, Sriram S, Lemmon J, Merel J, Wayne G, Tassa Y, Erez T, Wang Z, Eslami

SMA, Riedmiller M, Silver D. 2017. Emergence of Locomotion Behaviours in Rich

Environments. *arXiv:170702286 [cs]*.

Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R. 2018. Encoding of error and learning to

correct that error by the Purkinje cells of the cerebellum. *Nature Neuroscience* **21**:736–

743. doi:10.1038/s41593-018-0136-y

Hirashima M, Nozaki D. 2012. Distinct Motor Plans Form and Retrieve Distinct Motor Memories

for Physically Identical Movements. *Current Biology* **22**:432–436.

doi:10.1016/j.cub.2012.01.042

Houde JF, Jordan MI. 1998. Sensorimotor Adaptation in Speech Production. *Science*

279:1213–1216. doi:10.1126/science.279.5354.1213

Howard IS, Ingram JN, Franklin DW, Wolpert DM. 2012. Gone in 0.6 Seconds: The Encoding of

Motor Memories Depends on Recent Sensorimotor States. *J Neurosci* **32**:12756–12768.

doi:10.1523/JNEUROSCI.5909-11.2012

Howard IS, Wolpert DM, Franklin DW. 2015. The value of the follow-through derives from motor

learning depending on future actions. *Curr Biol* **25**:397–401.

doi:10.1016/j.cub.2014.12.037

Howard IS, Wolpert DM, Franklin DW. 2013. The effect of contextual cues on the encoding of

motor memories. *Journal of Neurophysiology* **109**:2632–2644.

doi:10.1152/jn.00773.2012

Hutter SA, Taylor JA. 2018. Relative sensitivity of explicit reaiming and implicit motor

adaptation. *Journal of Neurophysiology* **120**:2640–2648. doi:10.1152/jn.00283.2018

Hwang EJ, Smith MA, Shadmehr R. 2006. Dissociable effects of the implicit and explicit

memory systems on learning control of reaching. *Exp Brain Res* **173**:425–437.

doi:10.1007/s00221-006-0391-0

Izawa J, Criscimagna-Hemminger SE, Shadmehr R. 2012. Cerebellar Contributions to Reach

Adaptation and Learning Sensory Consequences of Action. *J Neurosci* **32**:4230–4239.

doi:10.1523/JNEUROSCI.6353-11.2012

Karniel A, Mussa-Ivaldi FA. 2002. Does the motor control system use multiple models and

context switching to cope with a variable environment? *Exp Brain Res* **143**:520–524.

doi:10.1007/s00221-002-1054-4

Kass RE, Raftery AE. 1995. Bayes Factors. *Journal of the American Statistical Association*

90:773–795. doi:10.1080/01621459.1995.10476572

Kawato M, Gomi H. 1992. A computational model of four regions of the cerebellum based on

feedback-error learning. *Biol Cybern* **68**:95–103. doi:10.1007/BF00201431

Khilkevich A, Zambrano J, Richards M-M, Mauk MD. 2018. Cerebellar implementation of

movement sequences through feedback. *eLife* **7**:e37443. doi:10.7554/eLife.37443

Kim HE, Morehead JR, Parvin DE, Moazzezi R, Ivry RB. 2018. Invariant errors reveal limitations

in motor correction rather than constraints on error sensitivity. *Communications Biology*

1:19. doi:10.1038/s42003-018-0021-y

Kim HE, Parvin DE, Ivry RB. 2019. The influence of task outcome on implicit motor learning.

eLife **8**:e39882. doi:10.7554/eLife.39882

Kim JJ, Thompson RE. 1997. Cerebellar circuits and synaptic mechanisms involved in classical

eyeblink conditioning. *Trends in Neurosciences* **20**:177–181. doi:10.1016/S0166-

2236(96)10081-3

Kitazawa S, Kohno T, Uka T. 1995. Effects of delayed visual information on the rate and amount

of prism adaptation in the human. *J Neurosci* **15**:7644–7652.

doi:10.1523/JNEUROSCI.15-11-07644.1995

Kojima Y, Iwamoto Y, Yoshida K. 2004. Memory of Learning Facilitates Saccadic Adaptation in

the Monkey. *J Neurosci* **24**:7531–7539. doi:10.1523/JNEUROSCI.1741-04.2004

Kojima Y, Soetedjo R, Fuchs AF. 2010. Changes in Simple Spike Activity of Some Purkinje

Cells in the Oculomotor Vermis during Saccade Adaptation Are Appropriate to

Participate in Motor Learning. *J Neurosci* **30**:3715–3727.

doi:10.1523/JNEUROSCI.4953-09.2010

Krakauer JW, Hadjiosif AM, Xu J, Wong AL, Haith AM. 2019. Motor LearningComprehensive

Physiology. American Cancer Society. pp. 613–663. doi:10.1002/cphy.c170043

Leow L-A, Marinovic W, Rugy A de, Carroll TJ. 2018. Task errors contribute to implicit

aftereffects in sensorimotor adaptation. *European Journal of Neuroscience* **48**:3397–

3409. doi:10.1111/ejn.14213

Lorenzetti FD, Mozzachiodi R, Baxter DA, Byrne JH. 2006. Classical and operant conditioning

differentially modify the intrinsic properties of an identified neuron. *Nature Neuroscience*

9:17–19. doi:10.1038/nn1593

Marr D. 1969. A theory of cerebellar cortex. *The Journal of Physiology* **202**:437–470.

doi:10.1113/jphysiol.1969.sp008820

Mazzoni P, Krakauer JW. 2006. An Implicit Plan Overrides an Explicit Strategy during

Visuomotor Adaptation. *J Neurosci* **26**:3642–3645. doi:10.1523/JNEUROSCI.5317-

05.2006

McDougle SD, Bond KM, Taylor JA. 2017. Implications of plan-based generalization in

sensorimotor adaptation. *Journal of Neurophysiology* **118**:383–393.

doi:10.1152/jn.00974.2016

McDougle SD, Bond KM, Taylor JA. 2015. Explicit and Implicit Processes Constitute the Fast

and Slow Processes of Sensorimotor Learning. *J Neurosci* **35**:9568–9579.

doi:10.1523/JNEUROSCI.5061-14.2015

McDougle SD, Ivry RB, Taylor JA. 2016. Taking Aim at the Cognitive Side of Learning in

Sensorimotor Adaptation Tasks. *Trends in Cognitive Sciences* **20**:535–544.

doi:10.1016/j.tics.2016.05.002

McDougle SD, Taylor JA. 2019. Dissociable cognitive strategies for sensorimotor learning.

Nature Communications **10**:40. doi:10.1038/s41467-018-07941-0

Morehead JR, Qasim SE, Crossley MJ, Ivry R. 2015. Savings upon Re-Aiming in Visuomotor

Adaptation. *J Neurosci* **35**:14386–14396. doi:10.1523/JNEUROSCI.1046-15.2015

Morehead JR, Taylor JA, Parvin DE, Ivry RB. 2017. Characteristics of Implicit Sensorimotor

Adaptation Revealed by Task-irrelevant Clamped Feedback. *Journal of Cognitive*

Neuroscience **29**:1061–1074. doi:10.1162/jocn_a_01108

Ohmae S, Medina JF. 2015. Climbing fibers encode a temporal-difference prediction error

during cerebellar learning in mice. *Nature Neuroscience* **18**:1798–1803.

doi:10.1038/nn.4167

Ostlund SB, Balleine BW. 2007. Orbitofrontal Cortex Mediates Outcome Encoding in Pavlovian

But Not Instrumental Conditioning. *J Neurosci* **27**:4819–4825.

doi:10.1523/JNEUROSCI.5443-06.2007

Osu R, Hirai S, Yoshioka T, Kawato M. 2004. Random presentation enables subjects to adapt

to two opposing forces on the hand. *Nature Neuroscience* **7**:111–112.

doi:10.1038/nn1184

Parvin DE, McDougle SD, Taylor JA, Ivry RB. 2018. Credit assignment in a motor decision

making task is influenced by agency and not sensorimotor prediction errors. *J Neurosci*

3601–17. doi:10.1523/JNEUROSCI.3601-17.2018

Pavlov IP. 1927. Conditioned reflexes, translated by GV Anrep. *London: Oxford.*

Popa LS, Streng ML, Hewitt AL, Ebner TJ. 2016. The Errors of Our Ways: Understanding Error

Representations in Cerebellar-Dependent Motor Learning. *Cerebellum* **15**:93–103.

doi:10.1007/s12311-015-0685-5

Rasmussen A, Jirenhed D-A, Hesslow G. 2008. Simple and Complex Spike Firing Patterns in

Purkinje Cells During Classical Conditioning. *Cerebellum* **7**:563. doi:10.1007/s12311-

008-0068-2

Raymond JL, Lisberger SG, Mauk MD. 1996. The Cerebellum: A Neuronal Learning Machine?

Science **272**:1126–1131. doi:10.1126/science.272.5265.1126

Reisman DS, Block HJ, Bastian AJ. 2005. Interlimb Coordination During Locomotion: What Can

be Adapted and Stored? *Journal of Neurophysiology* **94**:2403–2415.

doi:10.1152/jn.00089.2005

Rescorla R, Wagner A. 1972. A theory of Pavlovian conditioning: Variations in the effectiveness

of reinforcement and nonreinforcement. *Classical conditioning II: Current research and*

theory **2**:64–69.

Schlerf J, Ivry RB, Diedrichsen J. 2012. Encoding of Sensory Prediction Errors in the Human

Cerebellum. *J Neurosci* **32**:4913–4922. doi:10.1523/JNEUROSCI.4504-11.2012

Schlerf JE, Xu J, Klemfuss NM, Griffiths TL, Ivry RB. 2012. Individuals with cerebellar

degeneration show similar adaptation deficits with large and small visuomotor errors.

Journal of Neurophysiology **109**:1164–1173. doi:10.1152/jn.00654.2011

Schneiderman N, Gormezano I. 1964. Conditioning of the nictitating membrane of the rabbit as

a function of CS-US interval. *Journal of Comparative and Physiological Psychology*

57:188–195. doi:10.1037/h0043419

Schween R, Hegele M. 2017. Feedback delay attenuates implicit but facilitates explicit

adjustments to a visuomotor rotation. *Neurobiology of Learning and Memory* **140**:124–

133. doi:10.1016/j.nlm.2017.02.015

Schween R, McDougle SD, Hegele M, Taylor JA. 2020. Assessing explicit strategies in force

field adaptation. *Journal of Neurophysiology* **123**:1552–1565. doi:10.1152/jn.00427.2019

Shadmehr R, Krakauer JW. 2008. A computational neuroanatomy for motor control. *Exp Brain*

Res **185**:359–381. doi:10.1007/s00221-008-1280-5

Shadmehr R, Mussa-Ivaldi FA. 1994. Adaptive representation of dynamics during learning of a

motor task. *J Neurosci* **14**:3208–3224. doi:10.1523/JNEUROSCI.14-05-03208.1994

Shadmehr R, Smith MA, Krakauer JW. 2010. Error Correction, Sensory Prediction, and

Adaptation in Motor Control. *Annual Review of Neuroscience* **33**:89–108.

doi:10.1146/annurev-neuro-060909-153135

Sheahan HR, Franklin DW, Wolpert DM. 2016. Motor Planning, Not Execution, Separates Motor

Memories. *Neuron* **92**:773–779. doi:10.1016/j.neuron.2016.10.017

Sheahan HR, Ingram JN, Žalalytė GM, Wolpert DM. 2018. Imagery of movements immediately

following performance allows learning of motor skills that interfere. *Scientific Reports*

8:14330. doi:10.1038/s41598-018-32606-9

Smith MA, Ghazizadeh A, Shadmehr R. 2006. Interacting Adaptive Processes with Different

Timescales Underlie Short-Term Motor Learning. *PLOS Biology* **4**:e179.

doi:10.1371/journal.pbio.0040179

Smith MC, Coleman SR, Gormezano I. 1969. Classical conditioning of the rabbit's nictitating

membrane response at backward, simultaneous, and forward CS-US intervals. *Journal

of Comparative and Physiological Psychology* **69**:226–231. doi:10.1037/h0028212

Soetedjo R, Kojima Y, Fuchs AF. 2008. Complex Spike Activity in the Oculomotor Vermis of the

Cerebellum: A Vectorial Error Signal for Saccade Motor Learning? *Journal of

Neurophysiology* **100**:1949–1966. doi:10.1152/jn.90526.2008

Telgen S, Parvin D, Diedrichsen J. 2014. Mirror Reversal and Visual Rotation Are Learned and

Consolidated via Separate Mechanisms: Recalibrating or Learning De Novo? *J Neurosci*

34:13768–13779. doi:10.1523/JNEUROSCI.5306-13.2014

Tsay JS, Parvin DE, Ivry RB. 2020. Continuous Reports of Sensed Hand Position During

Sensorimotor Adaptation. *bioRxiv* 2020.04.29.068197. doi:10.1101/2020.04.29.068197

Verstynen T, Sabes PN. 2011. How Each Movement Changes the Next: An Experimental and

Theoretical Study of Fast Adaptive Priors in Reaching. *J Neurosci* **31**:10050–10059.

doi:10.1523/JNEUROSCI.6525-10.2011

Wainscott SK, Donchin O, Shadmehr R. 2005. Internal Models and Contextual Cues: Encoding

Serial Order and Direction of Movement. *Journal of Neurophysiology* **93**:786–800.

doi:10.1152/jn.00240.2004

Welch RB, Abel MR. 1970. The generality of the "target-pointing effect" in prism adaptation.

Psychon Sci **20**:226–227. doi:10.3758/BF03329034

Wilterson SA, Taylor JA. 2020. Implicit visuomotor adaptation remains limited after several days

of training. *bioRxiv* 711598. doi:10.1101/711598

Wolpert DM, Flanagan JR. 2001. Motor prediction. *Current Biology* **11**:R729–R732.

doi:10.1016/S0960-9822(01)00432-8

Wolpert DM, Ghahramani Z. 2000. Computational principles of movement neuroscience. *Nature*

Neuroscience **3**:1212–1217. doi:10.1038/81497

Wolpert DM, Ghahramani Z, Jordan MI. 1995. An internal model for sensorimotor integration.

Science **269**:1880–1882. doi:10.1126/science.7569931

Wolpert DM, Miall RC, Kawato M. 1998. Internal models in the cerebellum. *Trends in Cognitive*

Sciences **2**:338–347. doi:10.1016/S1364-6613(98)01221-2

1009

1010

  FIGURE CAPTIONS

Figure 1. Reaching Task. (A) Clamped-feedback induces implicit adaptation. (B) Adaptation

described as a conditioning process. (C) Participants reach to a single target (blue dot), moving

a cursor (white dot) from a start location (white circle). The target is displayed in a fixed location

for the entire experiment (location varied across participants), and the direction of the cursor

trajectory is fixed ("clamped") on all trials. After a random delay, a tone and/or a light (white frame)

is presented, serving as the movement imperative. This cue persists for the duration of the

movement.

1019

Figure 2. Experiment 1: Differential Conditioning. (A) Trial types. A 15° clamp (CW/CCW,

counterbalanced across participants) was associated with CS+ and a 0° clamp with CS-

(counterbalancing the associations with the tone and light across participants). (B) Experimental

protocol. CS+ (e.g., tone) and CS- (e.g. light) trials were randomly interleaved. Clamped feedback

(±15°/0°) was presented during the acquisition phase (white background), and no feedback was

provided during the probe phase (gray). (C) Mean hand angle time course (N=16). (D) Simulation

results for trial-by-trial change (Δ) in hand angle during acquisition. While the state-space model

predicts only an adaptation effect (main effect of trial *n-1*, dark vs light blue), the Rescorla-Wagner

model also predicts a Pavlovian effect (main effect of the presented CS on the current trial *n*, filled

vs empty bars). (E) Experimental results for trial-by-trial Δ hand angle during acquisition comport

with the Rescorla-Wagner model simulations but not the state-space model simulations (left

panel). The black outlined bar (right panel) presents the Pavlovian effect, i.e. the subtraction of

hand angle changes between CS+ and CS- trials. (F, G) Similar to (D) and (E), respectively, but

for the probe phase, showing a significant Pavlovian effect that is consistent with the Rescorla-

Wagner model. Shaded regions and error bars represent SEM. Dots represent individual

participants.

1036

Figure 3. Experiment 2: Compound Conditioning. (A, B) During acquisition, a 15° clamp was

associated with a compound CS: Tone and light were presented simultaneously as movement

imperatives. During the probe phase, no feedback was provided, and the CSs were presented

either together (Compound CS) or alone (Single CS; tone or light). (C) Mean hand angle time

course (N=22). (D) Simulation results for trial-by-trial Δ hand angle during the probe phase for the

compound CS (filled bar) and each of its elements (empty bars). Unlike the state-space model,

the Rescorla-Wagner model predicts a positive relative change for the compound CS on the

current trial n and a negative relative change for each element. (E) Experimental results for the Δ

hand angle during the probe are in line with the Rescorla-Wagner model. (F) Scatter plot showing

the relationship between individuals' responses to the elemental CSs (the dotted black line

represents the unity line). Shaded regions and error bars represent SEM. Dots represent

individual participants.

1049

1050 **Figure S1. Simulations of Pavlovian effects in differential conditioning.** Simulations of

1051 current trial CS effect, i.e. the difference between the change in hand angle on CS+ and CS- trials,

1052 for the acquisition (top panel) and probe (bottom panel) phases of Experiment 1, as predicted by

1053 the state-space (left) and Rescorla-Wagner (right) models. Contours represent mean±SEM of the

1054 Pavlovian effect from Experiment 1.

1055

1056 **Figure S2. Model comparison.** The sum of squares residuals (SSR) and Akaike Information

1057 Criterion (AIC) difference between Rescorla-Wagner (RW) and the state-space (SS) models for

1058 each participant of Experiment 1.

1059

1060   **Figure S3. Dynamics of adaptation and Pavlovian effects.** Time courses of the mean weights

1061   (regression β) of the adaptation (trial *n-1*, purple) and Pavlovian (trial *n*, cyan) effects in

1062   Experiment 1 as predicted by the Rescorla-Wagner model (left), and the actual experimental

1063   results (right). Shaded regions represent SEM.

Figure 1

Figure 2

Figure 3

Figure S1

Figure S2

Figure S3

Rescorla-Wagner simulation

Experimental results