Mapping Brain-Behavior Space Relationships Along the Psychosis Spectrum

Jie Lisa Ji[1,2], **Markus Helmer**[1], **Clara Fonteneau**[1], **Joshua B. Burt**[1,3], **Zailyn Tamayo**[1], **Jure Demšar**[4], **Brendan Adkinson**[1,2], **Aleksandar Savić**[5], **Katrin H. Preller**[6], **Flora Moujaes**[6], **Franz X. Vollenweider**[6], **William J. Martin**[7], **Grega Repovš**[4], **John D. Murray**[1,2,3], and **Alan Anticevic**[1,2,8,✉]

[1]Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA
[2]Interdepartmental Neuroscience Program, Yale University School of Medicine, New Haven, CT, USA
[3]Department of Physics, Yale University, New Haven, CT, USA
[4]Department of Psychology, University of Ljubljana, Ljubljana, Slovenia
[5]Department of Psychiatry, University of Zagreb, Zagreb, Croatia
[6]University Hospital for Psychiatry Zurich, Zurich, Switzerland
[7]The Janssen Pharmaceutical Companies of Johnson & Johnson
[8]Department of Psychology, Yale University School of Medicine, New Haven, CT, USA

Difficulties in advancing effective patient-specific therapies for psychiatric disorders highlight a need to develop a stable neurobiologically-grounded mapping between neural and symptom variation. This gap is particularly acute for psychosis-spectrum disorders (PSD). Here, in a sample of 436 cross-diagnostic PSD patients, we derived and replicated a dimensionality-reduced symptom space across hallmark psychopathology symptoms and cognitive deficits. In turn, these symptom axes mapped onto distinct, reproducible brain maps. Critically, we found that multivariate brain-behavior mapping techniques (e.g. canonical correlation analysis) do not produce stable results. Instead, we show that a univariate brain-behavioral space (BBS) can resolve stable individualized prediction. Finally, we show a proof-of-principle framework for relating personalized BBS metrics with molecular targets via serotonin and glutamate receptor manipulations and gene expression maps. Collectively, these results highlight a stable and data-driven BBS mapping across PSD, which offers an actionable path that can be iteratively optimized for personalized clinical biomarker endpoints.

psychosis spectrum | functional neuroimaging | symptoms mapping | molecular target selection | personalized patient selection | statistical learning | data reduction

Correspondence: *alan.anticevic@yale.edu & jielisa.ji@yale.edu*

Introduction

Mental health conditions cause profound disability with most treatments yielding limited efficacy across psychiatric symptoms (1–4). A key step towards developing more effective therapies for specific psychiatric symptoms is reliably mapping them onto underlying neural systems. This goal is particularly challenging because neuropsychiatric diagnostics still operate under "legacy" categorical constraints, which were not quantitatively informed by neural or symptom data. Critically, diagnostic systems in psychiatry [e.g. the Diagnostic and Statistical Manual of Mental Disorders (DSM) (5)] were built to aid clinical consensus, but were not designed to guide quantitative mapping of symptoms onto neural alterations (6, 7). Consequently, this system cannot optimally map onto patient-specific brain-behavioral alterations. This challenge is particularly evident along the psychosis spectrum disorders (PSD) where there is notable shared symptom

variation across what distinct DSM diagnostic categories, including schizophrenia (SZP), schizo-affective (SADP), bipolar disorder with psychosis (BPP). For instance, despite BPP being a distinct diagnosis, BPP patients exhibit similar but attenuated psychosis symptoms and neural alterations similar to SZP (e.g. thalamic functional connectivity (FC) (8)). It is essential to quantitatively map such shared clinical variation onto common neural alterations, to circumvent constraints for biomarker development (6, 7, 9) – a key goal for development of neurobiologically-informed personalized therapies (8, 9).

Recognizing the limitations of categorical frameworks, the NIMH's Research Domain Criteria (RDoC) initiative introduced dimensional mapping of functional domains on to neural circuits (10). This motivated cross-diagnostic multi-site studies for mapping PSD symptom and neural variation (11–14). Multivariate neuro-behavioral analyses across PSD and mood spectra reported brain-behavioral relationships across diagnoses, with the goal of informing individualized treatment (15). These studies attempted to address the challenge of moving beyond traditional *a priori* clinical scales, which provide composite scores (16) that may not optimally capture neural variation (7, 17). For instance, despite many data-driven dimensionality-reduction symptom studies (18–24), a common approach in PSD neuroimaging research is still to sum "positive" or "negative" psychosis symptoms into a single score for relating to neural measures (8, 25). Importantly, neural alterations in PSD may reflect a more complex weighted combination of symptoms than *a priori* composite scores.

While multivariate neuro-behavioral studies have addressed this, such studies face the risk of failing to replicate due to overfitting (26), arising from high dimensionality of behavioral and neural features and a comparatively limited number of subjects (27). Notably, current state-of-the-art large-scale clinical neuroimaging studies have target enrollment totals of ∼200-600 subjects (https://www.humanconnectome.org/disease-studies). We therefore wanted to test whether a stable, reproducible neuro-behavioral geometry can be derived with a sample size that is

on par with current large-scale studies.

We hypothesized that a linearly-weighted low-dimensional symptom solution (capturing key disease-relevant information) produces a stable and robust univariate brain-behavioral mapping. Indeed, recent work used dimensionality reduction methods successfully to compute a neural mapping across canonical SZP symptoms (23). However, it remains unknown if this approach generalizes across PSD. Moreover, it is unknown if incorporating cognitive assessment, a hallmark and untreated PSD deficit (17), explains neural feature variation that is distinct from canonical PSD symptoms. Finally, prior work has not tested if a low-dimensional symptom-neural mapping can be predictive at the single patient level – a prerequisite for individualized clinical endpoints.

To inform these gaps, we first tested two key questions: i) Can data-reduction methods reliably reveal symptom axes across PSD that include both canonical symptoms and cognitive deficits? ii) Do these lower-dimensional symptom axes map onto a reproducible brain-behavioral solution across PSD? Specifically, we combined fMRI-derived resting-state measures with psychosis and cognitive symptoms (28, 29) obtained from a public multi-site cohort of 436 PSD patients and 202 healthy individuals collected by the Bipolar-Schizophrenia Network for Intermediate Phenotypes (BSNIP-1) consortium across 6 sites in North America (11). The dataset included included 150 patients formally diagnosed with BPP, 119 patients diagnosed SADP, and 167 patients diagnosed with SZP (**Table S2**). This cohort enabled cross-site symptom-neural comparisons across multiple psychiatric diagnostic categories, which we then mapped onto specific neural circuits. First, we tested if dimensionality reduction of PSD symptoms revealed a stable solution for individual patient prediction. Next, we tested if this low-dimensional symptom solution yields novel, stable, and statistically robust neural mapping compared to canonical composite symptom scores or DSM diagnoses. We then tested if the computed symptom-neural mapping is reproducible across symptom axes and actionable for individual patient prediction. Finally, we anchor the derived symptom-relevant neural maps by computing their similarity against mechanistically-informed neural maps. We used independently collected pharmacological fMRI maps from healthy adults in response to putative PSD receptor treatment targets (glutamate via ketamine and serotonin via LSD) (30, 31). We also computed transcriptomic maps from the Allen Human Brain Atlas (AHBA) (32, 33) for genes implicated in PSD. The primary purpose of this paper is to derive a comprehensive, stable, and reproducible symptom-neural geometry across symptom domains in chronic PSD, that can be resolved at the single-subject level and mechanistically linked to molecular benchmarks. This necessitated the development of novel methods and rigorous testing (as we found that existing multivariate methods are drastically underpowered (27)) which we have also presented in the paper for transparency and reproducibility. This approach can be iteratively improved upon to further elucidate the neural underpinnings of PSD variation, and expanded to apply to broader research

questions and psychiatric spectra, symptom domains, and neural features.

To our knowledge, no study to date has mapped a data-reduced symptom geometry encompassing a transdiagnostic PSD cohort across combined cognitive and psychopathology symptom domains, with demonstrable statistical stability and reproducibility at the single subject level. Additionally, this symptom geometry can be mapped robustly onto neural data to achieve reproducible group-level effects as well as individual-level precision following feature selection. Furthermore, while other other studies have evaluated relationships between neural maps and complementary molecular datasets (e.g. transcriptomics), no study to our knowledge has benchmarked single-subject selected neural maps using reproducible neuro-behavioral features against both pharmacological neuroimaging maps and gene expression maps that may inform single subject selection for targeted treatment. Collectively, this study used data-driven dimensionality-reduction methods to derive stable orthogonal symptom dimensions across 436 PSD patients and mapped them onto novel and stable neuroimaging features. These effects were then benchmarked against molecular imaging targets, outlining an actionable path towards making personalized recommendations for clinical endpoints.

Methods

Overall Data Collection and Study Design. We used publicly available behavioral and neural data from the Bipolar-Schizophrenia Network on Intermediate Phenotypes (BSNIP) consortium (11). All data were obtained from the National Data Archive (NDA) repository (https://nda.nih.gov/edit_collection.html?id=2274). Participants were collected at 6 sites across the United States. Full recruitment details are provided in prior publications (11, 34, 35). In brief, participants were recruited through advertising in Baltimore MD, Boston MA, Chicago IL, Dallas TX, and Hartford CT. All assessments were standardized across sites. Participants were excluded if they had i) a history of seizures or head injury resulting in >10 minutes loss of consciousness, ii) positive drug screen on the day of testing, iii) a diagnosis of substance abuse in the past 30 days or substance dependence in the past 6 months, iv) history of serious medical or neurological disorder that would likely affect cognitive functioning, v), history of serious medical or neurological disorder that would likely affect cognitive functioning, vi) insufficient English proficiency, or vii) an age-corrected Wide-Range Achievement Test (4th edition) reading test standard score <65. Additionally, participants were required to have had no change in medication and been clinically stable over the past month. Participants completed a SCID interview and were given a diagnosis via consensus from study clinicians; participants with an Axis 1 clinical psychosis diagnosis were additionally given assessments including the Positive and Negative Symptom Scale (PANSS)(29). Note that apart from the measures used in this paper, the full BSNIP dataset includes a rich characterization of measures from multiple modalities, including elec-

trophysiological, eye tracking, structural and diffusion neuroimaging, as well as a number of clinical batteries, which are not quantified in this study. We used data from a total of 638 participants with complete behavioral and neural data after preprocessing and quality control, including 202 healthy controls (CON), 167 patients diagnosed with schizophrenia (SZP), 119 patients with schizoaffective disorder (SADP), and 150 patients with bipolar disorder with psychosis (BPP). For full demographic and clinical characteristics of the sample see **Table S2**.

Neural Data Acquisition and Preprocessing. Participants completed a neural magnetic resonance imaging (MRI) scan at 3T, including resting-state functional blood-oxygen-level-dependent imaging (BOLD) and a magnetization-prepared rapid gradient-echo (MP-RAGE) sequence for T1 weighted data. Full details on scanners and acquisition parameters used at each of the sites have previously been described (36) and are summarized here in **Table S3**. Neuroimaging data were preprocessed using the Human Connectome Project (HCP) minimal preprocessing pipeline (37), adapted for compatibility with "legacy" data, which are now featured as a standard option in the HCP pipelines provided by our team (https://github.com/Washington-University/HCPpipelines/pull/156). These modifications to the HCP pipelines were necessary as the BSNIP data did not include a standard field map and did not incorporate a T2w high-resolution image. The adaptations for single-band BOLD acquisition have previously been described in detail (38).

A summary of the HCP pipelines is as follows: the T1-weighted structural images were first aligned by warping them to the standard Montreal Neurological Institute-152 (MNI-152) brain template in a single step, through a combination of linear and non-linear transformations via the FMRIB Software Library (FSL) linear image registration tool (FLIRT) and non-linear image registration tool (FNIRT) (39). Next, FreeSurfer's recon-all pipeline was used to segment brain-wide gray and white matter to produce individual cortical and subcortical anatomical segmentations (40). Cortical surface models were generated for pial and white matter boundaries as well as segmentation masks for each subcortical grey matter voxel. Using the pial and white matter surface boundaries, a 'cortical ribbon' was defined along with corresponding subcortical voxels, which were combined to generate the neural file in the Connectivity Informatics Technology Initiative (CIFTI) volume/surface 'grayordinate' space for each individual subject (37). BOLD data were motion-corrected by aligning to the middle frame of every run via FLIRT in the initial NIFTI volume space. In turn, a brain-mask was applied to exclude signal from non-brain tissue. Next, cortical BOLD data were converted to the CIFTI gray matter matrix by sampling from the anatomically-defined gray matter cortical ribbon and subsequently aligned to the HCP atlas using surface-based nonlinear deformation (37). Subcortical voxels were aligned to the MNI-152 atlas using whole-brain non-linear registration and then the Freesurfer-defined subcortical segmentation applied to isolate the sub-cortical grayordinate portion of the CIFTI space.

After the HCP minimal preprocessing pipelines, movement scrubbing was performed (41). As done in our prior work (42), all BOLD image frames with possible movement-induced artifactual fluctuations in intensity were flagged using two criteria: frame displacement (the sum of the displacement across all six rigid body movement correction parameters) exceeding 0.5 mm (assuming 50 mm cortical sphere radius) and/or the normalized root mean square (RMS) (calculated as the RMS of differences in intensity between the current and preceding frame, computed across all voxels and divided by the mean intensity) exceeding 1.6 times the median across scans. Any frame that met one or both of these criteria, as well as the frame immediately preceding and immediately following, were discarded from further preprocessing and analyses. Subjects with more than 50% frames flagged using these criteria were excluded. Next, a high-pass filter (threshold 0.008 Hz) was applied to the BOLD data to remove low frequency signals due to scanner drift. In-house Matlab code was used to calculate the average variation in BOLD signal in the ventricles, deep white matter, and across the whole grey matter, as well as movement parameters. These signals, as well as their first derivatives to account for delayed effects, were then regressed out of the grey matter BOLD time series as nuisance variables (as any change in the BOLD signal due to these variables are persistent and likely not reflecting neural activity) (43). It should be noted that using global signal regression to remove spatially persistent artifact is highly controversial in neuroimaging (44, 45), but it remains the field-wide gold standard (though see other recent and emerging approaches at (46, 47)).

Behavioral Symptom and Cognitive Data. The behavioral measures analyzed included the Positive and Negative Syndrome Scale (PANSS), an assessment of psychosis symptom severity (29), and the Brief Assessment of Cognition in Schizophrenia (BACS) battery, which provided an assessment of cognitive functioning (48). Only control subjects with complete BACS measures were used for analyses; PANSS symptom scores were imputed for control subjects for whom the PANSS had not been administered under the assumption that these subjects were asymptomatic. Only patient subjects with complete PANSS and BACS measures were used in analyses. The BACS scores used here are presented as standardized Z-scores normalized to the mean and standard deviation of the control group for each BSNIP site, as done in prior work (11). The full PANSS battery is conventionally divided into three sub-scales: Positive symptom scale (7 items), Negative symptom scale (7 items), and General Psychopathology symptom scale (16 items). The BACS battery consists of 6 individual sub-scores (48). In total, this yielded 36 symptom variables per participant. Effects of symptom variation across assessment sites have been rigorously characterized in prior publications (49). Nevertheless, we explicitly tested for site effects in our analyses, described in detail below.

Principal Component Analysis of Behavioral Symptom Measures. The principal component analysis (PCA) of behavioral data was computed by including all 36 symptom variables across all N=436 patients. Variables were first scaled to have unit variance across patients before running the PCA. Significance of the derived principal components (PCs) was computed via permutation testing. For each permutation, patient order was randomly shuffled for each symptom variable before re-computing PCA. This permutation was repeated 5,000 times to establish the null model. PCs which accounted for a proportion of variance that exceeded chance (p<0.05 across all 5000 permutations) were retained for further analysis.

Stability and Reproducibility of the Symptom PCA. Stability of the symptom PCA solution was assessed with within-sample cross-validation. To evaluate if there were site differences that uniquely drove the PCA solution, we performed a leave-site-out cross-validation analysis. Specifically, we re-ran the PCA using all patients except those from one site, which was held out. This process was repeated for each of the 6 sites. To further evaluate the stability of the derived PCA solutions, we performed 1,000 runs of *k*-fold cross-validation for values of *k* between 2 and 10. For each *k*-fold run, the full sample of patients was randomly divided into equally-sized *k* sets and a PCA was re-computed using subjects in *k*-1 sets (as the left-out set was used as the held-out sample).

For each run of leave-site-out and *k*-fold cross-validations significance was assessed via permutation testing as described above. The number of significant PCs and the total proportion of variance explained by these significant PCs remained highly stable across all runs (see **Fig. 2A** and **Fig. S6** and **Fig. S5**). Additionally, we compared observed and predicted PCA scores for the held-out sample of patients. Predicted scores for the held-out sample of patients were computed by weighing their raw symptom scores with the loadings from the PCA computed in all other patients. Observed scores for held-out patients were obtained from the original PCA computed on the full sample presented in the main text. The similarity between predicted and observed scores was high for all five significant PCs across all runs of leave-site-out and *k*-fold cross-validation, exceeding *r*=0.9 in most analyses (see **Fig. 2B-C** and **Fig. S6** and **S5**). Notably, the PCA solution did not show medication status or dosage effects (**Fig. S7**).

We also assessed the similarity of the PCA loadings using leave-site-out and 1,000 runs of 5-fold cross-validation frameworks (**Fig. 2D**). Importantly, this cross-validation was designed to test if the observed loadings remained stable (as opposed to predicted patient-level scores). The loadings for significant PCs from each leave-site-out PCA solution as well as each run of the 5-fold cross-validation were highly correlated (**Fig. 2D**.

We also assessed the reproducibility of the PCA solution using independent split-half samples. The leave-site-out and *k*-fold cross-validation PCA analyses by definition use overlapping samples of patients for each iteration. Therefore, we additionally conducted a full split-half replication using entirely non-overlapping sets of patients in each iteration. For each split-half iteration, the full patient sample was randomly divided into two sets with equal proportions of each of the three diagnostic groups (BPP, SADP, SZP). Then, a PCA was computed using each of the split-half patient samples. The loadings from the two PCA solutions were then evaluated for reproducibility. This process was repeated 1,000 times. The loadings for significant PCs were again highly similar even when comparing PCA solutions derived from completely non-overlapping patient samples (**Fig. 2D**).

To predict individual patient PC scores for the leave-one-out analysis (**Fig. S27A**), a PCA was computed using all patients except one held-out patient (N=435). In turn, the derived loadings were then used to compute the predicted PC scores for the left-out patient. This process was repeated until predicted PC scores were calculated for every patient. Finally, the predicted score for each patient was evaluated for reproducibility relative to the observed score obtained from the PCA solution computed using the full N=436 sample of patients. In addition, we computed an independent component analysis (ICA) to evaluate the consistency of the behavioral data reduction solution across methods (see **Methods**).

Global Brain Connectivity Calculation. Following preprocessing, the functional connectivity (FC) matrix was calculated for each participant by computing the Pearson's correlation between every grayordinate in the brain with all other grayordinates. A Fisher's r-to-Z transform was then applied. Global brain connectivity (GBC) was calculate by computing every grayordinate's mean FC strength with all other grayordinates (i.e. the mean, per row, across all columns of the FC matrix). GBC is a data-driven summary measure of connectedness that is unbiased with regards to the location of a possible alteration in connectivity (50) and is therefore a principled way for reducing the number of neural features while assessing neural variation across the entire brain.

$$GBC(x) = \frac{1}{N} \sum_{y=1}^{N} r_{xy} \tag{1}$$

- where $GBC(x)$ denotes the GBC value at grayordinate x;

- where N denotes the total number of grayordinates;

- where $\sum_{y=1}^{N}$ denotes the sum from $y = 1$ to $y = N$;

- where r_{xy} denotes the correlation between the time-series of grayordinates x and y;

For parcel-wise GBC maps (described below) we first computed the mean BOLD signal within each parcel (see section below for parcellation details) for each participant and then computed the pairwise FC between all parcels. Finally, to obtain the parcellated GBC metric we computed the mean FC for each parcel and all other parcels. This order of operations (first parcellating the dense data series and then computing GBC) was chosen because it resulted in stronger statistical

values due to increased within-parcel signal-to-noise of the BOLD data (**Fig. 4A**).

$$GBC(p) = \frac{1}{N} \sum_{q=1}^{N} r_{pq} \qquad \qquad \textbf{(2)}$$

- where $GBC(p)$ denotes the GBC value at parcel p;

- where N denotes the total number of parcels;

- where $\sum_{y=1}^{N}$ denotes the sum from $q = 1$ to $q = N$;

- where r_{pq} denotes the correlation between the time-series of parcels p and q;

Neural Data Reduction via Functional Brain-wide Parcellation. Here we applied a recently developed Cole-Anticevic Brain Network Parcellation (CAB-NP) parcellation (51), which defines functional networks and regions across cortex and subcortex that leveraged the Human Connectome Project's Multi-Modal Parcellation (MMP1.0) (51, 52). The final published CAB-NP 1.0 parcellation solution can be visualized via the Brain Analysis Library of Spatial maps and Atlases (BALSA) resource (https://balsa.wustl.edu/rrg5v) and downloaded from the public repository (https://github.com/ColeLab/ColeAnticevicNetPartition). The cortex component of the parcellation solution is comprised of 180 bilateral parcels (a total of 360 across both left and right hemispheres), consistent with the Human Connectome Project's Multi-Modal Parcellation (MMP1.0) (52). The subcortex component is comprised of 358 parcels defined using resting-state functional BOLD covariation with the cortical network solution (51).

Mass Univariate Symptom-Neural Mapping. Behavioral scores (*a priori* and PCA) were quantified in relation to individual GBC variation (either dense or parcellated) via a mass univariate regression procedure. The resulting map of regression coefficients reflected the strength of the relationship between patients' behavioral PC score and GBC at every neural location, across all 436 patients. The greater the magnitude of the coefficient for a given location, the stronger the statistical relationship between GBC and the behavioral variation across patients. The coefficients were then Z-scored for each map. Significance of the maps was assessed via non-parametric permutation testing, 2000 random shuffles with TFCE (53) type-I error-protection computed via the Permutation Analysis of Linear Models program (54).

Cross-validation and Split-half Replication of Mass Univariate Symptom-Neural Mapping. Stability of the symptom-neural mapping was assessed following the same cross-validation logic as described for the symptom-driven PCA solutions. Specifically, five-fold cross-validation was performed by first randomly partitioning all patients (N=436) into 5 subsets. Regression of the behavioral PC scores onto GBC across patients was performed while holding out 1/5 of

the patient sample (N=349). The correlation between the resulting neural coefficient map was then computed with the neural map obtained from the full sample calculation. For leave-site-out cross-validation, all subjects except those from one site were used when calculating the symptom-neural regression model. The resulting maps were compared to the map from the full sample regression. This process was repeated 6 times, each time leaving out subjects in one of the 6 sites.

Reproducibility of the symptom-neural mapping was assessed with 1,000 iterations of split-half replication. For each iteration, the full sample of subjects was first randomly split into two halves (referred to as H1 and H2) with the proportion of subjects in each diagnostic group (BPP, SADP, SZP) preserved within each half. For each iteration we used the H1 PCA solution and loadings to compute the predicted PCA scores for H2. In turn, the observed H1 scores were computed from a PCA loadings on the same H1 half-sample of patients. These H1 scores were then regressed against parcellated GBC for patients in H2. This coefficient GBC map reflects the strength of the relationship between the predicted PC score and GBC across H2 patients. Finally, the GBC coefficient maps derived from the H1 observed and H2 predicted PCA scores were then correlated for each PC axis. This process was then repeated 1,000 times and evaluated for reproducibility.

Principal Component Analysis of Neural Data. To evaluate the consistency of the neural GBC, we computed a PCA solution for the parcellated neural GBC data for all 202 control subjects as well as for all 436 patients (**Fig. S20**). The resulting neural GBC-derived PCs capture the striking consistency of the neural variance components irrespective of clinical status, which highlights that the bulk of the neural variance is not symptom-relevant. As with the behavioral PCA, significance of the neural PCA solution was assessed via permutation testing (1,000 random shuffles of parcels within subject). To evaluate the effect of shared neural variance on the symptom-neural solution, we parsed out common neural PCs from the PSD GBC data. We first conducted a PCA using the parcellated GBC data from all 436 PSD and 202 CON (a matrix with dimensions 638 subjects x 718 parcels). This GBC-PCA resulted in 637 independent GBC-PCs. Since PCs are orthogonal to each other, we then partialled out the variance attributable to GBC-PC1 from the PSD data by reconstructing the PSD GBC matrix using only scores and coefficients from the remaining 636 GBC-PCs ($G\hat{B}C_{woPC1}$). We then reran the univariate regression as described in **Fig. 3**, using the same five symptom PC scores across 436 PSD. We also repeated the symptom-neural regression with the first 2 GBC-PCs partialled out of the PSD data ($G\hat{B}C_{woPC1-2}$), with the first 3 PCs parsed out ($G\hat{B}C_{woPC1-3}$), and with the first 4 neural PCs parsed out ($G\hat{B}C_{woPC1-4}$).

Canonical Correlation Analysis. Canonical correlation analysis (CCA) is a multivariate statistical technique which examines simultaneously the relationships between multiple independent variables and multiple dependent variables

by computing linear combinations of each variable set that maximizes the correlations between the two sets (**Fig. 5A**). Here, the two variates used were symptom and neural features across all subjects. Each feature was Z-scored prior to computing the CCA solution. Given the size of the 'dense' neural feature space reduced the number of neural features in a principled manner via the described parcellation. This reduced the number of neural features to 180 symmetrized cortical parcels (i.e. GBC was averaged for each pair of analogous parcels in the left and right hemispheres). Critically, corresponding cortical parcels in the left and right hemispheres of this parcellation have been shown to be highly similar (33, 37). First, we computed the CCA solution using all 36 item-level symptom measures as behavioral features. In turn, we computed an additional CCA solution using scores from the significant 5 PCs.

Each symptom PC is a weighted linear composite of the original symptom measures and each behavioral CV is a weighted linear composite of the symptom PCs. Therefore, to compute the loadings of the 36 symptom measures on each of the behavioral CVs (computed using the symptom PCs), we multiplied the matrix of loadings from the CCA with the matrix of loadings from the PCA (**Fig. 5J**).

CCA Cross-validation and Split-half Replication. Stability of the CCA solution was assessed via cross-validation. Five-fold cross-validation of the CCA was performed by first randomly partitioning all patients (N=436) into 5 subsets. The CCA was then performed between neural and behavioral features using all but one of the subsets. The results of this CCA was then compared to the full sample CCA. **Fig. S18A-D** shows the comparisons across all 5 five-fold cross-validation runs of the CCA compared to the full model, including neural factor loadings, behavioral factor loadings, and projected symptom measure loadings. For leave-site-out cross-validation, all subjects except those from one site were used in the CCA. The resulting outputs were then compared to those from the full sample CCA. This process was repeated for all 6 sites. These data are shown in **Fig. S18E-H**.

For leave-one-out cross-validation of the CCA, one subject was held out as a CCA was performed using neural and behavioral features from the other 435 subjects. The loadings matrices Ψ and Θ from the CCA were then used to calculate the "predicted" neural and behavioral latent scores for all 5 canonical modes for the holdout subject. This was repeated for every subject, such that predicted neural and behavioral latent score matrices ($\hat{N}$ and $\hat{B}$) were computed, of the same dimensions as (N and B) respectively. The corresponding CVs in $\hat{N}$ and $\hat{B}$ were then correlated across subjects, as shown in **Fig. 5M**. If the CCA solution is stable, these correlations should be comparable to the canonical correlations of the full CCA (**Fig. 5D**).

Reproducibility of the CCA solution was assessed via split-half replication. This was performed by first randomly splitting the full sample into two halves (H1 and H2) with the proportion of subjects in each diagnostic group (BPP, SADP, SZP) preserved within each half. A CCA was then performed separately in each half and the resulting outputs were then

compared to each other. This process was repeated 1,000 times to obtain mean and standard deviation performance metrics. These data are shown in **Fig. 5L**.

Multivariate CCA Power Analysis. Multivariate power analyses to estimate the minimum sample size needed to sufficiently power a CCA were computed using methods described in (27), using the Generative Modeling of Multivariate Relationships tool (*gemmr*, **https://github.com/murraylab/gemmr** (v0.1.2)). Briefly, a model was built by: 1) Generating synthetic datasets for the two input data matrices, by sampling from a multivariate normal distribution with a joint covariance matrix that was structured to encode CCA solutions with specified properties; 2) Performing CCAs on these synthetic datasets. Because the joint covariance matrix is known, the true values of estimated association strength, weights, scores, and loadings of the CCA, as well as the errors for these four metrics, can also be computed. In addition, statistical power that the estimated association strength is different from 0 is determined through permutation testing; 3) Varying parameters of the generative model (number of features, assumed true between-set correlation, within-set variance structure for both datasets) the required sample size n_{req} is determined in each case such that statistical power reaches 90% and all of the above described error metrics fall to a target level of 10%; and 4) Fitting and validating a linear model to predict the required sample size n_{req} from parameters of the generative model. This linear model was then used to calculate n_{req} for CCA in three data scenarios: i) 718 neural vs. 5 symptom features; ii) 180 neural vs. 5 symptom features; iii) 12 neural vs. 5 symptom features.

Independent Replication Dataset. To illustrate the generalizability of the single-subject prediction results across independent datasets and DSM disorders, we use an independently collected dataset consisting of 30 patients with a formal diagnosis of schizophrenia (SZP) and 39 patients diagnosed with obsessive compulsive disorder (OCD). These patients were recruited via clinician referral and regional advertising and assessed at the Connecticut Mental Health Center in New Haven, CT. Demographic and symptom data are shown in **Table S4**.

The behavioral assessment for the replication dataset included the PANSS and the Penn Computerized Neurocognitive Battery (PennCNB)(55). Items from the PennCNB were matched to items from the BACS as follows: BACS digital sequencing (working memory) -> LNB (N-back Total score); BACS symbol coding (attention) -> Penn Continuous Performance Test); BACS token motor task (motor) -> CTAP; BACS Tower of London (abstraction and mental flexibility) -> PCET or RAVEN (Penn Conditional Exclusion Test); BACS verbal fluency (verbal ability) -> SPVRT (Penn Verbal Reasoning Test); BACS verbal memory test -> PLLT (Penn List Learning Test) (56).

Neural data were collected using a Siemens 3T scanner with a 64 channel head coil at the Yale Center for Biomedi-

cal Imaging. Imaging acquisition parameters were aligned with those of the Human Connectome Project (HCP) (57). High-resolution T1w and T2w structural images were acquired in 224 AC-PC aligned slices, 0.8mm isotropic voxels. T1w images were collected with a magnetization-prepared rapid gradient-echo (MP-RAGE) pulse sequence (TR=2400 ms, TE=2.07 ms, flip angle=8o, field of view=256 x 256 mm). T2w images were collected with a SCP pulse sequence [TR=3200 ms, TE=564 ms, flip angle mode=T2 var, field of view=256 x 256 mm]. Resting-state BOLD images were collected with a multi-band accelerated fast gradient-echo, echo-planar sequence (acceleration factor=6, time repetition (TR)=800 ms, time echo (TE)=31.0 ms, flip angle=55o, field of view=210 x 210 mm, matrix=84 x 84, bandwidth=2290 Hz); 54 interleaved axial slices aligned to the anterior-posterior commissure (AC-PC) with 2.5mm isotropic voxels. Additionally, a pair of reverse phase-encoded spin-echo field maps (anterior-to-posterior and posterior-to-anterior) were acquired (voxel size=2.5 mm isotropic, TR=7220 ms, TE=73 ms, flip angle=90o, field of view=210 x 210 mm, bandwidth=2290 Hz). These neural data were preprocessed using the HCP minimal preprocessing pipeline as described in the section "Neural data acquisition and preprocessing" above, with the same motion scrubbing and nuisance signal regression parameters as were used for the BSNIP dataset.

Neural Parcel ΔGBC Feature Selection and Prediction. For the purposes of patient selection, we were focused on individual differences in neural features as they co-vary in relation to individual symptom scores. However, this individual variation in neural features may be small compared to the overall group mean. Hence for each patient, we compute the difference in each brain location (parcel) between the patient's actual GBC and the group mean GBC for that location. This is denoted by ΔGBC. Importantly, using a de-meaned GBC metric "standardizes" the data and helps to correct for possible differences in scanners/protocols across different datasets. Next, we developed an optimized univariate regression framework leveraging a "dot product" metric to relate a vector of neural features with a single symptom scalar value. This process for neural feature selection (results in **Fig. S27**) is shown as a systems flow diagram in **Fig. S25**. The observed dot product GBC metric ($dpGBC^{obs}$) is computed as follows:

$$dpGBC_i^{obs} = \begin{pmatrix} \Delta GBC_1^{obs} \\ \vdots \\ \Delta GBC_{P_{select}}^{obs} \end{pmatrix}_i \bullet \begin{pmatrix} \beta GBC_1^{obs} \\ \vdots \\ \beta GBC_{P_{select}}^{obs} \end{pmatrix}_{N-i} \tag{3}$$

- where for a given subject i $dpGBC_i^{obs}$ denotes the dot product GBC value of the two vectors $\Delta GBC_i^{obs} \cdot \beta GBC_{N-1}^{obs}$ across all P_{select} parcels,

- ΔGBC_i^{obs} is a vector of length P_{select} denoting a difference map of subject i's GBC^{obs} map relative to the group mean GBC^{obs} map, within a given number of parcels P_{select},

- the βGBC_{N-i}^{obs} vector denotes the PCA-to-GBC statistical group-level β map for a low-dimensional PC symptom score across selected parcels P_{select} for N subjects excluding subject i.

This calculation is then repeated for each subject i, resulting in a final vector $dpGBC^{obs} = [dpGBC_1^{obs}, \ldots, dpGBC_N^{obs}]$ for N subjects. There are several key properties of the $dpGBC^{obs}$ statistic: i) it is not inflated by individual GBC map similarity to the group map because each subject's ΔGBC^{obs} map is demeaned relative to the reference group computed independently of the left-out subject; ii) this statistic is not biased by the parcel number (which drops with iterative selection) because the resulting $dpGBC^{obs}$ value variation is quantified relative to the low-dimensional symptom score across all subjects (see Eq. 2). Put differently, the final evaluation considers the relationship between $dpGBC^{obs}$ and the low-dimensional PC symptom scores across individuals; iii) The dot product statistic can yield both positive and negative values – a property which some map similarity measures lack (e.g. η^2); iv) It is unbounded (unlike a correlation), which is key to maximize co-variation with low-dimensional symptom scores across individuals (see Eq. 2); v) The ΔGBC_i^{obs} map for a given individual is projected onto the basis set of the βGBC_{N-i}^{obs} map, which is independent of the left-out individual but directly related to the low-dimensional PC symptom score variance, thus maximizing the dot product optimization.

Next, we select $N-i$ individuals and compute a univariate regression where participants' low dimensional symptom scores are regressed onto the $dpGBC_{N-i}^{obs}$ values:

$$\begin{pmatrix} dpGBC_1^{obs} \\ \vdots \\ dpGBC_{N-i}^{obs} \end{pmatrix} = \alpha \begin{pmatrix} S_1^{obs} \\ \vdots \\ S_{N-i}^{obs} \end{pmatrix} + \epsilon \tag{4}$$

- where for $dpGBC^{obs}$ each element denotes the dot product value of the $\Delta GBC_i^{obs} \cdot \beta GBC_{N-1}^{obs}$ vectors per subject across P parcels,

- α denotes the regression coefficient in the univariate linear model,

- where for S^{obs} each element denotes the observed low-dimensional symptom score (e.g. *PC3* score) per subject,

- ϵ denotes the error term in the univariate linear model.

After the $dpGBC^{obs} = \alpha S^{obs} + \epsilon$ regression is computed on $N-i$ subjects it is applied to the left-out-subject i. This is repeated for all N subjects and the model is evaluated for the number of parcels that maximize two key dot product evaluation metrics (**Fig. 6B**):

Metric A

For $p = \{1, \ldots, P\}$ find p where

$$\Rightarrow A = \max_r \left[\begin{pmatrix} dpGBC_1 \\ \vdots \\ dpGBC_N \end{pmatrix}^{pred}, \begin{pmatrix} S_1 \\ \vdots \\ S_N \end{pmatrix}^{pred} \right] \quad (5)$$

Metric B

For $p = \{1, \ldots, P\}$ find p where

$$\Rightarrow B = \max_r \left[\begin{pmatrix} dpGBC_1 \\ \vdots \\ dpGBC_N \end{pmatrix}^{obs}, \begin{pmatrix} dpGBC_1 \\ \vdots \\ dpGBC_N \end{pmatrix}^{pred} \right] \quad (6)$$

- where A denotes the maximum r correlation value for the two vectors of $dpGBC^{pred}$ values and the predicted S^{pred} low-dimensional symptom scores (e.g. $PC3$ axis) for N subjects from the leave-one-out cross-validation,

- where B denotes the maximum r correlation value for the two vectors of obs and $pred$ $dpGBC$ values for N subjects.

In the initial step in the step-down model all $P = 718$ parcels are retained in the initial dot product calculation. For each iteration of P selected parcels, the least predictable parcel P (i.e. the parcel with the weakest value in the $PC3$ map) is eliminated from the map. Then, the step-down regression is repeated until $P = 1$.

Pharmacological Neuroimaging Acquisition in Healthy Volunteers - LSD. Methods for the lysergic acid diethylamide (LSD) neuroimaging study are described in detail in prior publications (30). The use of LSD in humans was authorized by the Swiss Federal Office of Public Health, Bern, Switzerland. The study protocol was approved by the Cantonal Ethics Committee of Zurich ($KEK - ZH_{No} : 2014_0496$). Participants received both written and oral descriptions of the study procedures and the effects and possible risks of LSD. All participants provided written informed consent in accordance with the Declaration of Helsinki. The study employed a fully double-blind, randomized, within-subject cross-over design with 3 conditions: (1) placebo + placebo (Pla) condition: placebo (179 mg Mannitol and Aerosil 1 mg po) after pretreatment with placebo (179 mg Mannitol and Aerosil 1 mg po); (2) Pla+LSD (LSD) condition: LSD (100 μg po) after pretreatment with placebo (179 mg Mannitol and Aerosil 1 mg po), or (3) Ketanserin+LSD (Ket+LSD) condition: LSD (100 μg po) after pretreatment with the 5-HT2A antagonist Ket (40 mg po). Data were collected for all subjects in a randomized counterbalanced order at three different sessions each two weeks apart. For all conditions, the first substance was administered 60 minutes before the second substance, and the first neural scan was conducted 75 minutes after the second administration, with a second scan conducted at 300 minutes post-administration. In the present study, only data

from the two neural scans for the LSD and Pla conditions were evaluated.

Briefly, neuroimaging data acquisition details for the LSD study are as follows. MRI data were acquired on a Philips Achieva 3.0T whole-body scanner (Best, The Netherlands). A 32-channel receiver head coil and MultiTransmit parallel radio frequency transmission was used. Images were acquired using a whole-brain gradient-echo planar imaging (EPI) sequence (repetition time=2,500 ms; echo time=27 ms; slice thickness=3 mm; 45 axial slices; no slice gap; field of view=240 mm^2; in-plane resolution=3 mm $\times$ 3 mm; sensitivity- encoding reduction factor=2.0). 240 volumes were acquired per resting state scan resulting in a scan duration of 10 mins. Additionally, two high-resolution anatomical images were acquired using T1-weighted and T2-weighted sequences. T1-weighted images were collected via a 3D magnetization-prepared rapid gradient-echo sequence (MP-RAGE) with the following parameters: voxel size=0.7 mm^3, time between two inversion pulses=3123 ms, inversion time=1055 ms, inter-echo delay=12 ms, flip angle=8°, matrix=320×335, field of view=224×235 mm, 236 sagittal slices. Furthermore T2-weighted images were collected using via a turbo spin-echo sequence with the following parameters: voxel size=0.7 mm^3, repetition time=2500 ms, echo time=415 ms, flip angle=90°, matrix=320 $\times$ 335, field of view=224 mm $\times$ 235 mm, 236 sagittal slices.

Pharmacological Neuroimaging Acquisition in Healthy Volunteers - Ketamine. Similar to the LSD study, the ketamine pharmacological neuroimaging protocol employed a within-subject design where all healthy volunteer participants underwent a single scanning session consisting of two infusions: i) placebo (saline solution) followed by ii) ketamine infusion. The study was approved by the Yale University institutional review board. Participants received both written and oral descriptions of the study procedures and the effects and possible risks of ketamine, and all participants provided written informed consent before beginning the study. Healthy volunteers were informed prior to scanning that they would undergo one placebo run and one ketamine run but were blinded to the order of administration. Because of the sustained effects of ketamine, this infusion was always the second of the two runs, consistent with prior work (42). A doctor and two nurses as well as a study coordinator remained present for the duration of the scan. A bolus of ketamine (0.3 mg/kg of bodyweight) or saline were delivered via infusion 5 sec after the start of the run and then continuously at a rate of 0.65 mg/kg per hour through the duration of the session. The sequence of scans in each for either the placebo or ketamine infusion was as follows: i) resting state (4.67 min); ii) blood draw (sham if saline condition); iii) a cognitive working memory task (total 14 min); iv) blood draw (sham if saline condition); v) a cognitive working memory task (total 14 min); vii) blood draw (sham if saline condition); vii) a cognitive working memory task(total 8.63 min). Data from the the cognitive working memory task were not used in the present study and are actively undergoing a distinct analysis. Participants were scanned at Yale University

on a Siemens Trio 3T whole-body scanner and 32-channel receiver head coil. High-resolution structural T1-weighted images were acquired using an MP-RAGE sequence and the following parameters: voxel size=0.8 mm^3, time between two inversion pulses=3123 ms, inversion time=1055 ms, inter-echo delay=12 ms, flip angle=8°, matrix=320×335, field of view=227×272 mm, 227 sagittal slices. T2-weighted images were acquired with the following parameters:voxel size=0.8 mm^3, repetition time=2500 ms, echo time=415 ms, flip angle=90°, matrix=320 × 335, field of view=227×272 mm, 227 sagittal slices. BOLD images were acquired using a whole-brain gradient-echo planar imaging (EPI) sequence (400 frames at TR=0.7 ms; TE=30 ms; slice thickness=2.5 mm; 54 axial slices; no slice gap; field of view=250 mm^2; in-plane resolution=2.5 mm × 2.5 mm). In addition, a field map and pair of spin-echo gradients were collected at the end of every scanning session.

Pharmacological Neuroimaging Processing and Analysis for LSD and Ketamine Samples. All data were preprocessed to be consistent with the BSNIP and replication dataset processing steps. Specifically, we used the HCP minimal preprocessing pipeline and with the same motion scrubbing and nuisance signal regression parameters as were used for the BSNIP dataset as described in the section "Neural data acquisition and preprocessing" above. Parcel-wise GBC maps were computed as described in the section "Resting-state functional connectivity and global brain connectivity" above, by first parcellating the dense time-series for each subject and then computing the parcel-level GBC. As with the BSNIP and replication analyses, parcel GBC was first calculated by parcellating the dense time-series for every subject and then computing GBC across all 718 parcels from the whole-brain functional parcellation (51, 52). Group-level GBC maps were computed for every participant for all conditions. We computed the contrast map for "LSD-Placebo" as well as "Ketamine-Placebo" conditions as a t-test between both the pharmacological scans versus placebo scans. The Z-scored t-contrast maps between pharmacological and placebo conditions were used as a pharmacological target map in relation to the PC-derived neural GBC maps **Fig. 7** and **Fig. S30**.

Neural Gene Expression Mapping. The gene mapping analyses in this study utilize the procedure described in (33). Briefly, we used cortical gene expression data from the publicly available Allen Human Brain Atlas (AHBA, RRID:SCR_007416) , mapped to cortex (33). Specifically, the AHBA quantified expression levels across 20,737 genes obtained from six postmortem human brains using DNA microarray probes sampled from hundreds of neuroanatomical loci. Recent studies demonstrated the ability to map expression of each gene onto neuroimaging-compatible templates (32, 33). Building on these innovations we mapped gene expression on to 180 symmetrized cortical parcels from the HCP atlas (52) in line with recently published methods (33). This yielded a group-level map for each gene where the value in each parcel reflected the average expression level of that gene in the AHBA dataset. These group-level maps were in turn as a gene expression target map in relation to the PC-derived neural GBC maps.

Data Availability. All primary results derive from data that is publicly available from sources described above. All symptom and neuroimaging data for the BSNIP-1 consortium was obtained via the NDA (https://nda.nih.gov/edit_collection.html?id=2274).

Code Availability. Code and core results presented in this paper are available at https://bitbucket.org/jielisa_ji/bbs_manuscript/src/master/.

Results

The key questions and results of this study are summarized in **Fig. S1** and an overview of the workflow is presented in **Fig. S2**. Additionally, for convenient reference, a glossary of the key terms and abbreviations used throughout the study are provided in **Table S1**.

Dimensionality-Reduced PSD Symptom Variation is Stable and reproducible. First, to evaluate PSD symptom variation we examined core PSD psychopathology metrics captured by two instruments: the Brief Assessment of Cognition in Schizophrenia (BACS) and the Positive and Negative Syndrome Scale (PANSS). We refer to these 36 items as "symptom measures" throughout the rest of the paper. We observed group mean differences across DSM diagnoses (**Fig. 1A**, p<0.05 Bonferroni corrected); however, symptom measure distributions revealed notable overlap that crossed diagnostic boundaries (11, 58). Furthermore, we observed marked collinearity between symptom measures across the PSD sample (**Fig. 1B**), indicating that a dimensionality-reduced solution may sufficiently capture meaningful varation in this symptom space. We hypothesized that such a dimensionality-reduced symptom solution may improve PSD symptom-neural mapping as compared to pre-existing symptom scales.

Here we report results from a principal component analysis (PCA) as it produces a deterministic solution with orthogonal axes (i.e. no *a priori* number of factors needs to be specified) and explains all variance in symptom measures. Results were highly consistent with prior symptom-reduction studies in PSD: we identified 5 PCs (**Fig. 1C**), which captured ∼50.93% of all variance (see **Methods** & **Fig. S3**) (23).

Critically, the key innovation here is the combined analysis across PSD diagnoses of canonical PSD symptoms and cognitive deficits, a fundamental PSD feature (17). The 5 PCs revealed few distinct boundaries between DSM categories (**Fig. 1D**). Furthermore, we observed substantial overlap in PC scores between controls and patients in all DSM diagnostic groups, with PC1 showing the greatest differentiation between PSD and the "normal-spectrum". This overlap is not unexpected, given that PSD symptoms are present to some extent in "healthy" individuals not meeting DSM diagnostic criteria (59–62).

Fig. 1E highlights loading configurations of symptom measures forming each PC. To aid interpretation, we assigned a

Fig. 1. Quantifying data-driven low-dimensional variation of cross-diagnostic psychosis spectrum disorder symptoms and cognitive deficits. (A) Distributions of symptom scores for each of the DSM diagnostic groups across core psychosis symptom meaures (PANSS positive, negative, and general symptoms tracking illness severity) and cognitive deficits (BACS composite cognitive performance). BPP: bipolar disorder with psychosis (yellow, N=150); SADP: schizo-affective disorder (orange, N=119); SZP - schizophrenia (red, N=167); All PSD patients (black, N=436); Controls (white, N=202). Bar plots show group means; error bars show standard deviations. **(B)** Correlations between 36 symptom measures across all PSD patients (N=436). **(C)** Screeplot shows the % variance explained by each of the principal components (PCs) from a PCA performed using all 36 symptom measures across 436 PSD patients. The size of each point is proportional to the variance explained. The first five PCs (green) survived permutation testing (p<0.05, 5,000 permutations). Together they capture 50.93% of all symptom variance (inset). **(D)** Distribution plots showing subject scores for the 5 significant PCs for each of the clinical groups, normalized relative to the control group. Note that control subjects (CON) were not used to derive the PCA solution; however, all subjects, including CON, can be projected into the data-reduced symptom geometry. **(E)** Loading profiles shown in dark grey for the 36 PANSS/BACS symptom measures on the 5 significant PCs. Each PC ("Global Functioning", "Cognitive Functioning", "Psychosis Configuration", "Affective Valence", "Agitation/Excitement") was named based on the pattern of loadings on symptom measures. See **Fig. S3G** for numerical loading values. The PSD group mean score on each symptom measure is also shown, in light grey (scaled to fit on the same radarplots). Note that the group mean configuration resembles the *PC1* loading profile most closely (as *PC1* explains the most variance in the symptom measures). **(F)** PCA solution shown in the coordinate space defined by the first three PCs. Colored arrows show *a priori* composite PANSS/BACS vectors projected into the *PC1-3* coordinate space. The *a priori* composite symptom vectors do not directly align with data-driven PC axes, highlighting that PSD symptom variation is not captured fully by any one aggregate *a priori* symptom score. Spheres denote centroids (i.e. center of mass) for each of the patient diagnostic groups and control subjects. Alternative views showing individual patients and controls projected into the PCA solution are in **Fig. S3A-F**.

Fig. 2. Dimensionality reduction of PSD symptom measures is highly stable and reproducible. (A) PCA solutions for leave-one-site out cross-validation (left) and 5-fold bootstrapping (right) explain a consistent total proportion of variance with a consistent number of significant PCs after permutation testing. Full results available in **Fig. S5** and **Fig. S6**. **(B)** Predicted versus observed single subject PC scores for all five PCs are shown for an example site (here Site 1). **(C)** Mean correlations between predicted and observed PC scores across all patients calculated via k-fold bootstrapping for k=2-10. For each k, patients were randomly split into k folds. For each fold, a subset of patients was held out and PCA was performed on the remaining patients. Predicted PC scores for the held-out patients were computed from the PCA obtained in all other retained patients. Original observed PC scores for the held-out patients were then correlated with the predicted PC scores. **(D)** Mean correlations between predicted and observed symptom measure loadings are shown for leave-one-site-out cross-validation (left), across 1,000 runs of 5-fold cross-validation (middle), and 1,000 split-half replications (right). For split-half replication, loadings were compared between the PCA performed in each independent half sample. Note: for panels **C-D** correlation values were averaged all k runs, all 6 leave-site-out-runs, or all 1,000 runs of the 5-fold cross-validation and split-half replications. Error bars indicate the standard error of the mean.

name for each PC based on its most strongly weighted symptom measures. This naming is qualitative but informed by the pattern of loadings of the original 36 symptom measures (**Fig. 1**). For example, *PC1* was highly consistent with a general impairment dimension (i.e. "Global Functioning"); *PC2* reflected more exclusively variation in cognition (i.e. "Cognitive Functioning"); *PC3* indexed a complex configuration of psychosis-spectrum relevant items (i.e. "Psychosis Configuration"); *PC4* generally captured variation mood and anxiety related items (i.e. "Affective Valence"); finally, *PC5* reflected variation in arousal and level of excitement (i.e. "Agitation/Excitation"). For instance, a generally impaired patient would have a highly negative *PC1* score, which would reflect low performance on cognition and elevated scores on most other symptomatic items. Conversely, an individual with a high positive *PC3* score would exhibit delusional, grandiose, and/or hallucinatory behavior, whereas a person with a negative *PC3* score would exhibit motor retardation, social avoidance, possibly a withdrawn affective state with blunted affect (63). Comprehensive loadings for all 5 PCs are shown in **Fig. S3G**. **Fig. 1F** highlights the mean of each of the 3 diagnostic groups (colored spheres) and healthy controls (black sphere) projected into a 3-dimensional orthogonal coordinate system for *PCs 1,2 & 3* (x,y,z axes respectively; alternative views of the 3-dimensional coordinate system with all patients projected are shown in **Fig. S3**). Critically, PC axes were not parallel with traditional aggregate symptom scales. For instance, *PC3* is angled at ∼45° to the dominant direction of PANSS Positive and Negative symptom variation (purple and blue arrows respectively in **Fig. 1F**).

Because *PC3* loads most strongly on to hallmark symptoms of PSD (including strong positive loadings across PANSS Positive symptom measures in the PANSS and strong negative loadings onto most Negative measures), we focus on this PC as an opportunity to quantify an innovative, fully data-driven dimension of symptom variation that is highly characteristic of the PSD patient population. Additionally, this bi-directional symptom axis captured shared variance from measures in other traditional symptoms factors, such the PANSS General factor and cognition. We found that the *PC3* result provided a powerful empirical demonstration of how using a data-driven dimensionality-reduced solution (via PCA) can reveal novel patterns intrinsic to the structure of PSD psychopathology.

Notably, independent component analysis (ICA), an alternative dimensionality reduction procedure which does not enforce component orthogonality, produced similar effects for this PSD sample, see **Supplementary Note 1** & **Fig. S4A**). Certain pairs of components between the PCA and ICA solutions appear to be highly similar and exclusively mapped

(IC5 and PC4; IC4 and PC5) (**Fig. S4B**). On the other hand, PCs 1-3 and ICs 1-3 do not exhibit a one-to-one mapping. For example, *PC3* appears to correlate positively with *IC2* and equally strongly negatively with *IC3*, suggesting that these two ICs are oblique to the PC and perhaps reflect symptom variation that is explained by a single PC. The orthogonality of the PCA solution forces the resulting components to capture maximally separated, unique symptom variance, which in turn map robustly on to unique neural circuits. We observed that the data may be distributed in such a way that in the ICA highly correlated independent components emerge, which do not maximally separate the symptom variance associate with neural variance. We demonstrate this by plotting the relationship between parcel beta coefficients for the $\beta_{PC3}GBC$ map versus the $\beta_{IC2}GBC$ and $\beta_{IC3}GBC$ maps (**Fig. S13G**). The sigmoidal shape of the distribution indicates an improvement in the Z-statistics for the $\beta_{PC3}GBC$ map relative to the $\beta_{IC2}GBC$ and $\beta_{IC3}GBC$ maps.

Next, we show that the PCA solution was highly ***stable*** when tested across sites, and k-fold cross-validations, and was ***reproducible*** in independent split-half samples. First, we show that the symptom-derived PCA solution remained highly robust across all sites (**Fig. 2A**) and 5-fold cross-validation iterations (see **Methods**). Notably, the total proportion of variance explained as well as the total number of significant PCs remained stable (**Fig. 2A**). Second, PCA loadings accurately and reliably computed the scores of subjects in a hold-out sample (**Fig. 2B**). Specifically, for each left-out site, we tested if single-subject predicted scores were similar to the originally observed scores of these same subjects in the full PCA (i.e. performed with all 436 subjects). This similarity was high for all sites (**Fig. S6**). Third, we verified that the predicted-versus-observed PCA score similarities also hold via a *k*-fold cross-validation (for *k*=2 to 10, **Fig. 2C** & **S5**). Finally, across all 5 PCs, predicted-to-observed similarity of PCA loadings was remarkably high. Moreover, PCA loadings were stable when testing via leave-site-out analysis and 5-fold cross-validation. We furthermore demonstrated the reproducibility of the solution using 1,000 independent split-half replications (**Fig. 2D**).For each run of split-half replication, PSD subjects were split into 2 sets and a PCA was performed independently in both sets. The loadings from both PCAs were then compared using a Pearson's correlation. Importantly, results were not driven by medication status or dosage (**Fig. S7**). Collectively, these data-reduction analyses strongly support a stable and reproducible low-rank PSD symptom geometry.

Dimensionality-Reduced PSD Symptom Geometry Reveals Novel and Robust Neuro-Behavioral Relationships. Next, we tested if the dimensionality-reduced symptom geometry can identify robust and novel patterns of neural variation across PSD. We chose to use global brain connectivity (GBC), a summary FC metric, to measure neural variance because it yields a parsimonious measure reflecting how globally coupled an area is to the rest of the brain (64) (see **Methods**). Furthermore, we selected GBC because: i) the metric is agnostic regarding the location of dysconnectivity

as it weights each area equally; ii) it yields an interpretable dimensionality-reduction of the full FC matrix; iii) unlike the full FC matrix or other abstracted measures, GBC produces a neural map, which can be related to other independent neural maps (e.g. gene expression or pharmacology maps, discussed below). Furthermore, GBC has been shown to be sensitive to altered patterns of global neural connectivity in PSD cohorts (65–68) as well as in models of psychosis (30, 69).

All 5 PCs captured unique patterns of GBC variation across the PSD (**Fig. S10**), which were not observed in CON (**Fig. S11**). Again we highlight the hallmark *"Psychosis Configuration"* dimension (i.e. *PC3*), here to illustrate the benefit of the low-rank PSD symptom geometry for symptom-neural mapping relative to traditional aggregate PANSS symptom scales. The relationship between total PANSS Positive scores and GBC across N=436 PSD patients ($\beta_{Pos}GBC$, **Fig. 3A**) was statistically modest (**Fig. 3B**) and no areas survived whole-brain type-I error protection (**Fig. 3C**, *p*<0.05). In contrast, regressing *PC3* scores onto GBC across N=436 patients revealed a robust symptom-neural $\beta_{PC3}GBC$ map (**Fig. 3E-F**), which survived whole-brain type-I error protection. Of note, the *PC3 "Psychosis Configuration"* axis is bidirectional whereby individuals who score either highly positively or negatively are symptomatic. Therefore, a high positive *PC3* score was associated with both reduced GBC across insular and superior dorsal cingulate cortices, thalamus, and anterior cerebellum and elevated GBC across precuneus, medial prefrontal, inferior parietal, superior temporal cortices and posterior lateral cerebellum – consistent with the default-mode network (70). A high negative *PC3* score would exhibit the opposite pattern. Critically, this robust symptom-neural mapping emerged despite no differences between mean diagnostic group *PC3* scores (**Fig. 3D**). These two diverging directions may be captured separately in the ICA solution, when orthogonality is not enforced (**Fig. S13**). Moreover, the *PC3* symptom-neural map exhibited improved statistical properties relative to other GBC maps computed from traditional aggregate PANSS symptom scales (**Fig. 3G-J**). Notably, the *PC2 – Cognitive Functioning* dimension, which captured a substantial proportion of cognitive performance-related symptom variance independent of other symptom axes, revealed a circuit that was moderately (anti-)correlated with other PC maps but strongly anti-correlated with the BACS composite cognitive deficit map (*r*=-0.81, **Fig. S9O**). This implies that the *PC2* map reflects unique neural circuit variance that is relevant for cognition, independent of the other *PC* symptom dimensions.

Univariate Neuro-Behavioral Map of Psychosis Configuration is Reproducible. After observing improved symptom-neural *PC3* statistics, we tested if these neural maps replicate. Recent attempts to derive stable symptom-neural mapping using multivariate techniques, while inferentially valid, have not replicated (26). This fundamentally relates to the tradeoff between the sample size needed to resolve multivariate neuro-behavioral solutions and the size of the feature space. To mitigate the feature size issue we re-computed the $\beta_{PC}GBC$ maps using a functionally-

Fig. 3. Dimensionality-reduced symptom variation reveals robust neuro-behavioral mapping. (A) Distributions of total PANSS Positive symptoms for each of the clinical diagnostic groups normalized relative to the control group (white=CON; black=all PSD patients; yellow=BPP; orange=SADP; red=SZP). **(B)** $\beta_{Positive}GBC$ map showing the relationship between the aggregate PANSS Positive symptom score for each patient regressed onto global brain connectivity (GBC) across all patients (N=436). **(C)** No regions survived non-parametric family-wise error (FWE) correction at p<0.05 using permutation testing with threshold-free cluster enhancement (TFCE). **(D)** Distributions of scores for *PC3 "Psychosis Configuration"* across clinical groups, again normalized to the control group. **(E)** $\beta_{PC3}GBC$ map showing the relationship between the *PC3 "Psychosis Configuration"* score for each patient regressed onto GBC across all patients (N=436). **(F)** Regions surviving p<0.05 FWE whole-brain correction via TFCE showed clear and robust effects. **(G)** Comparison between the *Psychosis Configuration* symptom score versus the aggregate PANSS Positive symptom score GBC map for every datapoint in the neural map (i.e. greyordinate in the CIFTI map). The sigmoidal pattern indicates an improvement in the Z-statistics for the *Psychosis Configuration* symptom score map (panel **E**) relative to the aggregate PANSS Positive symptom map (panel **B**). **(H)** A similar effect was observed when comparing the *Psychosis Configuration* GBC map relative to the PANSS Negative symptoms GBC map (**Fig. S9**). **(I)** Comparison of the variances for the *Psychosis Configuration*, PANSS Negative and PANSS Positive symptom map Z-scores. **(J)** Comparison of the ranges between the *Psychosis Configuration*, Negative and Positive symptom map Z-scores. Symptom-neural maps for all 5 PCs and all 4 traditional symptom scales (BACS and PANSS subscales) are shown in **Fig. S9**.

derived whole-brain parcellation via the recently-validated CAB-NP atlas (51, 52) (**Methods**). Here, a functional parcellation is a principled way of neural feature reduction (to 718 parcels) that can also appreciably boost signal-to-noise (51, 52). Indeed, parcelling the full-resolution "dense" resting-state signal for each subject prior to computing GBC statistically improved the group-level symptom-neural maps compared to parcelling after computing GBC (**Fig. 4A–C**, all maps in **Fig. S10**). Results demonstrate that the univariate symptom-neural mapping was indeed stable across 5-fold bootstrapping, leave-site-out, and split-half cross-validations (**Fig. 4D–N**, see **Supplementary Note 2**), yielding consistent symptom-neural *PC3* maps. Importantly, the symptom-neural maps computed using ICA showed comparable results

(**Fig. S13**).

A Multivariate PSD Neuro-Behavioral Solution Can be Computed but is Not Reproducible with the Present Sample Size. Several recent studies have reported "latent" neuro-behavioral relationships using multivariate statistics (15, 71, 72), which would be preferable because they simultaneously solve for maximal covariation across neural and behavioral features. Given the possibility of deriving a stable multivariate effect, here we tested if results improve with canonical correlation analysis (CCA) (73) (**Fig. 5A**).

To evaluate if the number of neural features affects the solution, we computed CCA using GBC from: i) 180 symmetrized cortical parcels; ii) 359 bilateral subcortex-only

Fig. 4. Parcellated symptom-neural GBC maps reflecting psychosis configuration are statistically robust and reproducible. (A) Z-scored *PC3 Psychosis Configuration* GBC neural map at the 'dense' (full CIFTI resolution) level. **(B, C)** Neural data parcellated using a whole-brain functional partition (51) *before* computing subject-level GBC yielded stronger statistical values in the Z-scored *Psychosis Configuration* GBC neural map as compared to when parcellation was performed *after* computing GBC for each subject. **(D)** Summary of similarities between all symptom-neural βGBC maps (PCs and traditional symptom scales) across 5-fold cross-validation. Boxplots show the range of *r* values between βGBC maps for each fold and the full model. **(E)** Normalized βGBC map from regression of individual patients' *PC3* scores onto parcellated GBC data, shown here for a subset of patients from Fold 1 out of 5 (N=349). The greater the magnitude of the coefficient for a parcel, the stronger the statistical relationship between GBC of that parcel and *PC3* score. **(F)** Correlation between the βGBC value of each parcel in the regression model computed using patients in Fold 1 and the full PSD sample (N=436) model indicates that the leave-one-fold-out βGBC map was highly similar to the βGBC map obtained from the full PSD sample model (*r*=0.924). **(G)** Summary of leave-one-site-out regression for all symptom-neural maps. Regression of PC symptom scores onto parcellated GBC data, each time leaving out subjects from one site, resulted in highly similar maps. This highlights that the relationship between *PC3* scores and GBC is robust and not driven by a specific site. **(H)** βGBC map for all PSD except one site. As an example, Site 3 is excluded here given that it recruited the most patients (and therefore may have the greatest statistical impact on the full model). **(I)** Correlation between the value of each parcel in the regression model computed using all patients minus Site 3, and the full PSD sample model. **(J)** Split-half replication of $\beta_{PC3}GBC$ map. Bar plots show the mean correlation across 1,000 runs; error bars show standard error. Note that the split-half effect for *PC1* was exceptionally robust. The split-half consistency for *PC3*, while lower, was still highly robust and well above chance. **(K)** $\beta_{PC3}GBC$ map from *PC3*-to-GBC regression for the first half (H1) patients, shown here for one exemplar run out of 1000 split-half validations. **(L)** Correlation across 718 parcels between the H1 predicted coefficient map (i.e. panel **K**) and the observed coefficient map for H1. **(M-N)** The same analysis as **K-L** is shown for patients in H2, indicating a striking consistency in the *Psychosis Configuration* $\beta_{PC3}GBC$ map correspondence.

Fig. 5. Multivariate symptom-neural feature mapping using canonical correlation analysis (CCA). (A) Schematic of CCA data (**B**, **N**), transformation (Ψ, Θ), and transformed latent (U, V) matrices. Here each column in **U** and **V** is referred to as a canonical variate (CV); each corresponding pair of CVs (e.g. U_1 and V_1) is referred to as a canonical mode. **(B)** CCA maximized correlations between the CVs (**U** and **V**) **(C)** Screeplot showing canonical modes obtained from 180 neural features (cortical GBC symmetrized across hemispheres) and 36 symptom measures ('180 vs. 36 CCA'). Inset illustrates the correlation (r=0.85) between the CV of the first mode, U_1 and V_1 (note that the correlation was not driven by a separation between diagnoses). Modes 9 and 12 remained significant after FDR correction. **(D)** CCA computed with 180 neural features and 5 PC symptom features ('180 vs. 5 CCA'). Here all modes remained significant after FDR correction. Dashed black line shows the null calculated via a permutation test with 5,000 shuffles; grey bars show 95% confidence interval. **(E)** Correlation between **B** and **NΘ** reflects how much of the symptom variation can be explained by the latent neural features. **(F)** Proportion of symptom variance explained by each of the neural CVs in the 180 vs. 36 CCA. Inset shows the total proportion of behavioral variance explained by the neural CVs. **(G)** Proportion of total symptom variance explained by each of the neural CVs in the 180 vs. 5 CCA. While CCA using symptom PCs has fewer dimensions and thus accounts for lower total variance (see inset), each neural variate explains a higher amount of symptom variance than in **F**, suggesting that CCA could be optimized by first obtaining a low-rank symptom solution. Dashed black line indicates the null calculated via a permutation test with 5,000 shuffles; grey bars show 95% confidence interval. Neural variance explained by symptom CVs are plotted in **Fig. S16**. **(H)** Distributions of *CV3* scores from the 180 vs. 5 CCA are shown here as an example of characterizing CV configurations. Scores for all diagnostic groups are normalized to CON. Additionally, **(I)** symptom canonical factor loadings, **(J)** loadings of the original 36 symptom measures, and **(K)** neural canonical factor loadings for *CV3* are shown. **(L)** Within-sample CCA cross-validation appeared robust (see **Fig. S18**). However, a split-half replication of the 180 vs. 5 CCA (using two independent non-overlapping samples) was not reliable. Bar plots show the mean correlation for each CV between the first half (H1) and the second half (H2) CCA, each computed across 1,000 runs. *Left:* split-half replication of the symptom PC loadings matrix Ψ; *Middle:* individual symptom measure loadings; *Right:* the neural loadings matrix Θ, which in particular was not stable. Error bars show the standard error of the mean. Scatterplot shows the correlation between *CV3* neural loadings for H1 vs. H2 for one example CCA run, illustrating lack of reliability. **(M)** Leave-one-subject-out cross-validation further highlights CCA instability.

parcels; iii) 192 symmetrized subcortical parcels; iv) 12 functional networks, all from the brain-wide cortico-subcortical CAB-NP parcellation (51, 52). We did not compute a solution using 718 bilateral whole-brain parcels, as this exceeded the number of samples and rendered the CCA insolvable. Notably, the 359 subcortex-only solution did not produce stable results according to any criterion, whereas the 192 symmetrized subcortical features (**Fig. S14**) and 12 network-level features (**Fig. S15**) solutions captured statistically modest effects relative to the 180 symmetrized cortical features (**Fig. 5B-D**). Therefore, we characterized the 180-parcel solution further. We examined two CCA solutions using these 180 neural features in relation to symptom scores: i) all 36 item-level symptom measure scores from the PANSS/BACS ('180 vs. 36 CCA', **Fig. 5C&F**); and ii) 5 PC symptom scores ('180 vs. 5 CCA', **Fig. 5D&G**, see **Methods**). Only 2 out of 36 modes for the 180 vs. 36 CCA solution survived permutation testing and false discovery rate (FDR) correction (**Fig. 5C**). In contrast, all 5 modes of the 180 vs. 5 CCA survived (**Fig. 5D**). Critically, we found that no single CCA mode computed on item-level symptom measures captured more variance than the CCA modes derived from PC symptom scores, suggesting that the PCA-derived dimensions capture more neurally-relevant variation than any one single clinical item (**Fig. 5E-G**). Additional CCA details are presented in **Supplementary Note 3** and **Fig. S16**.

We highlight here an example canonical variate, *CV3*, across both symptom and neural effects, with data for all *CVs* shown in **Fig. S17**. *CV3* scores across diagnostic groups normalized to controls are shown **Fig. 5H** in addition to how *CV3* loads onto each *PC* (**Fig. 5I**). The negative loadings on to *PCs 1, 4, and 5* and the high positive loadings on to *PC3* in **Fig. 5I** indicate that *CV3* captures some shared variation across symptom *PCs*. This can also be visualized by computing how *CV3* projects onto the original 36 symptom measures (**Fig. 5J**). Finally, the neural loadings for *CV3* are shown in **Fig. 5K**.

Lastly, we tested if the 180 vs. 5 CCA solution is stable and reproducible, as done with PC-to-GBC univariate results. The CCA solution was robust when tested with *k*-fold and leave-site-out cross-validation (**Fig. S18**) likely because these methods use CCA loadings derived from the full sample. However, the CCA loadings did not replicate in non-overlapping split-half samples (**Fig. 5L**, see **see Supplementary Note 4**). Moreover, a leave-one-subject-out cross-validation revealed that removing a single subject from the sample affected the CCA solution such that it did not generalize to the left-out subject (**Fig. 5M**). This is in contrast to the PCA-to-GBC univariate mapping, which was substantially more reproducible for all attempted cross-validations relative to the CCA approach. This is likely because substantially more power is needed to resolve a stable multivariate neuro-behavioral effect with this many features (26). Indeed, a multivariate power analysis using 180 neural features and 5 symptom features and assuming a true canonical correlation of $r = 0.3$ suggests that a minimal sample size of $N = 8,145$ is needed to sufficiently detect the effect (27), which is an or-

der of magnitude greater than the available sample size (**Fig. S19**). Therefore, we leverage the univariate symptom-neural result for subsequent subject-specific model optimization and comparisons to molecular neuroimaging maps.

A Major Proportion of Overall Neural Variance May Not be Relevant for Psychosis Symptoms. Most studies look for differences between clinical and control groups, but to our knowledge no study has tested whether both PSD and healthy controls actually share a major portion of neural variance that may be present across all people. If the bulk of the neural variance is similar across both PSD and CON groups then including this clinically-irrelevant neural signal might obscure clinical-relevant neuro-behavioral relationships. To test this, we examined the shared variance structure of the neural signal for all PSD patients (N=436) and all controls (N=202) independently by conducting a PCA on the GBC maps (see **Methods**). Patients' and controls' neural signals were highly similar for each of the first 3 neural PCs (>30% of all neural variance in each group)(**Fig. S20A–J**). These neural *PCs* may reflect a "core" shared symptom-irrelevant neural variance that generalizes across all people. These data suggest that the bulk of neural variance in PSD patients may actually not be symptom-relevant, which highlights the importance of optimizing symptom-neural mapping. Under this assumption, removing the shared neural variance between PSD and CON should not drastically affect the reported symptom-neural univariate mapping solution, because this common variance *does not map to clinical features*. Thus, we conducted a PCA using the parcellated GBC data from all 436 PSD and 202 CON (see **Methods**), which we will refer to as "GBC-PCA" to avoid confusion with the symptom/behavioral PCA. This GBC-PCA resulted in 637 independent GBC-PCs. Since PCs are orthogonal to each other, we then partialled out the variance attributable to GBC-PC1 from the PSD data by reconstructing the PSD GBC matrix using only scores and coefficients from the remaining 636 GBC-PCs ($\hat{GBC}_{woPC1}$). We then reran the univariate regression as described in **Fig. 3**, using the same five symptom PC scores across 436 PSD (**Fig. S21**). Removing the first PC of shared neural variance (which accounted for about 15.8% of the total GBC variance across CON and PSD) from PSD data attenuated the statistics slightly (not unexpected as the variance was by definition reduced) but otherwise did not strongly affect the univariate mapping solution. We repeated the symptom-neural regression next with the first two GBC-PCs partialled out of the PSD data **Fig. S22**, with the first three PCs parsed out **Fig. S23**, and with the first four neural PCs parsed out **Fig. S24**. The symptom-neural maps remain fairly robust, although the similarity with the original $\beta_{PC}GBC$ maps does drop as more common neural variance is parsed out.

Optimizing Symptom-Neural Mapping Features for Personalized Prediction via Dimensionality-reduced Symptoms. Above we demonstrated that symptom *PC* scores can be reliably computed across sites and cross-validation approaches (**Fig. 2**). Next we show that leave-

Fig. 6. Optimizing neural feature selection to inform single-subject prediction via a low-dimensional symptom solution. (A) Leave-one-out cross-validation for the symptom PCA analyses indicates robust individual score prediction. Top panel: Scatterplot shows the correlation between each subject's predicted *PC3* score from a leave-one-out PCA model and their observed *PC3* score from the full-sample PCA model, r=0.99. Bottom panel: Correlation between predicted and observed individual PC scores was above 0.99 for each of the significant PCs (see **Fig. 1**). The red line indicates r=1. **(B)** We developed a univariate step-down feature selection framework to obtain the most predictive parcels using a subject-specific approach via the $dpGBC$ index. Specifically, the 'observed' patient-specific $dpGBC^{obs}$ was calculated using each patient's ΔGBC^{obs} (i.e. the patient-specific GBC map vs. the group mean GBC for each each parcel) and the 'reference' symptom-to-GBC *PC3* map (described in **Fig. 4B**) $[dpGBC^{obs} = \Delta GBC^{obs} \cdot \beta_{PC3} GBC^{obs}]$. See **Methods** & **Fig. S25** for complete feature selection procedure details. In turn, we computed the predicted $dpGBC$ index for each patient by holding their data out of the model and predicting their score ($dpGBC^{pred}$). We used two metrics to evaluate the maximally predictive feature subset: i) The correlation between *PC3* symptom score and $dpGBC^{obs}$ across all N=436, which was maximal for $P = 39$ parcels [r=0.36, purple arrow]; ii) The correlation between $dpGBC^{obs}$ and $dpGBC^{pred}$, which also peaked at $P = 39$ parcels [r=0.35, purple arrow]. **(C)** The $P = 39$ maximally predictive parcels from the $\beta_{PC3} GBC^{obs}$ map are highlighted (referred to as the "selected" map). **(D)** Across all n=436 patients we evaluated if the selected parcels improve the statistical range of similarities between the ΔGBC^{obs} and the $\beta_{PC3} GBC^{obs}$ reference for each patient. For each subject the value on the X-axis reflects a correlation between their ΔGBC^{obs} map and the $\beta_{PC3} GBC^{obs}$ map across all 718 parcels; the Y-axis reflects a correlation between their ΔGBC^{obs} map and the $\beta_{PC3} GBC^{obs}$ map only within the "selected" 39 parcels. The marginal histograms show the distribution of these values across subjects. **(E)** Each DSM diagnostic group showed comparable correlations between predicted and observed $dpGBC$ values. The r-value shown for each group is a correlation between the $dpGBC^{obs}$ and $dpGBC^{pred}$ vectors, each of length N. **(F)** Scatterplot for a single patient with a positive behavioral loading ($PC3$ score$= 1.42$) and also with a high correlation between predicted ΔGBC^{pred} versus observed ΔGBC^{obs} values for the "selected" 39 parcels ($\rho = 0.825$). Right panel highlights the observed vs. predicted ΔGBC map for this patient, indicating that 94.9% of the parcels were predicted in the correct direction (i.e. in the correct quadrant).

one-subject-out cross-validation yields reliable effects for the low-rank symptom PCA solution (**Fig. 6A**). This stable single-subject PC score prediction provides the basis for testing if the derived neural maps can yield an individually-reliable subset of features. To this end, we developed a framework for univariate neural feature selection (**Fig. S25**) based on PC scores (e.g. *PC3* score). Specifically, we computed a dot product GBC metric ($dpGBC$) that provides an index of similarity between an individual ΔGBC topography relative to a "reference" group-level $\beta_{PC} GBC$ map (see **Methods** and **Fig. S25**). Using this $dpGBC$ index via a feature selection step-down regression, we found a subset of parcels for which the symptom-neural statistical association was maximal (**Fig. 6A**). For *PC3* we found $P = 39$ maximally predictive parcels out of the group neural $\beta_{PC} GBC$ map. Specifically, the relationship between *PC3* symptom scores and $dpGBC$ values across subjects was maximal (**Fig. 6B**, top panel, $r = 0.36$) as was the relationship between predicted $dpGBC^{pred}$ vs. observed $dpGBC^{obs}$ (**Fig. 6B**, bottom panel, $r = 0.35$) (see **Fig. S26** for all PCs). Importantly, the "subset" feature map (i.e. $[\beta_{PC3} GBC^{obs}]^{P=39}$, **Fig. 6C**) exhibited improved statistical properties relative to the full map (i.e. $[\beta_{PC3} GBC^{obs}]^{P=718}$, **Fig. 6D**). Furthermore, the relationship between observed vs. predicted subset feature maps (i.e. $r[dpGBC^{obs}, dpGBC^{pred}]$) was highly consistent across DSM diagnoses (**Fig. 6E**). Finally, a sin-

gle patient is highlighted for whom the correlation between predicted and observed subset feature maps was high (i.e. $r[\Delta GBC^{pred}, \Delta GBC^{obs}]$, **Fig. 6F**), demonstrating that the dimensionality-reduced symptom scores can be used to quantitatively optimize symptom-neural map features for individuals.

Single Patient Evaluation via Neuro-Behavioral Target Map Similarity. We showed a quantitative framework for computing a neuro-behavioral model at the single-patient level. This brain-behavior space (BBS) model was optimized along a single dimensionality-reduced symptom PC axis. Next, we tested a hybrid patient selection strategy by first imposing a PC-based symptom threshold, followed by a target neural similarity threshold driven by the most highly predictive symptom-neural map features. This is described in **Methods**, **Supplementary Note 5**, and **Fig. S27-S28**. We found that for patients with a high (either positive or negative) *PC* symptom score the symptom-neural relationship was robust (**Fig. S27A-D**), and these patients could be reliably selected for that particular dimension (**Fig. S27E,G**). Conversely, patients with a low absolute *PC* score showed a weak relationship with symptom-relevant neural features. This is intuitive because these individuals do not vary along the selected PC symptom axis. We also tested this patient selection strategy on an independent cross-diagnostic "replica-

Fig. 7. Leveraging subject-specific brain-behavioral maps for molecular neuroimaging target selection. (A) Data for two individual patients from the replication dataset are highlighted for PC3: X_{PC3} (blue) and Y_{PC3} (yellow). Both of these patients scored above the neural and behavioral thresholds for *PC3* defined in the "discovery" PSD dataset. Patient X_{PC3} loads highly negatively on the *PC5* axis and Patient Y_{PC3} loads highly positively. Density plots show the projected PC scores for Patients X_{PC3} and Y_{PC3} overlaid on distributions of PC scores from the discovery PSD sample. **(B)** Neural maps show cortical and subcortical ΔGBC^{obs} for the two patients X_{PC3} and Y_{PC3} specifically reflecting a difference from the mean *PC3*. The similarity of ΔGBC^{obs} and the $\beta_{PC3}GBC^{obs}$ map within the most predictive neural parcels for PC3 (outlined in green). Note that the sign of neural similarity to the reference *PC3* map and the sign of the *PC3* score is consistent for these two patients. **(C)** The selected *PC3* map (parcels outlined in green) is spatially correlated to the neural map reflecting the change in GBC after ketamine administration (ρ=0.76, **Methods**). Note that Patient X_{PC3} who exhibits ΔGBC^{obs} that is anti-correlated to the ketamine map also expresses depressive moods symptoms (panel **A**). This is consistent with the possibility that this person may clinically benefit from ketamine administration, which may elevate connectivity in areas where they show reductions (74). In contrast, Patient Y_{PC3} may exhibit an exacerbation of their psychosis symptoms given that their ΔGBC^{obs} is positively correlation with the ketamine map. **(D)** Data for two individual patients from the discovery dataset are highlighted for PC5: Q_{PC5} (blue) and Z_{PC5} (yellow). Note that no patients in the replication dataset were selected for *PC5* so both of these patients were selected from "discovery" PSD dataset for illustrative purposes. Patient Q_{PC5} loads highly negatively on the *PC5* axis and Patient Z_{PC5} loads highly positively. Density plots show the projected PC scores for Patients Q_{PC5} and Z_{PC5} overlaid on distributions of PC scores from the discovery PSD sample. **(E)** Neural maps show cortical and subcortical ΔGBC^{obs} for Patients Q_{PC5} and Z_{PC5}, which are highly negatively and positively correlated with the selected *PC5* map respectively. **(F)** The selected *PC5* map (parcels outlined in green) is spatially anti-correlated with the LSD response map (ρ=-0.44, see **Methods**), suggesting that circuits modulated by LSD (i.e. serotonin, in particular 5-HT2A) may be relevant for the *PC5* symptom expression. Here a serotonin receptor agonist may modulate the symptom-neural profile of Patient Q_{PC5}, whereas an antagonist may be effective for Patient Z_{PC5}.

tion" sample, which yielded consistent results (**Fig. S27F,H**). Collectively, these results show that data-driven symptom scores can pinpoint individual patients for whom neural variation strongly maps onto a target neural reference map. These data also highlight that both symptom and neural information for an independent patient can be quantified in the reference "discovery" BBS using their symptom data alone.

Subject-Specific PSD Neuro-Behavioral Features Track Neuropharmacological Map Patterns. Next, we use the personalized BBS selection in a proof-of-concept

framework for informing molecular mechanism of possible treatment response by relating subject-specific BBS to independently-acquired pharmacological neuroimaging maps. Here we examine two mechanisms implicated in PSD neuropathology via ketamine, a N-methyl-D-aspartate (NMDA) receptor antagonist (75), and lysergic acid diethylamide (LSD), primarily a serotonin receptor agonist (30, 76, 77). We first quantified individual subjects' BBS "locations" in the established reference neuro-behavioral geometry. The radarplot in **Fig. 7A** shows original symp-

toms whereas **Fig. 7B** shows ΔGBC^{obs} maps for two patients from the replication dataset (denoted here with X_{PC3} and Y_{PC3}, see **Fig. S12** for other example patients). Both of these patients exceeded the neural and behavioral BBS selection indices for *PC3* (defined independently in the "discovery" dataset, **Fig. S27C**). Furthermore, both patients exhibited neuro-behavioral variation in line with their expected locations in the BBS geometry. Specifically, patient X_{PC3} from the replication dataset scored highly negatively on the *PC3* axis defined in the "discovery" PSD sample (**Fig. 7A**). In contrast, patient Y_{PC3} scored positively on the *PC3* axis. Importantly, the correlation between the ΔGBC^{obs} map for each patient and the group-reference $\beta_{PC3}GBC$ was directionally consistent with their symptom PC score **Fig. 7B-C**).

We then tested if the single-subject BBS selection could be quantified with respect to a neural map reflecting glutamate receptor manipulation, a hypothesized mechanism underlying PSD symptoms (78). Specifically, we used an independently collected ketamine infusion dataset, collected in healthy adult volunteers during resting-state fMRI (79). As with the clinical data, here we computed a ΔGBC map reflecting the effect of ketamine on GBC relative to placebo (**Methods**). The maximally-predictive *PC3* parcels exhibited high spatial similarity with the ketamine map (ρ=0.76, see **Methods**), indicating that the ΔGBC pattern induced by ketamine tracks with the GBC pattern reflecting *PC3* symptom variation.

Critically, because X_{PC3} is negatively loaded on the *PC3* symptom axis, an NMDA receptor antagonist like ketamine may modulate symptom-relevant circuits in a way that reduces similarity with the *PC3* map. This may in turn have an impact on the *PC3*-like symptoms. Consistent with this hypothesis, X_{PC3} expresses predominantly depressive symptoms (**Fig. 7A**), and ketamine has been shown to act as an anti-depressant (74). This approach can be applied for patients that load along another axis, such as *PC5*. **Fig. 7D-E** shows the symptom and neural data for two patients whom met thresholds for *PC5* selection (**Fig. S27C**). Notably, the selected *PC5* map is anti-correlated with a ΔGBC map reflecting LSD vs. placebo effects (30) (ρ=-0.44, **Fig. 7F**). Hence areas modulated by LSD may map onto behavioral variation along *PC5*. Consequently, serotonergic modulation may be effective for treating Q_{PC5} and Z_{PC5}, via an antagonist or an agonist respectively. These differential similarities between pharmacological response maps and BBS maps (**Fig. S30**) can be refined for quantitative patient segmentation.

Group-Level PSD Neuro-Behavioral Features Track Neural Gene Expression Patterns. To further inform molecular mechanism for the derived BBS results, we compared results with neural gene expression profiles derived from the Allen Human Brain Atlas (AHBA) (32, 33)(**Fig. 8A**, **Methods**). Specifically, we tested if BBS cortical topographies, which reflect stable symptom-neural mapping along PSD, covary with the expression of genes implicated in PSD neuropathology. We focus here on serotonin receptor sub-

units (HTR1E, HTR2C, HTR2A), GABA receptor subunits (GABRA1, GABRA5), and the interneuron markers somatostatin (SST) and parvalbumin (PVALB). Serotonin agonists such as LSD have been shown to induce PSD-like symptoms in healthy adults (30) and the serotonin antagonism of "second-generation" antipsychotics are thought to contribute to their efficacy in targeting broad PSD symptoms (80–82). Abnormalities in GABAergic interneurons, which provide inhibitory control in neural circuits, may contribute to cognitive deficits in PSD (83–85) and additionally lead to downstream excitatory dysfunction that underlies other PSD symptoms (86, 87). In particular, a loss of prefrontal parvalbumin-expression fast-spiking interneurons has been implicated in PSD (88–91). **Fig. 8B** shows the distribution of correlations between the *PC3* map and the cortical expression patterns of 20,200 available AHBA genes (results for other PCs are shown in **Fig. S31**). Seven genes of interest are highlighted, along with their cortical expression topographies and their similarity with the *PC3* BBS map (**Fig. 8C-E**). This BBS-to-gene mapping can potentially reveal novel therapeutic molecular targets for neuro-behavioral variation. For example, the HTR1E gene, which encodes the serotonin 5-HT$_{1E}$ receptor, is highly correlated with the *PC3* BBS map. This could drive further development of novel selective ligands for this receptor, which are not currently available (92).

Discussion

We found a robust and reproducible symptom-neural mapping across the psychosis spectrum that emerged from a low-dimensional symptom solution. Critically, this low-rank symptom solution was predictive of a neural circuit pattern, which reproduced at the single-subject level. In turn, we show that the derived PSD symptom-neural feature maps exhibit spatial correspondence with independent pharmacological and gene expression neural maps that are directly relevant for PSD neurobiology. We demonstrate, for the first time in the literature, a transdiagnostic data-reduced PSD symptom geometry across hallmark psychopathology symptoms and cognitive measures, that shows stable and robust relationships with neural connectivity with singles subject-level precision. Critically, we anchor these symptom-neural maps to mechanistically-informed molecular imaging benchmarks that can be linked to relevant pathophysiology, and furthermore demonstrate how these methods can be combined to inform personalized patient selection decisions.

Deriving an Individually Predictive Low-Dimensional Symptom Representation Across the Psychosis Spectrum. Psychosis spectrum is associated with notable clinical heterogeneity such deficits in cognition as well as altered beliefs (i.e. delusions), perception (i.e. hallucinations), and affect (i.e. negative symptoms) (24). This heterogeneity is captured by clinical instruments that quantify PSD symptoms across dozens of specific questions and ratings. This yields a high-dimensional symptom space that is intractable for reliable mapping of neural circuits (27). Here we show that a low-rank solution captures principal axes of PSD symp-

Fig. 8. Psychosis spectrum symptom-neural maps track neural gene expression patterns computed from the Allen Human Brain Atlas (AHBA). (A) The symptom loadings and the associated neural map jointly reflect the *PC3* brain-behavioral space (BBS) profile, which can be quantitatively related to human cortical gene expression patterns obtained from the AHBA (33). **(B)** Distribution of correlation values between the *PC3* BBS map and ~20,000 gene expression maps derived from the AHBA dataset. Specifically, AHBA gene expression maps were obtained using DNA microarrays from six postmortem brains, capturing gene expression topography across cortical areas. These expression patterns were then mapped onto the cortical surface models derived from the AHBA subjects' anatomical scans and aligned with the Human Connectome Project (HCP) atlas, described in prior work and methods (33). Note that because no significant inter-hemispheric differences were found in cortical gene expression all results were symmetrized to the left hemisphere, resulting in 180 parcels. We focused on a select number of psychosis-relevant genes – namely genes coding for the serotonin and GABA receptor subunits and interneuron markers. Seven genes of interest are highlighted with dashed lines. Note that the expression pattern of HTR2C (green dashed line) is at the low negative tail of the entire distribution, i.e. highly anti-correlated with *PC3* BBS map. Conversely, GABRA1 and HTR1E are on the far positive end, reflecting a highly similar gene-to-BBS spatial pattern. **(C)** Upper panels show gene expression patterns for two interneuron marker genes, somatostatin (SST) and parvalbumin (PVALB). Positive (yellow) regions show areas where the gene of interest is highly expressed, whereas negative (blue) regions indicate low expression values. Lower panels highlight all gene-to-BBS map spatial correlations where each value is a symmetrized cortical parcel (180 in total) from the HCP atlas parcellation. **(D)** Gene expression maps and spatial correlations with the *PC3* BBS map for two GABA_A receptor subunit genes: GABRA1 and GABRA5. **(E)** Gene expression maps and spatial correlations with the *PC3* BBS map for three serotonin receptor subunit genes: HTR1E, HTR2C, and HTR2A.

tom variation, a finding in line with prior work in SZP (18–24, 93).

These results highlights two key observations: i) Existing symptom reduction studies (even those in SZP specifically) have not evaluated solutions that include cognitive impairment – a hallmark deficit across the psychosis spectrum (17). Here we show that cognitive performance captures a notable portion of the symptom variance independent of other axes. We observed that cognitive variation captured 10% of PSD sample variance even after accounting for "Global Functioning" psychopathology. ii) No study has quantified cognitive deficit variation along with core psychosis symptoms via dimensionality reduction across multiple PSD diagnoses. While existing studies have evaluated stability of data-reduced solutions within a single DSM category (18, 24, 94, 95), current results show that dimensionality-reduced PSD symptom solutions can be reproducibly obtained across DSM diagnoses.

Importantly, on each data-reduced symptom axis, some sub-

set of PSD patients received a score near zero. This does not imply that these patients were unimpaired; rather, the symptom configurations for these patients were orthogonal to variation along this specific axis. While in the current paper we demonstrate individual selection based on symptom scores along a single axis, it may also be possible to use a combination of several symptom dimensions to further inform individual treatment selection. For example, for patients with above-threshold symptom scores along several dimensions, a combination of these *PCs* may be more predictive of their neural features and in turn be a better indication of treatment selection for these individuals. Alternatively, the symptom dimensions can be used as exclusionary criteria for selectively picking out individuals along only one particular axis (e.g. filtering out patients above-threshold on *PCs 1, 3, 4,* and *5* to selectively study patients with only severe *PC2* cognitive deficits). Additionally, the ICA solution revealed oblique axes which are linear combinations of the *PCs*, e.g. *IC2* which is oblique to *PCs 1, 2,* and *3* (**Fig. S4**). However,

these *ICs* do not result in as robust or unique a neural circuit mapping as do the *PCs*, as shown in **Fig. S13**. The observation that PSD are associated with multiple complex symptom dimensions highlights an important intuition that may extend to other mental health spectra. Additionally, the PSD symptom axes reported here are neither definitive nor exhaustive. In fact, close to 50% of all clinical variance was not captured by the symptom PCA – an observation often overlooked in symptom data-reduction studies, which focus on attaining "predictive accuracy". Such studies rarely consider how much variance remains unexplained in the final data-reduced model and, relatedly, if the proportion of explained variance is reproducible across samples. This is a key property for reliable symptom-to-neural mapping. Thus, we tested if this reproducible low-dimensional PSD symptom space robustly mapped onto neural circuit patterns.

Leveraging a Robust Low-Dimensional Symptom Representation for Mapping Brain-Behavior Relationships. We show that the dimensionality-reduced symptom space improved the mapping onto neural circuit features (i.e. GBC), as compared to *a priori* item-level clinical symptom measures (**Fig. 3**). This symptom-neural mapping was highly reproducible across various cross-validation procedures and split-half replication (**Fig. 4**). The observed statistical symptom-neural improvement after dimensionality reduction suggests that data-driven clinical variation more robustly covaried with neural features. As noted, the low-rank symptom axes generalized across DSM diagnoses. Consequently, the mapping onto neural features (i.e. GBC) may have been restricted if only a single DSM category or clinical item was used. Importantly, as noted, traditional clinical scales are uni-directional (i.e. zero is asymptomatic, hence there is an explicit floor). Here, we show that data-driven symptom axes (e.g. *PC3*) were associated with bi-directional variation (i.e. no explicit floor effect). Put differently, patients who score highly on either end of these data-driven axes are severely symptomatic but in very different ways. If these axes reflect clinically meaningful phenomena at both tails, then they should more robustly map to neural feature variation, which is in line with reported effects. Therefore, by definition, the derived map for each of the *PCs* will reflect the neural circuitry that may be modulated by the behaviors that vary along that PC (but not others). For example, we named the *PC3* axis *"Psychosis Configuration"* because of its strong loadings onto conventional "positive" and "negative" PSD symptoms. This *PC3 "Psychosis Configuration"* showed strong positive variation along neural regions that map onto the canonical default mode network (DMN), which has frequently been implicated in PSD (96–101) suggesting that individuals with severe "positive" symptoms exhibit broadly elevated connectivity with regions of the DMN. On the contrary, this bi-directional *"Psychosis Configuration"* axis also showed strong negative variation along neural regions that map onto the sensory-motor and associative control regions, also strongly implicated in PSD (8, 25). The "bi-directionality" property of the *PC* symptom-neural maps may thus be desirable for identifying neural fea-

tures that support individual patient selection. For instance, it may be possible that *PC3* reflects residual untreated psychosis symptoms in this chronic PSD sample, which may reveal key treatment neural targets. In support of this circuit being symptom-relevant, it is notable that we observed a mild association between GBC and *PC* scores in the CON sample (**Fig. S11**). The symptom-neural mapping of dimensionality-reduced symptom scores also produced notable neural maps along other *PC* axes (**Fig. S10**). Higher *PC1* scores, indicating higher general functioning, may be associated with lower global connectivity in sensory/cingulate cortices and thalamus, but higher global connectivity with the temporal lobe. Higher *PC2* scores, indicating better cognitive functioning were positively associated with higher medial prefrontal and cerebellar GBC, and negatively with visual cortices and striatum. Importantly, the unique neural circuit variance shown in the $\beta_{PC2}GBC$ map suggest that there are PSD patients with more (or less) severe cognitive deficits independent of any other symptom axis, which would be in line with the observation that these symptoms are not treatable with antipsychotic medication (and therefore should not correlate with symptoms that are treatable by such medications; i.e. *PC3*). Of note, the statistics in the $\beta_{PC2}GBC$ map were relatively moderate, opening up the possibility that a more targeted measure of neural connectivity (e.g. seed FC, see below) would reveal a robust circuit map for this particular symptom dimension. Given the key role of cognitive deficits in PSD (17), this will be a particularly important direction to pursue, especially in prodromal or early-stage individuals where cognitive dysfunction has been shown to predict illness trajectories (102–105). The *PC4 – Affective Valence* axis was positively associated with higher GBC in cingulate and sensori-motor cortices and subcortically with the thalamus, basal ganglia, and anterior cerebellum, which may be implicated in the deficits in social functioning and affective aspects of this symptom dimension. Lastly, positive scores on *PC5*, which we named *"Agitation/Excitement"*, were associated with broad elevated global connectivity in the frontal associative and sensori-motor cortices, thalamus, and basal ganglia. Interestingly, however, the nucleus accumbens (as well as hippocampus, amygdala, cerebellum, and temporal lobe) appear to be negatively associated with *PC5* score (i.e. more severe grandiosity, unusual thought, delusional and disorganized thinking), illustrating another example of a bi-directional axis.

Deriving Individually Actionable Brain-Behavior Mapping Across the Psychosis Spectrum. Deriving a neurobehavioral mapping that is resolvable and stable at the individual patient level is a necessary benchmark for deploying symptom-neural "biomarkers" in a clinically useful way. Therefore, there is increasing attention placed on the importance of achieving reproducibility in the psychiatric neuroimaging literature (106–109), which becomes especially important for individualized symptom-neural mapping. Recently, several efforts have deployed multivariate methods to quantify symptom-neural relationships (15, 71, 72, 110–112), highlighting how multivariate techniques may perhaps

provide clinically innovative insights. However, such methods face the risk of overfitting for high-dimensional but underpowered datasets (26), as recently shown via formal generative modeling (27).

Here we attempted to use multivariate solutions (i.e. CCA) to quantify symptom and neural feature co-variation. In principle, CCA is well-suited to address the brain-behavioral mapping problem. However, symptom-neural mapping using CCA in our sample was not reproducible even when using a low-dimensional symptom solution and parcellated neural data as a starting point. Therefore, while CCA (and related multivariate methods such as partial least squares) are theoretically appropriate and may be helped by regularization methods such as sparse CCA, in practice many available psychiatric neuroimaging datasets may not provide sufficient power to resolve stable multivariate symptom-neural solutions (27). A key pressing need for forthcoming studies will be to use multivariate power calculators to inform sample sizes needed for resolving stable symptom-neural geometries at the single subject level. Of note, though we were unable to derive a stable CCA in the present sample, this does not imply that the multivariate neuro-behavioral effect may not be reproducible with larger effect sizes and/or sample sizes. Critically, this does highlight the importance of power calculations prior to computing multivariate brain-behavioral solutions (27).

Consequently, we tested if a low-dimensional symptom solution can be used in a univariate symptom-neural model to optimize individually predictive features. Indeed, we found that a univariate brain-behavioral space (BBS) relationship can result in neural features that are stable for individualized prediction. Critically, we found that if a patient exhibited a high PC symptom score, they were more likely to exhibit a topography of neural ΔGBC that was topographically similar to the $\beta_{PC}GBC$ map. This suggests that optimizing such symptom-neural mapping solutions (and ultimately extending them to multivariate frameworks) can inform cross-diagnostic patient segmentation with respect to symptom-relevant neural features. Importantly, this could directly inform patient identification based on neural targets that are of direct symptom relevance for clinical trial design.

Utilizing Independent Molecular Neuroimaging Maps to "Benchmark" Symptom-Relevant Neural Features. Selecting single patients via stable symptom-neural mapping of BBS solutions is necessary for individual patient segmentation, which may ultimately inform treatment indication. However, it is also critical to be able to relate the derived symptom-neural maps to a given mechanism. Here we highlight two ways to "benchmark" the derived symptom-neural maps by calculating their similarity against independent pharmacological neuroimaging and gene expression maps. We show a proof-of-principle framework for quantifying derived symptom-neural reference maps with two PSD-relevant neuropharmacological manipulations derived in healthy adults via LSD and ketamine. These analyses revealed that selecting single patients, via the derived symptom-neural mapping solution, can yield above-chance

quantitative correspondence to a given molecular target map. These data highlight an important effect: it is possible to construct a "strong inference" (113) evaluation of single patients' differential similarity to one molecular target map versus another. For instance, this approach could be applied to maps associated with already approved PSD treatments (such as clozapine, olanzapine, or chlorpromazine (114, 115)) to identify patients with symptom-neural configurations that best capture available treatment-covarying neural targets.

Relatedly, AHBA gene expression maps (33) may provide an *a priori* benchmark for treatment targets that may be associated with a given receptor profile. Here we show that identified BBS maps exhibit spatial correspondence with neural gene expression maps implicated in PSD – namely serotonin, GABA and interneuron gene expression. This gene-to-BBS mapping could be then used to select those patients that exhibit high correspondence to a given gene expression target. Collectively, this framework could inform empirically testable treatment selection methods (e.g. a patient may benefit from ketamine, but not serotonin agonists such as LSD/psilocybin). In turn, this independent molecular benchmarking framework could be extended to other approaches (e.g. positron emission tomography (PET) maps reflecting specific neural receptor density patterns (116, 117)) and iteratively optimized for quantitative patient-specific selection against actionable molecular targets.

Considerations for Generalizing Solutions Across Time, Severity and Mental Health Spectra. There are several constraints of the current result that require future optimization – namely the ability to generalize across time (early course vs. chronic patients), across a range of symptom severity (e.g. severe psychotic episode or persistent low-severity psychosis) and across distinct symptom spectra (e.g. mood). This applies to both the low-rank symptom solution and the resulting symptom-neural mapping. It is possible that the derived lower-dimensional symptom solution, and consequently the symptom-neural mapping solution, exhibits either time-dependent (i.e. state) or severity-dependent (i.e. trait) re-configuration. Relatedly, medication dose, type, and timing may also impact the solution. Another important aspect that will require further characterization is the possibility of oblique axes in the symptom-neural geometry. While orthogonal axes derived via PCA were appropriate here and similar to the ICA-derived axes in this solution, it is possible that oblique dimensions more clearly reflect the geometry of other psychiatric spectra and/or other stages in disease progression. For example, oblique components may better capture dimensions of neuro-behavioral variation in a sample of prodromal individuals, as these patients are exhibiting early-stage psychosis-like symptoms and may show signs of diverging along different trajectories.

Critically, these factors should constitute key extensions of an iteratively more robust model for individualized symptom-neural mapping across the PSD and other psychiatric spectra. Relatedly, it will be important to identify the 'limits' of a given BBS solution – namely a PSD-derived effect may not generalize into the mood spectrum (i.e. both the symptom

space and the resulting symptom-neural mapping is orthogonal). It will be important to evaluate if this framework can be used to initialize symptom-neural mapping across other mental health symptom spectra, such as mood/anxiety disorders.

These types of questions will require longitudinal and clinically diverse studies that start prior to the onset of full-blown symptoms (e.g. the North American Prodrome Longitudinal Study (NAPLS) (118, 119)). A corollary of this point is that ~50% of unexplained symptom variance in the current PCA solution necessitates larger samples with adequate power to map this subtle, but perhaps clinically essential, PSD variation.

Notably, the current cohort was adequately powered for symptom data reduction that drove univariate neural mapping. However, this sample was insufficiently powered for resolving stable multivariate symptom-neural relationships even with dimensionality-reduced symptom features. Consequently, the limited sample size necessitated choices for dimensionality reduction of the neural feature space in this study even for univariate analyses. While both parcellation and GBC constitute principled choices, symptom-relevant neural information may have been lost (which may be embedded in a higher-dimensional space). One obvious solution is to increase sample sizes (e.g. via datasets such as the UK Biobank (120)). However, in parallel, it will be critical to develop neurobiologically-informed feature space reduction and/or to optimize the stability of multivariate solutions via regularization methods. Another improvement would be to optimize neural data reduction sensitivity for specific symptom variation (121). We chose to use GBC for our initial geometry characterizations as it is a principled and assumption-free data-reduction metric that captures (dys)connectivity across the whole brain and generates neural maps (where each region can be represented by a value, in contrast to full functional connectivity matrices) that are necessary for benchmarking against molecular imaging maps. However, GBC is a summary measure that by definition does not provide information regarding connectivity between specific pairs of neural regions, which may prove to be highly symptom-relevant and informative. Thus symptom-neural relationships should be further explored with higher-resolution metrics, such as restricted GBC (65) which can summarize connectivity information for a specific network or subset of neural regions, or seed-based FC using regions implicated in PSD (e.g. thalamus (8, 99)).

Here we focused on the neural blood oxygen level dependent (BOLD) signal from fMRI. However, other modalities such as diffusion-weighted imaging (DWI), PET imaging, or electroencephalography (EEG) could be leveraged. Additional clinically-relevant information could be derived from genetics (such as polygenic risk scores (122–124)) or ecological momentary assessment (EMA) (125, 126), especially to parse state vs. trait biomarker variation.

Lastly, building on the proof-of-concept molecular neuroimaging comparisons, it will be imperative to eventually test such predictions in formal clinical trials. An actionable next step would be to optimize patient selection against existing treatments, which could result in higher success rates for drug development trials and potentially have massive impact for developing new interventions. Critically, the opportunity to develop, validate, and refine an individual-level quantitative framework could deliver a more rapid and cost-effective way of pairing patients with the right treatments.

Conclusions. We show that complex and highly heterogeneous PSD symptom variation can be robustly reduced into a low-rank symptom solution that is cross-diagnostic, individually predictive, generalizable and incorporates cognitive deficits. In turn, the derived PSD symptom axes robustly mapped onto distinct yet reproducible neural patterns, which were predictive at the single-patient level. Leveraging these stable results, we show a proof-of-principle framework for relating the derived symptom-relevant neural maps, at the individual patient level, with molecular targets implicated in PSD via LSD and ketamine neuro-pharmacological manipulations. Lastly, we used AHBA gene expression maps to show that identified PSD symptom-relevant neural maps covary with serotonin, GABA and interneuron neural gene expression patterns. This set of symptom-neural mapping results can be iteratively and quantitatively optimized for personalized treatment segmentation endpoints.

Competing Interests. J.L.J., C.F., J.B.B., G.R., Z.T., J.D., and J.D.M. consult for BlackThorn Therapeutics. A.A. consults and is a SAB member for BlackThorn Therapeutics Inc. W.J.M. is an employee of The Janssen Pharmaceutical Companies of Johnson & Johnson. J.L.J, J.D.M., and A.A. are co-inventors for the following pending patent: Anticevic A, Murray JD, Ji JL: Systems and Methods for Neuro-Behavioral Relationships in Dimensional Geometric Embedding (N-BRIDGE), PCT International Application No. PCT/US2119/022110, filed March 13, 2019. J.D.M., A.A., and W.J.M. are co-inventors for the following pending patent: Murray JD, Anticevic A, Martin, WJ: Methods and tools for detecting, diagnosing, predicting, prognosticating, or treating a neurobehavioral phenotype in a subject, U.S. Application No. 16/149,903 filed on October 2, 2018, U.S. Application or PCT International Application No. 18/054,009 filed on October 2, 2018.

ACKNOWLEDGEMENTS
We thank J.H. Krystal and I. Levy for useful discussions. This work was supported by the Heffter Research Institute (Grant No. 1-190420), Swiss Neuromatrix Foundation (Grant No. 2016-0111), Swiss National Science Foundation under the framework of Neuron Cofund (Grant No. 01EW1908), Slovenian Research Agency (ARRS) (Grant Nos. J7-8275, J7-6829, P3-0338 to GR), 1U01MH121766 (to AA), R01MH112746 (to JDM), SFARI Pilot Award (to JDM & AA), 5R01MH112189 (to AA), 5R01MH108590 (to AA), and by BlackThorn Therapeutics (J.D.M. & A.A.). Data were provided by the Bipolar-Schizophrenia Network on Intermediate Phenotypes (BSNIP) consortium via the National Institute of Mental Health Data Archive (NDA).

Bibliography

1. Mauricio Tohen, Eduard Vieta, Joseph Calabrese, Terence A Ketter, Gary Sachs, Charles Bowden, Philip B Mitchell, Franca Centorrino, Richard Risser, Robert W Baker, et al. Efficacy of olanzapine and olanzapine-fluoxetine combination in the treatment of bipolar i depression. *Archives of general psychiatry*, 60(11):1079–1088, 2003.
2. Joseph P McEvoy, Jeffrey A Lieberman, Diana O Perkins, Robert M Hamer, Hongbin Gu, Arthur Lazarus, Dennis Sweitzer, Christina Olexy, Peter Weiden, and Stephen D

Strakowski. Efficacy and tolerability of olanzapine, quetiapine, and risperidone in the treatment of early psychosis: a randomized, double-blind 52-week comparison. *American journal of Psychiatry*, 164(7):1050–1060, 2007.

3. Marije aan het Rot, Katherine A Collins, James W Murrough, Andrew M Perez, David L Reich, Dennis S Charney, and Sanjay J Mathew. Safety and efficacy of repeated-dose intravenous ketamine for treatment-resistant depression. *Biological Psychiatry*, 67(2):139–145, 2010.

4. Andrea Cipriani, Toshi A Furukawa, Georgia Salanti, Anna Chaimani, Lauren Z Atkinson, Yusuke Ogawa, Stefan Leucht, Henricus G Ruhe, Erick H Turner, Julian PT Higgins, et al. Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: a systematic review and network meta-analysis. *Focus*, 16(4):420–429, 2018.

5. American Psychiatric Association and American Psychiatric Association. Task Force on DSM-IV. *Diagnostic and Statistical Manual of Mental Disorders: DSM-IV*. DIAGNOSTIC AND STATISTICAL MANUAL OF MENTAL DISORDERS. American Psychiatric Association, 1994. ISBN 9780890420621.

6. Katharine A Phillips, Michael B First, and Harold Alan Pincus. *Advancing DSM: Dilemmas in psychiatric diagnosis*. American Psychiatric Pub, 2008.

7. Seth J Gillihan and Erik Parens. Should we expect" neural signatures" for dsm diagnoses? *Journal of Clinical Psychiatry*, 72(10):1383, 2011.

8. Alan Anticevic, Michael W Cole, Grega Repovs, John D Murray, Margaret S Brumbaugh, Anderson M Winkler, Aleksandar Savic, John H Krystal, Godfrey D Pearlson, and David C Glahn. Characterizing thalamo-cortical disturbances in schizophrenia and bipolar illness. *Cerebral cortex*, 24(12):3116–3130, 2013.

9. BJ Casey, Nick Craddock, Bruce N Cuthbert, Steven E Hyman, Francis S Lee, and Kerry J Ressler. Dsm-5 and rdoc: progress in psychiatry research? *Nature Reviews Neuroscience*, 14(11):810, 2013.

10. Thomas R Insel. The nimh research domain criteria (rdoc) project: precision medicine for psychiatry. *American Journal of Psychiatry*, 171(4):395–397, 2014.

11. Carol A Tamminga, Godfrey Pearlson, Matcheri Keshavan, John Sweeney, Brett Clementz, and Gunvant Thaker. Bipolar and schizophrenia network for intermediate phenotypes: outcomes across the psychosis continuum. *Schizophrenia bulletin*, 40(Suppl_2):S131–S137, 2014.

12. Nikolaos Koutsouleris, Lana Kambeitz-Ilankovic, Stephan Ruhrmann, Marlene Rosen, Anne Ruef, Dominic B Dwyer, Marco Paolini, Katharine Chisholm, Joseph Kambeitz, Theresa Haidl, et al. Prediction models of functional outcomes for individuals in the clinical high-risk state for psychosis or with recent-onset depression: a multimodal, multisite machine learning analysis. *JAMA psychiatry*, 75(11):1156–1172, 2018.

13. BJ Casey, Tariq Cannonier, May I Conley, Alexandra O Cohen, Deanna M Barch, Mary M Heitzeg, Mary E Soules, Theresa Teslovich, Danielle V Dellarco, Hugh Garavan, et al. The adolescent brain cognitive development (abcd) study: imaging acquisition across 21 sites. *Developmental cognitive neuroscience*, 32:43–54, 2018.

14. Adriana Di Martino, Chao-Gan Yan, Qingyang Li, Erin Denio, Francisco X Castellanos, Kaat Alaerts, Jeffrey S Anderson, Michal Assaf, Susan Y Bookheimer, Mirella Dapretto, et al. The autism brain imaging data exchange: towards a large-scale evaluation of the intrinsic brain architecture in autism. *Molecular psychiatry*, 19(6):659, 2014.

15. Andrew T Drysdale, Logan Grosenick, Jonathan Downar, Katharine Dunlop, Farrokh Mansouri, Yue Meng, Robert N Fetcho, Benjamin Zebley, Desmond J Oathes, Amit Etkin, et al. Resting-state connectivity biomarkers define neurophysiological subtypes of depression. *Nature medicine*, 23(1):28, 2017.

16. Lee J Cronbach and Paul E Meehl. Construct validity in psychological tests. *Psychological bulletin*, 52(4):281, 1955.

17. Deanna M Barch, Juan Bustillo, Wolfgang Gaebel, Raquel Gur, Stephan Heckers, Dolores Malaspina, Michael J Owen, Susan Schultz, Rajiv Tandon, Ming Tsuang, et al. Logic and justification for dimensional assessment of symptoms and related clinical phenomena in psychosis: relevance to dsm-5. *Schizophrenia research*, 150(1):15–20, 2013.

18. Mark van der Gaag, Anke Cuijpers, Tonko Hoffman, Mila Remijsen, Ron Hijman, Lieuwe de Haan, Berno van Meijel, Peter N van Harten, Lucia Valmaggia, Marc De Hert, et al. The five-factor model of the positive and negative syndrome scale i: confirmatory factor analysis fails to confirm 25 published five-factor solutions. *Schizophrenia research*, 85 (1-3):273–279, 2006.

19. Jean-Pierre Lindenmayer, Ruth Bernstein-Hyman, and Sandra Grochowski. A new five factor model of schizophrenia. *Psychiatric Quarterly*, 65(4):299–322, 1994.

20. Robin Emsley, Jonathan Rabinowitz, Martijn Torreman, RIS-INT-35 Early Psychosis Global Working Group, et al. The factor structure for the positive and negative syndrome scale (panss) in recent-onset psychosis. *Schizophrenia research*, 61(1):47–57, 2003.

21. Sonia Dollfus, Brian Everitt, Jean Marie Ribeyre, Francoise Assouly-Besse, Charlie Sharp, and Michel Petit. Identifying subtypes of schizophrenia by cluster analyses. *Schizophrenia bulletin*, 22(3):545–555, 1996.

22. Jack J Blanchard and Alex S Cohen. The structure of negative symptoms within schizophrenia: implications for assessment. *Schizophrenia bulletin*, 32(2):238–245, 2006.

23. Ji Chen, Kaustubh R Patil, Susanne Weis, Kang Sim, Thomas Nickl-Jockschat, Juan Zhou, André Aleman, Iris E Sommer, Edith J Liemburg, Felix Hoffstaedter, et al. Neurobiological divergence of the positive and negative schizophrenia subtypes identified on a new factor structure of psychopathology using non-negative factorization: An international machine learning study. *Biological Psychiatry*, 2019.

24. Jérémy Lefort-Besnard, Gaël Varoquaux, Birgit Derntl, Oliver Gruber, Andre Aleman, Renaud Jardri, Iris Sommer, Bertrand Thirion, and Danilo Bzdok. Patterns of schizophrenia symptoms: hidden structure in the panss questionnaire. *Transl Psychiatry*, 8(1):237, 10 2018. doi: 10.1038/s41398-018-0294-4.

25. Jie Lisa Ji, Caroline Diehl, Charles Schleifer, Carol A Tamminga, Matcheri S Keshavan, John A Sweeney, Brett A Clementz, S Kristian Hill, Godfrey Pearlson, Genevieve Yang, et al. Schizophrenia exhibits bi-directional brain-wide alterations in cortico-striato-cerebellar circuits. *Cerebral Cortex*, 29(11):4463–4487, 2019.

26. Richard Dinga, Lianne Schmaal, Brenda WJH Penninx, Marie Jose van Tol, Dick J Veltman, Laura van Velzen, Maarten Mennes, Nic JA van der Wee, and Andre F Marquand.

27. Markus Helmer, Shaun D Warrington, Ali-Reza Mohammadi-Nejad, Jie Lisa Ji, Amber Howell, Benjamin Rosand, Alan Anticevic, Stamatios N Sotiropoulos, and John D Murray. On stability of canonical correlation analysis and partial least squares with application to brain-behavior associations. *bioRxiv*, 2020. doi: 10.1101/2020.08.25.265546.

28. Carla M Canuso, Cynthia A Bossie, Young Zhu, Eriene Youssef, and David L Dunner. Psychotic symptoms in patients with bipolar mania. *Journal of Affective Disorders*, 111 (2-3):164–169, 2008.

29. Stanley R Kay, Abraham Fiszbein, and Lewis A Opler. The positive and negative syndrome scale (panss) for schizophrenia. *Schizophrenia bulletin*, 13(2):261–276, 1987.

30. Katrin H Preller, Joshua B Burt, Jie Lisa Ji, Charles H Schleifer, Brendan D Adkinson, Philipp Stämpfli, Erich Seifritz, Grega Repovs, John H Krystal, John D Murray, et al. Changes in global and thalamic brain connectivity in LSD-induced altered states of consciousness are attributable to the 5-HT2A receptor. *eLife*, 7:e35082, 2018.

31. Alan Anticevic, Philip R Corlett, Michael W Cole, Aleksandar Savic, Mark Gancsos, Yanqing Tang, Grega Repovs, John D Murray, Naomi R Driesen, Peter T Morgan, et al. N-methyl-d-aspartate receptor antagonist effects on prefrontal cortical connectivity better model early than chronic schizophrenia. *Biological Psychiatry*, 77(6):569–580, 2015.

32. Michael J Hawrylycz, Ed S Lein, Angela L Guillozet-Bongaarts, Elaine H Shen, Lydia Ng, Jeremy A Miller, Louie N Van De Lagemaat, Kimberly A Smith, Amanda Ebbert, Zackery L Riley, et al. An anatomically comprehensive atlas of the adult human brain transcriptome. *Nature*, 489(7416):391, 2012.

33. Joshua B Burt, Murat Demirtaş, William J Eckner, Natasha M Navejar, Jie Lisa Ji, William J Martin, Alberto Bernacchia, Alan Anticevic, and John D Murray. Hierarchy of transcriptomic specialization across human cortex captured by structural neuroimaging topography. *Nature Neuroscience*, 21(9):1251, 2018.

34. Shashwath A Meda, Zheng Wang, Elena I Ivleva, Gaurav Poudyal, Matcheri S Keshavan, Carol A Tamminga, John A Sweeney, Brett A Clementz, David J Schretlen, Vincent D Calhoun, et al. Frequency-specific neural signatures of spontaneous low-frequency resting state fluctuations in psychosis: evidence from bipolar-schizophrenia network on intermediate phenotypes (B-SNIP) consortium. *Schizophrenia bulletin*, 41(6):1336–1348, 2015.

35. Julia M Sheffield, Sridhar Kandala, Carol A Tamminga, Godfrey D Pearlson, Matcheri S Keshavan, John A Sweeney, Brett A Clementz, Dov B Lerman-Sinkoff, S Kristian Hill, and Deanna M Barch. Transdiagnostic associations between functional brain network integrity and cognition. *JAMA psychiatry*, 74(6):605–613, 2017.

36. Shashwath A Meda, Adrienne Gill, Michael C Stevens, Raymond P Lorenzoni, David C Glahn, Vince D Calhoun, John A Sweeney, Carol A Tamminga, Matcheri S Keshavan, Gunvant Thaker, et al. Differences in resting-state functional magnetic resonance imaging functional network connectivity between schizophrenia and psychotic bipolar probands and their unaffected first-degree relatives. *Biological Psychiatry*, 71(10):881–889, 2012.

37. Matthew F Glasser, Stamatios N Sotiropoulos, J Anthony Wilson, Timothy S Coalson, Bruce Fischl, Jesper L Andersson, Junqian Xu, Saad Jbabdi, Matthew Webster, Jonathan R Polimeni, et al. The minimal preprocessing pipelines for the human connectome project. *Neuroimage*, 80:105–124, 2013.

38. Jie Lisa Ji, Caroline Diehl, Charles Schleifer, Carol A Tamminga, Matcheri S Keshavan, John A Sweeney, Brett A Clementz, S Kristian Hill, Godfrey Pearlson, Genevieve Yang, Gina Creatura, John H Krystal, Grega Repovs, John Murray, Anderson Winkler, and Alan Anticevic. Schizophrenia Exhibits Bi-directional Brain-Wide Alterations in Cortico-Striato-Cerebellar Circuits. *Cerebral Cortex*, 06 2019. ISSN 1047-3211. doi: 10.1093/cercor/bhy306.

39. Mark Jenkinson, Peter Bannister, Michael Brady, and Stephen Smith. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage*, 17(2):825–841, 2002.

40. Martin Reuter, Nicholas J Schmansky, H Diana Rosas, and Bruce Fischl. Within-subject template estimation for unbiased longitudinal image analysis. *Neuroimage*, 61(4):1402–1418, 2012.

41. Jonathan D Power, Kelly Anne Barnes, Abraham Z Snyder, Bradley L Schlaggar, and Steven E Petersen. Steps toward optimizing motion artifact removal in functional connectivity mri; a reply to carp. *Neuroimage*, 76, 2013.

42. Alan Anticevic, Mark Gancsos, John D Murray, Grega Repovs, Naomi R Driesen, Debra J Ennis, Mark J Niciu, Peter T Morgan, Toral S Surti, Michael H Bloch, et al. Nmda receptor function in large-scale anticorrelated neural systems with implications for cognition and schizophrenia. *Proceedings of the National Academy of Sciences*, 109(41):16720–16725, 2012.

43. Jonathan D Power, Mark Plitt, Stephen J Gotts, Prantik Kundu, Valerie Voon, Peter A Bandettini, and Alex Martin. Ridding fmri data of motion-related influences: Removal of signals with distinct spatial and physical bases in multiecho data. *Proceedings of the National Academy of Sciences*, 115(9):E2105–E2114, 2018.

44. Jonathan D Power, Mark Plitt, Timothy O Laumann, and Alex Martin. Sources and implications of whole-brain fmri signals in humans. *Neuroimage*, 146:609–625, 2017.

45. Genevieve Y Yang, John D Murray, Matthew Glasser, Godfrey D Pearlson, John H Krystal, Charlie Schleifer, Grega Repovs, and Alan Anticevic. Altered global signal topography in schizophrenia. *Cerebral cortex*, 27(11):5156–5169, 2016.

46. Matthew F Glasser, Timothy S Coalson, Janine D Bijsterbosch, Samuel J Harrison, Michael P Harms, Alan Anticevic, David C Van Essen, and Stephen M Smith. Using temporal ica to selectively remove global noise while preserving global signal in functional mri data. *NeuroImage*, 181:692–717, 2018.

47. Kevin M Aquino, Ben D Fulcher, Linden Parkes, Kristina Sabaroedin, and Alex Fornito. Identifying and removing widespread signal deflections from fmri data: Rethinking the global signal regression problem. *bioRxiv*, page 662726, 2019.

48. Richard SE Keefe, Terry E Goldberg, Philip D Harvey, James M Gold, Margaret P Poe, and Leigh Coughenour. The brief assessment of cognition in schizophrenia: reliability, sensitivity, and comparison with a standard neurocognitive battery. *Schizophrenia research*, 68 (2-3):283–297, 2004.

49. Carol A. Tamminga, Elena I. Ivleva, Matcheri S. Keshavan, Godfrey D. Pearlson, Brett A.

Evaluating the evidence for biotypes of depression: Methodological replication and extension of. *NeuroImage: Clinical*, 22:101796, 2019.

Clementz, Bradley Witte, David W. Morris, Jeffrey Bishop, Gunvant K. Thaker, and John A. Sweeney. Clinical phenotypes of psychosis in the bipolar-schizophrenia network on intermediate phenotypes (b-snip). *American Journal of Psychiatry*, 170(11):1263–1274, 2013. doi: 10.1176/appi.ajp.2013.12101339. PMID: 23846857.

50. Michael W Cole, Genevieve J Yang, John D Murray, Grega Repovš, and Alan Anticevic. Functional connectivity change as shared signal dynamics. *Journal of neuroscience methods*, 259:22–39, 2016.

51. Jie Lisa Ji, Marjolein Spronk, Kaustubh Kulkarni, Grega Repovš, Alan Anticevic, and Michael W Cole. Mapping the human brain's cortical-subcortical functional network organization. *NeuroImage*, 185:35–57, 2019.

52. Matthew F Glasser, Timothy S Coalson, Emma C Robinson, Carl D Hacker, John Harwell, Essa Yacoub, Kamil Ugurbil, Jesper Andersson, Christian F Beckmann, Mark Jenkinson, et al. A multi-modal parcellation of human cerebral cortex. *Nature*, 536(7615):171, 2016.

53. Stephen M Smith and Thomas E Nichols. Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. *Neuroimage*, 44(1):83–98, 2009.

54. Anderson M Winkler, Gerard R Ridgway, Matthew A Webster, Stephen M Smith, and Thomas E Nichols. Permutation inference for the general linear model. *Neuroimage*, 92:381–397, 2014.

55. Tyler M Moore, Steven P Reise, Raquel E Gur, Hakon Hakonarson, and Ruben C Gur. Psychometric properties of the penn computerized neurocognitive battery. *Neuropsychology*, 29(2):235, 2015.

56. Ruben C Gur, Jan Richard, Paul Hughett, Monica E Calkins, Larry Macy, Warren B Bilker, Colleen Brensinger, and Raquel E Gur. A cognitive neuroscience-based computerized battery for efficient measurement of individual differences: standardization and initial construct validation. *Journal of neuroscience methods*, 187(2):254–262, 2010.

57. David C Van Essen, Stephen M Smith, Deanna M Barch, Timothy EJ Behrens, Essa Yacoub, Kamil Ugurbil, Wu-Minn HCP Consortium, et al. The wu-minn human connectome project: an overview. *Neuroimage*, 80:62–79, 2013.

58. Matcheri S Keshavan, David W Morris, John A Sweeney, Godfrey Pearlson, Gunvant Thaker, Larry J Seidman, Shaun M Eack, and Carol Tamminga. A dimensional approach to the psychosis spectrum between bipolar disorder and schizophrenia: the schizo-bipolar scale. *Schizophrenia research*, 133(1-3):250–254, 2011.

59. Hélène Verdoux and Jim van Os. Psychotic symptoms in non-clinical populations and the continuum of psychosis. *Schizophrenia research*, 54(1-2):59–65, 2002.

60. Roberto Nuevo, Somnath Chatterji, Emese Verdes, Nirmala Naidoo, Celso Arango, and José Luis Ayuso-Mateos. The continuum of psychotic symptoms in the general population: a cross-national study. *Schizophrenia bulletin*, 38(3):475–485, 2012.

61. Ian Kelleher and Mary Cannon. Psychotic-like experiences in the general population: characterizing a high-risk group for psychosis. *Psychological medicine*, 41(1):1–6, 2011.

62. NC Stefanis, M Hanssen, NK Smirnis, DA Avramopoulos, IK Evdokimidis, CN Stefanis, H Verdoux, and J Van Os. Evidence that three dimensions of psychosis have a distribution in the general population. *Psychological medicine*, 32(2):347–358, 2002.

63. Alan J Gelenberg. The catatonic syndrome. *The Lancet*, 307(7973):1339–1341, 1976.

64. Michael W Cole, Sudhir Pathak, and Walter Schneider. Identifying the brain's most globally connected regions. *Neuroimage*, 49(4):3132–3148, 2010.

65. Alan Anticevic, Margaret S Brumbaugh, Anderson M Winkler, Lauren E Lombardo, Jennifer Barrett, Phillip R Corlett, Hedy Kober, June Gruber, Grega Repovs, Michael W Cole, et al. Global prefrontal and fronto-amygdala dysconnectivity in bipolar i disorder with psychosis history. *Biological psychiatry*, 73(6):565–573, 2013.

66. Alex Fornito, Jong Yoon, Andrew Zalesky, Edward T Bullmore, and Cameron S Carter. General and specific functional connectivity disturbances in first-episode schizophrenia during cognitive control performance. *Biological psychiatry*, 70(1):64–72, 2011.

67. Avital Hahamy, Vince Calhoun, Godfrey Pearlson, Michal Harel, Nachum Stern, Fanny Attar, Rafael Malach, and Roy Salomon. Save the global: global signal connectivity as a tool for studying clinical populations with functional magnetic resonance imaging. *Brain connectivity*, 4(6):395–403, 2014.

68. Michael W Cole, Alan Anticevic, Grega Repovs, and Deanna Barch. Variable global dysconnectivity and individual differences in schizophrenia. *Biological psychiatry*, 70(1): 43–50, 2011.

69. Naomi R Driesen, Gregory McCarthy, Zubin Bhagwagar, Michael Bloch, Vincent Calhoun, Deepak C D'Souza, Ralitza Gueorguieva, George He, Ramani Ramachandran, Raymond F Suckow, et al. Relationship of resting brain hyperconnectivity and schizophrenia-like symptoms produced by the nmda receptor antagonist ketamine in humans. *Molecular psychiatry*, 18(11):1199–1204, 2013.

70. Michael D Fox, Abraham Z Snyder, Justin L Vincent, Maurizio Corbetta, David C Van Essen, and Marcus E Raichle. The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proceedings of the National Academy of Sciences*, 102 (27):9673–9678, 2005.

71. Cedric Huchuan Xia, Zongming Ma, Rastko Ciric, Shi Gu, Richard F Betzel, Antonia N Kaczkurkin, Monica E Calkins, Philip A Cook, Angel Garcia de la Garza, Simon N Vandekar, et al. Linked dimensions of psychopathology and connectivity in functional brain networks. *Nature communications*, 9(1):3003, 2018.

72. Meichen Yu, Kristin A Linn, Russell T Shinohara, Desmond J Oathes, Philip A Cook, Romain Duprat, Tyler M Moore, Maria A Oquendo, Mary L Phillips, Melvin McInnis, et al. Childhood trauma history is linked to abnormal brain connectivity in major depression. *Proceedings of the National Academy of Sciences*, 116(17):8582–8590, 2019.

73. David R Hardoon, Sandor Szedmak, and John Shawe-Taylor. Canonical correlation analysis: An overview with application to learning methods. *Neural computation*, 16(12):2639–2664, 2004.

74. Robert M Berman, Angela Cappiello, Amit Anand, Dan A Oren, George R Heninger, Dennis S Charney, and John H Krystal. Antidepressant effects of ketamine in depressed patients. *Biological Psychiatry*, 47(4):351–354, 2000.

75. John H Krystal, Laurence P Karper, John P Seibyl, Glenna K Freeman, Richard Delaney, J Douglas Bremner, George R Heninger, Malcolm B Bowers, and Dennis S Charney. Subanesthetic effects of the noncompetitive nmda antagonist, ketamine, in humans: psy-

chotomimetic, perceptual, cognitive, and neuroendocrine responses. *Archives of general psychiatry*, 51(3):199–214, 1994.

76. Javier González-Maeso, Noelia V Weisstaub, Mingming Zhou, Pokman Chan, Lidija Ivic, Rosalind Ang, Alena Lira, Maria Bradley-Moore, Yongchao Ge, Qiang Zhou, et al. Hallucinogens recruit specific cortical 5-ht2a receptor-mediated signaling pathways to affect behavior. *Neuron*, 53(3):439–452, 2007.

77. Christina T Egan, Katharine Herrick-Davis, Keith Miller, Richard A Glennon, and Milt Teitler. Agonist activity of LSD and lisuride at cloned 5ht2a and 5ht2c receptors. *Psychopharmacology*, 136(4):409–414, 1998.

78. Bita Moghaddam and Daniel Javitt. From revolution to evolution: the glutamate hypothesis of schizophrenia and its implication for treatment. *Neuropsychopharmacology*, 37(1):4–15, 2012.

79. Alan Anticevic, Mark Gancsos, John D. Murray, Grega Repovs, Naomi R. Driesen, Debra J. Ennis, Mark J. Niciu, Peter T. Morgan, Toral S. Surti, Michael H. Bloch, Ramachandran Ramani, Mark A. Smith, Xiao-Jing Wang, John H. Krystal, and Philip R. Corlett. Nmda receptor function in large-scale anticorrelated neural systems with implications for cognition and schizophrenia. *Proceedings of the National Academy of Sciences*, 109(41):16720–16725, 2012. ISSN 0027-8424. doi: 10.1073/pnas.1208494109.

80. Mark A Geyer and Franz X Vollenweider. Serotonin research: contributions to understanding psychoses. *Trends in pharmacological sciences*, 29(9):445–453, 2008.

81. H Y Meltzer, B W Massey, and M Horiguchi. Serotonin receptors as targets for drugs useful to treat psychosis and cognitive impairment in schizophrenia. *Current pharmaceutical biotechnology*, 13(8):1572–1586, 2012.

82. Anissa Abi-Dargham, Marc Laruelle, George K Aghajanian, Dennis Charney, and John Krystal. The role of serotonin in the pathophysiology and treatment of schizophrenia. *The Journal of neuropsychiatry and clinical neurosciences*, 9(1):1–17, 1997.

83. Francine M Benes and Sabina Berretta. Gabaergic interneurons: implications for understanding schizophrenia and bipolar disorder. *Neuropsychopharmacology*, 25(1):1–27, 2001.

84. Melis Inan, Timothy J. Petros, and Stewart A. Anderson. Losing your inhibition: Linking cortical gabaergic interneurons to schizophrenia. *Neurobiology of Disease*, 53:36–48, 2013. ISSN 0969-9961. doi: https://doi.org/10.1016/j.nbd.2012.11.013. What clinical findings can teach us about the neurobiology of schizophrenia?

85. Samuel J Dienel and David A Lewis. Alterations in cortical interneurons and cognitive function in schizophrenia. *Neurobiology of disease*, 131:104208, 2019.

86. John E Lisman, Joseph T Coyle, Robert W Green, Daniel C Javitt, Francine M Benes, Stephan Heckers, and Anthony A Grace. Circuit-based framework for understanding neurotransmitter and risk gene interactions in schizophrenia. *Trends in neurosciences*, 31(5): 234–242, 2008.

87. Anthony A Grace. Dysregulation of the dopamine system in the pathophysiology of schizophrenia and depression. *Nature Reviews Neuroscience*, 17(8):524, 2016.

88. John F Enwright III, Zhiguang Huo, Dominique Arion, John P Corradi, George Tseng, and David A Lewis. Transcriptome alterations of prefrontal cortical parvalbumin neurons in schizophrenia. *Molecular psychiatry*, 23(7):1606–1613, 2018.

89. Daniel J Lodge, Margarita M Behrens, and Anthony A Grace. A loss of parvalbumin-containing interneurons is associated with diminished oscillatory activity in an animal model of schizophrenia. *Journal of Neuroscience*, 29(8):2344–2354, 2009.

90. Clare L Beasley and Gavin P Reynolds. Parvalbumin-immunoreactive neurons are reduced in the prefrontal cortex of schizophrenics. *Schizophrenia research*, 24(3):349–355, 1997.

91. David A Lewis, Allison A Curley, Jill R Glausier, and David W Volk. Cortical parvalbumin interneurons and cognitive dysfunction in schizophrenia. *Trends in neurosciences*, 35(1): 57–67, 2012.

92. Sean L Kitson. 5-hydroxytryptamine (5-ht) receptor ligands. *Current pharmaceutical design*, 13(25):2621–2637, 2007.

93. Leonard White, Philip D Harvey, Lewis Opler, and JP Lindenmayer. Empirical assessment of the factorial structure of clinical symptoms in schizophrenia. *Psychopathology*, 30(5): 263–274, 1997.

94. Mark van der Gaag, Tonko Hoffman, Mila Remijsen, Ron Hijman, Lieuwe de Haan, Berno van Meijel, Peter N van Harten, Lucia Valmaggia, Marc De Hert, Anke Cuijpers, et al. The five-factor model of the positive and negative syndrome scale ii: a ten-fold cross-validation of a revised model. *Schizophrenia research*, 85(1-3):280–287, 2006.

95. Jean-Pierre Lindenmayer, Sandra Grochowski, and Ruth B Hyman. Five factor model of schizophrenia: replication across samples. *Schizophrenia research*, 14(3):229–234, 1995.

96. Susanna L Fryer, Scott W Woods, Kent A Kiehl, Vince D Calhoun, Godfrey Pearlson, Brian J Roach, Judith M Ford, Vinod H Srihari, Thomas H McGlashan, and Daniel H Mathalon. Deficient suppression of default mode regions during working memory in individuals with early psychosis and at clinical high-risk for psychosis. *Frontiers in psychiatry*, 4:92, 2013.

97. Alan Anticevic, Michael W Cole, John D Murray, Philip R Corlett, Xiao-Jing Wang, and John H Krystal. The role of default network deactivation in cognition and disease. *Trends in cognitive sciences*, 16(12):584–592, 2012.

98. Dost Öngür, Miriam Lundy, Ian Greenhouse, Ann K Shinn, Vinod Menon, Bruce M Cohen, and Perry F Renshaw. Default mode network abnormalities in bipolar disorder and schizophrenia. *Psychiatry Research: Neuroimaging*, 183(1):59–68, 2010.

99. Neil D Woodward, Baxter Rogers, and Stephan Heckers. Functional resting-state networks are differentially affected in schizophrenia. *Schizophrenia research*, 130(1-3):86–93, 2011.

100. Justin T Baker, Avram J Holmes, Grace A Masters, BT Thomas Yeo, Fenna Krienen, Randy L Buckner, and Dost Öngür. Disruption of cortical association networks in schizophrenia and psychotic bipolar disorder. *JAMA psychiatry*, 71(2):109–118, 2014.

101. Shashwath A Meda, Gualberto Ruaño, Andreas Windemuth, Kasey O'Neil, Clifton Berwise, Sabra M Dunn, Leah E Boccaccio, Balaji Narayanan, Mohan Kocherla, Emma Sprooten, et al. Multivariate analysis reveals genetic associations of the resting default mode network in psychotic bipolar disorder and schizophrenia. *Proceedings of the National Academy of Sciences*, 111(19):E2066–E2075, 2014.

102. Paolo Fusar-Poli, Giacomo Deste, Renata Smieskova, Stefano Barlati, Alison R Yung, Oliver Howes, Rolf-Dieter Stieglitz, Antonio Vita, Philip McGuire, and Stefan Borgwardt.

Cognitive functioning in prodromal psychosis: a meta-analysis. *Archives of general psychiatry*, 69(6):562–571, 2012.

103. E Bora, A Lin, SJ Wood, AR Yung, PD McGorry, and C Pantelis. Cognitive deficits in youth with familial and clinical high risk to psychosis: a systematic review and meta-analysis. *Acta Psychiatrica Scandinavica*, 130(1):1–15, 2014.

104. Emre Bora. Developmental trajectory of cognitive impairment in bipolar disorder: comparison with schizophrenia. *European Neuropsychopharmacology*, 25(2):158–168, 2015.

105. Kevin M Antshel, Wanda Fremont, Seetha Ramanathan, and Wendy R Kates. Predicting cognition and psychosis in young adults with 22q11. 2 deletion syndrome. *Schizophrenia bulletin*, 43(4):833–842, 2017.

106. Stephanie Noble, Dustin Scheinost, and R Todd Constable. A decade of test-retest reliability of functional connectivity: A systematic review and meta-analysis. *Neuroimage*, 203: 116157, 2019.

107. Joshua H Balsters, Dante Mantini, Matthew AJ Apps, Simon B Eickhoff, and Nicole Wenderoth. Connectivity-based parcellation increases network detection sensitivity in resting state fmri: an investigation into the cingulate cortex in autism. *NeuroImage: Clinical*, 11: 494–507, 2016.

108. Hengyi Cao, Sarah C McEwen, Jennifer K Forsyth, Dylan G Gee, Carrie E Bearden, Jean Addington, Bradley Goodyear, Kristin S Cadenhead, Heline Mirzakhanian, Barbara A Cornblatt, et al. Toward leveraging human connectomic data in large consortia: generalizability of fmri-based brain graphs across sites, sessions, and paradigms. *Cerebral Cortex*, 29(3):1263–1279, 2019.

109. Choong-Wan Woo, Luke J Chang, Martin A Lindquist, and Tor D Wager. Building better biomarkers: brain models in translational neuroimaging. *Nature Neuroscience*, 20(3):365, 2017.

110. Stephen M Smith, Thomas E Nichols, Diego Vidaurre, Anderson M Winkler, Timothy EJ Behrens, Matthew F Glasser, Kamil Ugurbil, Deanna M Barch, David C Van Essen, and Karla L Miller. A positive-negative mode of population covariation links brain connectivity, demographics and behavior. *Nature Neuroscience*, 18(11):1565, 2015.

111. Dominik A Moser, Gaelle E Doucet, Won Hee Lee, Alexander Rasgon, Hannah Krinsky, Evan Leibu, Alex Ing, Gunter Schumann, Natalie Rasgon, and Sophia Frangou. Multivariate associations among behavioral, clinical, and multimodal imaging phenotypes in patients with psychosis. *JAMA psychiatry*, 75(4):386–395, 2018.

112. Amanda L Rodrigue, Jennifer E McDowell, Neeraj Tandon, Matcheri S Keshavan, Carol A Tamminga, Godfrey D Pearlson, John A Sweeney, Robert D Gibbons, and Brett A Clementz. Multivariate relationships between cognition and brain anatomy across the psychosis spectrum. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 3(12): 992–1002, 2018.

113. John R Platt. Strong inference. *science*, 146(3642):347–353, 1964.

114. John Lally and James H MacCabe. Antipsychotic medication in schizophrenia: a review. *British medical bulletin*, 114(1):169–179, 2015.

115. S Miyamoto, GE Duncan, CE Marx, and JA Lieberman. Treatments for schizophrenia: a critical review of pharmacology and mechanisms of action of antipsychotic drugs. *Molecular psychiatry*, 10(1):79–104, 2005.

116. Lars Farde, Håkan Hall, Erling Ehrin, and Göran Sedvall. Quantitative analysis of d2 dopamine receptor binding in the living human brain by pet. *Science*, 231(4735):258–261, 1986.

117. Ryosuke Arakawa, Hiroshi Ito, Akihiro Takano, Hidehiko Takahashi, Takuya Morimoto, Takeshi Sassa, Katsuya Ohta, Motoichiro Kato, Yoshiro Okubo, and Tetsuya Suhara. Dose-finding study of paliperidone er based on striatal and extrastriatal dopamine d 2 receptor occupancy in patients with schizophrenia. *Psychopharmacology*, 197(2):229–235, 2008.

118. Jean Addington, Kristin S Cadenhead, Barbara A Cornblatt, Daniel H Mathalon, Thomas H McGlashan, Diana O Perkins, Larry J Seidman, Ming T Tsuang, Elaine F Walker, Scott W Woods, et al. North american prodrome longitudinal study (napls 2): overview and recruitment. *Schizophrenia research*, 142(1-3):77–82, 2012.

119. Larry J Seidman, Anthony J Giuliano, Eric C Meyer, Jean Addington, Kristin S Cadenhead, Tyrone D Cannon, Thomas H McGlashan, Diana O Perkins, Ming T Tsuang, Elaine F Walker, et al. Neuropsychology of the prodrome to psychosis in the napls consortium: relationship to family history and conversion to psychosis. *Archives of general psychiatry*, 67(6):578–588, 2010.

120. Cathie Sudlow, John Gallacher, Naomi Allen, Valerie Beral, Paul Burton, John Danesh, Paul Downey, Paul Elliott, Jane Green, Martin Landray, et al. Uk biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS medicine*, 12(3):e1001779, 2015.

121. Zarrar Shehzad, Clare Kelly, Philip T Reiss, R Cameron Craddock, John W Emerson, Katie McMahon, David A Copland, F Xavier Castellanos, and Michael P Milham. A multivariate distance-based analytic framework for connectome-wide association studies. *Neuroimage*, 93 Pt 1:74–94, Jun 2014. doi: 10.1016/j.neuroimage.2014.02.024.

122. Steven E Hyman. Diagnosis of mental disorders in light of modern genetics. *The conceptual evolution of DSM-5*, 3:18, 2011.

123. Michael C O'Donovan, Nick J Craddock, and Michael J Owen. Genetics of psychosis; insights from views across the genome. *Human genetics*, 126(1):3–12, 2009.

124. Hengyi Cao, Hang Zhou, and Tyrone D Cannon. Functional connectome-wide associations of schizophrenia polygenic risk. *Molecular Psychiatry*, pages 1–9, 2020.

125. Cheryl M Corcoran, Facundo Carrillo, Diego Fernández-Slezak, Gillinder Bedi, Casimir Klim, Daniel C Javitt, Carrie E Bearden, and Guillermo A Cecchi. Prediction of psychosis across protocols and risk cohorts using automated language analysis. *World Psychiatry*, 17(1):67–75, 2018.

126. Tara A Niendam, Laura M Tully, Ana-Maria Iosif, Divya Kumar, Kathleen E Nye, Jennifer C Denton, Lauren N Zakskorn, Taylor L Fedechko, and Katherine M Pierce. Enhancing early psychosis treatment using smartphone technology: a longitudinal feasibility and validity study. *Journal of Psychiatric Research*, 96:239–246, 2018.

127. A. Hyvarinen. Fast and robust fixed-point algorithms for independent component analysis. *IEEE Transactions on Neural Networks*, 10(3):626–634, May 1999. ISSN 1045-9227. doi: 10.1109/72.761722.

128. Nathaniel E. Helwig and Sungjin Hong. A critique of tensor probabilistic independent component analysis: Implications and recommendations for multi-subject fmri data analysis. *Journal of Neuroscience Methods*, 213(2):263–273, 3 2013. ISSN 0165-0270. doi: 10.1016/j.jneumeth.2012.12.009.

129. Rebekka Lencer, Andreas Sprenger, James L Reilly, Jennifer E McDowell, Leah H Rubin, Judith A Badner, Matcheri S Keshavan, Godfrey D Pearlson, Carol A Tamminga, Elliot S Gershon, et al. Pursuit eye movements as an intermediate phenotype across psychotic disorders: evidence from the b-snip study. *Schizophrenia research*, 169(1-3):326–333, 2015.

130. August B Hollingshead et al. Four factor index of social status. 1975.

131. Jaya L Padmanabhan, Neeraj Tandon, Chiara S Haller, Ian T Mathew, Shaun M Eack, Brett A Clementz, Godfrey D Pearlson, John A Sweeney, Carol A Tamminga, and Matcheri S Keshavan. Correlations between brain structure and symptom dimensions of psychosis in schizophrenia, schizoaffective, and psychotic bipolar i disorders. *Schizophrenia bulletin*, 41(1):154–162, 2015.

Supplementary Note 1: Independent Component Analysis (ICA) of Behavioral Symptom Data

We performed an independent component analysis (ICA) on the symptom data as a comparison to our PCA data-reduction solution. Specifically, an ICA was conducted on the behavioral data as an alternative dimensionality reduction method to the PCA. Five components were selected to allow for a direct comparison to be made to the results of the PCA **Fig. S4**. ICA decomposition was computed on the *436 x 36* (patients x symptom variables) data matrix using Hyvarinen's (127) FastICA algorithm, executed via the "icafast" package in R (128). Data were mean-centered before decomposition. After decomposition, the source signal estimate ("ICA score") for all subjects for each independent component (IC) was regressed on to the neural data across patients, as with *a priori* and PCA scores, described in the methods section. The proportion of variance explained by each independent component (IC) is shown in **Fig. S4A**. These percentages are comparable to those observed across all the PCs in **Fig. 1C**. **Fig. S4B** shows the correlations between subject scores across all five ICs and five PCs (N=436). Notably, certain pairs of components between the two solutions appear to be highly similar and exclusively mapped to each other (*IC5* and *PC4*; *IC4* and *PC5*). The correspondence between PCs 1-3 and ICs 1-3 appears to be more distributed as opposed to following a one-to-one mapping, likely because PCs are orthogonal to each other whereas ICs are not necessarily orthogonal. For example, *PC3* appears to correspond to *IC2*, but is negatively related to *IC3*, suggesting that these two ICs are oblique to the PC and perhaps share behavioral symptom variation in the ICA solution that is explained by a single *PC3*. Of note, *PC3* (the "Psychosis Configuration" axis) is a bi-directional axis whereby individuals who score either highly positively or highly negatively are highly symptomatic in different ways. The diverging nature of this axis may be not be captured well in the ICA solution, when orthogonality is not enforced. **Fig. S13F** shows the correlations across parcels (N=718 parcels) between the five IC maps (**Fig. S13A-E**) and five PC maps. Again, the neural pattern resembles that in **Fig. S13**, with certain pairs corresponding highly between the the ICA and PCA solutions. The coefficient maps for each of the five ICs are shown in **Fig. S13**. To compare the ICA solution to the PCA solution, we performed a correlation between the individual PCA and ICA scores across subjects, as well as between parcel coefficients of the 5 PCA maps and 5 ICA maps across parcels.

Supplementary Note 2: Establishing the Reproducibility of the Data-reduced Symptom-Neural Mapping Solution

Here we expand on the specific details regarding the cross-validation and reproducibility of the symptom-neural mapping result described in **Fig. 4**. Specifically, **Fig. 4D** shows a summary of a 5-fold bootstrapping symptom-neural map validation. Here the patient sample was randomly split into 5 folds and each fold was held out once as the regression was performed between symptom scores and parcel-level GBC in the other 4 folds. The resulting GBC coefficient map was then correlated with the full sample map performed in all 436 patients. Notably, the correlations between maps for each of the *k*-fold runs and the full model reached effects between r=0.8-0.9. The coefficient map for one such fold is shown in **Fig. 4E** for and the correlation between the coefficients in this map and the coefficients of the full regression model (performed in all 436 subjects) is shown in **Fig. 4F**. **Fig. 4G** shows a similar summary for leave-one-site-out validation for the regression of all five PC dimensions and traditional symptom scales onto neural GBC. Here, patients from one of the sites were left out while a regression was performed in all remaining patients. In turn, this effect was correlated with the coefficient map from the full patient sample. Again, the maps are highly similar for all sites, showing that the relationships between behavior and neural data are not driven by any one site in particular. Here, the coefficient map for one example is shown in **Fig. 4H** for PC3. Specifically, the results performed without Site 3 are shown because this is the site with the greatest number of patients and therefore may be the most likely to impact the final effect. The correlation between the coefficients in the resulting map without Site 3 and the coefficients of the full sample regression model are shown in the scatterplot in **Fig. 4I**. Finally, we performed a stratified split-half replication. Here, the full sample of patients was randomly divided into two halves (H1 & H2) with the proportion of each diagnostic group (BPP, SADP, SZP) preserved within each half. Then, PCA on the symptom data was computed in each half independently. As noted in the main text, this resulted in the "observed" PCA values for H1 and H2. In turn, we used loadings from H1 PCA to compute the "predicted" PC scores for subjects in H2, and vice versa. These "predicted" and "observed" PC scores were then independently regressed against parcelled GBC for subjects a given half. This yielded two maps for each PC: one for the "predicted" PC scores based on H1 data; and one for the "observed" PC scores from H1 data. These two symptom-neural maps were then correlated to evaluate reproducibility of the effects. This process was performed for both H1-to-H2 reproducibility as well as H2-to-H1 reproducibility. We performed 1,000 runs of the entire split-half replication analysis for the PC1-5 maps (**Fig. 4J**. Example data for *PC3* H1 and H2 are shown in **Fig. 4K-M**). These results highlight that the behavioral PC-to-GBC mapping yields a highly reproducible "brain-behavior" effect.

Supplementary Note 3: Multivariate Canonical Correlation Analysis of Symptom and Neural Features in PSD

CCA solves for transformation matrices Ψ and Θ such that the correlations between the linearly transformed 'latent' matrices U and V are maximal (see **Methods** for computational details). Matrix U contains the latent "behavioral" subject scores for all CV and is a linear composite of behavioral feature matrix B scaled by the loadings in Ψ; similarly, V is the latent neural score matrix and is the linear composite of neural features N transformed by Θ. Each column in U and V is known as a canonical variate (CV); each corresponding pair of canonical variates (e.g. $U1$ and $V1$) is referred to as a canonical mode. The correlations between the transformed latent matrices U and V are maximized (**Fig. 5B**), under the constraint that the canonical modes are orthogonal and independent to each other. Hence the correlation between each subsequent pair of CVs is computed from the residuals of the previous pair, until the number of columns in the smaller of the two matrices is reached.

In the main text we show that our CCA solution is not reproducible. We also examined whether the size of the neural feature space affects the CCA solution. To this end we computed CCA using:i) 180 symmetrized cortical parcels; ii) 359 bilateral subcortex-only parcels; iii) 192 symmetrized subcortical parcels; iv) 12 functional whole-brain networks, from the recently developed cortico-subcortical CAB-NP parcellation (51, 52). We show results for subcortical (192 neural features) **Fig. S14**, as well as network-level (12 features neural features) **Fig. S15**. We evaluated CCA significance via permutation testing whereby patient order was randomly shuffled 5,000 times (such that the association between behavioral and neural features was violated) and a CCA solution was computed for each shuffle; see **Methods**).

For each CCA solution, we examined how much symptom variance can be accounted for by the neural feature variance. Specifically, the correlation between behavioral features B and the latent neural matrix V (i.e. $N\Theta$) reflects the amount of variance in B that can be explained by the neural CVs in V (**Fig. 5E**). In the main text, **Fig. 5F** shows the proportion of behavioral variance explained by each of the neural CVs in a CCA performed between 180 neural features and all 36 symptom measures. **Fig. 5G** shows the result for the 5 PC-derived symptom dimensions. Given that the symptom PCs accounted for $\sim$50.93% of total symptom variance (see **Fig. 1**), the percentages here were scaled to reflect the total proportion of behavioral variance explained by each CV. The amount of neural variance explained by each behavioral CV is shown in **Fig. S17**.

Next, we examined the proportion of symptom variance explained by each neural canonical variate (CV). Because the PCs accounted for $\sim$50.93% of total symptom variance (**Fig. 1**), the percentages here were scaled accordingly. While the CCA using PC scores has fewer features than the one using item-level symptom measures, each neural CV explained more behavioral variance (**Fig. 5G** versus **5F**), suggesting that the PCs may capture neurally-relevant symptom variation with fewer features. Each CV has a pair of associated symptom and neural profiles, which can inform interpretation. This is shown for $CV3$ in **Fig. 5H–K**. Specifically, the latent scores for an example behavioral canonical variate (CV3) across diagnostic groups normalized to controls are shown in **Fig. 5H**. The profile of PC loadings onto $CV3$ are shown in **Fig. 5I**. Full results are shown in **Fig. S17**. CCA loadings are computed as the correlation of the latent matrix U (or V) and the input data matrix B (or N). These CCA loadings reflect the amount of variance in the original data that is extracted by the canonical variates. The high negative loadings of $CV3$ onto $PCs\ 1,4,5$ versus the high positive loadings of $CV3$ onto $PC3$ in **Fig. 5I** indicate that $CV3$ captures complex symptom variation. This can also be seen by the projecting 36 symptom measure loadings from a given PC onto $CV3$ **Fig. 5J**. Here the most positively loaded items were #7 "Delusions" (PANSS P1), #9 "Hallucinations" (PANSS P3), #21 "Somatic Concern" (PANSS G1) and #29 "Unusual Thought Content" (PANSS G9), while negatively-loaded items include #16 "Poor Rapport" ((PANSS N3), #19 "Lack of Spontaneity" (PANSS N6), and #28 "Uncooperativeness" (PANSS G8). Conversely, the corresponding neural factor loadings for $CV3$ are shown in **Fig. 5K**, illustrating GBC patterns associated with the $CV3$ profile. Data for all 5 CVs are shown in **Fig. S16**.

Interestingly, the cross-validation of the CCA effect was reproducible when using overlapping cross-validation strategies **Fig. S18**. However, as noted in the main text, the CCA solution was not reproducible with out-sample cross-validation strategies even though the effect passed p<0.05 permutation testing. This highlights the importance of rigorously testing for out-of-sample reproducibility of brain-behavioral effects across independent sub-samples (27).

Supplementary Note 4: Establishing the Reproducibility of the CCA Solution

Here we outline the details of the split-half replication for the CCA solution. Specifically, the full patient sample was randomly split (referred to as "H1" and "H2" respectively), while preserving the proportion of patients in each diagnostic group. Then, CCA was performed independently for H1 and H2. While the loadings for behavioral PCs and original symptom measures are somewhat similar (mean $r \approx 0.5$) between the two CCAs in each run, the neural loadings were not stable across H1 and H2 CCA solutions. Critically, CCA results did not perform well for leave-one-subject-out cross-validation (**Fig. 5M**). Here, one patient was held out while CCA was performed using all data from the remaining 435 patients. The loadings matrices Ψ and Θ from the CCA were then used to calculate the "predicted" neural and behavioral latent scores for all 5 CVs for the patient that was held out of the CCA solution. This process was repeated for every patient and the final result was evaluated for reproducibility. As described in the main text, this did not yield reproducible CCA effects (**Fig. 5M**).

Of note, CCA may yield higher reproducibility if the neural feature space were to be further reduced. As noted, our approach was to first parcellate the BOLD signal and then use GBC as a data-driven method to yield a neuro-biologically and quantitatively interpretable neural data reduction, and we additionally symmetrized the result across hemispheres. Nevertheless, in sharp contrast to the PCA univariate feature selection approach, the CCA solutions were still not stable in the present sample size of $N = 436$. Indeed, a multivariate power analysis (27) estimates that the following sample sizes will be required to sufficiently power a CCA between 180 neural features and 5 symptom features, at different levels of true canonical correlation (r_{true}):

- if $r_{true} = 0.1$: $n_{required} = 103,545$

- if $r_{true} = 0.3$: $n_{required} = 8,145$

- if $r_{true} = 0.5$: $n_{required} = 2,497$

To test if further neural feature space reduction may be improve reproducibility, we also evaluated CCA solutions with neural GBC parcellated according to 12 brain-wide functional networks derived from the recent HCP-driven network parcellation (51). Again, we computed the CCA for all 36 symptom measures as well as 5 *PCs* (**Fig. S15**). As with the parcel-level effects, the network-level CCA analysis produced significant results (for *CV1* when using 36 symptom measure scores and for all 5 CVs when using the 5 PC-derived scores). Here the result produced much lower canonical correlations (~0.3-0.5; however, these effects (for *CV1*) clearly exceeded the 95% confidence interval generated via random permutations, suggesting that they may reflect the true canonical correlation. We observed a similar result when we evaluated CCAs computed with neural GBC from 192 symmetrized subcortical parcels and 36 symptoms or 5 PCs (**Fig. S14**). In other words, data-reducing the neural signal to 12 functional networks likely averaged out parcel-level information that may carry symptom-relevant variance, but may be closer to capturing the true effect. Indeed, the power analysis suggests that the current sample size is closer to that needed to detect an effect with 12 vs. 5 features:

- if $r_{true} = 0.1$: $n_{required} = 8,855$

- if $r_{true} = 0.3$: $n_{required} = 696$

- if $r_{true} = 0.5$: $n_{required} = 213$

Note that we do not present a CCA conducted with parcels across the whole brain, as the number of variables would exceed the number of observations. However, the multivariate power analysis using 718 neural features and 5 symptom features estimates that the following sample sizes would be required to detect the following effects:

- if $r_{true} = 0.1$: $n_{required} = 421,608$

- if $r_{true} = 0.3$: $n_{required} = 33,166$

- if $r_{true} = 0.5$: $n_{required} = 10,168$

This analysis suggests that even the lowest bound of ~10k samples exceeds the present available sample size by two orders of magnitude.

Supplementary Note 5: Evaluating Individual Patients in Discovery and Replication Samples via Similarity to Target Brain-Behavior Spaces

We tested a hybrid neuro-behavioral patient selection strategy for single *PC* symptom axes by first imposing a PC-based symptom threshold, followed by a target neural similarity threshold driven by the most highly predictive symptom-neural map features. Specifically, the "neural similarity prediction index (NSPI)" computes a patient-specific Spearman's ρ between that patient's ΔGBC^{obs} and the group reference $\beta_{PC3}GBC^{obs}$ map using the maximally predictive $P = 39$ parcels (see **Fig. S28** for whole-brain results and alternative similarity metrics). **Fig. S27A** shows a significant relationship between each patient's *PC3* symptom score (X-axis) and the neural similarity index (Y-axis). In turn, **Fig. S27B** shows binned results, which provides a visual intuition for patient segmentation across both the neural and behavioral indices (**Fig. S27A**, right: binned by $\rho = 0.1$ & $PC3_{score} = 0.5$). For patients at either tail, the neuro-behavioral relationship was robust. Conversely, patients with a low absolute *PC3* score showed a weak relationship with symptom-relevant neural features. This is intuitive because these individuals do not vary along the selected PC symptom axis.

Fig. S27C shows the mean NSPI across subjects within each PC symptom bin along the X-axis. The resulting sigmoid captures that patients exhibit greater neural similarity if their PC symptom scores are more extreme. To evaluate if this relationship can yield a personalized patient selection, we computed the absolute NSPI **Fig. S27D**). The effect was approximated by a quadratic function, highlighting that patients with extreme *PC3* scores (either positive or negative) exhibited a stronger NSPI (i.e. personalized neural effects that strongly resembled the reference $(\beta_{PC3}GBC^{obs})^P = 39$ map). **Fig. S27E** shows the application of this neuro-behavioral selection procedure, demonstrating that PSD patients with extreme *PC3* scores (defined at the top/bottom 10th percentile of the "discovery" sample, $+2.17 < PC3_{score} < -2.41$) exhibit high NSPI values. We observed an inherent trade-off such that if the PC score threshold was raised then neural target similarity confidence goes up, but fewer patients will be selected. In the discovery PSD sample, all patients were accurately selected above the following neuro-behavioral thresholds: $90^{th}\%tile < PC3_{score} < 10^{th}\%tile$ and $|\rho| > 0.4$ (**Fig. S27E**, 34/436 patients selected, green line). We show consistent effects for when the selection was applied to *PC5* (**Fig. S27F**; results for all PCs in **Fig. S29**).

To test if the neuro-behavioral selection is generalizable, we used an independent cross-diagnostic sample of 30 patients diagnosed with SZP and 39 diagnosed with obsessive-compulsive disorder (OCD) (**Fig. S27G**, see **Methods** and **Table S4** for details). Applying the "discovery" selection thresholds yielded similar results for ∼6% of the cross-diagnostic "replication" sample for *PC3* (**Fig. S27G**, full analyses in **Fig. S28C**). Notably, no replication sample patients were selected along the neuro-behavioral thresholds for *PC5* (**Fig. S27H**). While there are SZP patients in the replication cross-diagnostic sample, few scored highly on *PC5* and none met the neural similarity threshold, emphasizing that not all patients within the same DSM-based diagnosis will exhibit variation along the same neuro-behavioral axis. Collectively, these results show that data-driven symptom scores can pinpoint individual patients for whom their neural variation strongly maps onto a target neural reference map. These data also highlight that both symptom and neural information for an independent patient can be quantified in the reference 'discovery' BBS using their symptom data alone.

Supplementary Tables

<div align="center">

Supplementary Table S1 | Glossary of Terms and Abbreviations

</div>

Abbreviation / Term	Definition
General & Behavioral Terms	
PSD	Psychosis spectrum disorder; or patients diagnosed with a PSD.
BPP	Bipolar disorder with psychosis; or patients diagnosed with BPP.
SADP	Schizoaffective disorder with psychosis; or patients diagnosed with SADP.
SZP	Schizophrenia with psychosis; or patients diagnosed with SZP.
CON	Control subjects.
BACS	Brief Assessment of Cognition in Schizophrenia.
PANSS	Positive and Negative Syndrome Scale.
Symptom measures	The 36 behavioral items from the BACS and PANSS.
PCA, PC	Principal component analysis, and principal component.
ICA, IC	Independent component analysis, and independent component.
Neural Terms	
FC	Functional connectivity (Fisher's r-to-Z transformed Pearson correlation values).
GBC	Global brain connectivity; computed for each brain location as the the mean FC across the whole brain.
$\beta_{PC}GBC$	The beta coefficient map from a mass univariate regression of PC scores on to GBC across subjects.
$\beta_{PC}GBC^{obs}$	The beta coefficient map from a mass univariate regression of observed (measured) symptom PC scores on to observed (measured) GBC across subjects.
$\beta_{PC}GBC^{pred}$	The beta coefficient map from a mass univariate regression of predicted symptom PC scores on to observed (measured) GBC across subjects.
GBC-PCA	A PCA performed on the neural GBC data across all N=638 PSD and CON subjects.
$\hat{GBC}_{woPC1}$	The GBC matrix reconstructed without the first PC of the GBC-PCA.
$\hat{GBC}_{woPC1-2}$	The GBC matrix reconstructed without the first PC of the GBC-PCA.
$\hat{GBC}_{woPC1-3}$	The GBC matrix reconstructed without the first two PCs of the GBC-PCA.
$\hat{GBC}_{woPC1-4}$	The GBC matrix reconstructed without the first three PCs of the GBC-PCA.
Multivariate Analyses Terms	
CCA	Canonical correlation analysis.
CV	Canonical variate from a CCA solution.
180 vs. 36 CCA	The CCA computed between 180 neural features and 36 symptom measures.
180 vs. 5 CCA	The CCA computed between 180 neural features and 5 symptom PCs.
N, B	The input neural and behavioral data matrices in a CCA, respectively.
$\hat{N}$, $\hat{B}$	The predicted neural and behavioral data matrices in the CCA cross-validation, respectively.
θ, Ψ	The neural and behavioral transformation matrices in a CCA, respectively.
U, V	The transfomed neural and behavioral data matrices in a CCA, respectively.
Single-Subject Prediction Analyses Terms	
ΔGBC	The difference between an individual's measured GBC map and the mean GBC map computed for the whole group; reflects individual differences in GBC, which may otherwise be conflated by the group mean.
$\beta_{PC}GBC^{obs}_{N-1}$	The beta coefficient map computed across N-1 subjects, using observed symptom PC and GBC data.
S^{obs}	The vector of observed symptom scores for a single PC (i.e. from a PCA computed using symptom measures), with length N-1 (as subject i is left out).
S^{pred}	The predicted symptom PC for subject i, using the loadings from a leave-one-out PCA computed on N-i subjects.
dpGBC	Dot product between a subject's ΔGBC map and a $\beta_{PC}GBC$ map.
$dpGBC^{obs}_i$	Dot product between a subject i's ΔGBC map and the $\beta_{PC}GBC^{obs}_{N-1}$ map computed without that subject.
$dpGBC^{pred}_i$	The predicted dpGBC for a subject i, predicted from a regression model using the $dpGBC^{obs}$ from all other subjects and the predicted S^{pred} for subject i.
P_{select}	The number of parcels in the subset of the brain map used in a particular analysis. P_{select}=718 for the full whole-brain map.
BBS	Brain-Behavior Space; the symptom and neural mapping for a particular dimension.

<div align="center">

Table S1. Glossary of the key terms and abbreviations used in the study.

</div>

Ji *et al.* | Brain-Behavioral Relationships in Psychosis

Supplementary Table S2 | BSNIP Dataset Clinical and Demographic Characteristics

Characteristic	HCS (N =202)		PSD (N=436)		BPP (N=150)		SADP (N=119)		SZP (N=167)	
	Mean	S.D.	Mean	S.D	Mean	S.D.	Mean	S.D.	Mean	S.D.
Age (years)	37.18	12.18	35.33	12.31	36.29	12.94	35.86	11.87	34.16	11.99
Gender (% male)	42.08	-	48.62	-	31.33	-	44.54	-	67.07	-
Parental Education (years)	13.35	3.42	13.67	3.43	14.16	3.44	13.05	3.95	13.62	2.93
Participant's Education	14.76	2.27	13.47	2.34	14.22	2.31	13.25	2.22	12.93	2.28
Handedness (% right)	85.64	-	85.78	-	83.33	-	88.24	-	86.23	-
Signal-to-noise (SNR)	231.03	82.70	218.34	93.44	229.78	100.09	230.93	93.83	198.10	83.13
% Frames Flagged	2.81	3.99	6.17	9.09	6.41	9.67	5.32	9.32	6.57	8.35
Medication (CPZ equivalents)	-	-	443.20	402.92	318.61	321.06	514.43	462.70	490.64	395.76
PANSS Positive Symptoms	7.03	0.30	15.66	5.35	12.87	4.35	18.19	5.20	16.37	5.15
PANSS Negative Symptoms	7.01	0.16	14.59	5.10	12.01	3.66	15.55	4.52	16.22	5.68
PANSS General Psychopathology	16.04	0.36	31.45	8.67	28.72	8.20	34.57	8.70	31.67	8.30
PANSS Total Psychopathology	30.08	0.61	61.67	16.32	53.59	13.69	68.22	15.72	64.26	16.04
BACS Cognitive Score (Z)	0.02	1.10	-1.22	1.20	-0.83	1.19	-1.30	1.10	-1.52	1.19

Table S2. Clinical and demographic details for the BSNIP sample. Clinical and demographic measures are shown at the group level for the BSNIP sample used for initial "discovery" analyses. Means and standard deviations were computed for each of the diagnostic groups. Abbreviations: CON, controls; PSD, psychosis spectrum disorder patients; BPP, bipolar disorder with psychosis; SADP, schizoaffective disorder with psychosis; SZP, schizophrenia; CPZ, chlorpromazine; PANSS, Positive and Negative Syndrome Scale for Schizophrenia, BACS, Brief Assessment of Cognition in Schizophrenia.

Supplementary Table S3 | BSNIP Dataset Scan Acquisition Parameters

BOLD	TR (ms)	TE (ms)	FA (deg.)	Slices (N)	Slice Order	Matrix (mm)	Voxel (mm)	Vendor
Baltimore	2210	30	70	36	I-A	64 x 64	3.4 x 3.4 x 3	Siemens Trio
Hartford	1500	27	70	29	S-A	64 x 64	3.4 x 3.4 x 5	Siemens Allegra
Detroit	1570 /1720	22	60	29	S-A	64 x 64	3.4 x 3.4 x 4	Siemens Trio
Dallas	1500	27	60	29	S-A	64 x 64	3.4 x 3.4 x 4	Philips
Chicago	1775	27	60	29	S-A	64 x 64	3.4 x 3.4 x 4	GE Signa HDX
Boston	3000	27	60	30	S-A	64 x 64	3.4 x 3.4 x 5	GE Signa HDX

T1w	TR (ms)	TE (ms)	FA (deg.)	Slices (N)	Slice Order	Matrix (mm)	Voxel (mm)	Vendor
Baltimore	2300	2.91	9	160	—	256 x 240	1 x 1 x 1.2	Siemens Trio
Hartford	2300	2.91	9	160	—	256 x 240	1 x 1 x 1.2	Siemens Allegra
Detroit	2300	2.94	9	160	—	256 x 240	1 x 1 x 1.2	Siemens Trio
Dallas	6.6	2.8	8	170	—	256 x 256	1 x 1 x 1.2	Philips
Chicago	6.98	2.84	8	166	—	256 x 256	1 x 1 x 1.2	GE Signa HDX
Boston	6.98	2.84	8	166	—	256 x 256	1 x 1 x 1.2	GE Signa HDX

Table S3. Scan acquisition parameters for the BSNIP dataset across sites. Scan acquisition details are shown for the 6 BSNIP acquisition sites. Adapted from (34).

Supplementary Table S4 I Replication Cross-diagnostic Dataset Clinical and Demographic Characteristics

Characteristic	CON (N=34)		PTT (N=69)		OCD (N=39)		SCZP (N=30)	
	Mean	S.D.	Mean	S.D	Mean	S.D.	Mean	S.D.
Age (years)	29.91	8.11	33.7	12.89	32.23	12.32	35.63	13.63
Gender (% male)	61.7	-	59.4	-	46.1	-	76.7	-
Parental Education (years)	15.05	3.31	13.98	3.45	14.02	3.78	13.93	3.01
Participant's Education	16.37	2.34	14.16	2.12	15.22	1.95	12.79	2.34
Handedness (% right)	93.9	-	91.3	-	84.6	-	100.0	-
Signal-to-noise (SNR)	53.45	5.59	51.86	5.63	52.29	5.28	51.33	6.08
% Frames Flagged	0.21	0.53	7.27	15.53	5.11	13.81	10.07	17.37
PANSS Positive Symptoms	7.58	1.06	11.40	4.23	8.97	2.74	14.56	6.16
PANSS Negative Symptoms	7.75	1.57	11.06	3.59	8.80	2.21	14.00	5.39
PANSS General Psychopathology	32.62	2.62	48.80	10.96	42.71	9.07	56.72	13.42
PANSS Total Psychopathology	47.95	5.25	71.26	18.78	60.48	14.02	85.28	24.97
BACS Cognitive Score (Z)	0.00	1.00	-0.67	1.45	-0.07	1.03	-1.45	1.55

Table S4. Clinical and demographic details for the independent replication cross-diagnostic sample. Clinical and demographic measures are shown at the group level for the independent clinical sample used for replication analyses presented in main text **Fig.8**. Means and standard deviations were computed for each of the diagnostic groups. Abbreviations: CON, controls; PTT, all patients; OCD, obsessive-compulsive disorder; SZP, schizophrenia; PANSS, Positive and Negative Syndrome Scale for Schizophrenia, BACS, Brief Assessment of Cognition in Schizophrenia.

Supplementary Figures

Question	Results	Corresponding Figures
Can data-reduction methods (e.g. principal component analysis, PCA) reliably map symptom axes across PSD that include both canonical symptoms and cognitive deficits?	The PCA on 36 behavioral measures and 436 PSD patients produced five significant components (PCs) spanning key symptom domains and oblique to conventional symptom factors and diagnostic categories.	Main Fig. 1 Supplementary Fig. S3-S4
Are the derived dimensionality-reduced symptom axes in PSD stable and replicable?	The behavioral PCA solution was stable across split-half replication, leave-site-out, and k-fold cross-validations, and not driven by medication effects.	Main Fig. 2 Supplementary Fig. S5-S7
Do dimensionality-reduced symptom axes yield stronger statistical neural maps across PSD, relative to conventional scales & diagnoses?	PC symptom scores yielded a much stronger statistical effect with neural features (as measured by global brain connectivity, GBC) across patients than did conventional symptom scores.	Main Fig. 3 Supplementary Fig. S8-S10
Are the PC brain-behavior maps stable and replicable?	PC brain-behavior maps are stable and reproducible across split-half, leave-site-out and k-fold cross-validations.	Main Fig. 4 Supplementary Fig. S11
Can a multivariate approach (e.g. canonical correlation analysis, CCA) be used to derive a stable neuro-behavioral mapping?	CCA produced a stable within-sample solution, but performed poorly on out-of-sample cross-validations.	Main Fig. 5 Supplementary Fig. S12-S16
Is the computed symptom-to-neural mapping actionable for individual patient selection?	A majority of neural variance in PSD patients is not symptom-relevant. However, brain-behavioral maps can be optimized to select specific patients along PC axes. This method can also be applied to select patients from independent datasets.	Main Fig. 6 & 7 Supplementary Fig. S17-S22
Can we inform molecular mechanism and treatment decisions by relating patient-specific brain-behavioral maps to independently-acquired pharmacological neuroimaging maps?	Pharmacological neuroimaging maps targeting mechanisms of interest (e.g. ketamine and LSD) can be used as independent benchmarks against which patient-specific brain-behavioral maps can be evaluated.	Main Fig. 7 Supplementary Fig. S23
Can we inform molecular mechanism and potential novel therapeutic targets by relating brain-behavior maps to gene expression maps?	Gene expression maps for genes implicated in PSD neuropathology (e.g. serotonin and GABA receptors, interneuron markers) can inform mechanism and potential novel targets relating to specific axes of neuro-behavioral variation.	Main Fig. 8 Supplementary Fig. S24

Fig. S1. Summary of the key questions, results, and supporting data in this study.

Fig. S2. Study workflow used study to quantify shared neural and behavioral variation in individuals diagnosed with PSD.
(A) Data from the BSNIP study were acquired from the National Institute of Health Data Archive (NDA). T1-weighted structural and resting-state BOLD neuroimaging data were obtained for a total of 638 individuals (202 controls, 150 patients with a diagnosis of bipolar disorder with psychosis, 119 with a diagnosis of schizoaffective disorder, and 167 patients with schizophrenia). Data were processed through the Human Connectome Project's (HCP) Minimal Preprocessing Pipeline with modifications made for "legacy" BOLD and T1w data, which are now featured as a standard option in the HCP pipelines provided by our team (https://github.com/Washington-University/HCPpipelines/pull/156) using Yale High Performance Computing clusters. **(B)** Symptom data were first normalized across the sample. The correlation matrix across all 436 PSD patients and 36 symptom measures was computed followed by dimensionality reduction (e.g. using PCA or ICA). The dimensionality-reduced solution was then cross-validated to assess stability and reproducibility across sites, *k*-fold cross-validations, leave-subject-out and split-half approaches. **(C)** In parallel, all neuroimaging data were processed as noted above. T1w and resting-state BOLD images were preprocessed using a modified version of the HCP Minimal Preprocessing Pipeline, including individual-subject registration of structural and function data, de-noising, and mapping of BOLD data on a hybrid surface-volume cortico-subcortical format (Connectivity Informatics Technology Initiative [CIFTI] format, see **Methods** for details). After registration to a standard CIFTI template BOLD data were parcellated at the individual subject level using the Cole-Anticevic Brain-wide Network Partition (CAB-NP), which is a functionally-defined network and parcel-level partition in the CIFTI space encompassing both cortex and subcortex (51, 52). Lastly, a global brain connectivity (GBC) map for each subject was computed by taking the mean functional connectivity of each parcel with all other parcels in the brain at the single subject level. **(D)** After symptom and neural data were fully processed in tandem, the symptom-to-neuroimaging mapping was quantified across subjects. Specifically, the relationship between data-reduced symptom scores and GBC was computed for each parcel across all patients. This produced a group-level symptom-neural map, which was subsequently cross-validated using leave-site-out, *k*-fold, leave subject-out as well as split-half approaches to assess reproducibility of the effect. Finally, the stable symptom-neural mapping result was further feature-optimized for single-subject prediction. This yielded a set of parcels for quantifying patient-specific symptom-neural prediction based on a cross-validated group reference map as well as comparison of the selected parcels with independent molecular neuroimaging maps (i.e. pharmacological (30, 42) and gene expression maps (33)).

Fig. S3. Alternative views of the behavioral PCA triplot. (A–C) Alternative views of the triplot in **Fig. 1F** showing the relationship between the three principal axes of variation in behavior and standard clinical symptom factors. Each point represents an individual subject projected into the geometry defined by the first 3 principal components (PC). Vectors show the projections of standard symptoms factors. **(D–F)** Alternative views of the triplot in panels **A–C**, where each sphere represents the mean of each a priori clinical group. Vectors show the projections of standard symptoms factors [PANSS positive (purple), negative (blue), general (pink) symptoms and BACS cognitive performance (green)]. BPP, bipolar disorder; SADP, schizoaffective disorder; SZP, schizophrenia; CON, controls. **(G)** Heatmap of the loadings of each of the 36 symptom measures on the five significant PCs (also seen in radarplot in **Fig. 1E**). Positive loadings are indicated in red; negative loadings are shown in blue. Each PC is named based on its most strongly loaded items.

Fig. S4. Independent component analysis (ICA) as an alternative method of dimensionality-reduction of symptom data. (A) Screeplot showing the total proportion of variance explained by each independent component (IC) in a five-component solution performed across all 36 behavioral measures in 436 patients. The size of each point is proportional to the variance explained by that IC. **(B)** Correlation matrix showing correlations of individual subject scores for the 5 significant principal components (PCs) from the PCA solution shown in **Fig. 1** and the five ICs from the ICA solution, across all 436 subjects. Neural maps for each IC are shown in **Fig. S13**.

Fig. S5. k-Fold cross-validation for symptom-driven principal component analysis (PCA). These results show a 5-fold cross-validation analysis to test the stability of the PCA solution. The full patient sample was first randomly split into 5 sets and patients were randomly assigned to one of 5 subsets. Each subset of patients was then used as an independent 'test sample' in a PCA that was derived from the other 4 subsets. Screeplot shows proportion of variance explained by each of the PCs in a PCA of all 36 behavioral measures, excluding a subset of 88 patients. The number of significant PCs determined via a permutation test and the total proportion of variance explained by these PCs are all comparable to the full model shown in the main text . To obtain a 'predicted' PC score for the 88 patients in the excluded subset, the loadings from the model obtained from the other 348 patients were used. The 'observed' PC scores are the scores from the full model of the same 88 patients. The scatterplot shows that the predicted and observed scores for *PC1* are highly correlated (*r*=0.999), suggesting that the PCA solution is stable and predictive at the individual-subject level. Similarly, predicted and observed scores are highly correlated for all five PCs. **(B–E)** The results of the PCA are also highly comparable and predictive for the other four folds.

Fig. S6. Leave-site-out cross-validation for symptom-driven principal component analysis (PCA). These results show a 6-fold leave-site-out cross-validation analysis to test the stability of the PCA solution when a given site is excluded from the model. The full patient sample was first split into 6 sets according to data collection site. Each held-out site was then used as an independent 'test sample' in a PCA that was derived from the other 5 sites. **(A)** Proportion of variance explained by each of the PCs in a PCA of all 36 behavioral measures, excluding one of the six sites at which data was collected (here we excluded the Boston site). The number of significant PCs determined via a permutation test and the total proportion of variance explained by these PCs are all comparable to the full model shown in **Fig. 1C**. To obtain a 'predicted' PC score for the 46 patients in the excluded site, the loadings from the model obtained from the other 390 patients were used. The 'observed' PC scores are the scores from the full model of the same 88 patients. The scatterplot shows that the predicted and observed scores for *PC1* are highly correlated (*r*=0.999), suggesting that the PCA solution is stable and robustly predictive at the individual patient level. Predicted and observed scores are highly correlated for all five PCs. **(B–F)** The results of the PCA are also highly comparable for the other 5 sites, suggesting that possible site differences in evaluating patient symptoms or patient sample composition are not impacting the obtained PCA solutions.

A Correlation between Medication Dosage (Chloropromazine Equivalent) and PC Scores

○ Patients not on antipsychotic medication (N=59)
● Patients on antipsychotic medication (N=279)

PC1 Score: ρ = -0.102, p = 0.087, N = 279
PC2 Score: ρ = -0.282, p = 1.33e-6, N = 279
PC3 Score: ρ = 0.104, p = 0.078, N = 279
PC4 Score: ρ = -0.066, p = 0.268, N = 279
PC5 Score: ρ = -0.144, p = 0.015, N = 279

CPZ Equivalent (mg)

B Differences in Mean PC Scores of Patients on No Medication vs Antipsychotic Medication

PC1 Score: t = 1.07, df = 77.80, p = 0.289
PC2 Score: t = 4.28, df = 80.31, p = 5.07e-5
PC3 Score: t = 0.807, df = 101.9, p = 0.421
PC4 Score: t = 2.31, df = 84.45, p = 0.023
PC5 Score: t = 0.390, df = 93.65, p = 0.698

Triplot of PC Scores for Patients on No Medication vs Antipsychotic Medication

C *View 1*
D *View 2*
E *View 3*

Fig. S7. PCA solution is not driven by medication status or dosage. Antipsychotic medication dosages were available for N=338 out of 436 PSD patients, including 59 patients not on antipsychotic medication. Antipsychotic dosages were converted to chlorpromazine (CPZ) equivalents ([129]). **(A)** Spearman's ρ between medication dosage (CPZ equivalents) and PC scores for each of the 5 significant principal components (PCs), for medicated patients (grey points). Patients not on antipsychotic medication are also shown (yellow points, CPZ=0mg), however they were not included in the calculation of Spearman's ρ as they contain no rank information. **(B)** Bar plots show the mean PC scores of unmedicated (yellow) versus medicated (grey) PSD patients for each of the 5 PCs. Error bars show standard error of the mean. Note that only PC2 "Cognitive Performance" scores appear to show a significant relationship with medication consistently; this could be because antipsychotic medication (particularly first-generation antipsychotics) are related to symptom variance across some PCA-derived symptom dimensions but do not effectively treat cognitive deficits. Also, cognitive deficits (such as reaction time and fluency) may be exacerbated due to the neuroleptic effects of first-generation antipsychotics. **(C–E)** Alternative views of the triplot (as seen in **Fig. S3A–C**) showing the relationship between the three principal axes of variation in symptoms and aggregate scores from the PANSS and BACS symptom scales (i.e. vectors show the projections of standard symptoms factors). Each point represents an individual patient projected into the geometry defined by the first 3 principal components (PC). Each patient is colored according to medication status (yellow=unmedicated, grey=with medication). As evident from this plot there is no apparent clustering of patients according to their medication status in the 3D PCA-derived space.

Fig. S8. Effects of age, socio-economic status, and sex on symptom PCA solution. (A) Correlations between symptom PC scores and age (years) across N=436 PSD. Pearson's correlation value and uncorrected p-values are reported above scatterplots. After Bonferroni correction, we observed a significant positive relationship between age and *PC3* score. This may be because older patients have been ill for a longer period of time and exhibit more severe symptoms along the positive *PC3* dimension. **(B)** Correlations between symptom PC scores and socio-economic status (SES) as measured by the Hollingshead Index of Social Position (130), across N=387 PSD with available data. The index is computed as (Hollingshead occupation score * 7) + (Hollingshead education score * 4); a higher score indicates lower SES (131). We observed a significant negative relationship between Hollingshead index and *PC1* and PC2 scores. Lower *PC1* and *PC2* scores indicate poorer general functioning and cognitive performance respectively, which is consistent with higher Hollingshead indices (i.e. lower-skilled jobs or unemployment and fewer years of education). **(C)** The Hollingshead index can be split into five classes, with 1 being the highest and 5 being the lowest SES class (130). Consistent with **(B)** we found a significant difference between the classes after Bonferroni correction for *PC1* and *PC2* scores. **(D)** Distributions of PC scores across Hollingshead SES classes show the overlap in scores. White lines indicate the mean score in each class. **(E)** Differences in PC scores between (M)ale and (F)emale PSD subjects. We found a significant difference between sexes in *PC2 – Cognitive Functioning*, *PC4 – Affective Valence*, and *PC5 – Agitation/Excitement* scores. **(F)** Distributions of PC scores across M and F subjects show the overlap in scores. White lines indicate the mean score for each sex.

Fig. S9. Similarity across *a priori*, categorical and PCA-derived brain-behavioral GBC maps. Z-scored maps of t-test for the difference in group mean GBC between traditional diagnostic groups: **(A)** all patients (PSD) versus all healthy controls (CON); **(B)** patients with bipolar disorder (BPP) versus CON; **(C)** patients with schizophrenia (SZP) versus CON; **(D)** patients with schizoaffective disorder (SADP) versus CON. **(E)** Z-scored map of the F-test for the difference in group mean GBC between patients in all 3 diagnostic groups (BPP, SZP, SADP). Z-scored map of the regression against GBC, across all patients (PSD), of traditional symptom/behavioral scales: **(F)** BACS cognitive composite performance score; **(G)** PANSS total negative symptom score; **(H)** PANSS total positive symptom score; **(I)** PANSS total general symptom score. Z-scored map of the regression against GBC, across all patients, of data-derived behavioral dimension scores: **(J)** *PC1* score; **(K)** PC2 score; **(L)** *PC3* score; **(M)** *PC4* score; **(N)** *PC5* score. **(O)** Correlation matrices showing the similarity between brain-behavioral maps in **A-N**. **(P)** Violin plots of the distribution of Z-values in all phenotype maps. Note that although there are strong correlations between the PC maps and the *a priori* symptom maps, the statistical properties of some of the PC maps are improved (i.e. the range of Z-scores is greater, with more extreme values), suggesting a stronger mapping between neural and behavioral variation.

Mapping GBC to PC Scores within PSD (N=436)

Fig. S10. Parcellated symptom-neural GBC maps across all PSD patients derived from PCA dimensionality reduction of symptom measures. (A) All of the maps shown here were parcellated at the single patient level using the Cole-Anticevic Brain Network Parcellation (CAB-NP) parcellation (51), which defines functional networks and regions across cortex and subcortex that leveraged the Human Connectome Project's Multi-Modal Parcellation (MMP1.0) (51, 52). The final published CAB-NP 1.0 parcellation solution can be visualized via the Brain Analysis Library of Spatial maps and Atlases (BALSA) resource (https://balsa.wustl.edu/rrg5v) and downloaded from the public repository (https://github.com/ColeLab/ColeAnticevicNetPartition). **(B–F)** Relationships across all patients (N=436) at each parcel location between global brain connectivity (GBC) and PC score for each of the significant PCs. Values shown in each brain parcel is the Z-scored regression coefficient of PC score on to parcel GBC, across all 436 subjects.

Fig. S11. Parcellated symptom-neural GBC maps across all CON subjects derived from PCA dimensionality reduction of PSD symptom measures. (A) As with PSD maps, all of the maps shown here were parcellated at the single patient level using the Cole-Anticevic Brain Network Parcellation (CAB-NP) parcellation (51), which defines functional networks and regions across cortex and subcortex that leveraged the Human Connectome Project's Multi-Modal Parcellation (MMP1.0) (51, 52). The final published CAB-NP 1.0 parcellation solution can be visualized via the Brain Analysis Library of Spatial maps and Atlases (BALSA) resource (https://balsa.wustl.edu/rrg5v) and downloaded from the public repository (https://github.com/ColeLab/ColeAnticevicNetPartition). **(B–F)** Relationships across all controls (N=202) at each parcel location between global brain connectivity (GBC) and PC score for each of the significant PCs. Values shown in each brain parcel is the Z-scored regression coefficient of PC score on to parcel GBC, across all 202 control subjects.

Fig. S12. Individual patients exhibit complex projections into the PCA-derived brain-behavioral space (BBS) geometry (A) The *PC3* group-level cortical map is shown here for comparison purposes. **(B)** Neural and behavioral profile for one PSD patients, "Patient P1". Neural data shows the △GBC for Patient P1 (i.e. GBC demeaned relative to group mean of the entire PSD patient sample) to reflect the pattern of changes in their GBC relative to the 'average' of the entire sample. The bar plot shows five behavioral PC scores for Patient P1. The radarplot shows original scores on the 36 individual symptom measures for Patient P1 (black circle indicates zero; positive values are indicated by outward deviation of the grey line). **(C–D)** Data is shown for two other individual Patients P2 & P3. **(E)**. Data is shown for one Control C1 participant. Note the difference between the symptom-neural profiles of Patient P1, a positive *PC3* 'loading' individual, and Patient P2, a negative *PC3* 'loading' individual, both of whom were diagnosed with SZP under a conventional categorical DSM approach. These data further illustrate the need to consider the complex relationships in the symptom-neural mapping and highlight that individual-level precision in BBS mapping can be obtained.

Fig. S13. Independent component analysis (ICA) as an alternative method of dimensionality-reduction for symptom-neural mapping. (A–E) Relationships across all patients (N=436) at each brain location between global brain connectivity (GBC) and IC score, for ICs 1-5. Values shown in each brain parcel is the Z-scored regression coefficient of IC score on to parcel GBC, across all 436 subjects. **(F) Scatterplots showing the relationships across parcels between the PC3 map and the IC2 and IC3 maps. Sigmoids and a greater range (max – min) indicate an improvement in the Z-statistics of the PC3 map relative to the IC maps. (G)** Correlation matrix showing correlations of individual parcel regression coefficients for the 5 significant PCs and the five ICs (shown in **A–E**), across all 718 neural parcels. Of note, *PC3 "Psychosis Configuration"* discovered via the PCA solution appears to be oblique to both *IC2* and *IC3*, suggesting that these two *ICs* may capture diverging tails of the *PC3* axis; however, the orthogonality of the PCA solution yielded maximally separated symptom dimensions that mapped on to unique neural variance, as shown by the superior statistics of the PC3 map versus the *IC2/IC3* maps.

Fig. S14. Canonical correlation analysis (CCA) of behavioral and subcortical neural features. (A) As noted in the main text, CCA maximizes correlations between canonical variates (CVs), i.e. matrices **U** and **V**. Here we evaluated a version of CCA using subcortical neural parcel features in relation to item-level symptom measures or dimensionality-reduced PC symptom measures. **(B)** Screeplot showing canonical modes for the CVs obtained from 192 subcortical neural features (GBC of parcels from a neurobiologically-derived functional parcellation of the subcortex) and 36 single-item PANSS and BACS symptom measures. Inset illustrates the correlation (r=0.85) between the CV of the first mode, **U_1** and **V_1** (note that the correlation was not driven by a separation between diagnoses). **(C)** CCA was obtained from 192 subcortical neural features and 5 low-dimensional symptom scores derived via the PCA analysis. Here all modes remained significant after FDR correction. Dashed black line shows the null calculated via a permutation test with 5,000 shuffles; grey bars show 95% confidence interval. **(D)** Correlation between the neural data matrix **N** and the behavioral data matrix weighted by the transformation matrix (**B**Ψ) reflects the amount of variance in **N** that can be explained by the final latent neural matrix **U**. Put differently, this transformation calculates how much of the neural variation can be explained by the latent behavioral features. **(E)** Proportion of symptom variance explained by each of the neural CVs in a CCA performed between 192 subcortical neural features and all 36 behavioral measures. Inset shows the total proportion of behavioral variance explained by the neural variates. **(F)** Proportion of total behavioral variance explained by each of the neural CVs in a CCA performed between 192 subcortical neural features and the 5 low-dimensional symptom scores derived via the PCA analysis. Dashed black line shows the null calculated via a permutation test with 5,000 shuffles; grey bars show 95% confidence interval. **(G)** Correlation between the behavioral data matrix **B** and the neural data matrix weighted by the transformation matrix (**N**Θ) reflects the amount of variance in **B** that can be explained by the final latent neural matrix **V**. Put differently, this transformation calculates how much of the symptom variation can be explained by the latent neural features. **(H)** Proportion of symptom variance explained by each of the neural CVs in a CCA performed between 192 subcortical neural features and all 36 behavioral measures. Inset shows the total proportion of behavioral variance explained by the neural variates. **(I)** Proportion of total behavioral variance explained by each of the neural CVs in a CCA performed between 192 subcortical neural features and the 5 low-dimensional symptom scores derived via the PCA analysis. While CCA using symptom PCs has fewer dimensions and thus lower total variance explained (see inset), each neural variate explains a higher amount of symptom variance than seen in **F**, suggesting that CCA could be further optimized by first obtained a principled low-rank symptom solution. Dashed black line shows the null calculated via a permutation test with 5,000 shuffles; grey bars show 95% confidence interval.

Fig. S15. Canonical correlation analysis (CCA) of behavioral and network-level neural features. (A) As noted in the main text, CCA maximizes correlations between canonical variates (CVs), i.e. matrices **U** and **V**. Here we evaluated a version of CCA using network-level neural parcel features in relation to item-level symptom measures or dimensionality-reduced PC symptom measures. **(B)** Screeplot showing canonical modes for the CVs obtained from 192 subcortical neural features (GBC of parcels from a neurobiologically-derived functional parcellation of the subcortex) and 36 single-item PANSS and BACS symptom measures. Inset illustrates the correlation (r=0.85) between the CV of the first mode, U_1 and V_1 (note that the correlation was not driven by a separation between diagnoses). **(C)** CCA was obtained from 192 subcortical neural features and 5 low-dimensional symptom scores derived via the PCA analysis. Here all modes remained significant after FDR correction. Dashed black line shows the null calculated via a permutation test with 5,000 shuffles; grey bars show 95% confidence interval. **(B)** Screeplot showing the correlations between the canonical variates of a CCA performed between 12 neural features (mean GBC of 12 whole-brain networks) and all 36 behavioral measures. Dashed black line shows the null calculated via a permutation test with 5,000 shuffles; grey bars show 95% confidence interval. Inset illustrates the correlation (r=0.48) between the canonical variates of the first mode, U_1 and V_1. Note that the correlation is not driven by a separation between categorical diagnoses). **(C)** Screeplot showing the correlations between the canonical variates of a CCA performed between 12 network neural features and the 5 behavioral principal components. Dashed black line shows the null calculated via a permutation test with 5,000 shuffles; grey bars show 95% confidence interval. Inset illustrates the correlation (r=0.37) between the canonical variates of the first mode, V_1 and V_1. Note that the strength of the correlations is greatly reduced compared to the parcel-level CCA shown in **Fig. 5. (D)** The correlation between the neural data N and the transformed behavioral data matrix **B**Ψ reflects the amount of variance in **N** that can be explained by behavioral canonical variates **V**. **(E)** Screeplot showing the proportion of neural variance explained by each of the behavioral canonical variates in a CCA performed between 12 network neural features and all 36 behavioral measures. Dashed black line shows the null calculated via a permutation test with 5,000 shuffles; grey bars show 95% confidence interval. Inset shows the total proportion of neural variance explained by the behavioral variates. **(F)** Screeplot of the proportion of neural variance explained by each of the behavioral canonical variates in a CCA performed between 12 network neural features and the 5 PCs of behavior. **(G)** The correlation between the behavioral data **B** and the transformed neural data matrix **N**Θ reflects the amount of variance in **B** that can be explained by neural canonical variates **U**. **(H)** Screeplot showing the proportion of behavioral variance explained by each of the neural canonical variates in a CCA performed between 12 network neural features and all 36 behavioral measures. Dashed black line shows the null calculated via a permutation test with 5,000 shuffles; grey bars show 95% confidence interval. Inset shows the total proportion of behavioral variance explained by the neural variates.t **(I)** Screeplot of the proportion of total behavioral variance explained by each of the neural canonical variates in a CCA performed between 12 network neural features and the 5 PCs of behavior. Note that although the CCA using behavioral PCs has far fewer dimensions, each neural variate explains a higher amount of total behavioral variance than neural variates in **H**, suggesting that the identified PCs of behavior capture variance with far fewer features.

Fig. S16. Canonical correlation analysis (CCA) results showing the amount of variance explained in the neural features via behavioral canonical variates. **(A)** The correlation between the neural data N and the transformed behavioral data matrix **B**Ψ reflects the amount of variance in **N** that can be explained by behavioral canonical variates. **(B)** Screeplot showing the proportion of neural variance explained by each of the behavioral canonical variates in a CCA performed between 180 neural features (symmetrized cortical parcel GBC) and all 36 behavioral measures. Dashed black line shows the null calculated via a permutation test with 5,000 shuffles; grey bars show 95% confidence interval. Inset shows the total proportion of neural variance explained by the behavioral variates. **(C)** Screeplot of the proportion of neural variance explained by each of the behavioral canonical variates in a CCA performed between 180 neural features (symmetrized cortical parcel GBC) and the 5 PCs of behavior.

Fig. S17. Canonical correlation analysis (CCA) symptom and neural configurations. (A) Distributions of scores for the first canonical variate (*CV1*) by diagnostic group (white: controls; black: all patients; yellow: bipolar disorder with psychosis; orange: schizoaffective disorder with psychosis; red: schizophrenia). All scores are normalized to controls. **(B)** Loadings of the PC-derived symptom scores for *CV1*. **(C)** Loadings of the original item-level symptom measures for *CV1*. **(D)** Loadings of each neural parcel for *CV1*. **(E–T)** The same analyses are shown for canonical variates 2-5. As noted in the main text, these CCA effects were not stable when tested via out-of-sample cross-validation methods (see **Fig. 5**).

Fig. S18. K-Fold cross-validation of canonical correlation analysis (CCA). In the main text we show that CCA effects do not reproduce when tested via out-of-sample cross-validation methods (see **Fig. 5**). In contrast, here we show that using within-sample K-fold methods yields effects that appear reproducible but in fact reflect an overfit CCA solution. Panels **(A-E)** illustrate the results from a 5-fold cross-validation analysis to test the reproducibility of the CCA solution. Subjects were first randomly assigned to one of 5 subsets. Each subset of subjects was then used as an independent 'test sample' in a CCA that was derived from subjects in the other 4 subsets. **(A)** Results of CCA performed in a subsample of N=349 subjects. Left: screeplot showing the correlations between the canonical variates of a CCA performed between 180 neural features (symmetrized cortical parcel GBC) and the 5 behavioral principal components in N=349 subjects. Middle: Screeplot of the proportion of neural variance explained by each of the behavioral canonical variates in a CCA performed between 180 neural features (symmetrized cortical parcel GBC) and the 5 PCs of behavior. Right: Screeplot of the proportion of total behavioral variance explained by each of the neural canonical variates in a CCA performed between 180 neural features (symmetrized cortical parcel GBC) and the 5 PCs of behavior. Dashed black line shows the null calculated via a permutation test with 5,000 shuffles; grey bars show 95% confidence interval. Note that the outcome of this CCA is highly similar to the full sample CCA shown in the main text, but it is not similar at all to the split-half cross-validation effects, which reveal how overfit CCA effects seem to be with this number of features **Fig. 5**. **(B)** Comparison of the first canonical variate (CV1) in a CCA performed in a subsample of N=349 subjects with *CV1* from the full model performed with N=436 subjects. Left: neural factor loadings for *CV1* obtained from the subsample (Fold 1) CCA and from the full sample CCA are highly correlated, at r=0.85. Middle: Behavioral factor loadings are also highly correlated between the Fold 1 CCA and the full CCA, at r=0.85. Right: Additionally, absolute values of individual symptom measure loadings associated with *CV1* are highly correlated between the Fold 1 CCA and the full CCA. **(C)** Summary of correlation values between Fold 1 CCA and full CCA results, as in panel **B**, across all 5 CVs. Note that the neural and behavioral factor loadings as well as individual symptom measure loadings are highly preserved between the sample CCA and the full model. **(D)** Summary of correlation values between all 5 subsample CCAs and the full sample CCA, for all 5 CVs. Each of the 5 CVs is plotted along the X-axis; each point represents the correlation between one of the 5-fold subsample CCAs and the full CCA, hence there are 5 points (in the 5-fold cross-validation) for each CV. **(E)** Results of a CCA performed in a subsample of N=357 subjects, with subjects from one site left out (leave-one-site-out, LOSO). Panels as described in **A**. **(F)** Comparison of the first canonical variate (CV1) in a LOSO CCA and the *CV1* from the full model performed with N=436 subjects. Panels as described in **B**. **(G)** Summary of correlation values between the LOSO CCA and full CCA results, as in panel **B**, across all 5 CVs. Panels as described in **C**. **(H)** Summary of correlation values between all 6 LOSO CCAs and the full sample CCA, for all 5 CVs. Panels as described in **D**; each point represents the correlation between one of the 6 LOSO CCAs and the full CCA.

Fig. S19. Multivariate power analysis for CCA. Sample sizes were calculated according to ([27](#)), see also **Methods** and https://gemmr.readthedocs.io/en/latest/. We computed the multivariate power analyses for three versions of CCA reported in this manuscript: i) 718 neural vs. 5 symptom features; ii) 180 neural vs. 5 symptom features; iii) 12 neural vs. 5 symptom features. **(A)** At different levels of features, the ratio of samples (i.e. subjects) required per feature to derive a stable CCA solution remains approximately the same across all values of r_{true}. As discussed in ([27](#)), at $r_{true} = 0.3$ the number of samples required per feature is about 40, which is much greater than the ratio of samples to features available in our dataset. **(B)** The total number of samples required (n_{req}) for a stable CCA solution given the total number of neural and symptom features used in our analyses, at different values of r_{true}. In general these required sample sizes are much greater than the N=436 (horizontal grey line) PSD in our present dataset, consistent with the finding that the CCA solutions computed using our data were unstable. Notably, the '12 vs. 5' CCA assuming $r_{true} = 0.3$ requires only ~700 subjects, which is closest to the N=436 used in the present sample. This may be in line with the observation of the CCA with 12 neural vs 5 symptom features (**Fig. S15C**) that the canonical correlation ($r = 0.38$ for CV1) clearly exceeds the 95% confidence interval, and may be closer to the true effect. However, to to confidently detect effects in such an analysis (particularly if r_{true} is actually less than 0.3), a larger sample would likely still be needed.

Fig. S20. Principal component analysis (PCA) of neural features in control and patient subjects. (A) Results of PCA performed on neural features (718 whole-brain parcel GBC) across all control subjects (N=202). **(B)** Results of PCA performed on neural features (718 whole-brain parcel GBC) across all patient subjects (N=436). **(C)** Note that the first neural PC in control subjects (panel A) and in patient subjects (panel **B**) are highly similar (*r*=0.90 across parcels). This may reflect a component of neural variance common across all humans. **(D–I)** The second and third PCs are also similar between control and patient subjects. **(J–L)** The fourth PC is markedly dissimilar between control and patient subjects, suggesting that this component may reflect neural variance which deviates in individuals with psychiatric symptoms. **(M–O)** The fifth PC is also highly dissimilar between controls and patients, also possibly reflecting diagnosis-relevant differences in neural variance. **(P)** Screeplot showing the total proportion of variance explained by the first 100 PCs from the PCA performed across all 718 neural features in 202 control subjects. The size of each point is proportional to the variance explained by that PC. The first 8 PCs (green) were determined to be significant using a permutation test. Inset shows the proportion of variance both accounted and not accounted for by the 8 significant PCs. Together, these 8 PCs capture 51.95% of the total variance in neural GBC in the sample. **(Q)** Screeplot showing the total proportion of variance explained by the first 100 PCs from the PCA performed across all 718 neural features in 436 patient subjects. The size of each point is proportional to the variance explained by that PC. The first 15 PCs (green) were determined to be significant using a permutation test. Inset shows the proportion of variance both accounted and not accounted for by the 15 significant PCs. Together, these five PCs capture 57.4% of the total variance in neural GBC in the sample. **(R)** Correlations between the first 15 neural PCs in controls (CON) and patients (PSD). The first 3 PCs are common across both control and patient subjects, possibly reflecting common components of human neural variance. After this, the PCs diverge, possibly reflecting differences in neural variance of healthy individuals and those with psychiatric symptoms.

Mapping Symptom PC Scores to GBC and Partial GBC Across PSD (N=436)

Fig. S21. Comparison between the PSD $\beta_{PC}GBC$ maps computed using GBC and GBC with the first neural PC parsed out. If a substantial proportion of neural variance is not be clinically relevant, then removing the shared neural variance between PSD and CON should not drastically affect the reported symptom-neural univariate mapping solution, because this common variance will not map to clinical features. We therefore performed a PCA on CON and PSD GBC to compute the shared neural variance (see **Methods**), and then parsed out the first GBC-PC from the PSD GBC data ($G\hat{B}C_{woPC1}$). We then reran the univariate regression as described in **Fig. 3**, using the same five symptom PC scores across 436 PSD. **(A)** The $\beta_{PC1}GBC$ map, also shown in **Fig. S10**. **(B)** The first GBC-PC accounted for about 15.8% of the total GBC variance across CON and PSD. Removing GBC-PC1 from PSD data attenuated the $\beta_{PC1}GBC$ statistics slightly (not unexpected as the variance was by definition reduced) but otherwise did not strongly affect the univariate mapping solution. **(C)** Correlation across 718 parcels between the two $\beta_{PC1}GBC$ map shown in **A** and **B**. **(D-O)** The same results are shown for $\beta_{PC2}GBC$ to $\beta_{PC5}GBC$ maps.

Mapping Symptom PC Scores to GBC and Partial GBC Across PSD (N=436)

Fig. S22. Comparison between the PSD $\beta_{PC}GBC$ maps computed using GBC and GBC with the first two neural PCs parsed out. We performed a PCA on CON and PSD GBC and then parsed out the first three GBC-PC from the PSD GBC data ($G\hat{B}C_{woPC1-2}$, see **Methods**). We then reran the univariate regression as described in **Fig. 3**, using the same five symptom PC scores across 436 PSD. **(A)** The $\beta_{PC1}GBC$ map, also shown in **Fig. S10**. **(B)** The second GBC-PC accounted for about 9.5% of the total GBC variance across CON and PSD. **(C)** Correlation across 718 parcels between the two $\beta_{PC1}GBC$ map shown in **A** and **B**. **(D-O)** The same results are shown for $\beta_{PC2}GBC$ to $\beta_{PC5}GBC$ maps.

Mapping Symptom PC Scores to GBC and Partial GBC Across PSD (N=436)

Fig. S23. Comparison between the PSD $\beta_{PC}GBC$ maps computed using GBC and GBC with the first three neural PCs parsed out. We performed a PCA on CON and PSD GBC and then parsed out the first three GBC-PC from the PSD GBC data ($\hat{GBC}_{woPC1-3}$, see **Methods**). We then reran the univariate regression as described in **Fig. 3**, using the same five symptom PC scores across 436 PSD. **(A)** The $\beta_{PC1}GBC$ map, also shown in **Fig. S10**. **(B)** The second GBC-PC accounted for about 9.5% of the total GBC variance across CON and PSD. **(C)** Correlation across 718 parcels between the two $\beta_{PC1}GBC$ map shown in **A** and **B**. **(D-O)** The same results are shown for $\beta_{PC2}GBC$ to $\beta_{PC5}GBC$ maps.

Mapping Symptom PC Scores to GBC and Partial GBC Across PSD (N=436)

Fig. S24. Comparison between the PSD $\beta_{PC}GBC$ maps computed using GBC and GBC with the first four neural PCs parsed out. We performed a PCA on CON and PSD GBC and then parsed out the first four GBC-PC from the PSD GBC data ($G\hat{B}C_{woPC1-4}$, see **Methods**). We then reran the univariate regression as described in **Fig. 3**, using the same five symptom PC scores across 436 PSD. **(A)** The $\beta_{PC1}GBC$ map, also shown in **Fig. S10**. **(B)** The second GBC-PC accounted for about 9.5% of the total GBC variance across CON and PSD. **(C)** Correlation across 718 parcels between the two $\beta_{PC1}GBC$ map shown in **A** and **B**. **(D-O)** The same results are shown for $\beta_{PC2}GBC$ to $\beta_{PC5}GBC$ maps.

Fig. S25. Workflow for optimizing the neural feature selection for a predictive symptom-neural subject specific model. We optimized the selection of a subset of neural parcels from the group βGBC_N^{obs} map (where N is the total number of subjects) that could be predicted via a single symptom score across subjects (e.g. for the *PC3* beta map and *PC3* symptom scores). (**START**) We start with P_{select} set as the full parcellated map,$P_{select} = \beta GBC_N^{obs}$. Hence initially there are 718 elements in the vector P_{select}, each representing the value of a parcel ($|P_{select}| = 718$). Next, since $|P_{select}| > 1$, the selection process enters the **A. Feature Selection Loop** (in red). Selects subset of parcels P_{select} (**A.1**) for which the predictive model will be evaluated. This loop iteratively removes parcels from the P_{select} vector (**A.2**) and each time passes the updated P_{select} as input in the Feature Evaluation Loop (in blue). **B. Feature Evaluation Loop** (shown in blue). Using a multi-step algorithm, this loop produces the set of metrics to be evaluated at the final output using a leave-one-subject out approach. **B.1**: For each subject i, this first step is to compute the "observed" dot product metric $dpGBC_i^{obs}$ for that subject. This is the dot product between ΔGBC_i^{obs} and βGBC_{N-i}^{obs} using the P_{select} features. ΔGBC_i^{obs} is the vector of the parcel-wise difference between subject i's GBC map and the group mean GBC map; βGBC_{N-i}^{obs} denotes group-level relationship between symptom variation and GBC excluding subject i. This group βGBC_{N-i}^{obs} is used as a reference for the dot-product. **B.2**: Next for subject i we compute their predicted symptom score S_i^{pred}. First, a symptom-prediction model is built using X_{N-i}^{obs} (all item-level symptom measures excluding subject i). This yields the "observed" model scores and coefficients excluding subject i (S_{N-i}^{obs}). The symptoms for excluded subject i are then used to obtain the predicted symptom score S_i^{pred}, as done in **Fig. 6A**. **B.3**: Lastly, for excluded subject i the loop computes the "predicted" dot product metric $dpGBC_i^{pred}$ to compare it to the "observed" value for model evaluation. Here, a regression model is computed relating $dpGBC_{N-i}^{obs}$ to S_{N-i}^{obs}, excluding subject i. This yields regression coefficients α, which are then used to predict the $dpGBC_i^{pred}$ for the excluded subject i using their predicted symptom scores S_i^{pred}. Once $i > N$ (i.e. $dpGBC_i^{pred}$ and S_i^{pred} have been computed for all subjects), the algorithm moves to step "C". **C. Feature Evaluation Metrics** (shown in purple). **C.1**: This step captures feature evaluation metrics across N subjects for the model that was computed using the selected features (P_{select}). Specifically, three vectors (outputs from "B") are computed: $dpGBC_N^{obs}$ (the vector of observed $dpGBC$ values for all N subjects); $dpGBC_N^{pred}$ (the vector of predicted $dpGBC$ values for all N subjects); and S_N^{pred} (the vector of predicted model symptom scores for all N subjects). **C.2**: Generates two feature evaluation metrics across the entire sample ($r(dpGBC_N^{obs}, S_N^{pred})$ and $r(dpGBC_N^{obs}, dpGBC_N^{pred})$). **C.3**: The feature evaluation metrics for the selected P_{select} are generated, such that they can be evaluated across all P_{select} iterations (red loop). **D. Final Feature Selection** (in green). This step evaluates the metrics computed for all P_{select} iterations, starting from from $|P_{select}| = 718$ to $|P_{select}| = 1$. **D.1**: The input is the vector of $r(dpGBC_N^{obs}, S_N^{pred})$ and the vector of $r(dpGBC_N^{obs}, S_N^{pred})$ for all 718 models. **D.2**: The most predictive feature subset is obtained by finding the model for which $r(dpGBC_N^{obs}, S_N^{pred})$ and $r(dpGBC_N^{obs}, dpGBC_N^{pred})$ were maximized, which leads to the final output features (**STOP**). *Note*: that the algorithm imposed a threshold of $|P_{select}| > 30$ to ensure that the correlation metrics were not unstable based on a small feature set.

Fig. S26. Optimizing neural feature selection to inform statistically reliable single-subject prediction via low-dimensional symptom scores only. (A) Individual PC-derived symptom scores for *PC1* can be robustly predicted, as shown by leave-one-out cross-validation for the symptom PCA analyses. Here for the leave-one-out cross-validation, a PCA was computed in all except one subject (N=435) and the PCA loadings were then used to compute the predicted PC scores for the left-out subject. This was repeated such that predicted scores were calculated for every subject, using loadings from a PCA computed in the other 435 subjects. Observed scores were obtained from a PCA computed using the full sample of subjects (N=436). Top panel: Scatterplot at left shows the correlation between each subject's predicted *PC1* score from leave-one-out PCA model and their observed *PC1* score from the full-sample PCA model, *r*=0.99. Red line shows perfect prediction, *r*=1. **(B)** Top panel indicates that PC symptom scores can yield robust neural feature selection. ΔGBC denotes an individual subject GBC demeaned parcel-wise relative to the group mean GBC. The process is described in detail in the **Methods** and in **Fig. S25**. The correlation between *PC1* behavioral score and [ΔGBC · *PC1* beta] across all 436 subjects was maximal for $P = 79$, at *r*=0.27 (purple arrow). Bottom panel: For each value of *p*, the regression was performed in 435 subjects and was then used to predict the [ΔGBC · *PC1* beta] of the selected parcels in the left-out subject. The mean correlation between the predicted [ΔGBC · *PC1* beta] and the actual observed [ΔGBC · *PC1* beta] across all 436 subjects was highest for $P = 79$, with *r*=0.19 across all subjects, highly consistent with the across-subject analysis in the top panel. **(C)** Comparing the correlation between ΔGBC_{Obs}, ΔGBC_{Pred} for the full map ($P = 718$) versus the selected map ($P = 79$) shows that the predictability of Δ GBC is improved by using the selected parcels in the selected map.s **(D–O)** Similar results for PCs 2-5. Note that for *PC4* and *PC5* a local maximum ($P = 50$ and $P = 31$ respectively) was selected after imposing a threshold of $P > 30$ to ensure that the correlation metrics used for evaluation were not made unstable from a small number of parcels in computing the $dpGBC$.

Fig. S27. Evaluating patient-specific similarity to derived symptom-neural targets via data-reduced symptom scores. (A) Scatterplot of *PC3* symptom score (X-axis) versus *PC3* neural similarity prediction index (NSPI, Y-axis) for all 436 PSD subjects. The NSPI is defined as the Spearman's correlation between ΔGBC^{obs} and the $\beta_{PC3}GBC^{obs}$ map of the maximally-predictable "selected" $P = 39$ parcels. Alternative metrics are shown in **Fig. S28. (B)** Bins across axes express subject counts within each cell as a heatmap, indicating a high similarity between symptom PC score and *PC3* NSPI for a number of patients. **(C)** Mean NSPI is computed for a given bin along the the X-axis to visualize patient clustering. Note the sigmoidal shape of the distribution reflecting greater neural similarity at more extreme values of the *PC3* score. **(D)** The absolute value of the mean NSPI reflects the magnitude irrespective of neural similarity direction. This highlights a quadratic effect, showing that patients with higher *PC3* symptom scores (either positive/negative) exhibited higher neural correspondence of their maps with the target neural reference map. **(E)** Using the NSPI and PC scores we demonstrate one possible brain-behavioral patient selection strategy. We first imposed a PC score symptom threshold to select patients at the extreme tails (i.e. outside of the $10th - 90th\%tile$ behavioral range [>+2.17 or <-2.41]). Note that this patient selection strategy excludes patients (shown in grey) below the PC symptom score threshold. This yielded n=38 patients. Next, for each patient we predicted the sign of their individual NSPI based on their individual PC3 score, which served as the basis for the neural selection. Next, at each NSPI threshold we evaluated the proportion of patients correctly selected until there were no inaccurately selected patients in at least one PC3 direction (green line or higher). The number of accurately (A) vs. inaccurately (I) selected patients within each bin is shown in red and blue respectively. Note that as the neural ρ threshold increases the A/I ratio increases. **(F)** The neural and behavioral thresholds defined in the "discovery" sample were applied to an independent "replication" dataset (N=69, see **Methods**), yielding a similar final proportion of accurately selected patients. **(G)** The same brain-behavioral patient selection strategy was repeated for *PC5* in the discovery sample (thresholds of $10^{th}\%tile$=-1.89 and $90^{th}\%tile$= +1.47; NSPI threshold of ρ=0.4 optimized for PC5). Results yielded similar A/I ratios as found for PC3. **(H)** The neural and behavioral thresholds for PC5 defined in the discovery sample were applied to the replication sample. Here the results failed to generalize due true clinical differences between the discovery and replication samples.

Fig. S28. Metrics quantifying the relationship between individual-level neural effects relative to the group mean and individual-level PC-derived symptom scores. The relationship between various symptom-neural similarity metrics and the behavioral *PC3* symptom scores across N=436 PSD patients are shown. The goal was to evaluate metrics that maximize the spread of the Y-axis, while maintaining a centered distribution of scores around zero to yield an interpretable symptom-neural variation pattern across subjects. Specifically, along the Y-axis we examined either the Pearson's correlation (top panels in A and B) or Spearman's ρ (bottom panels in A and B) between two neural maps at the individual patient level: i) the group-level $\beta_{PC3}GBC$ map and ii) an individual subjects' raw GBC map which can be expressed as either a raw map (left panels in A and B) or as ΔGBC^{obs} relative to the group mean map (right panels in A and B). **(A)** These four panels show group-level symptom-neural relationships for sample using the entire 718-parcel neural map. As noted, the neural similarity metrics on each Y-axis show the relationship between individual-patient neural GBC (or ΔGBC^{obs}) and the group-level $\beta_{PC3}GBC$ map across 718 parcels. Note that the *r* and *rho* values reported in the corner of each plot reflect the *group-level* relationship between the neural similarity metric and the individual-level PC3 symptom score computed across across 436 patients. **(B)** These four parcels show group-level symptom-neural relationships using only the selected 39 *PC3* parcels derived from the feature selection workflow (**Fig. S25**). The metric used in the main text is the ρ within selected parcels (shown with a light purple envelope) because the spread along Y-axis was maximized and the values centered around zero for both the x and the Y-axis. Furthermore, the ρ metric is desirable given violations of normality that the *r* metric assumes. **(C)** The neural-behavioral similarity plot for the Discovery BSNIP sample was binned by $\rho = 0.1$ & $PC3_{score} = 0.5$ to provide a visual intuition for patient segmentation across both the neural and symptom indices. For patients with highly positive or negative scores, the neuro-behavioral relationship was robust. Conversely, patients with a low absolute *PC3* score showed a weak relationship with symptom-relevant neural features. **(D)** The neural-behavioral similarity plot for the independent replication sample (see **Methods** for sample details) was binned by $\rho = 0.1$ & $PC3_{score} = 0.5$, using the same neural similarity index ($\rho[\Delta GBC^{obs}, \beta_{PC3}GBC^{obs}]$ within selected 39 parcels) and projected *PC3* symptom scores. Although the grid is far more sparse (total N=69 for this sample), a similar pattern between PC3 symptom score and neural similarity index emerges.

Fig. S29. Evaluating patient-specific similarity to derived symptom-neural targets via data-reduced symptom scores for all PCs. As noted in the main text for *PC3*, here we plot the *PC* symptom score (X-axis) versus the neural similarity prediction index (NSPI, **Fig. S27**) for all 436 PSD patients. The NSPI is defined as the Spearman's correlation between ΔGBC^{obs} and the $\beta_{PC3}GBC^{obs}$ map of the maximally-predictable "selected" P parcels (Y-axis for each plot). Alternative metrics were evaluated in **Fig. S28**. **(A)** A parameter sweep was performed for $0<|rho|<1$ in bins of 0.1, for all subjects with *PC1* score<-3.97 ("negative behavioral threshold" set at the 10th percentile ($P_{10-Disc}$) of the distribution and for all patients with *PC1* score>+3.60 ("positive behavioral threshold" set at the 90th percentile ($P_{90-Disc}$) of the distribution). The number of patients accurately selected within each bin is shown in red; the number of inaccurately selected patients is shown in blue. Note that this selection process excludes a number of patients (shown in grey) who do not meet the PC threshold cutoff. Note that as the neural ρ threshold increases the proportion of accurate to inaccurate selections increases and the number of inaccurately selected subjects decreases. For illustrative purposes, a neural threshold of $\rho=0.5$ is applied at the point where no inaccurate selections occur for the positive behavioral threshold. **(B)** The neural and symptom score thresholds defined in discovery patients were applied on an independent external "replication" dataset (N=69, see **Methods**). For the symptom score ($P_{10-Disc}$=-3.97 and $P_{90-Disc}$=+3.60) and neural (ρ=0.5) thresholds as those defined in the discovery sample, no patients are selected for *PC1*. **(C–D)** Similar data to **A–B** for *PC2*, using PC score thresholds of $P_{10-Disc}$=-2.43 and $P_{90-Disc}$=+2.32 and a PC map threshold of ρ=0.3 on a subset of $P=68$ optimized parcels. **(E–F)** Similar data to **A–B** for *PC3* using PC score thresholds of $P_{10-Disc}$=-2.41 and $P_{90-Disc}$=+2.17 and a PC map threshold of ρ=0.4 on a patients of $P=39$ optimized parcels. **(G–H)** Similar data to **A–B** for *PC4*, using PC score thresholds of $P_{10-Disc}$=-1.69 and $P_{90-Disc}$=+1.72 and a PC map threshold of ρ=0.2 on a subset of $P=50$ optimized parcels. **(I–J)** Similar data to **A–B** for *PC5*, using PC score thresholds of $P_{10-Disc}$=-1.89 and $P_{90-Disc}$=+1.47 and a PC map threshold of ρ=0.4 on a subset of $P=31$ optimized parcels.

Fig. S30. Examining most predictive parcels for all *PC*-driven symptom-neural effects and their relationship with pharmacological neuroimaging maps. (A) (Left) The selected group-level $\beta_{PC1}GBC$ map showing the 79 selected parcels derived from the feature selection workflow (**Fig. S25**). (Middle) The same parcels are highlighted from the group-level ΔGBC LSD map reflecting the pharmacological effect relative to placebo. (Right) the group-level ketamine ΔGBC map reflecting the pharmacological effect relative to placebo. (Bottom bar plot) Here we show four values: i) Spearman's ρ between the ketamine ΔGBC and $\beta_{PC1}GBC$ maps (KET,PC1); ii) Spearman's ρ between the LSD ΔGBC and $\beta_{PC1}GBC$ maps (LSD,PC1); iii) Absolute value of the difference for the (i) and (ii) (i.e. differential similarity; |(KET,PC1) - (LSD,PC1)|); iv) Spearman's ρ between the ketamine ΔGBC response and LSD ketamine ΔGBC response maps (KET,LSD). These relationships are quantified within both the selected 79 features (dark grey bars) and the full *P*=718 neural map (light grey bars). **(B–E)** These analyses are repeated and shown for *PCs 2-5*. Note that a strong differential similarity for a pair of pharmacological maps may indicate a strategy for patient treatment selection along a particular brain-behavioral axis. Specifically, in this framework patients loading on *PC3* behaviorally (or neurally) would be indicated as highly similar to neural variation induced by ketamine but not LSD. Such metrics may be clinically useful for selecting one pharmacological target over another in the process making individual patient segmentation decisions.

Fig. S31. Quantifying the relationship between *PC*-derived symptom-neural effects and expression maps for genes implicated in psychosis spectrum disorder (PSD) neurobiology. (A) The symptom loading profile (radarplot) and neural loading map derived in the main analyses are shown for the *PC1* BBS. **(B)** The distribution shows the Pearson's correlation coefficients between the *PC1* symptom-neural map and all 20,200 gene expression maps derived from the Allen Human Brain Atlas (AHBA, see **Methods** and for details on gene expression analyses see (33)). In this analysis we focused on a select number of PSD-relevant genes, as highlighted in the main text for PC3 specifically **Fig. 8**. Seven genes of interest are shown. Specifically, these include two interneuron marker genes, somatostatin (SST) and parvalbumin (PVALB); two GABA_A receptor subunit genes: GABRA1 and GABRA3; and three serotonin receptor subunit genes: HTR1E, HTR1F, and HTR2A. The percentile for each highlighted gene relative to the entire distribution of 20,200 genes is reported in square brackets. Note that the gene expression maps of HTR1F and PVALB are at the negative tail of the entire distribution, i.e. anti-correlated with *PC1* symptom-neural map. Conversely, GABRA3, SST, and HTR1E are on the far positive end, reflecting a highly similar spatial pattern of the gene expression maps with the *PC1* symptom-neural map. **(C–J)** *PC* symptom-neural map profiles and distributions of gene expression map similarities are also shown for PCs 2-5. Note that some of these exemplar genes show a strong spatial similarity (or anti-similarity) with the PC-derived neural maps relative to the entire distribution of all gene-PC map pairs.