Phosphorylation of luminal region of the SUN-domain protein Mps3 promotes nuclear envelope localization during meiosis

H. B. D. Prasada Rao[1], Takeshi Sato[2], Kiran Challa[3], Miki Shinohara[4], and Akira Shinohara*

Institute for Protein Research, Osaka University, Suita, Osaka 565-0871, Japan

Running head: Meiotic control of NE remodeling

Keywords: meiosis; CDK; DDK; Mps3; chromosome motion; Nuclear envelope (NE)

[1]Current address: National Institute for Animal Breeding, Hyderabad, India

[2]Current address: Kyoto Pharmaceutical University, Kyoto, Japan

[3]Current address: Friedrich Miescher Institute for Biomedical Research, Basel, Switzerland

[4]Current address: Graduate School of Agriculture, Kindai University, Nara, Japan

*Correspondence:

Akira Shinohara

Institute for Protein Research, Osaka University

3-2 Yamadaoka, Suita, Osaka 565-0871, Japan

Telephone: 81-6-6879-8624

FAX: 81-6-6879-8626

E-mail: ashino@protein.osaka-u.ac.jp

ORCID: 0000-0003-4207-8247

Summary

During meiosis, protein ensembles in the nuclear envelope (NE) containing

SUN- and KASH-domain proteins, called linker nucleocytoskeleton and

cytoskeleton (LINC) complex, promote chromosome motion. How LINC

complexes acquire the meiotic property is largely unknown. Here we

showed that cyclin-dependent protein kinase (CDK) and Dbf4-dependent

Cdc7 protein kinase (DDK) promote proper meiosis-specific localization of

yeast SUN-domain protein Mps3 on NE and control force-dependent

movement of chromosomes during meiosis. We also found a NE luminal

region of Mps3 juxtaposed to inner nuclear membrane (INM) is required for

meiosis-specific localization of Mps3 on NE. Negative charges introduced

by meiosis-specific non-canonical phosphorylation of the luminal region of

Mps3 changes its interaction with INM, which may induce NE localization

by promoting the formation of a canonical LINC complex with Mps3. Our

study reveals unique phosphorylation-dependent regulation on the

localization and function of Mps3 protein in meiotic NE remodeling.

Introduction

During meiosis, homologous chromosomes pair with each other. This pairing leads to juxtaposition of the chromosomes along their entire length, resulting in the formation of the synaptonemal complex (SC). SC is a proteinaceous structure comprising two chromosome axes called the axial/lateral element (AE/LE) flanking a central region with a ladder-like structure (Hawley and Gilliland, 2009; Zickler and Kleckner, 1999). AE formation is initiated in the leptotene stage of meiotic prophase I, accompanied with the assembly of the chromosome axis with multiple chromatin loops. In the zygotene stage, a part of homologous AE pairs each other to form a short SC. In the pachytene stage, SC is elongated along the entire chromosome. After the pachytene stage, SC dismantles in the diplotene stage before the onset of meiosis I.

Meiotic cells also exhibit a unique arrangement of chromosomes inside of the nucleus. In most organisms, telomeres attach to nuclear envelopes (NEs) in early meiotic prophase (or premeiotic phase) and occasionally cluster on one area of NE, particularly near the vicinity of the centrosome (a microtubule organizing center) (Zickler and Kleckner, 1998). Telomere clustering is seen in only the zygotene stage and is known as "telomere/chromosome bouquet." In the pachytene stage, telomere bouquets are resolved and telomeres are dispersed over the entire NE. The bouquet configuration of chromosomes is proposed to promote homologous pairing by restricting the homology search process from three dimension to two dimension (Zickler and Kleckner, 1998).

Recent studies revealed the very dynamic nature of meiotic chromosomes in various organisms from yeasts to mammals (Hiraoka and Dernburg, 2009; Koszul and Kleckner, 2009). The oscillated motion of a nuclei during meiotic prophase I has been well described in fission yeast (Chikashige et al., 2007). The nuclear movement is driven by cytoplasmic microtubule cables, along which SPB embedded in NE moves together with clustered telomeres. This telomere-led movement promotes pairing of homologous loci on chromosomes. Although less coordinated, chromosome motion throughout prophase I has been reported in budding yeast, nematodes, maize and mouse (Penkner et al., 2009; Sato et al., 2009a; Sheehan and Pawlowski, 2009).

Protein ensembles embedded in NE that connect telomeres in
nucleoplasm with the cytoskeleton in cytoplasm, which are referred to as the
LINC (linker of nucleocytoskeleton and cytoskeleton) complex, promote meiotic
chromosome motion (Hiraoka and Dernburg, 2009; Starr and Fridolfsson, 2010).
The LINC complex necessary for chromosome motion contains an inner nuclear
membrane (INM) protein harboring the SUN (Sad1/UNc-84) domain and an outer
nuclear membrane (ONM) protein with the KASH (Klarsicht, ANC-1, SYNE1
Homology) domain. SUN and KASH domains interact with each other in a space
between INM and ONM, luminal region or peri-nuclear space. SUN-domain
proteins bind to telomeres and/or nucleocytoskeleton in nucleoplasm while
KASH-domain proteins bind to cytoskeletons. LINC complexes transmit forces
generated by cytoskeleton to NEs for nuclear positioning and migration in somatic
cells and to telomeres for chromosome motion in meiotic cells (Starr & Fridolfsson
2010).

In *Saccharomyces cerevisiae*, telomeres are tethered to NEs as several
clusters during mitosis and move around NEs during meiotic prophase I. These
telomeres are transiently clustered during prophase I (Conrad et al., 2008; Koszul
et al., 2008; Trelles-Sticken et al., 2000; Trelles-Sticken et al., 1999). Telomere
movement on NE drives chromosome motion inside the nucleus (Koszul et al.,
2008; Scherthan et al., 2007). At least three distinct regulatory steps facilitate
chromosome motion in the budding yeast meiosis. First, meiosis-specific
components of the LINC complex, such as Ndj1 and Csm4, are expressed and
incorporated. Ndj1 is a meiosis-specific telomere-binding protein necessary for
telomere tethering to NEs (Chua and Roeder, 1997; Conrad et al., 1997),
whereas Csm4 is a meiosis-specific ONM protein (Conrad et al., 2007; Kosaka
et al., 2008; Wanat et al., 2008), which interacts with a unique class of KASH
protein, Mps2 (Chen et al., 2019; Lee et al., 2020). Second, the formation of actin
cables is specifically induced in the cytoplasm of meiotic cells (Koszul et al., 2008;
Taxis et al., 2006). Third, NE dynamics including NE remodeling and movement
occur prominently in meiotic prophase I (Conrad *et al.* 2007; this paper). NE
remodeling is accompanied by NE localization of a yeast SUN-domain protein,
Mps3/Nep98 (Conrad et al., 2007) as well as Mps2 (Lee et al., 2020). In mitosis,

Mps3 and Mps2 predominantly localizes to SPB as a component of the half-
bridge as well as SPB-interacting network (SPIN) (Chen et al., 2019) and play an
essential role in SPB duplication (Jaspersen et al., 2002; Nishikawa et al., 2003).
During meiotic prophase-I, in addition to SPB, Mps3 localizes as a distinct protein
ensembles on NE as a LINC complex (Conrad et al., 2007). In early meiotic
prophase I, Mps3 forms several distinct foci/patches on NE and later shows more
disperse staining around NE. Mps3 plays an important role in
telomere/chromosome dynamics (Conrad et al., 2007). However, how meiosis-
specific Mps3 localization on NE is induced is poorly understood.

Here we showed that cyclin-dependent Cdc28 kinase (CDK) and Dbf4-
dependent Cdc7 kinase (DDK) are necessary for proper telomere movement,
particularly for the resolution of telomere clusters. Defects in telomere movement
induced by CDK- and DDK-inactivation are associated with impaired meiosis-
specific NE localization of Mps3; resolution of Mps3 clusters. We also found that
a luminal region of Mps3, which is located near INM, is important for its NE
localization. This region is subject to phosphorylation during meiosis prophase I.
Phospho-defective *mps3* mutant protein showed reduced NE localization during
meiosis. Introduction of negative charged residues in this region suppresses the
defects in Mps3 localization in *cdk* and *ddk* mutants. In reconstituted liposomes,
a Mps3 peptide containing the luminal region with negatively charged residues
showed reduced affinity to the membrane relative to a control wild-type Mps3
peptide. Our results suggested that collaborative action of CDK and DDK and
Mps3 luminal region control Mps3 localization to NE, thus, the assembly of
meiosis-specific canonical LINC complex for chromosome motion and NE
remodeling during meiosis.
142

Results

Force-independent localization of Mps3 on NE during meiosis

We examined the localization of Mps3-GFP (Conrad et al., 2007) in various mutants. Mps3-GFP showed stage-specific distribution and movement on NE (Figure 1A, 1B). During meiotic prophase I, Mps3 shows unique localization patterns on NE. At 0 h, Mps3 predominantly localizes on SPB as a single focus on NE (Figure 1A) (Jaspersen et al., 2002; Nishikawa et al., 2003) with little movement (Figures 1A, 1C). During early prophase I including meiotic S phase (2–3 h), Mps3 formed 2–5 foci (including 1 focus in SPB) on NE with slow movement. After 3 h, the number of Mps3 foci increased to more than 5, and several Mps3 foci fused to form a patch (Figures 1A-1D), as shown previously (Conrad et al., 2007). During this stage, Mps3 foci/patches moved around NE with oscillated motion (Figure 1C). At 3-5 h, some Mps3 foci/patches localized to one area of NE with clustering. In late prophase I (e.g., at 5 and 6 h), Mps3 often covered most parts of NE and was associated with NE deformation and protrusion (Figure 1). These localizations were classified in four classes and quantified at each time point (Figure 1B). After 6 h, most Mps3 signals on NE disappeared, leaving two SPB-associate foci (Figure 1A).

Although Mps3 movement was dependent on Cms4 and partly on Ndj1 (Chua and Roeder, 1997; Conrad et al., 1997) (Conrad et al., 2007; Kosaka et al., 2008; Wanat et al., 2008), Mps3 localization on NE did not depend on Ndj1 and Csm4 (Figure 1D). At late times such as 5 h, the *ndj1* mutant showed full coverage of Mps3 on NE (Conrad et al., 2007) with NE deformation (distorted nucleus) and motion. The *csm4* mutant formed multiple Mps3 patches/foci on NE (Conrad et al., 2007; Kosaka et al., 2008; Wanat et al., 2008), but did not show complete coverage on NE with multiple foci/patches. The *csm4* mutant is also defective in clustering and motion of Mps3 foci/patches as well as NE deformation (with round nucleus). Moreover, we examined the effect of an actin polymerization inhibitor, latrunculin B (LatB), on Mps3 localization (Trelles-Sticken et al., 2005). Treatment of meiotic cells with LatB, that disrupted the cytoplasmic actin cables but not the actin patches (Figure S1B), showed normal distribution of Mps3 foci/patches on NE during prophase (Figure 1D). However, as shown previously

(Trelles-Sticken et al., 2005), dynamic movement of Mps3-GFP was completely
absent (Figure 1D) without deformation of nuclei. These defects are similar to
those in the *csm4* mutant. At one-hour after washing out LatB, the motion of Mps3
resumed. These results suggested that Mps3 dynamics during meiosis can be
functionally separated into two processes: force-independent NE localization and
force-dependent motion. The force-dependent motion requires actin
polymerization and Csm4, and party Ndj1. On the other hand, the force-
independent NE localization does not require action polymerization or Csm4
(Ndj1), suggesting unique localization mechanism of Mps3 on NE during meiosis.
185

CDK and DDK promote efficient Mps3 localization on NE during meiosis

Because Mps3-dependent telomere movement is initiated in early meiotic
prophase (Conrad et al., 2008; Koszul et al., 2008; Trelles-Sticken et al., 2000;
Trelles-Sticken et al., 1999), we analyzed the role of two protein kinases that are
active from early meiosis; CDK and DDK. Because CDK and DDK are essential
for vegetative growth, we used conditional mutant alleles of the catalytic subunits
of CDK and DDK (Cdc28 and Cdc7), *cdc28-as1* and *cdc7-as3*, which are
sensitive to specific inhibitors,1NM-PP1 and PP1, respectively (Benjamin et al.,
2003; Wan et al., 2006). Although CDK and DDK are also essential for DNA
replication and recombination in meiosis, telomere movement during meiosis,
thus Mps3 dynamics, is independent of replication and recombination (Trelles-
Sticken *et al.* 1999; Trelles-Sticken *et al.* 2000).

We performed time-lapse analysis of Mps3-GFP in the *cdc28-as1* and *cdc7-*
as3 in either absence or present of each specific inhibitor. To inactivate CDK,
1NM-PP1, that did not affect Mps3-GFP dynamics in wild-type cells (Figure S1A),
was added to the *cdc28-as1* at 0h. In the absence of the inhibitor, the mutant cells
showed normal Mps3 localization during meiosis (Figure 1E). CDK inactivation
induced impaired Mps3 localization (Figures 1B, 1E). We detected a few Mps3
foci/patches, which clustered on one area of NE. These loose clusters of Mps3
patches were persistent during prophase I (Figures 1B). On the other hand, upon
CDK inhibition, the formation of multiple Mps3 foci/patches around NE and fully
coverage of Mps3 on NE was largely diminished. Tracking motion of Mps3 foci
showed persistent clustering with restricted motion of the foci on one region of

NE under the condition (Figure 1F). Importantly, at 2 h after washing the inhibitor,

NE localization and motion of Mps3 was recovered. CDK inactivation after meiotic

S phase (the addition of the inhibitor at 3 h) interfered with disperse localization

of Mps3 throughout NE with loose clustering of Mps3 foci/patches. Therefore,

persistent CDK activity is necessary for the establishment and maintenance of

Mps3 disperse localization, particularly the resolution of clustering, and normal

movement on NE.

A similar defect in Mps3 localization on NE and movement was observed

with DDK inactivation (*cdc7-as3* + PP1) (Figure 1B, 1E). PP1 did not affect the

localization of Mps3-GFP in wild-type cells (Figure S1A). In its absence, the *cdc7-*

as3 cells showed normal localization and motion of Mps3 foci/patches (Figures

1B, 1E). DDK inhibition by PP1 from 0 h reduced NE localization of Mps3 with

accumulation as a loose cluster at late meiosis. Removal of the inhibitor restored

the NE localization of Mps3 in the *cdc7-as3* cells. The addition of PP1 at 3 h also

induced similar Mps3 localization defect to that at 0 h. Although CDK and DDK

are essential for pre-meiotic DNA replication and DSB formation, the inactivation

of either kinase after the DNA replication and DSB formation resulted in the

impaired localization, suggesting direct regulatory role of these kinases in Mps3

dynamics. Moreover, the defect in Mps3 localization under DDK inactivation is

similar to that under CDK inactivation, suggesting that CDK and DDK work in a

same pathway to control Mps3 dynamics during meiosis.

We also analyzed the Mps3-GFP localization in the *cdc7 bob-1-1* double

mutant, in which the *bob1-1* mutation in the helicase gene *MCM5* suppresses the

lethality of *cdc7* deletion (Jackson et al., 1993; Sasanuma et al., 2008), A small

fraction (~20%) of the *cdc7 bob-1* mutant cells showed a few Mps3 foci on NE

while the most of cells did not form some Mps3 foci on NE. These defect in the

cdc7 bob-1-1 double mutant is more severe than that in the *cdc7-as3* mutant with

PP1. Whole cell immunostaining of the Mps3-Flag protein showed that reduced

CDK and DDK activities hampered localization of Mps3, but not of Ndj1, to NE

during meiosis (Figure 1G), suggesting a critical role of the kinases in the coupling

of Mps3 to Ndj1 on NE, suggesting Mps3-independent pathway for Ndj1

localization on NE. Taken together, these results revealed that CDK and DDK

activities are necessary for proper Mps3 localization and efficient movement on

NE during meiosis.

CDK and DDK promote telomere dynamics during meiosis

To know the role of CDK and DDK in telomere movement, we analyzed the motion

a green fluorescent protein (GFP) fusion of a telomere-binding protein, Rap1, as

described (Conrad et al., 2008; Koszul et al., 2008; Trelles-Sticken et al., 2000;

Trelles-Sticken et al., 1999). During vegetative growth or pre-meiosis, Rap1-GFP

exhibited 1–4 foci on NE on a single focal plane with little movement (Figure 2A)

(Trelles-Sticken et al., 2005). Once the cells entered meiosis, Rap1 foci on NE

increased in number (up to 10). Rap1-GFP foci moved rapidly and were

occasionally clustered in a single area of NE (Figure 2A), which corresponds to

the telomere bouquet (Trelles-Sticken et al., 2005). This telomere clustering was

very transient (half-life less than 5 second) and was predominantly observed after

3–4 h of incubation in a sporulation medium (Figures 2A, 2E). Rap1 foci were

more dispersed during late meiotic prophase (4–6 h for the pachytene stage)

before the onset of meiosis I (MI; Figure 2A). The tracking analysis showed that

Rap1 foci move around a meiotic nucleus in an uncoordinated way as reported

(Figure 2D) (Trelles-Sticken et al., 2005).

CDK inactivation (*cdc28-as1* + 1NM-PP1) impaired meiotic Rap1

dynamics (Figures 2B, 2E). The *cdc28-as1* showed normal Rap1 dynamics

during meiosis (Figures 2B, 2E). With compromised CDK activity from 0 h, cells

with clustered Rap1 foci accumulated during prophase I (Figure 2E). A few Rap1

foci or patches showed polarized localization on one peripheral area of a nucleus.

indicating that CDK activity is required for meiotic telomere dynamics, particularly

for the resolution of telomere clusters (but not necessarily for clustering *per se*).

The 2D tracking confirmed the restricted motion of Rap1 foci in one nuclear area

(Figure 2D). Addition of an inhibitor during early prophase I (e.g., at 3 h) also

resulted in defective resolution of telomere clustering; washing off the inhibitor

restored telomere dynamics (Figure 2B). Treatment of wild-type cells with the

inhibitor did not affect Rap1-GFP dynamics, confirming the specificity of the

inhibitor (Figure S1C). These results indicated that persistent CDK activity during

meiotic prophase I promotes normal telomere dynamics. The inactivation of CDK,

which regulates actin dynamics in mitotic cells, does not affect the formation of

cytoplasmic actin cables during meiosis (Figure S2A).

Similar to CDK inactivation, Cdc7 inactivation (the treatment of the *cdc7-*
as3 with PP1) at 0 and 3 h impaired Rap1-GFP dynamics, which showed
accumulation of clusters with limited motion (Figures 2C, 2E); whereas the
treatment of the wild-type cells with PP1 did not affect Rap1 dynamics (Figure
S1C). This defective resolution of Rap1 cluster under DDK inhibition is similar to
that seen in CDK inhibition. Moreover, the *cdc7* null mutant with *bob1-1* mutation
exhibited persistent telomere clustering (Figures 2C, 2D, 2E). These results
indicated that persistent Cdc7 kinase activity is also required for meiotic telomere
dynamics.

Mps3 is phosphorylated during meiosis
Previously, it is shown that S70 of Mps3 is phosphorylated in late prophase-I (Li
et al., 2017). We also found that Mps3 is a phosphoprotein in middle prophase I.
On western blots, Mps3-Flag showed mobility shift, which is induced during
meiotic prophase I (4 and 6 h; Figure 3C). Mps3 mobility was reduced by the
treatment with a lambda phosphatase (Figure S2B). Moreover,
immunoprecipitation (IP) of Mps3-Flag from meiotic cells showed that Mps3
cross-reacted with anti-phosphoserine antibody (Figure S2C) whose reactivity
was largely diminished by treatment with the phosphatase.

Previous systematic analyses of CDK substrates *in vitro* identified Mps3
as a Cdc28–Clb2 substrate (Ubersax et al., 2003). Mps3 contains five potential
CDK sites (S/TPXK/R or S/TP), two of which seem to be candidates for CDK/DDK
phosphorylation (188–190 <u>TSS</u>PGK and 450–451 <u>TS</u>P; Figure 2A, S2A).
Because the 450–451 site is located in the SUN domain, which is not essential
for Mps3 localization on NE (Rao *et al.* 2011), we focused on T188-S189-S190
(hereafter, TSS; Figure 3B). We purified a GST fusion protein of an Mps3
fragment (residue 160–260) including TSS. *In vitro* Cdc28–Clb5 alone, but not
Cdc7–Dbf4 alone, promoted phosphorylation of the Mps3 fragment
(approximately 10-fold compared to the background; Figure S3E). Co-incubation
with CDK and DDK resulted in robust incorporation of ^{32}P in GST-Mps3, indicating
that CDK and DDK cooperate in Mps3 phosphorylation *in vitro* (30-fold compared
to the background). Triple alanine substitutions (T188A, S189A, S190A; referred

to as the *mps3-AAA* mutant) made the GST-Mps3 fragment a poor substrate by
these kinases. These results indicated that TSS is required for efficient CDK/DDK
phosphorylation *in vitro*.

Consistent with the *in vitro* result, the Mps3-AAA mutant protein showed a
reduced band shift and decreased reactivity to anti-phosphoserine antibody
(Figures 3C and S3F). Furthermore, we raised an antibody against a Mps3
peptide with 189-phosphoserine and 190-phosphoserine (186-
GAT**pSpS**PGKSF-195) and found that this phospho-specific antibody could
recognize immunoprecipitated Mps3-Flag specifically between 4–8 h in meiosis,
but not in mitosis (Figure 3D). Strong signals were observed during late prophase
I (4 and 6 h). This antibody did not recognize precipitated Mps3-AAA-Flag. These
results indicate that 189S and 190S are phosphorylated *in vivo* specifically during
meiosis.

Accidently, the Mps3 sequence around putative phosphorylation sites is
similar to that with phosphorylation sites on Mer2 (Henderson et al., 2006;
Sasanuma et al., 2008; Wan et al., 2008), including TSSP sequence (Figure 3B).
Mer2 is a meiosis-specific protein essential for DSB formation (Cool and Malone,
1992; Engebrecht et al., 1990). CDK and DDK phosphorylate the TSS sequence
of Mer2, which are necessary for meiotic DSB formation. We swapped a 10-
amino-acid region (GATSSPGKSF) containing the Mps3 phosphorylation sites
with the corresponding region of Mer2 (TETSSPFRST; Figure 3B). This *MPS3*
allele (*MPS3-MER2S*) conferred wild-type spore viability (95%; n=50 tetrads).
The *MPS3-MER2S-Flag* exhibited similar telomere (Rap1) clustering to the wild
type (Figures 3F and 3G). The Mps3-MER2S protein cross-reacted with
antibodies against Mer2 phospho-S30 antibody (Henderson *et al.* 2006), but not
with the anti-Mps3-phospho antibody (Figures 3D and 3E). The Mps3-MER2S-
Flag protein shows meiosis-specific localization on NE which often co-localized
with Ndj1 (Figure 3H). These suggest that residues 26-35 of Mer2 are functionally
equivalent to residues 185-194 of Mps3. Like Mps3, the region of Mer2 is rich in
serine and threonine (Figure 3B).

The *mps3-AAA* mutant is defective in Mps3 localization on NE
To investigate the role of the phosphorylated residues of Mps3, we constructed

and characterized phenotypes of the *mps3-AAA-FLAG* mutant (hereafter, *mps3-*
AAA; Figure 4). The *mps3-AAA* mutant showed little apparent defects in mitosis
such as SPB duplication (Figure S3A) seen in other *mps3* mutants (Jaspersen et
al., 2002; Nishikawa et al., 2003), suggesting that TSS residues of Mps3 are not
critical for the mitotic function of Mps3. The *mps3-AAA* mutant showed defects in
meiosis with reduced spore viability of 87.5%, which is slightly higher than that of
the *ndj1* mutant (75.8%) (Chua and Roeder, 1997; Conrad et al., 1997). The
spore viability of the *ndj1 mps3-AAA* double mutant is 72.3%, which is similar to
that of the *ndj1* mutant. The *mps3-AAA* mutant delayed the entry to meiosis I by
~1 h relative to the wild type (Figure 4A). The *ndj1 mps3-AAA* double mutant
showed ~2.5 h delayed entry into meiosis I, which is similar to the *ndj1* single
mutant, indicting the epistatic relationship of the *mps3-AAA* to the *ndj1*. The
mps3-AAA mutant shows normal progression of meiotic S-phase (Figure S3B).
Immuno-staining of Zip1, a component of central region of SC, revealed delayed
formation of full length SCs together in the *mps3-AAA* mutant with delayed
loading of Zip1 at early time points (Figures S3C, S3D), suggesting that Mps3 NE
localization is necessary for efficient synapsis.
We also analyzed chromosome dynamics during meiosis of the *mps3-AAA*
mutant cells. First, we checked Rap1-GFP movement in the *mps3-AAA* mutant.
Different from the wild-type cells, the *mps3-AAA* mutant was defective in
movement of Rap1-GFP and, thus of telomeres (Figures 4B,4C). In the mutant,
the number of Rap1 foci did not increase (Figure 4B) and little movement of Rap1-
GFP foci was observed during meiosis. Rap1 motion in meiotic prophase I of the
mps3-AAA mutant is restricted to one area of NE (Figure 4C). Therefore, we
concluded that T188, S189, and S190 of Mps3 are important for meiotic telomere
dynamics. Indeed, the *mps3-AAA* mutant was defective in chromosome motion
as shown by analysis of a GFP fusion of Zip1 (Koszul *et al.* 2008). During
prophase I, Zip1-GFP shows dynamic motion (Figure 4D). The *mps3-AAA* mutant
reduced velocity of Zip1 motion by approximately 3-fold relative to the wild type
(48±15 versus 130±31 nm/sec, Figure 4E).
Next, we examined the localization of the Mps3-AAA mutant protein in
meiosis as a GFP fusion protein (Figures 4F). *mps3-AAA-GFP* cells showed 85%
of spore viability, which was comparable to the mutant stain without the GFP tag.

Similar to the wild-type Mps3-GFP, the Mps3-AAA GFP fusion protein normally

resided on the SPB as a single focus during mitosis (Figure 4F). During meiosis,

the Mps3-AAA mutant protein showed reduced NE localization. On entry into

meiosis, similar to the wild-type Mps3, the Mps3-AAA mutant protein formed a

few (usually two or three small) foci on NE, but showed little increase in the

number of Mps3 foci. At late time points, different from wild-type Mps3, the mutant

protein did not form an Mps3 patch or cover on NE (Figures 4F, 4G). Later, such

as at 6 h, two or three foci of Mps3 showed a limited movement (Figures 4H, 4I).

These results suggested that T188, S189, and S190 of Mps3 are necessary for

meiosis-specific Mps3 localization and motion on NE, particularly efficient

localization on NE accompanied with the formation of large complexes.

Interestingly, the Mps3-AAA-Flag mutant protein still bound to Ndj1 and Csm4

like wild-type Mps3 protein (Figure S3E), which is consistent with the Mps3-AAA

protein retains N-terminal nucleoplasmic region and SUN domain for Ndj1 and

Csm4 binding, respectively.

The *mps3-AAA* mutant is deficient in NE growth

In addition to localization/movement defects, the *mps3-AAA* mutant showed a

defect in nuclear morphology. First, the meiotic nucleus of the *mps3-AAA* mutant

was smaller than that of the wild type (Figure 4B). The nucleus of the wild type

grew when the cell underwent meiosis and exhibited a 4-fold increase (in area; 2

folds in length) at 4 h (Figure 4J). The treatment with LatB did not affect the

expansion, indicating that meiosis-specific NE remodeling does not require actin

polymerization. The *mps3-AAA* mutant showed defective NE growth in prophase

I; nuclear area in the *mps3-AAA* mutant is similar between 0 and 4 h (Figure 4M).

These indicate that Mps3 luminal region plays a critical role in the growth of the

meiotic nucleus, possibly by controlling NE remodeling.

Second, a fraction of the *mps3-AAA* mutant cells showed a partial defect in

telomere tethering to NE during meiosis. In one or two Rap1-GFP foci, the *mps3-

AAA* mutant (40%; n=100) was localized to the nucleoplasm rather than the

nuclear periphery, whereas all Rap1-GFP foci were localized near NE in the wild

type (Figure 4B). The weak telomere tethering defect in the *mps3-AAA* mutant is

reminiscent of the defect observed with an *NDJ1* mutant, which also showed a

partial tethering defect (Figure 4B) (Conrad et al., 1997; Trelles-Sticken et al.,
2000).
409
Aspartate substitution of TSS in Mps3 suppress CDK and DDK defects
We constructed a version of Mps3 with three negative charges on the TSS
sequence (T188D, S189D, and S190D), referred to as *MPS3-DDD* (Figure 3B).
The *MPS3-DDD* allele did not show any mitotic defects. The *MPS3-DDD* cells
showed wild-type spore viability and enter into MI like wild-type cells (Figure S3E,
F). The *MPS3-DDD* cells also displayed similar Rap1-GFP dynamics to wild-type
cells with normal clustering during prophase I (Figures 5A, 5B). The Mps3-DDD-
GFP protein localized to SPB during vegetative growth and showed wild-type like
localization on NE during meiosis (Figure 5C and Figure S3G). Mps3-DDD-GFP
displayed movement that was similar to that of the wild-type protein (Figures 5C).
The aspartate substitutions of Mps3 suppressed the resolution defect in Rap1
clustering at late time points of meiosis by CDK and DDK inactivation (Figures
5B, 5C), although some clustering persisted under the conditions (Figure 5B).
Mps3-DDD-GFP showed NE localization and movement during meiosis even
under the conditions of either CDK or DDK inactivation (Figures 5C, 5D), which
is similar to that in wild type Mps3-GFP and Mps3-DDD-GFP cells. Clustering of
Mps3 induced by CDK/DDK inactivation was clearly ameliorated by the
introduction of Mps3-DDD-GFP (Figures 5C,5C). However, there are some
restricted motion of Mps3-DDD protein under CDK/DDK inactivation relative to
Mps3-DDD protein, suggesting CDK/DDK activities are still required for wild-type
levels of Mps3 motion (Figure 5E), which may reflect partial rescue of resolution
defect in Rap1 clustering. Importantly, NE localization and dynamics of Mps3-
DDD protein still depends on meiosis. In vegetative cells, Mps3-DDD cells shows
SPB localization of Mps3-DDD and normal Rap1 localization (Figure S3H, S3I).
These results suggested that Mps3-DDD protein does not depend on CDK and
DDK for its meiosis-specific dynamics on NE.
Surprisingly, Mps3-DDD, not wild-type Mps3, show NE localization in mitotic
cells only in the *cdc7 bob-1* mutant, but not in the *cdc7as3* with the inhibitor
(Figure 5C). This suggests that, different from meiosis, DDK negatively regulates
NE localization of Mps3 in vegetative cells.

Luminal region of Mps3 binds to membrane *in vitro*

To know the role of putative phosphorylation of Mps3 luminal region, we analyzed biophysical properties of a 55 amino acid Mps3 peptide (153-208 aa) containing both transmembrane (TM) and luminal regions (Figure 3A). The luminal region is referred to as juxtamembrane (JM) region (Figure 6A). We prepared two peptides, Mps3-wild type (WT) and Mps3-DDD, which contains three aspartates (Figure 6A). Synthesized Mps3 peptides were mixed with liposome comprised of POPC/POPS (10/3)(Sato et al., 2006) First, we checked conformation of each peptides in the bilayers by polarized FT-IR (Fourier-transform Infrared spectroscopy) and found both Mps3-WT and Mps3-DDD peptides contained similar content of α-helix and β-strand (Figure S4A). Dichroic ratio calculated for Mps3-WT and Mps3-DDD were 2.90 and 2.85, suggesting that both transmembrane regions are inserted into lipid bilayers with a similar tilt angle of ~30° relative to the membrane normal (Tamagaki et al., 2014). We cannot assign those peaks from β-strand to which residues in the JM regions are involved for the β-structure with this FT-IR experiment alone.

We next synthesized Mps3 TM-JM peptide with Alexa fluorophore-dye (Alexa-568) at the C-terminus of the JM region and analyzed spectrum of the dye in presence of different concentrations of negatively charged PIP_2 (phosphatidyl-inositol 4,5-bisphosphate), which modulates status of charge on the membrane surface (Sato et al., 2009b). Here we utilized PIP_2-involving model system to see if the JM gets affected by membrane although we are aware of the fact that it has not been known whether Msp3 interacts with PIP_2. This simple model system must provide a possible mechanism of how negative charge on the JM region caused by its phosphorylation affect its behavior against the membrane.

For the wild-type peptide, the addition of PIP_2 decreased the peak intensity of fluorophore at 600 nm up to ~40% (Figure 6B). The triplicate showed similar results. Decreased fluorescence could be due to quenching of the fluorescence from the dye association (self-quenching) by inter-peptide association (Figure 6C) as shown in the previous study on an ErbB2 peptide containing TM and JM regions (Matsushita et al., 2013). These results suggest that the JM region of the Mps3-WT peptide is sensitive to the change in the membrane component and

also that the JM regions interact with the acidic lipid bilayer. On the other hand, Mps3-DDD peptide decreased the intensity to ~20% (a half of the intensity from WT peptide). The decrease in the change in the fluorescence intensity must be due to two types of electrostatic repulsion, between side chains of the aspartates on each JM regions and/or between the aspartates and the acidic lipids (Figure 6C). These repulsions must place the fluorophores apart preventing from self-quenching. The latter repulsion can reduce the JM region of Mps3-DDD peptide binding to the bilayers compared from that of the Mps3-WT. These suggest the different conformation of Mps3-WT and -DDD peptides in respect of the interaction of JM region of Mps3 to the membrane.

To get more insight on behavior of the JM region with the membrane, we utilized a technique of molecular dynamic (MD) simulation. Since there is no reported structure on the TM-JM region of Mps3, we artificially constructed the initial structure of the TM-JM sequence as follow: using a software *Chimera* (Pettersen et al., 2004), R161-M181 is set to be α-helix as the TM region; the rest was set to be random structure. This constructed structure was embedded in membrane (POPC/POPS/PIP$_2$=8:2:1 with 80 nm in XY dimension) using *Membrane Builder* module in *CHARMM-GUI* (Jo et al., 2008). The peptide-membrane system was fully hydrated in 150 mM NaCl solution.

Firstly, we performed coarse-grained molecular dynamics simulations on the constructed TM-JM peptide in membrane. We ran simulations on two sequences, Mps3-WT and Mps3-DDD. Duration of each run was >3 µs and three sets were performed for each sequence. Results from all of the calculations showed that the JM region from Mps3-DDD dissociated from the membrane (Figure S4B). Conversely, that of Mps3-WT kept attached on the membrane (Figure S4B). Secondly, we performed all-atom simulation on three TM-JM sequence peptides, Mps3-WT, Mps3-DDD and Mps3-WT phosphorylated at T197-S198-S199. Duration of each calculation was >50 ns. In Figure 6D, snap-shots from trajectories for three sequences are shown. Results from the simulation, together with mass density profiles (Figure S4C), indicate that JM regions from Mps3-DDD and phosphorylated Mps3 dissociated from the membrane. On the other hand, the JM region from Mps3-WT kept attached on the membrane.

Discussion

Here we described the multilayered control of the dynamics of a NE protein with SUN domain, Mps3, during yeast meiosis. In both mitotic and meiotic cells, Mps3 is a major component of a half bridge as well as SPIN (SPB interacting network) (Chen et al., 2019; Lee et al., 2020) (Li et al., 2017). Meiosis induces the localization and motion of Mps3 on NE, which are accompanied with the formation of a large protein ensembles in NE as the LINC complex detected as foci/patches. The meiosis-specific LINC complex tethers telomeres and interacts with cytoplasmic actin cables. In a luminal region between INM and ONM, Mps3 binds to a KASH protein, Mps2, both during mitosis and meiosis, and to a meiosis-specific Mps2-binding protein, Csm4 (Chen et al., 2019; Lee et al., 2020). Mps2 and Csm4 cooperate for the binding to the motor Myo2 in cytoplasm. Myo2 on actin cables, which is induced in meiotic cells, generates forces for the motion of the LINC complex on NE, which in turn move chromosomes inside of meiotic cells (rapid prophase movement; RPM) as well as the clustering of telomeres in zygotene stage (Conrad et al., 2008; Koszul et al., 2008; Trelles-Sticken et al., 2000; Trelles-Sticken et al., 1999). While Mps3 localization on NE is independent of actin-generated force (and other meiosis specific factors such as Ndj1 and Csm4), Mps3 movement on NE is dependent of the force. Meiosis-specific Mps3 modulation promotes the formation of a LINC complex containing Mps3 for meiotic events such as chromosome pairing/synapsis, and chromosome motion as well as NE expansion.

Localization of Mps3 on NE in meiotic cells is divided into several distinct steps (Figures 1A and 7); 1) In addition to the SPB, at early times in prophase I, a few Mps3 foci appear on NE. 2) With increased numbers of the foci, patches of Mps3, large cohesive ensembles with multiple foci, are formed on NE, which sometimes associated with clustering in one area of NE. 3) Multiple foci and patches cover on entire NE. 4) Mps3 shows continuous coverage on NE (sometimes with a hole at nucleus-vacuole junction; NVJ). 5) During late prophase I or pro-metaphase I, most of Mps3 on NE disappear from NE, leaving two SPB-associating foci. At each stage, Mps3 foci/patches show RPM which is maximum at mid/late prophase I (Conrad et al., 2008; Koszul et al., 2008; Trelles-Sticken et al., 2000; Trelles-Sticken et al., 1999).

In this study, we found that two cell-cycle kinases, CDK and DDK, promote

proper Mps3 localization on NE during meiosis, particularly the resolution of

clustered Mps3 foci (and, thus, telomere clustering), suggesting that the

phosphorylation of a target protein(s) plays a role in resolution of Mps3 clusters.

Moreover, we showed that the JM region of Mps3 near INM, which is located in

lumen of NE, is critical for the NE localization of the protein. S179 and S180, and,

probably T178, of the JM region seems to be subject to meiosis-specific

phosphorylation. We also showed that introducing negative charges in the JM

region changes the binding ability of the region to the membrane in reconstituted

liposomes. These support the idea that Mps3 localization on NE is a part of a

novel meiotic regulatory system known as "NE remodeling." Probably, non-

canonical phosphorylation of the JM region is likely to promote the meiosis-

specific NE localization of Mps3.

Phosphorylation of NE luminal region

Although our results described here suggest one intriguing hypothesis that CDK

and DDK collaborate to catalyze the phosphorylation of the JM region of Mps3

for its NE localization, this is less likely for several reasons. It is unlikely that CDK

and DDK are responsible for direct phosphorylation reaction in lumen

compartments. These kinases are not localized to luminal regions of both

endoplasmic reticulum (ER) and NE since the catalytic subunits (Cdc28 and

Cdc7) and its regulatory subunits (Clb5,6 and Dbf4) lack a leader sequence

necessary for the luminal localization.

The protein phosphorylation of the luminal region between INM and ONM

is very unusual, and have not been reported in the past. On the other hand, recent

studies have shown that the phosphorylation of secreted proteins and

oligosaccharides of membrane proteins is catalyzed by an atypical kinase such

as Fam20C in the lumen of Golgi apparatus and ER in vertebrate cells

(Tagliabracci et al., 2012; Tagliabracci et al., 2013; Tagliabracci et al., 2015).

Fam20C, whose structure and sequence are divergent from typical kinases, uses

ATP as a donor of phosphate, but prefers Mn^{2+} rather than Mg^{2+} for the catalysis

(Tagliabracci et al., 2012; Tagliabracci et al., 2013; Tagliabracci et al., 2015).

Given a luminal region of NE is continuous to that of ER where INM/ONM proteins

are synthesized, it is possible that a similar activity to Fam20C family might be
present in the lumen between INM and ONM, which may catalyze the
phosphorylation of NE luminal region of Mps3. However, there is no report on the
presence of Fam20C orthologues or relative enzymes in lower eukaryotes such
as yeasts. Alternatively, the phosphorylation of the luminal region might be
mediated by a novel mechanism. Given membranes are rich in phospho-lipids,
the phosphate group in the lipids as a donor might be transferred to proteins by
a novel mechanism. We need further studies to identify a protein responsible for
the phosphorylation of JM region of Mps3.

A previous study showed that S70 in the nucleoplasmic region of Mps3 is
phosphorylated during late prophase I, which facilitates the removal of the N-
terminal region of Mps3 by proteosome-dependent cleavage (Li et al., 2017). This
removal may play a role in disappearance of Mps3 from NE in late prophase I
and for separation of duplicated SPBs (Li et al., 2017). During meiosis of *C.*
elegans, the N-terminal nucleoplasmic region of Sun-1 protein is phosphorylated
by Chk-1 and Cdk-1 (Penkner et al., 2009; Zuela and Gruenbaum, 2016). In M
phase of human cells, SUN1 protein is phosphorylated at its N-terminal region by
CDK and PLK (Patel et al., 2014). Together with results described here, these
suggest the importance of the phosphorylation in the regulation of dynamics
and/or regulation of SUN-domain proteins.

Th role of negative charges in JM region of Mps3
Biophysical analysis of synthetic peptides containing TM and JM regions of Mps3
in synthetic liposomes revealed that the JM region of Mps3 has a capability to
bind to the lipid bilayers. Importantly, introducing three negative charges in the
JM region weakens the binding to the membrane, resulting in a change of
geometry of JM region of Mps3, as extension, the luminal region of Mps3
including SUN domain. As a typical LINC complex, SUN domain of Mps3 in INM
binds to KASH domain of Msp2 in ONM. The connection would span ~20 nm
between INM and ONM (Chen et al., 2019; Lee et al., 2020). Recently, Jasperson
and her colleagues showed "atypical" configuration of Mps3-Mps2, in which not
only SUN-KASH interaction, but also N-terminal regions of both Mps3 and Mps2
bind to each other in SPIN which surrounds SPB (Chen et al., 2019; Lee et al.,

2020). Typical LINC could accommodate *trans*-membrane configuration in the
luminal region while atypical LINC would do *cis*-membrane configuration (Figure
7). These suggest that luminal regions of Mps3 and Mps2 are quite flexible to
accommodate two different SUN-KASH configurations. Our results support that
the switch from *cis*- to *trans*-membrane configurations might be regulated by the
binding of the JM region of Mps3 to the INM. While the interaction between JM
region of Mps3 and INM promotes *cis*-membrane form of noncanonical LINC
complex, negative charges induced by phosphorylation may reduce the
interaction between JM and INM, which in turn might accommodate a
conformation suitable for the *trans*-membrane form. Alternatively, negatively
charged JM region of Mps3 would be a binding site of the other protein which
may promote the configuration switch.
618
What is a target by CDK- and DDK in chromosome motion?
In this study, we showed that, in the budding yeast, two cell cycle kinases, CDK
and DDK, collaborate in meiotic telomere dynamics by regulating proper
localization of Mps3 on NE. As discussed above, it is unlikely that CDK and DDK
directly phosphorylate the luminal region of Mps3. Moreover, the comparison of
cdk/ddk mutants with the *mps3-AAA* mutant reveals the difference in meiotic
defects in Mps3 localization. CDK/DDK inactivation results in persistent telomere
clustering with Mps3 polarized localization. These indicate that CDK and DDK
activities promote the resolution of Mps3 foci/patches and/or telomeres rather
than the NE localization *per se*. Mps3 localization on NE seems be normal even
in the absence of CDK and DDK activities. Phenotypic similarity between *cdk* and
ddk mutants on Mps3 localization suggests a common target of these kinases for
the resolution of the clusters. It is well known that CDK phosphorylation primes
the DDK phosphorylation, which is seen in Mer2 phosphorylation during meiosis
(Henderson et al., 2006; Sasanuma et al., 2008; Wan et al., 2008). Interestingly,
telomere-resolution defects under CDK and DDK inactivation are similar to that
in the mutant of the *REC8* gene (Challa et al., 2016), which encodes a meiosis-
specific kleisin subunit of the cohesin complex (Klein et al., 1999) and cohesin
regulator, *RAD61/WPL1* (Challa et al., 2016). It is well known that Rec8 is
phosphorylated by DDK (and Polo and CK1 kinases), but not by CDK (Katis et

al., 2010) and that Rad61/Wpl1 is phosphorylated by DDK and Polo-like kinase

(PLK). Thus, there might be the other target in cohesin components by these

kinases. In mitotic yeast cells, Eco1, which catalyzes acetylation of Smc3 cohesin

subunit, is phosphorylated by CDK and DDK for the degradation (Seoane and

Morgan, 2017). Interestingly, Eco1 is shown to acetylate Mps3 in vegetative cells

(Ghosh et al., 2012).

Telomere clustering during prophase I persists with CDK or DDK inactivation,

suggesting the presence of a constraint on the resolution of the clustering. The

phosphorylation, thus negative charges, in JM region, might overcome the

constraint for the resolution. This is supported by the fact that the *MPS3-DDD*

partially rescued the resolution defect of telomere clustering with CDK or DDK

inactivation. Moreover, we found Mps3-DDD, not wild-type Mps3, show NE

localization in mitotic cells only in the *cdc7 bob-1* mutant (Figure 5C), suggesting

the presence of DDK-dependent constraints on NE localization of Mps3 during

mitosis.

Meiosis establishes Mps3 localization on NE

As described above, Mps3 localization on NE is regulated in distinct stages

(Figure 7). The *mps3-AAA* mutant still retains an ability to form early meiosis-

specific Mps3 foci on NE. And also Mps3-DDD protein needs meiosis for NE

localization during meiosis. These suggest that Mps3 localization on NE is

initiated and/or established during early meiotic prophase I. Given that Mps3

localization as an ensemble on NE is specific to meiosis, there might be an early

meiosis-specific factor or modification that regulates the establishment of the

Mps3 localization as an ensemble on NE upon the entry into meiosis. We

examined two meiosis-specific factors, Ndj1 and Csm4, both of which are

involved in chromosome motion and Mps3 dynamics, but it is unlikely that these

factors function in the establishment of Mps3 localization. Further work is

necessary to identify the factors necessary for the initiation event of NE

remodeling.

We showed that non-canonical phosphorylation of the luminal JM region

of Mps3 promotes the formation of multiple Mps3 foci/patches on NE. The *mps3-*

AAA mutant is deficient in the transition to early Mps3 foci to middle Mps3

foci/patches, suggesting the role of negative charges in JM region in the transition,

thus the maintenance of Mps3 localization on NE.

NE remodeling controls various nuclear events

Characterization of the *mps3-AAA* single mutant revealed that Mps3 localization

to NE (NE remodeling) play multiple roles during meiosis. As expected from

previous reports (Koszul et al., 2008; Scherthan et al., 2007), Mps3-dependent

NE remodeling is important for chromosome motion and thus chromosome

pairing/synapsis. Mps3 phosphorylation seems to be essential for the formation

of active NE-bound ensembles for chromosome motion via recruitment of various

proteins and promotion of telomere tethering. Moreover, we found that the *mps3-*

AAA single mutant is defective in meiosis-specific enlargement of nuclei, which

presumably is facilitated by NE expansion. NE-bound Mps3 may regulate NE

growth by directly promoting lipid synthesis on NE or importing lipids from the ER

(Sosa Ponce et al., 2020).

Materials and Methods

Strains and plasmids

All strains described here are derivatives of SK1 diploids and are described in Table S1. Strain construction is described in the Extended Experimental Procedures. GST fusion genes of a PCR-amplified Mps3 fragment (160–260 amino acid residues of the Mps3 protein) were constructed on pGEX-2T.

Anti-serum and antibodies

Anti-HA antibody (16B12; Babco), anti-Flag (M2, Sigma), anti-tubulin (MCA77G, Bio-Rad/Serotec, Ltd), anti-GFP (3E6; Molecular Probes), and rabbit anti-Dmc1 (Shinohara et al., 2000) were used for staining. Anti-Zip1 (rabbit) and anti-Red1 (chicken) were described previously (Shinohara et al., 2008). Rabbit anti-Rec8 antibody was described previously (Rao et al., 2011). Guiana pig anti-Hop1 serum was described in (Bani Ismail et al., 2014). The second antibodies for staining were Alexa-488 (Goat) and -594 (Goat) IgG used at a 1/2000 dilution (Themo Fishers).

10 amino acid of Mps3 peptide with 189-phosphoserine and 190-phosphoserine (186-GAT**pSpS**PGKSF-195) was synthesized and used for the immunization of rabbit. From obtained serum, antibody specific phosphor-peptide was affinity-purified using the phosphor-peptide (MBL Co. Ltd. Japan).

Cytology

Time-lapse images of Mps3-GFP, Rap1-GFP and Zip1-GFP were captured using a computer-assisted fluorescence microscope system (DeltaVision; Applied Precision). The objective lens was an oil immersion lens (100×; NA, 1.35). Image deconvolution was performed using an image workstation (SoftWorks; Applied Precision). Time-lapse image acquisition was performed every 0.5 to 1 s at a single focal plane. Meiotic cell cultures (~150μl) were placed on bottom-glass dishes (Matsunami) pre-coated with Concavalin A (10 mg/ml, Sigma)

The sectioned area of each nucleus with Rap1-GFP, which showed nuclear and dotty staining, was measured using the Velocity™ program (Applied Precision) with manual assignment of the rim of the sectioned area. Ninety-six

722 nuclei were counted. Tracking of Zip1-GFP/Mps3-GFP/Rap1-GFP was analyzed

723 using the ImageJ program (NIH) and plug-in MTrack2 (Nico Shuuman).

IP and Western blotting

Yeast cell lysates were prepared by the glass bead disruption method. The

lysates were incubated with magnetic beads (Dynal M260; GE Healthcare)

coated with anti-Flag antibody (M2, Sigma) for 12 h and washed extensively.

Bound proteins were eluted by adding the SDS sample buffer and were analyzed

on an SDS-PAGE gel, transferred to a nylon membrane (Millipore Co. Ltd), and

probed with specific antibodies.

Analyses of meiotic recombination

734 Time-course analyses of DNA events in meiosis and cell cycle progression were

735 performed as described previously (Shinohara et al., 1997; Storlazzi et al., 1996).

Southern Blotting

Southern blotting analysis was performed with the same procedure as in

(Storlazzi et al., 1995). For the *HIS4-LEU2* locus, genomic DNA prepared was

digested with *Xho*I (for crossover) and *Pst*I (for meiotic DSBs). For the *URA3-*

ARG4 locus, the DNA were digested with *Xho*I. Probes for southern blotting were

Probe 155 for crossover and Probe 291 for DSB detection on the *HIS4-LEU2* as

described in (Xu et al., 1995). Image gauge software (Fujifilm Co. Ltd., Japan)

was used for quantification for bands.

***In vitro* kinase assay**

For *in vitro* phosphorylation, 1 μg of purified GST-Mps3 or GST-Mps3-AAA was

incubated with γ-^{32}P-ATP (100 μM; 2.2 MBq) in the presence of CDK (~0.2μg)

and/or DDK (~0.2μg) in a 20 μl of buffer [20 mM Tris-HCl (pH 7.5), 10 mM MgCl$_2$,

10 mM β-glycerophosphate]. After 1-h incubation at 30°C, the reaction was

stopped by adding 5× SDS-PAGE sample buffer, and the products were analyzed

on 10% SDS-PAGE gel. Dried gels were analyzed using a phosphorimager BAS

2000 (Fuji Co. Ltd).

Synthesis of Mps3 peptides

Peptides with the TM and JM region of Mps3 (residue 153-208) were synthesized by solid-phase method with the following sequence: GEAKKLK<u>WRTYIFYGGLFFVFYFFGSFL</u>MTTVKNNDLESHSSGA**TSS**PGKSFSNL-Cys and GEAKKLK<u>WRTYIFYGGLFFVFYFFGSFL</u>MTTVKNNDLESHSSGA**DDD**PGKSFSNL-Cys (underlines indicate TM region). The synthetic peptides were purified by reverse-phase HPLC on a C4 column with gradient of 2-propanol over water containing 0.1% trifluoroacetic acid. For fluorescent-labeling, Alexa Fluor 568 C5-maleimide (Molecular probe) was introduced to the sulfide group on cysteine by mixing the peptides and the fluorescent dye in dimethylformamide under basic conditions.

Preparation of lipid bilayer with peptides

The Mps3 peptides were co-solubilized with 1-palmitoyl-2-oleoyl-*sn*-phosphatidycholine (POPC), 1-palmitoyl-2-oleoyl-*sn*-phosphatidyserine (POPS) and octyl-β-glucoside in trifluoroethanol (TFE). The peptide to lipid molar ratio was 1:100. The lipid concentration was 200 μM. The ratio of POPC and POPS was 10:3. The mixture was incubated for 90 min at 37°C. Then TFE was removed under stream of Ar. MES buffer (10 mM MES, 50 mM NaCl, and 5 mM DTT, pH 6.2) was added to the solid from the previous step and mixed at 37 °C for 6 h. The octyl- β -glucoside was removed by dialysis against the MES buffer.

Measurement of FT-IR

Polarized attenuated total reflection FT-IR spectra were obtained on a JASCO FT/IR-4700 spectrometer. Membranes containing Mps3 TM–JM peptides were layered on a germanium internal reflection element using a slow flow of nitrogen gas directed at an oblique angle to the IR plate to form an oriented multilamellar lipid–peptide film.1000 scans were acquired and averaged for each sample at a resolution of 4 cm^{-1}. The absorption of polarized light by the amide I bond yields the dichroic ratio defined as a ratio of absorption intensity for parallel, relative to perpendicular, polarized light. From the dichroic ratio, we estimated the tilt angle of the TM helix relative to the membrane normal based on the method described

by Smith and coworkers (Liu et al., 2004; Smith et al., 2002)) using a value of

41.8° for angle α between the helix director and the transition-dipole moment of

the amide I vibrational mode. For the FT-IR experiment, the amount of lipid used

per experiment was ~4 mg. The film on the ATR plate in our experiment was

assumed to be greater than ~10 μm. For calculating the dichroic ratio, the thick

film limit is applicable (Bechinger et al., 1999). Equations that we used for the

calculation of the dichroic ratios were based on this assumption.

Fluorescence spectroscopy

For experiments with PIP_2, TM-JM peptides were inserted into POPC/POPS

membrane as explained above. The peptide/lipid ratio was set to 1:1000 for

observing the fluorescence from Alexa568. The lipid concentration was 200–250

μM in MOPS buffer (1 mM MOPS, 0.1 M KCl, pH 7.0). Vesicles were formed by

extrusion of multilamellar vesicles through 200-nm polycarbonate filters. PIP_2

were added into the membranes by addition of PIP_2 micelles to the vesicle

solution. Fluorescence was measured within 1 hour to minimize PIP_2 hydrolysis.

The PIP_2 concentration was from 5 μM to 25 μM. Fluorescence measurement

was carried out on HITACHI F-2500 fluorescence spectrophotometer at the

excitation wavelength of 568 nm.

Coarse-grained MD simulation

All simulations were performed using GROMACS 2016 package (Abraham et al.,

2015) using the MARTINI force field with polarizable water (Marrink et al., 2007).

Periodic boundary conditions were applied in all directions, and the time step was

20 fs.   The temperature was controlled with the Berendsen temperature coupling

scheme (Berendsen et al., 1984)with a time constant of 1 ps, and the pressure

was controlled using the Parrinello–Rahman semi-isotropic barostat (Parrinello

and Rahman, 1981) with a time constant of 12 ps and a compressibility of 3×10^{-4}

bar^{-1}.   All the simulations were carried out in the isothermal–isobaric (NPT)

ensemble at a temperature of 303 K and pressure of 1.0 bar. The cut-off of the

Van der waals interactions were set to 1.1 nm.

All atom MD simulation

The NAMD software package (Phillips et al., 2005) was used to perform all-atom
MD simulations. The CHARMM36m force field (Huang et al., 2017) together with
TIP3P water model were used for all simulations.　All simulations in NPT
dynamics (constant particle number, pressure and temperature) were carried out
at 303 K with Langevin dynamics, and at 1 atm maintained by Nosè-Hoover
Langevin piston method. Full electrostatic interactions were treated by the
particle mesh Ewald (PME) approach with a grid spacing of less than 1 Å. The
cut-off radii of long-range electrostatic and van der Waals interactions were set
to be 12 Å, with a smoothing function applied from 10 Å.
830
831

Acknowledgments

We are grateful to Drs. Nancy Hollingsworth, Franz Klein, Kunihiro Ohta, Kazu

Nishikawa, Hisao Masai, and Hiroyuki Araki for providing the materials used in

this study. HBDPR was supported by the BMC program and a scholarship from

the Graduate School of Science, Osaka University, JASSO as well as MEXT. KC

KC were supported by the Institute for Protein Research. This work was

supported by a Grant-in-Aid from the JSPS KAKENHI Grant Number; 22125001,

22125002, 15H05973 and 16H04742, 19H00981 to A.S.

Competing interest

The authors declare no competing financial interest.

Author contributions

H.B.D.P.R., T.S., K.C., M.S. and A.S. designed the experiments. H.B.D.P.R., T.S.,

K.C., M.S. and A.S. carried out experiments. M.S provided materials. H.B.D.P.R.,

T.S., K.C. and A.S. analyzed the data. A.S. wrote the manuscript with inputs from

H.B.D.P.R., T.S., K.C. and M.S.

References

Abraham, M.J., T. Murtola, R. Schulz, S. Páll, J.C. Smith, B. Hess, and E. Lindahl.

2015. GROMACS: High performance molecular simulations through multi-

level parallelism from laptops to supercomputers. *SoftwareX*. 1-2:19-25.

Bani Ismail, M., M. Shinohara, and A. Shinohara. 2014. Dot1-dependent histone

H3K79 methylation promotes the formation of meiotic double-strand

breaks in the absence of histone H3K4 methylation in budding yeast. *PloS*

one. 9:e96648.

Bechinger, B., J.M. Ruysschaert, and E. Goormaghtigh. 1999. Membrane helix

orientation from linear dichroism of infrared attenuated total reflection

spectra. *Biophysical journal*. 76:552-563.

Benjamin, K.R., C. Zhang, K.M. Shokat, and I. Herskowitz. 2003. Control of

landmark events in meiosis by the CDK Cdc28 and the meiosis-specific

kinase Ime2. *Genes & development*. 17:1524-1539.

Berendsen, H.J.C., J.P.M. Postma, W.F.v. Gunsteren, A. DiNola, and J.R. Haak.

1984. Molecular dynamics with coupling to an external bath. *The Journal*

of Chemical Physics. 81:3684-3690.

Challa, K., M.S. Lee, M. Shinohara, K.P. Kim, and A. Shinohara. 2016.

Rad61/Wpl1 (Wapl), a cohesin regulator, controls chromosome

compaction during meiosis. *Nucleic Acids Res*. 44:3190-3203.

Chen, J., J.M. Gardner, Z. Yu, S.E. Smith, S. McKinney, B.D. Slaughter, J.R.

Unruh, and S.L. Jaspersen. 2019. Yeast centrosome components form a

noncanonical LINC complex at the nuclear envelope insertion site. *J Cell*

Biol. 218:1478-1490.

Chikashige, Y., T. Haraguchi, and Y. Hiraoka. 2007. Another way to move

chromosomes. *Chromosoma*. 116:497-505.

Chua, P.R., and G.S. Roeder. 1997. Tam1, a telomere-associated meiotic protein,

functions in chromosome synapsis and crossover interference. *Genes &*

development. 11:1786-1800.

Conrad, M.N., A.M. Dominguez, and M.E. Dresser. 1997. Ndj1p, a meiotic

telomere protein required for normal chromosome synapsis and

segregation in yeast. *Science (New York, N.Y.)*. 276:1252-1255.

Conrad, M.N., C.Y. Lee, G. Chao, M. Shinohara, H. Kosaka, A. Shinohara, J.A.

885    Conchello, and M.E. Dresser. 2008. Rapid telomere movement in meiotic
886      prophase is promoted by NDJ1, MPS3, and CSM4 and is modulated by
887      recombination. *Cell.* 133:1175-1187.

888  Conrad, M.N., C.Y. Lee, J.L. Wilkerson, and M.E. Dresser. 2007. MPS3 mediates
889      meiotic bouquet formation in Saccharomyces cerevisiae. *Proc Natl Acad*
890      *Sci U S A.* 104:8863-8868.

891  Cool, M., and R.E. Malone. 1992. Molecular and genetic analysis of the yeast
892      early meiotic recombination genes REC102 and REC107/MER2.
893      *Molecular and cellular biology.* 12:1248-1256.

894  Engebrecht, J., J. Hirsch, and G.S. Roeder. 1990. Meiotic gene conversion and
895      crossing over: their relationship to each other and to chromosome
896      synapsis and segregation. *Cell.* 62:927-937.

897  Ghosh, S., J.M. Gardner, C.J. Smoyer, J.M. Friederichs, J.R. Unruh, B.D.
898      Slaughter, R. Alexander, R.D. Chisholm, K.K. Lee, J.L. Workman, and S.L.
899      Jaspersen. 2012. Acetylation of the SUN protein Mps3 by Eco1 regulates
900      its function in nuclear organization. *Mol Biol Cell.* 23:2546-2559.

901  Hawley, R.S., and W.D. Gilliland. 2009. Homologue pairing: getting it right. *Nature*
902      *cell biology.* 11:917-918.

903  Henderson, K.A., K. Kee, S. Maleki, P.A. Santini, and S. Keeney. 2006. Cyclin-
904      dependent kinase directly regulates initiation of meiotic recombination.
905      *Cell.* 125:1321-1332.

906  Hiraoka, Y., and A.F. Dernburg. 2009. The SUN rises on meiotic chromosome
907      dynamics. *Dev Cell.* 17:598-605.

908  Huang, J., S. Rauscher, G. Nawrocki, T. Ran, M. Feig, B.L. de Groot, H.
909      Grubmüller, and A.D. MacKerell, Jr. 2017. CHARMM36m: an improved
910      force field for folded and intrinsically disordered proteins. *Nature methods.*
911      14:71-73.

912  Jackson, A.L., P.M. Pahl, K. Harrison, J. Rosamond, and R.A. Sclafani. 1993. Cell
913      cycle regulation of the yeast Cdc7 protein kinase by association with the
914      Dbf4 protein. *Molecular and cellular biology.* 13:2899-2908.

915  Jaspersen, S.L., T.H. Giddings, Jr., and M. Winey. 2002. Mps3p is a novel
916      component of the yeast spindle pole body that interacts with the yeast
917      centrin homologue Cdc31p. *J Cell Biol.* 159:945-956.

Jo, S., T. Kim, V.G. Iyer, and W. Im. 2008. CHARMM-GUI: A web-based graphical
user interface for CHARMM. *Journal of Computational Chemistry*.
29:1859-1865.

Katis, V.L., J.J. Lipp, R. Imre, A. Bogdanova, E. Okaz, B. Habermann, K. Mechtler,
K. Nasmyth, and W. Zachariae. 2010. Rec8 phosphorylation by casein
kinase 1 and Cdc7-Dbf4 kinase regulates cohesin cleavage by separase
during meiosis. *Dev Cell*. 18:397-409.

Klein, F., P. Mahr, M. Galova, S.B. Buonomo, C. Michaelis, K. Nairz, and K.
Nasmyth. 1999. A central role for cohesins in sister chromatid cohesion,
formation of axial elements, and recombination during yeast meiosis. *Cell*.
98:91-103.

Kosaka, H., M. Shinohara, and A. Shinohara. 2008. Csm4-dependent telomere
movement on nuclear envelope promotes meiotic recombination. *PLoS*
genetics. 4:e1000196.

Koszul, R., K.P. Kim, M. Prentiss, N. Kleckner, and S. Kameoka. 2008. Meiotic
chromosomes move by linkage to dynamic actin cables with transduction
of force through the nuclear envelope. *Cell*. 133:1188-1201.

Koszul, R., and N. Kleckner. 2009. Dynamic chromosome movements during
meiosis: a way to eliminate unwanted connections? *Trends in cell biology*.
19:716-724.

Lee, C.Y., C.G. Bisig, M.M. Conrad, Y. Ditamo, L. Previato de Almeida, M.E.
Dresser, and R.J. Pezza. 2020. Extranuclear Structural Components that
Mediate Dynamic Chromosome Movements in Yeast Meiosis. *Current*
biology : CB. 30:1207-1216.e1204.

Li, P., H. Jin, B.A. Koch, R.L. Abblett, X. Han, J.R. Yates, 3rd, and H.G. Yu. 2017.
Cleavage of the SUN-domain protein Mps3 at its N-terminus regulates
centrosome disjunction in budding yeast meiosis. *PLoS genetics*.
13:e1006830.

Liu, W., M. Eilers, A.B. Patel, and S.O. Smith. 2004. Helix packing moments
reveal diversity and conservation in membrane protein structure. *Journal*
of molecular biology. 337:713-729.

Marrink, S.J., H.J. Risselada, S. Yefimov, D.P. Tieleman, and A.H. de Vries. 2007.
The MARTINI Force Field: Coarse Grained Model for Biomolecular

Simulations. *The Journal of Physical Chemistry B*. 111:7812-7824.

Matsushita, C., H. Tamagaki, Y. Miyazawa, S. Aimoto, S.O. Smith, and T. Sato.
2013. Transmembrane helix orientation influences membrane binding of
the intracellular juxtamembrane domain in Neu receptor peptides. *Proc*
Natl Acad Sci U S A. 110:1646-1651.

Nishikawa, S., Y. Terazawa, T. Nakayama, A. Hirata, T. Makio, and T. Endo. 2003.
Nep98p is a component of the yeast spindle pole body and essential for
nuclear division and fusion. *J Biol Chem*. 278:9938-9943.

Parrinello, M., and A. Rahman. 1981. Polymorphic transitions in single crystals:
A new molecular dynamics method. *Journal of Applied Physics*. 52:7182-
7190.

Patel, J.T., A. Bottrill, S.L. Prosser, S. Jayaraman, K. Straatman, A.M. Fry, and S.
Shackleton. 2014. Mitotic phosphorylation of SUN1 loosens its connection
with the nuclear lamina while the LINC complex remains intact. *Nucleus*
(Austin, Tex.). 5:462-473.

Penkner, A.M., A. Fridkin, J. Gloggnitzer, A. Baudrimont, T. Machacek, A. Woglar,
E. Csaszar, P. Pasierbek, G. Ammerer, Y. Gruenbaum, and V. Jantsch.
2009. Meiotic chromosome homology search involves modifications of the
nuclear envelope protein Matefin/SUN-1. *Cell*. 139:920-933.

Pettersen, E.F., T.D. Goddard, C.C. Huang, G.S. Couch, D.M. Greenblatt, E.C.
Meng, and T.E. Ferrin. 2004. UCSF Chimera—A visualization system for
exploratory research and analysis. *Journal of Computational Chemistry*.
25:1605-1612.

Phillips, J.C., R. Braun, W. Wang, J. Gumbart, E. Tajkhorshid, E. Villa, C. Chipot,
R.D. Skeel, L. Kalé, and K. Schulten. 2005. Scalable molecular dynamics
with NAMD. *J Comput Chem*. 26:1781-1802.

Rao, H.B., M. Shinohara, and A. Shinohara. 2011. Mps3 SUN domain is important
for chromosome motion and juxtaposition of homologous chromosomes
during meiosis. *Genes Cells*. 16:1081-1096.

Sasanuma, H., K. Hirota, T. Fukuda, N. Kakusho, K. Kugou, Y. Kawasaki, T.
Shibata, H. Masai, and K. Ohta. 2008. Cdc7-dependent phosphorylation
of Mer2 facilitates initiation of yeast meiotic recombination. *Genes &*
development. 22:398-410.

Sato, A., B. Isaac, C.M. Phillips, R. Rillo, P.M. Carlton, D.J. Wynne, R.A. Kasad,
and A.F. Dernburg. 2009a. Cytoskeletal forces span the nuclear envelope
to coordinate meiotic chromosome pairing and synapsis. *Cell*. 139:907-
919.
Sato, T., P. Pallavi, U. Golebiewska, S. McLaughlin, and S.O. Smith. 2006.
Structure of the membrane reconstituted transmembrane-juxtamembrane
peptide EGFR(622-660) and its interaction with Ca2+/calmodulin.
Biochemistry. 45:12704-12714.
Sato, T., T.C. Tang, G. Reubins, J.Z. Fei, T. Fujimoto, P. Kienlen-Campard, S.N.
Constantinescu, J.N. Octave, S. Aimoto, and S.O. Smith. 2009b. A helix-
to-coil transition at the epsilon-cut site in the transmembrane dimer of the
amyloid precursor protein is required for proteolysis. *Proc Natl Acad Sci U*
S A. 106:1421-1426.
Scherthan, H., H. Wang, C. Adelfalk, E.J. White, C. Cowan, W.Z. Cande, and D.B.
Kaback. 2007. Chromosome mobility during meiotic prophase in
Saccharomyces cerevisiae. *Proc Natl Acad Sci U S A*. 104:16934-16939.
Seoane, A.I., and D.O. Morgan. 2017. Firing of Replication Origins Frees Dbf4-
Cdc7 to Target Eco1 for Destruction. *Current biology : CB*. 27:2849-
2855.e2842.
Sheehan, M.J., and W.P. Pawlowski. 2009. Live imaging of rapid chromosome
movements in meiotic prophase I in maize. *Proc Natl Acad Sci U S A*.
106:20989-20994.
Shinohara, M., S.L. Gasior, D.K. Bishop, and A. Shinohara. 2000. Tid1/Rdh54
promotes colocalization of rad51 and dmc1 during meiotic recombination.
Proceedings of the National Academy of Sciences of the United States of
America. 97:10814-10819.
Shinohara, M., S.D. Oh, N. Hunter, and A. Shinohara. 2008. Crossover
assurance and crossover interference are distinctly regulated by the ZMM
proteins during yeast meiosis. *Nat Genet*. 40:299-309.
Shinohara, M., E. Shita-Yamaguchi, J.M. Buerstedde, H. Shinagawa, H. Ogawa,
and A. Shinohara. 1997. Characterization of the roles of the
Saccharomyces cerevisiae RAD54 gene and a homologue of RAD54,
RDH54/TID1, in mitosis and meiosis. *Genetics*. 147:1545-1556.

Smith, S.O., C. Smith, S. Shekar, O. Peersen, M. Ziliox, and S. Aimoto. 2002.
Transmembrane interactions in the activation of the Neu receptor tyrosine
kinase. *Biochemistry*. 41:9321-9332.

Sosa Ponce, M.L., S. Moradi-Fard, V. Zaremberg, and J.A. Cobb. 2020. SUNny
Ways: The Role of the SUN-Domain Protein Mps3 Bridging Yeast Nuclear
Organization and Lipid Homeostasis. *Frontiers in genetics*. 11:136.

Starr, D.A., and H.N. Fridolfsson. 2010. Interactions Between Nuclei and the
Cytoskeleton Are Mediated by SUN-KASH Nuclear-Envelope Bridges.
Annu Rev Cell Dev Biol.

Storlazzi, A., L. Xu, L. Cao, and N. Kleckner. 1995. Crossover and noncrossover
recombination during meiosis: timing and pathway relationships. *Proc Natl*
Acad Sci U S A. 92:8512-8516.

Storlazzi, A., L. Xu, A. Schwacha, and N. Kleckner. 1996. Synaptonemal complex
(SC) component Zip1 plays a role in meiotic recombination independent
of SC polymerization along the chromosomes. *Proceedings of the National*
Academy of Sciences of the United States of America. 93:9043-9048.

Tagliabracci, V.S., J.L. Engel, J. Wen, S.E. Wiley, C.A. Worby, L.N. Kinch, J. Xiao,
N.V. Grishin, and J.E. Dixon. 2012. Secreted kinase phosphorylates
extracellular proteins that regulate biomineralization. *Science (New York,*
N.Y.). 336:1150-1153.

Tagliabracci, V.S., L.A. Pinna, and J.E. Dixon. 2013. Secreted protein kinases.
Trends in biochemical sciences. 38:121-130.

Tagliabracci, V.S., S.E. Wiley, X. Guo, L.N. Kinch, E. Durrant, J. Wen, J. Xiao, J.
Cui, K.B. Nguyen, J.L. Engel, J.J. Coon, N. Grishin, L.A. Pinna, D.J.
Pagliarini, and J.E. Dixon. 2015. A Single Kinase Generates the Majority
of the Secreted Phosphoproteome. *Cell*. 161:1619-1632.

Tamagaki, H., Y. Furukawa, R. Yamaguchi, H. Hojo, S. Aimoto, S.O. Smith, and
T. Sato. 2014. Coupling of transmembrane helix orientation to membrane
release of the juxtamembrane region in FGFR3. *Biochemistry*. 53:5000-
5007.

Taxis, C., C. Maeder, S. Reber, N. Rathfelder, K. Miura, K. Greger, E.H. Stelzer,
and M. Knop. 2006. Dynamic organization of the actin cytoskeleton during
meiosis and spore formation in budding yeast. *Traffic*. 7:1628-1642.

Trelles-Sticken, E., C. Adelfalk, J. Loidl, and H. Scherthan. 2005. Meiotic
telomere clustering requires actin for its formation and cohesin for its
resolution. *J Cell Biol*. 170:213-223.

Trelles-Sticken, E., M.E. Dresser, and H. Scherthan. 2000. Meiotic telomere
protein Ndj1p is required for meiosis-specific telomere distribution,
bouquet formation and efficient homologue pairing. *J Cell Biol*. 151:95-106.

Trelles-Sticken, E., J. Loidl, and H. Scherthan. 1999. Bouquet formation in
budding yeast: initiation of recombination is not required for meiotic
telomere clustering. *Journal of cell science*. 112 (Pt 5):651-658.

Ubersax, J.A., E.L. Woodbury, P.N. Quang, M. Paraz, J.D. Blethrow, K. Shah,
K.M. Shokat, and D.O. Morgan. 2003. Targets of the cyclin-dependent
kinase Cdk1. *Nature*. 425:859-864.

Wan, L., H. Niu, B. Futcher, C. Zhang, K.M. Shokat, S.J. Boulton, and N.M.
Hollingsworth. 2008. Cdc28-Clb5 (CDK-S) and Cdc7-Dbf4 (DDK)
collaborate to initiate meiotic recombination in yeast. *Genes &
development*. 22:386-397.

Wan, L., C. Zhang, K.M. Shokat, and N.M. Hollingsworth. 2006. Chemical
inactivation of cdc7 kinase in budding yeast results in a reversible arrest
that allows efficient cell synchronization prior to meiotic recombination.
Genetics. 174:1767-1774.

Wanat, J.J., K.P. Kim, R. Koszul, S. Zanders, B. Weiner, N. Kleckner, and E. Alani.
2008. Csm4, in collaboration with Ndj1, mediates telomere-led
chromosome dynamics and recombination during yeast meiosis. *PLoS
genetics*. 4:e1000188.

Xu, L., M. Ajimura, R. Padmore, C. Klein, and N. Kleckner. 1995. NDT80, a
meiosis-specific gene required for exit from pachytene in Saccharomyces
cerevisiae. *Mol Cell Biol*. 15:6572-6581.

Zickler, D., and N. Kleckner. 1998. The leptotene-zygotene transition of meiosis.
Annu Rev Genet. 32:619-697.

Zickler, D., and N. Kleckner. 1999. Meiotic chromosomes: integrating structure
and function. *Annu Rev Genet*. 33:603-754.

Zuela, N., and Y. Gruenbaum. 2016. Matefin/SUN-1 Phosphorylation on Serine
43 Is Mediated by CDK-1 and Required for Its Localization to Centrosomes

and Normal Mitosis in C. elegans Embryos. *Cells*. 5.

1084

    Figure legends

    Figure 1. Meiotic Mps3 localization and movement on NEs

    (A) Mps3-GFP localization was analyzed in wild-type cells (PRY64) at various

        time points during meiosis. Bottom color bars indicates classes of Mps3-

        localization in (B). Bar indicates 2 μm.

    (B) Based on Mps3-GFP localization, cells with Mps3-GFP were classified into

        four classes (A) and quantified: single Mps3 focus (blue), 2–5 foci (green),

        more than 5 foci/patches (orange), and coverage on NE (red). At each time

        point, more than 150 nuclei were counted.

    (C) Time-lapse analysis of Mps3-GFP in the wild-type strain (PRY64) at different

        time points in meiosis. A single focal plane of a cell was analyzed every 3 sec.

    (D) Time-lapse analysis of Mps3-GFP in various strains at different time points in

        meiosis. A single focal plane of a cell was analyzed every 3 sec. The inhibitor

        LatB (30 μM) was added at 0 h. While washing, the inhibitor was washed at

        5 h, and the cells were analyzed at 6 h. Wild type, PRY64; *ndj1*, PRY192;

        csm4, PRY198.

    (E) Time-lapse analysis of Mps3-GFP in the *cdc28-as1* mutant (PRY71) treated

        with or without the inhibitor 1NM-PP1 (0.5 μM) at 5 h during meiosis. The

        inhibitor was added at 0 h (second panels) or 3 h (third panels). While

        washing (fourth panels), the inhibitor was added at 0 h and washed at 3 h,

        and the cells were analyzed at 5 h. Time-lapse analysis of Mps3-GFP in the

        cdc7-as3 mutant (PRY260) treated with or without the inhibitor PP1 (15 μM).

        The *cdc7 bob1-1* mutant (PRY115) was analyzed at 6 h.

    (F) Tracking of Mps3-GFP in wild type, *cdc28-as1* (PRY71 with the inhibitor),

        cdc7-as3 (PRY260with the inhibitor), and *cdc7 bob1-1* (PRY115). Tracking

        was monitored for all Mps3 foci/patches in a single cell for 120 sec. Each line

        represents tracking of the foci at a single focal plane.

    (G) Whole-cell staining analysis of Mps3-Flag and Ndj1. Various cells were

        stained with anti-Flag for Mps3-Flag (red) and anti-HA for Ndj1 (green) at the

        nuclear periphery at indicated times. *cdc28-as1* (PRY13) and *cdc7-as3*

        (PRY72) cells were treated with their inhibitor from 0 h and examined at the

        indicated times. Wild type, HKY404.

1118

Figure 2. Meiotic telomere movement in *cdk* and *ddk* mutants

(A–C) Time-lapse analysis of Rap1-GFP in various strains (wild type [A; HKY167], *cdc28-as1* [B, PRY68], and *cdc7-as3* [C, PRY79] and *cdc7 bob1-1* [C, PRY116]) at different time points during meiosis. An image of a single focal plane of each mutant cell was taken every 3 sec. For the *cdc28-as1* and *cdc7-as3* mutants, the inhibitors (0.5 μM of 1NM-PP1 and 15 μM PP1, respectively) were added at 0 h (second panels) or 3 h (third panels). While washing (bottom panels), the inhibitor was added at 0 h and washed at 3 h, and the cells were analyzed at 5 h. Clustering of telomeres is shown by arrows. Bar indicates 2 μm.

(D) Tracking of Rap1-GFP in wild type (0 and 4h), *cdc28-as1* (with the inhibitor), and *cdc7-as3* (with the inhibitor). Tracking was monitored for all Rap1 foci in a single cell for 120 sec. Each line represents tracking of the center of foci at a single focal plane. Red dotted lines indicate the rim of each nucleus.

(E) Kinetics of telomere clustering during meiosis were quantified. At each time point, more than 150 nuclei were counted for clustering of telomeres (arrows).

Figure 3. Mps3 is phosphorylated in luminal region

(A) Schematic representation of the Mps3 protein. The putative transmembrane (TM) region and SUN domain are shown in green and gray, respectively.

(B) Sequence homology of an Mps3 region with a CDK/DDK phosphorylation site of the Mer2 protein. Identical amino acids are shown in red. Putative CDK or DDK phosphorylation sites are shown in yellow. The substitutions (*mps3-AAA, MPS3-DDD*, and *MPS3-MER2S*) for the putative phosphorylation sites of Mps3 are shown.

(C) Western blotting of Mps3 and tubulin at various times in meiosis. Cell lysates from the *MPS3-Flag* (right, HKY404) and *MPS3-AAA-Flag* (left, PRY163) strains were analyzed by western blotting using anti-Flag and anti-tubulin antibodies. The wild-type Mps3 protein shows band shift during meiosis for example at 4 and 6 h.

(D) Western blotting of the Mps3-Flag protein during meiosis. Wild-type (left, HKY404), *mps3-AAA* mutant (middle, PRY163) and *MPS3-MER2S* mutant

proteins (right, PRY514) were analyzed at different time points. Fractions
immunoprecipitated with anti-Flag antibody were probed with anti-Flag
antibody (top) as well as a phospho-specific antibody (bottom) that
recognized T189 and S190 phosphorylation.
(E) Western blotting of the Mps3-Flag protein during meiosis. Wild-type (left,
HKY404) and *MPS3-MER2s* proteins (right, PRY514) were analyzed at
different time points. Fractions immunoprecipitated with anti-Flag antibody
were probed with anti-Flag and anti-Mer2-phospho-S30 antibodies.
(F, G) Rap1-GFP dynamics at 4 h of meiosis analyzed in *MPS3-MER2s* (PRY518)
and wild-type (HKY167) strains. Kinetics of Rap1 clustering in each strain is
shown in G. Counting was performed as described in Figure 2E. Blue, wild
type; red, *MPS3-MER2S.*
(G) Localization of Ndj1-HA (green), Mps3-Mer2S-Flag (red), and DAPI (dark
blue) was analyzed by whole cell staining of PRY514. Bar indicates 2 μm.
1165

Figure 4. The *mps3-AAA* mutant impairs telomere and NE movement
(A) Meiosis division was analyzed by DAPI staining for various strains. More than
100 cells were counted for divisions. Spore viability of each strain is shown in
parenthesis. At least 100 asci were dissected for each strain.
(B) Time-lapse analysis of Rap1-GFP in the wild-type (HKY167) and *mps3-AAA*
(PRY138) strains at different time points in meiosis. A single focal plane of a
cell was analyzed every 3 s. A tethering defect was observed in the *mps3-*
AAA strains. Bar indicates 2 μm.
(C) Tracking of Rap1-GFP in wild type (left) and *mps3-AAA*. See Figure 2E for
quantification.
(D) Time-lapse analysis of Zip1-GFP in wild type (SEY672) and *mps3-AAA*
(PRY332).
(E) Tracing of Zip1-GFP shows a step size of each Zip1-GFP line. Step size per
given time is converted into the relative velocity of chromosomes. Blue bars;
wild type; red bars, *mps3-AAA.*
(F) Localization of wild-type Mps3-GFP (PRY64) and Mps3-AAA-GFP (PRY186)
proteins in a cell at different time points in meiosis.
(G) Kinetics of Mps3 distribution. Based on Mps3-GFP morphology, cells with

Mps3-GFP were classified into four classes and quantified: 2–5 foci (green),
more than 5 foci/patches (orange), and coverage on NE (red). See Figure 1B
for quantification.

(H) Time-lapse analysis of wild type and Mps3-AAA-GFP during meiosis.

(I) Tracking of Mps3-GFP in wild type (left) and *mps3-AAA*. See Figure 1F for
quantification.

(J) The maximum sectional area of each nucleus was measured using the
Velocity program. Each area of the 96 nuclei is ranked in the figure. Graphs
for wild type and *mps3-AAA* at 0 and 4 h as well as wild type treated with LatB
are shown.

(K) Interaction of Mps3-Flag with Ndj1-HA and Csm4 in the *mps3-AAA* mutant
was confirmed by immuno-precipitation using anti-Flag antibodies. IPed
fractions of cells with Mps3-Flag (left, HKY404) or Mps3-AAA-Flag (right,
PRY163) at each time point were probed with anti-Flag (top), anti-HA (middle)
and anti-Csm4 antibodies.

1199

Figure 5. Acid amino acids in lumen of Mps3 allele can suppress CDK and
DDK defects

(A) Time-lapse analysis of Rap1-GFP in a phosphomimetic allele of *MPS3*,
MPS3-DDD (PRY211), with a different combination of *cdc28-as1* (PRY211),
cdc7-as3 (PRY301)*, and *cdc7 bob1-1* (PRY272) mutations in meiosis. A
single focal plane of a cell was analyzed at every 3 s. Bar indicates 2 μm.

(B) Kinetics of clustering of telomere as Rap1-clusters was studied in *cdc7-as3*
(left, PRY309) and *cdc28-as1* (right, PRY303) in the presence of its specific
inhibitors. Wild type (blue, HKY167) and *MPS3-DDD* (PRY236) with the allele
was used.

(C) Time-lapse analysis of Mps3-DDD-GFP with different mutant alleles and in
treatment with a specific inhibitor at different time points in meiosis.

(D) Kinetics of Mps3. Based on Mps3-GFP morphology, cells with Mps3-GFP
were classified into four classes and quantified: single Mps3 focus (blue), 2–
3 foci (green), 4–5 foci (orange), and more than 5 foci (red). See Figure 2B
for quantification. Graphs for wild type, *cdc28-as1*, and *cdc7-as3* cells are the
same as in Figure 2B.

(E) Tracking of Mps3-GFP in wild type (left) and *mps3-AAA*. See Figure 2D for
quantification.
1219

Figure 6. Biophysical analysis of JM region of Mps3
(A) Amino acid sequences of Mps3 peptides used. Amino acid sequence in
transmembrane region is shown in green. Phosphorylated residues TSS and
its derivative DDD are shown in red.
(B) Fluorescence spectra from a synthetic liposome containing Alexa568-labelled
WT-Mps3 (left) peptide and Mps3-DDD peptide (right) were measured in the
presence of various concentrations of PIP_2.
(C) Schematic illustration of the binding of Mps3-WT peptide (left) and Mps3-DDD
peptide (right) to lipid bilayer with PIP_2. DDD residues may induce electric
repulsion wither between the peptides or between the peptide and the
membrane.
(D) Snapshots from all-atom MD simulation trajectories for Mps3-WT peptide (left),
Mps3-DDD peptide (center) and Mps3-WT peptide with phosphorylation
(right).
1234

Figure 7. A model of regulation of Mps3 on NE during meiosis
During mitosis, Mps3 forms a non-canonical LINC complex with *cis*-membrane
interaction with Mps2 as a component of the half bridge and SPIN (left). Upon the
induction of meiosis, unknown factor(s) promotes the formation of Mps3 foci on
NE. Then, the JM region of Mps3 is subject to non-canonical phosphorylation,
which in turn weaken the binding of the JM to INM, probably by electric repulsion
between the JM regions with negative charges and acid lipids. This process
promotes the formation of canonical LINC complex with the *trans*-membrane
configuration, in which Mps3 SUN domain binds to KASH domain of Mps2 and
Csm4. Csm4 may promote a structural change of luminal region of Mps2. The N-
terminal region of Mps3, which is located in nucleoplasm binds to a telomere-
binding protein, Ndj1. During middle and late prophase I, Mps3 forms a large
protein ensembles on NE, which is seen as a patch. Mps3 foci and patches are
clustered on one area of NE during zygotene stage, which corresponds with
telomere bouquet. The resolution of Mps3 clusters is promoted by CDK and DDK

activities. During mitosis, DDK and other factor(s) limits the NE localization of

Mps3.

1252

Figure 1. Rao et. al.

Figure 2. Rao et. al.

Figure 3 Rao et. al.

A

Mps3

173-199
150-157

1   452   620   682

SUN domain

TM domain

B

DDD   Mps3-DDD
AAA   Mps3-AAA

173   180   191   199

TTVKNNDLESHSSGATSSPGKS-FSNLQK   Mps3

STDVSEANSEHSLMITETSSPFRSIFSHSG   Mer2
20   30   39
180   191   199

TTVKNNDLESHSSTETSSPFRSIFSHQK   Mps3-Mer2S

C

MPS3-AAA-Flag   MPS3-Flag
0 2 4 6   0 2 4 6   (h)

WB: Anti-Flag — Mps3-Flag

WB: Anti-Tub — Tubulin

D

MPS3-Flag   non-tag   MPS3-AAA-Flag   MPS3-MER2S-Flag
0 2 4 6 8   4   0 2 4 6 8   0 2 4 6 8   (h)

WB: Anti-Flag — Mps3-Flag

WB: Anti-P-Mps3 — P-S189 P-S190 (Mps3)

E

MPS3-MER2S-Flag   MPS3-Flag
0 2 4 6   0 2 4 6   (h)

WB: Anti-Flag — Mps3-Flag

WB: Anti-Mer2-pS30 — P-S30 (Mer2)

F

Rap1-GFP

0   3   6   9   12   (sec)

Wild type (4 h)

MPS3-MER2S (4 h)

G

Rap1-clustering

Percent of cells

0   2   4   6

Time in SPM (h)

Wild type
MPS3-MER2S

H

Ndj1/Mps3/DAPI

Wild type (4 h)

MPS3-MER2S (4 h)

MPS3-MER2S (0 h)

Figure 4 Rao et. al.

Figure 5 Rao et. al.

Figure 6. Rao et. al.

A

Mps3-WT   NH2-GEAKKLKWRTYIFYGGLFFVFFFGSFLMTTVKNNDLESHSSGATSSPGKSFSNL

Mps3-DDD   NH2-GEAKKLKWRTYIFYGGLFFVFFFGSFLMTTVKNNDLESHSSGADDDPGKSFSNL

B

C

D

Figure 7 Rao et. al.

 Supplemental information on Rao et al.

 Supplemental Figure

1255

Figure S1. Meiotic telomere movement in the presence of the inhibitors

(A) To check the off-target of two inhibitors, 1NM-PP1 (for *cdc28-as1*) and PP1

(for *cdc7-as3*), wild type diploid cells (HKY404) with Mps3-GFP were grown

in the presence of each inhibitor (0.5 μM of 1NM-PP1, top panels; 15 μM PP1,

bottom panels). The inhibitors were added at 0 h (second panels) and time-

lapse images were examined in 4 h-incubation with SPM. The two inhibitors

did not affect rapid movement of Mps3-GFP foci on NEs. Bar indicates 2 μm.

(B) Cytoplasmic actin cables were visualized by Phalloidin (red) and DAPI

(purple) staining of *cdc28-as1* (PRY68) cells after 5 h of incubation in SPM

both without (left) or with (middle) its inhibitor 1NM-PP1 from 0 h. As a control,

the mutant cells were treated with 30 μM of latrunculin B (LatB) at 4 h and

analysed for actin staining at 5 h (right). Bar indicates 2 μm.

(C) To check the off-target of two inhibitors, 1NM-PP1 (for *cdc28-as1*) and PP1

(for *cdc7-as3*), wild type diploid cells with Rap1-GFP were grown in the

presence of each inhibitor (0.5 μM of 1NM-PP1, top panels; 15 μM PP1,

bottom panels). The inhibitors were added at 0 h (second panels) and time-

lapse images were examined in 4 h-incubation with SPM. The two inhibitors

did not affect rapid movement of Rap1-GFP foci on NEs. Bar indicates 2 μm.

1274

Figure S2. CDK and DDK are necessary for Mps3 localization on NE

(A) Schematic representation of the Mps3 protein. Five CDK consensus

sequences are shown by red lines. The putative transmembrane (TM) region

and SUN domain are shown in blue and gray, respectively.

(B) *In vivo* phosphorylation was confirmed by a decreased band shift of Mps3-

Flag protein purified from meiotic cell lysates (4 h incubation with SPM) using

the anti-Flag antibody. After immuno-precipitation, the precipitate was

incubated with 1 unit of CIP for one hour either in the presence or absence of

phosphatase inhibitors. Mps3-Flag was detected using anti-Flag (M2)

antibody.

(C) Affinity-purified-Mps3-Flag from meiotic cell (HKY407) lysates (4 h incubation

with SPM) using anti-Flag column was treated with CIP as described in (B)

and was probed with anti-Phospho-Serine antibody (Qiagen). During

incubation, Mps3 protein was degraded extensively (bottom).

(D) Reactivity of wild-type Mps3-Flag and Mps3-AAA mutant protein to anti-

phospho-serine antibody was examined by Western blotting. Wild-type Mps3-

Flag and Mps3-AAA mutant protein at each time point were IPed and probed

with the antibody. While wild-type Mps3-Flag shows an increase reactivity to

the antibody, Mps3-AAA-Flag did not increase the reactivity in meiosis (4 h)

compared to mitosis (0 h).

(E) *In vitro* phosphorylation analysis of GST-Mps3 fusion proteins. Purified wild-

type GST-Mps3 and GST-Mps3-AAA fragments (shown on the left by

Coomassie staining) were incubated with different combinations of purified

the Cdc28–Clb5 complex and/or Cdc7–Dbf4 kinase complexes in the

presence of γ-^{32}P-ATP. After 1-h incubation, proteins were fractionated on a

SDS-PAGE gel and analyzed using a phosphorimager for incorporation of ^{32}P

in the Mps3 fragments (right bottom) and Cdc28/Cdc7 protein (right top). Cdc7

was phosphorylated by CDK. Quantification is shown below the

autoradiogram.

1304

Figure S3. Meiotic phenotypes of *mps3-AAA* and *MPS3-DDD* mutants

(A) Whole cell staining of Mps3-Flag and Tubulin. Wild type (MSY832/833) and

the *mps3-AAA* mutant (PRY122) cells in mitosis were stained with anti-Flag

(red) and anti-tubulin (green). Representative images with short spindles with

two Mps3 foci are shown. Bar indicates 2 μm.

(B) FACS analysis of DNA contents in wild type and the *mps3-AAA* mutant cells

at indicated times. Cells at each time point were fixed and stained with

propidium iodide (PI) and analyzed by FACS after the sonication.

(C) Immunostaining analysis of meiotic chromosome proteins. For SC formation,

Zip1 localization (red) with DAPI (blue) image is also analyzed for wild type

(top) and *mps3-AAA* mutant strains (bottom) at 4 h.

(D) Kinetics of SC formation. SC elongation was analyzed by staining Zip1 protein.

Class I (blue) contains Zip1 dots, Class II (green) contains partial lines of Zip1,

and Class III (red) contains fully elongated Zip1 lines.

(E) Localization of Red1 (green) and Hop1 (red) was analyzed in the wild type
(top) and *mps3-AAA* mutant (bottom) at 4 h. Bars indicate 1 µm.

(F) Meiosis division was analyzed by DAPI staining for wild type and the *MPS3-*
DDD strains (PRY201).

(G) Localization of Mps3-DDD-GFP protein in wild type (PRY211) at different time
points. Bar indicates 2 µm.

(H) Localization of Mps3-DDD-GFP protein in wild type, *cdc28-as1* (with 1NM-
PP1; PRY301) and *cdc7-as3* (with PP1; PRY272) at 0 h. Bar indicates 2 µm.

(I) Time-lapse analysis of Rap1-GFP in the *mps3-DDD* strains with a different
combination of *cdc28-as1*, *cdc7-as3* or *cdc7 bob1-1* mutations in mitosis (0
h). A single-focal plane of a cell was analyzed at every 3 second. Bar indicates
2 µm.

1331

Figure S4. Structure and dynamics of JM region of Mps3

(A) Polarized attenuated total reflection FT-IR spectra are shown. Spectra (red)
a synthetic liposome containing Mps3-WT (left) peptide and Mps3-DDD
peptide (right) were obtained with light polarized parallel (top) or
perpendicular (bottom). Deconvoluted spectra into α-helix and β-sheet are
shown in blue.

(B) Snapshots from coarse-grained MD simulation trajectories for Mps3-WT
peptide (left), Mps3-DDD peptide (right).

(C) Mass density profiles relative to the center of the lipid bilayer for residues,
T197, S198 and S199 on Mps3-WT peptide (blue), Mps3-DDD peptide (red)
and Mps3-WT peptide with phosphorylation (green). Density profile for
phosphate groups on lipids are colored with yellow.

1344
1345

1346 **Table S1: Strain list**

Strain Name	Genotype
NKY1551	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", his4B-LEU2/his4X-LEU2-URA3, arg4-bgl/arg4-nsp*
MSY832/833	*MATa/α, ho::LYS2/", ura3/", leu2::hisG/", trp1::hisG/", lys2/"*
HKY404	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", MPS3-3FLAG::KanMX6/", NDJ1-3HA::KanMX6/ "*
PRY13	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", MPS3-3FLAG::KanMX6/", NDJ1-3HA::KanMX6/", cdc28-as1/"*
PRY72	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", MPS3-3FLAG::KanMX6/", NDJ1-3HA::KanMX6/",cdc7-as3-9MYC/"*
PRY 64	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", MPS3-3GFP::KanMX6/"*
PRY71	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", MPS3-3GFP::KanMX6/", cdc28-as1/"*
PRY260	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", MPS3-3GFP::KanMX6/", cdc7-as3-9MYC/"*
PRY115	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", MPS3-3GFP::KanMX6/", cdc7::KanMX6/", bob1-1/"*
PRY122	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188,S189A,S190A-PSTI/"*
PRY186	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188,S189A,S190A-PSTI-3GFP::KanMX6 /"*
PRY163	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188,S189A,S190A-PSTI-3FLAG::KanMX6 /", NDJ1-3HA::KanMX6/"*
PRY138	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188,S189A,S190A-PSTI/", Rap1-GFP::LEU2/"*
PRY201	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188D,S189D,S190D-BSPEI/"*
PRY211	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188D,S189D,S190D-BSPEI-3GFP::KanMX6/"*

PRY219	*MAT**a**/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188D,S189D,S190D-BSPEI-3FLAG::KanMX6/" NDJ1-3HA::KanMX6/"*
PRY236	*MAT**a**/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188D,S189D,S190D-BSPEI-3GFP::KanMX6/", Rap1-GFP::LEU2/"*
PRY301	*MAT**a**/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188D,S189D,S190D-BSPEI-3GFP::KanMX6/", cdc28-as1/"*
PRY272	*MAT**a**/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188D,S189D,S190D-BSPEI-3GFP::KanMX6/", cdc7-as3::9MYC/"*
HKY167	*MAT**a**/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", Rap1-GFP::LEU2/"*
PRY68	*MAT**a**/ α, ho::LYS2/", lys2/", leu2::hisG/", ura3/",, Rap1-GFP::LEU2/", cdc28-as1/"*
PRY79	*MAT**a**/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/",, Rap1-GFP::LEU2/", cdc7-as3::9MYC/"*
PRY116	*MAT**a**/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/",, Rap1-GFP::LEU2/", cdc7::KanMX6/", bob1-1/"*
PRY303	*MAT**a**/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188D,S189D,S190D-BSPEI/", Rap1-GFP::LEU2/", cdc28-as1/ "*
PRY309	*MAT**a**/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188D,S189D,S190D-BSPEI/", Rap1-GFP::LEU2/", cdc7-as3::9MYC/"*
PRY322	*MAT**a**/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188D,S189D,S190D-BSPEI/", Rap1-GFP::LEU2/", cdc7::KanMX6/", bob1-1/"*
PRY344	*MAT**a**/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", his4B-LEU2/his4X-LEU2-URA3, arg4-bgl/arg4-nsp, mps3-T188,S189A,S190A-PSTI/" rad50S::URA3/rad50S::URA3*
PRY192	*MAT**a**/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", MPS3-*

	3GFP::KanMX6/", ndj1::KanMX6/"
PRY198	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", MPS3-3GFP::KanMX6/", csm4::KanMX6/"*
PRY514	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", Mer2-Mps3 (swap)-SacI FLAG::KanMX6/", Rap1-GFP::LEU2/", NDJ1-HA/"*
SEY672	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", trp1::hisG/", ZIP1-GFP::KamMX6/"*
PRY332	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", trp1::hisG/", ZIP1-GFP::KamMX6/", mps3-T188,S189A,S190A-PSTI-3GFP::KanMX6 /"*
PRY311	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", lys2/", his4B-LEU2/his4X-LEU2-URA3, arg4-bgl/arg4-nsp, mps3-T188,S189A,S190A-PSTI /"*
PRY147	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", mps3-T188,S189A,S190A-PSTI /", ndj1::KamMX6/ ndj1::KamMX6*
HKY329	*MATa/α, ho::LYS2/", lys2/", leu2::hisG/", ura3/", trp1::hisG/", his4B-LEU2/his4X-LEU2-URA3, arg4-bgl/arg4-nsp, ndj1::KamMX6/"*

1347

Supplemental Figure S1 Rao et. al.

A

Mps3-GFP

0   3   6   9   12 (sec)

Wild type (5 h)
(add 1NM-PP1
at 0 h)

Wild type (5 h)
(add of PP1
at 0 h)

B

cdc28-as1 (5 h)

No addition	+ 1NM-PP1	+ LatB

actin
/DAPI

C

Rap1-GFP

0   3   6   9   12 (sec)

Wild type (5 h)
(add of 1NM-PP1
at 0 h)

Wild type (5 h)
(add of PP1
at 0 h)

Supplemental Figure S2 Rao et. al.

A

B

+	+	+	−	Mps3-Flag
+	+	−	+	CIP
−	+	−	+	Inhibitor

WB: Anti-Flag

— Mps3-Flag

C

+	+	+	−	Mps3-Flag
−	+	+	+	CIP
−	−	+	+	Inhibitor

WB: Anti-P-Ser

—Mps3-Flag

WB: Anti-Flag

—Mps3-Flag

F

Wild type *MPS3-AAA*

0   4   0   4   (h)

WB: Anti-P-Ser

—Mps3-Flag

WB: Anti-Flag

—Mps3-Flag

E

CBB   Autoradograph

+	+	+	−	+	+	+	−	GST-Mps3(160-260)
−	−	−	+	−	−	−	+	GST-Mps3-AAA(160-260)
+	−	+	+	+	−	+	+	DDK
−	+	+	+	−	+	+	+	CDK

— Cdc7
— Cdc28

— GST-Mps3(160-260)

Supplemental Figure S4 Rao et. al.