Phosphorylation of luminal region of the SUN-domain protein Mps3 promotes nuclear envelope localization during meiosis

H. B. D. Prasada Rao[1], Takeshi Sato[2], Kiran Challa[3], Yurika Fujita, Miki Shinohara[4], and Akira Shinohara*

Institute for Protein Research, Osaka University, Suita, Osaka 565-0871, Japan

Running head: Meiotic control of NE remodelling

Keywords: meiosis; phosphorylation; Mps3; chromosome motion; Nuclear envelope (NE); SUN domain

[1]Current address: National Institute for Animal Biotechnology, Hyderabad, India

[2]Current address: Kyoto Pharmaceutical University, Kyoto, Japan

[3]Current address: Paul Scherrer Institute, Villigen, Switzerland

[4]Current address: Graduate School of Agriculture, Kindai University, Nara, Japan

*Correspondence:

Akira Shinohara

Institute for Protein Research, Osaka University

3-2 Yamadaoka, Suita, Osaka 565-0871, Japan

Telephone: 81-6-6879-8624

FAX: 81-6-6879-8626

E-mail: ashino@protein.osaka-u.ac.jp

ORCID: 0000-0003-4207-8247

Abstract

During meiosis, protein ensembles in the nuclear envelope (NE) containing SUN- and KASH-domain proteins, called linker nucleocytoskeleton and cytoskeleton (LINC) complex, promote the chromosome motion. Yeast SUN-domain protein, Mps3, forms multiple meiosis-specific ensembles on NE, which show dynamic localisation for chromosome motion; however, the mechanism by which these Mps3 ensembles are formed during meiosis remains largely unknown. Here, we showed that the cyclin-dependent protein kinase (CDK) and Dbf4-dependent Cdc7 protein kinase (DDK) regulate meiosis-specific dynamics of Mps3 on NE, particularly by mediating the resolution of Mps3 clusters and telomere clustering. We also found that the luminal region of Mps3 juxtaposed to the inner nuclear membrane is required for meiosis-specific localisation of Mps3 on NE. Negative charges introduced by meiosis-specific phosphorylation in the luminal region of Mps3 alter its interaction with negatively charged lipids by electric repulsion in reconstituted liposomes. Phospho-mimetic substitution in the luminal region suppresses the localisation of Mps3 via the inactivation of CDK or DDK. Our study revealed multi-layered phosphorylation-dependent regulation of the localisation of Mps3 on NE for meiotic chromosome motion and NE remodelling.

Introduction

During meiosis, homologous chromosomes pair with each other. This pairing leads to juxtaposition of the chromosomes along their entire length, resulting in the formation of the synaptonemal complex (SC). SC is a proteinaceous structure comprising two chromosome axes called the axial/lateral element (AE/LE), flanking a central region with a ladder-like structure (Hawley & Gilliland, 2009; Zickler & Kleckner, 1999). AE formation is initiated in the leptotene stage of meiotic prophase I, accompanied by the assembly of the chromosome axis with multiple chromatin loops. In the zygotene stage, some homologous AE pairs form a short SC, while in the pachytene stage, the SC elongates along the entire chromosome. After the pachytene stage, the SC dismantles in the diplotene stage before the onset of meiosis I.

Meiotic cells also exhibit a unique arrangement of chromosomes in the nucleus. In most organisms, telomeres attach to nuclear envelopes (NEs) in early meiotic prophase (or premeiotic phase) and occasionally cluster in one area of NE, particularly near the vicinity of the centrosome (a microtubule organising centre) (Zickler & Kleckner, 1998). Telomere clustering is seen only in the zygotene stage and is known as "telomere/chromosome bouquet." In the pachytene stage, telomere bouquets are resolved, and telomeres are dispersed over the entire NE. The bouquet configuration of chromosomes is proposed to promote homologous pairing by restricting the homology search process from three dimensions to two dimensions (Zickler & Kleckner, 1998).

The dynamic nature of meiotic chromosomes in various organisms is conserved between yeasts and mammals (Hiraoka & Dernburg, 2009; Koszul & Kleckner, 2009; Burke, 2018; Lee et al., 2020b; Paouneskou & Jantsch, 2019). The oscillating motion of nuclei during meiotic prophase I has been well described in fission yeast (Chikashige et al., 1994). Nuclear movement is driven by cytoplasmic microtubule cables, along which the spindle pole body (SPB), an yeast centrosome equivalent, embedded in NE moves together with the clustered telomeres. This telomere-led movement promotes the pairing of homologous loci on chromosomes. Although less coordinated, chromosome motion throughout prophase I has been reported in budding yeast, nematodes, maize and mouse

(Lee et al., 2015; Penkner et al., 2009; Sheehan & Pawlowski, 2009; Shibuya,

Ishiguro, & Watanabe, 2014).

Protein ensembles embedded in NE that connect telomeres in the

nucleoplasm with the cytoskeleton in the cytoplasm, which are referred to as the

linker of nucleocytoskeleton and cytoskeleton (LINC) complex, promote meiotic

chromosome motion (Hiraoka & Dernburg, 2009; Starr & Fridolfsson, 2010)

(Jahed, Domkam, Ornowski, Yerima, & Mofrad, 2021; Lee et al., 2020b; Wong,

Loo, & Stewart, 2021). The LINC complex necessary for chromosome motion

contains an inner nuclear membrane (INM) protein harbouring the SUN

(Sad1/UNc-84) domain and an outer nuclear membrane (ONM) protein with the

KASH (Klarsicht, ANC-1, SYNE1 Homology) domain. The SUN and KASH

domains interact with each other in a space between the INM and ONM, luminal

region, or peri-nuclear space. SUN-domain proteins bind to telomeres and/or

nucleocytoskeletons in the nucleoplasm, while KASH-domain proteins bind to

cytoskeletons. LINC complexes transmit forces generated by the cytoskeleton to

NEs for nuclear positioning and migration in somatic cells and telomeres for

chromosome motion in meiotic cells (Starr & Fridolfsson 2010).

In *Saccharomyces cerevisiae*, telomeres are tethered to NEs as several

clusters during mitosis and move around NEs during meiotic prophase I. These

telomeres are transiently clustered during prophase I (Conrad et al., 2008; Koszul,

Kim, Prentiss, Kleckner, & Kameoka, 2008; Trelles-Sticken, Dresser, & Scherthan,

2000; Trelles-Sticken, Loidl, & Scherthan, 1999). Telomere movement on NE

drives chromosome motion inside the nucleus (Koszul et al., 2008; Scherthan et

al., 2007). At least three distinct regulatory steps facilitate chromosome motion

during budding yeast meiosis. First, meiosis-specific components of the LINC

complex, such as Ndj1 and Csm4, are expressed and incorporated. Ndj1 is a

meiosis-specific telomere-binding protein necessary for telomere tethering to

NEs (Chua & Roeder, 1997; Conrad, Dominguez, & Dresser, 1997), whereas

Csm4 is a meiosis-specific ONM protein (Conrad, Lee, Wilkerson, & Dresser,

2007; Kosaka, Shinohara, & Shinohara, 2008; Wanat et al., 2008), which

interacts with a unique class of KASH protein, Mps2 (Chen et al., 2019; Lee et

al., 2020a). Second, actin cable formation is specifically induced in the cytoplasm

of meiotic cells (Koszul et al., 2008; Taxis et al., 2006). Third, NE dynamics,

including NE remodelling and movement, occur prominently in meiotic prophase
I (Conrad *et al.* 2007; this paper). NE remodelling is accompanied by NE
localisation of a yeast SUN-domain protein, Mps3/Nep98 (Conrad et al., 2007),
and Mps2 (Lee et al., 2020a). During mitosis, Mps3 and Mps2 predominantly
localise to SPB as a component of the half-bridge as well as SPB-interacting
network (SPIN) (Chen et al., 2019) and play an essential role in SPB duplication
(Jaspersen, Giddings, & Winey, 2002; Nishikawa et al., 2003). During meiotic
prophase-I, in addition to SPB, Mps3 localises as a distinct protein ensemble on
NE as a LINC complex (Conrad et al., 2007). In early meiotic prophase I, Mps3
forms several distinct foci/patches on NE and later shows more dispersed
staining around the NE. Mps3 plays an important role in the
telomere/chromosome dynamics (Conrad et al., 2007). However, how meiosis-
specific Mps3 localisation in NE is induced is poorly understood.

Here, we showed that cyclin-dependent Cdc28 kinase (CDK) and Dbf4-
dependent Cdc7 kinase (DDK) are necessary for dynamic telomere movement,
particularly for the resolution of telomere clusters. Defects in telomere movement
induced by CDK and DDK inactivation are associated with impaired NE
localisation of Mps3 and resolution of Mps3 clusters. We also found that the
luminal region of Mps3, located near the INM, is important for NE localisation.
This region is subject to phosphorylation during prophase I in meiosis. Phospho-
defective *mps3* mutant protein showed reduced NE localisation during meiosis.
The introduction of negatively charged residues in this region suppresses the
defects in the Mps3 resolution in *cdk* and *ddk* mutants. In reconstituted liposomes,
an Mps3 peptide containing the luminal region with negatively charged residues
showed reduced affinity to the lipid membrane relative to the control wild-type
Mps3 peptide. Our results suggest that multiple phosphorylation events, such as
CDK and DDK-dependent phosphorylation as well as phosphorylation in the
luminal region, control the Mps3 localisation to NE, thereby also controlling the
assembly of meiosis-specific canonical LINC complex for chromosome motion
and NE remodelling during meiosis.

Results

Force-independent localization of Mps3 on NE during meiosis

We examined the localisation of the Mps3 fusion protein with a green fluorescent protein (GFP), Mps3-GFP (Conrad et al., 2007), in various mutants. Mps3-GFP showed stage-specific distribution and movement in NE during meiotic prophase I (Figure 1A, 1B). At 0 h, Mps3 predominantly localised as a single focus, probably on SPB (Figure 1A) (Jaspersen et al., 2002; Nishikawa et al., 2003) with little movement (Figure 1C; Video 1). During early prophase I, including meiotic S phase (2–3 h), Mps3 formed 2–5 foci (including 1 focus in SPB) on NE with slow movement. After 3 h, the number of Mps3 foci increased to more than 5, and several Mps3 foci fused to form a patch (Figure 1A-C; Videos 2-4), as shown previously (Conrad et al., 2007). During this stage, Mps3 foci/patches moved around the NE with oscillatory motion (Figure 1C, Figure S1A, B). At 3–5 h, some Mps3 foci/patches were localised to one area of NE with clustering. In late prophase I (e.g. at 5 and 6 h), Mps3 often covered most of the NE and was associated with NE deformation and protrusion (Figure 1C; Videos 3, 4). Mps3 localisation was classified into four classes and quantified at each time point (Figure 1B). After 7 h, most Mps3 signals on NE disappeared, leaving two SPB-associated foci (Figure 1A).

Although Mps3 movement is dependent on Cms4 (Conrad et al., 2007; Kosaka et al., 2008; Wanat et al., 2008) and partly on Ndj1 (Chua & Roeder, 1997; Conrad et al., 1997), Mps3 localisation on NE did not depend on Ndj1 and Csm4 (Figure 1D). At late times, such as 5 h, the *ndj1* mutant showed full coverage of Mps3 on NE (Conrad et al., 2007) with NE deformation (distorted nucleus) and motion. The *csm4* mutant formed multiple Mps3 patches/foci on NE (Conrad et al., 2007; Kosaka et al., 2008; Wanat et al., 2008), but did not show full coverage of NE. The *csm4* mutant is also defective in clustering and motion of Mps3 foci/patches, as well as NE deformation with a round nucleus (Figure 1D). Moreover, we examined the effect of an actin polymerisation inhibitor, latrunculin B (LatB), on Mps3 localization (Trelles-Sticken, Adelfalk, Loidl, & Scherthan, 2005). Treatment of meiotic cells with LatB, which disrupted the cytoplasmic actin cables but not the actin patches (Figure S1C), showed normal

distribution of Mps3 foci/patches on NE during prophase (Figure 1D). However,
as shown previously (Trelles-Sticken et al., 2005), the dynamic movement of
Mps3-GFP was largely absent (Figures 1D, S1B, S1D; Video 5) without the
deformation of nuclei. These defects were similar to those observed in the *csm4*
mutant (Figure S1B, D). One hour after washing out LatB, the motion of Mps3
resumed (Figure 1D). These results suggest that Mps3 dynamics during meiosis
can be functionally separated into two processes: force-independent NE
localisation and force-dependent motion. Force-dependent motion requires actin
polymerisation, Csm4, and a party, Ndj1. On the other hand, force-independent
NE localisation does not require action polymerisation or Csm4 (Ndj1),
suggesting a unique localisation mechanism of Mps3 on NE during meiosis.
194
CDK and DDK promote proper Mps3 localization on NE during meiosis
Because Mps3-dependent telomere movement is initiated in early meiotic
prophase (Conrad et al., 2008; Conrad et al., 2007), we analysed the role of two
protein kinases that are active during early meiosis: CDK and DDK. Because
CDK and DDK are essential for vegetative growth, we used conditional mutant
alleles of the catalytic subunits of CDK and DDK (Cdc28 and Cdc7), *cdc28-as1*,
and *cdc7-as3*, which are sensitive to specific inhibitors,1 NM-PP1 and PP1,
respectively (Benjamin, Zhang, Shokat, & Herskowitz, 2003; Wan, Zhang, Shokat,
& Hollingsworth, 2006). Although CDK and DDK are also essential for DNA
replication and recombination in meiosis (Benjamin et al., 2003; Sasanuma et al.,
2008), telomere movement during meiosis, Mps3 localisation, is independent of
replication and recombination (Conrad et al., 2007; Trelles-Sticken et al., 1999).
We performed time-lapse analysis of Mps3-GFP in *cdc28-as1* and *cdc7-as3*
in the absence or presence of each specific inhibitor (Figure 1E). To inactivate
CDK, 1 NM-PP1, which did not affect Mps3-GFP dynamics in wild-type cells or a
steady-state level of Mps3-Flag protein in *cdc28-as1* cells (Figure S1E, F), was
added to *cdc28-as1* at 0h. In the absence of the inhibitor, the mutant cells showed
normal Mps3 localisation during meiosis (Figure 1E). CDK inactivation impaired
Mps3 localisation and motion (Figure 1B, E). In the wild type, several Mps3
foci/patches are loosely clustered on one area of NE in a transient manner, which
reflects telomere bouquet (Conrad et al., 2007). With CDK inhibition, these loose

clusters of Mps3 foci were persistent during prophase I, which largely reduced
the spread of Mps3 foci on NE (Figure 1B, G). Tracking the motion of Mps3 foci
confirmed persistent clustering with restricted motion of the foci in one region of
NE under these conditions (Figure 1F).

We also measured the speed of Mps3 GFP focus/patches on NE (Figure
1H). Previous measurement of the motion of a single chromosome locus showed
heterologous nature of chromosome motion (Conrad et al., 2008). To avoid
biased results, we simply measured a step-size of a single Mps3-GFP focus or
patch in one-second interval through 20 seconds in multiple nuclei, and
measured an average velocity a maximum velocity during the measurement
(Figure 1H). Wild type Mps3-GFP at 5 h showed 0.19 and 0.25 μm/s in average
and maximum speeds, respectively. CDK inactivation slightly decreased the
average speed, but not the maximum speed (Figure 1H, Video 6).

Importantly, 2 h after washing the inhibitor, NE localisation and Mps3 motion
were recovered. CDK inactivation after meiotic S phase (the addition of the
inhibitor at 3 h) interfered with the dispersed localisation of Mps3 throughout NE
with loose clustering of Mps3 foci/patches (Figure 1E). Therefore, persistent CDK
activity is necessary for the establishment and maintenance of the dispersed
localisation of Mps3, and thereby the resolution of clustering. We concluded that
CDK plays a role in the resolution of Mps3 focus clustering (probably, not in
focus/patch formation *per se*) and a minor role in the motion of Mps3 on NE.

DDK inactivation (*cdc7-as3* + PP1) also induced a defect in Mps3
localisation on NE and motion (Figure 1B, E). PP1 did not affect the localisation
of Mps3-GFP in wild-type cells or a steady level of Mps3-Flag protein under DDK
inactivation conditions (Figure S1F, G). In its absence, *cdc7-as3* cells showed
normal localisation and motion of Mps3 foci/patches (Figure 1B, E). DDK
inhibition by PP1 from 0 h reduced the NE localisation of Mps3 with accumulation
of the loose cluster of the foci (Figure 1E, F). Removal of the inhibitor restored
the NE localisation of Mps3 in *cdc7-as3* cells (Figure 1E). The addition of PP1 at
3 h also induced a Mps3 localisation defect similar to that at 0 h. This shows that
persistent DDK activity is critical for the efficient resolution of Mps3 clusters on
NE. The resolution defect of Mps3 clusters in reduced DDK activity is similar to
that in response to decreased CDK activity, suggesting that CDK and DDK work

in the same pathway to control Mps3 dynamics during meiosis. In contrast to
reduced CDK, reduced DDK activity clearly decreased the velocity of Mps3 foci
on NE (Figure 1H, Video 7). Therefore, DDK appears to play a more critical role
in Mps3 motion than CDK.
We also analysed the Mps3-GFP localisation in the *cdc7 bob-1-1* double
mutant, in which the *bob1-1* mutation in the helicase gene *MCM5* suppresses the
lethality of the *cdc7* deletion (Jackson, Pahl, Harrison, Rosamond, & Sclafani,
1993; Sasanuma et al., 2008), A small fraction (~20%) of the *cdc7 bob-1* mutant
cells showed a few Mps3 foci on NE, whereas most of the cells did not form some
Mps3 foci on NE (Figure 1B, E, F). The velocity of Mps3 foci formed in the
complete absence of DDK was highly reduced and was similar to that in the *csm4*
mutant and in the presence of LatB (Figures 1H, S1D; Video 8). These defects in
the *cdc7 bob-1-1* double mutant were more severe than those in the *cdc7-as3*
mutant with PP1, confirming the importance of DDK in Mps3 localisation and
motion in NE.
Whole cell immunostaining of the Mps3-Flag and Ndj1-HA proteins showed
that reduced CDK and DDK activities hampered the localisation of Mps3, but not
of Ndj1, to NE during meiosis (Figure S1H), suggesting a critical role for the
kinases in the coupling of Mps3 to Ndj1 on NE. Double-staining of Mps3 and Ndj1
in wild-type meiosis showed distinct Mps3- and Ndj1-staining domains on NE
together with colocalized foci/patches of both. The reduced colocalization of Ndj1
and Mps3 on NE is clearly seen under CDK and DDK inactivation conditions.
Ndj1 was not required for Mps3 localisation on the NE (Figure 1D). These results
suggest the presence of an Mps3-independent pathway for Ndj1 localisation on
NE or telomeres, probably mediated by Ndj1 binding to Rap1/telomeres. Taken
together, these results revealed that CDK and DDK activities are necessary for
proper Mps3 localisation and efficient motion in NE during meiosis.
276

CDK and DDK promote telomere dynamics during meiosis
To determine the role of CDK and DDK in telomere movement, we analysed the
motion of GFP fusion of a telomere-binding protein, Rap1, as described
previously (Trelles-Sticken et al., 2005). During vegetative growth or pre-meiosis,
Rap1-GFP exhibited 1–4 foci on NE on a single focal plane with little movement

(Figure 2A, B; Video 9) (Trelles-Sticken et al., 2005). Once the cells entered

meiosis, Rap1 foci on NE increased in number (up to 10). Rap1-GFP foci moved

rapidly and were occasionally clustered in one area of NE (Figure 2A, B; Video

10), which corresponds to the telomere bouquet (Trelles-Sticken et al., 2005).

This telomere clustering was very transient (half-life less than 5 s) and was

predominantly observed after 3–4 h of incubation in sporulation medium (Figure

2A, D). Rap1 foci were more dispersed during late meiotic prophase (4–6 h for

the pachytene stage) before the onset of meiosis I (MI; Figure 2D). Tracking

analysis showed that Rap1 foci move around the meiotic nucleus in an

uncoordinated manner (Figure 2B).

CDK inactivation (*cdc28-as1* + 1 NM-PP1) impaired meiotic Rap1

dynamics (Figure 2B, E). With compromised CDK activity from 0 h, cells with

clustered Rap1 foci accumulated during prophase I (Figure 2D, E; Video 11),

indicating that CDK activity is required for meiotic telomere dynamics, particularly

for the resolution of telomere clusters (but not necessarily for clustering *per se*).

The 2D tracking confirmed the restricted motion of Rap1 foci in one nuclear area

(Figure 2B). The addition of the inhibitor during early prophase I (e.g. at 3 h) also

resulted in defective resolution of telomere clustering; washing off the inhibitor

restored telomere dynamics (Figure 2E). Treatment of wild-type cells with the

inhibitor did not affect Rap1-GFP dynamics, confirming the specificity of the

inhibitor (Figure S1I). These results indicate that persistent CDK activity during

meiotic prophase I promotes normal telomere dynamics. CDK inactivation of CDK

does not affect the formation of cytoplasmic actin cables during meiosis (Figure

S1C).

Similar to CDK inactivation, Cdc7 inactivation (the treatment of *cdc7-as3*

with PP1) at 0 and 3 h impaired Rap1-GFP dynamics, which showed

accumulation of Rap1 clusters with restricted motion (Figure 2B-D, F; Video 12);

however, treatment of wild-type cells with PP1 did not affect Rap1 dynamics

(Figure S1I). This defective resolution of the Rap1 cluster under DDK inhibition

was similar to that observed in CDK inhibition. Moreover, the *cdc7* null mutant

with *bob1-1* mutation exhibited persistent telomere clustering (Figures 2B, D, F,

and S13). These results indicate that persistent Cdc7 kinase activity is also

required for meiotic telomere dynamics. Although CDK and DDK inactivation

restricted Rap1 foci to one area, inactivation did not affect the velocity of Rap1

foci (Figure 2C). Since Mps3 foci are actually formed under CDK and DDK

inactivation conditions (Figure 1), the Mps3 complexes on NE are sufficient to

ensure some Rap1 motion. Alternatively, the Mps3-independent mechanism may

promote motion under reduced CDK or DDK activities, particularly in the *cdc7*

null mutant.

Mps3 is phosphorylated during meiosis

Previously, it was shown that S70 of SPB-associated Mps3 is phosphorylated in

late prophase-I (Li et al., 2017) (Figure 3A). We also found that Mps3 is a

phosphoprotein in middle prophase I (Figure 3). On immunoprecipitation (IP)

western blots, Mps3-Flag showed a mobility shift, which was induced during

meiotic prophase I (4 and 6 h; Figure 3B, left panel). Mps3 mobility in the IP

fractions was reduced by treatment with lambda phosphatase (Figure 3C).

Moreover, IP fractions of Mps3-Flag from meiotic cells showed that Mps3 cross-

reacted with anti-phosphoserine antibody (Figure 3D), whose reactivity was

largely diminished by treatment with phosphatase.

Previous systematic analyses of CDK substrates *in vitro* identified Mps3

as a Cdc28–Clb2 substrate (Ubersax et al., 2003). Mps3 contains six potential

CDK sites (S/TPXK/R or S/TP) located in the luminal region, two of which seem

to be candidates for CDK/DDK phosphorylation (188–190 <u>TSS</u>PGK and 450–451

<u>TS</u>P; Figure 3A, E). Because the 450–451 site is located in the SUN domain,

which is not essential for Mps3 localisation on NE (Rao, Shinohara, & Shinohara,

2011), we focused on T188-S189-S190 (hereafter, TSS; Figure 3E). We purified

a GST fusion protein of an Mps3 fragment (residues 160–260), including TSS. *In*

vitro Cdc28–Clb5 alone, but not Cdc7–Dbf4 alone, promoted phosphorylation of

the Mps3 fragment (approximately 10-fold compared to the background; Figure

3F). Co-incubation with CDK and DDK resulted in robust incorporation of ^{32}P in

GST-Mps3, indicating that CDK and DDK cooperate in Mps3 phosphorylation *in*

vitro (~30-fold compared to the background). Triple alanine substitutions (T188A,

S189A, S190A; referred to as the *mps3-AAA* mutant; Figure 3E) made the GST-

Mps3 fragment a poor substrate by these kinases. These results indicate that

TSS is required for efficient CDK/DDK phosphorylation *in vitro*.

Consistent with the *in vitro* results, the Mps3-AAA mutant protein showed a reduced band shift (Figure 3B, G) and decreased reactivity to anti-phosphoserine antibody (Figure 3H). Furthermore, we raised an antibody against a Mps3 peptide with 189-phosphoserine and 190-phosphoserine (186-GAT**pSpS**PGKSF-195) and found that this phospho-specific antibody could recognise immunoprecipitated Mps3-Flag specifically between 4 and 8 h in meiosis, but not in mitosis (Figure 3C, left panel). Strong signals were observed during late prophase I (4 h and 6 h). This antibody did not recognise precipitated Mps3-AAA-Flag (Figure 3C, middle panel). These results indicate that 189S and 190S are phosphorylated *in vivo*, specifically during meiosis.

The Mps3 sequence around the TSS site is similar to that of CDK/DDK-catalysed phosphorylation sites in Mer2 (Henderson, Kee, Maleki, Santini, & Keeney, 2006; Sasanuma et al., 2008; Wan et al., 2008), including the TSSP sequence (Figure 3E). Mer2 is a meiosis-specific protein essential for DSB formation (Cool & Malone, 1992; Engebrecht, Hirsch, & Roeder, 1990). CDK and DDK sequentially phosphorylate the TSS sequence of Mer2, which is essential for meiotic DSB formation (Sasanuma et al., 2008; Wan et al., 2008). We swapped a 10-amino-acid region (GATSSPGKSF) containing the Mps3 phosphorylation sites with the corresponding region of Mer2 (TETSSPFRST; Figure 3E). This *MPS3* allele (*MPS3-MER2S*) conferred wild-type spore viability (95%; n=50 tetrads; Figure S2A). Importantly, the Mps3-MER2S protein cross-reacted with an antibody against the Mer2 phospho-S30 antibody (Henderson *et al.* 2006) (Figure 3I), but not with the anti-Mps3-phospho antibody (Figure 3B, right panel). *MPS3-MER2S-Flag* exhibited similar telomere (Rap1) clustering to the wild type (Figure 3J, K). The Mps3-MER2S-Flag protein showed meiosis-specific localisation on NE which often co-localized with Ndj1 (Figure 3L). This suggests that residues 26-35 of Mer2 are functionally equivalent to residues 185-194 of Mps3. Rather, the sequence of Mps3, the composition of amino acids, seems to be critical for NE localisation. Similar to Mps3, the region of Mer2 is rich in serine and threonine (Figure 3E).

The *mps3-AAA* mutant is defective in Mps3 localization on NE

To investigate the role of the phosphorylated residues of Mps3, we constructed

and characterised the phenotypes of the *mps3-AAA-FLAG* mutant (hereafter,

mps3-AAA; Figures 3E, 4). The *mps3-AAA* mutant showed few defects in mitosis,

such as SPB duplication (Figure S2B) seen in other *mps3* mutants (Jaspersen et

al., 2002; Nishikawa et al., 2003), suggesting that the TSS residues of Mps3 are

not critical for the mitotic function of Mps3. The *mps3-AAA* mutant showed

defects in meiosis with a reduced spore viability of 87.5% (Figure 4A), which was

slightly higher than that of the *ndj1* mutant (75.8%) (Chua & Roeder, 1997;

Conrad et al., 1997). The spore viability of the *ndj1 mps3-AAA* double mutant was

72.3%, which is similar to that of the *ndj1* mutant. The *mps3-AAA* mutant delayed

entry into meiosis I by ~1 h relative to the wild type (Figure 4A). The *ndj1 mps3-*

AAA double mutant showed ~2.5 h delayed entry into meiosis I, which is similar

to the *ndj1* single mutant, indicating the epistatic relationship of *mps3-AAA* to *ndj1*.

The *mps3-AAA* mutant showed normal progression of the meiotic S-phase

(Figure S2C). Immunostaining of Zip1, a component of the central region of SC

(Sym, Engebrecht, & Roeder, 1993), showed a delay in both the loading of Zip1

and the formation of full-length SCs in the *mps3-AAA* mutant (Figure S2D, E),

suggesting that Mps3 NE localisation is necessary for timely synapsis.

We also analysed chromosome dynamics during meiosis in *mps3-AAA*

mutant cells. First, we checked the Rap1-GFP movement in the *mps3-AAA*

mutant. In contrast to wild-type cells, the *mps3-AAA* mutant was defective in the

localisation and motion of Rap1-GFP (Figure 4B, C). In the mutant, the number

of Rap1 foci did not increase (Figure 4B), and little movement of Rap1-GFP foci

was observed during meiosis (Figure 4C). Rap1 motion in meiotic prophase I of

the *mps3-AAA* mutant is restricted, and the velocity of each Rap1 focus is

decreased relative to the wild type (Figure 4C, D; Video 14). Therefore, we

concluded that T188, S189, and S190 of Mps3 are important for meiotic telomere

dynamics. Indeed, the *mps3-AAA* mutant was defective in chromosome motion,

as shown by the analysis of a GFP fusion of Zip1 (Koszul et al., 2008; White,

Cowan, Cande, & Kaback, 2004). During prophase I, Zip1-GFP showed dynamic

motion (Figure 4E; Videos S15, 16). The *mps3-AAA* mutant reduced the velocity

of Zip1 motion by approximately 3-fold relative to that of the wild type (48±15

versus 130±31 nm/s, Figure 4F).

Next, we examined the localisation of the Mps3-AAA mutant protein in

meiosis as a GFP fusion protein (Figure 4G). *mps3-AAA-GFP* cells showed 85%

spore viability, which was comparable to the mutant strain without the GFP tag.

Similar to the wild-type Mps3-GFP, the Mps3-AAA GFP fusion protein normally

resided on the SPB as a single focus during mitosis (Figure 4G). During meiosis,

the Mps3-AAA mutant protein showed reduced NE localisation. Upon entry into

meiosis, similar to the wild-type Mps3, the Mps3-AAA mutant protein formed a

few (usually two or three small) foci on NE, but showed little increase in the

number of Mps3 foci. At late time points, unlike wild-type Mps3, the mutant protein

did not form an Mps3 patch or cover on NE (Figure 4G, H, I). At 6 h, two or three

foci of Mps3 showed limited movement (Figure 4J, K, L; Video 17). These results

suggest that T188, S189, and S190 of Mps3 are necessary for meiosis-specific

Mps3 localisation and motion on NE, particularly for efficient localisation on NE

accompanied by the formation of large complexes. Interestingly, the Mps3-AAA-

Flag mutant protein still bound to Ndj1 and Csm4, similar to the wild-type Mps3

protein (Figure S2F), which is consistent with the Mps3-AAA protein retaining the

N-terminal nucleoplasmic region and SUN domain necessary for Ndj1 and Csm4

binding, respectively.

432   The *mps3-AAA* mutant is deficient in NE growth

In addition to localization/movement defects, the *mps3-AAA* mutant showed

defects in nuclear morphology. First, the meiotic nucleus of the *mps3-AAA* mutant

was smaller than that of the wild type (Figure 4B). The nucleus of the wild type

grew when the cell underwent meiosis and exhibited a 4-fold increase (in area; 2

folds in length) at 4 h (Figure 4M). Treatment with LatB did not affect the

expansion, indicating that meiosis-specific NE remodelling does not require actin

polymerisation. The *mps3-AAA* mutant showed defective NE growth in prophase

I; the nuclear area in the *mps3-AAA* mutant was similar between 0 and 4 h (Figure

4M). These results indicate that the Mps3 luminal region plays a critical role in

the growth of the meiotic nucleus, possibly by controlling NE remodelling.

Second, a fraction of the *mps3-AAA* mutant cells showed a partial defect

in telomere tethering to NE during meiosis. In one or two Rap1-GFP foci, the

mps3-AAA mutant (40%; n=100) was localised to the nucleoplasm rather than to

the nuclear periphery, whereas all Rap1-GFP foci were localised near the NE in

the wild type (Figure 4B). The weak telomere tethering defect in the *mps3-AAA*

mutant is reminiscent of the defect observed with an *NDJ1* mutant, which also

showed a partial tethering defect (Conrad et al., 1997; Trelles-Sticken et al.,

2000).

Aspartate substitution of TSS in Mps3 overcomes the CDK and DDK defects

We constructed a version of Mps3 with three negative charges on the TSS

sequence (T188D, S189D, and S190D), referred to as *MPS3-DDD* (Figure 3E).

The *MPS3-DDD* allele did not show any mitotic defects, such as colocalization

with tubulin and tubulin elongation (Figure S3A-C). The *MPS3-DDD* cells showed

wild-type spore viability (95%) and entered MI similar to wild-type cells (Figure

S3D). The *MPS3-DDD* cells also displayed similar Rap1-GFP dynamics to wild-

type cells with normal clustering during prophase I (Figure 5A, B). The Mps3-

DDD-GFP protein localised to SPB during vegetative growth (Figure S3A) and

showed wild-type-like localisation on NE during meiosis (Figures 5C, D, and S3E).

Mps3-DDD-GFP displayed movement similar to that of the wild-type protein

(Figure 5C, E, F; Video 18). The aspartate substitutions of Mps3 suppressed the

resolution defect in Rap1 clustering at late time points of meiosis by CDK and

DDK inactivation (Figure 5A, B), although some clustering persisted under these

conditions (Figure 5B). Mps3-DDD-GFP showed NE localisation (Figure 5D) and

movement during meiosis even under the conditions of either CDK or DDK

inactivation (Figure 5E, F; Videos 19-22), which is similar to that in wild-type

Mps3-GFP or Mps3-DDD-GFP cells. Clustering of Mps3 induced by CDK/DDK

inactivation was clearly ameliorated by the introduction of Mps3-DDD-GFP

(Figure 5C, E; Videos 23-25). DDD substitution also suppressed the motion

defects of Mps3 by DDK inactivation (Figure 5F). Importantly, NE localisation and

dynamics of Mps3-DDD protein still depend on meiosis, suggesting the presence

of additional regulation of meiosis-specific localisation of Mps3. In vegetative cells

(0 h, Figure S3E), Mps3-DDD showed SPB localisation and normal Rap1

localisation (Figures 5G, S3F). Surprisingly, Mps3-DDD, but not wild-type Mps3,

showed dispersed staining on NE with SPB in mitotic cells only in the *cdc7 bob-*

1 mutant with, but not in *cdc7as3* with the inhibitor (Figures 5H, S3G). This

suggests that, in contrast to meiosis, DDK negatively regulates the NE

localisation of Mps3 in vegetative cells.

The *mps3-S189A* and *mps3-S190A* mutants phenocopy the *mps3-AAA*

mutant

Although the *MPS3-DDD* mutation suppressed Mps3 localisation defects induced

by CDK or DDK inactivation, Mps3 localisation defects in the *mps3-AAA* mutant

(reduced localisation) are different from those under CDK/DDK inactivation

(persistent clustering). To clarify the relationship between the phosphorylation of

the luminal region of Mps3 with CDK and DDK, we created additional mutants in

the Mps3 TSS sequence, called *mps3-S189A and -S190A*, which has a single

alanine substitution of S189 and S190, respectively (Figure 6A), which are

phosphorylated *in vivo* (Figure 3). If CDK could phosphorylate S190 as the CDK

site, *mps3-S190A* would phenocopy the CDK defect in Mps3 localisation. And

also, if DDK could phosphorylate S189 as the DDK site, *mps3-S189A* would

phenocopy the DDK defect in Mps3 localisation. The localisation of GFP fusion

of Mps3-S189A and -S190A proteins was analysed as described above (Figure

6B). Both Mps3-S189A and -S190A proteins showed reduced NE localisation

compared to wild-type Mps3. In the localisation defects, both mutants were

weaker than the *mps3-AAA* mutant (Figure 6D). The *mps3-S189A* mutant was

more severe than the *mps3-S190A* mutant. Both *mps3* mutants also showed

reduced Mps3 motion (Figure 6E, F) and were also partially deficient in meiosis-

induced nuclear expansion (Figure 6G). The *mps3-S190A* mutant exhibited

weaker defects than the *mps3-S190A* mutant. This suggests that, although

weaker, the *mps3-S189A* and *-S190A* mutants shared similar defects to the

mps3-AAA mutant and were different from *cdk* or *ddk* mutants. Taken together,

these results suggest that CDK and DDK phosphorylation sites are not S189 or

S190 (see Discussion).

Luminal region of Mps3 binds to lipid bilayers *in vitro*

To determine the role of putative phosphorylation of the Mps3 luminal region, we

analysed the biophysical properties of a 55 amino acid Mps3 peptide (153-208

aa) containing both transmembrane (TM) and luminal regions (Figure 3A). The

luminal region is referred to as the juxtamembrane (JM) region (Figure 7A). We

prepared two peptides, Mps3-wild type (WT) and Mps3-DDD, which contain three aspartates (Figure 7A). Synthesised Mps3 peptides were mixed with liposomes comprising POPC/POPS (10/3)(Sato, Pallavi, Golebiewska, McLaughlin, & Smith, 2006) First, we checked the conformation of each peptide in the bilayers by polarised Fourier-transform infrared spectroscopy (FT-IR) and found that both Mps3-WT and Mps3-DDD peptides contained similar contents of α-helices and β-strands (Figure S4A). Dichroic ratio calculated for Mps3-WT and Mps3-DDD were 2.90 and 2.85, suggesting that both transmembrane regions are inserted into lipid bilayers with a similar tilt angle of ~30° relative to the membrane normal (Tamagaki et al., 2014). We cannot assign those peaks from the β-strand to which residues in the JM regions are involved in the β-structure with this FT-IR experiment alone.

Next, we synthesised Mps3 TM-JM peptide with Alexa fluorophore-dye (Alexa-568) at the C-terminus of the JM region and analysed the spectrum of the dye in the presence of different concentrations of negatively charged phosphatidyl-inositol 4,5-bisphosphate (PIP$_2$), which modulates the status of charge on the membrane surface (Sato et al., 2009). Here, we utilised a PIP$_2$-involving model system to determine whether JM is affected by the membrane, although we are aware of the fact that it is not known whether Msp3 interacts with PIP$_2$. This simple model system must provide a possible mechanism of how negative charge on the JM region caused by its phosphorylation affects its behaviour against the membrane.

For the wild-type peptide, the addition of PIP$_2$ decreased the peak intensity of the fluorophore at 600 nm up to ~40% (Figure 7B). The triplicates showed similar results. Decreased fluorescence could be due to quenching of the fluorescence from the dye association (self-quenching) by inter-peptide association (Figure 7C), as shown in a previous study on an ErbB2 peptide containing TM and JM regions (Matsushita et al., 2013). These results suggest that the JM region of the Mps3-WT peptide is sensitive to changes in the membrane component and that the JM regions interact with the acidic lipid bilayer. On the other hand, the Mps3-DDD peptide decreased the intensity to ~20% (half of the intensity from the WT peptide). The decrease in the change in the fluorescence intensity was due to electrostatic repulsion between the aspartates

and acidic lipids (Figure 7C). This repulsion must place the fluorophores apart,
preventing self-quenching. The repulsion can reduce the JM region of the Mps3-
DDD peptide binding to the bilayers compared to that of Mps3-WT. These results
suggest the different conformations of Mps3-WT and -DDD peptides with respect
to the interaction of the JM region of Mps3 with the membrane.

To gain more insight into the behaviour of the JM region with the
membrane, we utilised a molecular dynamic (MD) simulation technique. Since
there is no reported structure on the TM-JM region of Mps3, we artificially
constructed the initial structure of the TM-JM sequence as follows: using the
software *Chimaera* (Pettersen et al., 2004), R161-M181 is set to be α-helix as
the TM region; the rest was set to a random structure. This constructed structure
was embedded in a membrane (POPC/POPS/PIP$_2$=8:2:1 with 80 nm in XY
dimension) using *the Membrane Builder* module in *CHARMM-GUI* (Jo, Kim, Iyer,
& Im, 2008). The peptide-membrane system was fully hydrated in a 150 mM NaCl
solution.

First, we performed coarse-grained molecular dynamics simulations of the
constructed TM-JM peptide in the membrane. We ran simulations on two
sequences, Mps3-WT and Mps3-DDD. The duration of each run was >3 μs, and
three sets were performed for each sequence. The results from all the
calculations showed that the JM region from Mps3-DDD dissociated from the
membrane (Figure S4B). Conversely, that of Mps3-WT remained attached to the
membrane (Figure S4B). Second, we performed all-atom simulations on three
TM-JM sequence peptides, Mps3-WT, Mps3-DDD, and Mps3-WT
phosphorylated at T188-S189-S190. The duration of each calculation was >50
ns (Videos 26-28). In Figure 7D, snapshots from trajectories for the three
sequences are shown. Results from the simulation, together with mass density
profiles (Figure S4C), indicate that JM regions from Mps3-DDD and
phosphorylated Mps3 dissociated from the membrane. On the other hand, the
JM region from Mps3-WT remained attached to the membrane.

Discussion

Here, we describe the multi-layered control of the dynamics of an SUN-domain

protein, Mps3, during yeast meiosis. In both mitotic and meiotic cells, Mps3 is a

major component of a half-bridge as well as the SPB interacting network (SPIN)

(Chen et al., 2019; Lee et al., 2020a) (Li et al., 2017). Meiosis induces the

localisation and motion of Mps3 on NE, which is accompanied by the formation

of large protein ensembles of the LINC complex in NE, detected as foci/patches.

The meiosis-specific LINC complex tethers telomeres and interacts with

cytoplasmic actin cables (Lee et al., 2020b). In the luminal region between INM

and ONM, Mps3 binds to a KASH protein, Mps2, during both mitosis and meiosis,

and to a meiosis-specific Mps2-binding protein, Csm4 (Chen et al., 2019; Lee et

al., 2020a). Mps2 and Csm4 cooperate to bind to the motor Myo2 in the

cytoplasm. Myo2 on actin cables, whose formation is induced in meiotic cells,

generates forces for the motion of the LINC complex on NE, which in turn moves

chromosomes inside of meiotic cells (rapid prophase movement; RPM) as well

as the clustering of telomeres in the zygotene stage (Conrad et al., 2008; Koszul

et al., 2008; Trelles-Sticken et al., 2000; Trelles-Sticken et al., 1999). While Mps3

localisation on NE is independent of actin-generated force (and other meiosis-

specific factors, such as Ndj1 and Csm4), Mps3 movement on NE is dependent

on the force. Meiosis-specific regulation of Mps3 localisation on NE is a key event

in the formation of a LINC complex with Mps3 for chromosome pairing/synapsis

and chromosome motion.

Our studies revealed that, in addition to its role in chromosome dynamics,

Mps3 localisation in NE is important for NE remodelling during meiosis. The

characterisation of the *mps3-AAA* and *mps3-S189A* and *-S190A* mutants

revealed that Mps3 localisation to NE (NE remodelling) plays a unique role in NE

expansion in meiotic cells. NE-bound Mps3 may regulate NE growth by directly

promoting lipid synthesis on NE or by importing lipids from the ER (Sosa Ponce,

Moradi-Fard, Zaremberg, & Cobb, 2020).

Localisation of Mps3 on NE in meiotic cells is divided into several distinct

steps, which are coupled with distinct regulatory mechanisms (Figures 1A and

8A); 1-At early times in prophase I, in addition to the SPB, a few Mps3 foci appear

on NE. 2-With increased numbers of foci, patches of Mps3, large cohesive

ensembles, probably with multiple foci, are formed on NE, which shows transient
clustering in one area of NE. 3-Multiple foci and patches cover the entire NE. 4-
Mps3 shows nearly full coverage of NE, sometimes with a hole at the nucleus-
vacuole junction (NVJ). 5-During late prophase I or pro-metaphase I, most Mps3
on NE disappear from NE, leaving two SPB-associating foci. At each stage, Mps3
foci/patches show RPM which is the maximum at mid/late prophase I (Conrad et
al., 2008; Koszul et al., 2008; Trelles-Sticken et al., 2000; Trelles-Sticken et al.,
1999).

In this study, we found that two cell cycle kinases, CDK and DDK, control
proper Mps3 localisation on NE during meiosis, particularly by promoting the
resolution of clustered Mps3 foci (and, thus, telomere bouquets), suggesting that
the phosphorylation of a target protein(s) by these kinases plays a role in the
resolution of Mps3 clusters. Moreover, we showed that the JM region of Mps3
near the INM, which is located in the lumen of NEs, is critical for meiosis-specific
NE localisation of the protein. S179 and S180 of the JM region in Mps3 are
subject to meiosis-specific phosphorylation of Mps3 assembly on NEs. The
introduction of negative charges in the JM region changes the binding ability of
the region to the membrane in reconstituted liposomes. These results suggest
that Mps3 localisation in NE is subject to multi-layered regulation governed by
distinct phosphorylation.

Do CDK and DDK phosphorylate the lumen of Mps3?
During meiosis of *C. elegans*, the N-terminal nucleoplasmic region of Sun-1
protein is phosphorylated by Chk-1 and Cdk-1 (Penkner et al., 2009; Zuela &
Gruenbaum, 2016). In M phase of human cells, SUN1 protein is phosphorylated
at its N-terminal region by CDK and PLK (Patel et al., 2014). Together with results
described here, these suggest the importance of the phosphorylation in the
regulation of dynamics and/or regulation of SUN-domain proteins.

Our *in vitro* study (Figure 3F) showed that CDK and DDK phosphorylate
188-TSS-190 sequence of Mps3 in a collaborative manner, suggesting that CDK
and DDK collaborate to catalyze the phosphorylation of the JM region of Mps3
for its NE localization *in vivo*. However, this is less likely for several reasons. First,
it is unlikely that CDK and DDK are responsible for direct phosphorylation reaction

in NE lumen. These kinases are not localized to luminal regions of both

endoplasmic reticulum (ER) and NE since the catalytic subunits (Cdc28 and

Cdc7) and its regulatory subunits (Clb5,6 and Dbf4) lack a leader sequence

necessary for the luminal localization. Second, defects in Mps3 localization

induced by CDK/DDK inactivation are different from those in the *mps3-AAA*

mutant. Third, if CDK is a priming kinase on S190 in SSP sequence of Mps3, a

mutation in the CDK site (S190) could phenocopy the defect conferred by CDK

defect. But this is not the case (Figure 6). Forth, if DDK is the secondary kinase

on S-pS-P primed by CDK, S189 substitution would confer similar (or less) defect

to S190 substitution. However, the *mps3-S189A* is more defective than the *mps3-*

S190A. Rather, S190 and S189 phosphorylation seem to play an overlapping role

in Mps3 localization.

What is the target for CDK and DDK in Mps3 localization?

If CDK and DDK are unlikely to directly phosphorylate the luminal region of Mps3,

what are the target(s) of these kinases? Our results indicate that CDK and DDK

activities promote the resolution of Mps3 foci/patches and/or telomeres rather

than NE localisation *per se*. Mps3 localisation on NE seems normal, even in the

absence of CDK and DDK activities. Phenotypic similarity between *cdk* and *ddk*

mutants in Mps3 localisation suggests a common target of these kinases for the

resolution of the clusters. Interestingly, telomere-resolution defects under CDK

and DDK inactivation are similar to those in the mutant of the *REC8* gene (Challa,

Lee, Shinohara, Kim, & Shinohara, 2016; Conrad et al., 2007), which encodes a

meiosis-specific kleisin subunit of the cohesin complex (Klein et al., 1999) and

cohesin regulator, *RAD61/WPL1* (Challa et al., 2016). It is well known that Rec8

is phosphorylated by DDK (and Polo and CK1 kinases), but not by CDK (Katis et

al., 2010) and Rad61/Wpl1 is phosphorylated by DDK and Polo-like kinase (PLK)

{Challa, 2019 #221}. Thus, there might be another target in cohesin components

by the CDK-DDK kinase axis. In mitotic yeast cells, Eco1, which catalyses the

acetylation of the Smc3 cohesin subunit, is phosphorylated by CDK and DDK for

degradation (Seoane & Morgan, 2017). Interestingly, Eco1 acetylates Mps3 in

vegetative cells (Ghosh et al., 2012). Therefore, Eco1 may be a possible target

of CDK-DDK for telomere resolution.

Telomere clustering during prophase I persisted with CDK or DDK

inactivation, suggesting the presence of a constraint on the resolution of the

clustering. The phosphorylation, and thus negative charges in the JM region,

might overcome this constraint for resolution. This is supported by the fact that

MPS3-DDD partially rescued the resolution defect of telomere clustering with

CDK or DDK inactivation (Figure 4). Moreover, we found that Mps3-DDD, but not

wild-type Mps3, showed NE localisation in mitotic cells only in the *cdc7 bob-1*

mutant (Figure 5C), suggesting the presence of DDK-dependent constraints on

the NE localisation of Mps3 during mitosis (Figure 8A).

Phosphorylation in the NE luminal region of Mps3

Protein phosphorylation of the luminal region between INM and ONM is very

unusual and has not been reported in the past. However, recent studies have

shown that the phosphorylation of secreted proteins and oligosaccharides of

membrane proteins is catalysed by an atypical kinase, such as Family with

sequence similarity 20, member C (Fam20C), in the lumen of the Golgi apparatus

and ER in vertebrate cells (Tagliabracci et al., 2012; Tagliabracci, Pinna, & Dixon,

2013; Tagliabracci et al., 2015). Given that the luminal region of NE is consistent

with that of ER, where INM/ONM proteins are synthesised, it is possible that a

Fam20C family-like kinase might be present in the lumen between INM and ONM,

which may catalyse the phosphorylation of the NE luminal region of Mps3.

However, there are no reports on the presence of Fam20C orthologues or relative

enzymes in lower eukaryotes, such as yeasts. Alternatively, phosphorylation of

the luminal region may be mediated by a novel mechanism. Given that

membranes are rich in phospho-lipids, the phosphate group in the lipids might

act as a donor for transfer to proteins. Further studies are needed to identify the

protein responsible for the phosphorylation of the JM region of Mps3.

Th role of negative charges in JM region of Mps3

Biophysical analysis of synthetic peptides containing TM and JM regions of Mps3

in synthetic liposomes revealed that the JM region of Mps3 has the capability to

bind to the lipid bilayers. Importantly, the introduction of three negative charges

into the JM region weakens the binding to the membrane, resulting in a change

in the geometry of the JM region of Mps3, as an extension, the luminal region of

Mps3, including the SUN domain. As a typical LINC complex, the SUN domain of

Mps3 in INM binds to the KASH domain of Msp2 in ONM. The connection spans

~20 nm between the INM and ONM (Chen et al., 2019; Lee et al., 2020a).

Recently, Jasperson and her colleagues showed "atypical" configuration of Mps3-

Mps2 in SPB, in which not only SUN-KASH interaction, but also N-terminal

regions of both Mps3 and Mps2 bind to each other in SPIN which surrounds SPB

(Chen et al., 2019; Lee et al., 2020a). Typical LINC could accommodate *trans*-

membrane configuration in the luminal region, while atypical LINC would have *a*

cis-membrane configuration (Figure 8B). This suggests that the luminal regions

of Mps3 and Mps2 are quite flexible in accommodating two different SUN-KASH

configurations. Our results suggest that the switch from *cis*- to *trans*-membrane

configurations of the LINC complex might be regulated by the binding of the JM

region of Mps3 to the INM. While the interaction between the JM region of Mps3

and INM promotes the *cis*-membrane form of the noncanonical LINC complex,

negative charges induced by phosphorylation may reduce the interaction

between JM and INM, which may accommodate a conformation suitable for the

trans-membrane form. Alternatively, the negatively charged JM region of Mps3

would be a binding site for the other protein which may promote the configuration

switch. This idea should be tested in future studies.

Meiosis establishes Mps3 localization on NE

As described above, Mps3 localisation in NE is regulated at distinct stages

(Figure 8). The *mps3-AAA* mutant retains the ability to form early meiosis-specific

Mps3 foci on NE. The Mps3-DDD protein also requires meiosis for NE localisation

during meiosis. This suggests that Mps3 localisation on NE is initiated and/or

established during very early meiotic prophase I. Given that Mps3 localisation as

an ensemble on NE is specific to meiosis, there might be an early meiosis-specific

factor or modification that regulates the establishment of Mps3 localisation on NE

upon entry into meiosis. We examined two meiosis-specific factors, Ndj1 and

Csm4, both of which are involved in chromosome motion and Mps3 dynamics,

but it is unlikely that these factors function in the establishment of Mps3

localisation. Further studies are necessary to identify the factors necessary for

the initiation of NE remodelling.

Materials and Methods

Strains and plasmids

All strains described here are derivatives of SK1 diploids and are described in Table S1. *MPS3-FLAG* and *MPS3-GFP* diploid cells showed wild-type spore viability, but both showed ~2 h delay in the entry of meiosis I (Rao et al., 2011). *RAP1-GFP* and *NDJ1-HA* diploids showed wild-type spore viability and normal progression of meiosis (Kosaka et al., 2008; Rao et al., 2011; Trelles-Sticken et al., 1999). GST fusion genes of a PCR-amplified Mps3 fragment (160–260 amino acid residues of the Mps3 protein) were constructed on pGEX-2T (Cytiva, 28954653).

Anti-serum and antibodies

Anti-HA antibody (16B12; Babco), anti-Flag (M2, Sigma), anti-tubulin (MCA77G, Bio-Rad/Serotec, Ltd), anti-GFP (3E6; Molecular Probes), and rabbit anti-Dmc1 (M. Shinohara, Gasior, Bishop, & Shinohara, 2000) were used for staining. Anti-Zip1 (rabbit) and anti-Red1 (chicken) were described previously (M. Shinohara, Oh, Hunter, & Shinohara, 2008). Rabbit anti-Rec8 antibody was described previously (Rao et al., 2011). Guiana pig anti-Hop1 serum was described in (Bani Ismail, Shinohara, & Shinohara, 2014). The second antibodies for staining were Alexa-488 (Goat) and -594 (Goat) IgG used at a 1/2000 dilution (Themo Fishers).

10 amino acid of Mps3 peptide with 189-phosphoserine and 190-phosphoserine (186-GAT**pSpS**PGKSF-195) was synthesized and used for the immunization of rabbit. From obtained serum, antibody specific phosphor-peptide was affinity-purified using the phosphor-peptide (MBL Co. Ltd. Japan).

Cytology

Time-lapse images of Mps3-GFP, Rap1-GFP and Zip1-GFP were captured using a computer-assisted fluorescence microscope system (DeltaVision; Applied Precision). The objective lens was an oil immersion lens (100×; NA, 1.35). Image deconvolution was performed using an image workstation (SoftWorks; Applied Precision). Time-lapse image acquisition was performed every 0.5 to 1 s at a single focal plane. Meiotic cell cultures (~150μl) were placed on bottom-glass

dishes (Matsunami) pre-coated with Concavalin A (10 mg/ml, Sigma).

Tracking of Zip1-GFP/Mps3-GFP/Rap1-GFP was analyzed using the

ImageJ program (NIH) and plug-in MTrack2 (Nico Shuuman). Tracking and

velocity analysis of Mps3-GFP/Rap1-GFP were analyzed using Imaris software

(Oxford Instrument). The sectioned area of each nucleus with Rap1-GFP, which

showed nuclear and dotty staining, was measured using the Velocity^{TM} program

(Applied Precision) with manual assignment of the rim of the sectioned area.

Ninety-six nuclei were counted.

IP and Western blotting

Yeast cell lysates were prepared by the glass bead disruption method. The

lysates were incubated with magnetic beads (Dynal M260; GE Healthcare)

coated with anti-Flag antibody (M2, Sigma) for 12 h and washed extensively

(Sasanuma et al., 2013). Bound proteins were eluted by adding the SDS sample

buffer and were analyzed on an SDS-PAGE gel, transferred to a nylon membrane

(Millipore Co. Ltd), and probed with specific antibodies.

In vitro **kinase assay**

For *in vitro* phosphorylation, 1 μg of purified GST-Mps3 or GST-Mps3-AAA was

incubated with γ-^{32}P-ATP (100 μM; 2.2 MBq) in the presence of CDK (~0.2μg)

and/or DDK (~0.2μg) in a 20 μl of buffer [20 mM Tris-HCl (pH 7.5), 10 mM MgCl$_2$,

10 mM β-glycerophosphate]. After 1-h incubation at 30°C, the reaction was

stopped by adding 5× SDS-PAGE sample buffer, and the products were analyzed

on 10% SDS-PAGE gel. Dried gels were analyzed using a phosphorimager BAS

2000 (Fuji Co. Ltd).

Synthesis of Mps3 peptides

Peptides with the TM and JM region of Mps3 (residue 153-208) were synthesized

by solid-phase method with the following sequence:

GEAKKLK<u>WRTYIFYGGLFFVFYFFGSFL</u>MTTVKNNDLESHSSGA**TSS**PGKSFS

NL-Cys   and

GEAKKLK<u>WRTYIFYGGLFFVFYFFGSFL</u>MTTVKNNDLESHSSGA**DDD**PGKSF

SNL-Cys (underlines indicate TM region). The synthetic peptides were purified

by reverse-phase HPLC on a C4 column with gradient of 2-propanol over water

containing 0.1% trifluoroacetic acid. For fluorescent-labeling, Alexa Fluor 568 C5-

maleimide (Molecular probe) was introduced to the sulfide group on cysteine by

mixing the peptides and the fluorescent dye in dimethylformamide under basic

conditions.

Preparation of lipid bilayer with peptides

The Mps3 peptides were co-solubilized with 1-palmitoyl-2-oleoyl-*sn*-

phosphatidycholine (POPC), 1-palmitoyl-2-oleoyl-*sn*-phosphatidyserine (POPS)

and octyl-β-glucoside in trifluoroethanol (TFE). The peptide to lipid molar ratio

was 1:100. The lipid concentration was 200 μM. The ratio of POPC and POPS

was 10:3. The mixture was incubated for 90 min at 37°C.   Then TFE was

removed under stream of Ar.   MES buffer (10 mM MES, 50 mM NaCl, and 5 mM

DTT, pH 6.2) was added to the solid from the previous step and mixed at 37 °C

for 6 h. The octyl- β -glucoside was removed by dialysis against the MES buffer.

Measurement of FT-IR

Polarized attenuated total reflection FT-IR spectra were obtained on a JASCO

FT/IR-4700 spectrometer. Membranes containing Mps3 TM–JM peptides were

layered on a germanium internal reflection element using a slow flow of nitrogen

gas directed at an oblique angle to the IR plate to form an oriented multilamellar

lipid–peptide film.1000 scans were acquired and averaged for each sample at a

resolution of 4 cm^{-1}. The absorption of polarized light by the amide I bond yields

the dichroic ratio defined as a ratio of absorption intensity for parallel, relative to

perpendicular, polarized light. From the dichroic ratio, we estimated the tilt angle

of the TM helix relative to the membrane normal based on the method described

by Smith and coworkers (Liu, Eilers, Patel, & Smith, 2004; Smith et al.,

2002)) using a value of 41.8° for angle α between the helix director and the

transition-dipole moment of the amide I vibrational mode. For the FT-IR

experiment, the amount of lipid used per experiment was ~4 mg. The film on the

ATR plate in our experiment was assumed to be greater than ~10 μm. For

calculating the dichroic ratio, the thick film limit is applicable (Bechinger,

Ruysschaert, & Goormaghtigh, 1999). Equations that we used for the calculation
of the dichroic ratios were based on this assumption.
844

Fluorescence spectroscopy
For experiments with PIP$_2$, TM-JM peptides were inserted into POPC/POPS
membrane as explained above. The peptide/lipid ratio was set to 1:1000 for
observing the fluorescence from Alexa568. The lipid concentration was 200–250
μM in MOPS buffer (1 mM MOPS, 0.1 M KCl, pH 7.0). Vesicles were formed by
extrusion of multilamellar vesicles through 200-nm polycarbonate filters. PIP$_2$
were added into the membranes by addition of PIP$_2$ micelles to the vesicle
solution. Fluorescence was measured within 1 hour to minimize PIP$_2$ hydrolysis.
The PIP$_2$ concentration was from 5 μM to 25 μM. Fluorescence measurement
was carried out on HITACHI F-2500 fluorescence spectrophotometer at the
excitation wavelength of 568 nm.
856

Coarse-grained MD simulation
All simulations were performed using GROMACS 2016 package (Abraham,
2015) using the MARTINI force field with polarizable water (Marrink, Risselada,
Yefimov, Tieleman, & de Vries, 2007). Periodic boundary conditions were
applied in all directions, and the time step was 20 fs. The temperature was
controlled with the Berendsen temperature coupling scheme (Berendsen, 1984)
with a time constant of 1 ps, and the pressure was controlled using the Parrinello–
Rahman semi-isotropic barostat (Parrinello, 1981) with a time constant of 12 ps
and a compressibility of 3×10^{-4} bar $^{-1}$. All the simulations were carried out in
the isothermal–isobaric (NPT) ensemble at a temperature of 303 K and pressure
of 1.0 bar. The cut-off of the Van der waals interactions were set to 1.1 nm.
868

All atom MD simulation
The NAMD software package (Phillips et al., 2005) was used to perform all-atom
MD simulations. The CHARMM36m force field (Huang et al., 2017) together with
TIP3P water model were used for all simulations. All simulations in NPT dynamics
(constant particle number, pressure and temperature) were carried out at 303 K
with Langevin dynamics, and at 1 atm maintained by Nosè-Hoover Langevin

piston method. Full electrostatic interactions were treated by the particle mesh

Ewald (PME) approach with a grid spacing of less than 1 Å. The cut-off radii of

long-range electrostatic and van der Waals interactions were set to be 12 Å, with

a smoothing function applied from 10 Å.

Acknowledgments

We are grateful to Drs. Nancy Hollingsworth, Franz Klein, Kunihiro Ohta, Kazu
Nishikawa, Hisao Masai, and Hiroyuki Araki for providing the materials used in
this study. We thank Ms. Saeko Hashimoto for excellent technical assistance.
HBDPR was supported by the BMC program and a scholarship from the
Graduate School of Science, Osaka University, JASSO as well as MEXT. KC
were supported by the Institute for Protein Research. This work was supported
by a Grant-in-Aid from the JSPS KAKENHI Grant Number; 22125001, 22125002,
15H05973,16H04742, and 19H00981 to A.S.

Competing interest

The authors declare no competing financial interest.

Author contributions

H.B.D.P.R., T.S., K.C., M.S. and A.S. designed the experiments. H.B.D.P.R., T.S.,
K.C., M.S. and A.S. carried out experiments. T.S. performed liposome
experiments and MD simulation. M.S. provided materials. H.B.D.P.R., T.S., K.C.,
Y.F., and A.S. analyzed the data. A.S. wrote the manuscript with inputs from
H.B.D.P.R., T.S., K.C. and M.S.

References

Abraham, M. J., T. Murtola, R. Schulz, S. Páll, J.C. Smith, B. Hess, and E. Lindahl. (2015). GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX., 1-2*, 19-25.

Bani Ismail, M., Shinohara, M., & Shinohara, A. (2014). Dot1-dependent histone H3K79 methylation promotes the formation of meiotic double-strand breaks in the absence of histone H3K4 methylation in budding yeast. *PLoS One, 9*(5), e96648. doi:10.1371/journal.pone.0096648

Bechinger, B., Ruysschaert, J. M., & Goormaghtigh, E. (1999). Membrane helix orientation from linear dichroism of infrared attenuated total reflection spectra. *Biophys J, 76*(1 Pt 1), 552-563. doi:10.1016/S0006-3495(99)77223-1

Benjamin, K. R., Zhang, C., Shokat, K. M., & Herskowitz, I. (2003). Control of landmark events in meiosis by the CDK Cdc28 and the meiosis-specific kinase Ime2. *Genes Dev, 17*(12), 1524-1539. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12783856

Berendsen, H. J. C., J.P.M. Postma, W.F.v. Gunsteren, A. DiNola, and J.R. Haak. (1984). Molecular dynamics with coupling to an external bath. *The Journal of Chemical Physics., 81*, 3684-3690.

Burke, B. (2018). LINC complexes as regulators of meiosis. *Curr Opin Cell Biol, 52*, 22-29. doi:10.1016/j.ceb.2018.01.005

Challa, K., Lee, M. S., Shinohara, M., Kim, K. P., & Shinohara, A. (2016). Rad61/Wpl1 (Wapl), a cohesin regulator, controls chromosome compaction during meiosis. *Nucleic Acids Res, 44*(7), 3190-3203. doi:10.1093/nar/gkw034

Chen, J., Gardner, J. M., Yu, Z., Smith, S. E., McKinney, S., Slaughter, B. D., . . . Jaspersen, S. L. (2019). Yeast centrosome components form a noncanonical LINC complex at the nuclear envelope insertion site. *J Cell Biol, 218*(5), 1478-1490. doi:10.1083/jcb.201809045

Chikashige, Y., Ding, D. Q., Funabiki, H., Haraguchi, T., Mashiko, S., Yanagida,

M., & Hiraoka, Y. (1994). Telomere-led premeiotic chromosome movement in fission yeast. *Science, 264*(5156), 270-273. doi:10.1126/science.8146661

Chua, P. R., & Roeder, G. S. (1997). Tam1, a telomere-associated meiotic protein, functions in chromosome synapsis and crossover interference. *Genes Dev, 11*(14), 1786-1800. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9242487

Conrad, M. N., Dominguez, A. M., & Dresser, M. E. (1997). Ndj1p, a meiotic telomere protein required for normal chromosome synapsis and segregation in yeast. *Science, 276*(5316), 1252-1255. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9157883

Conrad, M. N., Lee, C. Y., Chao, G., Shinohara, M., Kosaka, H., Shinohara, A., . . . Dresser, M. E. (2008). Rapid telomere movement in meiotic prophase is promoted by NDJ1, MPS3, and CSM4 and is modulated by recombination. *Cell, 133*(7), 1175-1187. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18585352

Conrad, M. N., Lee, C. Y., Wilkerson, J. L., & Dresser, M. E. (2007). MPS3 mediates meiotic bouquet formation in Saccharomyces cerevisiae. *Proc Natl Acad Sci U S A, 104*(21), 8863-8868. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17495028

Cool, M., & Malone, R. E. (1992). Molecular and genetic analysis of the yeast early meiotic recombination genes REC102 and REC107/MER2. *Mol Cell Biol, 12*(3), 1248-1256. doi:10.1128/mcb.12.3.1248-1256.1992

Engebrecht, J., Hirsch, J., & Roeder, G. S. (1990). Meiotic gene conversion and crossing over: their relationship to each other and to chromosome synapsis and segregation. *Cell, 62*(5), 927-937. doi:10.1016/0092-8674(90)90267-i

Ghosh, S., Gardner, J. M., Smoyer, C. J., Friederichs, J. M., Unruh, J. R., Slaughter, B. D., . . . Jaspersen, S. L. (2012). Acetylation of the SUN

protein Mps3 by Eco1 regulates its function in nuclear organization. *Mol*
Biol Cell, 23(13), 2546-2559. doi:10.1091/mbc.E11-07-0600

Hawley, R. S., & Gilliland, W. D. (2009). Homologue pairing: getting it right. *Nat*
Cell Biol, 11(8), 917-918. Retrieved from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=19648975

Henderson, K. A., Kee, K., Maleki, S., Santini, P. A., & Keeney, S. (2006). Cyclin-
dependent kinase directly regulates initiation of meiotic recombination.
Cell, 125(7), 1321-1332. Retrieved from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=16814718

Hiraoka, Y., & Dernburg, A. F. (2009). The SUN rises on meiotic chromosome
dynamics. *Dev Cell, 17*(5), 598-605. Retrieved from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=19922865

Huang, J., Rauscher, S., Nawrocki, G., Ran, T., Feig, M., de Groot, B. L., . . .
MacKerell, A. D., Jr. (2017). CHARMM36m: an improved force field for
folded and intrinsically disordered proteins. *Nat Methods, 14*(1), 71-73.
doi:10.1038/nmeth.4067

Jackson, A. L., Pahl, P. M., Harrison, K., Rosamond, J., & Sclafani, R. A. (1993).
Cell cycle regulation of the yeast Cdc7 protein kinase by association with
the Dbf4 protein. *Mol Cell Biol, 13*(5), 2899-2908.
doi:10.1128/mcb.13.5.2899-2908.1993

Jahed, Z., Domkam, N., Ornowski, J., Yerima, G., & Mofrad, M. R. K. (2021).
Molecular models of LINC complex assembly at the nuclear envelope. *J*
Cell Sci, 134(12). doi:10.1242/jcs.258194

Jaspersen, S. L., Giddings, T. H., Jr., & Winey, M. (2002). Mps3p is a novel
component of the yeast spindle pole body that interacts with the yeast
centrin homologue Cdc31p. *J Cell Biol, 159*(6), 945-956. Retrieved from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=12486115

Jo, S., Kim, T., Iyer, V. G., & Im, W. (2008). CHARMM-GUI: a web-based graphical
user interface for CHARMM. *J Comput Chem, 29*(11), 1859-1865.

doi:10.1002/jcc.20945

Katis, V. L., Lipp, J. J., Imre, R., Bogdanova, A., Okaz, E., Habermann, B., . . .
Zachariae, W. (2010). Rec8 phosphorylation by casein kinase 1 and Cdc7-
Dbf4 kinase regulates cohesin cleavage by separase during meiosis. *Dev*
Cell, *18*(3), 397-409. Retrieved from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=20230747

Klein, F., Mahr, P., Galova, M., Buonomo, S. B., Michaelis, C., Nairz, K., &
Nasmyth, K. (1999). A central role for cohesins in sister chromatid
cohesion, formation of axial elements, and recombination during yeast
meiosis. *Cell, 98*(1), 91-103. doi:10.1016/S0092-8674(00)80609-1

Kosaka, H., Shinohara, M., & Shinohara, A. (2008). Csm4-dependent telomere
movement on nuclear envelope promotes meiotic recombination. *PLoS*
Genet, *4*(9), e1000196. Retrieved from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=18818742

Koszul, R., Kim, K. P., Prentiss, M., Kleckner, N., & Kameoka, S. (2008). Meiotic
chromosomes move by linkage to dynamic actin cables with transduction
of force through the nuclear envelope. *Cell, 133*(7), 1188-1201. Retrieved
from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=18585353

Koszul, R., & Kleckner, N. (2009). Dynamic chromosome movements during
meiosis: a way to eliminate unwanted connections? *Trends Cell Biol,*
19(12), 716-724. Retrieved from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=19854056

Lanz, M. C., Yugandhar, K., Gupta, S., Sanford, E. J., Faça, V. M., Vega, S., . . .
Smolka, M. B. (2021). In-depth and 3-dimensional exploration of the
budding yeast phosphoproteome. *EMBO Rep, 22*(2), e51121.
doi:10.15252/embr.202051121

Lee, C. Y., Bisig, C. G., Conrad, M. M., Ditamo, Y., Previato de Almeida, L.,
Dresser, M. E., & Pezza, R. J. (2020a). Extranuclear Structural

Components that Mediate Dynamic Chromosome Movements in Yeast
Meiosis. *Curr Biol, 30*(7), 1207-1216 e1204.
doi:10.1016/j.cub.2020.01.054

Lee, C. Y., Bisig, C. G., Conrad, M. N., Ditamo, Y., Previato de Almeida, L.,
Dresser, M. E., & Pezza, R. J. (2020b). Telomere-led meiotic chromosome
movements: recent update in structure and function. *Nucleus, 11*(9), 111-
116. doi:10.1080/19491034.2020.1769456

Lee, C. Y., Horn, H. F., Stewart, C. L., Burke, B., Bolcun-Filas, E., Schimenti, J.
C., . . . Pezza, R. J. (2015). Mechanism and regulation of rapid telomere
prophase movements in mouse meiotic chromosomes. *Cell Rep, 11*(4),
551-563. doi:10.1016/j.celrep.2015.03.045

Li, P., Jin, H., Koch, B. A., Abblett, R. L., Han, X., Yates, J. R., 3rd, & Yu, H. G.
(2017). Cleavage of the SUN-domain protein Mps3 at its N-terminus
regulates centrosome disjunction in budding yeast meiosis. *PLoS Genet,*
13(6), e1006830. doi:10.1371/journal.pgen.1006830

Liu, W., Eilers, M., Patel, A. B., & Smith, S. O. (2004). Helix packing moments
reveal diversity and conservation in membrane protein structure. *J Mol Biol,*
337(3), 713-729. doi:10.1016/j.jmb.2004.02.001

Marrink, S. J., Risselada, H. J., Yefimov, S., Tieleman, D. P., & de Vries, A. H.
(2007). The MARTINI force field: coarse grained model for biomolecular
simulations. *J Phys Chem B, 111*(27), 7812-7824. doi:10.1021/jp071097f

Matsushita, C., Tamagaki, H., Miyazawa, Y., Aimoto, S., Smith, S. O., & Sato, T.
(2013). Transmembrane helix orientation influences membrane binding of
the intracellular juxtamembrane domain in Neu receptor peptides. *Proc*
Natl Acad Sci U S A, 110(5), 1646-1651. doi:10.1073/pnas.1215207110

Nishikawa, S., Terazawa, Y., Nakayama, T., Hirata, A., Makio, T., & Endo, T.
(2003). Nep98p is a component of the yeast spindle pole body and
essential for nuclear division and fusion. *J Biol Chem, 278*(11), 9938-9943.
Retrieved from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=12493774

Paouneskou, D., & Jantsch, V. (2019). Meiotic chromosome movement: what's
lamin got to do with it? *Nucleus, 10*(1), 1-6.

doi:10.1080/19491034.2019.1572413

Parrinello, M., and A. Rahman. (1981). Polymorphic transitions in single crystals: A new molecular dynamics method. *Journal of Applied Physics., 52*, 7182-7190.

Patel, J. T., Bottrill, A., Prosser, S. L., Jayaraman, S., Straatman, K., Fry, A. M., & Shackleton, S. (2014). Mitotic phosphorylation of SUN1 loosens its connection with the nuclear lamina while the LINC complex remains intact. *Nucleus, 5*(5), 462-473. doi:10.4161/nucl.36232

Penkner, A. M., Fridkin, A., Gloggnitzer, J., Baudrimont, A., Machacek, T., Woglar, A., . . . Jantsch, V. (2009). Meiotic chromosome homology search involves modifications of the nuclear envelope protein Matefin/SUN-1. *Cell, 139*(5), 920-933. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19913286

Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C., & Ferrin, T. E. (2004). UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem, 25*(13), 1605-1612. doi:10.1002/jcc.20084

Phillips, J. C., Braun, R., Wang, W., Gumbart, J., Tajkhorshid, E., Villa, E., . . . Schulten, K. (2005). Scalable molecular dynamics with NAMD. *J Comput Chem, 26*(16), 1781-1802. doi:10.1002/jcc.20289

Rao, H. B. D. P., Shinohara, M., & Shinohara, A. (2011). Mps3 SUN domain-dependent chromosome motion controls juxtaposition of homologous chromosomes during meiosis. *Genes Cells, 16*, in press.

Sasanuma, H., Hirota, K., Fukuda, T., Kakusho, N., Kugou, K., Kawasaki, Y., . . . Ohta, K. (2008). Cdc7-dependent phosphorylation of Mer2 facilitates initiation of yeast meiotic recombination. *Genes Dev, 22*(3), 398-410. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18245451

Sasanuma, H., Tawaramoto, M. S., Lao, J. P., Hosaka, H., Sanda, E., Suzuki, M., . . . Shinohara, A. (2013). A new protein complex promoting the assembly of Rad51 filaments. *Nat Commun, 4*, 1676.

doi:10.1038/ncomms2678

Sato, T., Pallavi, P., Golebiewska, U., McLaughlin, S., & Smith, S. O. (2006). Structure of the membrane reconstituted transmembrane-juxtamembrane peptide EGFR(622-660) and its interaction with Ca2+/calmodulin. *Biochemistry, 45*(42), 12704-12714. doi:10.1021/bi061264m

Sato, T., Tang, T. C., Reubins, G., Fei, J. Z., Fujimoto, T., Kienlen-Campard, P., . . . Smith, S. O. (2009). A helix-to-coil transition at the epsilon-cut site in the transmembrane dimer of the amyloid precursor protein is required for proteolysis. *Proc Natl Acad Sci U S A, 106*(5), 1421-1426. doi:10.1073/pnas.0812261106

Scherthan, H., Wang, H., Adelfalk, C., White, E. J., Cowan, C., Cande, W. Z., & Kaback, D. B. (2007). Chromosome mobility during meiotic prophase in Saccharomyces cerevisiae. *Proc Natl Acad Sci U S A, 104*(43), 16934-16939. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17939997

Seoane, A. I., & Morgan, D. O. (2017). Firing of Replication Origins Frees Dbf4-Cdc7 to Target Eco1 for Destruction. *Curr Biol, 27*(18), 2849-2855 e2842. doi:10.1016/j.cub.2017.07.070

Sheehan, M. J., & Pawlowski, W. P. (2009). Live imaging of rapid chromosome movements in meiotic prophase I in maize. *Proc Natl Acad Sci U S A, 106*(49), 20989-20994. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19926853

Shibuya, H., Ishiguro, K., & Watanabe, Y. (2014). The TRF1-binding protein TERB1 promotes chromosome movement and telomere rigidity in meiosis. *Nat Cell Biol, 16*(2), 145-156. doi:10.1038/ncb2896

Shinohara, A., Gasior, S., Ogawa, T., Kleckner, N., & Bishop, D. K. (1997). Saccharomyces cerevisiae recA homologues RAD51 and DMC1 have both distinct and overlapping roles in meiotic recombination. *Genes Cells, 2*(10), 615-629. doi:10.1046/j.1365-2443.1997.1480347.x

Shinohara, M., Gasior, S. L., Bishop, D. K., & Shinohara, A. (2000). Tid1/Rdh54 promotes colocalization of rad51 and dmc1 during meiotic recombination.

Proc Natl Acad Sci U S A, 97(20), 10814-10819. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe d&dopt=Citation&list_uids=11005857

Shinohara, M., Oh, S. D., Hunter, N., & Shinohara, A. (2008). Crossover assurance and crossover interference are distinctly regulated by the ZMM proteins during yeast meiosis. *Nat Genet, 40*(3), 299-309. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe d&dopt=Citation&list_uids=18297071

Smith, S. O., Smith, C., Shekar, S., Peersen, O., Ziliox, M., & Aimoto, S. (2002). Transmembrane interactions in the activation of the Neu receptor tyrosine kinase. *Biochemistry, 41*(30), 9321-9332. doi:10.1021/bi012117l

Sosa Ponce, M. L., Moradi-Fard, S., Zaremberg, V., & Cobb, J. A. (2020). SUNny Ways: The Role of the SUN-Domain Protein Mps3 Bridging Yeast Nuclear Organization and Lipid Homeostasis. *Front Genet, 11*, 136. doi:10.3389/fgene.2020.00136

Starr, D. A., & Fridolfsson, H. N. (2010). Interactions Between Nuclei and the Cytoskeleton Are Mediated by SUN-KASH Nuclear-Envelope Bridges. *Annu Rev Cell Dev Biol*. Retrieved from http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe d&dopt=Citation&list_uids=20507227

Storlazzi, A., Xu, L., Schwacha, A., & Kleckner, N. (1996). Synaptonemal complex (SC) component Zip1 plays a role in meiotic recombination independent of SC polymerization along the chromosomes. *Proc Natl Acad Sci U S A, 93*(17), 9043-9048. doi:10.1073/pnas.93.17.9043

Sym, M., Engebrecht, J. A., & Roeder, G. S. (1993). ZIP1 is a synaptonemal complex protein required for meiotic chromosome synapsis. *Cell, 72*(3), 365-378. doi:10.1016/0092-8674(93)90114-6

Tagliabracci, V. S., Engel, J. L., Wen, J., Wiley, S. E., Worby, C. A., Kinch, L. N., . . . Dixon, J. E. (2012). Secreted kinase phosphorylates extracellular proteins that regulate biomineralization. *Science, 336*(6085), 1150-1153. doi:10.1126/science.1217817

Tagliabracci, V. S., Pinna, L. A., & Dixon, J. E. (2013). Secreted protein kinases. *Trends Biochem Sci, 38*(3), 121-130. doi:10.1016/j.tibs.2012.11.008

Tagliabracci, V. S., Wiley, S. E., Guo, X., Kinch, L. N., Durrant, E., Wen, J., . . .
Dixon, J. E. (2015). A Single Kinase Generates the Majority of the
Secreted Phosphoproteome. *Cell, 161*(7), 1619-1632.
doi:10.1016/j.cell.2015.05.028
Tamagaki, H., Furukawa, Y., Yamaguchi, R., Hojo, H., Aimoto, S., Smith, S. O., &
Sato, T. (2014). Coupling of transmembrane helix orientation to membrane
release of the juxtamembrane region in FGFR3. *Biochemistry, 53*(30),
5000-5007. doi:10.1021/bi500327q
Taxis, C., Maeder, C., Reber, S., Rathfelder, N., Miura, K., Greger, K., . . . Knop,
M. (2006). Dynamic organization of the actin cytoskeleton during meiosis
and spore formation in budding yeast. *Traffic, 7*(12), 1628-1642. Retrieved
from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=17118118
Trelles-Sticken, E., Adelfalk, C., Loidl, J., & Scherthan, H. (2005). Meiotic
telomere clustering requires actin for its formation and cohesin for its
resolution. *J Cell Biol, 170*(2), 213-223. Retrieved from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=16027219
Trelles-Sticken, E., Dresser, M. E., & Scherthan, H. (2000). Meiotic telomere
protein Ndj1p is required for meiosis-specific telomere distribution,
bouquet formation and efficient homologue pairing. *J Cell Biol, 151*(1), 95-
106. Retrieved from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=11018056
Trelles-Sticken, E., Loidl, J., & Scherthan, H. (1999). Bouquet formation in
budding yeast: initiation of recombination is not required for meiotic
telomere clustering. *J Cell Sci, 112 (Pt 5)*, 651-658. Retrieved from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=9973600
Ubersax, J. A., Woodbury, E. L., Quang, P. N., Paraz, M., Blethrow, J. D., Shah,
K., . . . Morgan, D. O. (2003). Targets of the cyclin-dependent kinase Cdk1.
Nature, 425(6960), 859-864. Retrieved from

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=14574415

Wan, L., Niu, H., Futcher, B., Zhang, C., Shokat, K. M., Boulton, S. J., &
Hollingsworth, N. M. (2008). Cdc28-Clb5 (CDK-S) and Cdc7-Dbf4 (DDK)
collaborate to initiate meiotic recombination in yeast. *Genes Dev, 22*(3),
386-397.   Retrieved   from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=18245450

Wan, L., Zhang, C., Shokat, K. M., & Hollingsworth, N. M. (2006). Chemical
inactivation of cdc7 kinase in budding yeast results in a reversible arrest
that allows efficient cell synchronization prior to meiotic recombination.
Genetics, 174(4), 1767-1774.   Retrieved   from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=17057233

Wanat, J. J., Kim, K. P., Koszul, R., Zanders, S., Weiner, B., Kleckner, N., & Alani,
E. (2008). Csm4, in collaboration with Ndj1, mediates telomere-led
chromosome dynamics and recombination during yeast meiosis. *PLoS*
Genet, 4(9), e1000188.   Retrieved   from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=18818741

White, E. J., Cowan, C., Cande, W. Z., & Kaback, D. B. (2004). In vivo analysis
of synaptonemal complex formation during yeast meiosis. *Genetics,*
167(1), 51-63. doi:10.1534/genetics.167.1.51

Wong, X., Loo, T. H., & Stewart, C. L. (2021). LINC complex regulation of genome
organization and function. *Curr Opin Genet Dev, 67*, 130-141.
doi:10.1016/j.gde.2020.12.007

Zickler, D., & Kleckner, N. (1998). The leptotene-zygotene transition of meiosis.
Annu Rev Genet, 32, 619-697.   Retrieved   from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe
d&dopt=Citation&list_uids=9928494

Zickler, D., & Kleckner, N. (1999). Meiotic chromosomes: integrating structure
and function. *Annu Rev Genet, 33*, 603-754. Retrieved from
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMe

d&dopt=Citation&list_uids=10690419

Zuela, N., & Gruenbaum, Y. (2016). Matefin/SUN-1 Phosphorylation on Serine 43

Is Mediated by CDK-1 and Required for Its Localization to Centrosomes

and Normal Mitosis in C. elegans Embryos. *Cells, 5*(1).

doi:10.3390/cells5010008

1238

1239

Figure legends

Figure 1. Meiotic Mps3 localization and movement on NEs

(A) Mps3-GFP localization was analyzed in wild-type cells (PRY64) at various time points during meiosis. Representative images at each time are shown. Bottom color bars indicate classes of Mps3-localization in (B). Bar indicates 2 μm.

(B) Based on Mps3-GFP localization, cells with Mps3-GFP were classified into four classes (A) and quantified: single Mps3 focus (blue), 2–5 foci (green), more than 5 foci/patches (orange), and coverage of the Mps3 signal on NE (red). At each time point, more than 100 nuclei were counted. The graphs are a representative of two independent time courses.

(C) Time-lapse analysis of Mps3-GFP in the wild-type strain (PRY64) at different time points in meiosis. A single focal plane of a cell every 3 sec is shown. See Videos 1-4.

(D) Time-lapse analysis of Mps3-GFP in various strains at different time points in meiosis. A single focal plane of a cell was analyzed every 3 sec. The inhibitor LatB (30 μM) was added at 0 h. See supplemental movie S5. While washing, the inhibitor was washed at 5 h, and the cells were analyzed at 6 h. Wild type, PRY64; *ndj1*, PRY192; *csm4*, PRY198.

(E) Time-lapse analysis of Mps3-GFP in the *cdc28-as1* mutant (PRY71) treated with or without the inhibitor 1NM-PP1 (0.5 μM) at 5 h during meiosis. The inhibitor was added at 0 h (second panels) or 3 h (third panels). While washing (fourth panels), the inhibitor was added at 0 h and washed at 3 h, and the cells were analyzed at 5 h. Time-lapse analysis of Mps3-GFP in the *cdc7-as3* mutant (PRY260) treated with or without the inhibitor PP1 (15 μM). The *cdc7 bob1-1* mutant (PRY115) was analyzed at 6 h. See Videos 6-8.

(F) Tracking of Mps3-GFP in wild type, *cdc28-as1* (PRY71 with the inhibitor), *cdc7-as3* (PRY260with the inhibitor), and *cdc7 bob1-1* (PRY115). Tracking was monitored for all Mps3 foci/patches in a single cell for 20 sec at 4 or 5 h in SPM. Each line represents tracking of the foci at a single focal plane.

(G) Percentages of cells with different classes of Mps3-GFP were quantified at 5 h under different conditions (triplicates, Error bars show standard deviation;

s.d.); single Mps3 focus (blue), 2–5 foci (green), more than 5 foci/patches

(orange), and coverage of the Mps3 signal on NE (red).

(H) Velocity of Mps3-GFP foci or patches was quantified. Time-lapse images

were taken for every second in 20 seconds. Mps3-GFP foci or patches were

identified as shown in Methods and followed for their tracks. For each track,

an average velocity and maximum velocity were calculated. More than 20

cells were analyzed for the quantification. Red lines show mean with s.d.. *P*-

values were calculated using Man-Whitney's *U*-test.

1281

Figure 2. Meiotic telomere movement in *cdk* and *ddk* mutants

(A) Time-lapse analysis of Rap1-GFP in wild type (HKY167) at different time

points during meiosis. An image of a single focal plane of each mutant cell

was taken every 3 sec. Clustering of telomeres is shown by yellow arrows.

Bar indicates 2 μm. See Videos 9, 10.

(B) Tracking of Rap1-GFP in wild type (0 and 4h), *cdc28-as1* (with the inhibitor),

and *cdc7-as3* (with the inhibitor). Tracking was monitored for all Rap1 foci in

a single cell for 20 sec. Each line represents tracking of the center of foci at a

single focal plane.

(C) Velocity of Rap1-GFP foci or patches were quantified. Time-lapse images

were taken for every second in 20 seconds. Rap1-GFP foci or patches were

identified as shown in Methods and followed for their tracks. For each track,

an average velocity and maximum velocity were calculated. More than 20

cells were analyzed for the quantification. Red lines show mean with s.d.. *P*-

values were calculated using Man-Whitney's *U*-test.

(D) Kinetics of telomere clustering during meiosis were quantified. At each time

point, more than 150 nuclei were counted for clustering of telomeres (arrows

in A, E, F). The graphs are a representative of two independent time course.

(left) *cdc28-as1* with 1NM-PP1, blue closed circles; *cdc28-as1* without 1NM-

PP1, blue closed circles; *cdc7-as3* with PP1, red closed circles; *cdc7-as3*

without PP1, red closed circles. (right) wild type, blue closed circles; *cdc7*

bob1-1, red closed circles.

(E) Time-lapse analysis of Rap1-GFP in various strains (*cdc28-as1*, PRY68) at

different time points during meiosis. An image of a single focal plane of each

mutant cell was taken every 3 sec. The inhibitors (final 0.5 μM of 1NM-PP1)

were added at 0 h (second panels) or 3 h (third panels). While washing

(bottom panels), the inhibitor was added at 0 h and washed at 3 h, and the

cells were analyzed at 5 h. Clustering of telomeres is shown by yellow arrows.

Bar indicates 2 μm. See Video 11.

(F) Time-lapse analysis of Rap1-GFP in various strains (*cdc7-as3,* PRY79; *cdc7*

bob1-1 PRY116) at different time points during meiosis. An image of a single

focal plane of each mutant cell was taken every 3 sec. For the *cdc7-as3*

mutant, the inhibitors (final 15 μM PP1, respectively) were added at 0 h

(second panels) or 3 h (third panels). While washing (bottom panels), the

inhibitor was added at 0 h and washed at 3 h, and the cells were analyzed at

5 h. Clustering of telomeres is shown by yellow arrows. Bar indicates 2 μm.

See Videos 12, 13.

1319

Figure 3. Mps3 is phosphorylated in luminal region

(A) Schematic representation of the Mps3 protein. The putative transmembrane

(TM) region (dark green) and SUN domain (green) are shown with three

known phosphorylation sites (red circles) in nucleoplasmic region (Lanz et al.,

2021). The position of a CDK consensus site (S/TP) is shown in red lines with

aa sequence on the bottom. All of the six sites are located in a luminal region

of Mps3.

(B) Western blotting of the Mps3-Flag protein during meiosis. Wild-type (left,

HKY404), *mps3-AAA* mutant (middle, PRY163) and *MPS3-MER2S* mutant

proteins (right, PRY514) were analyzed at different time points. Fractions

immunoprecipitated with anti-Flag antibody were probed with anti-Flag

antibody (top) as well as a phospho-specific antibody (bottom) that

recognized T189 and S190 phosphorylation.

(C) *In vivo* phosphorylation was confirmed by a decreased band shift of Mps3-

Flag protein purified from meiotic cell lysates (4 h incubation with SPM) using

the anti-Flag antibody. After immuno-precipitation, the precipitate was

incubated with 1 unit of CIP for one hour either in the presence or absence

of phosphatase inhibitors. Mps3-Flag was detected using anti-Flag (M2)

antibody.

(D) Affinity-purified-Mps3-Flag from meiotic cell lysates (4 h incubation with SPM)
using anti-Flag column was treated with CIP as described in (A) and was
probed with anti-Phospho-Serine antibody (Qiagen). During incubation, Mps3
protein was degraded extensively (bottom).

(E) Sequence homology of an Mps3 region with a CDK/DDK phosphorylation site
of the Mer2 protein. Identical amino acids are shown in red. Putative CDK or
DDK phosphorylation sites are shown in orange. Unique amino acid
sequence in Mer2 is shown in pale blue. The substitutions (*mps3-AAA,*
MPS3-DDD, and *MPS3-MER2S*) for the putative phosphorylation sites of
Mps3 are shown on the top.

(F) *In vitro* phosphorylation analysis of GST-Mps3 fusion proteins. Purified wild-
type GST-Mps3 and GST-Mps3-AAA fragments (shown on the left by
Coomassie staining) were incubated with different combinations of purified
the Cdc28–Clb5 complex and/or Cdc7–Dbf4 kinase complexes in the
presence of γ-^{32}P-ATP. After 1-h incubation, proteins were fractionated on a
SDS-PAGE gel and analyzed using a phosphorimager for incorporation of ^{32}P
in the Mps3 fragments (right bottom) and Cdc28/Cdc7 protein (right top).
Cdc7 was phosphorylated by CDK. Quantification is shown below the
autoradiogram.

(G) Western blotting of Mps3 and tubulin at various times in meiosis. Cell lysates
from the *MPS3-Flag* (right, HKY404) and *MPS3-AAA-Flag* (left, PRY163)
strains were analyzed by western blotting using anti-Flag and anti-tubulin
antibodies. The wild-type Mps3 protein shows band shift during meiosis for
example at 4 and 6 h.

(H) Reactivity of wild-type Mps3-Flag and Mps3-AAA mutant protein to anti-
phospho-serine antibody was examined by Western blotting. wild type Mps3-
Flag and Mps3-AAA mutant protein at each time point were IPed and probed
with the antibody. While wild type Mps3-Flag shows an increase reactivity to
the antibody, Mps3-AAA-Flag did not increase the reactivity in meiosis (4 h)
compared to mitosis (0 h).

(I) Western blotting of the Mps3-Flag protein during meiosis. Wild-type (left,
HKY404) and *MPS3-MER2S* proteins (right, PRY514) were analyzed at
different time points. Fractions immunoprecipitated with anti-Flag antibody

were probed with anti-Flag and anti-Mer2-phospho-S30 antibodies.

(J) Rap1-GFP dynamics at 4 h of meiosis analyzed in *MPS3-MER2s* (PRY518)

and wild-type (HKY167) strains.

(K) Kinetics of Rap1 clustering in each strain is shown in G. Counting was

performed as described in Figure 2E. Blue, wild type; red, *MPS3-MER2S.*

(L) Localization of Ndj1-HA (green), Mps3-Mer2S-Flag (red), and DAPI (dark

blue) was analyzed by whole cell staining of PRY514. Bar indicates 2 μm.

1379

Figure 4. The *mps3-AAA* mutant impairs telomere and NE movements

(A) Meiosis division was analyzed by DAPI staining for various strains. More than

100 cells were counted for divisions. The graphs are a representative of two

independent time course. Spore viability of each strain is shown in

parenthesis. At least 100 asci were dissected for each strain.

(B) Time-lapse analysis of Rap1-GFP in the wild-type (HKY167) and *mps3-AAA*

(PRY138) strains at different time points in meiosis. A single focal plane of a

cell was analyzed every 3 s. A tethering defect was observed in the *mps3-*

AAA strains. Bar indicates 2 μm. See supplemental movie S14 for *mps3-AAA.*

(C) Tracking of Rap1-GFP in wild type and *mps3-AAA* for 20 sec. See Methods

in Figure 2B.

(D) Velocity of Rap1-GFP foci or patches were quantified. See Methods in Figure

2C. Red lines show mean with s.d.. *P*-values were calculated using Man-

Whitney's *U*-test.

(E) Time-lapse analysis of Zip1-GFP in wild type (SEY672) and *mps3-AAA*

(PRY332). *mps3-AAA.* See Videos 15, 16.

(F) Tracing of Zip1-GFP shows a step size of each Zip1-GFP line. Step size per

given time is converted into the relative velocity of chromosomes. *n*=35. Blue

bars; wild type; red bars, *mps3-AAA.*

(G) Localization of wild-type Mps3-GFP (PRY64) and Mps3-AAA-GFP (PRY186)

proteins in a cell at different time points in meiosis.

(H) Kinetics of Mps3 distribution. Based on Mps3-GFP morphology, cells with

Mps3-GFP were classified into four classes and quantified: 2–5 foci (green),

more than 5 foci/patches (orange), and coverage on NE (red). See Figure 1B

for quantification.

(I) Percentages of cells with different classes of Mps3-GFP were quantified at 5 h in different conditions (triplicates, Error bars show standard deviation; s.d.); single Mps3 focus (blue), 2–5 foci (green), more than 5 foci/patches (orange), and coverage of the Mps3 signal on NE (red).

(J) Time-lapse analysis of wild type and Mps3-AAA-GFP during meiosis. See Video 17 for Mps3-AAA-GFP.

(K) Tracking of Mps3-GFP in wild type and *mps3-AAA* for 20 sec. See Figure 1F for quantification.

(L) Velocity of Mps3-GFP foci or patches were quantified. Methods are shown in Figure 1H. More than 20 cells were analyzed for the quantification. Red lines show mean with s.d.. *P*-values were calculated using Man-Whitney's *U*-test.

(M) The maximum sectional area of each nucleus was measured using the Velocity program. Each area of the 96 nuclei is ranked in the figure. Graphs for wild type and *mps3-AAA* at 0 and 4 h as well as wild type treated with LatB are shown.

Figure 5. Acid amino acids in lumen of Mps3 allele can suppress CDK and DDK defects

(A) Time-lapse analysis of Rap1-GFP in a phosphomimetic allele of *MPS3*, *MPS3-DDD* (PRY211), with a different combination of *cdc28-as1* (PRY211), *cdc7-as3* (PRY301)*,* and *cdc7 bob1-1* (PRY272) mutations in meiosis. A single focal plane of a cell was analyzed at every 3 s. Bar indicates 2 μm. See Videos 19-22.

(B) Kinetics of clustering of telomere as Rap1-clusters was studied in *cdc7-as3* (left, PRY309) and *cdc28-as1* (right, PRY303) in the presence of its specific inhibitors. Wild type (blue, HKY167) and *MPS3-DDD* (PRY236) with the allele was used.

(C) Time-lapse analysis of Mps3-DDD-GFP with different mutant alleles and in treatment with a specific inhibitor at different time points in meiosis. See Videos 18, 23-25.

(D) Kinetics of Mps3. Based on Mps3-GFP morphology, cells with Mps3-GFP were classified into four classes and quantified: single Mps3 focus (blue), 2–3 foci (green), 4–5 foci (orange), and more than 5 foci (red). See Figure 2B

for quantification. Graphs for wild type, *cdc28-as1*, and *cdc7-as3* cells are the

same as in Figure 1B.

(E) Tracking of Mps3-GFP in *mps3-DDD*, *cdc28-as1 mps3-DDD* (with 1NM-PP1),

cdc7-as3 mps3-DDD (with PP1), and *cdc7 bob1-1mps3-DDD* for 20 sec.

(F) Velocity of Mps3-GFP foci or patches were quantified. Methods are shown in

Figure 1H. More than 20 cells were analyzed for the quantification. Red lines

show mean with s.d.. *P*-values were calculated using Man-Whitney's *U*-test.

(G) Time-lapse analysis of Mps3-DDD-GFP with different mutant alleles during

vegetative growth.

1447

Figure 6. Biophysical analysis of JM region of Mps3

(A) Amino acid sequences of Mps3 with amino acid substitution in the *mps3-*

S189A and *mps3-S190A* mutants.

(B) Localization of wild-type Mps3-GFP (PRY64), Mps3-S189A-GFP

(KSY407/409) Mps3-S190A-GFP (KSY220/221) and proteins in a cell at

different time points in meiosis.

(C) Kinetics of Mps3 distribution. Based on Mps3-GFP morphology, cells with

Mps3-GFP were classified into four classes and quantified: 2–5 foci (green),

more than 5 foci/patches (orange), and coverage on NE (red). See Figure 1B

for quantification.

(D) Percentage of cells with different classes of Mps3-GFP were quantified at 5 h

in different conditions (triplicates, Error bars show standard deviation; s.d.);

single Mps3 focus (blue), 2–5 foci (green), more than 5 foci/patches (orange),

and coverage of the Mps3 signal on NE (red).

(E) Tracking of Mps3-GFP in wild type (left), *mps3-S189A* (middle) and *mps3-*

S190A (right).

(F) Velocity of Mps3-GFP foci or patches were quantified in different *mps3*

mutants. Methods are shown in Figure 1H. More than 20 cells were analyzed

for the quantification. Red lines show mean with s.d.. *P*-values were

calculated using Man-Whitney's *U*-test.

(G) The maximum sectional area of each nucleus was measured using the

Velocity program. Each area of the 96 nuclei is ranked in the figure.

1470

Figure 7. Biophysical analysis of JM region of Mps3

(A) Amino acid sequences of Mps3 peptides used. Amino acid sequence in transmembrane region is shown in green. Phosphorylated residues TSS and its derivative DDD are shown in blue and red, respectively.

(B) Fluorescence spectra from a synthetic liposome containing Alexa568-labelled WT-Mps3 (left) peptide and Mps3-DDD peptide (right) were measured in the presence of various concentrations of PIP_2. The graphs are a representative of three independent time course.

(C) Schematic illustration of the binding of Mps3-WT peptide (left) and Mps3-DDD peptide (right) to lipid bilayer with PIP_2 (red). DDD residues may induce electric repulsion between the peptide and the membrane. Transmembrane region of the Mps3 peptide is shown by blue a-helix. Peptides with WT JM region and with DDD substitution are shown blue and red lines, respectively. Purple and black stars show normal and quenched fluorophores, respectively.

(D) Snapshots from all-atom MD simulation trajectories for Mps3-WT peptide (left), Mps3-DDD peptide (center) and Mps3-WT peptide with phosphorylation at TSS sequence (right). See Videos 26-28.

Figure 8. A model of regulation of Mps3 on NE during meiosis

(A) Multiple phosphorylation regulates Mps3 localisation and motion in NE during meiosis. In mitotic cells, Mps3 is mainly located in the SPB. In some strain, disperse weak Mps3 signal is detected on NE, which might be regulated by DDK. Upon entry into meiosis, Mps3 forms a few foci on NE in early prophase I, and phosphorylation of the luminal region of Mps3 and, probably DDK, promotes localisation of more Mps3 proteins as focus/patch on NE. Mps3 foci/patches transiently form a cluster of some Mps3 foci by an unknown mechanism, and CDK and DDK and Rec8-cohesin promote dissociation of Mps3 and Rap1(telomere) clusters. It is likely that positive feedback on Mps3 phosphorylation would increase the localisation of Mps3 on NEs, which results in full NE coverage of Mps3.

(B) A hypothetical model of the formation of Mps3-containing LINC complex. During mitosis, Mps3 forms a non-canonical LINC complex with *cis-*membrane interactions with Mps2 as a component of the half-bridge and

SPIN (left). Upon induction of meiosis, unknown factors promote the formation

of Mps3 foci on NE. Then, the JM region of Mps3 is subject to non-canonical

phosphorylation, which might in turn weaken the binding of JM to INM,

probably by electric repulsion between the JM regions with negative charges

and acid lipids. This process promotes the formation of the canonical LINC

complex with the *trans*-membrane configuration, in which the Mps3 SUN

domain binds to the KASH domain of Mps2 and Csm4. Csm4 may promote

structural changes in the luminal region of Mps2. The N-terminal region of

Mps3, which is located in the nucleoplasm, binds to a telomere-binding protein,

Ndj1. During middle and late prophase I, Mps3 forms a large protein ensemble

on the NE, which is seen as a patch.

1515

Supplemental Figures

Figure S1. Meiotic movements of Mps3 and Rap1 in the presence of the inhibitors

(A) Tracking of Mps3-GFP in wild type. Tracking was monitored for all Mps3 foci/patches in a single cell (PRY64) for 120 sec. Each line with different colors, representing a single focus/patch, show tracking of the objects at a single focal plane.

(B) Tracking of Mps3-GFP in wild type. Tracking was monitored for all Mps3 foci/patches in a single cell (PRY64) for 20 sec. See (A).

(C) Cytoplasmic actin cables were visualized by Phalloidin (red) and DAPI (purple) staining of *cdc28-as1* (PRY68) cells after 5 h of incubation in SPM both without (left) or with (middle) its inhibitor 1NM-PP1 from 0 h. As a control, the mutant cells were treated with 30 μM of latrunculin B (LatB) at 4 h and analysed for actin staining at 5 h (right). Bar indicates 2 μm.

(D) Velocity of Mps3-GFP foci or patches were quantified. Methods are shown in Figure 1H. More than 20 cells were analyzed for the quantification. Red lines show mean with s.d.. *P*-values were calculated using Man-Whitney's *U*-test.

(E) Western blotting analysis of Mps3-Flag protein. Cell lysates from the *cdc28-as1* (PRY13) cells at different times in meiosis. The *cdc28-as1* cells treated with or without 1NM-PP1 from 0 h. Tubulin is a control.

(F) To check the off-target of two inhibitors, 1NM-PP1 and PP1, wild-type diploid cells (HKY404) with Mps3-GFP were grown in the presence of each inhibitor (0.5 μM of 1NM-PP1, top panels; 15 μM PP1, bottom panels). The inhibitors were added at 0 h (second panels) and time-lapse images were examined in 5 h-incubation with SPM. The two inhibitors did not affect rapid movement of Mps3-GFP foci on NEs. Bar indicates 2 μm.

(G) Western blotting analysis of Mps3-Flag protein. Cell lysates from the *cdc7-as3* (PRY72) cells at 0 and 5 h in meiosis. The *cdc7-as3* cells treated with or without PP1 from 0 h and collected at 5h. Tubulin is a control.

(H) Whole-cell staining analysis of Mps3-Flag and Ndj1. Various cells were stained with anti-Flag for Mps3-Flag (red) and anti-HA for Ndj1 (green) at the nuclear periphery at indicated times. *cdc28-as1* (PRY13) and *cdc7-as3*

(PRY72) cells were treated with their inhibitor from 0 h and examined at the
indicated times. Wild type, HKY404.

(I) To check the off-target of two inhibitors, 1NM-PP1 and PP1, wild-type diploid
cells (HKY167) with Rap1-GFP were grown in the presence of each inhibitor
(0.5 μM of 1NM-PP1, top panels; 15 μM PP1, bottom panels). The inhibitors
were added at 0 h (second panels) and time-lapse images were examined in
5 h-incubation with SPM. The two inhibitors did not affect rapid movement of
Rap1-GFP foci on NEs. Bar indicates 2 μm.

1557

Figure S2. CDK and DDK are necessary for Mps3 localization on NE

(A) Image of a part of a tetrad dissection plate of spores from *MPS3-MER2S-Flag*
diploid cells. After the dissection, the plate was incubated at 30C for two days.

(B) Whole cell staining of Mps3-Flag and Tubulin. Wild type (MSY832/833) and
the *mps3-AAA* mutant (PRY122) cells in mitosis were stained with anti-Flag
(red) and anti-tubulin (green). Representative images with short spindles with
two Mps3 foci are shown. Bar indicates 2 μm.

(C) FACS analysis of DNA contents in wild type and the *mps3-AAA* mutant cells
at indicated times. Cells at each time point were fixed and stained with
propidium iodide (PI) and analyzed by FACS after the sonication.

(D) Immunostaining analysis of meiotic chromosome proteins. For SC formation,
Zip1 localization (red) with DAPI (blue) image is also analyzed for wild type
(top) and *mps3-AAA* mutant strains (bottom) at 4 h.

(E) Kinetics of SC formation. SC elongation was analyzed by staining Zip1 protein.
Class I (blue) contains Zip1 dots, Class II (green) contains partial lines of Zip1,
and Class III (red) contains fully elongated Zip1 lines. At each time point, more
than 50 nuclei were counted. The graphs are a representative of two
independent time course.

(F) Interaction of Mps3-Flag with Ndj1-HA and Csm4 in the *mps3-AAA* mutant
was confirmed by immuno-precipitation using anti-Flag antibodies. IPed
fractions of cells with Mps3-Flag (left, HKY404) or Mps3-AAA-Flag (right,
PRY163) at each time point were probed with anti-Flag (top), anti-HA (middle)
and anti-Csm4 antibodies.

1581

Figure S3. Meiotic phenotypes of *MPS3-DDD* mutants

(A) Whole cell staining of Mps3-Flag and Tubulin. Wild type (MSY832/833) and the *mps3-DDD* mutant (PRY122) cells during vegetative growth were stained with anti-Flag (red) and anti-tubulin (green). Representative images with short spindles with two Mps3 foci are shown. Bar indicates 2 μm.

(B) Different classes of tubulin staining (red) in vegetative yeast cells.

(C) Cells with different tubulin classes (B) were counted (n=30) and percentages of each class in wild type and the *MPS3-DDD* strains (PRY201) are shown (n=3). Error bars show s.d.

(D) Meiosis I division was analyzed by DAPI staining for wild type and the *MPS3-DDD* strains (PRY201). At each time point, more than 100 nuclei were counted. The graphs are a representative of two independent time course.

(E) Localization of Mps3-DDD-GFP protein in *MPS3-DDD* (PRY201) at different time points. Bar indicates 2 μm.

(F) Localization of Mps3-DDD-GFP protein in wild type, *cdc28-as1* (with 1NM-PP1; PRY301) and *cdc7-as3* (with PP1; PRY272) at 0 h. Bar indicates 2 μm.

(G) An image of Rap1-GFP in the *mps3-DDD* strains with a different combination of *cdc28-as1*, *cdc7-as3* or *cdc7 bob1-1* mutations in mitosis (0 h). A single-focal plane of a cell was analyzed at every 3 second. Bar indicates 2 μm.

Figure S4. Structure and dynamics of JM region of Mps3

(A) Polarized attenuated total reflection FT-IR spectra are shown. Spectra (red) a synthetic liposome containing Mps3-WT (left) peptide and Mps3-DDD peptide (right) were obtained with light polarized parallel (top) or perpendicular (bottom). Deconvoluted spectra into α-helix and β-sheet are shown in blue.

(B) Snapshots from coarse-grained MD simulation trajectories for Mps3-WT peptide (left), Mps3-DDD peptide (right).

(C) Mass density profiles relative to the center of the lipid bilayer for residues, T188, S189 and S190 on Mps3-WT peptide (blue), Mps3-DDD peptide (red) and Mps3-WT peptide with phosphorylation (green). Density profile for phosphate groups on lipids are colored with yellow.

1615   **Supplemental movie Information**

1616   Each Video (Videos 1-25) was taken every one second in a single focal plane for

1617   120 s. The speed is 9.1 fold.

1618

Video 1. Mps3-GFP in a wild-type cell at 0 h in SPM.

Video 2. Mps3-GFP in a wild-type cell at 4 h in SPM.

Video 3. Mps3-GFP in a wild-type cell at 5 h in SPM, showing deformation.

Video 4. Mps3-GFP in a wild-type cell at 5 h in SPM, showing protrusion.

Video 5. Mps3-GFP in a wild-type cell treated with LatB (0 h) at 5 h in SPM.

Video 6. Mps3-GFP in *cdc28-as1* cells treated with 1NM-PP1 (0 h) at 5 h in SPM.

Video 7. Mps3-GFP in a *cdc7-as3* cell treated with PP1 (0 h) at 5 h in SPM.

Video 8. Mps3-GFP in a *cdc7 bob1-1* cell at 5 h in SPM.

Video 9. Rap1-GFP in a wild-type cell at 0 h in SPM.

Video 10. Rap1-GFP in a wild-type cells at 4 h in SPM.

Video 11. Rap1-GFP in a *cdc28-as1* cell treated with 1NM-PP1 (0 h) at 5 h in
SPM.

Video 12. Rap1-GFP in a *cdc7-as3* cell treated with PP1 (0 h) at 5 h in SPM.

Video 13. Rap1-GFP in a *cdc7 bob1-1* cell at 5 h in SPM.

Video 14. Rap1-GFP in *mps3-AAA* cells at 5 h in SPM.

Video 15. Zip1-GFP in a wild-type cell at 5 h in SPM.

Video 16. Zip1-GFP in *mps3-AAA* cells at 6 h in SPM.

Video 17. Mps3-AAA-GFP at 5 h in SPM.

Video 18. Mps3-DDD-GFP at 5 h in SPM.

Video 19. Rap1-GFP in *MPS3-DDD* cells at 5 h in SPM.

Video 20. Rap1-GFP in a *cdc28-as1 MPS3-DDD* cell treated with 1NM-PP1 (0 h)
at 5 h in SPM.

Video 21. Rap1-GFP in a *cdc7-as3 MPS3-DDD* cell treated with PP1 (0 h) at 5 h
in SPM.

Video 22. Rap1-GFP in a *cdc7 bob1-1 MPS3-DDD* cell at 5 h in SPM.

Video 23. Mps3-DDD -GFP in a *cdc28-as1 MPS3-DDD* cell treated with 1NM-
PP1 (0 h) at 5 h in SPM.

Video 24. Mps3-DDD -GFP in a *cdc7-as3 MPS3-DDD* cell treated with PP1 (0 h)
at 5 h in SPM.

Video 25. Mps3-DDD -GFP in a *cdc7 bob1-1 MPS3-DDD* cell at 5 h in SPM.

Video 26. MD simulation of wild-type Mps3 peptide.

Video 27. MD simulation of wild-type Mps3-DDD peptide.

Video 28. MD simulation of wild-type Mps3-phosphorylated peptide.

1652

Figure 1. Rao et. al.

Figure 2. Rao et. al.

Figure 3 Rao et. al.

Figure 4 Rao et. al.

Figure 5 Rao et. al.

Figure 6 Rao et. al.

Figure 7. Rao et. al.

A

| | Trans-membran domain | Juxtamembrane region |

Mps3-WT   NH2-GEAKKLKWRTYIFYGGLFFVFFFGSFLMTTVKNNDLESHSSGATSSPGKSFSNL

Mps3-DDD   NH2-GEAKKLKWRTYIFYGGLFFVFFFGSFLMTTVKNNDLESHSSGADDDPGKSFSNL

B

Mps3-WT

Mps3-DDD

0 μM PIP2
4.8 μM PIP2
9.6 μM PIP2
14.4 μM PIP2
19.2 μM PIP2
24.0 μM PIP2

C

wild type

Mps3-DDD

Electrostatic repulsion between negative charges

PIP₂ addition

D

wild type

Mps3-DDD

wild type phosphorylated

Table S1: Strain list

Strain Name	Genotype
NKY1551	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, his4B-LEU2/his4X-LEU2-URA3, arg4-bgl/arg4-nsp*
MSY832/833	*MATa/α, ho::LYS2, ura3, leu2::hisG, trp1::hisG, lys2*
HKY404	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3,lys2/lys2,MPS3-3FLAG::KanMX6/ MPS3-3FLAG::KanMX6,NDJ1-3HA::KanMX6/NDJ1-3HA::KanMX6*
PRY13	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, MPS3-3FLAG::KanMX6/ MPS3-3FLAG::KanMX6, NDJ1-3HA::KanMX6/NDJ1-3HA::KanMX6, cdc28-as1/cdc28-as1*
PRY72	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, MPS3-3FLAG::KanMX6/MPS3-3FLAG::KanMX6, NDJ1-3HA::KanMX6/NDJ1-3HA::KanMX6, cdc7-as3-9MYC/cdc7-as3-9MYC*
PRY 64	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, MPS3-3GFP::KanMX6/MPS3-3GFP::KanMX6*
PRY71	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3,MPS3-3GFP::KanMX6/MPS3-3GFP::KanMX6, cdc28-as1/cdc28-as1*
PRY260	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, MPS3-3GFP::KanMX6/MPS3-3GFP::KanMX6, cdc7-as3-9MYC/cdc7-as3-9MYC*

PRY115	MAT*a*/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, MPS3-3GFP::KanMX6/MPS3-3GFP::KanMX6, cdc7::KanMX6/cdc7::KanMX6, bob1-1/bob1-1
PRY122	MAT*a*/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, mps3-AAA(T188A S189A S190A-PSTI)/ mps3-AAA (T188A S189A S190A-PSTI)
PRY186	MAT*a*/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, mps3-AAA(T188A S189A S190A-PSTI)-3GFP::KanMX6/mps3-AAA (T188A S189A S190A-PSTI) -3GFP::KanMX6
PRY163	MAT*a*/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, mps3-3A(T188A S189A S190A-PSTI)-3FLAG::KanMX6/ mps3-3A(T188A S189A S190A-PSTI)-3FLAG::KanMX6 ,NDJ1-3HA ::KanMX6/ NDJ1-3HA::KanMX6
PRY138	MAT*a*/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, mps3-AAA(T188A S189A S190A-PSTI)/ mps3-AAA (T188A S189A S190A-PSTI), Rap1-GFP::LEU2/Rap1-GFP::LEU2
PRY201	MAT*a*/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, MPS3-DDD(T188D S189D S190D–BSPEI)/MPS3-DDD(T188D S189D S190D–BSPEI)
PRY211	MAT*a*/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, MPS3-DDD(T188D S189D S190D–BSPEI)-3GFP::KanMX6/ MPS3-DDD(T188D S189D S190D–BSPEI)-

3GFP::KanMX6

PRY219   *MAT**a**/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG,*

         ura3/ura3, MPS3-DDD(T188D S189D S190D–BSPEI)-

         3FLAG::KanMX6/ MPS3-DDD(1 T188D S189D S190D–

         BSPEI)-3FLAG::KanMX6, NDJ1-3HA ::KanMX6/NDJ1-

         3HA::KanMX6

PRY236   *MAT**a**/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG,*

         ura3/ura3, MPS3-DDD(T188D S189D S190D–BSPEI)/MPS3-

         DDD(T188D S189D S190D–BSPEI), Rap1-

         GFP::LEU2/Rap1-GFP::LEU2

PRY301   *MAT**a**/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG,*

         ura3/ura3, MPS3-DDD(T188D S189D S190D–BSPEI)-

         3GFP::KanMX6/ MPS3-DDD(T188D S189D S190D–BSPEI)-

         3GFP::KanMX6, cdc28-as1/cdc28-as1

PRY272   *MAT**a**/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG,*

         ura3/ura3, MPS3-DDD(T188D S189D S190D–BSPEI)-

         3GFP::KanMX6/ MPS3-DDD(T188D S189D S190D–BSPEI)-

         3GFP::KanMX6, cdc7-as3::9MYC/cdc7-as3::9MYC

HKY167   *MAT**a**/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG,*

         ura3/ura3, RAP1-GFP::LEU2/RAP1-GFP::LEU2

PRY68   *MAT**a**/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG,*

         ura3/ura3, RAP1-GFP::LEU2/RAP1-GFP::LEU2, cdc28-

         as1/cdc28-as1

PRY79   *MAT**a**/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG,*

	ura3/ura3, RAP1-GFP::LEU2/RAP1-GFP::LEU2, cdc7-as3-9MYC/cdc7-as3-9MYC
PRY116	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, RAP1-GFP::LEU2/RAP1-GFP::LEU2, cdc7::KanMX6/ cdc7::KanMX6, bob1-1/bob1-1*
PRY303	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, MPS3-DDD(T188D S189D S190D–BSPEI)-3GFP::KanMX6/ MPS3-DDD(T188D S189D S190D–BSPEI)-3GFP::KanMX6, Rap1-GFP::LEU2/Rap1-GFP::LEU2, cdc28-as1/cdc28-as1*
PRY309	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, MPS3-DDD(T188D S189D S190D–BSPEI)-3GFP::KanMX6/ MPS3-DDD(T188D S189D S190D–BSPEI)-3GFP::KanMX6, Rap1-GFP::LEU2/Rap1-GFP::LEU2, cdc7-as3::9MYC/cdc7-as3::9MYC*
PRY322	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, MPS3-DDD(T188D S189D S190D–BSPEI)-3GFP::KanMX6/ MPS3-DDD(T188D S189D S190D–BSPEI)-3GFP::KanMX6, Rap1-GFP::LEU2/Rap1-GFP::LEU2, cdc7::KanMX6/ cdc7::KanMX6, bob1-1/bob1-1*
PRY192	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, MPS3-3GFP::KanMX6/MPS3-3GFP::KanMX6 ndj1::KanMX6/ndj1::KanMX6*
PRY198	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, MPS3-3GFP::KanMX6/MPS3-3GFP::KanMX6,*

	csm4::KanMX6/csm4::KanMX6
PRY514	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, Mps3-MER2S (swap)-SacI-Flag/ Mps3-MER2S (swap)-SacI-Flag, Rap1-GFP::LEU2/Rap1-GFP::LEU2, NDJ1-HA/NDJ1-HA*
PRY518	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, Mps3-MER2S (swap)-SacI-Flag/ Mps3-MER2S (swap)-SacI-Flag, Rap1-GFP::LEU2/Rap1-GFP::LEU2, Ndj1-HA/ Ndj1-HA*
KSY220/221	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, mps3-S190A (-PSTI)-3GFP::KanMX6/mps3-S190A(-PSTI)-3GFP::KanMX6*
KSY407/409	*MATa/α, ho::LYS2/ho::LYS2, lys2/lys2, leu2::hisG/leu2::hisG, ura3/ura3, mps3-S189A-(BSPEI)-3GFP::KanMX6/mps3-S189A-(BSPEI)-3GFP::KanMX6*

Supplemental Figure S4 Rao et. al.