Individualized cyclic mechanical loading improves callus properties during the remodelling

phase of fracture healing in mice as assessed from time-lapsed *in vivo* imaging

Esther Wehrle[1], Graeme R Paul[1], Duncan C Tourolle né Betts[1], Gisela A Kuhn[1], Ralph Müller[1*]

[1] Institute for Biomechanics, ETH Zurich, Zurich, Switzerland.

Corresponding author:

Ralph Müller, PhD

Institute for Biomechanics

ETH Zurich

Leopold-Ruzicka-Weg 4

8093 Zurich, Switzerland

Email: ram@ethz.ch

Abstract

Fracture healing is regulated by mechanical loading. Understanding the underlying

mechanisms during the different healing phases is required for targeted mechanical

intervention therapies. Here, the influence of individualized cyclic mechanical loading on the

remodelling phase of fracture healing was assessed in a mouse femur defect model. After

bridging of the defect, a loading group (n=10) received individualized cyclic mechanical

loading (8-16 N, 10 Hz, 5 min, 3x/week) based on computed strain distribution in the callus

using animal-specific real-time micro-finite element analysis. Controls (n=10) received 0 N

treatment at the same post-operative time-points. By registration of consecutive scans,

structural and dynamic callus morphometric parameters were followed in three callus sub-

volumes and the adjacent cortex showing that the remodelling phase of fracture healing is

highly responsive to cyclic mechanical loading with changes in dynamic parameters leading

to significantly larger callus formation and mineralization. Loading-mediated maintenance of

callus remodelling was associated with distinct effects on Wnt-signalling-associated

molecular targets Sclerostin and Rankl in callus sub-regions and the adjacent cortex. Given

these distinct local protein expression patterns induced by cyclic mechanical loading, the

femur defect loading model with individualized load application seems suitable to

understand the local spatio-temporal mechano-molecular regulation of the different

fracture healing phases.

Introduction

Mechanical loading is a key factor for normal progression of the fracture healing process.

Delayed fracture repair and non-union formation is a major issue in orthopaedic surgery,

with an incidence of 5-10% and a high socioeconomic burden [1,2] and has been partially

attributed to low and inadequate mechanical loading of the defect region [3,4], affecting all

consecutive phases (inflammation, repair, remodelling) of the fracture repair process. The

local mechanical conditions in the fracture healing area (for details see tissue differentiation

hypothesis by Claes & Heigele [5]) are crucial for the repair process by determining molecular

and cellular reactions, the tissue to be formed and the type of ossification. We and others

have previously shown that externally applied mechanical stimuli, either applied locally to

the fracture site via dynamic fixators [6] or globally via whole body vibration platforms [7,8], can

improve fracture healing. In contrast, inadequate mechanical stimuli resulting in a too low or

too high interfragmentary motion were not successful or even impaired the healing process

[9-11]; for review see [12]). In order to understand the underlying mechanisms, refined and well

controlled *in vivo* loading models are needed. Recent studies particularly indicate that

profound understanding of the local spatio-temporal mechanical regulation of the fracture

healing process seems crucial for translation of this knowledge to the clinics, aiming at the

development of safe, targeted and individualized mechanical intervention therapies.

Whereas many preclinical fracture healing studies focused on improving the early healing

phases (inflammation, repair) with the aim of increasing bone formation and achieving

earlier bone union, recent studies also indicate a potential to improve fracture healing

outcome via modulation of callus remodelling [13-15]. Given the strong influence of mechanical

loading on normal bone remodelling, the application of controlled mechanical loading might

be suitable to also improve and accelerate the remodelling phase during fracture healing.

However, preclinical fracture healing studies with application of loading have so far shown

contradictory results [12] and have not been able to completely capture underlying

mechanisms due to restrictions in study design: (I) cross-sectional setup with only end-point

analyses, (II) effect assessment not specific to healing phases, (III) same load for all animals

irrespective of individual callus properties and healing progression. We aim to overcome

current limitations by combining state-of-the-art time-lapsed *in vivo* imaging with a novel

precisely controlled femur defect loading system in mice. By applying time-lapsed *in vivo*

micro-CT and animal-specific real-time micro-finite element analysis (micro-FEA), the

developed femur defect loading model allows scaling of loading settings based on strain

distribution in the callus [16], thereby considering individual differences in healing speed. To

follow the healing progression in each animal, a recently developed longitudinal *in vivo*

micro-CT based approach was applied, which was previously shown to capture the different

healing phases and to discriminate physiological and impaired healing patterns [17], without

significant radiation-associated effects on callus properties [18]. By registering consecutive

scans of the defect region for each animal and implementing a two-threshold approach, data

on bone turnover and mineralization kinetics can be obtained, which is particularly

important for targeting callus remodelling under different conditions.

The objective of this study was to assess the influence of individualized cyclic mechanical

loading on the remodelling phase of fracture healing via longitudinal *in vivo* micro-CT

imaging. We hypothesized, that individualized cyclic mechanical loading improves structural

and dynamic callus properties during the remodelling phase of fracture healing.

Results

By combining a novel *in vivo* femur defect loading model (Fig. 1 and Supplementary Fig. 1) with a recently established time-lapsed *in vivo* micro-CT based monitoring approach [18,19], the influence of individualized cyclic mechanical loading on callus properties was assessed during the remodelling phase of fracture healing. A loading group (n=10) received individualized cyclic mechanical loading (8-16 N, 10 Hz, 5 min, 3x/week) whereas controls (n=10) received 0N for 5 min at the same post-operative time-points. By registration of consecutive scans, structural and dynamic callus parameters were followed in three callus sub-volumes (defect centre: DC, defect periphery: DP, cortical fragment periphery: FP) and the adjacent cortical fragments (FC) over time (Fig. 2 and 3), for details on methods see Tourolle et al. 2019 [19]; for detailed study design see Supplementary Table S1).

General physical observation

All mice recovered rapidly from surgery. During the healing period, the animals' body weight did not significantly change compared to pre-operative values without significant differences between or within the two groups (see Supplementary Fig. S2). We also assessed the social interaction between mice (animals sitting in groups vs. separated from others) and the nesting behaviour (start of nest building after surgery). Specifically, all animals started to build nests on the day of surgery and were found in groups in the nest in the morning of post-operative day 1. Throughout the further healing period social interaction between mice and nesting behaviour did not differ from pre-surgical observations and was similar for animals of the scan and control groups.

Volumes of interest (VOI) for evaluation by time-lapsed *in vivo* micro-CT

In order to exclude bias in the further micro-CT analyses, we compared the size of the different VOIs (depicted in Fig. 3; for details on VOI generation see Tourolle et al., 2020 [19]). The VOIs encompassed the following volume for the control (n=10) and the loading group (n=10): $1.14\pm0.13mm^3$ vs. $1.04\pm0.16mm^3$ for the defect centre (DC), $6.68\pm0.77mm^3$ vs. $6.35\pm0.82mm^3$ for the defect periphery (DP), $2.28\pm0.17mm^3$ vs. $2.31\pm0.16mm^3$ for the cortical fragments (FC), $15.44\pm0.70mm^3$ vs. $15.84\pm0.99mm^3$ for the fragment periphery (FP). No significant differences in volume were detected in any of the VOIs between groups.

Pre-loading healing patterns monitored by time-lapsed *in vivo* micro-CT

In the pre-loading healing period (week 0 to week 3) similar and physiological healing patterns were observed for the control group (n=10) and the loading group (n=10) in all defect VOIs (DC, DP and FP) and the adjacent cortex (FC; Fig. 2 and 3). Registration of consecutive *in vivo* micro-CT scans allowed to capture distinct callus characteristics indicative of the different healing phases (inflammation, repair, remodelling; significant weekly differences in bone parameters indicated in Fig. 3). From week 0-1 to week 1-2 in both groups a significant 4x to 14x increase in bone formation was detected in the three callus VOIs DC (control group: p=0.0007; loading group: p=0.0006; Fig. 3a), DP (control group: p<0.0001; loading group: p<0.0001; Fig. 3d) and FP (control group: p=0.0012; loading group: p=0.0006; Fig. 3j), indicating progression from the inflammation to the reparative phase. This led to a significant gain in bone volume (BV/TV) by week 2 in the callus VOIs DC (control group: p<0.0001; loading group: p<0.0001; Fig. 3b), DP (control group: p<0.0001; loading group: p<0.0001; Fig. 3e) and FP (control group: p<0.0001; loading group: p<0.0001;

Fig. 3k). From week 1-2 to week 2-3 a significant 3x to 9x increase in resorptive activities was

seen in the callus VOIs DC (control group: p=0.0023; loading group: p<0.0001; Fig. 3a), DP

(control group: p=0.0499; loading group: p=0.0162; Fig. 3d) and FP (control group: p=0.0326;

loading group: p=0.0070; Fig. 3j), indicating the progression from the repair to the

remodelling phase. From week 2 to week 3, the highly mineralized bone fraction in the callus

VOIs significantly increased (p<0.0001 in DC, DP, FP) in both groups, indicating callus

maturation (Fig. 3c, 3f and 3l).

From week 0 to week 3, in both groups, strong resorptive activities (Fig. 2 and 3g) were seen

in the adjacent cortical fragments (FC). BRR showed significantly weekly increases whereas

no significant weekly changes in bone formation were observed in this VOI. The strong

resorptive activities let to decreased bone volume by week 3 reaching statistical significance

in the loading group (p<0.0001). Furthermore, the fraction of highly mineralized bone

significantly decreased from week 1 to week 3 in both groups (p<0.0001), indicating cortical

reorganisation.

Individualized cyclic mechanical loading improves callus properties during the remodelling

phase of fracture healing

Individualized cyclic mechanical loading (8-16N, 10 Hz, 300 cycles, 3x/week) started after

bridging of the femur defect (week 3; Table 1) at the beginning of the remodelling phase

(loading group, n=10). RT-micro-FE allowed determination of initial loads (8-12N) based on

individual callus properties, weekly load-scaling (8-16N) and matching of strain distribution

in the callus for the 4-week loading period. Control animals received 0N loading at the same

post-operative time-points. Already after 1 week, the first loading-associated effects on

callus properties and the adjacent cortical bone were seen: Specifically, loading reduced BRR

in all VOIs in week 3-4 compared to pre-loading values in week 2-3 (Fig. 3a, d, g, j) reaching

statistical significance in DC (p<0.0001). In contrast, in the control group, BRR further

increased in all VOIS during the same period with statistical significance seen in the

peripheral VOIs (p=0.0066 for DP, p=0.0126 for FP). From week 3-4 to week 6-7, the applied

individualized mechanical loading was also able to maintain bone formation at significantly

higher levels compared to controls, while bone resorption was significantly reduced during

the same period. This led to a linear increase in osseous callus volume in loaded animals

from week 3 to week 7 in all callus VOIs (DC: +41 %, DP: +130 %, FP: +111 %) and the

adjacent cortex (FC: +20 %), whereas it significantly declined in controls (DC: - 28 %, DP: - 44

%, FP: -27 %, FC: -30 %). In week 6-7, no significant differences in bone turnover were seen

anymore indicating load adaptation. After 7 weeks, loaded animals had significantly more

bone in the defect (2x for DC, p<0.0001; 4x for DP, p<0.0001; 3x for FP, p<0.0001) with a

significantly larger fraction of highly mineralized bone compared to controls (+8% for DC,

p=0.0006; +15% for DP, p<0.0015; +10% for FP, p<0.0001). Three-point bending also showed

a 46% higher relative bending stiffness (healed femur vs. contralateral femur) in the loaded

animals (107%) compared to controls (73%; n=2/group). In addition, according to the

standard clinical evaluation of X-rays, the number of bridged cortices per callus was

evaluated in two perpendicular planes and animals with ≥3 bridged cortices were

categorized as healed. From week 4 to week 7 all defects in both groups were categorized as

healed (Table 1), indicating that the applied forces during cyclic mechanical loading did not

lead to re-fracture of the callus.

Loading also affected the bone turnover in the adjacent cortex (FC). Already after 1 week,

loaded animals showed significantly lower cortical bone resorption compared to controls

(Fig. 3 j). Whereas cortical volume significantly declined (p<0.0001) in control animals from

week 3 to week 7 by 27%, it significantly increased (p<0.0001) by 20% in the loaded animals

during the same time period (Fig. 3k). From week 4 to week 7 loading was also associated

with significantly higher cortical mineralization (+4% in week 4 to +54% in week 7) compared

to controls (week 4: p=0.0459, week 5 – week 7: p<0.0001; Fig. 3l)).

In week 7, the FC VOI comprised 46% and 34% of the osseous tissue in the total VOI (TOT)

for the control and loading group, respectively. In the DC VOI, 20% (control group) and 17%

(loading group) of the total bone volume were seen. In the two peripheral VOIs, 9% (control

group) and 14% (loading group) of the total osseous tissue were seen in the DP and 26%

(control group) and 35% (loading group) in the FP VOI. Loading was associated with a shift in

the bone distribution from the cortical fragments to the peripheral VOIs.

Ex vivo, 3-point-bending testing was performed on healed femora and contralateral femora

of two animals from the loading and the control group. The relative bending stiffness (healed

femur vs. contralateral femur) was 107% in the loading group and 73% in the control group,

indicating loading-associated improved functional healing outcome.

Histology

In order to also visualize the tissue composition of the callus, we performed end point

histological stainings of serial sections from one animal of each group. Histology supported

our micro-CT findings with complete cortical bridging seen in the animal from the loading

and from the control group (Fig. 4 B-I). However, the loaded animal showed a larger callus

with some cartilage fractions suggesting ongoing endochondral ossification processes,

whereas in the control animal the callus was largely remodelled with restoration of the

medullary cavity indicating proceeding towards the end of healing (Fig. 4 b-e). Furthermore,

differences in the shape of the cell nuclei were observed between the osteocytes in the

fracture callus (round shape) of the loaded animal in comparison to the cell nuclei seen in

the cortical bone of the same animal (ellipsoidal shape) and the cell morphology seen in the

control animals (Fig. 5 a+b), suggesting an association between the local mechanical

environment and cellular morphology. In regions with lower strains visualized by micro-FE

(endosteal fracture callus; Fig. 4a and 5b), rounder cell nuclei were observed, compared to

ellipsoidal nuclei shape in higher strained regions (cortex; Fig. 4a and 5b). To capture

potential underlying mechano-molecular targets, we performed immunohistochemistry of

Sclerostin (inhibitor of the mechano-responsive and osteoanabolic Wnt signalling pathway)

and Rankl (negatively-regulated target gene of Wnt signalling) which have both previously

been associated with the mechanical regulation of bone adaptation and healing [8,20-24].

Sclerostin staining seemed less abundant in the fracture callus compared to the cortical bone

of the loaded animal (Fig. 4 f and 5b) with no region-specific differences in staining patterns

being seen in the control animal (Fig. 4g and Fig. 5b), suggesting that Wnt-signalling

contributes to the loading-mediated osteoanabolic effects on fracture healing seen in this

study. Surprisingly, at the same time Rankl, which is down-regulated by Wnt-signalling, was

more abundantly and stronger expressed in the callus of the loaded animal compared to the

control animal (Fig. 4 h+i and Fig. 5b), indicating that not only Wnt-signalling, but also other

molecular factors and signalling pathways have to be involved in the observed mechanically-

mediated effects on callus properties during the remodelling phase of fracture healing. In

line with previous studies, strong Rankl expression was seen in lowly strained callus regions

as visualized via micro-FE analysis (Fig. 4a and 5b), indicating ongoing bone remodelling.

Discussion

In this study, the effect of individualized cyclic mechanical loading on the remodelling phase

of fracture healing was assessed in a novel femur defect loading model in mice using a

recently developed *in vivo* micro-CT based monitoring approach. Given the strong mechano-

responsiveness of bone cells, several studies have applied mechanical loading (e.g. globally

via vibration platforms, or locally via external fixators) to improve fracture healing showing

contradictory results. Whereas some studies saw a loading-mediated improvement of

fracture healing, others were not successful or even reported an impairment of the fracture

healing process [9-11]. One limitation of these studies was the application of the same load to

all animals irrespective of individual differences in healing progression, which bears the risk

of re-fracturing lowly mineralized bone after initial bridging of the defect. In order to allow

for individualized mechanical loading we now developed a precisely controlled *in vivo*

loading model based on external fixation of femur defects in mice. By combining state-of-

the-art time-lapsed *in vivo* imaging with individualized mechanical loading protocols adapted

to animal-specific callus properties, this allows to precisely characterize spatio-temporal

effects of mechanical loading on the fracture callus. Given the high mechano-responsiveness

of bone cells during normal bone remodelling and as recent studies also indicate a potential

to improve fracture healing outcome via modulation of callus remodelling [13-15], we focused

this study on the effect of mechanical loading on callus properties during the remodelling

phase of fracture healing.

Via weekly *in vivo* micro-CT imaging, we saw, that cyclic mechanical loading linearly

increased the osseous callus volume during the remodelling phase of fracture healing, so

that by the study end, loaded animals had significantly more bone in all callus VOIs (2x for

DC, p<0.0001; 4x for DP, p<0.0001; 3x for FP, p<0.0001) with a significantly larger fraction of

highly mineralized bone compared to controls (+8% for DC, p=0.0006; +15% for DP,

p<0.0015; +10% for FP, p<0.0001). This is supported by a recent study from Liu et al. [25]

describing additional loading-mediated bone formation during the remodelling phase in a

uni-cortical tibia defect model in mice. However, they failed to show statistical significance

between the loading and the control group in histomorphometry and micro-CT analyses,

which they attributed to the high variability within groups. As the study applied the same

load to all animals irrespective of healing progression, differences in the induced strains

might have caused the high variability within groups. We addressed this by using our

recently developed real-time micro-FE based loading approach [16], which takes into account

the individual strain distribution within the callus for determining individual loading settings

at weekly post-operative intervals. So far, no study using a complete defect model assessed

the effect of local cyclic mechanical loading (e.g. via external fixators, tibia/ulna-loading)

when exclusively applied during the remodelling phase of the healing process. An earlier

tibia defect loading study (defect size: 3 mm) in sheep by Goodship et al. [6] applied cyclic

mechanical loading (25 µm interfragmentary movement, 30 Hz) throughout the healing

period (week 0 - week 10, 5x per week) and reported a significantly larger callus with higher

mineralization in loaded animals compared to controls. However, as the loading application

was not restricted to specific post-operative time frames, it was not possible to evaluate

whether loading during a specific fracture healing period was sufficient for the observed

improvement in fracture healing or if cyclic mechanical loading needs to be applied

throughout the healing process to achieve such an effect. We now showed, that the

remodelling phase is highly responsive to cyclic mechanical loading applied after initial

cortical bridging of the defect, with pronounced effects on callus size and mineralization. Via

limiting the loading application to the earlier healing period (day 7 – day 19), Gupta et al. [26]

aimed at assessing (histomorphometry and radiography on day 42) whether there is a

sustained effect of cyclic mechanical loading (2%, 10% and 30% strain, 0.5 Hz, 500 cycles) on

fracture healing outcome. They found a larger bone fraction in the callus and higher cortical

bridging scores in the loaded compared to the control animals, indicating, that loading

effects persisted after a 4-week non-loading period. In contrast to these studies, Schwarz et

al. [27] did not observe any loading-mediated improvement on fracture healing in a critically-

sized femur defect model in rats (defect size: 5mm). However, in this study loading (6 cycles

of constant loading at 500 μm IFM and 40s rest phase), was only applied once per week,

which indicates a critical role of the type of loading (constant vs. cyclic) and the frequency of

loading sessions per week. Whether cyclic mechanical loading in contrast to constant

mechanical loading is able to prevent non-union formation in critically-sized defects needs to

be assessed in future studies.

Via registration of longitudinal *in vivo* micro-CT scans (for details see Tourolle et al. 2019 [19]),

we were now for the first time able to also include dynamic parameters such as bone

formation and resorption in the monitoring of loading-mediated effects on fracture healing.

Compared to histology-based determination of the bone formation rate via *in vivo*

application of fluorescent dyes (e.g. calcein, alizarin), our micro-CT-based monitoring

approach with registration of consecutive scans also allows for assessment of the bone

resorption rate in individual animals, which was shown to be particularly suited for the

characterization of the remodelling phase during the fracture healing process [17,18]. The most

striking loading-mediated effect in the current study was the prompt and significant

reduction in bone resorption rate (week 3-4 vs. week 2-3), which was seen in the central and

periosteal VOIs (DC, FP) as early as 1 week after loading (Fig. 3A, 3J). Compared to control

animals, in which bone resorption still increased during this early remodelling period (week

3-4), loaded animals showed significantly lower bone resorption rates in all callus VOIs (DC,

DP, FP) in this first week of loading, indicating a strong mechano-responsiveness of the

fracture callus during this healing period. Most of the previous studies either assessed the

loading effects only after a longer loading period of minimum 2 weeks [27,28] or they did not

see significant loading-mediated effects on the fracture callus after shorter treatment

periods [8,10,26]. So far only one study by Leung et al. [29] was able to detect a loading-mediated

improvement of fracture healing (larger callus diameter) after a short loading period of 1

week. We were now able to strengthen these cross-sectional structural callus findings with

our longitudinal micro-CT data obtained for single animals. In respect to the underlying

mechanisms, we showed for the first time, that not only bone formation but also bone

resorption can be significantly modulated via cyclic mechanical loading during fracture

healing, indicating a potential for mechanical intervention therapies to improve impaired

fracture healing conditions associated with imbalanced bone remodelling. Furthermore,

implementation of our two-threshold approach showed cyclic mechanical loading to be also

effective in significantly advancing callus mineralization compared to control animals. Via

combining our novel loading model with time-lapsed *in vivo* imaging and a two-threshold

approach, we were able to characterize the spatio-temporal effects of loading during the

remodelling phase of fracture healing. Furthermore, the applied micro-FE-based loading

approach also allowed for spatio-temporal visualization of strains induced in the fracture

callus (Fig. 4a+b). In order to also visualize the tissue composition of the callus, we

performed end point histological stainings of serial sections from one animal of each group.

Histology supported our micro-CT findings with complete cortical bridging seen in both

groups. However, the loaded animal showed a larger callus with some cartilage fractions

suggesting ongoing endochondral ossification processes, whereas in the control animal the

callus was largely remodelled with restoration of the medullary cavity indicating proceeding

towards the end of healing. Furthermore, differences in the shape of the cell nuclei were

seen between the osteocytes in the fracture callus (round shape) of the loaded animal in

comparison to the cell nuclei seen in the cortical bone of the same animal (ellipsoidal shape)

and the cell morphology seen in the control animals. This is particularly interesting, as

previous *in vitro* studies found, that round cells are more sensitive to mechanical loading

compared to flat and elongated cells [30-32], suggesting that repeated cyclic mechanical loading

might be able to induce round cellular morphology of (early) osteocytes, thereby

maintaining the remodelling phase of fracture healing eventually leading to higher

mineralization of the newly formed tissue compared to controls. To capture potential

underlying molecular targets, we performed immunohistochemistry of Sclerostin (inhibitor

of the mechano-responsive and osteoanabolic Wnt signalling pathway) and Rankl

(negatively-regulated target gene of Wnt signalling) which have both previously been

associated with the mechanical regulation of bone adaptation and healing [8,20-24]. Whereas

Sclerostin staining seemed less abundant in the fracture callus compared to the cortical bone

of the loaded animal (Fig. 4f, Fig. 5b), no region-specific differences in staining patterns were

seen in the control animal (Fig. 4g, Fig. 5b). This might indicate that Wnt-signalling

contributes to the loading-mediated osteoanabolic effects on fracture healing seen in this

study. However, at the same time we saw more and stronger Rankl staining in the callus of

the loaded animal compared to the control animal (Fig. 4 h+i). This finding precludes that the

loading-mediated increases in Rankl expression seen in this study were solely mediated via

Wnt-signalling but rather an interplay with further signalling pathways must have triggered

this strong response in Rankl production. Via micro-FE analysis, we saw high Rankl

expression in the lowly strained endosteal fracture callus, indicating ongoing bone

remodelling. In order to shed light into the complex interplay between a multitude of

mechano-responsive molecules and signalling pathway, novel advents in RNAsequencing

including spatial transcriptomic approaches might, in future studies, open interesting

possibilities to understand the spatio-temporal mechanomics of fracture healing. In non-

fractured bone, some of these molecular technologies (e.g. microarrays, [33-35]) have already

been successfully applied to study mechano-molecular effects in well-controlled and

characterized loading models (e.g. ulna, tibia and tail loading) in rodents. Targets (e.g.

Sclerostin, Estrogen signalling) found via these comprehensive studies were then assessed in

detail in further studies using either models with local [36-38] or global load application [39]. As

the main underlying mechanisms of bone adaptation to loading were identified using local

models of loading (ulna, tibia ..), this indicates that comprehensive spatio-temporal

mechano-molecular analyses in fracture healing will require tightly regulated models with

local application of mechanical loading being more controllable and less prone to variation

induced by confounding factors compared to models with global loading application.

In summary, using our recently developed femur defect loading model in combination with

time-lapsed *in vivo* micro-CT and a two-threshold approach, we showed that the remodelling

phase of fracture healing is highly responsive to cyclic mechanical loading with changes in

dynamic callus parameters leading to larger callus formation and mineralization. Loading-

mediated maintenance of callus remodelling was associated with distinct effects on the Wnt-

signalling-associated molecular targets Sclerostin and Rankl in callus sub-regions and the

adjacent cortex as assessed via immunohistochemistry. Given these distinct local protein

expression patterns induced by external cyclic mechanical loading, our tightly-controlled

femur defect loading model with individualized load application could be used in future

studies to precisely understand the local spatio-temporal mechano-molecular regulation of

the different fracture healing phases. With our real-time micro-FE based approach for

individualized load application considering differences in defect geometry and healing

progression, the femur defect loading model could also be used to study the bone

regeneration capacity of biomaterials under load application. With further advents in

transcriptomics and registration techniques for different imaging technologies, the tightly

controlled femur defect loading model could widen our knowledge on the spatio-temporal

regulation fracture mechanomics relevant for application of mechanical intervention

therapies to clinical settings for improvement of impaired healing conditions.

Methods

Study design

By combining a novel *in vivo* femur defect loading model with a recently established time-

lapsed *in vivo* micro-CT based monitoring approach, the influence of individualized cyclic

mechanical loading on callus properties was assessed during the remodelling phase of fracture healing. All mice received a femur defect and post-operative micro-CT scans (vivaCT 40, ScancoMedical, Brüttisellen, Switzerland) were performed. After bridging of the femur defect (week 3), real-time micro-FE allowed determination of initial loads (8N-12N), and weekly load-scaling and matching of strain distribution in the callus for the 4 week loading period. The loading group (n=10) received individualized cyclic mechanical loading (8-16N, 10 Hz, 5 min, 3x/week) whereas controls (n=10) received 0N for 5 min at the same post-operative time-points. By registration of consecutive scans, structural and dynamic callus parameters were followed in three callus sub-volumes (defect centre: DC, defect periphery: DP, cortical fragment periphery: FP) and the adjacent cortical fragments (FC) over time (Figure 1, for details on methods see Tourolle et al. 2020[19]; for detailed study design see Supplementary Fig. S1).

Animals

All animal procedures were approved by the authorities (licence number: 36/2014; Kantonales Veterinäramt Zürich, Zurich, Switzerland). We confirm that all methods were carried out and reported in accordance with relevant guidelines and regulations (Swiss Animal Welfare Act and Ordinance (TSchG, TSchV) and ARRIVE guidelines). To study adult fracture healing female 12 week-old C57BL/6J mice were purchased from Janvier (Saint Berthevin Cedex, France) and housed in the animal facility of the ETH Phenomics Center (EPIC; 12h:12h light-dark cycle, maintenance feed (3437, KLIBA NAFAG, Kaiseraugst, Switzerland), 5 animals/cage) for 8 weeks. At an age of 20 weeks, all animals received a femur defect by performing an osteotomy with a 0.66mm Gigli wire saw as previously

described (group 1: control group, n=10; group 2: loading group, n=10; housing after

surgery: 2-3 animals/cage; for details on study design see Supplementary Table 1). All defect

surgeries were performed by the same veterinarian. Perioperative analgesia (25 mg/L,

Tramal®, Gruenenthal GmbH, Aachen, Germany) was provided via the drinking water two

days before surgery until the third post-operative day. For surgery and micro-CT scans,

animals were anaesthetized with isoflurane (induction/maintenance: 5%/1-2%

isoflurane/oxygen). Perioperative handling, and monitoring, micro-CT imaging and loading

application was performed by the surgeon.

Loading fixator assembly and group assignment

The four parts of the loading fixators (n=20) were assembled as depicted in Suppl. Fig. 1. To

allow optimal identification throughout the experiments, one side part was engraved with a

fixator-specific number (Suppl. Fig 1A, 1B). The stiffness of each fixator was measured using

a Zwick testing machine and the fixators were assigned to the two groups, to allow similar

distributions of fixator stiffness in the loading and control group (Suppl. Fig 1C).

Femur osteotomy

In all animals an external fixator (Mouse ExFix, RISystem, Davos, Switzerland; mean stiffness:

17N/mm; Suppl. Fig. 1) was positioned at the craniolateral aspect of the right femur and

attached using four mounting pins. First, the most proximal pin was inserted approximately

2mm proximal to the trochanter, followed by placement of the most distal and the inner

pins. Subsequently, a femur defect was created using a Gigli wire (diameter: 0.66mm).

Time-lapsed _in vivo_ micro-CT

Immediate post-surgery correct positioning of the fixator and the defect was visualized using

a vivaCT 40 (Scanco Medical AG, Brüttisellen, Switzerland) (isotropic nominal resolution: 10.5

µm; 2 stacks of 211 slices; 55 kVp, 145 µA, 350 ms integration time, 500 projections per

180°, 21 mm field of view (FOV), scan duration ca. 15 min). Subsequently, the fracture callus

and the adjacent bone between the inner pins of the fixator were scanned weekly using the

same settings. Scans were registered consecutively using a branching scheme (registration of

whole scan for bridged defects; separate registration of the two fragments for unbridged

defects)[5]. Subsequently, morphometric indices (bone volume - BV, bone volume/total

volume – BV/TV, bone formation rate – BFR, bone resorption rate - BRR) were computed

(threshold: 395mg HA/cm^3; for details on methods see Tourolle et al., 2020 [19]). To assess

mineralization progression, a second threshold (645mg HA/cm^3) was applied and the ratio

between highly and lowly mineralized tissue (BV_{645}/BV_{395}) was calculated. The two selected

thresholds are included in our recently developed multidensity threshold approach [19].

According to the standard clinical evaluation of X-rays, the number of bridged cortices per

callus was evaluated in two perpendicular planes (UCT Evaluation V6.5-1, Scanco Medical

AG, Brüttisellen, Switzerland). A ''healed fracture'' was considered as having a minimum of

at least three bridged cortices per callus.

For evaluation, four volumes of interest (VOIs) were defined, which were created

automatically from the post-operative measurement (Fig. 1): defect centre (DC), defect

periphery (DP), cortical fragment centre (FC), and fragment periphery (FP). Data were

normalised to the central VOIs: DC/DC, DP/DC, FC/FC, FP/FC.

Individualized cyclic mechanical loading

From week 4 to week 7, individualized cyclic loading (8-16N, 10Hz, 3000cycles; 3x/week;

controls - 0N) was applied via the external loading fixator (Suppl. Fig 1B) based on computed

strain distribution in the callus using animal-specific real-time micro-finite element (RT-

microFE) analysis (for detailed description of methods see [16]. Briefly, after weekly micro-CT

measurements of each animal, the images were pre-processed using threshold-binning to

create a high-resolution multi-density FE mesh, which was then solved on a supercomputer

within the same anaesthetic session for each mouse. 2D and 3D visualizations of the callus

were generated and a strain histogram was plotted. This information allowed load scaling

based on the strains induced in the callus in individualized animals. Specifically, the

distribution was scaled to achieve a median effective microstrain of 700. To assess regions

that posed a failure risk the 99th percentile strains of each simulation was visualised in both

image based and histogram forms. If more than 50 voxels exceeded 1% then the optimised

load was reduced by 1 N.

Mechanical testing

469 3 point bending tests were performed on a Zwick compression tester (ZwickRoell GmbH &

470 Co, Ulm, Germany). Tests were conducted on a setup of 8 mm support span and up to 3 N.

471 Tests were performed quasi-statically with a cross-head speed of 0.4 mm/min. Stiffness was

472 then calculated from the linear region of the resultant force-displacement curve. Each

473 sample was tested three times and the mean stiffness was taken.

Histology

Histological stainings were performed in one animal per group. On day 49, femora were excised, the femoral head was removed and the samples were placed in 4% neutrally buffered formalin for 24 hours and subsequently decalcified in 12.5% EDTA for 10-14 days. The samples were embedded in paraffin and the complete fracture callus was cut in 10 μm longitudinal serial sections. Every 10th section was stained with Safranin-O: Weigert's iron haematoxylin solution (HT1079, Sigma-Aldrich, St. Louis, MO) - 4min, 1:10 HCl-acidified 70% ethanol - 10s, tap water - 5min, 0.02% Fast Green (F7258, Sigma-Aldrich, St. Louis, MO) - 3min, 1% acetic acid - 10s, 0.1% Safranin-O (84120, Fluka, St. Louis, MO) - 5min. Images were taken with Slide Scanner Pannoramic 250 (3D Histech, Budapest, Hungary) at 20x magnification. Sections in between were stained for Sclerostin and Rankl.

For immunohistochemical staining of Sclerostin, nonspecific sites were blocked (1% BSA/PBS + 1% rabbit serum) for 60 min at room temperature. Subsequently, the sections were incubated with the primary antibody against Sclerostin (AF1589, R&D Systems, Minneapolis, MN; 1:150 in 1%BSA/PBS + 0.2% rabbit serum) overnight at 4°C. To detect the primary antibody, a secondary biotinylated rabbit anti-goat-IgG antibody (BAF017, R&D Systems, Minneapolis, MN) was added for 1 h at room temperature. For signal amplification, the slides were incubated with avidin-biotin complex (PK-6100 Vector Laboratories, Burlingame, CA) for 30 min. Diaminobenzidine (Metal Enhanced DAB Substrate Kit, 34065 ThermoFisher Scientific, Waltham, MA) was used as detection substrate. Counterstaining was performed with FastGreen (F7258, Sigma-Aldrich, St. Louis, MO).

For immunohistochemical staining of Rankl, slides were placed into TBS-Tween buffer and inserted into the Dako-Autostainer (Dako, Ft. Collins, USA) using the following staining

protocol: incubation with primary antibody against Rankl (1:100; ab 9957, Abcam,

Cambridge, UK) at 4°C overnight, rinsing with TBS-Tween, peroxidase blocking for 10min at

room temperature, rinsing with TBS-Tween, Envision+System HPR Rabbit (Dako K4003) for

60 min at room temperature, rinsing with TBS-Tween and incubation with DAB (Dako

K3468) for 10 min at room temperature. The slides were rinsed in A. dest and

counterstained with FastGreen for 2sec.

For both immunohistochemical stainings, species-specific IgG was used as isotype control.

Images were taken with Slide Scanner Pannoramic 250 (3D Histech, Budapest, Hungary) at

40x magnification.

Statistics

Data were tested for normal distribution (Shapiro-Wilk-Test) and homogeneity of variance

(Levene-Test). Depending on the test outcome, group comparisons (loading vs. control

group) of data derived at single time points (VOI size) were performed by two-tailed

Student's t-test or Mann-Whitney U-test (IBM SPSS Statistics Version 23). For statistical

evaluation of repeated measurements two-way ANOVA with Geisser-Greenhouse correction

and Bonferroni correction (GraphPad Prism 8) were performed. The level of significance was

set at $p < 0.05$.

Acknowledgements

We thank Marco Hitz, Peter Schwilch, Angad Malhotra and Jianhua Zhang for technical and

surgery assistance. We thank Christopher M. O'Neill for the micro-CT and micro-FE images

depicted in Figure 4. The authors gratefully acknowledge support from the EU (BIODESIGN

FP7-NMP-2012-262948 and ERC Advanced MechAGE ERC-2016-ADG-741883). E. Wehrle

received funding from the ETH Postdoctoral Fellowship Program (MSCA-COFUND, FEL-

25_15-1).

Author Contributions Statement

The study was designed by E.W., G.A.K., and R.M.. The experiments were performed by

E.W. and G.R.P. Data analyses were performed by E.W., G.R.P. and D.C.B.. The manuscript

was written by E.W. and reviewed and approved by all authors.

Data availability

All necessary data generated or analysed during the present study are included in this

published article and its Supplementary Information files (preprint available on BioRxiv

(BIORXIV/). Additional information related to this paper may be requested from the authors.

Competing Interests

The authors declare no competing interests.

References

1   Einhorn, T. A. & Gerstenfeld, L. C. Fracture healing: mechanisms and interventions. *Nat Rev*
Rheumatol **11**, 45-54, doi:10.1038/nrrheum.2014.164 (2015).
2   Gomez-Barrena, E. *et al.* Bone fracture healing: Cell therapy in delayed unions and
nonunions. *Bone* **70**, 93-101, doi:10.1016/j.bone.2014.07.033 (2015).
3   Bottlang, M. *et al.* Effects of Construct Stiffness on Healing of Fractures Stabilized with
Locking Plates. *J Bone Joint Surg Am* **92a**, 12-22, doi:10.2106/Jbjs.J.00780 (2010).
4   Steiner, M., Claes, L., Ignatius, A., Simon, U. & Wehner, T. Numerical Simulation of Callus
Healing for Optimization of Fracture Fixation Stiffness. *Plos One* **9**, doi:ARTN
e10137010.1371/journal.pone.0101370 (2014).

5   Claes, L. E. & Heigele, C. A. Magnitudes of local stress and strain along bony surfaces predict
the course and type of fracture healing. *J Biomech* **32**, 255-266, doi:Doi 10.1016/S0021-
9290(98)00153-5 (1999).
6   Goodship, A. E., Lawes, T. J. & Rubin, C. T. Low-Magnitude High-Frequency Mechanical
Signals Accelerate and Augment Endochondral Bone Repair: Preliminary Evidence of Efficacy.
J Orthop Res **27**, 922-930, doi:10.1002/jor.20824 (2009).
7   Shi, H. F., Cheung, W. H., Qin, L., Leung, A. H. C. & Leung, K. S. Low-magnitude high-frequency
vibration treatment augments fracture healing in ovariectomy-induced osteoporotic bone.
Bone **46**, 1299-1305, doi:10.1016/j.bone.2009.11.028 (2010).
8   Wehrle, E. *et al.* The impact of low-magnitude high-frequency vibration on fracture healing is
profoundly influenced by the oestrogen status in mice. *Dis Model Mech* **8**, 93-104,
doi:10.1242/dmm.018622 (2015).
9   Stuermer, E. K. *et al.* Musculoskeletal Response to Whole-Body Vibration During Fracture
Healing in Intact and Ovariectomized Rats. *Calcified Tissue Int* **87**, 168-180,
doi:10.1007/s00223-010-9381-0 (2010).
10 Wehrle, E. *et al.* Distinct Frequency Dependent Effects of Whole-Body Vibration on Non-
Fractured Bone and Fracture Healing in Mice. *J Orthop Res* **32**, 1006-1013,
doi:10.1002/jor.22629 (2014).
11 Wolf, S. *et al.* Effects of high-frequency, low-magnitude mechanical stimulus on bone healing.
Clin Orthop Relat R, 192-198 (2001).
12 Wang, J., Leung, K. S., Chow, S. K. H. & Cheung, W. H. The Effect of Whole Body Vibration on
Fracture Healing - a Systematic Review. *Eur Cells Mater* **34**, 108-127,
doi:10.22203/eCM.v034a08 (2017).
13 Bao, Q. W. *et al.* An appropriate Wnt/beta-catenin expression level during the remodeling
phase is required for improved bone fracture healing in mice. *Sci Rep-Uk* **7**, doi:ARTN
269510.1038/s41598-017-02705-0 (2017).
14 Hegde, V., Jo, J. E., Andreopoulou, P. & Lane, J. M. Effect of osteoporosis medications on
fracture healing. *Osteoporosis Int* **27**, 861-871, doi:10.1007/s00198-015-3331-7 (2016).
15 Schindeler, A., McDonald, M. M., Bokko, P. & Little, D. G. Bone remodeling during fracture
repair: The cellular picture. *Semin Cell Dev Biol* **19**, 459-466,
doi:10.1016/j.semcdb.2008.07.004 (2008).
16 Paul, G. R., Wehrle, E., Tourolle né Betts, D. C., Kuhn, G. & Müller, R. Reducing variance in a
mouse defect healing model: Real-time Finite Element Analysis allows homogenization of
tissue scale strains *bioRxiv*, doi:10.1101/2020.09.02.274878 (2020).
17 Wehrle, E. *et al.* In vivo micro-CT based approach for discrimination of physiological and
impaired healing patterns in mouse femur defect models. . *European Cell and Materials* **31**,
168 (2016).
18 Wehrle, E. *et al.* Evaluation of longitudinal time-lapsed in vivo micro-CT for monitoring
fracture healing in mouse femur defect models. *Sci Rep-Uk* **9**, doi:ARTN
1744510.1038/s41598-019-53822-x (2019).
19 Tourolle né Betts D.C., W. E., Pau G.R., Kuhn G.A., Christen P., Hofmann S, Müller R. The
association between mineralised tissue formation and the mechanical local in vivo
environment: Time-lapsed quantification of a mouse defect healing model. *Sci Rep-Uk* **10**,
doi:https://doi.org/10.1038/s41598-020-57461-5 (2020).
20 Li, C. *et al.* Sclerostin Antibody Treatment Enhances Fracture Healing and Increases Bone
Mass and Strength in Non-Fractured Bones in an Adult Rat Closed Femoral Fracture Model.
Osteoporosis Int **22**, 11-12 (2011).
21 Mcdonald, M. M. *et al.* Bone volume is not altered with unloading in the homozygous
sclerostin knockout mouse. *Bone* **50**, S94-S94, doi:10.1016/j.bone.2012.02.282 (2012).

22   Ominsky, M. S. *et al.* Inhibition of Sclerostin by Monoclonal Antibody Enhances Bone Healing
and Improves Bone Density and Strength of Nonfractured Bones. *Journal of Bone and*
Mineral Research **26**, 1012-1021, doi:10.1002/jbmr.307 (2011).
23   Robling, A. G. *et al.* Mechanical stimulation of bone in vivo reduces osteocyte expression of
Sost/sclerostin. *J Biol Chem* **283**, 5866-5875, doi:10.1074/jbc.M705092200 (2008).
24   Suen, P. K. *et al.* Sclerostin Monoclonal Antibody Enhanced Bone Fracture Healing in an Open
Osteotomy Model in Rats. *J Orthop Res* **32**, 997-1005, doi:10.1002/jor.22636 (2014).
25   Liu, C. *et al.* Effects of mechanical loading on cortical defect repair using a novel
mechanobiological model of bone healing. *Bone* **108**, 145-155,
doi:10.1016/j.bone.2017.12.027 (2018).
26   Gupta, R. R. *et al.* Axial strain enhances osteotomy repair with a concomitant increase in
connexin43 expression. *Bone Res* **3**, doi:ARTN 1500710.1038/boneres.2015.7 (2015).
27   Schwarz, C. *et al.* Mechanical Load Modulates the Stimulatory Effect of BMP2 in a Rat
Nonunion Model. *Tissue Eng Pt A* **19**, 247-254, doi:10.1089/ten.tea.2012.0265 (2013).
28   Tufekci, P. *et al.* Early mechanical stimulation only permits timely bone healing in sheep. *J*
Orthop Res **36**, 1790-1796, doi:10.1002/jor.23812 (2018).
29   Leung, K. S. *et al.* Low-Magnitude High-Frequency Vibration Accelerates Callus Formation,
Mineralization, and Fracture Healing in Rats. *J Orthop Res* **27**, 458-465,
doi:10.1002/jor.20753 (2009).
30   van Oers, R. F. M., Wang, H. & Bacabac, R. G. Osteocyte Shape and Mechanical Loading. *Curr*
Osteoporos Rep **13**, 61-66, doi:10.1007/s11914-015-0256-1 (2015).
31   Vatsa, A. *et al.* Osteocyte morphology in fibula and calvaria - Is there a role for
mechanosensing? *Bone* **43**, 452-458, doi:10.1016/j.bone.2008.01.030 (2008).
32   Wu, V. *et al.* Osteocyte morphology and orientation in relation to strain in the jaw bone. *Int J*
Oral Sci **10**, doi:Unsp 210.1038/S41368-017-0007-5 (2018).
33   McKenzie, J. A., Bixby, E. C. & Silva, M. J. Differential Gene Expression from Microarray
Analysis Distinguishes Woven and Lamellar Bone Formation in the Rat Ulna following
Mechanical Loading. *Plos One* **6**, doi:ARTN e2932810.1371/journal.pone.0029328 (2011).
34   Wasserman, E. *et al.* Differential load-regulated global gene expression in mouse trabecular
osteocytes. *Bone* **53**, 14-23, doi:10.1016/j.bone.2012.11.017 (2013).
35   Zaman, G. *et al.* Loading-related regulation of gene expression in bone in the contexts of
estrogen deficiency, lack of estrogen receptor alpha and disuse. *Bone* **46**, 628-642,
doi:10.1016/j.bone.2009.10.021 (2010).
36   Lara-Castillo, N. *et al.* In vivo mechanical loading rapidly activates beta-catenin signaling in
osteocytes through a prostaglandin mediated mechanism. *Bone* **76**, 58-66,
doi:10.1016/j.bone.2015.03.019 (2015).
37   Melville, K. M. *et al.* Effects of Deletion of ER in Osteoblast-Lineage Cells on Bone Mass and
Adaptation to Mechanical Loading Differ in Female and Male Mice. *Journal of Bone and*
Mineral Research **30**, 1468-1480, doi:10.1002/jbmr.2488 (2015).
38   Robling, A. G. *et al.* Sost, independent of the non-coding enhancer ECR5, is required for bone
mechanoadaptation. *Bone* **92**, 180-188, doi:10.1016/j.bone.2016.09.001 (2016).
39   Moura, M. L. A. *et al.* Estrogen therapy associated with mechanical vibration improves bone
microarchitecture and density in osteopenic female mice. *J Anat* **233**, 715-723,
doi:10.1111/joa.12893 (2018).
40   Paul, G. R., Schwilch, P., Wehrle, E., Kuhn, G. A. & Müller, R. Development of a micro-CT
mouse holder for a femur defect mouse model. *bioRxiv*, doi:doi:
https://doi.org/10.1101/2020.09.10.289553 (2020).

 Figure Legends

Figure 1. Loading system for femur defects in mice. a. Loading fixator (RISystem, Davos,

Switzerland) consisting of four PEEK parts connected via two transverse Titanium pins. b.

Insertion of loading fixator into loading device with clamping of loading adapter (1) and

removal of transverse pin (2). c. Loading device consisting of mouse holder with anaesthesia

inlet (1), heated ground plate (2), clamp for fixator (3) and actuation system with loading

chuck (4), load cell electronics box (5), electro-magnetic actuator (6), actuator control

electronic box (7). The mouse holder can be used for the loading device and the CT scanner

without the need to change the position of the mouse. For detailed description and

drawings see Paul et al. [40].

Figure 2. Representative images (full image and cut; threshold: 645 mg HA/cm^3) of the

defect region from animals of the loading group (week 0-7) and the control group (week 0-

7). Visualization of bone formation (orange) and resorption (blue) via registration of micro-

CT scans from weeks 1-6 to weeks 0-5.

Figure 3. Micro-CT based evaluation of bone parameters in the loading group (red) and the

control group (blue) using different VOIS (pre-loading healing period highlighted in lighter

color; loading period highlighted in darker color): defect center (DC; a-c), defect periphery

(DP; d-f), cortical fragment center (FC; g-i), cortical fragment periphery (FP; j-l). a, d, g, j:

Bone formation rate (solid line) and bone resorption rate (dashed line) given in percent per

day. b, e, h, k: Bone volume (BV) normalized to TV (DC for DC and DP, FC for FC+FP). c, f, i, l:

Degree of bone mineralization given as ratio of bone volume with a density ≥645 mg HA/cm^3

to the total osseous volume (threshold ≥395 mg HA/cm^3). n=10 per group; * indicates $p <$

0.05 determined by two-way ANOVA with Geisser-Greenhouse correction and Bonferroni

correction.

Figure 4. Micro-CT and histological analyses of consecutive callus sections. a, b: Micro-CT

images of fracture callus sections corresponding to sections stained with Safranin-O for one

animal from the loading (a) and the control group (b): CT image (threshold: 395 mg HA/cm^3),

visualization of regions of bone formation and resorption via registration of CT image from

weeks 6 and 7, visualization of strains on CT section via micro-FE analysis for week 7 and

week 6. c-j. Consecutive longitudinal sections of fractured femora 7 weeks after defect

surgery of one animal from the loading group and the control group stained for Safranin-O

(c-f), Sclerostin (g, h), Rankl (i, j). Scale bar = 100 µm (c, d), scale bar = 500 µm (e-j).

Figure 5. Visualization of local cellular distribution patterns and protein expression in the

fracture callus and adjacent cortex. a. Longitudinal Safranin-O stained sections of fractured

femora 7 weeks after defect surgery of one animal from the loading group and one animal

from the control group. Boxes indicate regions depicted in b (1: callus; 2: cortex); Scale bar =

100 µm. b. Magnification of callus region (left) and cortical region (right) stained for

Safranin-O, Sclerostin and Rankl; Scale bar = 50 µm.

694 **Tables**

695 **Table 1.** Fracture healing outcome assessed by cortical bridging (healed fracture ≥ 3 bridged
696 cortices, threshold 395 mg HA/cm^3).

		week							
	Group (n=10)	0	1	2	3	4	5	6	7
Healed fractures per group	Control/Loading	$^0/_0$	$^0/_0$	$^5/_6$	$^9/_{10}$	$^{10}/_{10}$	$^{10}/_{10}$	$^{10}/_{10}$	$^{10}/_{10}$

Figure 1

RISystem, Davos, Switzerland

Figure 2

Group	week 0	week 0-1	week 1-2	week 2-3	week 3-4	week 4-5	week 5-6	week 6-7
Loading								
Control								

Figure 3

Figure 4

Loading

Control

a

CT week 7   CT week 6-7

b

CT week 7   CT week 6-7

Formation
Resorption
Quiescent

FE week 7   FE week 6

FE week 7   FE week 6

Saf-O   c   d

Saf-O   e   f

Sclerostin   g   h

Rankl   i   j

Figure 5