Evolutionary dynamics of transposable elements in bdelloid rotifers

Reuben W. Nowell[1,2,*], Christopher G. Wilson[1,2], Pedro Almeida[2,3], Philipp H. Schiffer[4], Diego Fontaneto[5], Lutz Becks[6,7], Fernando Rodriguez[8], Irina R. Arkhipova[8], and Timothy G. Barraclough[1,2,*]

[1]Department of Zoology, University of Oxford, 11a Mansfield Road, Oxford, United Kingdom. [2]Department of Life Sciences, Imperial College London, Silwood Park Campus, Ascot, Berkshire, United Kingdom.[3]Division of Biosciences, University College London, London, United Kingdom. [4]Institute of Zoology, Section Developmental Biology, Wormlab, University of Cologne, Köln, Germany. [5]National Research Council of Italy, Water Research Institute, Verbania Pallanza, Italy. [6]Community Dynamics Group, Department of Evolutionary Ecology, Max Planck Institute for Evolutionary Biology, Plön, Germany. [7]Aquatic Ecology and Evolution, University of Konstanz, Germany. [8]Josephine Bay Paul Center for Comparative Molecular Biology and Evolution, Marine Biological Laboratory, Woods Hole, MA

*Corresponding authors: RWN: reuben.nowell@zoo.ox.ac.uk; TGB: tim.barraclough@zoo.ox.ac.uk

Abstract

Transposable elements (TEs) are often considered as selfish genomic parasites whose ability to spread autonomously is facilitated by sexual reproduction in their hosts. In populations that no longer reproduce sexually, however, the long-term fate of TEs and their hosts is unclear. One possibility is that an asexual lineage may eventually rid itself of deleterious TEs, via increased selection for removal or domestication. Alternatively, TEs may proliferate unabated until the host is driven extinct. Here, we test these ideas using whole-genome sequencing of 26 single individuals from eight species of bdelloid rotifers, a class of invertebrate animals in which males are thus far unknown. We find that all species encode diverse class I retrotransposons and class II DNA transposons, with evidence for recent and ongoing activity. Bdelloid TE content does not appear unusual or deficient when compared to the variation observed in sexual rotifers or other animals. While theory predicts that vertically transmitted class I retrotransposable elements specifically should be scarce after millions of generations of presumed asexuality, we find them in all bdelloid genomes, at lower levels than in sexual relatives, but not so low as to be unprecedented among metazoans more broadly. We consider several possible explanations for these observations. We find no evidence for the hypotheses that cryptic recombination affects TE dynamics in bdelloids, or that TE activity is restrained by unusual adaptations for DNA repair in bdelloid species that tolerate extended desiccation. However, we find that bdelloid genomes share a large expansion and diversification of the Argonaute and RNA-dependent RNA polymerase (RdRP) gene families, which both function in RNAi-mediated TE suppression in other eukaryotes. Indeed, bdelloids may encode the largest expansion of RdRP genes among any eukaryote investigated so far. Thus, it is possible that the deleterious effects of active TEs are mitigated by enhanced cellular defence mechanisms in bdelloids, which might help to compensate for the consequences of long-term asexuality and lateral influx of new TEs.

Introduction

Transposable elements (TEs) are repeated sequences of DNA that can mobilize and replicate themselves within genomes [1–3]. TEs are divided into two major categories: class I retrotransposons, which use a 'copy-and-paste' replication mechanism via a reverse-transcribed RNA intermediate, and class II DNA transposons, which use 'cut-and-paste' replication with a

DNA intermediate. Both classes are ancient and diverse—retrotransposons are found in some bacteria and nearly all eukaryotes, while DNA transposons are found across the tree of life [4–6]. Although TE replications are occasionally beneficial [7], the vast majority are deleterious for the host [8,9]. Costs include insertional mutations that disrupt genes [10], cellular costs of replicating and expressing excess DNA [11], and increased risk of chromosomal abnormalities due to ectopic recombination between homologous TE sequences interspersed through the genome [12,13]. Despite this, by replicating autonomously as selfish elements, TEs can accumulate to large numbers within genomes—for example, TEs comprise 46% of the human genome, including over 1 million (~11%) nonautonomous *Alu* retroelements [14,15]. TE numbers vary greatly, however, even between closely related species. In vertebrates, for example, TEs span an order of magnitude, from below 6% to over 50% of the genome [16], with similarly large variation observed within and between other groups such as arthropods [17], nematodes [18], and fungi [19]. Explaining this variation is vital to understanding the mechanisms affecting TE spread and control.

Sexual reproduction has long been thought to play a major role in TE dynamics within eukaryotes. On the one hand, sexual reproduction and outcrossing decouples the fate of TEs from other host genes, allowing them to jump into new genomic backgrounds and to behave as selfish genomic parasites [1,2,20]. On the other hand, sex enables the efficient removal of deleterious insertions from populations through recombination and segregation [21–23]. The risk of chromosome abnormalities due to ectopic recombination, arguably the main cost of high TE loads in eukaryotes [9,24], also occurs during chromosome pairing at meiosis. Sex therefore plays opposing roles—it permits spread and selfish behaviour of TEs, and yet it facilitates and strengthens selection against high loads. Variation in TE content among taxa might thus result from shifts in the balance of these different opposing forces.

By this logic, the loss of sexual reproduction should affect TE dynamics dramatically. Since asexual lineages generally arise from sexual species [25], it is likely that they initially harbor many active TEs [26,27]. All else being equal, the loss of recombination will limit the ability of selection to remove deleterious insertions from a fully linked host genome, and so the load of TEs should accumulate. At the same time, the fate of TEs is immediately coupled to that of the host genome, resulting in intensified selection for inactivation, excision or domestication of the elements [2,22,27–29]. The genomes of asexual lineages whose TEs continued to replicate unchecked would become overrun, potentially leading to extinction of the lineage and the TEs themselves. While some TEs could be maintained by horizontal transfer between species (especially class II DNA elements [6,30]) or by having beneficial effects (as in bacteria, [31,32]), other TEs—particularly the class I LINE-like (i.e. non–long terminal repeat [LTR]) retrotransposons—are thought to be transmitted almost exclusively vertically [5,33,34], and therefore depend strongly on sex for their persistence.

Models of the population genetics of vertically transmitted TEs in asexuals predict one of two outcomes: either TEs accumulate within lineages faster than they can be removed, overrunning each lineage in turn and driving the population extinct, or, conversely, TE removal outweighs proliferation and the population eventually purges itself entirely of deleterious TEs [27,35,36]. These predictions are difficult to test empirically, however, because the time required for a population to arrive at either extinction or redemption is expected to be on the order of millions of generations [27], too long to observe directly and beyond the lifespan of most asexual lineages [37,38].

Here, we test these ideas in a well-known group of asexual animals, the bdelloid rotifers. These
microscopic invertebrates appear to have reproduced without males or meiosis for tens of
millions of years, diversifying into hundreds of species within limno-terrestrial and freshwater
habitats globally [39,40]. Bdelloids sampled from nature (and those reared in the laboratory)
consist entirely of parthenogenetic females, and neither males nor hermaphrodites are described
for any species despite centuries of close observation by naturalists [39,41,42]. Genetic and
genomic evidence for their proposed ancient and obligate asexuality remains uncertain, however.
Initial evidence of long-term asexuality [43,44] has been refuted by later studies or confounded
by alternative explanations [45–47]. Some recent studies have proposed alternative modes of
inter-individual genetic exchange, but these suggestions would require exotic mechanisms
unknown in other animals [44,48], or rates of sex that are difficult to reconcile with the lack of
observed males [49]. While the precise nature of reproduction in bdelloids remains an open
question, nonetheless they provide a unique test-case for models of TE evolution when
conventional sex is absent or strikingly rare.
115
Initial PCR-based surveys of five bdelloid genomes found no evidence of class I
retrotransposons from either the LTR or LINE-like superfamilies, but did reveal a diverse array
of class II DNA transposons, mostly at low copy number [50]. The presence of class II TEs in
bdelloids might be explained by horizontal transfer, which is thought to occur more frequently
for class II TEs with DNA intermediates [6,30,33,34] (but see [51]). The apparent lack of
retrotransposons contrasted sharply, however, with their near ubiquity in other taxa. At the time,
the absence of class I TEs appeared consistent with the view that long-term asexual evolution in
bdelloids had caused the loss of parasitic elements that depended on sexual transmission
[22,27,50,52].
125
Another unusual aspect of bdelloid physiology was suggested to contribute to their seemingly
low TE complement. In most bdelloid species (but not all), individuals can survive complete
desiccation at any life stage via a process called anhydrobiosis ('life without water'). Desiccation
causes double-strand breakages (DSBs) in DNA, but bdelloids are able to repair these and
recover to an unusual degree [53–55]. It was proposed that anhydrobiosis might influence TE
evolution in two ways [27,52,56]. First, DSB repair could aid TE removal, either via gene
conversion from a homologous chromosome lacking the TE insertion, or excision of mis-paired
regions. Second, the pairing of homologous chromosomes, if required during DSB-repair, could
provide a context for ongoing selection against chromosomal abnormalities caused by ectopic
recombination. In either case, anhydrobiosis would decrease the number of TEs, potentially
helping to explain the low overall TE content encoded in bdelloid genomes.
137
These early ideas were transformed by more detailed studies of the model bdelloid species
Adineta vaga, which used refined methods and genome-scale data to discover a variety of
retrotransposon families. These include an endonuclease-deficient *Penelope*-like element (PLE)
designated *Athena* [57,58], which is itself incorporated within much larger and highly unusual
retroelements called *Terminons* [59]; another PLE that has retained its endonuclease [60], LTR
retrotransposons (*Juno, Vesta, TelKA* and *Mag* [61,62]), and LINE-like retrotransposons (*R4, R9,*
Hebe, RTE, Tx1 and *Soliton* [44,63,64]). In total, TEs accounted for 2.2% of the 217 Mb genome
(~4.8 Mb) [44], rising to ~4% on inclusion of the recently discovered giant *Terminon* elements
[59]. The conclusion that bdelloids lack class I TEs therefore no longer holds, and the predicted
effects of asexuality and anhydrobiosis on TE evolution remain open questions that are amenable
to testing with comparative genomics. Specifically, the comparison of desiccating species with
those few bdelloid rotifer lineages that are unable to survive desiccation would indicate whether
anhydrobiosis does limit TE numbers as hypothesized. Also, comparisons within populations

could shed light on the activity of TEs and whether bdelloids possess cryptic forms of
recombination that could aid in TE removal or facilitate TE spread.
153
Here, we test the predicted effects of asexuality and anhydrobiosis on TE evolution by
comparing 42 rotifer genomes belonging to 15 taxonomic species. Our sample includes both
desiccating and nondesiccating bdelloids, and eight monogonont rotifers [65–67], a separate
class that alternates sexual and asexual reproduction and cannot survive desiccation as adults.
Results are set in context by comparison to published genomes from an acanthocephalan [68]
(now classified with rotifers in the Phylum Syndermata) and a range of other animal phyla. We
ask six questions raised by theory. (1) How does the abundance and diversity of TEs in bdelloids
differ from that in other animals, including sexual rotifers? (2) Are TEs recently and currently
active in all bdelloids? (3) Can we detect signatures of recombination affecting bdelloid TEs, as
might be expected if bdelloids do have cryptic forms of sexual reproduction? (4) Do desiccating
species contain fewer TEs than nondesiccating species, as previously theorized? (5) Are TEs in
bdelloids under the same selective constraints as in other taxa? (6) Finally, do bdelloids possess
alternative molecular mechanisms that might help to keep their TEs under control, as might be
expected if the ability to control TEs by sexual reproduction is absent or reduced?
168
169

Results and Discussion

171
High-quality population genomics data for bdelloid rotifers. To quantify variation in repeat
content within and between bdelloid species, we generated *de novo* whole-genome assemblies
for 31 rotifer samples encompassing nine species (**Fig 1a**; **Table 1**; S1 Data). Three of these
assemblies were generated using 10x Genomics linked-read data (for *Adineta steineri*, *Rotaria*
sordida, and *Rotaria* sp. 'Silwood-1'), while 26 are from single-individual samples. In order to
capture as many potential repeats as possible, we generated two assemblies for each sample: a
'reference' assembly, with a focus on quality and contiguity, and a 'maximum haplotype'
(maxhap) assembly that included small or highly similar contigs that might be derived from
recent TE duplications or other sources of copy number variation, at the expense of contiguity.
181
Reference genomes showed an expected scaffold size (AU, see Materials and methods) ranging
from 21.1 kb (*Didymodactylos carnosus*) to 702.3 kb (*R.* sp. 'Silwood-1') and BUSCO scores
that indicated 89–98% of 303 core eukaryote genes were completely recovered, increasing to 96–
99% if fragmented copies are included (**Table 1**). General genome characteristics such as
genome size (assembly span), the proportion of G + C nucleotides (GC%), the number of coding
genes (CDS), and the level of homologous divergence (number of SNPs identified within CDS)
were within the range expected from previous analyses of bdelloid genomes [44,46] (**Fig 1b–c**;
Table 1; S1 Fig). Intragenomic collinearity and synonymous divergence of coding regions in the
A. steineri, *R. sordida* and *R.* sp. 'Silwood-1' maximum haplotype assemblies reveal the
characteristic signature of degenerate tetraploidy that has been found in all bdelloid species
examined to date (**Fig 1d**).
193
Compared to the reference set, maxhap assemblies generally showed increased span (mean
increase = 17.9 Mb ± 21.5 standard deviation [SD]) and were substantially more fragmented, as
expected (S1 Table). Nonetheless, BUSCO completeness scores remained high, with 76% to
98% of genes completely recovered (increasing to 95–98% if fragmented copies are included),
indicating that the majority of core genes are successfully captured (S1 Table). The BUSCO
duplication metric ('D') does not increase greatly between reference and maxhap assemblies,
which shows that the additional sequences retained in the maxhap assemblies do not contain

complete extra copies of core genes. Thus, the maxhap assemblies are not fully haplotype-
resolved representations of the genome, except in the case of the three 10x assemblies.

Table 1. Assembly statistics for one monogonont and 30 bdelloid rotifer reference assemblies presented in this study.

Sample ID	Species name	SZ (Mb)	NN	N50 (kb)	L50	AU (kb)	GC (%)	Gaps (kb)	Coverage (X)	Genome BUSCO score	CDS	Proteome BUSCO score	GenBank accession
Bc_PSC1	*Brachionus calyciflorus* (Monogonont)	116.7	14,869	18.5	1,692	26.6	25.6	78	186	C:96%[S:93%,D:3%],F:2%	24,404	C:98%[S:93%,D:5%],F:1%	GCA_904009505.1
Ar_ARIC003	*Adineta ricciae*	135.6	4,302	283.8	129	388	35.5	65	89	C:97%[S:58%,D:39%],F:2%	49,015	C:97%[S:52%,D:45%],F:1%	GCA_904047305.1
As_10x_p	*Adineta steineri*	171.1	8,257	200.1	163	394.5	29	206	198	C:95%[S:67%,D:33%],F:2%	50,321	C:97%[S:58%,D:38%],F:2%	
As_ASTE804	*Adineta steineri*	160.3	9,359	158.1	265	214.9	29.1	152	62	C:95%[S:74%,D:22%],F:2%	47,222	C:98%[S:74%,D:24%],F:2%	GCA_904047245.1
As_ASTE805	*Adineta steineri*	156.3	9,008	169.6	245	226.4	29.2	129	65	C:98%[S:77%,D:21%],F:1%	43,986	C:99%[S:72%,D:26%],F:1%	GCA_904047255.1
As_ASTE806	*Adineta steineri*	160.3	7,597	168.2	257	222.5	29.2	145	82	C:96%[S:72%,D:24%],F:2%	45,930	C:98%[S:74%,D:24%],F:2%	GCA_904047275.1
Dc_DCAR505	*Didymodactylos carnosus*	323.6	87,048	7.8	11,656	10.5	33.5	41	21	C:86%[S:69%,D:17%],F:8%	46,286	C:88%[S:71%,D:18%],F:9%	GCA_904047325.1
Dc_DCAR706	*Didymodactylos carnosus*	368.8	78,356	12	7,695	19.1	33.5	13	76	C:95%[S:70%,D:25%],F:2%	46,863	C:95%[S:71%,D:25%],F:2%	GCA_904053135.1
Rd_10x_p	*Rotaria sordida*	272.5	16,571	64.5	843	193.8	30.8	395	91	C:94%[S:77%,D:19%],F:2%	44,299	C:95%[S:69%,D:26%],F:2%	
Rd_RSOR408	*Rotaria sordida*	252.9	20,315	57.6	1,246	75.5	30.4	291	39	C:94%[S:76%,D:19%],F:3%	40,501	C:97%[S:73%,D:24%],F:2%	GCA_904053145.1
Rd_RSOR410	*Rotaria sordida*	252.6	19,518	60.9	1,179	80	30.4	252	42	C:95%[S:77%,D:18%],F:3%	40,474	C:98%[S:74%,D:24%],F:2%	GCA_904053495.1
Rd_RSOR504	*Rotaria sordida*	251.3	22,067	53.1	1,338	69.8	30.4	369	39	C:94%[S:78%,D:16%],F:3%	41,085	C:96%[S:73%,D:23%],F:3%	GCA_904047315.1
Rg_MAG1	*Rotaria magnacalcarata*	178.7	19,184	42	1,077	62.4	32	402	58	C:97%[S:81%,D:16%],F:1%	40,318	C:99%[S:76%,D:22%],F:1%	GCA_903995865.1
Rg_MAG2	*Rotaria magnacalcarata*	181.1	22,216	39.7	1,141	61	32	433	63	C:98%[S:81%,D:17%],F:1%	40,289	C:99%[S:74%,D:26%],F:0%	GCA_903995855.1
Rg_MAG3	*Rotaria magnacalcarata*	180.9	22,132	40.7	1,142	60	32	508	60	C:96%[S:80%,D:17%],F:1%	40,740	C:99%[S:77%,D:22%],F:0%	GCA_903995845.1
Rg_RM15	*Rotaria magnacalcarata*	174	18,391	46.5	966	67.3	32	430	55	C:96%[S:80%,D:16%],F:2%	38,283	C:99%[S:77%,D:22%],F:1%	GCA_903995885.1
Rg_RM9	*Rotaria magnacalcarata*	173.8	19,520	44	999	64.7	31.9	594	51	C:96%[S:80%,D:16%],F:1%	38,404	C:98%[S:76%,D:22%],F:1%	GCA_903995825.1
Rp_RPSE411	*Rotaria* sp. 'Silwood-2'	296.5	30,050	102.5	381	691.1	31	247	35	C:93%,[S:72%,D:21%],F:4%	48,378	C:95%[S:72%,D:23%],F:4%	GCA_904053095.1
Rp_RPSE503	*Rotaria* sp. 'Silwood-2'	285.6	33,174	78.3	449	627.2	31.3	446	34	C:91%,[S:75%,D:17%],F:5%	48,269	C:92%[S:72%,D:20%],F:7%	GCA_904053125.1
Rp_RPSE809	*Rotaria* sp. 'Silwood-2'	271.1	28,589	101.6	350	681.4	31	377	27	C:93%,[S:76%,D:17%],F:4%	47,010	C:95%[S:74%,D:22%],F:4%	GCA_904053085.1
Rp_RPSE812	*Rotaria* sp. 'Silwood-2'	264.1	34,498	80.8	403	616.2	31.1	428	27	C:89%,[S:74%,D:15%],F:8%	47,040	C:90%[S:73%,D:17%],F:8%	GCA_904053105.1
Rs_AK11	*Rotaria socialis*	149.2	6,303	111.3	370	150.2	31.8	442	39	C:97%,[S:80%,D:17%],F:1%	34,844	C:99%[S:75%,D:24%],F:1%	GCA_903995835.1
Rs_AK15	*Rotaria socialis*	147.4	5,030	134.7	305	177.6	31.8	423	37	C:96%,[S:79%,D:18%],F:1%	34,140	C:98%[S:76%,D:23%],F:1%	GCA_903995795.1
Rs_AK16	*Rotaria socialis*	147.4	4,720	139.5	296	180.3	31.8	332	43	C:97%,[S:80%,D:18%],F:0%	33,717	C:99%[S:76%,D:23%],F:1%	GCA_903995805.1
Rs_AK27	*Rotaria socialis*	149.9	5,952	123.7	343	159.8	31.8	458	36	C:97%,[S:80%,D:17%],F:0%	34,369	C:99%[S:75%,D:24%],F:1%	GCA_903995815.1
Rs_RS1	*Rotaria socialis*	151.1	6,254	124.9	334	166.2	31.8	490	40	C:97%,[S:80%,D:17%],F:0%	33,937	C:99%[S:77%,D:22%],F:1%	GCA_903995875.1

		SZ	NN	N50	L50	Gaps	GC	AU		BUSCO (genome)		BUSCO (transcriptome/protein)	Accession
Rw_10x_p	*Rotaria* sp. 'Silwood-1'	310.4	16,995	211.8	211	126.2	31.1	534	53	C:95%,[S:76%,D:20%],F:1%	44,241	C:97%[S:73%,D:24%],F:1%	
Rw_RSIL801	*Rotaria* sp. 'Silwood-1'	268.4	28,548	136.5	288	687.3	30.8	472	45	C:94%,[S:77%,D:17%],F:4%	41,574	C:95%[S:75%,D:21%],F:5%	GCA_904053155.1
Rw_RSIL802	*Rotaria* sp. 'Silwood-1'	249.9	21,286	153.4	238	702.3	30.7	451	42	C:92%,[S:76%,D:16%],F:4%	39,577	C:94%[S:76%,D:18%],F:4%	GCA_904053505.1
Rw_RSIL804	*Rotaria* sp. 'Silwood-1'	247.6	25,643	118.3	287	660.4	30.8	667	34	C:94%,[S:78%,D:16%],F:3%	41,139	C:96%[S:78%,D:19%],F:3%	GCA_904053115.1
Rw_RSIL806	*Rotaria* sp. 'Silwood-1'	294.1	29,968	132.4	333	681.9	30.8	500	31	C:95%,[S:79%,D:16%],F:2%	48,259	C:97%[S:78%,D:19%],F:2%	GCA_904054495.1

Sequence statistics codes: **SZ**, total sequence length (Mb); **NN**, number of sequences; **N50**, N50 scaffold length (kb); **L50**, N50 index; **AU**, expected scaffold size (area under 'Nx' curve, kb).

Gaps: Undetermined bases (Ns) introduced during scaffolding.

BUSCO score based on eukaryote set ($n = 303$); BUSCO codes: **C**, complete; **S**, complete and single copy; **D**, complete and duplicated; **F**, fragmented.

Abbreviations: GC, guanine + cytosine; BUSCO, Benchmarking Universal Single-Copy Orthologs.

210
To these new data, we added published genomes for seven monogonont taxa (one from the
Brachionus calyciflorus species complex [65] and six from four species of the *Brachionus*
plicatilis species complex, namely *B. asplanchnoidis*, *B. plicatilis* sensu stricto (HYR1), *B.*
rotundiformis, and *B.* sp. 'Tiscar' [66,67]) and four bdelloids (*A. vaga*, *Adineta ricciae*, *Rotaria*
magnacalcarata and *Rotaria macrura* [44,46]), yielding a total of 42 rotifer genomes. Of these,
11 samples belong to nondesiccating bdelloid species (five individuals each from *R.*
magnacalcarata and *Rotaria socialis*, and the previously published genome of *R. macrura*).
218
219

a

b

c

d

220
221
Fig 1. Genome properties of sequenced rotifers. (a) Bdelloid rotifer morphology; scale bar indicates
100 μm. (*i*) Individual from an undescribed species of *Rotaria* (*R.* sp. 'Silwood-1'), showing eyes (E) and
foot (F) with two spurs and three toes. (*ii*) Further image of *R.* sp. 'Silwood-1' with anterior–posterior (A–P)
axis marked. (*iii*) Two individuals of *A. steineri* in phase contrast. **(b)** Cumulative assembly span for six
bdelloid species with population genomics data (*n* > 2). 10x Genomics haploid ('pseudohap') and diploid
('megabubbles') assemblies for *A. steineri*, *R. sordida* and *R.* sp. 'Silwood-1' are indicated with diamond
and circle symbols, respectively. The four previously published genomes for *A. vaga* ('Av2013', GenBank
accession GCA_000513175.1) and *A. ricciae* ('Ar2018', GCA_900240375.1), *R. macrura* ('Rc2018',
GCA_900239685.1) and *R. magnacalcarata* ('Rg2018', GCA_900239745.1) are indicated in grey, for
comparison. **(c)** Intragenomic divergence, measured as the number of SNPs detected in coding regions

(CDS). Boxplots show the median (band), interquartile range (box) and minimum/maximum values
(whiskers). Underlying data are shown as jittered points. Desiccation-tolerant species are in orange,
intolerant species in blue. Species abbreviations: **As**, *A. steineri*; **Rd**, *R. sordida*; **Rw**, *R.* sp. 'Silwood-1';
Rp, *Rotaria* sp. 'Silwood-2'; **Rg**, *R. magnacalcarata*; **Rs**, *R. socialis*. **(d)** Genome structure in *A. steineri*,
R. sordida and *R.* sp. 'Silwood-1' haplotype-resolved ('megabubbles') assemblies. Each point represents
a collinear block of genes, plotted by average pairwise synonymous (K_S, *X*-axis) and collinearity score
(see Materials and methods and S1 note) on the *Y*-axis. Separation into two distinct clusters representing
homologous (squares) and homoeologous (circles) relationships among gene copies is consistent with
ancestral tetraploidy, with homoeologous copies derived from a putative ancient genome duplication.
241
242
Abundant and diverse TEs in bdelloid genomes. To ascertain the repeat content of bdelloid
genomes relative to other taxa in a consistent manner, we used the RepeatModeler and
RepeatMasker pipelines to identify and classify repeats across genomes. The total proportion of
the genome classified as repetitive ranged from ~19% to 45% across bdelloid genera, with
variation within and between species (**Fig 2a**; S2 Fig; S3 Fig; S2 Data; S3 Data). Most of these
are simple or unclassified repeats that do not belong to major TE superfamilies. While the
precise nature of these unclassified repeats is not elucidated, an appreciable fraction (~7–27%,
mean = 17%) are also annotated as protein-coding and thus may be derived from gene
expansions or other duplications, while a further small fraction (< 1%) are accounted for by an
additional survey of small, nonautonomous class II TEs called miniature inverted-repeats
(MITEs) (S2 Data). The proportion of the genome accounted for by known TEs was much
smaller, ranging from 2.4% to 7.3% (mean = 4.9% ± 1.2 standard deviations [SD], median =
5.1%) in bdelloids. Broken down by class and superfamily, the mean values are: class I total =
2.09% ± 0.75 (PLEs = 0.59% ± 0.14; LTRs = 0.68% ± 0.26; and LINEs = 0.82% ± 0.47); class II
total = 2.79 ± 0.8 (DNA transposons = 2.49% ± 0.77; rolling circles = 0.30 ± 0.11). These results
are in broad agreement with previous estimates of TE content in bdelloids [44,46,47,59].
259
We first compare these values to closely related sexual lineages in the phylum Syndermata, the
monogonont rotifers and acanthocephalans. The most striking difference relates to the recently
published genome of the obligately sexual acanthocephalan *Pomphorhynchus laevis* [68], which
encodes a substantially greater proportion of repeats than either bdelloids or monogononts. In
agreement with Mauer et al., we find ~66% of the *P. laevis* genome to be composed of repeats.
The large majority are class I retrotransposons (~71% of the total TE content) from the LINE
(~52%) and PLE (~15%) superfamilies, and there are relatively few DNA transposons (~1.3%)
(**Fig 2b**). There is increasing evidence that acanthocephalans may be the closest relatives to the
Class Bdelloidea [69–71]. However, all members of the Acanthocephala (also known as the
thorny-headed worms) are obligate endoparasites and are highly differentiated in both
morphological and molecular terms from other syndermatans. It is unclear whether these features
may contribute to the high repeat content and other unusual genome characteristics of *P. laevis*
[68].
273
In monogononts, the mean values for class I TEs is 5.2% ± 1.5 SD and for class II TEs it is 2.5%
± 1.0 SD. Thus, monogononts are slightly more TE-rich than bdelloids, but also substantially
more variable between species (**Fig 2a**; S3 Fig). Repeat content differs between bdelloids and
monogononts in two main ways, both in regard to the composition of class I retrotransposons.
First, monogononts encode substantially more LINE-like retroelements than bdelloids, making
up (on average) approximately 50% and 16% of the total TE content in each clade respectively.
The frequency of LINEs is of particular interest, because this class of TEs is thought to be least
likely to undergo horizontal transfer and thus the most dependent on sex for transmission
[5,33,34]. Second, *Penelope*-like elements (PLEs) have increased in proportion in all bdelloids
relative to monogononts, from ~1% in monogononts to ~12% in bdelloids on average.

Interestingly, a high proportion of PLEs is also seen in the acanthocephalan *P. laevis*, while a
high proportion of LINEs is found in both *D. carnosus* isolates (~35% of total TE content), a
deeply branching lineage sister to all other bdelloid taxa included in the analyses. Thus,
assuming that acanthocephalans are the closest relatives to bdelloids, the most parsimonious
explanation for these broad-scale patterns is that the expansion of PLEs occurred in the ancestor
to bdelloids and acanthocephalans, whereas the contraction of LINEs has occurred more
recently, confined to a subset of bdelloid genera.
291
To put these differences in TE content among rotifers in a broader context, we applied the same
repeat-finding pipeline to animals from a range of more distantly related protostome phyla: three
insects, a nematode, two tardigrades, five molluscs, two annelids, a brachiopod, platyhelminth
and orthonectid [72–83] (S2 Table). As expected, both the abundance and diversity of TEs varied
widely across taxa (**Fig 2b**). Total TE content ranged from 0.8% (the insect *Polypedilum*
vanderplanki) to ~24% (octopus and platyhelminth). Although the acanthocephalan genome
appears to be particularly rich in TEs, bdelloids (except for *D. carnosus*) have modest amounts
of TEs, including class I TEs specifically, while in monogononts these amounts vary greatly
among species. All bdelloids encode relatively more TEs than both *Polypedilum* species but
fewer than *Drosophila melanogaster*, *Caenorhabditis elegans*, annelid worms and some
molluscs, and are intermediate with respect to other taxa. Note that TE proportion in molluscs,
nematodes and flatworms is known to be highly variable (e.g. [18]), while bdelloids display
much less variability barring the early-branching *D. carnosus*.
305
306

a

307
308
309 **Fig 2. Repeat content and diversity in rotifer genomes. (a)** Maximum likelihood phylogeny of eight
310 monogonont (square symbols on tips) and 34 bdelloid (circles) genomes based on the concatenated
311 alignment of a subset of core eukaryotic (BUSCO) genes. Orange and blue tip colours indicate
312 desiccating and nondesiccating taxa, respectively. Species codes in tip names are: **Bc**, *Brachionus*

calyciflorus; **Br**, *B. rotundiformis*; **Bt**, *B.* sp. 'Tiscar'; **Bp**, *B. plicatilis* HYR1; **Ba**, *B. asplanchnoidis*; **Dc**, *Didymodactylos carnosus*; **Av**, *Adineta vaga*; **Ar**, *A. ricciae*; **As**, *A. steineri*; **Rs**, *Rotaria socialis*; **Rg**, *R. magnacalcarata*; **Rc**, *R. macrura*; **Rd**, *R. sordida*; **Rw**, *R.* sp. 'Silwood-1'; **Rp**, *R.* sp. 'Silwood-2'. Repeat content is shown as the genome proportion (%) broken down by TE superfamily (middle panel), and relative proportion (%) of total known (i.e., classified) TEs (right panel), where colours represent TE superfamilies (see legend) and shades of colour represent different TE families within each superfamily. **(b)** Equivalent repeat content analysis in 17 protostome animal genomes, including the model species *D. melanogaster* and *C. elegans*, the recently published acanthocephalan rotifer *P. laevis*, and selected other species from across the protostome group. Two further examples of desiccating (orange 'D') and nondesiccating (blue 'ND') species pairs are shown: the insects *P. nubifer* and *P. vanderplanki* and the tardigrades *H. exemplaris* and *R. varieornatus*.

These results show that bdelloid species encode an abundant diversity of both class I and II TEs, and, set against the repeat content of other rotifers and animals, do not appear particularly deficient or unusual with regard to the proportion of nucleotides encoding transposable elements. Although the bdelloids do have lower frequencies of class I TEs (~16% of total TE content) than the monogononts (~50%) or the acanthocephalan (~52%), as predicted by theory for elements dependent on vertical transmission, they still possess them in numbers that are comparable to sexual organisms (e.g. *C. elegans*). Furthermore, the basally divergent bdelloid *D. carnosus* did not show the same magnitude of decrease in LINEs (~35%), indicating that different dynamics may be at play in different lineages. Thus, it is clear that the most simplistic expectations of TE evolution under the hypothesis of long-term asexuality (i.e., either runaway proliferation or complete elimination) are not met, necessitating an evaluation of possible explanations that may align theory with observation.

TE transposition is recent and ongoing. One possible explanation is that TEs in bdelloid genomes do not replicate autonomously or are otherwise inactivated or 'fossilised' within their host genome. To investigate this, we first generated divergence 'landscapes' for identified TE copies within each genome, using the de novo RepeatMasker results. TE landscapes measure the amount of sequence divergence between each TE copy and a consensus derived from all copies in its family [84]. Histograms of the resulting Kimura distances (K-values [85]) provide insights into the evolutionary history of TE activity [16,86,87].

TE landscapes for the three diploid (10x Genomics) assemblies of *A. steineri*, *R.* sp. 'Silwood-1' and *R. sordida* show that TE divergence is bimodal but strongly zero-inflated (**Fig 3**). A large number of TE copies have very low or no divergence from the consensus (K-value ≤ 1%). Assuming a molecular clock for nucleotide substitutions within duplicated TEs, such elements represent recent duplications that are highly similar to their progenitor copy, consistent with recent transposition of an active element. In proportion, most of these belong to class II DNA transposon superfamilies (in red), but the spike of zero divergence is also present for class I retrotransposons (in blue and green). An older, broader mode is seen around a K-value of 20–30% that probably reflects historical TE transpositions and/or a signal from the tetraploid genome structure present in all bdelloids sequenced to date. The same pattern was observed in the haplotype-resolved assemblies of *A. vaga* [44] and *A. ricciae* [46] (**Fig 3b**; S4 Fig), and was generally present but less pronounced in the other 'maxhap' assemblies depending on the repeat pipeline applied (S4 Fig; S5 Fig).

Fig 3. TE divergence landscapes for selected genomes. The *X*-axes show the level of divergence (Kimura substitution level, CpG adjusted) between each identified TE copy and the consensus sequence for that TE family (the inferred ancestral copy). Thus, if newly arising TE copies evolve neutrally, the amount of divergence is a proxy for the time since its duplication, with older copies accumulating more substitutions and appearing further to the right. The *Y*-axis shows the proportion of the genome occupied by each bin. Colours represent TE superfamilies (see legend) and shades of colour represent different TE families within each superfamily. Data are shown for the 10x Genomics diploid ('megabubbles') assemblies of *A. steineri*, *R.* sp. 'Silwood-1' and *R. sordida* compared to the published genome of *A. vaga*. Note different scales on some *Y*-axes.

To evaluate recent TE activity further, we developed a simple method to identify insertion sites for LTR retrotransposons (LTR-Rs) and assess their presence or absence in related individuals. Because most LTR-Rs insert into random genome locations [5,8], the neighbouring genome sequence provides a unique marker for a given insertion event [88]. We constructed a library of such insertion markers ('LTR-tags') for all 'full-length' LTR-Rs (i.e. those with long-terminal repeats present at both the 5' and 3' ends of the element) detected in our genomes, and then searched for their presence or absence in the other samples. For a given LTR-tag identified in genome *A*, the presence of a contiguous alignment in genome *B* indicates that the same insertion is shared between *A* and *B*.

For a set of 161 high-confidence and non-redundant LTR-Rs identified in the single-individual samples, alignment contiguity for each LTR-tag versus each of the other genomes was scored using a read-mapping approach (see Materials and methods), resulting in a pairwise matrix of presence/absence scores (**Fig 4a**; S4 Data). High scores for LTR insertion-site presence

389 correlated strongly with the phylogeny, resulting in an average score of ~0.9 within species
390 compared to < 0.1 between species and a clear visual signal along the diagonal of Fig 4a. Very
391 few LTR insertion sites were shared between bdelloid species. While some absences could
392 reflect loss rather than gain, the restriction of nearly all LTR insertion sites to single species
393 indicates that they have been gained during the separate evolutionary history of that species.
394
LTR-R insertions also vary between individuals within the same species, indicating recent
transposition events and the potential for ongoing fitness consequences for the host. One case-
study is illustrated for *R. magnacalcarata* (**Fig 4b**; S6 Fig). The individuals RM9 and RM15
share an LTR-R insertion that is not present in conspecifics. Aligning the regions of the genome
assemblies containing these LTR-tags indicates that an 8.1 kb LTR-R has inserted into a protein-
coding sequence in the lineage leading to RM9 and RM15. It has introduced a premature stop
codon to a gene that encodes a protein (7,479 residues) of unknown function but with partial
similarity to midasin, an ATPase essential to ribosome biosynthesis in several model eukaryotes
[89,90]. In RM9 and RM15, the predicted product is substantially truncated (to 6,025 residues)
by the element insertion. Despite the potential fitness consequences, RM9 and RM15 have
evidently persisted for some time since, because they differ at approximately 0.5% of single-
nucleotide sites across the 8.1 kb LTR element itself. A possible explanation is that both the
RM9 and RM15 assemblies also contain a scaffold with an empty insertion site, indicating an
intact version of the coding sequence spanned by the LTR insertion (represented in Fig. 4b by
the partial matches on scaffolds 16719 and 07127, respectively). If the insertion is hemizygous,
an uninterrupted homologous copy of the affected gene might mask or reduce the effect of the
mutation. Given the degenerate tetraploid structure of bdelloid genomes, further homoeologous
copies (i.e. derived from whole-genome duplication) might provide additional functional
redundancy and help perform critical functions of interrupted genes.
414
415

416
417
Fig 4. LTR insertion-site polymorphism in bdelloid species. (a) Columns represent 161 LTR-Rs
identified across bdelloid samples, arranged by genome of origin (see colours at bottom and side).
Support for the presence of a given LTR-R at a specific insertion site in each genome is scored from 0
(absent, yellow) to 1 (present, dark blue), where a score < 0.5 is strong evidence for absence (see
Materials and methods for details). Arrows demark the location of the LTR-R example shown in b. **(b)**
Nucleotide alignment of region around an LTR-R insertion (blue) identified in RM9 (scaffold 05348) and
RM15 (scaffold 07890), alongside their putative homologous scaffolds (scaffolds 16719 and 07127
respectively) that do not show the insertion. Scaffolds from Rg2018 (RMAG00928) and MAG1 are also
shown for comparison. Predicted CDS with similarity to Pol and Env proteins are shown in red and light
blue. The LTR-R is most likely a member of the *TelKA* family, based on sequence similarity.

428
429
430 Thus, these data contradict the idea that bdelloid TEs are inactive. All TE superfamilies in other
431 bdelloids show a substantial fraction of copies at low-divergence, indicative of recent
432 proliferation. Moreover, there are multiple cases of insertion-site polymorphism within species,
433 and at least one case where a recent retroelement insertion into a protein-coding sequence seems
434 likely to have potential fitness consequences.
435
No evidence that cryptic recombination helps to limit the spread of LTR-Rs. Another
possible explanation for the apparent discrepancy between bdelloid TE profiles and theory is that
bdelloids in fact possess cryptic inter-individual recombination, either through undetected sex or
some alternative form of gene transfer. We therefore tested for a signature of recombination
among polymorphic LTR-R insertion sites within species. Under strict clonality, the pattern of
presence and absence across LTR-R loci should be nested and compatible with only mutational
gain and loss at each site. In contrast, in a sexual, outcrossing population, variation should be
shuffled among loci. LTR-Rs provide a powerful test of these predictions because random
insertion makes independent origins of the same LTR-R insertion site highly unlikely.
445
In every species with multiple samples, we found that variation in polymorphic TEs is perfectly
nested, with a consistency index in parsimony reconstruction of 1. Furthermore, in the two
species with multiple parsimony-informative characters, *R. socialis* and *R. magnacalcarata*, we
found a significantly positive index of association of presence and absences among LTR-R
insertion sites, as expected with clonal inheritance (S3 Table; S6 Fig). Approximate Bayesian
Computation with simulations of expected patterns under varying frequencies of sexual
reproduction showed that strictly clonal evolution could not be rejected (S7 Fig). While this test
uses a restricted set of markers, and so should not be viewed as a test of recombination for the
whole genome or species, it does support clonal inheritance of LTR-R loci and finds no evidence
that inter-individual recombination helps to limit their spread. Nevertheless, local LTR-LTR
recombination within genomes, leading to solo LTR formation, may act to bring the copy
number down [44].
458
No evidence for lower TE loads in desiccating bdelloids. The desiccation hypothesis posits
that TE numbers may be kept in check via the action of DSB-repair processes during recovery
from desiccation. Our study includes 11 nondesiccating bdelloid samples encompassing three
obligately aquatic species (*R. macrura*, *R. magnacalcarata* and *R. socialis*), while the remaining
samples were isolated from ephemeral ponds or moss and must undergo frequent cycles of
desiccation and rehydration to survive. Contrary to the prediction that TE load should be reduced
in desiccating species, there is little overall difference in TE proportions between desiccating and
nondesiccating lineages (mean = 4.8% ± 1.3% *vs.* 5.0% ± 0.9% respectively). Broken down by
TE superfamily, desiccating taxa have relatively more DNA transposons, simple, low complexity
and unclassified repeats, and relatively fewer PLE, LTR and LINE-like retroelements (S3 Fig;
S8a Fig), with the biggest differences seen between *Rotaria* lineages (S8b Fig). However,
perhaps unsurprisingly given only two independent shifts in desiccation ability within our sample
(see phylogeny in Fig 2a), results from a Bayesian mixed-effects modelling approach that
controlled for phylogenetic relationships showed no significant correlations between desiccation
ability and TE load, for either overall proportion or for any individual TE superfamily (S4
Table). For most TE superfamilies the strength of the phylogenetic signal (λ) was close to 1 (S9
Fig), consistent with a high fit of the data to the phylogeny under a Brownian motion model as
would be expected if TE load evolves neutrally along branches of the phylogeny.
477

Two further comparisons of desiccating versus nondesiccating species among our wider sample of animals also present contrasting results. In chironomid midges, the desiccation-tolerant *P. vanderplanki* encodes substantially fewer TEs than its nondesiccating sister species *P. nubifer*, as predicted (0.8% and 2.2% respectively, although this rises to ~11% in both species when all repeats are included). In tardigrades, however, the desiccation tolerant *Ramazzottius varieornatus* encodes a greater proportion of TEs than *Hypsibius exemplaris* (4.3% and 2.8%, respectively), which does not survive desiccation without extensive conditioning [81], although the trend is reversed when all repeats are included due to a large fraction of simple repeats in *H. exemplaris*. We therefore find no consistent evidence for the hypothesised link between anhydrobiosis and TE load in bdelloids or beyond. The overall effect of desiccation on TEs might be dual: while repair of a DSB within a TE via non-homologous end-joining would likely result in its inactivation (thus acting to reduce TE load), an efficient DSB repair system would enhance repair of DSBs that arise during transposition of cut-and-paste DNA TEs that leave a DSB behind upon excision (thus allowing an increased TE load).

Bdelloids experience similar selective constraints on TEs as do other species. A fourth hypothesis is that the selective environment for TEs is different in bdelloids than in other animals, thereby shifting their TE profiles compared to simple theory. For instance, bdelloids might tolerate insertions within genes unusually well, owing to redundancy arising from tetraploidy or multiple gene copies [45,91,92]. First, we explored the genomic 'environment' of TE insertions and their potential effects on genome function. Differences in the location of TE insertions might reveal differential costs and benefits compared to other taxa. To do this, we first compiled a high-confidence list of class I retrotransposons by searching for proteins with significant similarity to the reverse transcriptase (RT) domain found in all retrotransposons. Phylogenies of the resulting alignments showed a diverse array of RTs in all species, most of them full-length (in terms of conserved subdomain presence) and clustered within the three primary retrotransposon superfamilies of PLEs, LTRs and LINEs (**Fig 5a**; S10 Fig; S5 Data). Many (but not all) clustered within families previously identified in *A. vaga*. The elevated LINE content in *D. carnosus* in comparison to other bdelloids is mostly due to the expanded Soliton clade and to the presence of CR1-Zenon and Tad/I/Outcast clades, the latter being characterized by the presence of the RNase H domain.

We then characterised surrounding genome features for these TEs. In 50 kb windows surrounding each class I TE identified above, we counted the occurrence and span of three features of interest: other (non-TE) genes, other (non-focal) TEs, and the telomeric repeat "TGTGGG" (identified from *A. vaga* [58] and supported in other rotifers, see S2 Note). Relative to a set of core metazoan (BUSCO) genes, the regions surrounding PLE, LINE and LTR TEs all showed significant decreases in gene density, but significant increases in both TE and sub-telomeric repeat density (**Fig 5b–d**; S5 Table). In contrast, using a linear mixed-effects modelling approach (see Materials and methods), there were no significant differences between the monogonont *B. calyciflorus* and bdelloids, or between desiccating and nondesiccating bdelloid species (S5 Table). These results are consistent with previous findings that TEs are mainly confined to sub-telomeric regions of bdelloid genomes [56], a bias that is presumably due to either selection against insertions at or near functioning genes and/or strong specificity in TE insertion site. Thus, it appears that most TE insertions are costly in bdelloid rotifers, as in other taxa, and that selection leads to their concentration outside of gene-rich regions.

526
527
Fig 5. Phylogenetic diversity and genomic context of reverse transcriptase genes. (a) For each
phylogeny, coloured branches represent identified rotifer-encoded RT copies and grey branches
represent the core RT sequences from which the HMM was built (see Materials and methods and S9 Fig
for core RT tree details). Colours indicated the major superfamilies. Previously characterised
retrotransposons are indicated on the *A. vaga* tree (**He**, *Hebe*; **So**, *Soliton*; **RTE**, *RTE*; **Tx1**, *Tx1*; **R9/4**, *R9*
and *R4*; **Ve**, *Vesta*; **Ju**, *Juno*; **Te**, *TelKA*; **Ma**, *Mag*; **At**, *Athena*; **Pe**, *Penelope*). All phylogenies are rooted
on the branch separating the bacterial retrons. Scale bar represents 0.6 amino acid substitutions per site.
Desiccating and nondesiccating species are indicated with orange and blue, as previously. Species
codes: **Bc**, *B. calyciflorus* PSC1; **Dc**, *D. carnosus* DCAR706, **Av**, *A. vaga* Av2013; **Ar**, *A. ricciae*
ARIC003; **As**, *A. steineri* ASTE805; **Rg**, *R. magnacalcarata* MAG3; **Rs**, *R. socialis* AK11; **Rc**, *R. macrura*
Rc2018; **Rd**, *R. sordida* RSOR408; **Rw**, *R.* sp. 'Silwood-1' RSIL806; **Rp**, *R.* sp. 'Silwood-2' RPSE503.
The genomic context in which RT genes reside is then described based on proximity to three other
features: **(b)** 'other' genes (that do not overlap with any TE annotation), **(c)** other TEs, and **(d)** telomeric
repeats ("TGTGGG"; that do not overlap with any coding region) as identified in *A. vaga*. For each plot, a
25 kb window is drawn around the focal RT gene and the total span (kb) of each feature within the
window is counted, broken down per sample ID (coloured boxes) per TE superfamily (*X*-axis groups).
Boxplots show the median (band), interquartile range (box) and minimum/maximum values (whiskers;
outliers not plotted). The equivalent data for BUSCO genes (metazoan set) are also shown for
comparison. The same representative samples are used in (b–d) as for (a).
547
548

As a second source of selective constraints, we tested for evidence of selection against ectopic
recombination (ER). ER is argued to be a major cost of TEs in sexual taxa, but its effects derive
from chromosomal abnormalities during meiosis, which should be lacking in bdelloids. Because
the rate of ER increases with both the number of elements and their length [12], the strength of
selection is expected to be strongest against longer TEs at higher copy number [9,24,93]. Two
testable predictions arise: first, that bdelloids should have longer TEs than sexual taxa (under the
hypothesis that ER is absent in bdelloids because of a lack of meiosis), and second, that
nondesiccating bdelloids should have longer TEs than desiccating bdelloids (under the
hypothesis that ER may still occur when chromosomes pair during the repair of DSBs).
However, comparisons of TE length distributions provide no evidence for these predictions—
there was no significant difference between monogononts and bdelloids, or between desiccating
and nondesiccating bdelloids (**Fig 6a**; S6 Table; S6 Data). Thus, while the precise estimation of
TE lengths will no doubt improve with increasing assembly contiguity, our current data do not
show any indication of changes in TE length linked to asexuality (when compared to
monogononts) or desiccation ability within bdelloids. It is interesting to note that this finding
also applies to the PLEs, which include the *Athena* elements that comprise the unusually large
and complex *Terminon* elements found at bdelloid telomeres [59,94]. In this case, it is possible
that selection may still act on individual *Athena* elements but does not affect *Terminons* per se
because their structural diversity and genomic location does not make them targets of ER.
568
A final prediction of selection against ER is that there should be a negative correlation between
TE frequency and length, as is observed in *Drosophila* [24] and humans [93]. For both
monogononts and bdelloids, the majority of identified TEs are short (< 1 kb), which are
presumably partial matches or degraded copies. Nonetheless, we observe a sharp decline in copy
number as mean TE length increases above ~0.5 kb, and a distinct lack of longer elements at
higher copy numbers (**Fig 6b**). In vertebrates, previous work has suggested a lower threshold of
~0.6–1 kb under which ectopic recombination does not operate [93,95]. Thus, the observed
patterns in rotifers are consistent with the hypothesis that longer elements above a certain length
threshold are selected against more strongly due to the deleterious effects of ectopic
recombination. This finding is contrary to a similar analysis performed on TEs in nematodes,
which did not recover the expected relationship [18]. However, the pattern is the same in both
desiccating and nondesiccating bdelloid representatives as well as the monogonont *B.*
calyciflorus, and to some extent the acanthocephalan *P. laevis* (S11 Fig), suggesting that
selection against longer TEs at higher copy number is a general feature in Syndermata,
regardless of desiccation ability.
584
585

Fig 6. TE length dynamics. (a) Distribution of TE length for selected syndermatan samples decomposed into the major TE superfamilies (DNA transposons, LINE-like, LTR and PLE retrotransposons). Boxplots show the median (band), interquartile range (box) and minimum/maximum values (whiskers; outliers are shown in grey). Species codes: **Bc**, *B. calyciflorus* PSC1 (monogonont); **Dc**, *D. carnosus* DCAR706; **As**, *A. steineri* ASTE804; **Av**, *A. vaga* Av2013; **Ar**, *A. ricciae* Ar2018; **Rd**, *R. sordida* RSOR408; **Rp**, *R.* sp. 'Silwood-2' RPSE411; **Rw**, *R.* sp. 'Silwood-1' RSIL801 (desiccating bdelloids); **Rc**, *R. macrura* Rc2018; **Rg**, *R. magnacalcarata* MAG1; **Rs**, *R. socialis* AK11 (nondesiccating bdelloids). An equivalent plot including the acanthocephalan *P. laevis* is shown in S10 Fig. **(b)** Relationship between mean TE length per TE family (*X*-axis) and copy number (i.e., the number of TEs identified within each family; *Y*-axis). The same set of individuals are shown as for (a). A dashed line is drawn at 0.6 kb, given as the length threshold under which the rate of homologous ectopic recombination is negligible in mice.

Large expansion of TE silencing pathways in bdelloids. Having rejected a range of hypotheses to reconcile theory and observation of TE levels in bdelloid rotifers, we finally looked for expansions and/or diversifications in the molecular pathways that defend against TEs. We characterised copy number variation for three well-known gene families with direct roles in TE suppression via RNA interference (RNAi). (1) Argonaute proteins of both the Ago and Piwi subfamilies, the core effectors of RNAi gene-silencing that form complexes with various classes of small RNA [96,97]; (2) Dicer, an RNase III–family protein that cleaves double-stranded RNA (dsRNA) molecules from 'target' genes into shorter fragments that are subsequently incorporated into Argonaute complexes [98,99]; and (3) RNA-dependent RNA polymerase (RdRP), an RNA replicase that synthesises secondary small interfering RNAs (siRNAs) that amplify the silencing response [98,100].

Based on hidden Markov model (HMM) matches of key domains to the predicted proteomes of the Illumina maxhap assemblies (in which homologous copies are largely collapsed), we detect an average of 21.5 putative Argonaute copies, 3.9 Dicer copies and 37.3 RdRP copies in bdelloid genomes (**Fig 7a**; S4 Data). These expansions are substantially larger than previously uncovered from the diploid assembly of *A. vaga* (eight Dicers, 23 Ago/Piwi, 20 RdRP per diploid genome) [44], particularly after correcting for the assembly resolution (i.e. diploid vs haploid assembly; see Materials and methods), perhaps due to the increased sensitivity of the HMM-based approach, or to different degrees of pseudogenization and/or homolog collapse. Phylogenies of identified copies revealed a number of divergent clades, particularly in the Argonaute and RdRP families (**Fig 7b–d**; S12 Fig; S7 Data; S8 Data), that might indicate both expansion and

diversification of these proteins. Even accounting for degenerate tetraploidy in bdelloids, the
RdRP domain in particular is significantly expanded relative to monogononts and other
eukaryotes (**Fig 7e–g**; S9 Data). This expansion is not found in the monogononts *B. calyciflorus*
or *B. plicatilis* HYR1, nor is there evidence for it in the (unannotated) acanthocephalan genome
(S4 Data), suggesting that the expansions seen in Ago and (particularly) RdRP are unique to
bdelloids, including the basally divergent *D. carnosus*.
629
The expansion of bdelloid-specific Ago genes (denoted 'BDAGOs' in **Fig 7**) is comparable to
the 'worm-specific' Ago clade (WAGOs) found in nematodes [101–103]. Intriguingly, the
majority of nematode species (although not *C. elegans*) have apparently lost their Piwi-like
orthologue, the branch of the Argonaute family that is usually involved in TE suppression in
most other animals [104], but instead mediate TE silencing using a Dicer/RdRP pathway [105].
The primary function of RdRP is to amplify RNAi responses via the production of small
interfering RNAs (siRNAs) [98], but this family is found only rarely in animal genomes [100]. In
C. elegans, RdRP-generated siRNAs (known as '22G-RNAs') also play important roles in the
recognition of 'self' versus 'non-self' RNA and multigenerational (i.e., inherited) epigenetic
memory [103,106]. However, an analysis of TE variation across the nematode phylum found no
effect of RNAi pathway differences (in terms of presence/absence of RNAi genes) among
species, concluding that TE content is mediated primarily by the stochastic action of genetic drift
[18].
643
Why do bdelloids possess such a marked expansion of gene silencing machinery? One
explanation may be that it was required as part of the successful long-term transition to
asexuality, if other mechanisms usually acting in sexual populations were no longer operating.
Intriguingly, it has been shown in *A. vaga* that piwi-interacting small RNAs (piRNAs) target
both TEs and putatively 'foreign' genes (i.e. non-metazoan genes gained via horizontal gene
transfer [HGT]) [107], the latter of which are unusually frequent in bdelloid genomes
[44,46,108–110]. Thus, it may be the case that bdelloids have an enhanced RNAi system to
defend against invasion from horizontally transferred TEs, particularly if the level of exposure or
rate of import is higher relative to other animals. Furthermore, the ability of some RNAi
pathways to distinguish self from non-self may be needed to maintain genome integrity over
longer timescales, if such mechanisms are operating. Further work on the precise functions of the
divergent Ago and RdRP clades is required to explore these possibilities.
656
657

Fig 7. Expansion of TE silencing pathways in bdelloid rotifers. (a) Copy number variation for RNAi gene families Argonaute (Ago/Piwi, red), RNA-dependent RNA polymerase (RdRP, blue) and Dicer (green) in bdelloids compared to other protostome groups. Proteins are identified based on the presence of key identifying domains (see Materials and methods). Species codes for rotifers: **Bc**, *B. calyciflorus*; **Bp**, *B. plicatilis* HYR1; **Dc**, *D. carnosus*; **As**, *A. steineri*; **Rg**, *R. magnacalcarata*; **Rs**, *R. socialis*; **Rc**, *R. macrura*; **Rd**, *R. sordida*; **Rw**, *R.* sp. 'Silwood-1'; **Rp**, *R.* sp. 'Silwood-2'. Maximum likelihood unrooted phylogenies are then shown for **(b)** Argonaute, **(c)** RdRP and **(d)** Dicer gene copies identified in *A. steineri*, *R. sordida* and *R.* sp. 'Silwood-1' 10x haploid ('pseudohap') assemblies, aligned with orthologs from representative species from across the eukaryotes. Blue symbols indicate copies identified in the monogonont *B. plicatilis*, and letters on tips show selected reference species to aid visual orientation: 'C', *C. elegans*; 'H', human; 'D', *D. melanogaster*; 'N' *N. crassa*; 'A', *A. thaliana*. Some clade names are also shown where relevant; 'WAGO' indicates the worm-specific cluster of Ago genes in the Argonaute phylogeny. 'BDAGO' I and II and 'RDR' I and II indicate putative bdelloid-specific clades of Argonaute and RdRP proteins, respectively. **(e)** Comparative protein-domain abundance plot. Each point represents a Pfam domain ID, with (log) average abundance (i.e., count) in the reference eukaryote set shown on the *X*-axis and (log) abundance in the monogonont *B. plicatilis* on the *Y*-axis (see Materials and methods). The positions of the PAZ, PIWI (red up/down triangles), RdRP (blue square) and Dicer (green circle) domains are highlighted. Dashed line indicates the 1-1 relationship. **(f)** Equivalent plot for bdelloids,

where the *Y*-axis shows the (log) average abundance for the *A. steineri*, *R. sordida* and *R.* sp. 'Silwood-1'
10x haploid assemblies. Note that the average abundance for all Pfam entries is shifted above the 1-1
line due to the ancient genome duplication in all bdelloids, such that many genes are found in double
copy (i.e., homoeologs) even in a 'haploid' representation. **(g)** Comparative protein–domain abundance
plot for bdelloids versus eukaryotes (see Materials and methods). Entries to the right of the distribution
are overrepresented in bdelloids with respect to eukaryotes. The shaded area represents the 5% and
95% quantiles of the distribution, and the scores for the PAZ, PIWI, Dicer and RdRP domains are
indicated (see legend).

Conclusions

We show that all bdelloids encode a rich diversity of TEs from both class I (retroelements) and
class II (DNA transposons), and thus reject the idea that bdelloids are deficient or unusual in
their TE content or diversity. Moreover, a substantial fraction of these elements has been active
relatively recently within populations. This finding is at odds with the original predictions of
population genetic theory for TEs in asexuals. One possible resolution is that theory is missing
some component or assumption. It is possible that parameter space exists that permits
intermediate levels of TEs in an asexual population, perhaps sustained by high rates of horizontal
transfer, which there is evidence for in bdelloids. The HGT idea has some support from the
increased prevalence of class II DNA transposons in bdelloids, given their greater propensity for
horizontal transfer, but we found that the basally divergent lineage leading to *D. carnosus* had a
profile more like the monogonont rotifers with regard to non-LTR retrotransposon abundance,
despite sharing other features with the rest of bdelloids. Alternatively, some TEs might have
been co-opted to provide beneficial functions, which is hypothesised to explain the unusually
large and complex *Terminon* repeats. Other TEs may have evolved strong site-specificity to
neutral genome regions to mitigate negative effects of transposition. This idea is supported by
the preference shown for insertions into gene-poor regions that are probably at or near the
telomeres, although it seems unlikely that the full complement of bdelloid TEs have accumulated
in this way.

We also ask what forces may be acting to suppress TE activity in bdelloid populations. A major
finding is that an abundance and diversity of RNAi gene silencing pathways, characterised by a
large expansion of Argonaute and RdRP genes, appears to be a unique feature of bdelloid
genomes. The precise origins and functions of these divergent Ago and RdRP clades are yet to
be elucidated, but it seems likely that such an extended arsenal of TE defence genes offers
enhanced protection against the deleterious effects of TE activity, particularly if bdelloid
populations cannot keep TEs in check through sexual processes but are still exposed to new
invasions via HGT. This enhanced RNAi system may then provide a more deterministic action
against TEs, as opposed to stochastic forces (such as genetic drift) that may predominate in other
animal groups and may potentially explain the greater uniformity of TE proportions in bdelloids
relative to monogononts.

An alternative resolution is that the assumption of no recombination and strict clonality is not
met in bdelloids. Previous work, for example, proposed that intra-individual recombination
during the repair of DSBs caused by desiccation could provide a mechanism to keep TE numbers
in check. We found no evidence here that overall TE loads were lower in desiccating species
than nondesiccating species, a finding not limited to bdelloids, or for differences in activity or
rate of turnover between them. It remains possible that an equivalent mechanism, such as mitotic
recombination, or unusual DNA repair mechanisms operating in nondesiccating species as well,
could still act to limit TE proliferation and facilitate elimination of inserted copies, maintaining

an equilibrium between TE spread and removal by excision and selection. Finally, there could be
some hidden mechanism of inter-individual recombination that facilitates TE removal. We found
no evidence for its action here, but further work is needed for a final answer on the conundrum
of bdelloid asexuality.
733
734
Materials and methods
736
Rotifer sampling and culture. For most samples, individual rotifers were collected from
permanent and temporary freshwater habitats around Imperial College London's Silwood Park
campus, Ascot, UK between May 2015 and February 2019. Three samples (*R. magnacalcarata*
RM9 and RM15, and *R. socialis* RS1) were collected from a freshwater spring in Fontaneto
d'Agogna, Italy in 2016 (see S1 Data for details). Although we focused on the genera *Adineta*
and *Rotaria*, we also included two individuals from the desiccation-tolerant species
Didymodactylos carnosus. Preliminary phylogenetic data had identified this as a distant outgroup
to the focal genera, useful in rooting phylogenetic trees and as a further independent datapoint to
test the generality of conclusions about bdelloids. A total of 26 samples were submitted for
single-individual, whole genome sequencing; for these, DNA was extracted using either a Chelex
preparation (Bio-Rad InstaGene Matrix) or a QIAamp DNA Micro Kit (Qiagen), and whole-
genome amplified using a REPLI-g Single Cell kit (Qiagen) before sequencing on either
Illumina NextSeq500 at the Department of Biochemistry, University of Cambridge (Cambridge,
UK), or Illumina HiSeq X at Edinburgh Genomics, University of Edinburgh (Edinburgh, UK).
For *A. ricciae* ARIC003, DNA was extracted from ~200 animals descended from a single
individual before whole-genome amplification. For *B. calyciflorus* PSC1, individuals for DNA
extractions were derived from an individual isolate from a laboratory stock population previously
isolated from field-collected resting eggs [111]. DNA was extracted from ~5000 starved
individuals using a phenol-chloroform protocol and sequenced on the Illumina NextSeq500 at
the Max Planck Institute for Evolutionary Biology. Three 10x Genomics Chromium 'linked
reads' libraries were generated for *A. steineri*, *Rotaria* sp. 'Silwood-1' and *R. sordida*; for these,
high molecular weight DNA was extracted from thousands of animals reared clonally from a
single wild-caught animal, without whole-genome amplification, using the Chromium
Demonstrated Protocol "HMW gDNA Extraction from Single Insects"
(https://support.10xgenomics.com/permalink/7HBJeZucc80CwkMAmA4oQ2). Linked-read
libraries were constructed at the Centre for Genomics Research, Liverpool, UK, before
sequencing on the HiSeq X at Edinburgh Genomics. Further details on rotifer sampling, DNA
extraction and sequencing are provided in S1 Data.
765
Biological replicates. To check the repeatability of the whole-genome amplification (WGA),
sequencing, assembly and analysis pipelines, we included several samples that were either
biological replicates of the same rotifer clone, or where high-quality genomes were available for
the same clone from unamplified source material. Specifically, for *Rotaria* sp. 'Silwood-2' we
isolated two consecutive offspring from the same wild-caught mother and conducted WGA,
sequencing, assembly and analysis for these sisters independently (as Rp_RPSE411 and
Rp_RPSE503). From the same clonal laboratory line of *Rotaria* sp. 'Silwood-1' that was used
for 10x Genomics DNA preparation, we isolated two more individuals and processed each
independently using the WGA workflow (as Rw_RSIL801 and Rw_RSIL802). Finally, we
applied the WGA method to DNA from *A. ricciae*, for which a previous assembly was available
from unamplified DNA [46] on the same clonal culture and included this replicate in
downstream analyses alongside the earlier reference assembly.
778

779 **Data filtering and genome assembly.** We generated two assembly versions for each of the
780 single-individual rotifer samples. The 'reference' assemblies were scaffolded and polished to
781 result in haploid assemblies with improved contiguity. The 'maximum haplotig' ('maxhap')
782 assemblies instead retained highly similar contigs that might otherwise be removed during
783 assembly polishing. Our pipeline is outlined as follows.
784
For the Illumina libraries, raw sequence data were filtered for low quality bases and adapter
sequence using BBTools v38.73 'bbduk' (https://sourceforge.net/projects/bbmap/), and error
corrected using BBTools 'tadpole'. Data quality was inspected manually using FastQC v0.11.5
[112] aided by MultiQC [113] visualisation. For the *A. steineri*, *R.* sp. 'Silwood-1' and *R.*
sordida linked-read libraries, data were assembled into haploid ('pseudohap') and diploid
('megabubbles') genome representations using the 10x Genomics proprietary software
Supernova v2.1.1 [114] and further scaffolded with ARKS v1.0.4 [115]. All raw sequencing data
are deposited in the relevant International Nucleotide Sequence Database Collaboration (INSDC)
databases under the Study ID PRJEB39843 (see S1 Data for run accessions and counts for raw
and filtered data).
795
For the single-individual samples, an initial assembly was generated using SPAdes v3.13.0 [116]
with default settings. Contaminating reads from non–target organisms, identified based on
aberrant GC content, read coverage and/or taxonomic annotation, were then identified and
removed using BlobTools v1.1.1 [117,118]. For *R. magnacalcarata* and *R. socialis* samples,
resultant haplotigs were then collapsed using Redundans [119] with default settings before
scaffolding and gap filling with SSPACE v3.0 and GapCloser v1.12 respectively [120,121]. For
A. steineri, *R.* sp. 'Silwood-1' and *R. sordida* single-individual samples, the scaffolding step was
performed with RaGOO v1.1 [122,123], using the matching 10x Genomics 'pseudohap'
assembly as a reference (contigs from *R.* sp. 'Silwood-2' were scaffolded using the *R.* sp.
'Silwood-1' 10x reference), specifying the '-C' parameter to prevent concatenation of unaligned
contigs. Scaffolded assemblies were subjected to further rounds of BlobTools to remove any
additional sequences derived from non-target organisms. These assemblies were designated the
reference set described above.
809
For the maxhap assemblies, filtered fastq files were first generated by mapping the original
(trimmed and error-corrected) sequencing reads to each reference genome, using the
'outm=filtered_R#.fq' functionality of BBTools 'bbmap', and then reassembled with SPAdes,
increasing the final kmer value to 121. Assembly metrics were summarised using 'calN50.js'
(https://github.com/lh3/calN50), which reports the 'expected scaffold size' (AU) as an alternative
metric of assembly contiguity that is less biased than N50 (defined as the area under the
cumulative genome span versus contig length graph, equivalent to the expected scaffold size for
a randomly chosen assembly location [124]). Gene-completeness scores for core eukaryotic ($n =$
303) and metazoan ($n = 978$) genes were calculated for all assemblies using BUSCO v3.0.2
[125] with default settings. Reference and maxhap assemblies for *B. calyciflorus* PSC1 and *D.*
carnosus are the same, due to a lack of appropriate data for scaffolding.
821
Gene prediction. Gene prediction was performed on reference assemblies using one of three
approaches, depending on the availability of RNA-seq data. For *B. calyciflorus*, *A. ricciae*, and
all *R. magnacalcarata*, *R. socialis*, and *R. sordida* assemblies, published RNA-seq data
[109,110,126] were downloaded from NCBI Sequence Read Archive (SRA), quality-trimmed
using BBTools 'bbduk' with default settings and aligned to the genomic scaffolds using STAR
v2.7.3a [127] with the option '--twoPassMode Basic'. Aligned BAM files were then provided to
BRAKER v2.1.2 [128–132] with default settings for gene prediction. For *A. steineri*, *R.* sp.

'Silwood-1' and *R*. sp. 'Silwood-2' assemblies, RNA-seq data from a related species (*A. ricciae*
and *R. magnacalcarata* respectively) were used instead, aligned using BBTools 'bbmap' with the
options 'maxindel=200k minid=0.5', before gene prediction with BRAKER as above. Finally,
for the distantly related *D. carnosus*, BRAKER was run using gene-model parameters estimated
from BUSCO analysis of the genomic scaffolds. The quality of predicted proteins was assessed
using BUSCO in protein mode. Intragenomic divergence between homologous gene copies and
collinearity was calculated as for Nowell *et al.* (2018) and described in S1 Note. Genome
assemblies and gene predictions were converted to EMBL format using EMBLmyGFF3 v2
[133], and are deposited at DDBJ/ENA/GenBank under the Study ID PRJEB39843 (see **Table 1**
and S1 Table for individual GenBank accessions).
839
Rotifer phylogeny. Evolutionary relationships among new genomes and published genomes of
rotifers were determined using a core-genome phylogenomics approach based on the BUSCO
eukaryotic gene set. For genomes from species with very high intragenomic homologous
divergence (*A. ricciae* and *A. vaga*), redundancy was removed by selecting the copy with the
highest BUSCO score for all BUSCO genes with multiple copies, using the script
BUSCO_collapse_multicopy.pl' (https://github.com/reubwn/scripts). One-to-one co-orthologs
found in at least 95% of the samples were then identified using the script
'BUSCO_phylogenomics.py' (https://github.com/jamiemcg/BUSCO_phylogenomics). Protein
sequences were aligned using Clustalo [134] and concatenated in Geneious R9 [135]. The full
alignment was checked by eye and sections with ambiguous alignment within the bdelloid clade
were removed across all sequences to avoid aligning potential paralogs or homoeologs.
Translation errors arising from annotation issues in specific bdelloid genomes were identified by
obvious mismatches to the consensus of closely related genomes, and the affected residues were
deleted in the affected genome only. Potential alignment issues within the monogonont clade
were less obvious owing to the substantial genetic divergence from bdelloids and the smaller
number of genomes and replicates, so corrections were less stringent. A maximum-likelihood
phylogeny was then estimated using IQ-TREE v1.6.12, with automatic model selection
(VT+F+I+G4) [136,137]. Branching support was assessed using SH-aLRT and ultrafast
bootstrap sampling ('-alrt 1000 -bb 5000') [138,139].
859
Repeat annotation and TE dynamics. TEs and other repeats were identified using the
RepeatModeler and RepeatMasker pipelines. For each sample, a *de novo* repeat library was
generated directly from the assembled nucleotides using RepeatModeler2 [140] and combined
with a database of 12,662 protostome repeats from Repbase v23.08 [141] and 278 additional TEs
manually curated from the *A. vaga* genome [44]. Repeats and TEs were then detected and
classified using RepeatMasker v4.1.0 [84], and resultant outputs were post-processed using the
'One code to find them all' Perl script [142]. The breakdown of TE superfamilies in the final
database was 4,145 DNA transposons (including 300 rolling circles), 5,523 LTRs, 2,583 LINEs
(including SINEs), 227 PLEs, and 165 simple or low-complexity repeats. TE content (expressed
as a proportion of genome size) was mapped onto the phylogeny using 'contMap' in the Phytools
v0.6-99 package in R v3.6.0 [143,144]. There is no module for the detection of class II MITEs in
RepeatMasker; for these, the separate program Generic Repeat Finder (GRF) was run using
default parameters. TE dynamics were investigated by constructing Kimura 2-parameter
divergence [85] landscapes using the utility scripts in the RepeatMasker package and plotted
using custom scripts (see below). Selected assemblies were also submitted to the REPET v2.5
'TEdenovo' [145,146] TE detection and annotation pipeline with default parameters, for
comparison. In addition, for *D. carnosus* and *R. sordida* (using 10x Genomics) reference
assemblies, we increased the parameter 'minNbSeqPerGroup' from 3 to 5 to evaluate
contribution from tetraploid genes, which was judged to be negligible. Although

REPET*denovo* TE consensus sequences are automatically classified using Wicker's TE
classification [147], RepeatMasker was additionally applied for further TE classification,
detection and landscape divergence plot building.
882
The presence or absence of specific LTR retrotransposon (LTR-R) insertions in our population
data was inferred using a read-mapping approach. Specifically, the presence of a given insertion
was scored based on the alignment score of the 'best' read that mapped continuously and
contiguously across the LTR-genome boundary. First, full-length LTR-Rs (i.e. those with
annotated 5' and 3' LTR regions) were identified from each reference assembly using
LTR_retriever v2.8 [148]. Three filters were then applied to remove false positives. Candidates
that showed an overlap with a predicted gene in the 5' or 3' LTR itself or an 'N' base within 150
bases upstream or downstream of its genomic location that might indicate local mis-assembly
were removed. Candidates also required supporting evidence of LTR homology from a separate
RepeatMasker annotation of the reference assembly. For each remaining LTR-R, a library of
'LTR-tags' was then generated by extracting a 100 bp sequence that spanned 50 bases into the
genomic (i.e., non-TE) region of the insertion site from both the 5' and 3' terminal repeated
regions. Thus, each pair of 'LTR-tags' represents an insertion of a particular LTR into a specific
location in the focal genome, and a score is calculated based on the alignment information
contained in the CIGAR string of the 'best' read (i.e. with the highest number of alignment
matches) from the SAM mapping file: $S_i = ((M_{Li} - X_{Li}) + (M_{Ri} - X_{Ri}))/200$, where M_{Li} is the
number of alignment matches for the left-hand tag for LTR i, penalised by the number of
mismatches X_{Li}, with equivalent scoring for the right-hand tag. Since tag length is 100 bases, the
maximum score for a perfect alignment is 200, or 1 after normalisation. The number of mapped
reads is also recorded to provide an estimate of coverage (but note that S_i is taken from the best
read only). Sequencing reads from all single-individual rotifer samples were aligned to the
filtered LTR-tag set using BBTools 'bbmap' with the parameters 'minid=0.5 local=t' and scored
using the above system. Because orthologous LTR-Rs may be identified from searches started in
different genomes, we identified these cases by reconstructing the phylogeny of the LTR-tags
and any with pairwise sequence divergence less than 0.1 were collapsed to yield a condensed
final matrix.
909
The LTR-tag case-study in Fig. 4b was selected for closer investigation in the draft assemblies
after consideration of several examples, because it illustrates variability for an element insertion
site within a species and indicates that Class I TEs can insert in coding regions, with potential
fitness consequences. The LTR-tags were mapped to the RM15 draft assembly using Geneious
Prime v2020.1.2 [135], and were found to match an element annotated by LTR_retriever,
containing four predicted genes. In the RM9 draft assembly, only the left-hand tag was mapped,
as the scaffold ended before the inserted element was fully assembled. For the same reason, the
element in RM9 had not been annotated as such by LTR_retriever, but the sequence is nearly
identical (99.7%) to the insertion in RM15 along its aligned length (except that the annotations
predicted three element-associated genes rather than four). The scaffolds were trimmed to the
focal gene and aligned, and the region was used as a BLASTn query against local databases for
two other *R. magnacalcarata* reference genomes where the LTR-tag was absent: MAG1 and
Rg2018. In each case, this provided the location of a closely similar but uninterrupted copy of
the focal gene, although the annotations of the gene's structure differed slightly among genomes.
These scaffolds were trimmed and aligned against the copies from RM9 and RM15, using the
Geneious alignment tool with default settings, except that the gap extension penalty was reduced
from 3 to 0.2 to enable the algorithm to handle the element insertion. Local features were
manually reannotated to illustrate the interpretation provided in the text. To investigate the
potential function of the interrupted gene, the copy from MAG1 (g37061) was translated and

used as a BLASTp query against the NCBI RefSeq Protein Database [149]. A region of
approximately 1000 residues was found to have weak similarity (~25% pairwise identity) to
proteins annotated as midasins, from a range of eukaryotes. As a final step, the intact gene from
MAG1 was used as a BLASTn query against the full draft genomes of RM9 and RM15, which
revealed a separate scaffold in each case, containing a partial copy of the gene in which the
coding sequence was intact across the junction spanned by the LTR-tag, and the element
insertion was absent.
936
Recombination analyses. We tested sexual versus clonal patterns of variation in LTR presence
and absences. First, we calculated consistency indices (CI) with parsimony reconstruction of the
binary matrix. LTR-tags with scores > 0.875 were coded as present (i.e. no more than half of the
genome context or LTR region from both left and right LTR-tags was missing) and < 0.875
coded as absent (alternative thresholds led to the same qualitative results). A CI = 1 indicates
perfect nesting with no homoplasy, whereas a score less than 1 is expected if variation is shuffled
among loci and not tree-like. Next, we calculated the index of association and ran permutations
to test for significant linkage disequilibrium of the LTR-tag data relative to a null model of
random shuffling (expected in a fully outcrossing sexual population). We used the modified
index of association by Agapow and Burt (2001) [150] that corrects for an effect of the number
of loci on the index, and ran permutations using the 'ia' function in the Poppr v2.8.5 library
[151] in R. Data were coded as diploid and codominant presence/absence data (because of the
lack of diploid assemblies in the population-level data). Finally, for *R. magnacalcarata* and *R.*
socialis we ran simulations with the FacSexCoalescent simulator of Hartfield et al. (2018) [152]
to generate 50000 datasets with the same number of individuals and sampled binary loci as
observed, but with frequencies of sexual versus asexual reproduction within the populations
varying from 10^{-7} (i.e. negligible) to 1 (i.e. obligate sexual). We estimate the posterior
distribution of the frequency of sex for our observed samples using Approximate Bayesian
Computation on the simulated datasets implemented in the abc package in R.
956
Reverse transcriptase survey. A hidden Markov model (HMM) approach was used to survey
the predicted rotifer proteomes for proteins encoding the reverse transcriptase (RT) domain
(Pfam ID PF00078). First, a HMM was constructed from an alignment of 51 RT domains from
across the tree of life [57], supplemented with 67 bdelloid-specific retroelements
[44,57,58,61,63,64] (S9 Fig). Alternative transcripts were first removed from predicted
proteomes and proteins with a significant match (E-value ≤ 1e−5) were identified and inserted
into the core RT alignment using HMMER v3.2.1 'hmmsearch' and 'hmmalign', respectively
(http://hmmer.org/). Maximum likelihood phylogenies were then constructed using IQ-TREE as
above, specifying the root of the phylogeny to be on the branch leading to the bacterial retrons
[57]. Trees were manipulated using FigTree v1.4.4 (https://github.com/rambaut/figtree),
colouring the identified RT-encoding rotifer proteins based on their phylogenetic position. The
span accounted for by genome 'features' (other genes, other TEs and telomeric repeats) in a 25
kb window around each identified RT-containing protein was summarised using BEDTools
v2.29.2 'intersect' and 'groupby' [153]. Other genes were counted as predicted coding regions
that did not overlap with any TE annotation. Genomic locations of the telomeric hexamer
'TGTGGG' [58] were identified using EMBOSS 'fuzznuc' [154], excluding any hexamer that
overlapped with a predicted CDS. Note that the telomeric repeat for *Brachionus* is not known,
but the sequence above was among the most frequent G-rich hexamers identified in the PSC1
genome (see S2 Note).
976
977 **TE silencing machinery survey.** A similar HMM based approach was used to characterise gene
978 copy evolution of three key pathways involved in RNAi gene-silencing. Putative Argonaute

proteins were identified based on the presence of both the PAZ and PIWI domains (Pfam IDs PF02170 and PF02171 respectively), putative Dicer proteins were identified based on the presence of both PAZ and Dicer (PF03368) domains, and putative RdRP proteins were identified based on the presence of the RdRP domain (PF05183). Stockholm files were downloaded from Pfam [155] and aligned to the proteomes using HMMER (E-value $\leq$ 1e−5) as above. Reference proteomes from a selection of eukaryotic species to represent the diversity and distribution of Argonaute, Dicer and RdRP proteins were downloaded (June 2020) from UniProt and subjected to the same procedure: *Arabidopsis thaliana* (UP000006548), *Oryza sativa* (UP000007015), *Neurospora crassa* (UP000001805), *Schizosaccharomyces pombe* (UP000002485), *Laccaria bicolor* (UP000001194), *Dictyostelium discoideum* (UP000002195), *D. melanogaster* (UP000000803), *C. elegans* (UP000001940), *H. exemplaris* (UP000192578), *H. robusta* (UP000015101), *L. gigantea* (UP000030746), *S. haematobium* (UP000054474), *B. plicatilis* (UP000276133), *Branchiostoma floridae* (UP000001554) and *Homo sapiens* UP000005640. Proteins were aligned using either 'hmmalign' from the HMMER package or Clustalo, and ML phylogenies were constructed using IQ-TREE as above.

Comparative protein–domain abundance plots were constructed using counts of Pfam entries parsed directly from InterProScan5 [156] annotation of predicted proteomes. The 'abundance score' was computed as the (log) ratio of domain counts in bdelloids divided by the domain counts in eukaryotes, corrected for inflation in bdelloids due to the ancient whole-genome duplication by dividing the former by two. This correction is likely to be conservative, since many loci have lost one branch of the ancient duplication (i.e. tetraploidy is degenerate). To check that the putative RdRP expansion was indeed eukaryotic in origin, rather than viral, the HMMs for four viral RdRP families (PF00680, PF00978, PF00998 and PF02123) were downloaded from Pfam and submitted to the same search protocol, with zero hits to bdelloid proteomes recorded.

Statistical analyses. To assess differences in TE content between desiccating and non-desiccating rotifer species, we ran Bayesian linear mixed-effects models of TE content (as a percentage of genome span) including desiccation as a two-level fixed factor and sample ID as a random intercept term. The BUSCO gene phylogeny was used to account for non-independence among species. A separate model was run for each TE superfamily (including 'unclassified' TEs) as well as for all TEs combined. Inverse-Wishart priors were used for the random and residual variances, and models were run for 42,0000 iterations with a burn-in of 20,000 and a thinning interval of 200. This resulted in 2,000 stored samples of the posterior with minimal autocorrelation in all cases (< 0.2) [157]. Models were run using the MCMCglmm v2.29 [158] package in R. The phylogenetic signal, defined as the proportion of the total variance in TE content attributable to the phylogeny [159], was estimated from the MCMCglmm model output using the formula: $\lambda = \sigma P^2/(\sigma P^2 + \sigma R^2)$.

The density of genomic features surrounding BUSCO genes versus RT-containing genes were compared using linear mixed effects models with species as a random effect and gene class (BUSCO, PLE, LINE or LTR) as a fixed effect, run using lme4 [160] v1.1-21 in R. TE length distributions were compared using the same approach, with species as a random effect and binary factors specifying monogononts versus bdelloid, and desiccating versus nondesiccating species.

Code availability. All TE analysis scripts used in this study are available at https://github.com/reubwn/te-evolution.

Acknowledgements

Genome sequencing was performed by the UK Natural Environment Research Council (NERC) Biomolecular Analysis Facility at the Centre for Genomic Research (CGR) at the University of Liverpool (NBAF-Liverpool) and the DNA Sequencing Facility in the Biochemistry Department at the University of Cambridge. The authors wish to thank the following: Christiane Hertz-Fowler, Pia Koldkjær and John Kenny (CGR), Shilo Dickens and Nataliya Scott (Cambridge). Matthew Arno and Colin Sharp (Edinburgh), and Steven Van Belleghem (University of Puerto Rico) for support with the planning and execution of various aspects of genome sequencing and/or assembly, Tom Smith and Anita Kristiansen for rotifer sampling, and Mike Tristem for helpful discussions on detecting LTR polymorphisms.

References

1. Orgel LE, Crick FH. Selfish DNA: the ultimate parasite. Nature. 1980;284: 604–607. doi:10.1038/284604a0

2. Hickey DA. Selfish DNA: a sexually-transmitted nuclear parasite. Genetics. 1982;101: 519–531.

3. Charlesworth B, Charlesworth D. The population dynamics of transposable elements. Genet Res. 1983;42: 1–27. doi:10.1017/S0016672300021455

4. Doolittle RF, Feng D-F, Johnson MS, McClure MA. Origins and evolutionary relationships of retroviruses. Q Rev Biol. 1989;64: 1–30.

5. Eickbush TH, Malik HS. Origins and Evolution of Retrotransposons. In: Craig NL, Lambowitz AM, Craigie R, Gellert M, editors. Mobile DNA II. American Society of Microbiology; 2002. pp. 1111–1144. doi:10.1128/9781555817954.ch49

6. Robertson HM. Evolution of DNA Transposons in Eukaryotes. In: Craig NL, Lambowitz AM, Craigie R, Gellert M, editors. Mobile DNA II. American Society of Microbiology; 2002. pp. 1093–1110. doi:10.1128/9781555817954.ch48

7. Capy P, Gasperi G, Biémont C, Bazin C. Stress and transposable elements: co-evolution or useful parasites? Heredity. 2000;85 (Pt 2): 101–106. doi:10.1046/j.1365-2540.2000.00751.x

8. Burt A, Trivers R. Genes in Conflict: The Biology of Selfish Genetic Elements. Harvard University Press; 2009.

9. Bourgeois Y, Boissinot S. On the Population Dynamics of Junk: A Review on the Population Genomics of Transposable Elements. Genes. 2019;10. doi:10.3390/genes10060419

10. Finnegan DJ. Transposable elements. Curr Opin Genet Dev. 1992;2: 861–867.

11. Nuzhdin SV. Sure facts, speculations, and open questions about the evolution of transposable element copy number. Genetica. 1999;107: 129–137.

12. Montgomery E, Charlesworth B, Langley CH. A test for the role of natural selection in the stabilization of transposable element copy number in a population of *Drosophila melanogaster*. Genet Res . 1987;49: 31–41. doi:10.1017/s0016672308009634

13. Langley CH, Montgomery E, Hudson R, Kaplan N, Charlesworth B. On the role of unequal exchange in the containment of transposable element copy number. Genet Res. 1988;52: 223–235. doi:10.1017/s0016672300027695

14. Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, et al. Initial sequencing and analysis of the human genome. Nature. 2001;409: 860–921. doi:10.1038/35057062

15. Craig NL. A Moveable Feast: An Introduction to Mobile DNA. In: Craig NL, Chandler M, Gellert M, Lambowitz AM, Rice PA, Sandmeyer S, editors. Mobile DNA III. American Society of Microbiology; 2015. pp. 3–39. doi:10.1128/microbiolspec.MDNA3-0062-2014

16. Chalopin D, Naville M, Plard F, Galiana D, Volff J-N. Comparative analysis of transposable elements highlights mobilome diversity and evolution in vertebrates. Genome Biol Evol. 2015;7: 567–580. doi:10.1093/gbe/evv005

17. Petersen M, Armisén D, Gibbs RA, Hering L, Khila A, Mayer G, et al. Diversity and evolution of the transposable element repertoire in arthropods with particular reference to insects. BMC Evol Biol. 2019;19: 11. doi:10.1186/s12862-018-1324-9

18. Szitenberg A, Cha S, Opperman CH, Bird DM, Blaxter ML, Lunt DH. Genetic drift, not life history or RNAi, determine long-term evolution of transposable elements. Genome Biol Evol. 2016;8: 2964–2978. doi:10.1093/gbe/evw208

19. Castanera R, López-Varas L, Borgognone A, LaButti K, Lapidus A, Schmutz J, et al. Transposable elements
versus the fungal genome: Impact on whole-genome architecture and transcriptional profiles. PLoS Genet.
2016;12: e1006108. doi:10.1371/journal.pgen.1006108
20. Doolittle WF, Sapienza C. Selfish genes, the phenotype paradigm and genome evolution. Nature. 1980;284:
601–603. doi:10.1038/284601a0
21. Charlesworth B, Langley CH. The population genetics of *Drosophila* transposable elements. Annu Rev Genet.
1989;23: 251–287. doi:10.1146/annurev.ge.23.120189.001343
22. Wright S, Finnegan D. Genome evolution: sex and the transposable element. Curr Biol. 2001;11: R296-9.
doi:10.1016/S0960-9822(01)00168-3
23. Schaack S, Choi E, Lynch M, Pritham EJ. DNA transposons and the role of recombination in mutation
accumulation in *Daphnia pulex*. Genome Biol. 2010;11: R46. doi:10.1186/gb-2010-11-4-r46
24. Petrov DA, Aminetzach YT, Davis JC, Bensasson D, Hirsh AE. Size matters: non-LTR retrotransposable
elements and ectopic recombination in *Drosophila*. Mol Biol Evol. 2003;20: 880–892.
doi:10.1093/molbev/msg102
25. Schwander T, Crespi BJ. Twigs on the tree of life? Neutral and selective models for integrating
macroevolutionary patterns with microevolutionary processes in the analysis of asexuality. Mol Ecol. 2009;18:
28–42. doi:10.1111/j.1365-294X.2008.03992.x
26. Charlesworth B, Langley CH. The evolution of self-regulated transposition of transposable elements.
Genetics. 1986;112: 359–383.
27. Dolgin ES, Charlesworth B. The fate of transposable elements in asexual populations. Genetics. 2006;174:
817–827. doi:10.1534/genetics.106.060434
28. Bast J, Jaron KS, Schuseil D, Roze D, Schwander T. Asexual reproduction reduces transposable element load
in experimental yeast populations. Elife. 2019;8. doi:10.7554/eLife.48548
29. Fujita MK, Singhal S, Brunes TO, Maldonado JA. Evolutionary Dynamics and Consequences of
Parthenogenesis in Vertebrates. Annu Rev Ecol Evol Syst. 2020. doi:10.1146/annurev-ecolsys-011720-
114900
30. Schaack S, Gilbert C, Feschotte C. Promiscuous DNA: horizontal transfer of transposable elements and why it
matters for eukaryotic evolution. Trends Ecol Evol. 2010;25: 537–546. doi:10.1016/j.tree.2010.06.001
31. Basten CJ, Moody ME. A branching-process model for the evolution of transposable elements incorporating
selection. J Math Biol. 1991;29: 743–761. doi:10.1007/bf00160190
32. Edwards RJ, Brookfield JFY. Transiently beneficial insertions could maintain mobile DNA sequences in
variable environments. Mol Biol Evol. 2003;20: 30–37. doi:10.1093/molbev/msg001
33. Silva JC, Loreto EL, Clark JB. Factors that affect the horizontal transfer of transposable elements. Curr Issues
Mol Biol. 2004;6: 57–71.
34. Peccoud J, Loiseau V, Cordaux R, Gilbert C. Massive horizontal transfer of transposable elements in insects.
Proc Natl Acad Sci U S A. 2017. doi:10.1073/pnas.1621178114
35. Boutin TS, Le Rouzic A, Capy P. How does selfing affect the dynamics of selfish transposable elements? Mob
DNA. 2012;3: 5. doi:10.1186/1759-8753-3-5
36. Startek M, Le Rouzic A, Capy P, Grzebelus D, Gambin A. Genomic parasites or symbionts? Modeling the
effects of environmental pressure on transposition activity in asexual populations. Theor Popul Biol. 2013;90:
145–151. doi:10.1016/j.tpb.2013.07.004
37. Neiman M, Meirmans S, Meirmans PG. What can asexual lineage age tell us about the maintenance of sex?
Ann N Y Acad Sci. 2009;1168: 185–200. doi:10.1111/j.1749-6632.2009.04572.x
38. Jaron KS, Bast J, Nowell RW, Rhyker Ranallo-Benavidez T, Robinson-Rechavi M, Schwander T. Genomic
features of asexual animals. bioRxiv. 2019. p. 497495. doi:10.1101/497495
39. Mark Welch DB, Ricci C, Meselson M. Bdelloid Rotifers: Progress in Understanding the Success of an
Evolutionary Scandal. Lost Sex. Springer, Dordrecht; 2009. pp. 259–279. doi:10.1007/978-90-481-2770-2_13
40. Robeson MS, King AJ, Freeman KR, Birky CW Jr, Martin AP, Schmidt SK. Soil rotifer communities are
extremely diverse globally but spatially autocorrelated locally. Proc Natl Acad Sci U S A. 2011;108: 4406–
4410. doi:10.1073/pnas.1012678108
41. Hudson CT, Gosse PH. The Rotifera or wheel-animalcules. Longmans, Green; 1886.
42. Donner J. Ordnung Bdelloidea. Akademie Verlag, Berlin; 1965. p. 297.
43. Mark Welch D, Meselson M. Evidence for the evolution of bdelloid rotifers without sexual reproduction or
genetic exchange. Science. 2000;288: 1211–1215. doi:10.1126/science.288.5469.1211
44. Flot J-F, Hespeels B, Li X, Noel B, Arkhipova I, Danchin EGJ, et al. Genomic evidence for ameiotic
evolution in the bdelloid rotifer *Adineta vaga*. Nature. 2013;500: 453–457. doi:10.1038/nature12326
45. Mark Welch DB, Mark Welch JL, Meselson M. Evidence for degenerate tetraploidy in bdelloid rotifers. Proc
Natl Acad Sci U S A. 2008;105: 5145–5149. doi:10.1073/pnas.0800972105
46. Nowell RW, Almeida P, Wilson CG, Smith TP, Fontaneto D, Crisp A, et al. Comparative genomics of
bdelloid rotifers: Insights from desiccating and nondesiccating species. PLoS Biol. 2018;16: e2004830.

doi:10.1371/journal.pbio.2004830

47.   Simion P, Narayan J, Houtain A, Derzelle A, Baudry L, Nicolas E, et al. Homologous chromosomes in
asexual rotifer *Adineta vaga* suggest automixis. bioRxiv. 2020. p. 2020.06.16.155473.
doi:10.1101/2020.06.16.155473

48.   Signorovitch A, Hur J, Gladyshev E, Meselson M. Allele sharing and evidence for sexuality in a
mitochondrial clade of bdelloid rotifers. Genetics. 2015;200: 581–590. doi:10.1534/genetics.115.176719

49.   Vakhrusheva OA, Mnatsakanova EA, Galimov YR, Neretina TV, Gerasimov ES, Ozerova SG, et al.
Recombination in a natural population of the bdelloid rotifer *Adineta vaga*. bioRxiv. 2018. p. 489393.
doi:10.1101/489393

50.   Arkhipova I, Meselson M. Transposable elements in sexual and ancient asexual taxa. Proc Natl Acad Sci U S
A. 2000;97: 14473–14477. doi:10.1073/pnas.97.26.14473

51.   Zhang H-H, Peccoud J, Xu M-R-X, Zhang X-G, Gilbert C. Horizontal transfer and evolution of transposable
elements in vertebrates. Nat Commun. 2020;11: 1362. doi:10.1038/s41467-020-15149-4

52.   Arkhipova I, Meselson M. Deleterious transposable elements and the extinction of asexuals. Bioessays.
2005;27: 76–85. doi:10.1002/bies.20159

53.   Ricci C. Anhydrobiotic capabilities of bdelloid rotifers. Hydrobiologia. 1998;387–388: 321–326.
doi:10.1023/A:1017086425934

54.   Gladyshev E, Meselson M. Extreme resistance of bdelloid rotifers to ionizing radiation. Proc Natl Acad Sci U
S A. 2008;105: 5139–5144. doi:10.1073/pnas.0800966105

55.   Hespeels B, Knapen M, Hanot-Mambres D, Heuskin A-C, Pineux F, Lucas S, et al. Gateway to genetic
exchange? DNA double-strand breaks in the bdelloid rotifer *Adineta vaga* submitted to desiccation. J Evol
Biol. 2014;27: 1334–1345. doi:10.1111/jeb.12326

56.   Gladyshev EA, Arkhipova IR. Genome structure of bdelloid rotifers: shaped by asexuality or desiccation? J
Hered. 2010;101 Suppl 1: S85-93. doi:10.1093/jhered/esq008

57.   Arkhipova IR, Pyatkov KI, Meselson M, Evgen'ev MB. Retroelements containing introns in diverse
invertebrate taxa. Nat Genet. 2003;33: 123–124. doi:10.1038/ng1074

58.   Gladyshev EA, Arkhipova IR. Telomere-associated endonuclease-deficient *Penelope*-like retroelements in
diverse eukaryotes. Proc Natl Acad Sci U S A. 2007;104: 9352–9357. doi:10.1073/pnas.0702741104

59.   Arkhipova IR, Yushenova IA, Rodriguez F. Giant reverse transcriptase-encoding transposable elements at
telomeres. Mol Biol Evol. 2017. doi:10.1093/molbev/msx159

60.   Arkhipova IR, Yushenova IA, Rodriguez F. Endonuclease-containing *Penelope* retrotransposons in the
bdelloid rotifer *Adineta vaga* exhibit unusual structural features and play a role in expansion of host gene
families. Mob DNA. 2013;4: 19. doi:10.1186/1759-8753-4-19

61.   Gladyshev EA, Meselson M, Arkhipova IR. A deep-branching clade of retrovirus-like retrotransposons in
bdelloid rotifers. Gene. 2007;390: 136–145. doi:10.1016/j.gene.2006.09.025

62.   Rodriguez F, Kenefick AW, Arkhipova IR. LTR-retrotransposons from bdelloid rotifers capture additional
ORFs shared between highly diverse retroelement types. Viruses. 2017;9. doi:10.3390/v9040078

63.   Gladyshev EA, Arkhipova IR. Rotifer rDNA-specific R9 retrotransposable elements generate an exceptionally
long target site duplication upon insertion. Gene. 2009;448: 145–150. doi:10.1016/j.gene.2009.08.016

64.   Gladyshev EA, Arkhipova IR. A subtelomeric non-LTR retrotransposon *Hebe* in the bdelloid rotifer *Adineta*
vaga is subject to inactivation by deletions but not 5' truncations. Mob DNA. 2010;1: 12. doi:10.1186/1759-
8753-1-12

65.   Kim H-S, Lee B-Y, Han J, Jeong C-B, Hwang D-S, Lee M-C, et al. The genome of the freshwater
monogonont rotifer *Brachionus calyciflorus*. Mol Ecol Resour. 2018;18: 646–655. doi:10.1111/1755-
0998.12768

66.   Han J, Park JC, Choi B-S, Kim M-S, Kim H-S, Hagiwara A, et al. The genome of the marine monogonont
rotifer *Brachionus plicatilis*: Genome-wide expression profiles of 28 cytochrome P450 genes in response to
chlorpyrifos and 2-ethyl-phenanthrene. Aquat Toxicol. 2019;214: 105230. doi:10.1016/j.aquatox.2019.105230

67.   Blommaert J, Riss S, Hecox-Lea B, Mark Welch DB, Stelzer CP. Small, but surprisingly repetitive genomes:
transposon expansion and not polyploidy has driven a doubling in genome size in a metazoan species
complex. BMC Genomics. 2019;20: 466. doi:10.1186/s12864-019-5859-y

68.   Mauer K, Hellmann SL, Groth M, Fröbius AC, Zischler H, Hankeln T, et al. The genome, transcriptome, and
proteome of the fish parasite *Pomphorhynchus laevis* (Acanthocephala). PLoS One. 2020;15: e0232973.
doi:10.1371/journal.pone.0232973

69.   Wey-Fabrizius AR, Herlyn H, Rieger B, Rosenkranz D, Witek A, Welch DBM, et al. Transcriptome data
reveal Syndermatan relationships and suggest the evolution of endoparasitism in Acanthocephala via an
epizoic stage. PLoS One. 2014;9: e88618. doi:10.1371/journal.pone.0088618

70.   Sielaff M, Schmidt H, Struck TH, Rosenkranz D, Mark Welch DB, Hankeln T, et al. Phylogeny of
Syndermata (syn. Rotifera): Mitochondrial gene order verifies epizoic Seisonidea as sister to endoparasitic
Acanthocephala within monophyletic Hemirotifera. Mol Phylogenet Evol. 2016;96: 79–92.

doi:10.1016/j.ympev.2015.11.017

71.   Laumer CE, Fernández R, Lemer S, Combosch D, Kocot KM, Riesgo A, et al. Revisiting metazoan phylogeny
with genomic sampling of all phyla. Proc Biol Sci. 2019;286: 20190831. doi:10.1098/rspb.2019.0831

72.   C. elegans Sequencing Consortium. Genome sequence of the nematode *C. elegans*: a platform for
investigating biology. Science. 1998;282: 2012–2018. doi:10.1126/science.282.5396.2012

73.   Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD, Amanatides PG, et al. The genome sequence of
Drosophila melanogaster. Science. 2000;287: 2185–2195. doi:10.1126/science.287.5461.2185

74.   Young ND, Jex AR, Li B, Liu S, Yang L, Xiong Z, et al. Whole-genome sequence of *Schistosoma*
haematobium. Nat Genet. 2012;44: 221–225. doi:10.1038/ng.1065

75.   Zhang G, Fang X, Guo X, Li L, Luo R, Xu F, et al. The oyster genome reveals stress adaptation and
complexity of shell formation. Nature. 2012;490: 49–54. doi:10.1038/nature11413

76.   Simakov O, Marletaz F, Cho S-J, Edsinger-Gonzales E, Havlak P, Hellsten U, et al. Insights into bilaterian
evolution from three spiralian genomes. Nature. 2013;493: 526–531. doi:10.1038/nature11696

77.   Gusev O, Suetsugu Y, Cornette R, Kawashima T, Logacheva MD, Kondrashov AS, et al. Comparative
genome sequencing reveals genomic signature of extreme desiccation tolerance in the anhydrobiotic midge.
Nat Commun. 2014;5: 4784. doi:10.1038/ncomms5784

78.   Albertin CB, Simakov O, Mitros T, Wang ZY, Pungor JR, Edsinger-Gonzales E, et al. The octopus genome
and the evolution of cephalopod neural and morphological novelties. Nature. 2015;524: 220–224.
doi:10.1038/nature14668

79.   Luo Y-J, Takeuchi T, Koyanagi R, Yamada L, Kanda M, Khalturina M, et al. The *Lingula* genome provides
insights into brachiopod evolution and the origin of phosphate biomineralization. Nat Commun. 2015;6: 8301.
doi:10.1038/ncomms9301

80.   Mikhailov KV, Slyusarev GS, Nikitin MA, Logacheva MD, Penin AA, Aleoshin VV, et al. The genome of
Intoshia linei affirms orthonectids as highly simplified spiralians. Curr Biol. 2016;26: 1768–1774.
doi:10.1016/j.cub.2016.05.007

81.   Hashimoto T, Horikawa DD, Saito Y, Kuwahara H, Kozuka-Hata H, Shin-I T, et al. Extremotolerant
tardigrade genome and improved radiotolerance of human cultured cells by tardigrade-unique protein. Nat
Commun. 2016;7: 12808. doi:10.1038/ncomms12808

82.   Yoshida Y, Koutsovoulos G, Laetsch DR, Stevens L, Kumar S, Horikawa DD, et al. Comparative genomics of
the tardigrades *Hypsibius dujardini* and *Ramazzottius varieornatus*. PLoS Biol. 2017;15: e2002266.
doi:10.1371/journal.pbio.2002266

83.   Adema CM, Hillier LW, Jones CS, Loker ES, Knight M, Minx P, et al. Whole genome analysis of a
schistosomiasis-transmitting freshwater snail. Nat Commun. 2017;8: 15451. doi:10.1038/ncomms15451

84.   Smit AFA, Hubley R, Green P. RepeatMasker Open-4.0. 2013-2015. Available: http://www.repeatmasker.org

85.   Kimura M. A simple method for estimating evolutionary rates of base substitutions through comparative
studies of nucleotide sequences. J Mol Evol. 1980;16: 111–120. doi:10.1007/BF01731581

86.   Kapusta A, Suh A. Evolution of bird genomes-a transposon's-eye view. Ann N Y Acad Sci. 2017;1389: 164–
185. doi:10.1111/nyas.13295

87.   Shao F, Han M, Peng Z. Evolution and diversity of transposable elements in fish genomes. Sci Rep. 2019;9:
15399. doi:10.1038/s41598-019-51888-1

88.   Belshaw R, Pereira V, Katzourakis A, Talbot G, Paces J, Burt A, et al. Long-term reinfection of the human
genome by endogenous retroviruses. Proc Natl Acad Sci U S A. 2004;101: 4894–4899.
doi:10.1073/pnas.0307800101

89.   Garbarino JE, Gibbons IR. Expression and genomic analysis of midasin, a novel and highly conserved AAA
protein distantly related to dynein. BMC Genomics. 2002;3: 18. doi:10.1186/1471-2164-3-18

90.   Li P-C, Li K, Wang J, Zhao C-Z, Zhao S-Z, Hou L, et al. The AAA-ATPase MIDASIN 1 functions in
ribosome biogenesis and is essential for embryo and root development. Plant Physiol. 2019;180: 289–304.
doi:10.1104/pp.18.01225

91.   Hur JH, Van Doninck K, Mandigo ML, Meselson M. Degenerate tetraploidy was established before bdelloid
rotifer families diverged. Mol Biol Evol. 2009;26: 375–383. doi:10.1093/molbev/msn260

92.   Eyres I, Frangedakis E, Fontaneto D, Herniou EA, Boschetti C, Carr A, et al. Multiple functionally divergent
and conserved copies of alpha tubulin in bdelloid rotifers. BMC Evol Biol. 2012;12: 148. doi:10.1186/1471-
2148-12-148

93.   Song M, Boissinot S. Selection against LINE-1 retrotransposons results principally from their ability to
mediate ectopic recombination. Gene. 2007;390: 206–213. doi:10.1016/j.gene.2006.09.033

94.   Arkhipova IR, Yushenova IA. Giant transposons in eukaryotes: Is bigger better? Genome Biol Evol. 2019;11:
906–918. doi:10.1093/gbe/evz041

95.   Cooper DM, Schimenti KJ, Schimenti JC. Factors affecting ectopic gene conversion in mice. Mamm Genome.
1998;9: 355–360. doi:10.1007/s003359900769

96.   Höck J, Meister G. The Argonaute protein family. Genome Biol. 2008;9: 210. doi:10.1186/gb-2008-9-2-210

97.   Juliano C, Wang J, Lin H. Uniting germline and stem cells: the function of Piwi proteins and the piRNA
pathway in diverse organisms. Annu Rev Genet. 2011;45: 447–469. doi:10.1146/annurev-genet-110410-
132541

98.   Ghildiyal M, Zamore PD. Small silencing RNAs: an expanding universe. Nat Rev Genet. 2009;10: 94–108.
doi:10.1038/nrg2504

99.   de Jong D, Eitel M, Jakob W, Osigus H-J, Hadrys H, Desalle R, et al. Multiple dicer genes in the early-
diverging metazoa. Mol Biol Evol. 2009;26: 1333–1340. doi:10.1093/molbev/msp042

100.   Zong J, Yao X, Yin J, Zhang D, Ma H. Evolution of the RNA-dependent RNA polymerase (RdRP) genes:
duplications and possible losses before and after the divergence of major eukaryotic groups. Gene. 2009;447:
29–39. doi:10.1016/j.gene.2009.07.004

101.   Yigit E, Batista PJ, Bei Y, Pang KM, Chen C-CG, Tolia NH, et al. Analysis of the *C. elegans* Argonaute
family reveals that distinct Argonautes act sequentially during RNAi. Cell. 2006;127: 747–757.
doi:10.1016/j.cell.2006.09.033

102.   Shi Z, Montgomery TA, Qi Y, Ruvkun G. High-throughput sequencing reveals extraordinary fluidity of
miRNA, piRNA, and siRNA pathways in nematodes. Genome Res. 2013;23: 497–508.
doi:10.1101/gr.149112.112

103.   Buck AH, Blaxter M. Functional diversification of Argonautes in nematodes: an expanding universe. Biochem
Soc Trans. 2013;41: 881–886. doi:10.1042/BST20130086

104.   Aravin AA, Hannon GJ, Brennecke J. The Piwi-piRNA pathway provides an adaptive defense in the
transposon arms race. Science. 2007;318: 761–764. doi:10.1126/science.1146484

105.   Sarkies P, Selkirk ME, Jones JT, Blok V, Boothby T, Goldstein B, et al. Ancient and novel small RNA
pathways compensate for the loss of piRNAs in multiple independent nematode lineages. PLoS Biol. 2015;13:
e1002061. doi:10.1371/journal.pbio.1002061

106.   Shirayama M, Seth M, Lee H-C, Gu W, Ishidate T, Conte D Jr, et al. piRNAs initiate an epigenetic memory of
nonself RNA in the C. elegans germline. Cell. 2012;150: 65–77. doi:10.1016/j.cell.2012.06.015

107.   Rodriguez F, Arkhipova IR. Multitasking of the piRNA silencing machinery: targeting transposable elements
and foreign genes in the bdelloid rotifer *Adineta vaga*. Genetics. 2016;203: 255–268.
doi:10.1534/genetics.116.186734

108.   Gladyshev EA, Meselson M, Arkhipova IR. Massive horizontal gene transfer in bdelloid rotifers. Science.
2008;320: 1210–1213. doi:10.1126/science.1156407

109.   Boschetti C, Carr A, Crisp A, Eyres I, Wang-Koh Y, Lubzens E, et al. Biochemical diversification through
foreign gene expression in bdelloid rotifers. PLoS Genet. 2012;8: e1003035.
doi:10.1371/journal.pgen.1003035

110.   Eyres I, Boschetti C, Crisp A, Smith TP, Fontaneto D, Tunnacliffe A, et al. Horizontal gene transfer in
bdelloid rotifers is ancient, ongoing and more frequent in species from desiccating habitats. BMC Biol.
2015;13: 90. doi:10.1186/s12915-015-0202-9

111.   Becks L, Agrawal AF. The effect of sex on the mean and variance of fitness in facultatively sexual rotifers. J
Evol Biol. 2011;24: 656–664. doi:10.1111/j.1420-9101.2010.02199.x

112.   Andrews S. FastQC: A quality-control tool for high-throughput sequence data. 2015. Available:
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

113.   Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for multiple tools and
samples in a single report. Bioinformatics. 2016;32: 3047–3048. doi:10.1093/bioinformatics/btw354

114.   Weisenfeld NI, Kumar V, Shah P, Church DM, Jaffe DB. Direct determination of diploid genome sequences.
Genome Res. 2017;27: 757–767. doi:10.1101/gr.214874.116

115.   Coombe L, Zhang J, Vandervalk BP, Chu J, Jackman SD, Birol I, et al. ARKS: chromosome-scale scaffolding
of human genome drafts with linked read kmers. BMC Bioinformatics. 2018;19: 234. doi:10.1186/s12859-
018-2243-x

116.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome
assembly algorithm and its applications to single-cell sequencing. J Comput Biol. 2012;19: 455–477.
doi:10.1089/cmb.2012.0021

117.   Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. Nat Methods.
2015;12: 59–60. doi:10.1038/nmeth.3176

118.   Laetsch DR, Blaxter ML. BlobTools: Interrogation of genome assemblies. F1000Res. 2017;6.
doi:10.12688/f1000research.12232.1

119.   Pryszcz LP, Gabaldón T. Redundans: an assembly pipeline for highly heterozygous genomes. Nucleic Acids
Res. 2016;44: e113. doi:10.1093/nar/gkw294

120.   Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. Scaffolding pre-assembled contigs using SSPACE.
Bioinformatics. 2011;27: 578–579. doi:10.1093/bioinformatics/btq683

121.   Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, et al. SOAPdenovo2: an empirically improved memory-
efficient short-read de novo assembler. Gigascience. 2012;1: 18. doi:10.1186/2047-217X-1-18

122. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018;34: 3094–3100.
doi:10.1093/bioinformatics/bty191

123. Alonge M, Soyk S, Ramakrishnan S, Wang X, Goodwin S, Sedlazeck FJ, et al. RaGOO: fast and accurate
reference-guided scaffolding of draft genomes. Genome Biol. 2019;20: 224. doi:10.1186/s13059-019-1829-6

124. Salzberg SL, Phillippy AM, Zimin A, Puiu D, Magoc T, Koren S, et al. GAGE: A critical evaluation of
genome assemblies and assembly algorithms. Genome Res. 2012;22: 557–567. doi:10.1101/gr.131383.111

125. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing genome
assembly and annotation completeness with single-copy orthologs. Bioinformatics. 2015;31: 3210–3212.
doi:10.1093/bioinformatics/btv351

126. Hanson SJ, Stelzer C-P, Welch DBM, Logsdon JM Jr. Comparative transcriptome analysis of obligately
asexual and cyclically sexual rotifers reveals genes with putative functions in sexual reproduction, dormancy,
and asexual egg production. BMC Genomics. 2013;14: 412. doi:10.1186/1471-2164-14-412

127. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq
aligner. Bioinformatics. 2013;29: 15–21. doi:10.1093/bioinformatics/bts635

128. Stanke M, Schöffmann O, Morgenstern B, Waack S. Gene prediction in eukaryotes with a generalized hidden
Markov model that uses hints from external sources. BMC Bioinformatics. 2006;7: 62. doi:10.1186/1471-
2105-7-62

129. Stanke M, Diekhans M, Baertsch R, Haussler D. Using native and syntenically mapped cDNA alignments to
improve de novo gene finding. Bioinformatics. 2008;24: 637–644. doi:10.1093/bioinformatics/btn013

130. Barnett DW, Garrison EK, Quinlan AR, Strömberg MP, Marth GT. BamTools: a C++ API and toolkit for
analyzing and managing BAM files. Bioinformatics. 2011;27: 1691–1692. doi:10.1093/bioinformatics/btr174

131. Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. BRAKER1: Unsupervised RNA-Seq-Based
Genome Annotation with GeneMark-ET and AUGUSTUS. Bioinformatics. 2016;32: 767–769.
doi:10.1093/bioinformatics/btv661

132. Hoff KJ, Lomsadze A, Borodovsky M, Stanke M. Whole-Genome Annotation with BRAKER. Methods Mol
Biol. 2019;1962: 65–95. doi:10.1007/978-1-4939-9173-0_5

133. Norling M, Jareborg N, Dainat J. EMBLmyGFF3: a converter facilitating genome annotation submission to
European Nucleotide Archive. BMC Res Notes. 2018;11: 584. doi:10.1186/s13104-018-3686-x

134. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of high-quality
protein multiple sequence alignments using Clustal Omega. Mol Syst Biol. 2011;7: 539.
doi:10.1038/msb.2011.75

135. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious Basic: an integrated and
extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics.
2012;28: 1647–1649. doi:10.1093/bioinformatics/bts199

136. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic algorithm for
estimating maximum-likelihood phylogenies. Mol Biol Evol. 2015;32: 268–274. doi:10.1093/molbev/msu300

137. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: fast model selection
for accurate phylogenetic estimates. Nat Methods. 2017;14: 587–589. doi:10.1038/nmeth.4285

138. Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and methods to
estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol. 2010;59:
307–321. doi:10.1093/sysbio/syq010

139. Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. UFBoot2: Improving the Ultrafast Bootstrap
Approximation. Mol Biol Evol. 2018;35: 518–522. doi:10.1093/molbev/msx281

140. Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, et al. RepeatModeler2 for automated
genomic discovery of transposable element families. Proc Natl Acad Sci U S A. 2020;117: 9451–9457.
doi:10.1073/pnas.1921046117

141. Bao W, Kojima KK, Kohany O. Repbase Update, a database of repetitive elements in eukaryotic genomes.
Mob DNA. 2015;6: 11. doi:10.1186/s13100-015-0041-9

142. Bailly-Bechet M, Haudry A, Lerat E. "One code to find them all": a perl tool to conveniently parse
RepeatMasker output files. Mob DNA. 2014;5: 13. doi:10.1186/1759-8753-5-13

143. Revell LJ. phytools: an R package for phylogenetic comparative biology (and other things). Methods Ecol
Evol. 2012;3: 217–223. doi:10.1111/j.2041-210X.2011.00169.x

144. R Core Team. R: A language and environment for statistical computing. 2016. Available: https://www.R-
project.org/

145. Quesneville H, Bergman CM, Andrieu O, Autard D, Nouaud D, Ashburner M, et al. Combined evidence
annotation of transposable elements in genome sequences. PLoS Comput Biol. 2005;1: 166–175.
doi:10.1371/journal.pcbi.0010022

146. Flutre T, Duprat E, Feuillet C, Quesneville H. Considering transposable element diversification in de novo
annotation approaches. PLoS One. 2011;6: e16526. doi:10.1371/journal.pone.0016526

147. Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, et al. A unified classification system for

1386    eukaryotic transposable elements. Nat Rev Genet. 2007;8: 973–982. doi:10.1038/nrg2165

1387 148. Ou S, Jiang N. LTR_retriever: A Highly Accurate and Sensitive Program for Identification of Long Terminal
1388    Repeat Retrotransposons. Plant Physiol. 2018;176: 1410–1422. doi:10.1104/pp.17.01310

1389 149. Pruitt KD, Tatusova T, Maglott DR. NCBI reference sequences (RefSeq): a curated non-redundant sequence
1390    database of genomes, transcripts and proteins. Nucleic Acids Res. 2007;35: D61-5. doi:10.1093/nar/gkl842

1391 150. Agapow P-M, Burt A. Indices of multilocus linkage disequilibrium. Mol Ecol Notes. 2001;1: 101–102.
1392    doi:10.1046/j.1471-8278.2000.00014.x

1393 151. Kamvar ZN, Tabima JF, Grünwald NJ. Poppr: an R package for genetic analysis of populations with clonal,
1394    partially clonal, and/or sexual reproduction. PeerJ. 2014;2: e281. doi:10.7717/peerj.281

1395 152. Hartfield M, Wright SI, Agrawal AF. Coalescence and Linkage Disequilibrium in Facultatively Sexual
1396    Diploids. Genetics. 2018;210: 683–701. doi:10.1534/genetics.118.301244

1397 153. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics.
1398    2010;26: 841–842. doi:10.1093/bioinformatics/btq033

1399 154. Rice P, Longden I, Bleasby A. EMBOSS: the European Molecular Biology Open Software Suite. Trends
1400    Genet. 2000;16: 276–277. doi:10.1016/s0168-9525(00)02024-2

1401 155. El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, et al. The Pfam protein families database in
1402    2019. Nucleic Acids Res. 2019;47: D427–D432. doi:10.1093/nar/gky995

1403 156. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5: genome-scale protein
1404    function classification. Bioinformatics. 2014;30: 1236–1240. doi:10.1093/bioinformatics/btu031

1405 157. Garamszegi LZ, editor. Modern Phylogenetic Comparative Methods and Their Application in Evolutionary
1406    Biology: Concepts and Practice. Springer, Berlin, Heidelberg; 2014. doi:10.1007/978-3-662-43550-2

1407 158. Hadfield JD. MCMC Methods for Multi-Response Generalized Linear Mixed Models: The MCMCglmm R
1408    Package. J Stat Softw. 2010;33. doi:10.18637/jss.v033.i02

1409 159. de Villemereuil P, Nakagawa S. General Quantitative Genetic Methods for Comparative Biology. In:
1410    Garamszegi LZ, editor. Modern Phylogenetic Comparative Methods and Their Application in Evolutionary
1411    Biology: Concepts and Practice. Berlin, Heidelberg: Springer Berlin Heidelberg; 2014. pp. 287–303.
1412    doi:10.1007/978-3-662-43550-2_11

1413 160. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. Journal of
1414    Statistical Software, Articles. 2015;67: 1–48. doi:10.18637/jss.v067.i01