The potential of parrotfish faeces in replenishing reefs with coral-associated microbiome

Trigal M. Velásquez-Rodríguez[1], Catalina Zuluaga-Arias[1], Sandra M. Montaño-Salazar[1,3], John M.

González[2] & Juan Armando Sánchez[1]*

[1] Laboratorio de Biología Molecular Marina-Biommar, Departamento de Ciencias Biológicas,

Universidad de los Andes, Bogotá, Colombia.

[2] Laboratorio de Ciencias Básicas Médicas, Facultad de Medicina, Universidad de los Andes, Bogotá,

Colombia

[3] Facultad de Ciencias, Universidad Nacional de Colombia, Bogotá, Colombia

Contact info

*Correspondence author: juansanc@uniandes.edu.co

Keywords

Sparisoma viride, coral microbiome, Symbiodinaceae, environmental reservoirs

Abstract

Sparisoma viride is the most abundant parrotfish in the Caribbean and is considered as the most important

excavator due to corallivore behaviour. Parrotfishes are a keystone group that favour the growth and

resilience of coral reefs removing macroalgae and structuring the benthic communities. The microbial

symbiotic communities are involved in multiple functions related to nutrition and immunity maintaining

corals health. Because *S. viride* scrape coral tissues, the skeleton and the algae on coral, it could be

important reservoirs or vectors of microorganism for the corals through the faeces dispersion, however,

the role of parrotfishes as reservoirs are poorly studied. Establishing microbial communities present in

parrotfish faeces will contribute to understand the ecological impact of parrotfishes in coral resilience. We

investigated the composition of disseminated bacteria and the extent to which the cell integrity of

dinoflagellate photosymbionts (Symbiodiniaceae) is maintained in the faeces and compare with sediments

and water column controls. Then, we analysed diversity and structure of bacterial communities at family

level and search similarities between faeces of the study and coral associated microbiome reported in the

literature. Similar levels of structural integrity and photosynthetic health of Symbidiodinaceae cells were

found in both faeces and reef sediments. Besides, the sediments microbiome echoes the parrotfish faecal

microbiome by sharing high diversity and a similar bacterial community composition. Several bacterial

families were present in parrotfish faeces and in coral microbiome reported in the literature highlighting

the dispersal potential of parrotfishes replenishing coral reefs. Despite the sampling limitations, these

findings uncover the potential role of the excavator parrotfish in enriching environmental reservoirs,

especially reef sediments, with coral-associated bacteria and photosynthetic microalgae. Parrotfishes

could reinforce the coral microbiome and facilitate coral symbiont acquisition, key features critical to

maintaining the fitness of one of the most threatened ecosystems on Earth. This finding could be

considered as a first step in uncovering a mechanism for reef-microbiome maintenance.

Introduction

The coral microbiome includes species-specific symbiotic interactions with unicellular dinoflagellates of

the family Symbiodiniaceae, endolithic green algae, bacteria, archaea, fungi, protists and viruses (Ricci et

al. 2019). From the surface mucus layer into the skeleton, coral-associated bacterial communities and

photosynthetic dinoflagellates support coral homeostasis (Ainsworth et al. 2015). Bacteria occupy specific

niches in the coral according to their function at the intracellular and perialgal space (Hernandez-Agreda

et al. 2017), where critical roles in catabolizing organic matter and recycling nutrients are performed

(Bourne et al. 2016). Particularly, bacteria located in the coral mucus supply nutrition while protecting

against pathogens and UV light, which is an emerging and dynamic process from the constant

environmental interaction (Bourne et al. 2016). In addition, they produce diffusible and non-diffusible

antibacterial compounds that, along with physical exclusion, regulate other bacterial populations influx

(Shnit-Orland and Kushmaro 2009).

Acquisition of symbiotic dinoflagellates and bacteria by corals may occur mainly by horizontal

transmission from the environment, occasionally by vertical transfer to the gametes from the parent

colony, or by a mixture of both (Hernandez-Agreda et al. 2017). An essential element for the uptake

process is the availability of microbial components outside the host (Littman et al. 2009), especially in

environmental reservoirs such as the water column (Takabayashi et al. 2012), benthic macroalgae beds

(Porto et al. 2008) and sediments (Nitschke et al. 2016). Pioneering observations found corallivorous fish

delivering coral symbionts to the environment (Muller-Parker 1984). In the Caribbean, the occurrence of

Symbiodinaceae cells in faeces of the stoplight parrotfish *Sparisoma viride*, with densities up to 38.3×10^3

cells ml^{-1}, has supported this notion (Porto et al. 2008; Castro-Sanguino and Sánchez 2012).

S. viride is the main excavating parrotfish of the Caribbean. This feeding behaviour leads to the ingestion

of symbionts while foraging on rock and live coral (Sanchez et al. 2004; Francini-Filho et al. 2008).

Notably, its primary substratum to feed is the epilithic algal matrix found on dead coral, and live coral is

approximately 1% of the total fish diet (Bonaldo et al. 2014). It is important to remark, which is also well-

documented in the literature, that *S. viride* targets only endolithic material for feeding (Bruggemann et al.

1994; Clements et al. 2016; Nicholson and Clements 2020). The ingestion of photoautotrophs, like

Symbiodinaceae, should be considered as a by-product. Additionally, *S. viride* forages other benthic
species such as sponges (Radax et al. 2012; Burkepile et al. 2019a). After excavating, the grinding action
of *S. viride* pharyngeal jaws and long gut pass transforms calcium carbonate in fine sediment (Perry et al.
2015), which contains living Symbiodinaceae (Castro-Sanguino and Sánchez 2012) and even viable
macroalgae (Vermeij et al. 2013).
74
Coral reefs face unprecedented rates of degradation and increasing stressful scenarios (Hughes et al.
2017). Uncovering processes and mechanisms to enhance ecological resilience is crucial in addressing this
crisis (Chung et al. 2019). Moreover, abundance declining in key species such as *S. viride* (Near
Threatened - IUCN Red List) is threatening the ecological input for local resilience. On this basis, we
hypothesized that the dispersal of coral microbiome components is influenced by *S. viride* faeces
suggesting a coral reef microbiome maintenance mechanism, which is important to maintain the
environmental reservoirs of microbial communities that are key on coral reefs.
82
83 **Materials and Methods**
84
Data and code availability
The qiime2R and R code are available on the Github repository
(https://github.com/smmontanos/Microbiome-analysis-of-Parrotfish.git) with a readme. The procedures
are explained in the section of Methods Details.
Original raw sequences of the bacterial communities associated with the samples of the study were
deposited in NCBI short read sequences (SRA) under the Bioproject PRJNA623459.
91
Study Area
The fieldwork was carried out in the Archipelago of San Andrés, Old Providence, and Santa Catalina
(Southwestern Caribbean), which is composed of oceanic islands, submerged banks, and atolls located
approximately 800 km northwest of Colombia. This territory is part of the Seaflower Biosphere Reserve
declared by UNESCO in 2005 and is considered the largest Marine Protected Area (MPA) in the
Caribbean. Seaflower MPA is relevant at global scale as it contains over 200.000 ha of productive open-
ocean reefs and extended reefs associated with diverse ecosystems (The CaMPAM Mapping Project,
2017), whereby it is regarded as the third largest barrier reef worldwide. Specifically, this study was
developed around San Andrés Island which has 15 km of long barrier reef (The CaMPAM Mapping
Project, 2017). The sampling was conducted at Nirvana Diving Site in 12° 30′ 05″N and 81° 43′ 56″ W
and West Point reef at the site located in the coordinates 12º31´15.45"N, 81º43'48.60"W along the

leeward reef of the island ("Fig. S1"). Sampling was possible thanks to the research permit Nº00254

(2017) issued by Autoridad Nacional de Licencias Ambientales (ANLA), Ministerio de Medio Ambiente

y Desarrollo Sostenible, Colombia

Symbiodinaceae Faeces sampling

Observational samplings of focal male adult individuals (terminal phase) of *S. viride* were conducted

using SCUBA diving in a total area of 5.000 m^2, consisting of patch reefs (depth 3 to 12 m). The

standardized time for focal observations was 10 minutes per individual; preferred individuals were those

foraging near to the bottom due to the consistent quality of the faeces released. During this time, one pellet

faeces sample per individual was collected immediately after released on substrate using 10 ml syringes,

intensely procuring pellet material only. A total of 40 pellet faeces samples were collected. Additionally,

sediment control (seabed) samples were taken (60 cm horizontally to the pellet faeces fall place). Each

syringe was disposed in a Ziploc bag marked with the sample type and safely kept. After the immersion,

samples were fixed in formalin 10 % to preserve cell integrity. This sampling was developed during June

2016. Underwater time corresponded to 24 diving hours.

Faeces filtration

In the laboratory, each sample was filtered using a plastic Büchner funnel coupled with a vacuum pump

and a previously weighted Whatman filter paper grade 4 (25-30um mesh size). Sample scattered on the

filter paper was washed with 0.22 um Filtered Sea Water (FSW) until recovering a final volume of 50 ml.

After, the filter was dried at room temperature, and its new weight with the sample sediment retained on it

was registered. Symbiodinaceae cells size ranged from 5 to 20 um*,* and therefore, it was expected to

recover them the filtrate. Each 50 ml falcon tube was centrifuged at 3100 rpm for 5 min. Subsequently, the

pellet was resuspended into 250 ul of FSW. The difference between the blank filter paper and the dried

sediment sample on filter paper was considered as the collected faeces' weight dry sediment. This process

was applied to the control samples as well. Filtered samples were protected from light incidence due to the

possible autofluorescence degradation of Symbiodinaceae cells.

Microscopy verification and enumeration

Microscopic observation to verify the presence of Symbiodinaceae cells was carried out on an Improved

Neubauer haemocytometer (Hausser Scientific, USA), under a compound light microscope Olympus

CX21, using 40X magnification. Symbiodinaceae recognition was based on previous observations of

tissue samples and zooxanthellae extruded from *Pterogorgia guadalupensis*, and from published

identification photos on scientific papers. Thus, cells were identified using characters as unicellular

circular shape, golden-brownish colour, elongated and reticulated chloroplast and large pyrenoid body

(MR. Nitschke, doctoral dissertation and personal communication, 2015).

Viability quantification by flow cytometry

FCM Standardization. Quantification of viable cells (high fluorescence) and degrading/unviable cells

(low fluorescence) of Symbiodinaceae in *S. viride* faeces was carried out by flow cytometry (FCM). First,

it is important to note that viability was not based on fluorescence intensity only. Although Chl

autofluorescence may prevent the separation between viable and apoptotic cells (Strychar et al. 2004), low

Chl fluorescence is related to stressed or damaged Symbiodinaceae (Gierz et al. 2020), which cell

morphology can be accurately spotted using FCM measurements emitted by particles digitalized as: first,

thanks to the forward scatter (FSC) or the signal related to the cell size and shape; and second, due to side

scatter (SSC) or the signal indicative of cell granularity. Based on FSC and FSC, parameters are specific

to each particle type, FCM identifies cell populations composed of the same particle types. Moreover,

FCM identifies cells populations based on: first, particles light scattering principles, and second the

excitation and emission of fluorescence; both as the response to a light beam. In detail, a cell solution

enters into the sample injection probe, in which there is a stable stream (generally PBS solution) that

forces each particle to pass individually. During the travel, each particle passes through a flow cell, the

site in which is stimulated using laser illumination (488 nm or 633 nm, BD FACs Canto II, BD). As a

result, the particle emits a response expressed in two measures of light scattering that are detected and

converted into electrical signals by a light detector.

For FCM standardization purposes, preliminary assays were developed using two different samples types;

first *Pterogorgia guadalupensis*, a common zooxanthellate octocoral in the study area, extracts from live

tissue, and second, *P. guadalupe*nsis zooxanthellae cultures aliquots. In the beginning, the process

consisted in differentiating cellular components from debris particles based on FSC and SSC. Both

parameters indicated heterogeneous size of cellular components in samples. In the next stage, cellular

components were distinguished using fluorescence emission. For this, propidium iodide (PI) staining was

performed to both types of samples as a way of promoting fluorescence resolution for cell viability

quantifications; however, there were not observed significative differences in fluorescence, in comparison

to unstained samples. Consequently, it was concluded that Symbiodinaceae autofluorescence signal was

pertinent as a viability quantification criterion. In this way, a red fluorescence threshold set (670 nm long-

pass), related to Symbiodinaceae autofluorescence was established after testing the efficiency of three

different PMT (520 nm long-pass, 590 nm long-pass and 670 nm long-pass) in cells differentiation.

Finally, cells populations in faeces and sediment samples were gated and counted using FCS, SSC and

fluorescence parameters between particles. Subsequently, sorting (FACs ARIA II, BD) was carried out on

the individual defined populations. After, each sorted population was observed using an Improved

Neubauer haemocytometer and photographic images were taken. Additionally, fluorescence microscopy

observations were done to corroborate light microscopy findings. Based on the sorted population results,

the FCM settings were considered standardized for the *S.viride* faeces viability quantification.

FCM conditions for viability quantification *S.viride* faeces samples and sediment controls filtered were

analyzed using a BD FACS Canto II cytometer, coupled with the FACS Diva 6.1. Software. A sample

volume of 200 ul was used for each run, and a total of 1500-2000 cell events were registered. FSC and

SSC were collected using a 488nm band-pass filter, with blue light of origin. Additionally,

Symbiodinaceae red autofluorescence signals (chlorophyll a) were collected via 670nm band-pass filter.

As a conclusion, FSC and SSC, in combination with the Symbiodinaceae endogenous autofluorescence

were used to discriminate zooxanthellae population from macroalgae fragments, non-algal cells and high

quantity of debris in *S.viride* faeces. The same parameters functioned for the differentiation of

Symbiodinaceae viable cells and degrading/unviable cells. Normalization of data was made for assuring

an accurate comparison of cell size, fluorescence patterns and viability percentages between populations

in the total samples. Details used for the FCM analysis, under logarithmic criterion, were

(parameter/voltage): FSC/220, SSC/222, FITC/560, PE/500, PE/500 and PE-Cy7/600.

Statistical Analysis for Symbiodinaceae cells

Viability quantification data were analysed for normality. Multifactorial ANOVA was used for examining

significance of Symbiodinaceae viability vs. non-viability in *S. viride* faeces, and in contrast to sediment

controls. Additionally, the Person Correlation Coefficient was calculated for testing the correlation

between healthy and non-healthy Symbiodinaceae populations vs. milligrams of faeces dispersed.

FCM Method

During the standardization process, three cell populations were gated: P1 (red), P2 (green) and P3 (blue)

in a dot plot ("Fig.S2"). Additionally, it was determined an ungated population containing mainly

sediment, which hereinafter will be referred to as P4 (black). According to the populations' distribution,

P2 was identified as the one characterized by higher fluorescence signal emission, size and granular cells

content. Besides, P3 showed a lower fluorescence signal than P2, and granular cells too; P1 presented the

lowest fluorescence signal vs. P2 and P3; and P4 exhibited non-cellular components or debris as it was

previously mentioned. Additionally, fluorescence microscopy observations to sorted populations were

coherent with the previous light microscopy findings. In this sense, P2 contained Symbiodinaceae cells

that presented the highest fluorescence emission; P3 presented zooxanthellae cells with less fluorescence

emission in contrast to P2; and finally, P1 and P4 components emitted the lesser fluorescence ("Fig.S4").

Light and fluorescence observations in sorted populations supported the validation of FCM standardized

parameters. *S. viride* faeces and control sediment samples were observed under light microscopy to

register Symbiodinaceae presence. Zooxanthellae cells were confirmed in 100% of the samples and

controls collected. In all cases, cells were observed coccoid in shape with 11-13 um of diameter. Dividing

cells stages were evidenced in faeces samples. Intracellular composition patterns varied from well-

developed and elongated chloroplasts and defined pyrenoid body to few recognizable organelle's

presence. Additionally, the colour pattern observed corresponded to golden brownish cells to translucent

appearance.

Viability quantification by flow cytometry

Accordingly, to FCM standardized settings, the three gated populations and the ungated population

validated were identified and quantified in all *S. viride* faeces and sediment controls (figure 1 in the main

text). An entire Symbiodinaceae population per sample was defined as the sum of P2 (viable) and P3

(degrading/unviable) events. Thus, 22855 healthy events were registered in P2 vs. 7500 healthy ones in

P3. Regarding faeces or sediment control mg^{-1}, a total of 62 healthy events mg^{-1} corresponding to P2 were

detected, and in contrast, 32 healthy events mg^{-1} in P3 were evidenced. This fact means a proportion of 2

healthy cells events registered in faeces samples to 1 healthy cell event in sediment control. Conversely,

P1 and P4 data were not included in the analyses due to the absence of zooxanthellae in them. In this

context, Symbiodinaceae viability calculation per sample was estimated based on the relation between P2

events and the total cells population frequency, expressed as a percentage. The same calculation was done

for the estimation of the percentage of degrading/unviable cells (see raw data in "Table S1").

Microbiota in faeces sampling

The microbial component was sampled in West point reef. The sampling was developed in August 2019.

Faeces samples were collected in situ through SCUBA diving. Focal *Sparisoma viride* individuals were

followed until they defecated. One sample was collected per individual using a 50 mL syringe

immediately after the deposition and stored in an individual Ziploc bag. Additionally, sediment control

(seabed) samples were collected right next to the faeces deposition. Lastly, three water controls and three

sand channel samples were taken aleatorily within the area of study. Water samples were filtered in a 0.2

microns filter. All samples and filters from water samples were transfer to individual testing tubes with

ethanol at 70 % to preserve bacteria until DNA extractions in laboratory conditions. A total of 12 faeces

samples, 12 sediment samples, three water column samples (controls) and three sand channels samples

(controls) were collected.

DNA extraction and 16S rRNA gene Sequencing

DNA extraction was made using the Quick DNA Fecal/Soil Microbe Kits by Zymo Research according to

manufacture instructions. DNA extractions were sent to the Iowa State University (DNA facility) for

library preparation and sequencing on the Illumina Miseq Platform. The desired region for amplification

was the V4 hypervariable region of the 16S rRNA gene using the primer pairs 515F and 806R(Caporaso et

al. 2011).

16S rRNA based community analysis

Qiime 2 version 2018.4 was used for bioinformatics analysis with different implemented plugins (Bolyen

et al. 2019). Amplicon sequence variants (ASVs) were chosen to work with because they provide better

resolution than OTUs based methods(Callahan et al. 2017). First, DADA2 plugin(Callahan et al. 2016)

was used for quality control and noise reduction (-p-trim-left-r 0 –p-trim-left-r 0 –p-trunc-len-f 238 –p-

trunc-len-r 232). These parameters were chosen according to the reads quality distribution along the length

of the sequence. To assess which microbial taxa were present in the samples, we used to q2-feature-

classifier plug in to classify our ASVs taxonomically. The reference database used was SILVA 132,

specific for 515F and 806R region (http://qiime.org/home_static/dataFiles.html). Next, mitochondria,

chloroplast and eukaryotic sequences were removed and a filter of minimum frequency of 17 for the

ASVs was applied. These filtered sequences were used to construct the phylogenetic tree, which was made

using FastTreeMP with MAFFT alignment implemented in qiime2. The bacterial composition of the

environmental controls, water column and sand channels, as well as of sediments and *S. viride* faeces,

were graphed using the produced table by the script *qiime taxa collapsed* at family level. Subsequently,

the result was graphed using sunburst multilevel pie charts in excel. In order to analyse the diversity of

each treatment (*S. viride* faeces, reef sediments, water column and sand bank), we used the package

phyloseq (McMurdie and Holmes 2013), implemented in R v 3.5. In addition, we examined the

differences between samples richness (Chao1) using ANOVA and Tukey test ("Table S2"). The richness

bloxplots were graphed using the package ggplot2(Wickham 2009). For beta diversity a non-metric

multidimentional scaling (NMDS) analysis was done using Bray Curtis dissimilarities. A PERMANOVA

was performed (999 permutations) in phyloseq to determine differences in the composition of bacterial

communities and PERMADISP was performed (999 permutations) to test of multivariate dispersions.

Finally, we listed corals and sponges in which *S. viride* usually feeds on, and included the bacterial

families associated with those species based on literature data ("Table S3"). This information, along with

our findings of the faecal microbiome of *S. viride,* were used to elaborate the Venn diagram found in

Figure 3.B.

Results and discussion

S. viride faeces enriches reef sediments with viable Symbiodinaceae cells

We confirmed the presence of Symbiodinaceae cells in 100 % of _S. viride_ faeces and sediment controls,

with an average density of 3328 cells ml^{-1} for the former, and 1269 cells ml^{-1} for the latter (See Fig. 1 and

Table S1, Fig. S2-4). In general, viable cells in _S. viride_ faeces corresponded to 25.97 % ± 1.50, whereas

degrading/unviable cells accounted for 74.03 % ± 1.50 (Fig. 1). Similarly, at sediment controls 22.24 % ±

5.36 were viable cells while 77.76 % ± 5.36 corresponded to degrading/unviable cells. The percentages of

viable and degrading/unviable cells were significantly different in _S. viride_ faeces (F=481.2, _p_<0.001).

Contrastingly, no significant differences in the percentage of viable cells were found between _S. viride_

faeces and sediment controls (F=0.1, _p_>0.7; Fig. 1). Noticeable, no correlation was observed between the

percentages obtained in _S. viride_ faeces and control samples vs. dry mass collected (R^2 =0.023, F=1.067,

_p_>0.307), which means that the cell integrity values were not explained by the sample's mass. Based on

these patterns, we propose that _S. viride_ may act as a key vector to enrich reef sediments with

Symbiodinaceae cells. Therefore, we suggest that _S. viride´s_ faeces are crucial for maintaining the main

environmental reservoir of dinoflagellates coral symbionts, in accordance to the critical role of sediments

for the uptake of symbionts by corals (Adams et al. 2009; Nitschke et al. 2016).

S. viride disperses a highly diverse bacterial community in reef sediments

We analysed 16S rRNA to characterize the bacterial community in _S. viride_ faeces and controls. The

sediments and feces showed the highest bacterial diversity followed by controls, sand channels and water

column samples (Fig. 2a). Alpha diversity index demonstrated that richness of bacterial communities was

significantly different between the four types of samples (ANOVA F=9.1, _p_<0.001, Table S2) where

sediments and feces present similar richness (p=0.308) and feces differ from controls, the sand channels

(p=0.03) and water column (p=0.01) samples. The differences in bacterial composition were assessed

using Bray-Curtis dissimilarities in nMDS (Fig. 2b). Particularly, _S. viride_ and sediments grouped together

and were significantly different from controls (PERMANOVA, F=5.14, R2=0.33, _p_=0.001). Moreover,

non-significant differences in the homogeneity of multivariate dispersion were showed between types of

samples (PERMADISP, p >0.05).

Additionally, Proteobacteria was the most abundant phylum in all the samples with Gammaproteobacteria

and Alphaproteobacteria as the dominant classes. This dominance was followed by the phyla

Bacteroidetes, Fusobacteria, Firmicutes and Cyanobacteria in both _S. viride_ faeces and sediments, and

contrastingly, by Firmicutes and Planctomycetes in controls. Regarding the identification of bacterial

families of Gammaproteobacteria, *S. viride* faeces were dominated by Vibrionaceae (42.1 %) and

followed by Rhodobacteraceae (6.33 %), Fusobacteriaceae (5.53 %), Shewanellaceae (4.48 %) and

Woeseiaceae (4.29 %). In contrast, *S. viride* faeces were deficient in Alphaproteobacteria. The

Gammaproteobacteria in sediments belonging to Rhodobacteraceae (14.52 %) and Vibrionaceae (14.36

%) were the most abundant families while Halieaceae (6.31 %) and Woeseiaceae (6.06 %) were less

represented. We also found the family Cyclobacteriaceae from Bacteroidetes phylum (6.97 %) in the

sediments. Although *S. viride* faeces and sediments differed in families abundance, there were similarities

in the overall families identified for both types of samples (Figure 3a). Contrarily, the water column ("Fig.

S5") showed higher abundance of bacteria belonging to Phycisphaeraceae (24.07 %), Sphingomonadaceae

(17.01 %), Pseudohongiellaceae (13.9 %), Lactobacillaceae (12.86 %), Pseudomonadaceae (11.2 %),

while a lesser representation of Vibrionaceae (7.47 %) and, sand channels were dominated by

Desulfobulbaceae (11.48 %), Rhodobacteraceae (11.07 %), Sphingomonadaceae (9.41 %) and

Arenicellaceae (8.58 %), that differed from faeces and sediments. In fact, it was notable the presence of

common amplicon sequence variants (ASVs) across the identified phyla in both faeces and sediments, as

it is showed in the phylogeny of the complete bacterial community (Figure 2.c). On this basis, we

hypothesize that *S. viride* feaces may support the enrichment of bacteria in reef sediments.

***S. viride* faeces contain a bacterial community commonly associated with the microbiome of foraged**

species

S.viride faeces had high abundances of Gammaproteobacteria ASVs, which are usually found in

association with coral tissue (Bourne and Munn 2005)*.* In particular, this bacterial class is dominant in at

least ten coral species excavated by *S.viride* (Sanchez et al. 2004; Rotjan and Lewis 2005; Mumby 2009;

Castro-Sanguino and Sánchez 2012; Bonaldo et al. 2014; Burkepile et al. 2019b)*.* Eight bacterial families

in *S. viride* faeces are associated to coral microbiomes (Fig. 3.b).

Among the Gammaproteobacteria class, the family Rhodobacteraceae , which are prevalent in the

microbiome of *Orbicell*a (Meyer et al. 2016) and *Agaricia* (Gonzalez-Zapata et al. 2018), were identified

in both *S. viride* faeces and sediments. As diazotrophs, Rhodobacteraceae members are involved in the

fixation of nitrogen in association within living coral tissue and the surface mucus layer (Lesser et al.

2018). Similarly, other Gammaproteobacteria metabolise bioproducts from Symbiodinaceae such

dimethylsulfoniopropionate (DMSP) and dimethylsufide (DMS) (Raina et al. 2009). As a result, we

hypothesize that *S. viride* faeces replenish reef sediments with key bacteria supporting the metabolism of

Symbiodinaceae symbionts. This input of Gammaproteobacteria in reefs is relevant in shaping the

bacterial community and producing sulphur based antimicrobial functions (Peixoto et al. 2017).

Vibrionaceae was the most abundant family identified in *S. viride* faeces, and it is commonly found in the

microbiome of several corals and sponges (Dunlap and Pawlik 1996, 1998) (Table S3).

Other families found in *S. viride* faeces were Alteromonadaceae, Flavobacteriaceae and

Pseudoalteromonadaceae, which are common bacterial members associated with *Porites* corals (Table

S3). Similary, Verrucomicrobiaceae, Vibrionaceae and Rhodobacteraceae are bacterial families present

when the mucus is old in *Porites*, probably *S. viride* fed on this coral and disperse through the faeces,

however, it have to tested in further studies. On the other hand, *Agaricia* associates closely to

Vibrionaceae and Rhodobacteraceae (Gonzalez-Zapata et al. 2018), and *Siderastrea, Porites* and

Pseudodiploria maintain beneficial relationships with Alteromonadaceae (Table S3) that also are present

in parrotfishes of the study, suggest that parrotfish could replenish the coral reef with these symbiont

bacteria.

Besides, *Epulopiscium* bacteria was also found in *S. viride* faeces. Although, this result suggests a

potential positive feedback of this bacteria to *S. viride* digestion, it should be interpreted with caution in

parrotfishes. *Epulopiscium* has a major role in surgeonfish supporting the digestion of marine algae

(Ngugi et al. 2017). Parrotfishes are not hindgut fermenters, so the presence of *Epulopiscium* in the *S.*

viride could be ingested from sediments and deserves further study (Clements et al. 2014).

Understanding the ecological feedback of *S. viride´*s microbial dispersal

Our findings suggest that *S. viride* is a vector of symbiotic dinoflagellates, which constantly provides

inputs of viable cells. Moreover, a role in replenishing reef sediments is hypothesize based on identifying

the same ratio of viable cells:unviable cells in both sediments and faeces. We also found similarities in the

bacterial community structure between *S. viride* faeces and reefs sediments. Both of them had diverse

bacterial communities and sharing composition. Likewise, our results suggested that the *S. viride* spreads a

bacterial community commonly associated to the microbiome of species in which the fish feeds on.

Parrotfish release solid pellets of fine sediment than remain for a short time after being defecated (Goatley

and Bellwood 2010). Dispersal occurs in territories of approximately 100 to 300 m^2 (medium sized), and

in a daily feeding range of 50 to 800 m^2 (Bruggemann et al. 1996). The dispersal of Symbiodinaceae cells

and bacteria allows us to start thinking about the resemblance of ecological attributes between *S.*

viride and herbivorous in terrestrial ecosystems. The relevance of enriching reef sediments relies on coral

health and resilience. Sediments enhance the coral uptake of Symbiodinaceae cells (Adams et al. 2009),

which is relevant for coral recruitment and health recovery after bleaching events (Adams et al. 2009; Ali

375 et al. 2019), and facilitate the enrichment of the coral's ecosphere with beneficial microbes and therefore,

376 supports coral physiological responses from microbial exchange processes (Weber et al. 2019).

379 **Figure 1. *S. viride* enriches reef sediments with viable Symbiodinaceae cells**

381 Symbiodinaceae enumeration in *S. viride* faeces and reef sediments. We gated two distinctive populations

382 of Symbiodinaceae cells in both faeces and sediments. Viable cells showed an intact cell membrane with

383 spherical nucleus, well developed and elongated chloroplasts, defined pyrenoid bodies and high

384 fluorescence signal. Conversely, degrading/unviable cells exhibited traits such as a fragmented cell

385 membrane, undefined cytoplasmatic material, and a lower fluorescence. A. Cell density in *S. viride* faeces

386 vs. sediment control. B. Boxplot showing significant differences between the percentages of viable cells

387 and degrading/unviable in *S. viride* faeces and sediment controls. No significant differences were observed

388 in the percentage of viable cells in *S. viride* faeces and sediment controls. C-D- Flow cytometry-FCM dot

389 plots displaying the three gated populations identified during standardization in: C. *S. viride* faeces, and D.

390   Sediment controls. Both types of samples, faeces and sediment controls, presented viable cells (high

391   fluorescence) in P2 and degrading/unviable cells in P3.).

Figure 2. *S. viride* **disperses a highly diverse bacterial community in reef sediments**

A) Microbial community richness in *S. viride* faeces (blue), sediments (light purple), water column (green) and sand channels (pink) assessed with ANOVA (P-value= 0.003), B) Non metrical multidimensional scaling (nMDS) using Bray Curtis dissimilarities of bacterial communities present in the four types of samples. C) Phylogenetic tree of bacterial community was constructed using maximum likelihood with FasTreeMP and MAFFT alignments. The types of samples are indicated with dots (colours correspond to the part A) above each ASVs tips of the tree. Bacterial phyla are observed as regions above each clade of the tree with different colours. In the case of Proteobacteria, the phylogeny shows the phylum divide in three-class taxonomic levels.

Figure 3. *S. viride* **faeces contain a bacterial community commonly associated with the microbiome of foraged species**

A) Bacterial composition at phylum and family level using pie charts. Relative abundances of ASVs classified at family and phylum levels are indicated for two types of control samples which are water column (WC) and sand channels (SC); and in the lower part for the microbiota associated with reefs sediments (S) and *S. viride* faeces (PF). B) Venn diagram showing the shared families between *S. viride* faeces (in the left) found in the study and bacterial families of corals (*Agaricia, Orbicella, Siderastrea,*

Porites, Diploria and *Pseudodiploria* where parrotfish usually feed) based on literature data (See

supplementary table S3) on the right.

Acknowledgments

Thanks to Nacor Bolaños from Corporación para el Desarrollo Sostenible del Archipiélago de San Andrés,

Providencia y Santa Catalina (CORALINA), and Buconos Diving Center team for the assistance during

field surveys; Natalia Bolaños from the Laboratorio de Ciencias Básicas Médicas at the Universidad de los

Andes supported the FCM analysis process. Humberto Ibarra and his team from the Centro de

Microscopía at the Universidad de los Andes contributed with microscopy methods and the members of

Laboratorio de Biología Molecular Marina team (BIOMMAR) at the Universidad de los Andes for their

valuable assistance.

Conflicts of interest

The authors declare that the research was conducted in the absence of any commercial or financial

relationships that could be construed as a potential conflict of interest.

Reference list

Adams LM, Cumbo VR, Takabayashi M (2009) Exposure to sediment enhances primary acquisition of Symbiodinium by asymbiotic coral larvae. Mar Ecol Prog Ser 377:149–156

Ainsworth TD, Krause L, Bridge T, Torda G, Raina J-B, Zakrzewski M, Gates RD, Padilla-Gamiño JL, Spalding HL, Smith C, Woolsey ES, Bourne DG, Bongaerts P, Hoegh-Guldberg O, Leggat W (2015) The coral core microbiome identifies rare bacterial taxa as ubiquitous endosymbionts. ISME J 9:2261–2274

Ali A, Kriefall NG, Emery LE, Kenkel CD, Matz MV, Davies SW (2019) Recruit symbiosis establishment and Symbiodiniaceae composition influenced by adult corals and reef sediment. Coral Reefs 38:405–415

Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H, Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ, Brown CT, Callahan BJ, Caraballo-Rodríguez AM, Chase J, Cope EK, Da Silva R, Diener C, Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M, Estaki M, Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B, Holmes S, Holste H, Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L, Kaehler BD, Kang KB, Keefe CR, Keim P, Kelley ST, Knights D, Koester I, Kosciolek T, Kreps J, Langille MGI, Lee J, Ley R, Liu Y-X, Loftfield E, Lozupone C, Maher M, Marotz C, Martin BD, McDonald D, McIver LJ, Melnik AV, Metcalf JL, Morgan SC, Morton JT, Naimey AT, Navas-Molina JA, Nothias LF, Orchanian SB, Pearson T, Peoples SL, Petras D, Preuss ML, Pruesse E, Rasmussen LB, Rivers A, Robeson MS, Rosenthal P, Segata N, Shaffer M, Shiffer A, Sinha R, Song SJ, Spear JR, Swafford AD, Thompson LR, Torres PJ, Trinh P, Tripathi A, Turnbaugh PJ, Ul-Hasan S, van der Hooft JJJ, Vargas F, Vázquez-Baeza Y, Vogtmann E, von Hippel M, Walters W, Wan Y, Wang M, Warren J, Weber KC, Williamson CHD, Willis AD, Xu ZZ, Zaneveld JR, Zhang Y, Zhu Q, Knight R, Caporaso JG (2019) Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol 37:852–857

Bonaldo RM, Hoey AS, Bellwood DR (2014) The ecosystem roles of parrotfishes on tropical reefs. Oceanogr Mar Biol Annu Rev 52:81–132

Bourne DG, Morrow KM, Webster NS (2016) Insights into the Coral Microbiome: Underpinning the
Health and Resilience of Reef Ecosystems. Annu Rev Microbiol 70:317–340
Bourne DG, Munn CB (2005) Diversity of bacteria associated with the coral Pocillopora damicornis from
the Great Barrier Reef. Environ Microbiol 7:1162–1174
Bruggemann J, van Oppen M, Breeman A (1994) Foraging by the stoplight parrotfish Sparisoma viride. I.
Food selection in different, socially determined habitats. Mar Ecol Prog Ser 106:41–55
Bruggemann JH, Van Kessel AM, Van Rooij JM, Breeman AM (1996) Bioerosion and sediment ingestion
by the Caribbean parrotfish Scarus vetula and Sparisoma viride: implications of fish size, feeding mode
and habitat use. Mar Ecol Prog Ser 134:59–71
Burkepile DE, Adam TC, Roycroft M, Ladd MC, Munsterman KS, Ruttenberg BI (2019a) Species-
specific patterns in corallivory and spongivory among Caribbean parrotfishes. Coral Reefs 38:417–423
Burkepile DE, Adam TC, Roycroft M, Ladd MC, Munsterman KS, Ruttenberg BI (2019b) Species-
specific patterns in corallivory and spongivory among Caribbean parrotfishes. Coral Reefs 38:417–423
Callahan BJ, McMurdie PJ, Holmes SP (2017) Exact sequence variants should replace operational
taxonomic units in marker-gene data analysis. ISME J 11:2639
Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP (2016) DADA2: High-
resolution sample inference from Illumina amplicon data. Nat Methods 13:581–583
Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, Fierer N, Knight R
(2011) Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. Proc Natl
Acad Sci 108:4516–4522
Castro-Sanguino C, Sánchez JA (2012) Dispersal of Symbiodinium by the stoplight parrotfish Sparisoma
viride. Biol Lett 8:282–286
Chung AE, Wedding LM, Green AL, Friedlander AM, Goldberg G, Meadows A, Hixon MA (2019)
Building Coral Reef Resilience Through Spatial Herbivore Management. Front Mar Sci 6:98
Clements KD, Angert ER, Montgomery WL, Choat JH (2014) Intestinal microbiota in fishes: what's
known and what's not. Mol Ecol 23:1891–1898
Clements KD, German DP, Piché J, Tribollet A, Choat JH (2016) Integrating ecological roles and trophic
diversification on coral reefs: multiple lines of evidence identify parrotfishes as microphages. Biol J Linn
Soc
Dunlap M, Pawlik JR (1996) Video-monitored predation by Caribbean reef fishes on an array of
mangrove and reef sponges. Mar Biol 126:117–123
Dunlap M, Pawlik JR (1998) Spongivory by Parrotfish in Florida Mangrove and Reef Habitats. Mar Ecol
19:325–337
Francini-Filho RB, Moura RL, Ferreira CM, Coni EO (2008) Live coral predation by parrotfishes
(Perciformes: Scaridae) in the Abrolhos Bank, eastern Brazil, with comments on the classification of
species into functional groups. Neotropical Ichthyol 6:191–200
Gierz S, Ainsworth TD, Leggat W (2020) Diverse symbiont bleaching responses are evident from 2-
degree heating week bleaching conditions as thermal stress intensifies in coral. Mar Freshw Res 71:1149
Goatley CHR, Bellwood DR (2010) Biologically mediated sediment fluxes on coral reefs: sediment
removal and off-reef transportation by the surgeonfish Ctenochaetus striatus. Mar Ecol Prog Ser 415:237–
245
Gonzalez-Zapata FL, Bongaerts P, Ramírez-Portilla C, Adu-Oppong B, Walljasper G, Reyes A, Sanchez
JA (2018) Holobiont Diversity in a Reef-Building Coral over Its Entire Depth Range in the Mesophotic
Zone. Front Mar Sci 5:
Hernandez-Agreda A, Gates RD, Ainsworth TD (2017) Defining the Core Microbiome in Corals'
Microbial Soup. Trends Microbiol 25:125–140
Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD, Baird AH, Babcock RC,
Beger M, Bellwood DR, Berkelmans R, Bridge TC, Butler IR, Byrne M, Cantin NE, Comeau S, Connolly
SR, Cumming GS, Dalton SJ, Diaz-Pulido G, Eakin CM, Figueira WF, Gilmour JP, Harrison HB, Heron
SF, Hoey AS, Hobbs J-PA, Hoogenboom MO, Kennedy EV, Kuo C, Lough JM, Lowe RJ, Liu G,
McCulloch MT, Malcolm HA, McWilliam MJ, Pandolfi JM, Pears RJ, Pratchett MS, Schoepf V, Simpson

T, Skirving WJ, Sommer B, Torda G, Wachenfeld DR, Willis BL, Wilson SK (2017) Global warming and
recurrent mass bleaching of corals. Nature 543:373–377

Lesser MP, Morrow KM, Pankey SM, Noonan SHC (2018) Diazotroph diversity and nitrogen fixation in
the coral Stylophora pistillata from the Great Barrier Reef. ISME J 12:813–824

Littman RA, Willis BL, Pfeffer C, Bourne DG (2009) Diversities of coral-associated bacteria differ with
location, but not species, for three acroporid corals on the Great Barrier Reef. FEMS Microbiol Ecol
68:152–163

McMurdie PJ, Holmes S (2013) phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. PLOS ONE 8:e61217

Meyer JL, Rodgers JM, Dillard BA, Paul VJ, Teplitski M (2016) Epimicrobiota Associated with the
Decay and Recovery of Orbicella Corals Exhibiting Dark Spot Syndrome. Front Microbiol 7:

Muller-Parker G (1984) Dispersal of Zooxanthellae on Coral Reefs by Predators on Cnidarians. Biol Bull
167:159–167

Mumby PJ (2009) Herbivory versus corallivory: are parrotfish good or bad for Caribbean coral reefs?
Coral Reefs 28:683–690

Ngugi DK, Miyake S, Cahill M, Vinu M, Hackmann TJ, Blom J, Tietbohl MD, Berumen ML, Stingl U
(2017) Genomic diversification of giant enteric symbionts reflects host dietary lifestyles. Proc Natl Acad
Sci 114:E7592–E7601

Nicholson GM, Clements KD (2020) Resolving resource partitioning in parrotfishes (Scarini) using
microhistology of feeding substrata. Coral Reefs 39:1313–1327

Nitschke MR, Davy SK, Ward S (2016) Horizontal transmission of Symbiodinium cells between adult and
juvenile corals is aided by benthic sediment. Coral Reefs 35:335–344

Peixoto RS, Rosado PM, Leite DC de A, Rosado AS, Bourne DG (2017) Beneficial Microorganisms for
Corals (BMC): Proposed Mechanisms for Coral Health and Resilience. Front Microbiol 8:

Perry CT, Kench PS, O'Leary MJ, Morgan KM, Januchowski-Hartley F (2015) Linking reef ecology to
island building: Parrotfish identified as major producers of island-building sediment in the Maldives.
Geology 43:503–506

Porto I, Granados C, Restrepo JC, Sanchez JA (2008) Macroalgal-associated dinoflagellates belonging to
the genus Symbiodinium in Caribbean reefs. PLoS One 3:

Radax R, Rattei T, Lanzen A, Bayer C, Rapp HT, Urich T, Schleper C (2012) Metatranscriptomics of the
marine sponge Geodia barretti: tackling phylogeny and function of its microbial community. Environ
Microbiol 14:1308–1324

Raina J-B, Tapiolas D, Willis BL, Bourne DG (2009) Coral-Associated Bacteria and Their Role in the
Biogeochemical Cycling of Sulfur. Appl Environ Microbiol 75:3492–3501

Ricci F, Marcelino VR, Blackall LL, Kühl M, Medina M, Verbruggen H (2019) Beneath the surface:
community assembly and functions of the coral skeleton microbiome. Microbiome 7:159

Rotjan RD, Lewis SM (2005) Selective predation by parrotfishes on the reef coral Porites astreoides. Mar
Ecol Prog Ser 305:193–201

Sanchez JA, Gil MF, Chasqui LH, Alvarado EM (2004) Grazing dynamics on a Caribbean reef-building
coral. Coral Reefs 23:578–583

Shnit-Orland M, Kushmaro A (2009) Coral mucus-associated bacteria: a possible first line of defense.
FEMS Microbiol Ecol 67:371–380

Strychar KB, Sammarco PW, Piva TJ (2004) Apoptotic and necrotic stages of Symbiodinium
(Dinophyceae) cell death activity: bleaching of soft and scleractinian corals. Phycologia 43:768–777

Takabayashi M, Adams LM, Pochon X, Gates RD (2012) Genetic diversity of free-living Symbiodinium
in surface water and sediment of Hawai'i and Florida. Coral Reefs 31:157–167

Vermeij MJ, van der Heijden RA, Olthuis JG, Marhaver KL, Smith JE, Visser PM (2013) Survival and
dispersal of turf algae and macroalgae consumed by herbivorous coral reef fishes. Oecologia 171:417–425

Weber L, Gonzalez-Díaz P, Armenteros M, Apprill A (2019) The coral ecosphere: A unique coral reef
habitat that fosters coral–microbial interactions. Limnol Oceanogr 64:2373–2388

Wickham H (2009) ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York

561