Endozoochory by parrotfish replenishes reefs with coral-associated microbiome

Trigal M. Velásquez-Rodríguez[1], Catalina Zuluaga-Arias[1], Sandra M. Montaño-Salazar[1,3], John M. González[2] & Juan Armando Sánchez[1]*

[1] Laboratorio de Biología Molecular Marina-Biommar, Departamento de Ciencias Biológicas, Universidad de los Andes, Bogotá, Colombia.

[2] Laboratorio de Ciencias Básicas Médicas, Facultad de Medicina, Universidad de los Andes, Bogotá, Colombia

[3] Facultad de Ciencias, Universidad Nacional de Colombia, Bogotá, Colombia

Contact info

*Correspondence author: juansanc@uniandes.edu.co

Abstract

Parrotfish are a keystone group that reduces competition between algae and corals while shaping the dynamics and resilience of coral reefs. We investigated the structure of disseminated coral-associated bacteria and the extent to which the cell integrity of dinoflagellate photosymbionts (family Symbiodiniaceae) is maintained in the faeces. Similar levels of structural integrity and photosynthetic health of Symbidiodinaceae cells were found in both faeces and reef sediments. Besides, the sediments microbiome echoes the parrotfish faecal microbiome by sharing high diversity and a similar bacterial community composition. These findings uncover the role of excavator parrotfish endozoochory in enriching environmental reservoirs, especially reef sediments, with coral-associated bacteria and photosynthetic microalgae. These processes reinforce the coral microbiome and facilitate coral symbiont acquisition, key features critical to maintaining the health of the most threatened ecosystems on earth. Excavator parrotfish are comparable to herbivorous seed dispersers in terrestrial ecosystems based on the same endozoochorous strategy.

Keywords

Parrotfish, coral microbiome, Symbiodinaceae, endozoochory**,** environmental reservoirs

Introduction

The coral microbiome includes species-specific mutualistic or competitive interactions with unicellular dinoflagellates of the family Symbiodiniaceae, endolithic green algae, bacteria, archaea, fungi, protists and viruses (Ricci et al., 2019). From the surface mucus layer into the skeleton, coral-associated bacterial communities and photosynthetic dinoflagellates support coral homeostasis (Ainsworth et al., 2015). Bacteria occupy specific niches in the coral according to their mutualistic function at the intracellular and

perialgal space (Hernandez-Agreda et al., 2017)*,* where critical roles in catabolizing organic matter and

recycling nutrients are performed (Bourne et al., 2016)*.* Particularly, bacteria located in the coral mucus

supply nutrition while protecting against pathogens and UV light, which is an emerging and dynamic

process from the constant environmental interaction (Bourne et al., 2016)*.* In addition, they produce

diffusible and non-diffusible antibacterial compounds that, along with physical exclusion, regulate other

bacterial populations influx (Shnit-Orland and Kushmaro, 2009)*.*

Acquisition of symbiotic dinoflagellates and bacteria by corals may occur mainly by horizontal

transmission from the environment, occasionally by vertical transfer to the gametes from the parent

colony, or by a mixture of both (Hernandez-Agreda et al., 2017)*.* An essential element for the uptake

process is the availability of microbial components outside the host (Littman et al., 2009), especially in

environmental reservoirs such as the water column (Takabayashi et al., 2012)*,* benthic macroalgae beds

(Porto et al., 2008) and sediments (Nitschke et al., 2016). Pionnering observations found corallivorous fish

delivering coral symbionts to the environment (Parker, 1984). In the Caribbean, the occurrence of

Symbiodinaceae cells in faeces of the stoplight parrotfish *Sparisoma viride*, with densities up to 38.3×10^{3}

cells ml^{-1}, has supported this notion (Castro-Sanguino and Sánchez, 2012a; Porto et al., 2008).

S. viride is the main excavating parrotfish of the Caribbean*.* This feeding behaviour leads to the ingestion

of symbionts while foraging on rock and live coral (Francini-Filho et al., 2008; Sanchez et al., 2004)*.*

Notably, its primary substratum to feed is the epilithic algal matrix found on dead coral, and live coral is

approximately 1% of the total fish diet (Bonaldo et al., 2014)*.* Additionally, *S. viride* forages other benthic

species such as sponges (Burkepile et al., 2019a; Radax et al., 2012). After excavating, the grinding

action of *S. viride* pharyngeal jaws and long gut pass transforms calcium carbonate in fine sediment

(Perry et al., 2015), which contains living Symbiodinaceae (Castro-Sanguino and Sánchez, 2012a) and

even viable macroalgae (Vermeij et al., 2013)*.*

Coral reefs face unprecedented rates of degradation and increasing stressful scenarios (Hughes et al.,

2017). Uncovering processes and mechanisms to enhance ecological resilience is crucial in addressing

this crisis (Chung et al., 2019). Moreover, abundance declining in key species such as *S. viride* (Near

Threatened - IUCN Red List) is threating the ecological input for local resilience. On this basis, we

investigated the dispersal of coral microbiome components in *S. viride* faeces*.*

Results and discussion

***S. viride* endozoochory enriches reef sediments with viable Symbiodinaceae cells**

We confirmed the presence of Symbiodinaceae cells in 100% of *S. viride* faeces and sediment controls,

with an average density of 3328 cells ml^{-1} for the former, and 1269 cells ml^{-1} for the latter (See Fig. 1 and

Table S1, Fig. S2-4). In general, viable cells in *S. viride* faeces corresponded to 25.97%±1.50, whereas

degrading/unviable cells accounted for 74,03%±1,50 (Fig. 1). Remarkably similar, at sediment controls

22,24%±5,36 were viable cells while 77,76%±5,36 corresponded to degrading/unviable cells. The

percentages of viable and degrading/unviable cells were significantly different in *S. viride* faeces (F=481,2,

p<0,001). This pattern was also observed in sediment controls. Contrastingly, no significant differences in

the percentage of viable cells were found between *S. viride* faeces and sediment controls (F=0,1, *p*>0,7;

Fig. 1). Noticeable, no correlation was observed between the percentages obtained in *S. viride* faeces and

control samples vs. dry mass collected (R^2 =0,023, F=1,067, *p*>0,307), which means that the cell integrity

values were not explained by the sample's mass. Based on these patterns, we propose that *S. viride* may

act as a key vector to enrich or "fertilize" reef sediments with Symbiodinaceae cells. Therefore, we

suggest *S. viride* endozoochory for maintaining the main environmental reservoir of dinoflagellates coral

symbionts, in accordance to the critical role of sediments for the uptake of symbionts by corals (Adams et

al., 2009; Nitschke et al., 2016).

S. viride disperses a highly diverse bacterial community in reef sediments

S. viride faeces showed the highest bacterial diversity along with sediments as compared to the sand

channels, and it was least in the water column (Fig. 2a). Alpha diversity index demonstrated that richness

of bacterial communities was significantly different between the four types of samples (ANOVA F=9.1,

p<0.001, Table S2). The differences in bacterial composition were assessed using Bray-Curtis

dissimilarities in nMDS (Fig. 2b). Particularly, *S. viride* and sediments grouped together and were

significantly different from controls (PERMANOVA, F=5.14, R2=0.33, *p*=0.001). Additionally,

Proteobacteria was the most abundant phylum in all the samples with Gammaproteobacteria and

Alphaproteobacteria as the dominant classes. This dominance was followed by the phylum Bacteroidetes,

Fusobacteria, Firmicutes and Cyanobacteria in both *S. viride* faeces and sediments, and contrastingly, by

Firmicutes and Planctomycetes in controls.

S. viride faeces contain a bacterial community commonly associated with the microbiome of

foraged species

S.viride faeces had high abundances of Gammaproteobacteria ASVs, which are usually found in

association with coral tissue (Bourne and Munn, 2005)*.* In particular, this bacterial class is dominant in at

least ten coral species excavated by *S.viride* (Bonaldo et al., 2014; Burkepile et al., 2019b; Castro-

Sanguino and Sánchez, 2012a; Mumby, 2009; Rotjan and Lewis, 2005; Sanchez et al., 2004)*.* In the

figure 3.b, we recognized eight bacterial families in *S. viride* faeces that are associated to coral

microbiomes.

Among the Gammaproteobacteria class, the family Rhodobacteraceae , which are prevalent in the

microbiome of *Orbicell*a (Meyer et al., 2016) and *Agaricia* (Gonzalez-Zapata et al., 2018), were identified

in both *S. viride* faeces and sediments. As diazotrophs, Rhodobacteraceae members are involved in the

fixation of nitrogen in association within living coral tissue and the surface mucus layer (Lesser et al.,

2018). Similarly, other Gammaproteobacteria metabolise bioproducts from Symbiodinaceae such

dimethylsulfoniopropionate (DMSP) and dimethylsufide (DMS) (Raina et al., 2009). Therefore, we

highlight that *S. viride* endozoochory replenishes reef sediments with key bacteria to support the

metabolism of Symbiodinaceae symbionts. This input of Gammaproteobacteria in reefs is relevant in

shaping the bacterial community and producing sulphur based antimicrobial functions for the host (Peixoto

et al., 2017). Vibrionaceae was the most abundant family identified in *S. viride* faeces, and it is commonly

found in the microbiome of several coral genera and sponges (Dunlap and Pawlik, 1998, 1996) (see Table

S3).

Understanding the ecological feedback of *S. viride* endozoochory

We analysed faecal pellets of *S. viride* to understand how the endozoochory of coral microbial

components operate. Our findings showed that *S. viride* is a vector of symbiotic dinoflagellates which

constantly provides inputs of viable cells in reefs. Moreover, a role in replenishing reef sediments is

suggested based on identifying the same ratio of viable cells:unviable cells in both sediments and faeces.

We also found similarities in the bacterial community structure between *S. viride* faeces and reefs

sediments. Both of them have a highly diverse bacterial community and share a similar bacterial

composition. Likewise, our results demonstrated that the *S. viride* spreads a bacterial community

commonly associated with the microbiome of species in which the fish feeds on.

Parrotfish release solid pellets of fine sediment than remain for a short time after being defecated (Goatley

and Bellwood, 2010). Dispersal occurs in territories of approximately 100 to 300 m^2 (medium sized), and in

a daily feeding range of 50 to 800 m^2 (Bruggemann et al., 1996). The endozoochory of Symbiodinaceae

cells and bacteria gives *S. viride* comparable ecological attributes to herbivorous seed dispersers in

terrestrial ecosystems. Endozoochory is the primary seed dispersal strategy in forest trees (Guy Haim et

al., 2017). The relevance of enriching reef sediments relies on coral health and resilience. Sediments

enhance the coral uptake of Symbiodinaceae cells (Adams et al., 2009), which is relevant for coral

recruitment and health recovery after bleaching events (Adams et al., 2009; Ali et al., 2019), and facilitate

the enrichment of the coral's ecosphere with beneficial microbes and therefore, supports coral

physiological responses from microbial exchange processes (Weber et al., 2019).

Declaration of Interests

The authors declare no competing interests.

Figure 1. _S. viride_ endozoochory enriches reef sediments with viable Symbiodinaceae cells

Symbiodinaceae enumeration in _S. viride_ faeces and reef sediments. We gated two distinctive populations of Symbiodinaceae cells in both faeces and sediments. Viable cells showed an intact cell membrane with spherical nucleus, well developed and elongated chloroplasts, defined pyrenoid bodies and high fluorescence signal. Conversely, degrading/unviable cells exhibited traits such as a fragmented cell membrane, undefined cytoplasmatic material, and a lower fluorescence. A. Cell density in _S. viride_ faeces vs. sediment control. B. Boxplot showing significant differences between the percentages of viable cells and degrading/unviable in _S. viride_ faeces and sediment controls. No significant differences were observed in the percentage of viable cells in _S. viride_ faeces and sediment controls. C-D- Flow cytometry-FCM dot plots displaying the three gated populations identified during standardization in: C. _S. viride_ faeces, and D. Sediment controls. Both types of samples, faeces and sediment controls, presented viable cells (high fluorescence) in P2 and degrading/unviable cells in P3.).

Figure 2. *S. viride* disperses a highly diverse bacterial community in reef sediments

A) Microbial community richness in *S. viride* faeces (blue), sediments (light purple), water column (green) and sand channels (pink) assessed with ANOVA (P-value= 0,003), B) Non metrical multidimensional scaling (nMDS) using Bray Curtis dissimilarities of bacterial communities present in the four types of samples. C) Phylogenetic tree of bacterial community was constructed using maximum-likelihood with FasTreeMP and MAFFT alignments. The types of samples are indicated with dots (colours correspond to the part A) above each ASVs tips of the tree. Bacterial phyla are observed as regions above each clade of the tree with different colours. In the case of Proteobacteria, the phylogeny shows the phylum divide in three-class taxonomic levels.

Figure 3. *S. viride* **faeces contain a bacterial community commonly associated with the microbiome of foraged species**

A) Bacterial composition at phylum and family level using pie charts. Relative abundances of ASVs classified at family and phylum levels are indicated for two types of control samples which are water column (WC) and sand channels (SC); and in the lower part for the microbiota associated with reefs sediments (S) and *S. viride* faeces (PF). B) Venn diagram showing the shared families between *S. viride* faeces, corals and sponges based on results from this study and literature data.

190 **Methods**
191
Data and code availability
The qiime2R and R code are available on the Github repository
(https://github.com/smmontanos/Microbiome-analysis-of-Parrotfish.git) with a readme. The procedures are
explained in the section of Methods Details.
Original raw sequences of the bacterial communities associated with the samples of the study were
deposited in NCBI short read sequences (SRA) under the Bioproject PRJNA623459.
Study Area
The fieldwork was carried out in the Archipelago of San Andrés, Old Providence, and Santa Catalina
(Southwestern Caribbean), which is composed of oceanic islands, submerged banks, and atolls located
approximately 800 km northwest of Colombia. This territory is part of the Seaflower Biosphere Reserve
declared by UNESCO in 2005 and is considered the largest Marine Protected Area (MPA) in the
Caribbean. Seaflower MPA is relevant at global scale as it contains over 200.000 ha of productive open-
ocean reefs and extended reefs associated with diverse ecosystems (The CaMPAM Mapping Project,
2017), whereby it is regarded as the third largest barrier reef worldwide. Specifically, this study was
developed around San Andrés Island which has 15 km of long barrier reef (The CaMPAM Mapping
Project, 2017). The sampling was conducted at Nirvana Diving Site in 12° 30′ 05″N and 81° 43′ 56″ W and
West Point reef at the site located in the coordinates 12º31´15.45"N, 81º43'48.60"W along the leeward
reef of the island ("Fig. S1"). Sampling was possible thanks to the research permit Nº00254 (2017) issued
by Autoridad Nacional de Licencias Ambientales (ANLA), Ministerio de Medio Ambiente y Desarrollo
Sostenible, Colombia
Symbiodinaceae Faeces sampling
Observational samplings of focal male adult individuals (terminal phase) of *S. viride* were conducted using
SCUBA diving in a total area of $5.000m^2$, consisting of patch reefs (depth 3 to 12m). The standardized
time for focal observations was 10 minutes per individual; preferred individuals were those foraging near
to the bottom due to the consistent quality of the faeces released. During this time, one pellet faeces
sample per individual was collected immediately after released on substrate using 10 ml syringes. A total
of 40 pellet faeces samples were collected. Additionally, sediment control samples were taken (60 cm
horizontally to the pellet faeces fall place). Each syringe was disposed in a Ziploc bag marked with the
sample type and safely kept. Generally, pellet faeces were collected on sediment bottom; and
occasionally on live and dead coral surfaces. After the immersion, samples were fixed in formalin 10% to
preserve cell integrity. This sampling was developed during June 2016. Underwater time corresponded to
24 diving hours.
Faeces filtration
In the laboratory, each sample was filtered using a plastic Büchner funnel coupled with a vacuum pump
and a previously weighted Whatman filter paper grade 4 (25-30um mesh size). Sample scattered on the
filter paper was washed with 0.22 um Filtered Sea Water (FSW) until recovering a final volume of 50 ml.

After, the filter was dried at room temperature, and its new weight with the sample sediment retained on it

was registered. Symbiodinaceae cells size ranged from 5 to 20 um, and therefore, it was expected to

recover them the filtrate. Each 50 ml falcon tube was centrifuged at 3100 rpm for 5 min. Subsequently, the

pellet was resuspended into 250 ul of FSW. The difference between the blank filter paper and the dried

sediment sample on filter paper was considered as the collected faeces' weight dry sediment. This

process was applied to the control samples as well. Filtered samples were protected from light incidence

due to the possible autofluorescence degradation of Symbiodinaceae cells.

Microscopy verification and enumeration

Microscopic observation to verify the presence of Symbiodinaceae cells was carried out on an Improved

Neubauer haemocytometer (Hausser Scientific, USA), under a compound light microscope Olympus

CX21, using 40X magnification. *Symbiodinaceae* recognition was based on previous observations of

tissue samples and zooxanthellae extruded from *Pterogorgia guadalupensis*, and from published

identification photos on scientific papers. Thus, cells were identified using characters as unicellular circular

shape, golden-brownish colour, elongated and reticulated chloroplast and large pyrenoid body (MR.

Nitschke, personal communication, 2015).

Viability quantification by flow cytometry

FCM Standardization Quantification of viable cells (high fluorescence) and degrading/unviable cells (low

fluorescence) of Symbiodinaceae in *S. viride* faeces was carried out by flow cytometry (FCM). FCM

identifies cells populations based on: first, particles light scattering principles, and second the excitation

and emission of fluorescence; both as the response to a light beam. In detail, a cell solution enters into the

sample injection probe, in which there is a stable stream (generally PBS solution) that forces each particle

to pass individually. During the travel, each particle passes through a flow cell, the site in which is

stimulated using laser illumination (488nm or 633nm, BD FACs Canto II, BD). As a result, the particle

emits a response expressed in two measures of light scattering that are detected and converted into

electrical signals by a light detector. These measurements emitted by particles are digitalized as: first, the

forward scatter (FSC) or the signal related to the cell size and shape; and second, the side scatter (SSC)

or the signal indicative of cell granularity. Based on FSC and FSC parameters are specific to each particle

type, FCM identifies cell populations composed of the same particle types.

Additionally, FCM use the fluorescence emission as an identification criterion with an added fluorochrome

or autofluorescence. After the laser beam crosses particles, it excites their fluorochromes and light is

emitted at different wavelengths. These wavelengths are split using optical filters in specific colors and

detected by optical detectors known as photomultiplier tubes (PMT). As a result, the PMT generate an

electrical pulse converted into readable signals by computer software. Those signals are expressed as

cell events of a population in a dot plot.

For FCM standardization purposes, preliminary assays were developed using two different samples types;

first *Pterogorgia guadalupensis*, a common zooxanthellate octocoral in the study area, extracts from live

tissue, and second, *P. guadalupe*nsis zooxanthellae cultures aliquots. In the beginning, the process

consisted in differentiating cellular components from debris particles based on FSC and SSC. Both

parameters indicated heterogeneous size of cellular components in samples. In the next stage, cellular

components were distinguished using fluorescence emission. For this, propidium iodide (PI) staining was

performed to both types of samples as a way of promoting fluorescence resolution for cell viability

quantifications; however, there were not observed significative differences in fluorescence, in comparison

to unstained samples. Consequently, it was concluded that Symbiodinaceae autofluorescence signal was

pertinent as a viability quantification criterion. In this way, a red fluorescence threshold set (670nm long-

pass), related to Symbiodinaceae autofluorescence was established after testing the efficiency of three

different PMT (520nm long-pass, 590nm long-pass and 670nm long-pass) in cells differentiation. Finally,

cells populations in faeces and sediment samples were gated and counted using FCS, SSC and

fluorescence parameters between particles. Subsequently, sorting (FACs ARIA II, BD) was carried out on

the individual defined populations. After, each sorted population was observed using an Improved

Neubauer haemocytometer and photographic images were taken. Additionally, fluorescence microscopy

observations were done to corroborate light microscopy findings. Based on the sorted population results,

the FCM settings were considered standardized for the *S. viride* faeces viability quantification.

FCM conditions for viability quantification *S. viride* faeces samples and sediment controls filtered were

analyzed using a BD FACS Canto II cytometer, coupled with the FACS Diva 6.1. Software. A sample

volume of 200 ul was used for each run, and a total of 1500-2000 cell events were registered. FSC and

SSC were collected using a 488nm band-pass filter, with blue light of origin. Additionally, Symbiodinaceae

red autofluorescence signals (chlorophyll a) were collected via 670nm band-pass filter. As a conclusion,

FSC and SSC, in combination with the *Symbiodinaceae* endogenous autofluorescence were used to

discriminate zooxanthellae population from macroalgae fragments, non-algal cells and high quantity of

debris in *S. viride* faeces. The same parameters functioned for the differentiation of Symbiodinaceae viable

cells and degrading/unviable cells. Normalization of data was made for assuring an accurate comparison

of cell size, fluorescence patterns and viability percentages between populations in the total samples.

Details used for the FCM analysis, under logarithmic criterion, were (parameter/voltage): FSC/220,

SSC/222, FITC/560, PE/500, PE/500 and PE-Cy7/600.

Statistical Analysis for Symbiodinaceae cells

Viability quantification data were analysed for normality. Multifactorial ANOVA was used for examining

significance of Symbiodinaceae viability vs. non-viability in *S. viride* faeces, and in contrast to sediment

controls. Additionally, the Person Correlation Coefficient was calculated for testing the correlation between

healthy and non-healthy Symbiodinaceae populations vs. milligrams of faeces dispersed.

FCM Method

During the standardization process, three cell populations were gated: P1 (red), P2 (green) and P3 (blue)

in a dot plot ("Fig.S2"). Additionally, it was determined an ungated population containing mainly sediment,

which hereinafter will be referred to as P4 (black). According to the populations' distribution, P2 was

identified as the one characterized by higher fluorescence signal emission, size and granular cells

content. Besides, P3 showed a lower fluorescence signal than P2, and granular cells too; P1 presented

the lowest fluorescence signal vs. P2 and P3; and P4 exhibited non-cellular components or debris as it

was previously mentioned. Additionally, fluorescence microscopy observations to sorted populations were coherent with the previous light microscopy findings. In this sense, P2 contained *Symbiodinaceae* cells that presented the highest fluorescence emission; P3 presented zooxanthellae cells with less fluorescence emission in contrast to P2; and finally, P1 and P4 components emitted the lesser fluorescence ("Fig.S4"). Light and fluorescence observations in sorted populations supported the validation of FCM standardized parameters. *S. viride* faeces and control sediment samples were observed under light microscopy to register *Symbiodinaceae* presence. Zooxanthellae cells were confirmed in 100% of the samples and controls collected. In all cases, cells were observed coccoid in shape with 11-13 um of diameter. Dividing cells stages were evidenced in faeces samples. Intracellular composition patterns varied from well-developed and elongated chloroplasts and defined pyrenoid body to few recognizable organelle's presence. Additionally, the colour pattern observed corresponded to golden brownish cells to translucent appearance.

Viability quantification by flow cytometry

Accordingly, to FCM standardized settings, the three gated populations and the ungated population validated were identified and quantified in all *S. viride* faeces and sediment controls (figure 1 in the main text). An entire Symbiodinaceae population per sample was defined as the sum of P2 (viable) and P3 (degrading/unviable) events. Thus, 22855 healthy events were registered in P2 vs. 7500 healthy ones in P3. Regarding faeces or sediment control mg^{-1}, a total of 62 healthy events mg^{-1} corresponding to P2 were detected, and in contrast, 32 healthy events mg^{-1} in P3 were evidenced. This fact means a proportion of 2 healthy cells events registered in faeces samples to 1 healthy cell event in sediment control. Conversely, P1 and P4 data were not included in the analyses due to the absence of zooxanthellae in them. In this context, *Symbiodinaceae* viability calculation per sample was estimated based on the relation between P2 events and the total cells population frequency, expressed as a percentage. The same calculation was done for the estimation of the percentage of degrading/unviable cells (see raw data in "Table S1").

Microbiota in faeces sampling

The fieldwork was carried out in San Andrés island at West Point reef at the site located in the following coordinates 12º31´15.45"N, 81º43'48.60"W. The sampling was developed in August 2019. Faeces samples were collected in situ through SCUBA diving. The reef's depth was approximately 5-7 meters. Focal *Sparisoma viride* individuals where followed until they defecated. Individuals foraging near the bottom where preferred because it facilitated the recollection of the complete faeces deposited by the fish. We followed a method similar to (Castro-Sanguino and Sánchez, 2012b). One sample was collected per individual using a 50 mL syringe immediately after the deposition and stored in an individual Ziploc bag marked with the respective number sample. Additionally, sediment control samples were collected right next to the faeces deposition. Lastly, three water controls and three sand channel samples were taken aleatorily within the area of study. Water samples where filtered in a 0,2 microns filter. All samples and filters from water samples were transfer to individual testing tubes with ethanol at 70% to preserve

bacteria until DNA extractions in laboratory conditions. A total of 12 faeces samples, 12 sediment

samples, three water column samples (controls) and three sand channels samples (controls) were

collected.

DNA extraction and 16S rRNA gene Sequencing

DNA extraction was made using the Quick DNA Fecal/Soil Microbe Kits by Zymo Research according to

manufacture instructions. DNA extractions were sent to the Iowa State University (DNA facility) for library

preparation and sequencing on the Illumina Miseq Platform. The desired region for amplification was the

V4 hypervariable region of the 16S rRNA gene using the primer pairs 515F and 806R(Caporaso et al.,

2011).

16S rRNA based community analysis

Quiime 2 version 2018.4 was used for bioinformatics analysis with different implemented plugins(Bolyen

et al., 2019). Amplicon sequence variants (ASVs) were chosen to work with because they provide better

resolution than OTUs based methods(Callahan et al., 2017). First, DADA2 plugin(Callahan et al., 2016)

was used for quality control and noise reduction (-p-trim-left-r 0 –p-trim-left-r 0 –p-trunc-len-f 238 –p-trunc-

len-r 232). These parameters were chosen according to the reads quality distribution along the length of

the sequence. To asses which microbial taxa were present in the samples, we used to q2-feature-

classifier plug in to classify our ASVs taxonomically. The reference database used was SILVA 132,

specific for 515F and 806R region (http://qiime.org/home_static/dataFiles.html). Next, mitochondria,

chloroplast and eukaryotic sequences were removed and a filter of minimum frequency of 17 for the ASVs

was applied. These filtered sequences were used to construct the phylogenetic tree, which was made

using FastTreeMP with MAFFT alignment implemented in qiime2. The bacterial composition of the

environmental controls, water column and sand channels, as well as of sediments and *S. viride* faeces,

were graphed using the produced table by the script *qiime taxa collapsed* at family level. Subsequently,

the result was graphed using sunburst multilevel pie charts in excel. In order to analyse the diversity of

each treatment (*S. viride* faeces, reef sediments, water column and sand bank), we used the package

phyloseq (McMurdie and Holmes, 2013), implemented in R v 3.5. In addition, we examined the differences

between samples richness (Chao1) using ANOVA and Tukey test ("Table S2"). The richness bloxplots

were graphed using the package ggplot2(Wickham, 2009). For beta diversity a non-metric

multidimentional scaling (NMDS) analysis was done using Bray Curtis dissimilarities. A PERMANOVA was

performed to determine differences in the composition of bacterial communities. Finally, we listed corals

and sponges in which *S. viride* usually feeds on, and included the bacterial families associated with those

species based on literature data ("Table S3"). This information, along with our findings of the faecal

microbiome of *S. viride,* were used to elaborate the Venn diagram found in Figure 3.B.

Regarding the identification of bacterial families of Gammaproteobacteria, *S. viride* faeces were

dominated by Vibrionaceae (42,1%) and followed by Rhodobacteraceae (6,33%), Fusobacteriaceae

(5,53%), Shewanellaceae (4,48%) and Woeseiaceae (4,29%). In contrast, *S. viride* faeces were deficient

in Alphaproteobacteria. The Gammaproteobacteria in sediments belonging to Rhodobacteraceae

(14,52%) and Vibrionaceae (14,36%) were the most abundant families while Halieaceae (6,31%) and
Woeseiaceae (6,06%) were less represented. We also found the family Cyclobacteriaceae from
Bacteroidetes phylum (6,97%) in the sediments. It is worth noting that although *S. viride* faeces and
sediments differed in families' abundance, there were similarities in the overall families identified for both
types of samples (Figure 3a). Contrarily, the water column ("Fig. S5") showed higher abundance of
bacteria belonging to Phycisphaeraceae (24,07%), Sphingomonadaceae (17,01%), Pseudohongiellaceae
(13,9%), Lactobacillaceae (12,86%), Pseudomonadaceae (11,2%), while a lesser representation of
Vibrionaceae (7,47%) and, sand channels were dominated by Desulfobulbaceae (11,48%),
Rhodobacteraceae (11,07%), Sphingomonadaceae (9,41%) and Arenicellaceae (8,58%), that differed
from faeces and sediments. In fact, it was notable the presence of common amplicon sequence variants
(ASVs) across the identified phyla in both faeces and sediments, as it is showed in the phylogeny of the
complete bacterial community (Figure 2.c). On this basis, we hypothesize that *S. viride* endozoochory may
support the enrichment of bacteria in reef sediments.
393
Other families found in *S. viride* faeces were Alteromonadaceae, Flavobacteriaceae and
Pseudoalteromonadaceae, which are common bacterial members associated with *Porites* corals (Table
S3). *Porites* exhibits cyclical mucus aging and shedding that may follow lunar periodicity (Coffroth, 1991;
Glasl et al., 2016), where new mucus has a high abundance of *Endozoicomonas* while old mucus, ready
to be shed, presents a high abundance of bacteria belonging to the Verrucomicrobiaceae and
Vibrionaceae families; contrastingly, there is an increase in Rhodobacteraceae when the mucus starts
turning old (Glasl et al., 2016). Because aged mucus remains up to 3 days before being released, it is
possible that *S. viride* was feeding on the old mucus of this coral, which should be tested in further
studies. On the other hand, *Agaricia* associates closely to Vibrionaceae and Rhodobacteraceae
(Gonzalez-Zapata et al., 2018), and *Siderastrea, Porites* and *Pseudodiploria* maintain beneficial
relationships with Alteromonadaceae (Table S3). Besides, *Epulopiscium* bacteria was also found in *S.
viride* faeces. This result suggests a potential positive feedback of this bacteria to *S. viride* digestion. This
fact is based on the role of *Epulopiscium* as a symbiont coevolving with surgeonfish that supports the
digestion of marine algae (Ngugi et al., 2017). On this basis, *Epulopiscium* could develop a similar
function in the *S. viride* gut microbiome.
409
410
411
412 **References**
413

Adams LM, Cumbo VR, Takabayashi M. 2009. Exposure to sediment enhances primary
acquisition of Symbiodinium by asymbiotic coral larvae. *Mar Ecol Prog Ser* **377**:149–156.
doi:10.3354/meps07834

Ainsworth TD, Krause L, Bridge T, Torda G, Raina J-B, Zakrzewski M, Gates RD, Padilla-
Gamiño JL, Spalding HL, Smith C, Woolsey ES, Bourne DG, Bongaerts P, Hoegh-Guldberg
O, Leggat W. 2015. The coral core microbiome identifies rare bacterial taxa as ubiquitous
endosymbionts. *ISME J* **9**:2261–2274. doi:10.1038/ismej.2015.39

Alex A, Antunes A. 2015. Pyrosequencing characterization of the microbiota from Atlantic
intertidal marine sponges reveals high microbial diversity and the lack of co-occurrence
patterns. *PLoS One* **10**.

Ali A, Kriefall NG, Emery LE, Kenkel CD, Matz MV, Davies SW. 2019. Recruit symbiosis
establishment and Symbiodiniaceae composition influenced by adult corals and reef
sediment. *Coral Reefs* **38**:405–415. doi:10.1007/s00338-019-01790-z

Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H, Alm
EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ,
Brown CT, Callahan BJ, Caraballo-Rodríguez AM, Chase J, Cope EK, Da Silva R, Diener C,
Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M, Estaki M,
Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann
B, Holmes S, Holste H, Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L,
Kaehler BD, Kang KB, Keefe CR, Keim P, Kelley ST, Knights D, Koester I, Kosciolek T,
Kreps J, Langille MGI, Lee J, Ley R, Liu Y-X, Loftfield E, Lozupone C, Maher M, Marotz
C, Martin BD, McDonald D, McIver LJ, Melnik AV, Metcalf JL, Morgan SC, Morton JT,
Naimey AT, Navas-Molina JA, Nothias LF, Orchanian SB, Pearson T, Peoples SL, Petras D,
Preuss ML, Pruesse E, Rasmussen LB, Rivers A, Robeson MS, Rosenthal P, Segata N,
Shaffer M, Shiffer A, Sinha R, Song SJ, Spear JR, Swafford AD, Thompson LR, Torres PJ,
Trinh P, Tripathi A, Turnbaugh PJ, Ul-Hasan S, van der Hooft JJJ, Vargas F, Vázquez-Baeza
Y, Vogtmann E, von Hippel M, Walters W, Wan Y, Wang M, Warren J, Weber KC,
Williamson CHD, Willis AD, Xu ZZ, Zaneveld JR, Zhang Y, Zhu Q, Knight R, Caporaso JG.
2019. Reproducible, interactive, scalable and extensible microbiome data science using
QIIME 2. *Nat Biotechnol* **37**:852–857. doi:10.1038/s41587-019-0209-9

Bonaldo RM, Hoey AS, Bellwood DR. 2014. The ecosystem roles of parrotfishes on tropical
reefs. *Oceanogr Mar Biol Annu Rev* **52**:81–132.

Bonthond G, Merselis DG, Dougan KE, Graff T, Todd W, Fourqurean JW, Rodriguez-Lanetty
M. 2018. Inter-domain microbial diversity within the coral holobiont Siderastrea siderea from
two depth habitats. *PeerJ* **6**:e4323. doi:10.7717/peerj.4323

Bourne DG, Morrow KM, Webster NS. 2016. Insights into the Coral Microbiome: Underpinning
the Health and Resilience of Reef Ecosystems. *Annu Rev Microbiol* **70**:317–340.
doi:10.1146/annurev-micro-102215-095440

Bourne DG, Munn CB. 2005. Diversity of bacteria associated with the coral Pocillopora
damicornis from the Great Barrier Reef. *Environ Microbiol* **7**:1162–1174.
doi:10.1111/j.1462-2920.2005.00793.x

Bruggemann JH, Van Kessel AM, Van Rooij JM, Breeman AM. 1996. Bioerosion and sediment
ingestion by the Caribbean parrotfish Scarus vetula and Sparisoma viride: implications of fish
size, feeding mode and habitat use. *Mar Ecol Prog Ser* **134**:59–71.

Burkepile DE, Adam TC, Roycroft M, Ladd MC, Munsterman KS, Ruttenberg BI. 2019a.
Species-specific patterns in corallivory and spongivory among Caribbean parrotfishes. *Coral*
Reefs **38**:417–423. doi:10.1007/s00338-019-01808-6

Burkepile DE, Adam TC, Roycroft M, Ladd MC, Munsterman KS, Ruttenberg BI. 2019b.
Species-specific patterns in corallivory and spongivory among Caribbean parrotfishes. *Coral*
Reefs **38**:417–423. doi:10.1007/s00338-019-01808-6

Callahan BJ, McMurdie PJ, Holmes SP. 2017. Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. *ISME J* **11**:2639.

Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016. DADA2:
High-resolution sample inference from Illumina amplicon data. *Nat Methods* **13**:581–583.
doi:10.1038/nmeth.3869

Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, Fierer N,
Knight R. 2011. Global patterns of 16S rRNA diversity at a depth of millions of sequences
per sample. *Proc Natl Acad Sci* **108**:4516–4522. doi:10.1073/pnas.1000080107

Cárdenas A, Rodriguez-R LM, Pizarro V, Cadavid LF, Arévalo-Ferro C. 2012. Shifts in bacterial
communities of two caribbean reef-building coral species affected by white plague disease.
ISME J **6**:502–512. doi:10.1038/ismej.2011.123

Castro-Sanguino C, Sánchez JA. 2012a. Dispersal of Symbiodinium by the stoplight parrotfish
Sparisoma viride. *Biol Lett* **8**:282–286.

Castro-Sanguino C, Sánchez JA. 2012b. Dispersal of Symbiodinium by the stoplight parrotfish
Sparisoma viride. *Biol Lett* **8**:282–286.

Chung AE, Wedding LM, Green AL, Friedlander AM, Goldberg G, Meadows A, Hixon MA.
2019. Building Coral Reef Resilience Through Spatial Herbivore Management. *Front Mar*
Sci **6**:98. doi:10.3389/fmars.2019.00098

Coffroth MA. 1991. Cyclical mucous sheet formation on poritid corals in the San Blas Islands,
Panama. *Mar Biol* **109**:35–40. doi:10.1007/BF01320229

Daniels C, Baumgarten S, Yum LK, MIchell CT, Bayer T, Arif C, Roder C, Weil E, Voolstra
CR. 2015. Metatranscriptome analysis of the reef-building coral Orbicella faveolata indicates
holobiont response to coral disease. *Front Mar Sci* **2**. doi:10.3389/fmars.2015.00062

Dunlap M, Pawlik JR. 1998. Spongivory by Parrotfish in Florida Mangrove and Reef Habitats.
Mar Ecol **19**:325–337. doi:10.1111/j.1439-0485.1998.tb00471.x

Dunlap M, Pawlik JR. 1996. Video-monitored predation by Caribbean reef fishes on an array of
mangrove and reef sponges. *Mar Biol* **126**:117–123. doi:10.1007/BF00571383

Dunphy CM, Gouhier TC, Chu ND, Vollmer SV. 2019. Structure and stability of the coral
microbiome in space and time. *Sci Rep* **9**:1–13. doi:10.1038/s41598-019-43268-6

Ezzat L, Lamy T, Maher RL, Munsterman KS, Landfield KM, Schmeltzer ER, Clements CS,
Vega Thurber RL, Burkepile DE. 2020. Parrotfish predation drives distinct microbial
communities in reef-building corals. *Anim Microbiome* **2**:5. doi:10.1186/s42523-020-0024-0

Fiore CL, Labrie M, Jarett JK, Lesser MP. 2015. Transcriptional activity of the giant barrel
sponge, Xestospongia muta Holobiont: molecular evidence for metabolic interchange. *Front*
Microbiol **6**. doi:10.3389/fmicb.2015.00364

Frade PR, Roll K, Bergauer K, Herndl GJ. 2016. Archaeal and Bacterial Communities Associated
with the Surface Mucus of Caribbean Corals Differ in Their Degree of Host Specificity and
Community Turnover Over Reefs. *PLoS ONE* **11**. doi:10.1371/journal.pone.0144702

Francini-Filho RB, Moura RL, Ferreira CM, Coni EO. 2008. Live coral predation by parrotfishes
(Perciformes: Scaridae) in the Abrolhos Bank, eastern Brazil, with comments on the
classification of species into functional groups. *Neotropical Ichthyol* **6**:191–200.

Glasl B, Herndl GJ, Frade PR. 2016. The microbiome of coral surface mucus has a key role in
mediating holobiont health and survival upon disturbance. *ISME J* **10**:2280–2292.
doi:10.1038/ismej.2016.9

Goatley CHR, Bellwood DR. 2010. Biologically mediated sediment fluxes on coral reefs:
sediment removal and off-reef transportation by the surgeonfish Ctenochaetus striatus. *Mar*
Ecol Prog Ser **415**:237–245. doi:http://dx.doi.org/10.3354/meps08761

Gonzalez-Zapata FL, Bongaerts P, Ramírez-Portilla C, Adu-Oppong B, Walljasper G, Reyes A,
Sanchez JA. 2018. Holobiont Diversity in a Reef-Building Coral over Its Entire Depth Range
in the Mesophotic Zone. *Front Mar Sci* **5**. doi:10.3389/fmars.2018.00029

Guy□Haim T, Hyams□Kaphzan O, Yeruham E, Almogi□Labin A, Carlton JT. 2017. A novel
marine bioinvasion vector: Ichthyochory, live passage through fish. *Limnol Oceanogr Lett*
2:81–90. doi:10.1002/lol2.10039

Hernandez-Agreda A, Gates RD, Ainsworth TD. 2017. Defining the Core Microbiome in Corals'
Microbial Soup. *Trends Microbiol* **25**:125–140. doi:10.1016/j.tim.2016.11.003

Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD, Baird AH,
Babcock RC, Beger M, Bellwood DR, Berkelmans R, Bridge TC, Butler IR, Byrne M, Cantin
NE, Comeau S, Connolly SR, Cumming GS, Dalton SJ, Diaz-Pulido G, Eakin CM, Figueira
WF, Gilmour JP, Harrison HB, Heron SF, Hoey AS, Hobbs J-PA, Hoogenboom MO,
Kennedy EV, Kuo C, Lough JM, Lowe RJ, Liu G, McCulloch MT, Malcolm HA, McWilliam
MJ, Pandolfi JM, Pears RJ, Pratchett MS, Schoepf V, Simpson T, Skirving WJ, Sommer B,

Torda G, Wachenfeld DR, Willis BL, Wilson SK. 2017. Global warming and recurrent mass
bleaching of corals. *Nature* **543**:373–377. doi:10.1038/nature21707

Kellogg CA, Piceno YM, Tom LM, DeSantis TZ, Gray MA, Andersen GL. 2014. Comparing
Bacterial Community Composition of Healthy and Dark Spot-Affected Siderastrea siderea in
Florida and the Caribbean. *PLoS ONE* **9**. doi:10.1371/journal.pone.0108767

Lesser MP, Morrow KM, Pankey SM, Noonan SHC. 2018. Diazotroph diversity and nitrogen
fixation in the coral Stylophora pistillata from the Great Barrier Reef. *ISME J* **12**:813–824.
doi:10.1038/s41396-017-0008-6

Littman RA, Willis BL, Pfeffer C, Bourne DG. 2009. Diversities of coral-associated bacteria
differ with location, but not species, for three acroporid corals on the Great Barrier Reef.
FEMS Microbiol Ecol **68**:152–163.

McMurdie PJ, Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive Analysis
and Graphics of Microbiome Census Data. *PLOS ONE* **8**:e61217.
doi:10.1371/journal.pone.0061217

Meyer JL, Rodgers JM, Dillard BA, Paul VJ, Teplitski M. 2016. Epimicrobiota Associated with
the Decay and Recovery of Orbicella Corals Exhibiting Dark Spot Syndrome. *Front
Microbiol* **7**. doi:10.3389/fmicb.2016.00893

Morrow KM, Moss AG, Chadwick NE, Liles MR. 2012. Bacterial Associates of Two Caribbean
Coral Species Reveal Species-Specific Distribution and Geographic Variability. *Appl
Environ Microbiol* **78**:6438–6449. doi:10.1128/AEM.01162-12

Mumby PJ. 2009. Herbivory versus corallivory: are parrotfish good or bad for Caribbean coral
reefs? *Coral Reefs* **28**:683–690.

Ngugi DK, Miyake S, Cahill M, Vinu M, Hackmann TJ, Blom J, Tietbohl MD, Berumen ML,
Stingl U. 2017. Genomic diversification of giant enteric symbionts reflects host dietary
lifestyles. *Proc Natl Acad Sci* **114**:E7592–E7601. doi:10.1073/pnas.1703070114

Nitschke MR, Davy SK, Ward S. 2016. Horizontal transmission of Symbiodinium cells between
adult and juvenile corals is aided by benthic sediment. *Coral Reefs* **35**:335–344.

Parker GM. 1984. Dispersal of zooxanthellae on coral reefs by predators on cnidarians. *Biol Bull*
167:159–167. doi:10.2307/1541344

Peixoto RS, Rosado PM, Leite DC de A, Rosado AS, Bourne DG. 2017. Beneficial
Microorganisms for Corals (BMC): Proposed Mechanisms for Coral Health and Resilience.
Front Microbiol **8**. doi:10.3389/fmicb.2017.00341

Perry CT, Kench PS, O'Leary MJ, Morgan KM, Januchowski-Hartley F. 2015. Linking reef
ecology to island building: Parrotfish identified as major producers of island-building
sediment in the Maldives. *Geology* **43**:503–506.

Pita L, Fraune S, Hentschel U. 2016. Emerging Sponge Models of Animal-Microbe Symbioses.
Front Microbiol **7**. doi:10.3389/fmicb.2016.02102

Porto I, Granados C, Restrepo JC, Sanchez JA. 2008. Macroalgal-associated dinoflagellates
belonging to the genus Symbiodinium in Caribbean reefs. *PLoS One* **3**.

Pratte ZA, Richardson LL. 2018. Microbiome dynamics of two differentially resilient corals. *Dis*
Aquat Organ **131**:213–226. doi:10.3354/dao03289

Radax R, Rattei T, Lanzen A, Bayer C, Rapp HT, Urich T, Schleper C. 2012.
Metatranscriptomics of the marine sponge Geodia barretti: tackling phylogeny and function
of its microbial community. *Environ Microbiol* **14**:1308–1324. doi:10.1111/j.1462-
2920.2012.02714.x

Raina J-B, Tapiolas D, Willis BL, Bourne DG. 2009. Coral-Associated Bacteria and Their Role
in the Biogeochemical Cycling of Sulfur. *Appl Environ Microbiol* **75**:3492–3501.
doi:10.1128/AEM.02567-08

Ricci F, Marcelino VR, Blackall LL, Kühl M, Medina M, Verbruggen H. 2019. Beneath the
surface: community assembly and functions of the coral skeleton microbiome. *Microbiome*
7:159.

Rosales SM, Sinigalliano C, Gidley M, Jones PR, Gramer LJ. 2019. Oceanographic habitat and
the coral microbiomes of urban-impacted reefs. *PeerJ* **7**:e7552. doi:10.7717/peerj.7552

Rotjan RD, Lewis SM. 2005. Selective predation by parrotfishes on the reef coral Porites
astreoides. *Mar Ecol Prog Ser* **305**:193–201. doi:10.3354/meps305193

Sanchez JA, Gil MF, Chasqui LH, Alvarado EM. 2004. Grazing dynamics on a Caribbean reef-
building coral. *Coral Reefs* **23**:578–583.

Shnit-Orland M, Kushmaro A. 2009. Coral mucus-associated bacteria: a possible first line of
defense. *FEMS Microbiol Ecol* **67**:371–380.

Takabayashi M, Adams LM, Pochon X, Gates RD. 2012. Genetic diversity of free-living
Symbiodinium in surface water and sediment of Hawai 'i and Florida. *Coral Reefs* **31**:157–
167.

Vermeij MJ, van der Heijden RA, Olthuis JG, Marhaver KL, Smith JE, Visser PM. 2013.
Survival and dispersal of turf algae and macroalgae consumed by herbivorous coral reef
fishes. *Oecologia* **171**:417–425.

Weber L, Gonzalez☐Díaz P, Armenteros M, Apprill A. 2019. The coral ecosphere: A unique
coral reef habitat that fosters coral–microbial interactions. *Limnol Oceanogr* **64**:2373–2388.
doi:10.1002/lno.11190

Wichels A, Würtz S, Döpke H, Schütt C, Gerdts G. 2006. Bacterial diversity in the breadcrumb
sponge Halichondria panicea (Pallas). *FEMS Microbiol Ecol* **56**:102–118.
doi:10.1111/j.1574-6941.2006.00067.x

Wickham H. 2009. ggplot2: Elegant Graphics for Data Analysis, Use R! New York: Springer-
Verlag.

598