Title: Alpha fluctuations regulate the accrual of visual information to awareness

Abbreviated title: Alpha-regulates information accrual to awareness

Authors

Mireia Torralba[1], Alice Drew[1], Alba Sabaté San José[2], Luis Morís Fernández[1], and Salvador Soto-Faraco[1,3]

[1]Multisensory Research Group, Center for Brain and Cognition, University of Pompeu Fabra, Barcelona,

Spain, ZIP: 08002.

[2]Universitat de Barcelona, Barcelona, Spain, ZIP: 08007.

[3]Catalan Institution for Research and Advanced Studies (ICREA), Barcelona, Spain, ZIP: 08010.

Corresponding author: Mireia Torralba Cuello (mireia.torralba@upf.edu), ZIP: 08002

Competing interest

The authors declare no competing financial interests.

Acknowledgments

This research was supported by the Ministerio de Economia y Competitividad (PSI2016-75558-P

AEI/FEDER), AGAUR Generalitat de Catalunya (2017 SGR 1545). This project has been co-funded with

50% by the European Regional Development Fund under the framework of the FEDER Operative

Programme for Catalunya 2014-2020, with a grant of 1,527,637.88€. We want to thank Dr. Márta Szabina

Pápai and Indre Pileckyte for helping in the piloting of the Experiments and in EEG data collection.

Abstract:

Endogenous brain processes play a paramount role in shaping up perceptual phenomenology, as illustrated by the alternations experienced by humans (and other animals) when watching perceptually ambiguous, static images. Here, we hypothesised that endogenous alpha fluctuations in the visual cortex pace the accumulation of sensory information leading to perceptual outcomes. We addressed this hypothesis using binocular rivalry combined with visual entrainment and electroencephalography in humans (42 female, 40 male). The results revealed a correlation between the individual frequency of alpha oscillations in the occipital cortex and perceptual alternation rates experienced during binocular rivalry. In subsequent experiments we show that regulating endogenous brain activity via entrainment produced corresponding changes in perceptual alternation rate, which were observed only in the alpha range but not at lower entrainment frequencies. Overall, rhythmic alpha stimulation resulted in faster perceptual alternation rates, compared to arrhythmic or no stimulation. These findings support the notion that visual information is accumulated via alpha cycles to promote the emergence of conscious perceptual representations. We suggest that models of binocular rivalry incorporating posterior alpha as a pacemaker can provide an important advance in the comprehension of the dynamics of visual awareness.

Key words

Binocular rivalry, sensory entrainment, alpha activity, EEG, individual variability, discrete perception.

Significance statement:

Mainstream theories in cognitive neuroscience agree that endogenous brain processes play a paramount role in shaping our perceptual experience of sensory inputs. In vision, endogenous fluctuations in the alpha rhythm have been pointed out to regulate visual inputs to perception. In support of this hypothesis, here we used EEG recordings and visual entrainment to demonstrate that inter-individual differences in the speed of endogenous alpha fluctuations in the brain are causally related to the accrual of visual information to awareness. These findings provide, for the first time, evidence for alpha-gated regulation of the dynamics of alternations in conscious visual perception.

 Introduction

There is ample agreement that our perception of the world is not built from incoming sensory input alone,

but is also determined by endogenous brain processes (Friston, 2012). The impact of these endogenous

processes is well illustrated by spontaneous fluctuations in conscious perception when watching ambiguous

images, a phenomenon known as bistable perception (Sterzer et al., 2009). Ambiguous images lead

observers to experience alternations between different perceptual interpretations of an otherwise static input

and have been used by neuroscientists to investigate the neural correlates of fluctuations in perceptual

awareness. Here, we address the hypothesis that alpha oscillatory activity in the occipital cortex plays a

causal role in these perceptual fluctuations.

We focused on binocular rivalry (BR), a phenomenon that arises when two dissimilar images are presented

to different eyes. BR has been characterised as an intrinsically dynamic process resulting from competition

and mutual inhibition between different neural populations representing the rival images (Mamassian and

Goutcher, 2005). There is still controversy about the specific brain areas involved in this competition

(Leopold and Logothetis, 1999; Block, 2019; Phillips and Morales, 2020), however, evidence suggests that

the occipital cortex is involved in resolving perceptual ambiguity (Polonsky et al., 2000; Lee et al., 2007;

de Jong et al., 2016). In some models, BR competition is not solely based on mutual inhibition but also on

attentional modulation (Li et al., 2017).

BR is characterised by the natural alternation rate (NAR) between dominance periods, a stochastic process

strongly dependant on properties of the rival stimuli (Sobel and Blake, 2002; Brascamp et al., 2005; Kang,

2009). Interestingly, the NAR is relatively stable within individuals and genetically hereditary (Miller et

al., 2010). Additionally, although inter-individual variability in BR dynamics is large (Kleinschmidt et al.,

2012), alternation rates within an individual correlate across different forms of bistable perception

(Schmack et al., 2013) and stimuli (Carter and Pettigrew, 2003).

Recently, several studies have linked BR with brain oscillatory activity in different frequency bands

(Doesburg et al., 2005; Piantoni et al., 2010), and in particular to occipital alpha activity (8-14 Hz). Alpha

oscillations are strongly related to attention (Davidson et al., 2018; Katyal et al., 2019) and, more relevantly,

alpha activity in the occipital cortex is known to be related to rhythmic fluctuations in visual perception

(Vanrullen et al., 2011). For example, the gating by inhibition hypothesis (Jensen and Mazaheri, 2010)

contends that the alpha rhythm imposes a discrete sampling of sensory input via an alternation between

moments of heightened neural excitability and bouts of pulsed inhibition, cycling every 100 ms

approximately. In line with this hypothesis, variations in the speed of alpha oscillations correlate with inter-

individual variability in the temporal resolution of visual perception (Samaha and Postle, 2015) and

processing speed in general (Clark et al., 2007). The results of entrainment studies support the causal role

of alpha oscillations in modulating perception (Cecere et al., 2015; Ronconi et al., 2018). An interesting

property of the cortical alpha rhythm is that the individual alpha frequency (IAF) reflects systemic

properties of the brain, is highly heritable, and relates to cognitive functioning (Grandy et al., 2013).

In this study, we focus on establishing a connection between alpha endogenous activity speed and typical

BR alternation times. We propose that cortical alpha oscillations play a key role in the competition between

alternative representations during BR. In particular, we hypothesise that the alpha rhythm provides phasic

windows for the cyclical re-evaluation of the competing stimuli representations and, consequentially,

evidence accumulation takes place in a phasic manner. This discrete, alpha gated, evidence accumulation

eventually leads to perceptual switches. According to this hypothesis, faster alpha oscillations will correlate

positively with the NAR in a BR task and visual entrainment will causally modulate BR dynamics in a

frequency-specific way.

Methods

Subjects

Participants received 10€/ hour in return for their participation. They all provided written informed consent

prior to the study and were naive to the purpose of the experiment. The study was run in accordance with

the Declaration of Helsinki and the experimental protocol approved by the local ethics committee CIEC

Parc de Salut Mar (Universitat Pompeu Fabra, Barcelona, Spain).

Subject inclusion criteria in all the experiments were: normal or corrected to normal vision, not being under

medication and displaying a single peak in the alpha range during the eyes-closed pre-screening EEG

recording (see *IAF estimation*). In order to minimise alternations unrelated to endogenous activity (Kanai

et al., 2005), we excluded participants with a high percentage of EEG artefacts such as blinks, eye

movements or motor contamination in general (with a high percentage defined as more than 5% in

Experiment 1 and 10% in Experiments 2 and 3, applied separately in each of the 4 experimental conditions).

For Experiments 2 and 3, due to the reduced availability of data points per condition (5 blocks per condition compared to 9 in Experiment 1) we established an additional criterion to ensure a reliable estimation of the NAR (Supplementary materials and methods: *Inclusion criteria*): In Experiments 2 and 3 we included only subjects with 50 or more percepts in each experimental condition which, according to simulation, ensured a deviation in median estimation <3.5%. Please note that in Experiment 1, all subjects had far more than 50 percepts (a percept is the period of time from seeing one pure colour to seeing the next pure colour).

Thirty-one participants (15 female, aged 18-34, mean age 23) participated in Experiment 1. Four participants were excluded for having more than 5% of the segmented epochs contaminated by visual and movement artefacts (final sample N=27). Twenty-nine participants (16 female, aged 18-31, mean age 24) participated in Experiment 2. One participant was excluded for presenting a double peak in the alpha range, two participants were discarded due to excessive artefacts and one participant was discarded due to insufficient time for estimation of the NAR (final sample N=25). Thirty-four participants (19 female, aged 19-32, mean age 24) participated in Experiment 3. One participant was excluded for presenting a double peak in the alpha range, two participants were discarded due to an insufficient number of percepts and one participant was discarded due to a technical problem with the recordings (final sample N=30).

Apparatus and Stimuli

Participants were presented with one Gabor patch to each eye, 11.5° in diameter, Michelson contrast 0.1, opposed orientations (±45° off vertical) and different colours (red, RGB= [1 0 0] and green, RGB adjusted individually to match the luminance of the red Gabor patch). Both patches were surrounded by a black annulus and contained a central fixation cross identical in each of the patches (Fig. 1). The images were presented on a 19.8-inch CRT monitor (120 Hz refresh rate) with a grey background (10.7 cd/m^2) placed 80 cm from participants' eyes. Each image was presented to a different eye by means of stereoscope mirror glasses. Visual stimuli were created and presented using the Matlab toolbox Psychtoolbox (RRID: SCR_002881), version 3.

Visual stimulation for entrainment, when used, consisted of pulsed Gaussian increases in Michelson contrast from 0.1 to 0.6, of 58 ms duration, presented simultaneously to the images displayed to the left and

right eyes. Different entrainment frequencies were then achieved by varying the offset to onset time

between pulses, whilst maintaining the slope of contrast increases constant (Fig. 1).

Figure 1: A: Example of the Gabor patches used in the experimental protocol. B: Contrast of both

Gabor patches was modulated simultaneously with different temporal patterns (low and high

frequency rhythmic modulation, fast arrhythmic modulation). C: Participants continuously reported

the conscious percept (red, green, mixed or none) while EEG was concurrently recorded. D: Image

of our experimental setup. E: Target behavioural measure was individual characteristic duration of

conscious percept (NAR). F: Behavioural measures were complemented with endogenous oscillatory

activity recorded during the BR task.

Procedure

The procedure was identical for all three experiments (Supplementary Fig. 1), with the only variation

between experiments being the visual entrainment stimulation conditions (see below) and number of blocks

performed.

The experimental session started with the participant's IAF estimation during a 5-minute rest block, with

eyes closed. We then ran a protocol to match the two rival stimuli in terms of luminance for subjective

equality (Cavanagh et al., 1987) (Supplementary materials and methods: *Objective isoluminance*

adjustment). Next, the stereoscope mirrors were calibrated for the rival stimuli to appear at the same retinal

location of each eye. Finally, two training blocks (120 s each) on the BR task were performed prior to the

experimental BR runs.

All experiments contained the same rival images and were organised in 120 s blocks of a BR task. The eye

of presentation of the red and green stimuli was alternated from block to block. After each block, there was

a pause and participants were given the chance to rest before starting the next trial. The experimental

conditions used in each individual experiment were characterised in terms of the visual stimulation used:

• No entrainment (0 Hz).

• Rhythmic entrainment: periodic contrast increments with a frequency of presentation of 3 Hz,

IAF, IAF-2Hz and IAF+2HZ, depending on the condition.

• Arrhythmic visual stimulation (arrIAF): non-periodic contrast increments. In this condition,

the latencies between contrast increments were drawn from an exponential distribution centred at

subjects' 1/IAF in order to match the number of contrast increments during the corresponding

rhythmic IAF condition of the same subject.

In Experiment 1, subjects ran 9 blocks of the 0 Hz condition. In experiments 2 and 3 subjects ran 5 blocks

of each experimental condition: 3 Hz, IAF-2Hz, IAF and IAF+2Hz for Experiment 2 and 0 Hz, 3 Hz, IAF

and arrIAF for Experiment 3. (Supplementary Figure S1). The order of conditions was intermixed pseudo-

randomly between blocks (Supplementary Figure S1).

During the task, participants were instructed to fixate their gaze on the central fixation cross and report their

current percept (red, green, mixed or null) by key press with the left and right index of each hand. Response

mapping for the green stimulus and the red stimulus was counter-balanced between participants. To report

a mixed (piecemeal) percept, participants were instructed to press both keys simultaneously and to release

both keys if none of the three aforementioned percepts were apparent (e.g., null percept).

Predictions

In this study, we performed a series of three experiments, each of them designed to test specific predictions

derived from our hypothesis. The predictions and associated experiments are as follows.

NAR correlates positively with IAF. If alpha endogenous activity paces information accrual, one first

expected result is that faster individual alpha oscillations will lead the individual to experience a faster

natural alternation rate (NAR) in a BR task (Fig. 2). Experiment 1 was designed to test this prediction: EEG

endogenous activity was concurrently recorded while participants performed a BR task. The experiment

was pre-registered (https://osf.io/n5c8g/) before analysing any data. Recently, this prediction has been

independently supported in a study showing a correlation between NAR and IAF (Katyal et al., 2019).

Figure 2: Binocular rivalry can be represented as a double well potential, each of the wells (here

named R and G, for Red and Green respectively) represent the excitability of the competing neural

populations (the larger the excitability, the deepest the well). The grey disk represents the neural

population with stronger evidence. A: In some models, neural noise triggers perceptual switches, therefore the percepts could happen at any point in time. B: Other models predict that the excitability of the neural population sensitive to current conscious percept decreases over time, due to adaptation. As a result, as time decreases the probability of a switch increases, but the percept durations predicted by this model are highly irregular. C: At present, it is acknowledged that both noise and adaptation play a role in BR, therefore predicting stochastic percept durations (due to noise) and low probability of very long percept durations (due to adaptation). D: Top, sketch of a possible time course of the competition between neural populations during a BR task, as time evolves both noise and adaptation affect the instantaneous competition process. Middle: Alpha activity periodically samples the outcome of the competition process, and, as result, the conscious percept (colored line) is determined over time. Bottom: A slower alpha rhythm samples competition process more slowly and, as a result, perceptual switches are more spaced in time.

Alpha activity is causal to BR dynamics. Furthermore, our hypothesis implies that alpha oscillatory activity is causal to BR dynamics. Therefore, we anticipated that entraining relevant endogenous alpha activity with rhythmic visual stimulation should cause changes in perceptual alternation rate. Based on this, our second prediction is that entraining brain activity slightly below the individual's alpha rhythm will result in the individual experiencing slower BR alternation rates, whereas entraining brain activity slightly above the alpha rhythm will speed up alternation rates. Experiment 2 was designed to test this second prediction: we measured the NAR in a BR protocol while the observers' brain activity was entrained at 3Hz, IAF-2Hz, IAF or IAF+2Hz.

The effect of entrainment on BR dynamics is frequency specific and limited to periodic stimulation. One further implication of the hypothesis is frequency specificity. That is, only oscillatory frequencies neighbouring the alpha range are relevant to the re-evaluation of competing stimuli, and therefore entraining at rhythms outside the alpha range will not affect BR dynamics. Following this frequency specificity assumption, we predict that entraining brain activity at a frequency far from the alpha range (e.g., 3 Hz) will not impact competition dynamics and, therefore, the alternation rates under non-alpha rhythmic entrainment will be equivalent to those without visual entrainment. Finally, according to our hypothesis, the periodicity of neural fluctuations is consequential for BR dynamics. That is, modulations of BR dynamics imposed by entrainment depend on the temporal (rhythmic) structure of the stimulation and

cannot be explained solely by accumulated changes in the amount of energy (i.e. increases in competing

stimuli strength). Following this hypothesis, we assume that entraining with rhythmic (alpha) activity will

promote periodic sampling of the competing stimuli, compared to comparable amounts of stimulation

without a rhythmic structure. Hence, our fourth main prediction is that the NAR in BR, observed under

arrhythmic stimulation, will be slower than the NAR observed under rhythmic (alpha) entrainment. A final

prediction, following from the same hypothesis, is that entraining alpha activity might increase the NAR as

compared to no-entrainment. This is because alpha entraining can sustain alpha activity over and above its

naturally occurring ongoing dynamics, which is typically characterised by intermittent bouts (Rusiniak et

al., 2018). Experiment 3 was designed to test these two predictions: we measured the NAR while observers

were entrained with either 3Hz, rhythmic (IAF), comparable arrhythmic trains, or not entrained at all. This

experiment was registered prior to any data analysis. At the time of the registration the data of 6 subjects

had been collected (https://osf.io/w8m4q/) but none had been analysed.

It should be emphasised that the aim of this study, outlined in these hypotheses, is merely to probe the role

of alpha oscillations in rivalry dynamics. We make no claim as to the underlying cause of these fluctuations,

which is a question beyond the ambitions of the present study. Regardless of their cause, perceptual

fluctuations are a phenomenological effect of BR attributed to competition dynamics, and we set out to

investigate the role of alpha oscillations in this visual competition. The mechanisms underlying perceptual

fluctuations during BR have been addressed in a parallel study currently in preparation (Drew et al., 2019).

Sample size estimation

The sample size for each of the three experiments was calculated to attain a 95% statistical power in the

comparison of interest: IAF-NAR correlation in Experiment 1, IAF-2Hz compared to IAF+2Hz in

Experiment 2, and IAF compared to arrhythmic stimulation in Experiment 3 (Supplementary materials and

methods: *Sample Size estimation*).

Behavioural analysis

Percepts shorter than 300 ms (estimated latency of the motor evoked potential from key presses (Halgren,

1990)) were filtered out from the analysis. The NAR was obtained as the inverse of the median time between

reporting two different pure percepts (complete transitions from red to green or green to red) (Fig. 1).

EEG recording

EEG data were acquired using active electrodes (actiCAP, BrainProducts GmbH, Munich, Germany) with Brain Vision Recorder (BrainProducts GmbH, Munich, Germany) at a sampling rate of 500 Hz. The ground electrode was placed at AFz and online reference on the tip of the nose. Four external electrodes were used: left and right mastoid for offline re-referencing and horizontal and vertical electrooculogram for monitoring eye movements and blinks. Impedance was kept below 10 kΩ. In Experiment 1, 64 electrodes placed according to the 10-10 system were used for the recording. In Experiments 2 and 3, we recorded from a subset of 21 electrodes. The ROI (parieto-occipital electrodes), defined a priori, consisted of a cluster of 8 electrodes: PO9, PO3, POz, PO4, PO10, O1, Oz, O2 in Experiment 1, PO7, PO3, POz, PO4, PO8, O1, Oz, O2, in Experiments 2 and 3.

EEG preprocessing and analyses

Pre-processing. EEG preprocessing and analysis was performed using Matlab's toolbox Fieldtrip (RRID: SCR_004849) and custom-written code. The EEG data was band-pass filtered 1-70 Hz (Butterworth filter, order 2, two-pass); a notch filter was used to remove the 50 Hz line-noise. The data recorded for IAF calculation with eyes closed was processed without further analysis. The EEG data recorded during task performance was further visually inspected to mark visual or muscular artefacts. For the IAF measurement during the task (Experiments 1 and 3), data fragments containing artefacts were discarded from the analysis; for the evoked activity measurement (Experiments 2 and 3), it was not (please note that only participants with a very low percentage of artefacts (<10%) were included in the final sample). All analyses were performed in the parieto-occipital region of interest only.

IAF estimation. The IAF at rest was measured from continuous EEG recordings (300 s) using the Welch method (1 second segments, 10% overlap, 0.25 to 40 Hz in steps of 0.25 Hz). The power spectrum was averaged across the electrodes of interest (see ROI above) and background noise was estimated by fitting a $1/f^{\alpha}$ curve to the power spectrum (Haegens et al., 2014). IAF at rest was used only as a screening: participants presenting a single peak in the spectrum exceeding background noise in the 5 to 15 Hz range were accepted to proceed with the experiment. The calculation of IAF during task was performed in the 0 Hz (no entrainment) condition (Experiments 1 and 3). Data were segmented into epochs according to participants' reports (green or red percept, between key presses to avoid motor contamination). Pre-

processed epochs were subsequently divided into non-overlapping segments of 1 s duration , demeaned and

zero-padded to 10 s (Haegens et al., 2014). The power spectrum was calculated using a STFT (Hanning

window, 4 to 30 Hz in steps of 0.1 Hz) and background noise correction using a $1/f^{\alpha}$ fit was performed as

in the estimation of IAF at rest (see above).

SSVEP and phase-locking measures. In order to determine that endogenous activity was indeed entrained

during our experiments we performed several tests: we calculated the power spectrum of steady-state visual

evoked potentials (SSVEP) for all the rhythmic conditions (Experiments 2 and 3) to test that we observed

a peak of activity at the entrained frequencies, both at group and individual level (Supplementary materials

and methods: *Entrainment at single subject level*) and also verified the degree of phase locking of neural

signal to external stimulation.

SSVEPs were obtained for each of the rhythmic entrainment conditions by aligning the 5 blocks of each

entrainment condition (four conditions in Experiment 2, and two in Experiment 3) to stimulus presentation.

SSVEPs were then averaged across the electrodes of the ROI and the power spectrum was calculated with

the same method described above for the IAF at rest, but using a window length of 2 seconds (in order to

capture at least 5 cycles of the slowest entrainment activity, 3 Hz, in each of the windows of analysis). We

normalised frequencies relative to the IAF prior to averaging the power spectrum across participants.

To calculate phase locking we used cross-coherence (XCOH) (Keitel et al., 2017) to assess the extent to

which neural activity was synchronised to rhythmic stimulation. We calculated XCOH between external

rhythmic stimulation (train of Gaussian contrast increments) and each of the channels of the ROI for each

of the blocks (we discarded the initial and final 5 seconds of each block, resulting in 110 s of continuous

data). We used 1 second non-overlapping segments (Hanning window) and 2048 points to calculate the

FFT. Coherence was averaged across trials and electrodes of the ROI for each participant and experimental

condition. Frequencies were normalised to IAF cycles and, finally, the value of coherence at each of the

entraining frequencies of interest was selected and averaged across participants.

For the IAF condition in Experiment 3, we applied an additional entropy analysis adapted from Notbohm

et al. (Notbohm et al., 2016), in order to show that rhythmic stimulation results in a stronger phase

synchrony than arrhythmic stimulation. The time series of visual stimulation and concurrently recorded

EEG were band-passed around the frequency of stimulation (see above) for rhythmic and arrhythmic

conditions. Segments of data containing artefacts were discarded. For clean data, we obtained the phase

difference between visual stimulation and concurrent EEG and calculated modified entropy as described

by Notbohm et al. (Notbohm et al., 2016), using N=80 bins for entropy calculation. In case of entrainment,

modified entropy would be larger for the rhythmic condition compared to the arrhythmic condition.

Pre-registered statistical tests

For all statistical tests, the alpha level was set to 0.05. The correlation between the NAR and IAF was tested

by means of a Pearson correlation (right tailed). The impact of alpha speed on the NAR was tested by means

of a paired t-test (left tailed) for IAF-2Hz compared to IAF+2Hz. A one tailed t-test (right) was used to test

the hypothesis that IAF entrainment results in a faster NAR than arrhythmic entrainment.

A "two one sided tests" (TOST) (Lakens, 2017) was used to assess frequency specificity (following the

prediction that only entrainment at or around alpha impacts BR dynamics). A TOST is significant when

both p-values for left and right differences are below the set significance alpha level and the CI is contained

within the pre-defined equivalence interval. For this test, the NAR at 3 Hz was normalised with respect to

the 0 Hz NAR, and the equivalence interval was set to [-10 10] % before collecting data for Experiment 3

(https://osf.io/w8m4q/) based on simulations (Supplementary Materials and methods).

Exploratory and control statistical tests (not pre-registered analyses)

In order to extend the evaluation of the potential correlation between IAF and NAR across different datasets,

the p-values from correlations in Experiments 1 and 3 were combined by means of the Stouffer method

(Vanrullen, 2016).

The entrainment measure for Experiments 1 and 3 (EEG phase-alignment to external stimulation) was

assessed by testing the significance of XCOH peaks at each of the entrainment frequencies. We applied the

statistical test described by Biltoft et al (Biltoft and Pardyjak, 2009) using 2048 points for FFT estimation,

5 blocks of data, and 110 seconds of duration each at 500 Hz sampling rate, which resulted in 147 degrees

of freedom. In this range of parameters, any coherence peak larger than 0.0405 would be significant.

Lastly, the comparison regarding neural signal synchrony between rhythmic and arrhythmic stimulation

conditions in Experiment 3 was assessed by means of a paired one-tailed Wilcoxon test (Fay and Proschan,

2010).

<u>Results</u>

According to behavioural responses in the BR task, the proportion of time spent in each of the two rival

percepts was balanced in all three experiments. On average, participants reported a red percept 44±8% of

the total recorded time, green 50±7% and no percept 6±6%, with similar proportions for all experimental

conditions. For the calculation of the NAR, we used, on average, 231±71 percepts per condition for

Experiment 1, 179±66 for Experiment 2 and 171±85 for Experiment 3.

Although IAF at rest can sometimes differ from IAF measured during task (Haegens et al., 2014), we did

not observe these differences here for Experiment 1 (IAF_{rest} – IAF_{task}=0.14±1.1 Hz, t(26)=0.77, p-

value=0.4) nor for Experiment 3 (IAF_{rest}-IAF_{task}=0.14±0.9 Hz, t(29)=1.30,p-value=0.2). The differences

were evaluated by means of a paired t-test (two-tailed). In Experiment 2, as all conditions included

entrainment, IAF could not be measured during the task without interference and this test was not

performed.

The speed of endogenous alpha activity correlates with the speed of perceptual alternations

In line with our first prediction, in Experiment 1 we observed a significant correlation between IAF during

the task and NAR (Fig. 3) (r=0.35, p-value=0.037). Visual inspection of the correlation plot opened the

question of whether significance might be driven by two potential outliers with low IAF frequencies. From

an electrophysiological point of view, the IAFs of these two subjects fall within the usual range of IAFs

reported in the literature, 7 to 14 Hz (Haegens et al., 2011; Smulders et al., 2018), and therefore provide

informative points in the analysis. Nevertheless, according to the Matlab toolbox Robust Correlation

(Pernet et al., 2013), these two points were classified as bivariate outliers. In order to discard that the

significant correlation observed in the main analysis was due to outliers, we calculated the robust correlation

coefficient with the above-mentioned toolbox, following the recommendations of Pernet et al (Pernet et al.,

2013). Data were distributed normally (Henze-Zirkler normality test p-value = 0.21) and included two

bivariate outliers. We used the skipped Pearson correlation, which estimates the strength of the linear

correlation ignoring outliers but taking into account the overall structure of the data, as recommended for

this situation (Pernet et al., 2013). The correlation parameters obtained were in line with those obtained

with the main (pre-registered) analysis (r=0.35, p-value=0.037).

Note that Experiment 3, which was designed to test the effect of periodicity and frequency specificity, also

contained a no-stimulation condition, and therefore afforded an estimation of the correlation between IAF

and NAR, but with substantially fewer trials. Here, we observed a correlation in the expected direction but

that was not significant (r=0.19, p-value=0.15), perhaps not surprisingly given the reduced statistical power.

As the dataset in Experiment 1, data in Experiment 3 were distributed normally (p-values for the Henze-

Zirkler normality test p-value = 0.21) and included six bivariate outliers. The skipped Pearson correlation

returned a similar estimate (r=0.23 p-value=0.11 for Experiment 3).

Taken individually, the evidence for the correlation between IAF and NAR supports the initial hypothesis,

but it is not particularly strong. In order to garner more support, we used combined statistics and meta-

analysis, in post-hoc confirmatory analyses. First, we found that the combined statistics of Experiment 1

and 3 (Stouffer method) rendered a significant effect (p-value=0.024). In addition, we performed a meta-

analysis using the four independent datasets that, as far as we know, have tested IAF-NAR correlation:

(Katyal et al., 2019), data from a previous experiment of our group (Pápai et al., 2018), and data from

Experiments 1 and 3 in the current study. The meta-analysis was performed with the R package "meta",

using Fisher's z transformation of correlations. In order to homogenise datasets between studies we

transformed all behavioural data to natural alternation times (NAT=1/NAR), the method used by Katyal et

al. (please note that contrary to the NAR-IAF correlation, the expected direction for the NAT-IAF

correlation is negative). For the same reason, we used the mean of the percept duration distributions instead

of the median. This meta-analysis provided further confirmation of the correlation between BR dynamics

and endogenous IAF activity (r=-0.44, p-value<0.001). Lastly, in order to take into account publication bias

in the meta-analysis, we used the p-uniform method (van Assen et al., 2014). This method assesses the

distribution of the sum of independent uniformly distributed random variables using the p-method (Irwin-

Hall distribution, as implemented in the p-uniform package for R https://cran.r-

project.org/web/packages/puniform/). After correcting for publication bias, the correlation was, once more,

confirmed (r=-0.47, p-value=0.011). Altogether, the data supports our first hypothesis regarding the

prediction that alpha frequency is correlated with the natural alternation between percepts, across

individuals.

Figure 3: Natural alternation rate as a function of IAF during task for Experiments 1 (red) and 3

(green). Dotted lines correspond to the linear fit of NAR as a function of IAF.

The frequency of endogenous activity is causally related to the rate of alternations in conscious perception

In order to make sure that rhythmic visual stimulation successfully entrained endogenous oscillatory

activity, in Experiment 2, we evaluated SSVEP and cross coherence (XCOH) at group level. The group-

averaged SSVEP displayed a peak in the corresponding frequency of each of the entrainment conditions

(Fig. 4. A), and the cross-coherence measure presented a significant peak (3 Hz: XCOH=0.1097, p-

value=0.0002; IAF-2Hz: XCOH=0.135, p-value=0.00003; IAF: XCOH=0.0841, p-value=0.0017;

IAF+2Hz: XCOH=0.1133, p-value=0.0002) (Fig. 5. B and C).

Figure 4: Entrainment in Experiment 2. A: Group average SSVEP. The frequencies are normalised with respect to rest IAF (the one used during IAF entrainment) and 1/f background noise correction was applied prior to the group averaging. Vertical dotted lines represent the target frequency at each condition (only IAF is exactly equal to one in the normalised plot, shaded areas indicate the lower and upper bounds for 3 Hz, IAF-2Hz and IAF+2Hz normalised values for each participant). B: Cross-coherence between the EEG signal and the rhythmic entrainment at the frequencies corresponding to each entraining condition. Dots represent individual values, black lines are the mean value for each condition and the dark shaded area corresponds to the standard error of the mean. The red dotted line presents the value of the coherence at a significance level 0.05. C: Cross-coherence spectrum for each of the experimental conditions.

In Experiment 2, the NAR at 3 Hz was slower than any of the other conditions (Fig. 5. A and Table 1). In the condition of interest, as expected, we observed a significant decrease in the rate of perceptual alternations during the slow alpha entrainment condition (IAF-2Hz) with respect to the fast alpha entrainment condition (IAF+2Hz) (p-value=0.03, t(24)=-1.942, d=0.39) (Fig. 6. C). Entraining at the IAF produced NARs between the slow IAF-2Hz and the fast IAF+2Hz entrainment, as one would expect, but it was not significantly different from either of the two (Supplementary results: *Non-planned contrasts*). This finding confirms our second prediction, regarding the causal modulation of perceptual alternations by visual entrainment.

Figure 5: Effect of entrainment on perceptual alternations. A: NAR for Experiment 2 at the different experimental conditions. B: NAR for Experiment 3. C: NAR differences in the conditions of interest (IAF-2 Hz compared to IAF +2 Hz, 0 Hz compared to 3 Hz and IAF compared to arrhythmic IAF). IAF-2Hz was significantly slower than IAF+2 Hz, whereas 0 Hz and 3 Hz were equivalent (for a 10% equivalence interval) and the IAF rhythmic condition was significantly faster than the arrhythmic condition. Solid lines correspond to the group mean of each condition; dark shaded areas correspond to the mean ± standard error of the mean. Light shaded areas correspond to the distribution of the individual data and dots correspond to individual data points.

Rhythmic entrainment boosts the re-evaluation of competing stimuli

In Experiment 3, group averaged SSVEPs displayed peaks at each of the entraining conditions (Fig. 6. A). Endogenous activity was successfully entrained, as confirmed by the significant cross-coherence in all experimental conditions (3 Hz: XCOH=0.1251, p-value=0.0001; IAF: XCOH=0.0711, p-value=0.0048) (Fig. 6. B and C) and the higher values of normalised entropy observed in the IAF compared to arrIAF condition (p-value=0.000001) (Fig. 6. D). As would be expected, the cross-coherence for the arrhythmic condition was not significant (Fig. 6. B and C) (arrIAF: XCOH=0.016 p-value=0.32).

The comparison of interest here was between the rate of perceptual alternations during rhythmic (at IAF) and arrhythmic (arrIAF) entrainment. In the arrhythmic condition, intervals between pulses were randomly picked from an exponential distribution of latencies centred at the inverse of IAF. IAF entrainment resulted

421   in significantly faster NAR than arrhythmic entrainment (p-value=0.0005, t(29)=3.659, d=0.67) (Fig. 5. C),

422   indicating that periodicity plays a key role in BR dynamics. This confirms the third main prediction,

423   regarding periodicity.

Figure 6: Entrainment in Experiment 3. A: Group average SSVEP. The frequencies are normalised with respect to rest IAF (the one used during IAF entrainment) and 1/f background noise correction was applied prior to the group averaging. Vertical dotted lines represent the target frequency at each condition (only IAF is exactly equal to one in the normalised plot, shaded area indicate the lower and upper bounds for 3 Hz, normalised values for each participant). B: Values of coherence between neural signal and rhythmic stimulation at the frequencies corresponding to each entraining condition. Dots represent individual values, black lines are the mean value for each condition and the dark shaded area corresponds to the standard error of the mean. The red dotted line presents the value of the cross-coherence at a significance level 0.05. C: Cross-coherence spectrum for each of the experimental conditions. D: Difference between normalised entropy at IAF and arrIAF conditions. Dots represent individual values, black thick line is the mean value and dark grey shaded area is the standard error of the mean.

In other words, although arrhythmic and rhythmic alpha entrainment always resulted in faster perceptual

alternation rates compared to low-frequency (3 Hz) or no visual entrainment (0 Hz) (Fig. 5. B and

Supplementary results: *Non-planned contrasts*)), the increase in alternation rate in IAF entrainment cannot

be solely attributed to an increase in the number or rate of pulses per unit of time (Brascamp et al., 2015).

Instead, a mechanism involving periodicity is required to fully explain the result.

Modulations in perceptual alternation rate are specific to entrainment in the alpha band

Further confirming the frequency-specificity implication of the hypothesis, we observed that the speeding

up of perceptual alternations was constrained to the alpha band. Conditions that did not involve stimulation

in the spectral vicinity of alpha always produced slower perceptual alternation rates. What is more, 3 Hz

entrainment produced an alternation rate equivalent to no stimulation at all, as measured with two one sided

tests (TOST (Lakens, 2017)) for an equivalence interval of 10% (p-value$_{left}$=0.007, p-value$_{right}$=0.0001, CI=

[-0.07 0.03]) (Fig. 5. C).

<u>Discussion</u>

We addressed the hypothesis that alpha fluctuations in the occipital cortex mediate the accumulation of

visual information, leading to perceptual representations. Our findings, so far, seem to stand in good

agreement with this hypothesis. First, we confirmed that the speed of fluctuations in cortical alpha activity

correlates with the dynamics of perceptual fluctuations in BR, a result also recently observed in another

independent study (Katyal et al., 2019). The evidence provided by each individual experiment was weak to

moderate, but aggregated results from four experiments indicate a robust relationship between Alpha

frequency and alternation rate. We further proposed that endogenous alpha rhythm provides temporal

windows for the re-evaluation of competing visual representations based on visual input. Consequently,

inducing faster endogenous activity should accelerate perceptual switches, a result we observed in

Experiment 2. Moreover, our results suggest a causal relationship between alpha activity and awareness

dynamics, since modulation of endogenous brain rhythms resulted in consequential changes in conscious

perception. That is, speeding up or slowing down alpha cortical activity with entrainment alters the rate of

perceptual alternations accordingly. Critically, this causal modulation is restricted to the alpha range and

its effectiveness is subject to the periodicity of stimulation.

These findings are consistent with the pacemaker role of alpha activity in perceptual alternations, which we

propose here. We build from two well-known principles. First, that reciprocal inhibition between the neural

populations selective of the competing stimuli is the mechanism causing perceptual alternations in BR

(Tong et al., 2006). Second, that the activation of the neural population selective of the dominant conscious

percept tends to decrease over time due to adaptation (Shpiro et al., 2009), making perceptual switches

more likely with time. In this context, inhibition plays an active role: by stifling the neural population with

lower excitability, it increases the signal to noise ratio corresponding to the alternative population

(Klimesch, 2012). We hypothesise that this process takes place in a phasic manner: each alpha cycle opens

a window for re-evaluation of the competing neural signals. In each of these windows, the weight of each

neural population is evaluated, and the outcome of each evaluation determines the resulting conscious

percept. Successive alpha cycles lead to the progressive accumulation of evidence toward one or the other

percept.

Could this accumulation happen with changes in excitability regardless of rhythmic fluctuations?

(Brascamp et al., 2015) In Experiment 3, we tested whether local, non-rhythmic changes in stimulus

contrast could account for the increase in perceptual alternation rate during entrainment. Although

arrhythmic stimulation increased alternation rates with respect to no-stimulation, these changes could not

account for the effect in full. Rhythmic IAF stimulation resulted in faster alternations than arrhythmic

stimulation. Additionally, if local changes in contrast had been responsible for speeding up switch

dynamics, a difference between the no-stimulation condition and 3 Hz stimulation condition would have

been expected. Yet, these two conditions rendered equivalent, and low, alternation rates.

Our manipulation could have affected the interplay between noise and adaptation, reflected in the

coefficient of variation. Studies using anti-phase stimulation have reported drops in the coefficient of

variation (Kim et al., 2006), implying a decrease in the contribution of noise to BR dynamics. In our study,

the analysis of the coefficient of variation implied that the visual stimulation protocol used here did not

modify the basic mechanism of neural competition. Therefore, one can assume that the competition

mechanism remained unchanged and that only the frequency and probability of windows favouring the re-

evaluation of perceptual representations were modified.

Sensory entrainment at or around the IAF produced a dramatic increase in perceptual alternations compared

to 3Hz or no-stimulation. That is, alpha entrainment produces a sizable effect, even if there is already an

ongoing rhythm at that frequency. What is the origin of this increase? In line with the well-known fact that

endogenous alpha occurs in bursts rather than in a sustained fashion (Rusiniak et al., 2018), we speculate

that alpha-gated re-evaluation mostly occurs within these bursts. That is, in BR the re-evaluation of

competing stimuli might not unfold homogenously over time, but intermittently. Incidentally, this may lead

to the stochastic dynamics that characterises the phenomenon. In our protocol, entrainment at the natural

alpha rhythm must have increased alpha burst episodes, explaining the faster alternation rate under IAF,

compared to no entrainment. During these alpha bursts, in order to reduce stimulus complexity, information is compressed into discrete packages (Vanrullen and Koch, 2003). The dynamics of the system alternate between intervals where the probability of switch is low and recurring intervals of alpha activity where competition between conflicting representations is steep. Therefore, by increasing alpha burst episodes, the probability of stimulus re-evaluation increases and, as a result, faster alternation rates can be expected. The role of alpha in the gating by inhibition hypothesis (Jensen and Mazaheri, 2010) is compatible with the hypothesis laid out here. According to the former, sensory information is coded in gamma activity embedded within the high excitability windows of slower alpha cycles. One could argue that, in fact, slower alpha cycles may allow for more nested gamma cycles, hence lead to more accumulation of evidence per alpha cycle, predicting a result contrary to what we observed. However, variations in the speed of gamma may compensate for the different number of cycles per second across alpha speeds. Indeed, several findings suggest that the length of gamma cycles vary between subjects, and subjects with faster alpha also present faster gamma (Muthukumaraswamy et al., 2009; Baumgarten et al., 2018). We speculate that the number of gamma bursts per alpha cycle is approximately constant, therefore faster alpha rhythms would allow for an increased number of gamma bursts per second, in line with our hypothesis and compatible with gating by inhibition.

Regarding frequency specificity, we cannot exclude the possibility that the NAR would increase with entrainment faster than alpha, as we did not stimulate at frequencies above the alpha range. However, high frequency stimulation may not be an effective test for specificity, given that alpha activity could still be entrained as a sub-harmonic of higher frequencies, as some studies suggest (Herrmann, 2001; Herrmann et al., 2016). Instead, several independent results favour a frequency specificity interpretation. From our study, we can conclude that there is a lower frequency bound of at least 3Hz, to modulate NAR effectively. In addition, evidence from O'Shea and Crassini (O'shea and Crassini, 1984) provide indirect support for the existence of a higher bound. O'Shea and Crassini presented flickering stimuli at different frequencies during a BR protocol, and observed a pattern of NAR in line with our predictions: maximum NARs were observed in the alpha range and, additionally, subjects reported that flickering at faster frequencies (above the alpha range) disappeared, a result consistent with discrete sampling. However, methodological differences (i.e., only 3 subjects were tested and stimuli were presented for on-off periods) prevent a conclusive inference.

It is possible that the physical stimulation based on contrast changes used here resulted in a modulation of

attention, which has a well-known effect on BR (Zhang et al., 2011; Brascamp and Blake, 2012).

Compatible with our hypothesis, attention has been proposed to periodically sample rivalling stimuli during

BR (Davidson et al., 2018), and it is possible that the increase in NAR observed in the arrhythmic

stimulation condition is due to a global increase in arousal. Our findings can be framed in terms of existing

theoretical models of BR which incorporate attention (Li et al., 2017). According to these models, attention

recurrently (or rhythmically, in our hypothesis) regulates the imbalance between the two rival stimuli with

a slow time constant, triggered by a mutual inhibition process that has a fast(er) time constant. We can

interpret this model in terms of a slow rhythm (alpha) and a fast rhythm. This faster rhythm could be gamma,

which has also been found to correlate with BR dynamics in previous studies (Fesi and Mendola, 2015).

The concept of alpha discrete sampling can also be framed within Bayesian models of BR (Hohwy et al.,

2008; Gershman et al., 2012). It has been proposed that the continuous input of inconsistent stimuli creates

a situation in which neither of the stimuli representations has both high prior and high likelihood. As a

result, periodic sampling determines which of the two representations sways the perceptual outcome at any

given time (Hohwy et al., 2008).

The present findings provide a framework for the interpretation of inter-individual differences in BR

dynamics: in this framework, endogenous alpha cycles are a representative time constant for each subject

that regulates the competition process between neural populations representative of each stimuli. Alpha acts

as a pacemaker for information accumulation mediating the transition to awareness by setting the tempo

for the competition process. According to this interpretation, our results are in line with recent studies

suggesting that alpha activity may not regulate objective aspects of perception, such as perceptual

sensitivity, but awareness and confidence about perceptual properties (Lange et al., 2013; Benwell et al.,

2017; Samaha et al., 2017).

Data availability

The datasets generated during and/or analysed during the current study are available from the corresponding

author on reasonable request.

References

Baumgarten TJ, Neugebauer J, Oeltzschner G, Füllenbach ND, Kircheis G, Häussinger D, Lange J,

Wittsack HJ, Butz M, Schnitzler A (2018) Connecting occipital alpha band peak frequency, visual

temporal resolution, and occipital GABA levels in healthy participants and hepatic encephalopathy

patients. NeuroImage Clin 20:347–356.

Benwell CSY, Tagliabue CF, Veniero D, Cecere R, Savazzi S, Thut G (2017) Prestimulus EEG Power

Predicts Conscious Awareness But Not Objective Visual Performance. Eneuro 4:ENEURO.0182-

17.2017.

Biltoft CA, Pardyjak ER (2009) Spectral coherence and the statistical significance of turbulent flux

computations. J Atmos Ocean Technol 26:403–410.

Block N (2019) What Is Wrong with the No-Report Paradigm and How to Fix It. Trends Cogn Sci

23:1003–1013.

Brascamp JW, Blake R (2012) Inattention Abolishes Binocular Rivalry: Perceptual Evidence. Psychol Sci

23:1159–1167.

Brascamp JW, Ee R Van, Pestman WR, Berg AV Van Den (2005) Distributions of alternation rates in

various forms of bistable perception. :287–298.

Brascamp JW, Klink PC, Levelt JM (2015) The ' laws ' of binocular rivalry : 50 years of Levelt ' s

propositions. 109:20–37.

Carter OL, Pettigrew JD (2003) A common oscillator for perceptual rivalries? Perception 32:295–305.

Cavanagh P, MacLeod DI, Anstis SM (1987) Equiluminance: spatial and temporal factors and the

contribution of blue-sensitive cones. J Opt Soc Am A, Opt Image sicence 4:1428–1438.

Cecere R, Rees G, Romei V (2015) Individual Differences in Alpha Frequency Drive Crossmodal

Illusory Perception. Curr Biol 25:231–235

Clark CR, Veltmeyer MD, Hamilton RJ, Simms E, Paul R, Hermens D, Gordon E (2007) Spontaneous

alpha peak frequency predicts working memory performance across the age span. Int J

Psychophysiol 53:1–9.

Davidson MJ, Alais D, Boxtel JJA Van (2018) Attention periodically samples competing stimuli during

binocular rivalry. :1–25.

de Jong MC, Hendriks RJM, Vansteensel MJ, Raemaekers M, Verstraten FAJ, Ramsey NF, Erkelens CJ,

Leijten FSS, van Ee R (2016) Intracranial Recordings of Occipital Cortex Responses to Illusory

Visual Events. J Neurosci 36:6297–6311.

Doesburg SM, Kitajo CAK, Ward LM (2005) Increased gamma-band synchrony precedes switching of

conscious perceptual objects in binocular rivalry. 16:1–4.

Drew A, Torralba M, Ruzzoli M, Morís Fernández L, Sabaté A, Szabina Pápai M, Soto-Faraco S (2019)

Neural Evidence of Cognitive Conflict During Binocular Rivalry. SSRN Electron J:25–27.

Fay MP, Proschan MA (2010) Wilcoxon-Mann-whitney or t-test? Stat Surv 4:1–37.

Fesi JD, Mendola JD (2015) Individual Peak Gamma Frequency Predicts Switch Rate in Perceptual

Rivalry. Hum Brain Mapp 576:566–576.

Friston K (2012) Prediction, perception and agency. Int J Psychophysiol 83:248–252.

Gershman SJ, Vul E, Tenenbaum JB (2012) Multistability and perceptual inference. Neural Comput.

Grandy TH, Werkle-Bergner M, Chicherio C, Schmiedek F, Lövdén M, Lindenberger U (2013) Peak

individual alpha frequency qualifies as a stable neurophysiological trait marker in healthy younger

and older adults. Psychophysiology 50:570–582.

Haegens S, Cousijn H, Wallis G, Harrison PJ, Nobre AC (2014) Inter- and intra-individual variability in

alpha peak frequency. Neuroimage 92:46–55.

Haegens S, Nácher V, Luna R, Romo R, Jensen O (2011) α-Oscillations in the monkey sensorimotor

network influence discrimination performance by rhythmical inhibition of neuronal spiking. Proc

Natl Acad Sci U S A 108:19377–19382.

Halgren E (1990) Human Evoked Potentials. Neurophysiol Tech II:147–276.

Herrmann CS (2001) Human EEG responses to 1-100 Hz flicker: Resonance phenomena in visual cortex

and their potential correlation to cognitive phenomena. Exp Brain Res 137:346–353.

Herrmann CS, Murray MM, Ionta S, Hutt A, Lefebvre J (2016) Shaping intrinsic neural oscillations with

periodic stimulation. J Neurosci 36:5328–5337.

Hohwy J, Roepstorff A, Friston K (2008) Predictive coding explains binocular rivalry: An

epistemological review. Cognition 108:687–701.

Jensen O, Mazaheri A (2010) Shaping functional architecture by oscillatory alpha activity: gating by

inhibition. Front Hum Neurosci 4:186.

Kanai R, Moradi F, Shimojo S, Verstraten FAJ (2005) Perceptual alternation induced by visual transients.

Perception.

Kang M (2009) Size matters : A study of binocular rivalry dynamics. 9:1–11.

Katyal S, He S, He B, Engel SA (2019) Frequency of alpha oscillation predicts individual differences in

perceptual stability during binocular rivalry. Hum Brain Mapp:1–12.

Keitel C, Thut G, Gross J (2017) Visual cortex responses reflect temporal structure of continuous quasi-

rhythmic sensory stimulation. Neuroimage 146:58–70.

Kim YJ, Grabowecky M, Suzuki S (2006) Stochastic resonance in binocular rivalry. Vision Res 46:392–

406.

Kleinschmidt A, Sterzer P, Rees G (2012) Variability of perceptual multistability: From brain state to

individual trait. Philos Trans R Soc B Biol Sci.

Klimesch W (2012) α-band oscillations, attention, and controlled access to stored information. Trends

Cogn Sci 16:606–617.

Lakens D (2017) Equivalence Tests: A Practical Primer for t Tests, Correlations, and Meta-Analyses. Soc

Psychol Personal Sci 8:355–362.

Lange J, Oostenveld R, Fries P (2013) Reduced occipital alpha power indexes enhanced excitability

rather than improved visual perception. J Neurosci 33:3212–3220.

Lee SH, Blake R, Heeger DJ (2007) Hierarchy of cortical responses underlying binocular rivalry. Nat

Neurosci 10:1048–1054.

Leopold DA, Logothetis NK (1999) Multistable phenomena : changing views in perception. 6613:254–

264.

Li H, Rankin J, Rinzel J, Carrasco M, Heeger DJ (2017) Attention model of binocular rivalry.

Mamassian P, Goutcher R (2005) Temporal dynamics in bistable perception. J Vis 5:7.

Miller SM, Hansell NK, Ngo TT, Liu GB, Pettigrew JD, Martin NG, Wright MJ (2010) Genetic

contribution to individual variation in binocular rivalry rate. 107:9–13.

Muthukumaraswamy SD, Edden RAE, Jones DK, Swettenham JB, Singh KD (2009) Resting GABA

concentration predicts peak gamma frequency and fMRI amplitude in response to visual stimulation

in humans. Proc Natl Acad Sci U S A 106:8356–8361.

Notbohm A, Kurths J, Herrmann CS (2016) Modification of Brain Oscillations via Rhythmic Light

Stimulation Provides Evidence for Entrainment but Not for Superposition of Event-Related

Responses. Front Hum Neurosci 10.

O'shea RP, Crassini B (1984) Binocular rivalry occurs without simultaneous presentation of rival stimuli.

Percept Psychophys 36:266–276.

Pápai MS, Cuello MT, Bayer M, Soto-faraco S (2018) Occipital Alpha modulates the processing of

suppressed visual stimuli during binocular rivalry..

Pernet CR, Wilcox R, Rousselet GA (2013) Robust correlation analyses: False positive and power

validation using a new open source matlab toolbox. Front Psychol 3:1–18.

Phillips I, Morales J (2020) The Fundamental Problem with No-Cognition Paradigms. Trends Cogn Sci

24:165–167.

Piantoni G, Kline KA, Eagleman DM (2010) Beta oscillations correlate with the probability of perceiving

rivalrous visual stimuli. 13:1–11.

Polonsky A, Blake R, Braun J, Heeger DJ (2000) Neuronal activity in human primary visual cortex

correlates with perception during binocular rivalry. :1153–1159.

Ronconi L, Busch NA, Melcher D (2018) Alpha-band sensory entrainment alters the duration of temporal

windows in visual perception. Sci Rep 8:1–10.

Rusiniak M, Wróbel A, Cieśla K, Pluta A, Lewandowska M, Wójcik J, Skarżyński PH, Wolak T (2018)

The relationship between alpha burst activity and the default mode network. Acta Neurobiol

Exp:92–106.

Samaha J, Iemi L, Postle BR (2017) Prestimulus alpha-band power biases visual discrimination

confidence, but not accuracy. Conscious Cogn 54:47–55.

Samaha J, Postle BR (2015) The Speed of Alpha-Band Oscillations Predicts the Temporal Resolution of

Visual Perception Report The Speed of Alpha-Band Oscillations Predicts the Temporal Resolution

of Visual Perception. Curr Biol 25:2985–2990.

Schmack K, Sekutowicz M, Rössler H, Brandl EJ, Müller DJ, Sterzer P (2013) The influence of

dopamine-related genes on perceptual stability. Eur J Neurosci.

Shpiro A, Moreno-Bote R, Rubin N (2009) Balance between noise and adaptation in competition models

of perceptual bistability. J Comput Neurosci 27:37–54.

Smulders FTY, ten Oever S, Donkers FCL, Quaedflieg CWEM, van de Ven V (2018) Single-trial log

transformation is optimal in frequency analysis of resting EEG alpha. Eur J Neurosci 48:2585–

2598.

Sobel K V., Blake R (2002) How context influences predominance during binocular rivalry. Perception.

Sterzer P, Kleinschmidt A, Rees G (2009) The neural bases of multistable perception. Trends Cogn Sci

13:310–318.

van Assen MALM, van Aert RCM, Wicherts JM (2014) Meta-Analysis Using Effect Size Distributions of

Only Statistically. Pychological Methods.

Vanrullen R (2016) How to evaluate phase differences between trial groups in ongoing

electrophysiological signals. :1–24.

Vanrullen R, Busch NA, Drewes J, Dubois J (2011) Ongoing EEG phase as a trial-by-trial predictor of

perceptual and attentional variability. Front Psychol 2:1–9.

Vanrullen R, Koch C (2003) Is perception discrete or continuous ? Trends Cogn Sci 7:207–213.

Zhang P, Jamison K, Engel S, He B, He S (2011) Binocular Rivalry Requires Visual Attention. Neuron

71:362–369.

683