Title: Alpha fluctuations regulate the accrual of visual information to awareness

Abbreviated title: Alpha-regulates information accrual to awareness

Authors

Mireia Torralba[1], Alice Drew[1], Alba Sabaté San José[2], Luis Morís Fernández[1], and Salvador Soto-

Faraco[1,3]

[1]Multisensory Research Group, Center for Brain and Cognition, University of Pompeu Fabra, Barcelona,

Spain, ZIP: 08002.

[2]Universitat de Barcelona, Barcelona, Spain, ZIP: 08007.

[3]Catalan Institution for Research and Advanced Studies (ICREA), Barcelona, Spain, ZIP: 08010.

Corresponding author: Mireia Torralba Cuello (mireia.torralba@upf.edu), ZIP: 08002

Number of pages: 21

Number of figures: 6

Number of tables: 1

Number of words: Abstract 190, Introduction 620, Discussion 1462

Competing interest

The authors declare no competing financial interests.

Acknowledgments

This research was supported by the Ministerio de Economia y Competitividad (PSI2016-75558-P

AEI/FEDER), AGAUR Generalitat de Catalunya (2017 SGR 1545). We want to thank Dr. Márta Szabina

Pápai and Indre Pileckyte for helping in the piloting of the Experiments and in EEG data collection.

Abstract:

Endogenous brain processes play a paramount role in shaping up perceptual phenomenology, as

illustrated by the alternations experienced by humans (and other animals) when watching perceptually

ambiguous, static images. Here, we hypothesised that endogenous alpha fluctuations in the visual cortex

pace the accumulation of sensory information leading to perceptual outcomes. We addressed this

hypothesis using binocular rivalry combined with visual entrainment and electroencephalography in

humans (42 female, 40 male). The results revealed a correlation between the individual frequency of

alpha oscillations in the occipital cortex and perceptual alternation rates experienced during binocular

rivalry. In subsequent experiments we show that regulating endogenous brain activity via entrainment

produced corresponding changes in perceptual alternation rate, which were observed only in the alpha

range but not at lower entrainment frequencies. Overall, rhythmic alpha stimulation resulted in faster

perceptual alternation rates, compared to arrhythmic or no stimulation. These findings support the notion

that visual information is accumulated via alpha cycles to promote the emergence of conscious perceptual

representations. We suggest that models of binocular rivalry incorporating posterior alpha as a pacemaker

can provide an important advance in the comprehension of the dynamics of visual awareness.

Key words

Binocular rivalry, sensory entrainment, alpha activity, EEG, individual variability, discrete perception.

Significance statement:

Mainstream theories in cognitive neuroscience agree that endogenous brain processes play a paramount

role in shaping our perceptual experience of sensory inputs. In vision, endogenous fluctuations in the

alpha rhythm have been pointed out to regulate visual inputs to perception. In support of this hypothesis,

here we used EEG recordings and visual entrainment to demonstrate that inter-individual differences in

the speed of endogenous alpha fluctuations in the brain are causally related to the accrual of visual

information to awareness. These findings provide, for the first time, evidence for alpha-gated regulation

of the dynamics of alternations in conscious visual perception.

    Introduction

There is ample agreement that our perception of the world is not built from incoming sensory input alone,

but is also determined by endogenous brain processes (Friston, 2012). The impact of these endogenous

processes is well illustrated by spontaneous fluctuations in conscious perception when watching

ambiguous images, a phenomenon known as bistable perception (Sterzer et al., 2009). Ambiguous images

lead observers to experience alternations between different perceptual interpretations of an otherwise

static input and have been used by neuroscientists to investigate the neural correlates of fluctuations in

perceptual awareness. Here, we address the hypothesis that alpha oscillatory activity in the occipital

cortex plays a causal role in these perceptual fluctuations.

We focused on binocular rivalry (BR), a phenomenon that arises when two dissimilar images are

presented to different eyes. BR has been characterised as an intrinsically dynamic process resulting from

competition and mutual inhibition between different neural populations representing the rival images

(Mamassian and Goutcher, 2005). There is still controversy about the specific brain areas involved in this

competition (Leopold and Logothetis, 1999; Block, 2019; Phillips and Morales, 2020), however, evidence

suggests that the occipital cortex is involved in resolving perceptual ambiguity (Polonsky et al., 2000; Lee

et al., 2007; de Jong et al., 2016). In some models, BR competition is not solely based on mutual

inhibition but also on attentional modulation (Li et al., 2017).

BR is characterised by the natural alternation rate (NAR) between dominance periods, a stochastic

process strongly dependant on properties of the rival stimuli (Sobel and Blake, 2002; Brascamp et al.,

2005; Kang, 2009). Interestingly, the NAR is relatively stable within individuals and genetically

hereditary (Miller et al., 2010). Additionally, although inter-individual variability in BR dynamics is large

(Kleinschmidt et al., 2012), alternation rates within an individual correlate across different forms of

bistable perception (Schmack et al., 2013) and stimuli (Carter and Pettigrew, 2003).

Recently, several studies have linked BR with brain oscillatory activity in different frequency bands

(Doesburg et al., 2005; Piantoni et al., 2010), and in particular to occipital alpha activity (8-14 Hz). Alpha

oscillations are strongly related to attention (Davidson et al., 2018; Katyal et al., 2019) and, more

relevantly, alpha activity in the occipital cortex is known to be related to rhythmic fluctuations in visual

perception (Vanrullen et al., 2011). For example, the gating by inhibition hypothesis (Jensen and

Mazaheri, 2010) contends that the alpha rhythm imposes a discrete sampling of sensory input via an

alternation between moments of heightened neural excitability and bouts of pulsed inhibition, cycling

every 100 ms approximately. In line with this hypothesis, variations in the speed of alpha oscillations

correlate with inter-individual variability in the temporal resolution of visual perception (Samaha and

Postle, 2015) and processing speed in general (Clark et al., 2007). The results of entrainment studies

support the causal role of alpha oscillations in modulating perception (Cecere et al., 2015; Ronconi et al.,

2018). An interesting property of the cortical alpha rhythm is that the individual alpha frequency (IAF)

reflects systemic properties of the brain, is highly heritable, and relates to cognitive functioning (Grandy

et al., 2013).

In this study, we focus on establishing a connection between alpha endogenous activity speed and typical

BR alternation times. We propose that cortical alpha oscillations play a key role in the competition

between alternative representations during BR. In particular, we hypothesise that the alpha rhythm

provides phasic windows for the cyclical re-evaluation of the competing stimuli representations and,

consequentially, evidence accumulation takes place in a phasic manner. This discrete, alpha gated,

evidence accumulation eventually leads to perceptual switches. According to this hypothesis, faster alpha

oscillations will correlate positively with the NAR in a BR task and visual entrainment will causally

modulate BR dynamics in a frequency-specific way.

Methods

Subjects

Participants received 10€/ hour in return for their participation. They all provided written informed

consent prior to the study and were naive to the purpose of the experiment. The study was run in

accordance with the Declaration of Helsinki and the experimental protocol approved by the local ethics

committee CIEC Parc de Salut Mar (Universitat Pompeu Fabra, Barcelona, Spain).

Subject inclusion criteria in all the experiments were: normal or corrected to normal vision, not being

under medication and displaying a single peak in the alpha range during the eyes-closed pre-screening

EEG recording (see *IAF estimation*). In order to minimise alternations unrelated to endogenous activity

(Kanai et al., 2005), we excluded participants with a high percentage of EEG artefacts such as blinks, eye

movements or motor contamination in general (with a high percentage defined as more than 5% in

Experiment 1 and 10% in Experiments 2 and 3, applied separately in each of the 4 experimental

conditions). For Experiments 2 and 3, due to the reduced availability of data points per condition (5

blocks per condition compared to 9 in Experiment 1) we established an additional stability criterion. This

criterion ensured a sufficient amount of data a priori for a stable estimation of the NAR, based on our

simulations with empirical data from Experiment 1. According to this, in Experiments 2 and 3 we decided

to include only subjects with 50 or more percepts in each experimental condition which, according to

simulation, ensured a deviation in median estimation <3.5%. Please note that in Experiment 1, all subjects

had far more than 50 percepts (a percept is the period of time from seeing one pure colour to seeing the

next pure colour).

Thirty-one participants (15 female, aged 18-34, mean age 23) participated in Experiment 1. Four

participants were excluded for having more than 5% of the segmented epochs contaminated by visual and

movement artefacts (final sample N=27). Twenty-nine participants (16 female, aged 18-31, mean age 24)

participated in Experiment 2. One participant was excluded for presenting a double peak in the alpha

range, two participants were discarded due to excessive artefacts and one participant was discarded due to

insufficient time for estimation of the NAR (final sample N=25). Thirty-four participants (19 female,

aged 19-32, mean age 24) participated in Experiment 3. One participant was excluded for presenting a

double peak in the alpha range, two participants were discarded due to an insufficient number of percepts

and one participant was discarded due to a technical problem with the recordings (final sample N=30).

Apparatus and Stimuli

Participants were presented with one Gabor patch to each eye, 11.5° in diameter, Michelson contrast 0.1,

opposed orientations (±45° off vertical) and different colours (red, RGB=[1 0 0] and green, RGB adjusted

individually to match the luminance of the red Gabor patch). Both patches were surrounded by a black

annulus and contained a central fixation cross identical in each of the patches (Fig. 1). The images were

presented on a 19.8-inch CRT monitor (120 Hz refresh rate) with a grey background (10.7 cd/m^2) placed

80 cm from participants' eyes. Each image was presented to a different eye by means of stereoscope

mirror glasses. Visual stimuli were created and presented using the Matlab toolbox Psychtoolbox (RRID:

SCR_002881), version 3.

Visual stimulation for entrainment, when used, consisted of pulsed Gaussian increases in Michelson

contrast from 0.1 to 0.6, of 58 ms duration, presented simultaneously to the images displayed to the left

and right eyes. Different entrainment frequencies were then achieved by varying the offset to onset time

between pulses, whilst maintaining the slope of contrast increases constant (Fig. 1).

---- FIGURE 1 SHOULD BE PLACED HERE --

Procedure

The procedure was identical for all three experiments (Fig. 2), with the only variation between

experiments being the visual entrainment stimulation conditions (see below) and number of blocks

performed.

The experimental session started with the participant's IAF estimation during a 5-minute rest block, with

eyes closed. We then ran a protocol to match the two rival stimuli in terms of luminance for subjective

equality (Cavanagh et al., 1987). Next, the stereoscope mirrors were calibrated for the rival stimuli to

appear at the same retinal location of each eye. Finally, two training blocks (120 s each) on the BR task

were performed prior to the experimental BR runs.

All experiments contained the same rival images and were organised in 120 s blocks of a BR task. The

eye of presentation of the red and green stimuli was alternated from block to block. After each block,

there was a pause and participants were given the chance to rest before starting the next trial. The

experimental conditions used in each individual experiment were characterised in terms of the visual

stimulation used:

• No entrainment (0 Hz).

• Rhythmic entrainment: periodic contrast increments with a frequency of presentation of 3 Hz,

IAF, IAF-2Hz and IAF+2HZ, depending on the condition.

• Arrhythmic visual stimulation (arrIAF): non-periodic contrast increments. In this condition,

the latencies between contrast increments were drawn from an exponential distribution centred at

subjects' 1/IAF in order to match the number of contrast increments during the corresponding

rhythmic IAF condition of the same subject.

In Experiment 1, subjects ran 9 blocks of the 0 Hz condition. In experiments 2 and 3 subjects ran 5 blocks

of each experimental condition: 3 Hz, IAF-2Hz, IAF and IAF+2Hz for Experiment 2 and 0 Hz, 3 Hz, IAF

and arrIAF for Experiment 3. (Fig. 2). The order of conditions was intermixed pseudo-randomly between

blocks (Fig. 2).

During the task, participants were instructed to fixate their gaze on the central fixation cross and report

their current percept (red, green, mixed or null) by key press with the left and right index of each hand.

Response mapping for the green stimulus and the red stimulus was counter-balanced between

participants. To report a mixed (piecemeal) percept, participants were instructed to press both keys

simultaneously and to release both keys if none of the three aforementioned percepts were apparent (e.g.,

null percept).

Predictions

In this study, we performed a series of three experiments, each of them designed to test specific

predictions derived from our hypothesis. The predictions and associated experiments are as follows.

NAR correlates positively with IAF. If alpha endogenous activity paces information accrual, one first

expected result is that faster individual alpha oscillations will lead the individual to experience a faster

natural alternation rate (NAR) in a BR task. Experiment 1 was designed to test this prediction: EEG

endogenous activity was concurrently recorded while participants performed a BR task. The experiment

was pre-registered (https://osf.io/n5c8g/) before analysing any data. Recently, this prediction has been

independently supported in a study showing a correlation between NAR and IAF (Katyal et al., 2019).

Alpha activity is causal to BR dynamics. Furthermore, our hypothesis implies that alpha oscillatory

activity is causal to BR dynamics. Therefore, we anticipated that entraining relevant endogenous alpha

activity with rhythmic visual stimulation should cause changes in perceptual alternation rate. Based on

this, our second prediction is that entraining brain activity slightly below the individual's alpha rhythm

will result in the individual experiencing slower BR alternation rates, whereas entraining brain activity

slightly above the alpha rhythm will speed up alternation rates. Experiment 2 was designed to test this

second prediction: we measured the NAR in a BR protocol while the observers' brain activity was

entrained at 3Hz, IAF-2Hz, IAF or IAF+2Hz.

The effect of entrainment on BR dynamics is frequency specific and limited to periodic stimulation. One

further implication of the hypothesis is frequency specificity. That is, only oscillatory frequencies

neighbouring the alpha range are relevant to the re-evaluation of competing stimuli, and therefore

entraining at rhythms outside the alpha range will not affect BR dynamics. Following this frequency

specificity assumption, we predict that entraining brain activity at a frequency far from the alpha range

(e.g., 3 Hz) will not impact competition dynamics and, therefore, the alternation rates under non-alpha

rhythmic entrainment will be equivalent to those without visual entrainment. Finally, according to our

hypothesis, the periodicity of neural fluctuations is consequential for BR dynamics. That is, modulations

of BR dynamics imposed by entrainment depend on the temporal (rhythmic) structure of the stimulation

and cannot be explained solely by accumulated changes in the amount of energy (i.e. increases in

competing stimuli strength). Following this hypothesis, we assume that entraining with rhythmic (alpha)

activity will promote periodic sampling of the competing stimuli, compared to comparable amounts of

stimulation without a rhythmic structure. Hence, our fourth main prediction is that the NAR in BR,

observed under arrhythmic stimulation, will be slower than the NAR observed under rhythmic (alpha)

entrainment. A final prediction, following from the same hypothesis, is that entraining alpha activity

might increase the NAR as compared to no-entrainment. This is because alpha entraining can sustain

alpha activity over and above its naturally occurring ongoing dynamics, which is typically characterised

by intermittent bouts (Rusiniak et al., 2018). Experiment 3 was designed to test these two predictions: we

measured the NAR while observers were entrained with either 3Hz, rhythmic (IAF), comparable

arrhythmic trains, or not entrained at all. This experiment was registered prior to any data analysis. At the

time of the registration the data of 6 subjects had been collected (https://osf.io/w8m4q/) but none had

been analysed.

It should be emphasised that the aim of this study, outlined in these hypotheses, is merely to probe the

role of alpha oscillations in rivalry dynamics. We make no claim as to the underlying cause of these

fluctuations, which is a question beyond the ambitions of the present study. Regardless of their cause,

perceptual fluctuations are a phenomenological effect of BR attributed to competition dynamics, and we

set out to investigate the role of alpha oscillations in this visual competition. The mechanisms underlying

perceptual fluctuations during BR have been addressed in a parallel study currently in preparation (Drew

et al., 2019).

Sample size estimation

The sample size for each of the three experiments was calculated to attain a 95% statistical power in the comparison of interest: IAF-NAR correlation in Experiment 1, IAF-2Hz compared to IAF+2Hz in Experiment 2, and IAF compared to arrhythmic stimulation in Experiment 3.

In Experiment 1, we used data from a previous experiment (Pápai et al., 2018) to estimate the sample size by means of a Monte Carlo randomisation procedure. We sampled with repetition data for sample sizes from 5 to 30 subjects (1000 repetitions). We approximated the statistical power as the proportion of significant correlations observed for a given sample size. The minimum required sample size to obtain a significant true effect 95% of the time for the effect size in (Pápai et al., 2018) was 25 subjects.

In Experiment 2, we used data from Experiment 1 to estimate the effect size of the condition of interest (IAF-2Hz vs IAF+2Hz). First, we estimated that the NAR changed 0.0195 Hz per each Hz in IAF. Consequently, we expected a difference of $\Delta NAR = 0.078$ Hz between slow and fast conditions in Experiment 2. We also assumed that the standard deviation of both slow and fast NAR in Experiment 2 would be equal to the standard deviation in NAR measured in Experiment 1 ($\sigma_{NAR} = 0.069$). Then, we used G*Power (RRID: SCR_013726) to calculate the effect size of the difference between matched pairs, assuming uncorrelated populations ($\rho = 0$, which would result in smaller effect sizes).

$$d = \frac{\Delta NAR}{\sigma}, \text{ where } \sigma = \sigma_{NAR}\sqrt{2 \cdot (1 - \rho)}$$

The estimated effect size was d=0.799. Given this effect size d=0.799, the statistical power for a one tailed t-test of two means (matched pairs) is above 95% for 20 subjects, but we selected a sample size of 25 to be able to capture smaller effect sizes (down to d=0.68).

In Experiment 3, there is no previous data comparing rhythmic (IAF) vs. arrhythmic entrainers. In order to have an intuition about the effect size we could expect, we based our calculation on the empirical effect sizes observed in Experiment 2 (N=25), that ranged between d=0.39 (fast vs. slow alpha entrainment) and d=1.88 for the contrast between 3 Hz and IAF+2Hz. We used G*Power (RRID: SCR_013726) to calculate the sample size required to observe a medium effect (d>0.5) with 95% power. We selected 30 participants, which provided 95% statistical power for effects d≥0.62.

Behavioural analysis

Percepts shorter than 300 ms (estimated latency of the motor evoked potential from key presses (Halgren, 1990)) were filtered out from the analysis. The NAR was obtained as the inverse of the median time between reporting two different pure percepts (complete transitions from red to green or green to red) (Fig. 1).

EEG recording

EEG data were acquired using active electrodes (actiCAP, BrainProducts GmbH, Munich, Germany) with Brain Vision Recorder (BrainProducts GmbH, Munich, Germany) at a sampling rate of 500 Hz. The ground electrode was placed at AFz and online reference on the tip of the nose. Four external electrodes were used: left and right mastoid for offline re-referencing and horizontal and vertical electrooculogram for monitoring eye movements and blinks. Impedance was kept below 10 kΩ. In Experiment 1, 64 electrodes placed according to the 10-10 system were used for the recording. In Experiments 2 and 3, we recorded from a subset of 21 electrodes. The ROI (parieto-occipital electrodes), defined a priori, consisted of a cluster of 8 electrodes: PO9, PO3, POz, PO4, PO10, O1, Oz, O2 in Experiment 1, PO7, PO3, POz, PO4, PO8, O1, Oz, O2, in Experiments 2 and 3.

EEG preprocessing and analyses

Pre-processing. EEG preprocessing and analysis was performed using Matlab's toolbox Fieldtrip (RRID: SCR_004849) and custom-written code. The EEG data was band-pass filtered 1-70 Hz (Butterworth filter, order 2, two-pass); a notch filter was used to remove the 50 Hz line-noise. The data recorded for IAF calculation with eyes closed was processed without further analysis. The EEG data recorded during task performance was further visually inspected to mark visual or muscular artefacts. For the IAF measurement during the task (Experiments 1 and 3), data fragments containing artefacts were discarded from the analysis; for the evoked activity measurement (Experiments 2 and 3), it was not (please note that only participants with a very low percentage of artefacts (<10%) were included in the final sample). All analyses were performed in the parieto-occipital region of interest only.

IAF estimation. The IAF at rest was measured from continuous EEG recordings (300 s) using the Welch method (1 second segments, 10% overlap, 0.25 to 40 Hz in steps of 0.25 Hz). The power spectrum was

averaged across the electrodes of interest (see ROI above) and background noise was estimated by fitting a $1/f^{\alpha}$ curve to the power spectrum (Haegens et al., 2014). IAF at rest was used only as a screening: participants presenting a single peak in the spectrum exceeding background noise in the 5 to 15 Hz range were accepted to proceed with the experiment. The calculation of IAF during task was performed in the 0 Hz (no entrainment) condition (Experiments 1 and 3). Data were segmented into epochs according to participants' reports (green or red percept, between key presses to avoid motor contamination). Pre-processed epochs were subsequently divided into non-overlapping segments of 1 s duration , demeaned and zero-padded to 10 s (Haegens et al., 2014). The power spectrum was calculated using a STFT (Hanning window, 4 to 30 Hz in steps of 0.1 Hz) and background noise correction using a $1/f^{\alpha}$ fit was performed as in the estimation of IAF at rest (see above).

SSVEP and phase-locking measures. In order to determine that endogenous activity was indeed entrained during our experiments we performed several tests: we calculated the power spectrum of steady-state visual evoked potentials (SSVEP) for all the rhythmic conditions (Experiments 2 and 3) to test that we observed a peak of activity at the entrained frequencies. We also verified the degree of phase locking of neural signal to external stimulation.

SSVEPs were obtained for each of the rhythmic entrainment conditions by aligning the 5 blocks of each entrainment condition (four conditions in Experiment 2, and two in Experiment 3) to stimulus presentation. SSVEPs were then averaged across the electrodes of the ROI and the power spectrum was calculated with the same method described above for the IAF at rest, but using a window length of 2 seconds (in order to capture at least 5 cycles of the slowest entrainment activity, 3 Hz, in each of the windows of analysis). We normalised frequencies relative to the IAF prior to averaging the power spectrum across participants.

To calculate phase locking we used cross-coherence (XCOH) (Keitel et al., 2017) to assess the extent to which neural activity was synchronised to rhythmic stimulation. We calculated XCOH between external rhythmic stimulation (train of Gaussian contrast increments) and each of the channels of the ROI for each of the blocks (we discarded the initial and final 5 seconds of each block, resulting in 110 s of continuous data). We used 1 second non-overlapping segments (Hanning window) and 2048 points to calculate the FFT. Coherence was averaged across trials and electrodes of the ROI for each participant and

experimental condition. Frequencies were normalised to IAF cycles and, finally, the value of coherence at

each of the entraining frequencies of interest was selected and averaged across participants.

For the IAF condition in Experiment 3, we applied an additional entropy analysis adapted from Notbohm

et al. (Notbohm et al., 2016), in order to show that rhythmic stimulation results in a stronger phase

synchrony than arrhythmic stimulation. The time series of visual stimulation and concurrently recorded

EEG were band-passed around the frequency of stimulation (see above) for rhythmic and arrhythmic

conditions. Segments of data containing artefacts were discarded. For clean data, we obtained the phase

difference between visual stimulation and concurrent EEG and calculated modified entropy as described

by Notbohm et al. (Notbohm et al., 2016), using N=80 bins for entropy calculation. In case of

entrainment, modified entropy would be larger for the rhythmic condition compared to the arrhythmic

condition.

Pre-registered statistical tests

For all statistical tests, the alpha level was set to 0.05. The correlation between the NAR and IAF was

tested by means of a Pearson correlation (right tailed). The impact of alpha speed on the NAR was tested

by means of a paired t-test (left tailed) for IAF-2Hz compared to IAF+2Hz. A one tailed t-test (right) was

used to test the hypothesis that IAF entrainment results in a faster NAR than arrhythmic entrainment.

A two one sided tests (TOST) (Lakens, 2017) was used to assess frequency specificity (following the

prediction that only entrainment at or around alpha impacts BR dynamics). A TOST is significant when

both p-values for left and right differences are below the set significance alpha level and the CI is

contained within the pre-defined equivalence interval. For this test, the NAR at 3 Hz was normalised with

respect to the 0 Hz NAR, and the equivalence interval was set to [-10 10] % before collecting data for

Experiment 3 (https://osf.io/w8m4q/) based on simulations. In these simulations, we estimated the

magnitude of the differences expected between NAR values from a single population due to sampling

error using data from Experiment 1 (27 subjects, 9 blocks per subject) to randomly generate two datasets

(A and B) per subject (4 blocks per dataset). In order to reduce variability between participants, the NAR

of each participant's datasets A and B was normalised relative to the NAR of dataset A. We then

performed a permutation test, in which we sampled 30 participants from the sample (the sample size

selected for experiment 3) with repetition. We performed 1000 random samplings and calculated the

proportion of successful TOST at different equivalence intervals: 0 to 20%, in steps of 1%. We selected a

10% equivalence interval, as the statistical power for a sample size of 30 subjects was above 95%.

Exploratory and control statistical tests (not pre-registered analyses)

In order to extend the evaluation of the potential correlation between IAF and NAR across different

datasets, the p-values from correlations in Experiments 1 and 3 were combined by means of the Stouffer

method (Vanrullen, 2016).

The entrainment measure for Experiments 1 and 3 (EEG phase-alignment to external stimulation) was

assessed by testing the significance of XCOH peaks at each of the entrainment frequencies. We applied

the statistical test described by Biltoft et al (Biltoft and Pardyjak, 2009) using 2048 points for FFT

estimation, 5 blocks of data, and 110 seconds of duration each at 500 Hz sampling rate, which resulted in

147 degrees of freedom. In this range of parameters, any coherence peak larger than 0.0405 would be

significant.

Lastly, the comparison regarding neural signal synchrony between rhythmic and arrhythmic stimulation

conditions in Experiment 3 was assessed by means of a paired one-tailed Wilcoxon test (Fay and

Proschan, 2010).

Results

According to behavioural responses in the BR task, the proportion of time spent in each of the two rival

percepts was balanced in all three experiments. On average, participants reported a red percept 44±8% of

the total recorded time, green 50±7% and no percept 6±6%, with similar proportions for all experimental

conditions. For the calculation of the NAR, we used, on average, 231±71 percepts per condition for

Experiment 1, 179±66 for Experiment 2 and 171±85 for Experiment 3.

Although IAF at rest can sometimes differ from IAF measured during task (Haegens et al., 2014), we did

not observe these differences here for Experiment 1 (IAF_{rest} − IAF_{task}=0.14±1.1 Hz, t(26)=0.77, p-

value=0.4) nor for Experiment 3 (IAF_{rest}-IAF_{task}=0.14±0.9 Hz, t(29)=1.30,p-value=0.2). The differences

were evaluated by means of a paired t-test (two-tailed). In Experiment 2, as all conditions included

entrainment, IAF could not be measured during the task without interference and this test was not

performed.

The speed of endogenous alpha activity correlates with the speed of perceptual alternations

In line with our first prediction, in Experiment 1 we observed a significant correlation between IAF

during the task and NAR (Fig. 3) (r=0.35, p-value=0.037). Visual inspection of the correlation plot

opened the question of whether significance might be driven by two potential outliers with low IAF

frequencies. From a electrophysiological point of view, the IAFs of these two subjects fall within the

usual range of IAFs reported in the literature, 7 to 14 Hz (Haegens et al., 2011; Smulders et al., 2018),

and therefore provide informative points in the analysis. Nevertheless, according to the Matlab toolbox

Robust Correlation (Pernet et al., 2013), these two points were classified as bivariate outliers. In order to

discard that the significant correlation observed in the main analysis was due to outliers, we calculated the

robust correlation coefficient with the above-mentioned toolbox, following the recommendations of

Pernet et al (Pernet et al., 2013). Data were distributed normally (Henze-Zirkler normality test p-value =

0.21) and included two bivariate outliers. We used the skipped Pearson correlation, which estimates the

strength of the linear correlation ignoring outliers but taking into account the overall structure of the data,

as recommended for this situation (Pernet et al., 2013). The correlation parameters obtained were in line

with those obtained with the main (pre-registered) analysis (r=0.35, p-value=0.037).

Note that Experiment 3, which was designed to test the effect of periodicity and frequency specificity,

also contained a no-stimulation condition, and therefore afforded an estimation of the correlation between

IAF and NAR, but with substantially fewer trials. Here, we observed a correlation in the expected

direction but that was not significant (r=0.19, p-value=0.15), perhaps not surprisingly given the reduced

statistical power. As the dataset in Experiment 1, data in Experiment 3 were distributed normally (p-

values for the Henze-Zirkler normality test p-value = 0.21) and included six bivariate outliers. The

skipped Pearson correlation returned a similar estimate (r=0.23 p-value=0.11 for Experiment 3).

Taken individually, the evidence for the correlation between IAF and NAR supports the initial

hypothesis, but it is not particularly strong. In order to garner more support, we used combined statistics

and meta-analysis, in post-hoc confirmatory analyses. First, we found that the combined statistics of

Experiment 1 and 3 (Stouffer method) rendered a significant effect (p-value=0.024). In addition, we

performed a meta-analysis using the four independent datasets that, as far as we know, have tested IAF-NAR correlation: (Katyal et al., 2019), data from a previous experiment of our group (Pápai et al., 2018), and data from Experiments 1 and 3 in the current study. The meta-analysis was performed with the R package "meta", using Fisher's z transformation of correlations. In order to homogenise datasets between studies we transformed all behavioural data to natural alternation times (NAT=1/NAR), the method used by Katyal et al. (please note that contrary to the NAR-IAF correlation, the expected direction for the NAT-IAF correlation is negative). For the same reason, we used the mean of the percept duration distributions instead of the median. This meta-analysis provided further confirmation of the correlation between BR dynamics and endogenous IAF activity (r=-0.44, p-value<0.001). Lastly, in order to take into account publication bias in the meta-analysis, we used the p-uniform method (van Assen et al., 2014). This method assesses the distribution of the sum of independent uniformly distributed random variables using the p-method (Irwin-Hall distribution, as implemented in the p-uniform package for R https://cran.r-project.org/web/packages/puniform/). After correcting for publication bias, the correlation was, once more, confirmed (r=-0.47, p-value=0.011). Altogether, the data supports our first hypothesis regarding the prediction that alpha frequency is correlated with the natural alternation between percepts, across individuals.

---- FIGURE 3 SHOULD BE PLACED HERE ---

The frequency of endogenous activity is causally related to the rate of alternations in conscious perception

In order to make sure that rhythmic visual stimulation successfully entrained endogenous oscillatory activity, in Experiment 2, we evaluated SSVEP and cross coherence (XCOH) at group level. The group-averaged SSVEP displayed a peak in the corresponding frequency of each of the entrainment conditions (Fig. 4. A), and the cross-coherence measure presented a significant peak (3 Hz: XCOH=0.1097, p-value=0.0002; IAF-2Hz: XCOH=0.135, p-value=0.00003; IAF: XCOH=0.0841, p-value=0.0017; IAF+2Hz: XCOH=0.1133, p-value=0.0002) (Fig. 4. B and C).

---- FIGURE 4 SHOULD BE PLACED HERE ---

In Experiment 2, the NAR at 3 Hz was slower than any of the other conditions (Fig. 5. A and Table 1). In

the condition of interest, as expected, we observed a significant decrease in the rate of perceptual

alternations during the slow alpha entrainment condition (IAF-2Hz) with respect to the fast alpha

entrainment condition (IAF+2Hz) (p-value=0.03, t(24)=-1.942, d=0.39) (Fig. 5. C). Entraining at the IAF

produced NARs between the slow IAF-2Hz and the fast IAF+2Hz entrainment, as one would expect, but

it was not significantly different from either of the two (Table 1). This finding confirms our second

prediction, regarding the causal modulation of perceptual alternations by visual entrainment.

---- FIGURE 5 SHOULD BE PLACED HERE --

Rhythmic entrainment boosts the re-evaluation of competing stimuli

In Experiment 3, group averaged SSVEPs displayed peaks at each of the entraining conditions (Fig. 6. A).

Endogenous activity was successfully entrained, as confirmed by the significant cross-coherence in all

experimental conditions (3 Hz: XCOH=0.1251, p-value=0.0001; IAF: XCOH=0.0711, p-value=0.0048)

(Fig. 6. B and C) and the higher values of normalised entropy observed in the IAF compared to arrIAF

condition (p-value=0.000001) (Fig. 6. D). As would be expected, the cross-coherence for the arrhythmic

condition was not significant (Fig. 6. B and C) (arrIAF: XCOH=0.016 p-value=0.32).

The comparison of interest here was between the rate of perceptual alternations during rhythmic (at IAF)

and arrhythmic (arrIAF) entrainment. In the arrhythmic condition, intervals between pulses were

randomly picked from an exponential distribution of latencies centred at the inverse of IAF. IAF

entrainment resulted in significantly faster NAR than arrhythmic entrainment (p-value=0.0005,

t(29)=3.659, d=0.67) (Fig. 5. C), indicating that periodicity plays a key role in BR dynamics. This

confirms the third main prediction, regarding periodicity.

---- FIGURE 6 SHOULD BE PLACED HERE --

In other words, although arrhythmic and rhythmic alpha entrainment always resulted in faster perceptual

alternation rates compared to low-frequency (3 Hz) or no visual entrainment (0 Hz) (Fig. 5. B and

Table1), the increase in alternation rate in IAF entrainment cannot be solely attributed to an increase in

the number or rate of pulses per unit of time (Brascamp et al., 2015). Instead, a mechanism involving

periodicity is required to fully explain the result.

Modulations in perceptual alternation rate are specific to entrainment in the alpha band

Further confirming the frequency-specificity implication of the hypothesis, we observed that the speeding up of perceptual alternations was constrained to the alpha band. Conditions that did not involve stimulation in the spectral vicinity of alpha always produced slower perceptual alternation rates. What is more, 3 Hz entrainment produced an alternation rate equivalent to no stimulation at all, as measured with two one sided tests (TOST (Lakens, 2017)) for an equivalence interval of 10% (p-value$_{left}$=0.007, p-value$_{right}$=0.0001, CI= [-0.07 0.03]) (Fig. 5. C).

Reality check: Coefficient of variation

In Experiment 3, we performed a post-hoc analysis on the coefficient of variation for each of the entraining conditions. The CV is the ratio between the standard deviation of perceptual alternations and mean alternation time, and it is informative about the interplay between adaptation and noise: values close to 0 indicate that adaptation dominates BR dynamics, whereas values closer to 1 indicate that the noise is driving the competition. We used a TOST (Lakens, 2017) to ascertain whether entrainment conditions would result in a different balance between adaptation and competition, compared to no entrainment (0 Hz). The equivalence interval for the CV was determined with a statistical power above 95% for a sample size of 30 subjects. No significant differences were seen between CV at 0 Hz stimulation and the rest of conditions based on the equivalence interval derived from Experiment 1 (22%), (0 Hz - 3Hz: p$_{left}$=0.00004, p$_{right}$=0.01, CI= [-0.05 0.17], 0 Hz- IAF: p$_{left}$=0.03, p$_{right}$=0.003, CI= [-0.19 0.11], 0 Hz- arrIAF: p$_{left}$=0.009, p$_{right}$=0.0015, CI= [-0.16 0.10]). This result suggests that the balance between noise and endogenous contributions to BR dynamics was equivalent across conditions.

Exploratory analysis: correlation between alpha power fluctuations and percept duration

The series of experiments reported in this manuscript was specifically designed to test how the speed of an individual's alpha activity (e.g., the frequency peak) related to their perceptual alternations. However, it is possible that an effect of oscillatory power could coexist with the frequency effect we have observed. In particular, we believe that the balance in alpha-gated selective inhibition (Klimesch, 2012) of the neural populations responding to each of the stimuli is the most likely candidate to account for a power-percept correlation. Strong inhibition (high alpha power) in the suppressed population will result in a

decreased probability of perceptual switch, whereas strong inhibition of the dominant population will lead

to increased probability of perceptual switch. Unfortunately, we cannot track percept-selective power

with our experimental design via frequency tagging of each eye (Jamison et al., 2015) or via topography.

Please note that frequency tagging studies have shown that the topographical distributions corresponding

to each eye are the same (Petruk et al., 2018).

However, in our study we can measure overall power in non-entrainment conditions. In order to explore if

overall alpha power fluctuations may have had an impact on percept duration in our study, we compared

alpha power for long and short percepts. This was done for Experiment 1, with more data available in the

no entrainment condition. In order to avoid motor contamination, we discarded data segments from -500

ms to +300 ms with respect to key presses. From the remaining data, we then retained epochs that were

300 ms or longer for analysis. These data epochs were binned in terciles according to percept duration,

and each epoch divided into 300 ms segments without overlap. The power of each segment was estimated

by means of a STFT, Hanning taper, in the electrodes of the ROI. Power for long and short percepts was

transformed to dB (relative to the power of all segments), and compared using a 2-tailed paired t-test (p-

value=0.53 t(26)=0.627). Given the different number of segments between long and short percepts,

(812±300 vs 140 ±70), we repeated the test sampling segments from the most populated condition via

bootstraping (75% segments of short percepts, 1000 iterations) to equate the number of segments

(104±50). The power difference between long and short percepts was still non-significant (p-value=0.47,

t(26)=0.734).

This null result does not preclude the possibility that alpha power fluctuations have an impact on

alternation rate, since this study was not intended to measure such effects.

<u>Discussion</u>

We addressed the hypothesis that alpha fluctuations in the occipital cortex mediate the accumulation of

visual information, leading to perceptual representations. Our findings, so far, seem to stand in good

agreement with this hypothesis. First, we confirmed that the speed of fluctuations in cortical alpha activity

correlates with the dynamics of perceptual fluctuations in BR, a result also recently observed in another

independent study (Katyal et al., 2019). We further proposed that endogenous alpha rhythm provides

temporal windows for the re-evaluation of competing visual representations based on visual input.

Consequently, inducing faster endogenous activity should accelerate perceptual switches, a result we

observed in Experiment 2. Moreover, our results suggest a causal relationship between alpha activity and

awareness dynamics, since modulation of endogenous brain rhythms resulted in consequential changes in

conscious perception. That is, speeding up or slowing down alpha cortical activity with entrainment alters

the rate of perceptual alternations accordingly. Critically, this causal modulation is restricted to the alpha

range and its effectiveness is subject to the periodicity of stimulation.

These findings are consistent with the pacemaker role of alpha activity in perceptual alternations, which

we propose here. We build from two well-known principles. First, that reciprocal inhibition between the

neural populations selective of the competing stimuli is the mechanism causing perceptual alternations in

BR (Tong et al., 2006). Second, that the activation of the neural population selective of the dominant

conscious percept tends to decrease over time due to adaptation (Shpiro et al., 2009), making perceptual

switches more likely with time. In this context, inhibition plays an active role: by stifling the neural

population with lower excitability, it increases the signal to noise ratio corresponding to the alternative

population (Klimesch, 2012). We hypothesise that this process takes place in a phasic manner: each alpha

cycle opens a window for re-evaluation of the competing neural signals. In each of these windows, the

weight of each neural population is evaluated, and the outcome of each evaluation determines the

resulting conscious percept. Successive alpha cycles lead to the progressive accumulation of evidence

toward one or the other percept.

Could this accumulation happen with changes in excitability regardless of rhythmic fluctuations?

(Brascamp et al., 2015) In Experiment 3, we tested whether local, non-rhythmic changes in stimulus

contrast could account for the increase in perceptual alternation rate during entrainment. Although

arrhythmic stimulation increased alternation rates with respect to no-stimulation, these changes could not

account for the effect in full. Rhythmic IAF stimulation resulted in faster alternations than arrhythmic

stimulation. Additionally, if local changes in contrast had been responsible for speeding up switch

dynamics, a difference between the no-stimulation condition and 3 Hz stimulation condition would have

been expected. Yet, these two conditions rendered equivalent, and low, alternation rates.

Our manipulation could have affected the interplay between noise and adaptation, reflected in the

coefficient of variation. Studies using anti-phase stimulation have reported drops in the coefficient of

variation (Kim et al., 2006), implying a decrease in the contribution of noise to BR dynamics. In our

study, the analysis of the coefficient of variation implied that the visual stimulation protocol used here did

not modify the basic mechanism of neural competition. Therefore, one can assume that the competition

mechanism remained unchanged and that only the frequency and probability of windows favouring the

re-evaluation of perceptual representations were modified.

Sensory entrainment at or around the IAF produced a dramatic increase in perceptual alternations

compared to 3Hz or no-stimulation. That is, alpha entrainment produces a sizable effect, even if there is

already an ongoing rhythm at that frequency. What is the origin of this increase? In line with the well-

known fact that endogenous alpha occurs in bursts rather than in a sustained fashion (Rusiniak et al.,

2018), we speculate that alpha-gated re-evaluation mostly occurs within these bursts. That is, in BR the

re-evaluation of competing stimuli might not unfold homogenously over time, but intermittently.

Incidentally, this may lead to the stochastic dynamics that characterises the phenomenon. In our protocol,

entrainment at the natural alpha rhythm must have increased alpha burst episodes, explaining the faster

alternation rate under IAF, compared to no entrainment. During these alpha bursts, in order to reduce

stimulus complexity, information is compressed into discrete packages (Vanrullen and Koch, 2003). The

dynamics of the system alternate between intervals where the probability of switch is low and recurring

intervals of alpha activity where competition between conflicting representations is steep. Therefore, by

increasing alpha burst episodes, the probability of stimulus re-evaluation increases and, as a result, faster

alternation rates can be expected. The role of alpha in the gating by inhibition hypothesis (Jensen and

Mazaheri, 2010) is compatible with the hypothesis laid out here. According to the former, sensory

information is coded in gamma activity embedded within the high excitability windows of slower alpha

cycles. One could argue that, in fact, slower alpha cycles may allow for more nested gamma cycles, hence

lead to more accumulation of evidence per alpha cycle, predicting a result contrary to what we observed.

However, variations in the speed of gamma may compensate for the different number of cycles per

second across alpha speeds. Indeed, several findings suggest that the length of gamma cycles vary

between subjects, and subjects with faster alpha also present faster gamma (Muthukumaraswamy et al.,

2009; Baumgarten et al., 2018). We speculate that the number of gamma bursts per alpha cycle is

approximately constant, therefore faster alpha rhythms would allow for an increased number of gamma

bursts per second, in line with our hypothesis and compatible with gating by inhibition.

Regarding frequency specificity, we cannot exclude the possibility that the NAR would increase with

entrainment faster than alpha, as we did not stimulate at frequencies above the alpha range. However,

high frequency stimulation may not be an effective test for specificity, given that alpha activity could still

be entrained as a sub-harmonic of higher frequencies, as some studies suggest (Herrmann, 2001;

Herrmann et al., 2016). Instead, several independent results favour a frequency specificity interpretation.

From our study, we can conclude that there is a lower frequency bound of at least 3Hz, to modulate NAR

effectively. In addition, evidence from O'Shea and Crassini (O'shea and Crassini, 1984) provide indirect

support for the existence of a higher bound. O'Shea and Crassini presented flickering stimuli at different

frequencies during a BR protocol, and observed a pattern of NAR in line with our predictions: maximum

NARs were observed in the alpha range and, additionally, subjects reported that flickering at faster

frequencies (above the alpha range) disappeared, a result consistent with discrete sampling. However,

methodological differences (i.e., only 3 subjects were tested and stimuli were presented for on-off

periods) prevent a conclusive inference.

It is possible that the physical stimulation based on contrast changes used here resulted in a modulation of

attention, which has a well-known effect on BR (Zhang et al., 2011; Brascamp and Blake, 2012).

Compatible with our hypothesis, attention has been proposed to periodically sample rivalling stimuli

during BR (Davidson et al., 2018), and it is possible that the increase in NAR observed in the arrhythmic

stimulation condition is due to a global increase in arousal. Our findings can be framed in terms of

existing theoretical models of BR which incorporate attention (Li et al., 2017). According to these

models, attention recurrently (or rhythmically, in our hypothesis) regulates the imbalance between the two

rival stimuli with a slow time constant, triggered by a mutual inhibition process that has a fast(er) time

constant. We can interpret this model in terms of a slow rhythm (alpha) and a fast rhythm. This faster

rhythm could be gamma, which has also been found to correlate with BR dynamics in previous studies

(Fesi and Mendola, 2015). The concept of alpha discrete sampling can also be framed within Bayesian

models of BR (Hohwy et al., 2008; Gershman et al., 2012). It has been proposed that the continuous input

of inconsistent stimuli creates a situation in which neither of the stimuli representations has both high

prior and high likelihood. As a result, periodic sampling determines which of the two representations

sways the perceptual outcome at any given time (Hohwy et al., 2008).

The present findings provide a framework for the interpretation of inter-individual differences in BR

dynamics: in this framework, endogenous alpha cycles are a representative time constant for each subject

that regulates the competition process between neural populations representative of each stimuli. Alpha

acts as a pacemaker for information accumulation mediating the transition to awareness by setting the

tempo for the competition process. According to this interpretation, our results are in line with recent

studies suggesting that alpha activity may not regulate objective aspects of perception, such as perceptual

sensitivity, but awareness and confidence about perceptual properties (Lange et al., 2013; Benwell et al.,

2017; Samaha et al., 2017).

<u>Data availability</u>

The datasets generated during and/or analysed during the current study are available from the

corresponding author on reasonable request.

<u>References</u>

Baumgarten TJ, Neugebauer J, Oeltzschner G, Füllenbach ND, Kircheis G, Häussinger D, Lange J,
Wittsack HJ, Butz M, Schnitzler A (2018) Connecting occipital alpha band peak frequency, visual
temporal resolution, and occipital GABA levels in healthy participants and hepatic encephalopathy
patients. NeuroImage Clin 20:347–356 Available at: https://doi.org/10.1016/j.nicl.2018.08.013.
Benwell CSY, Tagliabue CF, Veniero D, Cecere R, Savazzi S, Thut G (2017) Prestimulus EEG Power
Predicts Conscious Awareness But Not Objective Visual Performance. Eneuro 4:ENEURO.0182-
17.2017.
Biltoft CA, Pardyjak ER (2009) Spectral coherence and the statistical significance of turbulent flux
computations. J Atmos Ocean Technol 26:403–410.
Block N (2019) What Is Wrong with the No-Report Paradigm and How to Fix It. Trends Cogn Sci
23:1003–1013 Available at: https://doi.org/10.1016/j.tics.2019.10.001.
Brascamp JW, Blake R (2012) Inattention Abolishes Binocular Rivalry: Perceptual Evidence. Psychol Sci
23:1159–1167.
Brascamp JW, Ee R Van, Pestman WR, Berg AV Van Den (2005) Distributions of alternation rates in
various forms of bistable perception. :287–298.
Brascamp JW, Klink PC, Levelt JM (2015) The ' laws ' of binocular rivalry : 50 years of Levelt ' s
propositions. 109:20–37.
Carter OL, Pettigrew JD (2003) A common oscillator for perceptual rivalries? Perception 32:295–305.
Cavanagh P, MacLeod DI, Anstis SM (1987) Equiluminance: spatial and temporal factors and the
contribution of blue-sensitive cones. J Opt Soc Am A, Opt Image sicence 4:1428–1438.
Cecere R, Rees G, Romei V (2015) Individual Differences in Alpha Frequency Drive Crossmodal
Illusory Perception. Curr Biol 25:231–235 Available at:
http://dx.doi.org/10.1016/j.cub.2014.11.034.
Clark CR, Veltmeyer MD, Hamilton RJ, Simms E, Paul R, Hermens D, Gordon E (2007) Spontaneous
alpha peak frequency predicts working memory performance across the age span. Int J
Psychophysiol 53:1–9.
Davidson MJ, Alais D, Boxtel JJA Van (2018) Attention periodically samples competing stimuli during
binocular rivalry. :1–25.
de Jong MC, Hendriks RJM, Vansteensel MJ, Raemaekers M, Verstraten FAJ, Ramsey NF, Erkelens CJ,
Leijten FSS, van Ee R (2016) Intracranial Recordings of Occipital Cortex Responses to Illusory
Visual Events. J Neurosci 36:6297–6311.
Doesburg SM, Kitajo CAK, Ward LM (2005) Increased gamma-band synchrony precedes switching of
conscious perceptual objects in binocular rivalry. 16:1–4.
Drew A, Torralba M, Ruzzoli M, Morís Fernández L, Sabaté A, Szabina Pápai M, Soto-Faraco S (2019)
Neural Evidence of Cognitive Conflict During Binocular Rivalry. SSRN Electron J:25–27.
Fay MP, Proschan MA (2010) Wilcoxon-Mann-whitney or t-test? Stat Surv 4:1–37.
Fesi JD, Mendola JD (2015) Individual Peak Gamma Frequency Predicts Switch Rate in Perceptual
Rivalry. Hum Brain Mapp 576:566–576.
Friston K (2012) Prediction, perception and agency. Int J Psychophysiol 83:248–252 Available at:
http://dx.doi.org/10.1016/j.ijpsycho.2011.11.014.

Gershman SJ, Vul E, Tenenbaum JB (2012) Multistability and perceptual inference. Neural Comput.

Grandy TH, Werkle-Bergner M, Chicherio C, Schmiedek F, Lövdén M, Lindenberger U (2013) Peak
individual alpha frequency qualifies as a stable neurophysiological trait marker in healthy younger
and older adults. Psychophysiology 50:570–582 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/23551082 [Accessed December 4, 2014].

Haegens S, Cousijn H, Wallis G, Harrison PJ, Nobre AC (2014) Inter- and intra-individual variability in
alpha peak frequency. Neuroimage 92:46–55 Available at:
http://dx.doi.org/10.1016/j.neuroimage.2014.01.049.

Haegens S, Nácher V, Luna R, Romo R, Jensen O (2011) α-Oscillations in the monkey sensorimotor
network influence discrimination performance by rhythmical inhibition of neuronal spiking. Proc
Natl Acad Sci U S A 108:19377–19382 Available at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3228466&tool=pmcentrez&rendertype
=abstract [Accessed August 20, 2014].

Halgren E (1990) Human Evoked Potentials. Neurophysiol Tech II:147–276.

Herrmann CS (2001) Human EEG responses to 1-100 Hz flicker: Resonance phenomena in visual cortex
and their potential correlation to cognitive phenomena. Exp Brain Res 137:346–353.

Herrmann CS, Murray MM, Ionta S, Hutt A, Lefebvre J (2016) Shaping intrinsic neural oscillations with
periodic stimulation. J Neurosci 36:5328–5337.

Hohwy J, Roepstorff A, Friston K (2008) Predictive coding explains binocular rivalry: An
epistemological review. Cognition 108:687–701.

Jensen O, Mazaheri A (2010) Shaping functional architecture by oscillatory alpha activity: gating by
inhibition. Front Hum Neurosci 4:186 Available at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2990626&tool=pmcentrez&rendertype
=abstract [Accessed July 11, 2014].

Kanai R, Moradi F, Shimojo S, Verstraten FAJ (2005) Perceptual alternation induced by visual transients.
Perception.

Kang M (2009) Size matters : A study of binocular rivalry dynamics. 9:1–11.

Katyal S, He S, He B, Engel SA (2019) Frequency of alpha oscillation predicts individual differences in
perceptual stability during binocular rivalry. Hum Brain Mapp:1–12.

Keitel C, Thut G, Gross J (2017) Visual cortex responses reflect temporal structure of continuous quasi-
rhythmic sensory stimulation. Neuroimage 146:58–70 Available at:
http://dx.doi.org/10.1016/j.neuroimage.2016.11.043.

Kim YJ, Grabowecky M, Suzuki S (2006) Stochastic resonance in binocular rivalry. Vision Res 46:392–
406.

Kleinschmidt A, Sterzer P, Rees G (2012) Variability of perceptual multistability: From brain state to
individual trait. Philos Trans R Soc B Biol Sci.

Klimesch W (2012) α-band oscillations, attention, and controlled access to stored information. Trends
Cogn Sci 16:606–617 Available at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3507158&tool=pmcentrez&rendertype
=abstract [Accessed July 10, 2014].

Lakens D (2017) Equivalence Tests: A Practical Primer for t Tests, Correlations, and Meta-Analyses. Soc
Psychol Personal Sci 8:355–362.

Lange J, Oostenveld R, Fries P (2013) Reduced occipital alpha power indexes enhanced excitability
rather than improved visual perception. J Neurosci 33:3212–3220 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/23407974 [Accessed December 3, 2014].

Lee SH, Blake R, Heeger DJ (2007) Hierarchy of cortical responses underlying binocular rivalry. Nat
Neurosci 10:1048–1054.

Leopold DA, Logothetis NK (1999) Multistable phenomena : changing views in perception. 6613:254–
264.

Li H, Rankin J, Rinzel J, Carrasco M, Heeger DJ (2017) Attention model of binocular rivalry.

Mamassian P, Goutcher R (2005) Temporal dynamics in bistable perception. J Vis 5:7.

Miller SM, Hansell NK, Ngo TT, Liu GB, Pettigrew JD, Martin NG, Wright MJ (2010) Genetic
contribution to individual variation in binocular rivalry rate. 107:9–13.

Muthukumaraswamy SD, Edden RAE, Jones DK, Swettenham JB, Singh KD (2009) Resting GABA
concentration predicts peak gamma frequency and fMRI amplitude in response to visual stimulation
in humans. Proc Natl Acad Sci U S A 106:8356–8361.

Notbohm A, Kurths J, Herrmann CS (2016) Modification of Brain Oscillations via Rhythmic Light
Stimulation Provides Evidence for Entrainment but Not for Superposition of Event-Related
Responses. 10.

O'shea RP, Crassini B (1984) Binocular rivalry occurs without simultaneous presentation of rival stimuli.

Percept Psychophys 36:266–276.
Pápai MS, Cuello MT, Bayer M, Soto-faraco S (2018) Occipital Alpha modulates the processing of
suppressed visual stimuli during binocular rivalry. Available at:
http://www.ub.edu/bridgingattentionprediction/content/occipital-alpha-modulates-processing-
suppressed-visual-stimuli-during-binocular-rivalry.
Pernet CR, Wilcox R, Rousselet GA (2013) Robust correlation analyses: False positive and power
validation using a new open source matlab toolbox. Front Psychol 3:1–18.
Phillips I, Morales J (2020) The Fundamental Problem with No-Cognition Paradigms. Trends Cogn Sci
24:165–167 Available at: https://doi.org/10.1016/j.tics.2019.11.010.
Piantoni G, Kline KA, Eagleman DM (2010) Beta oscillations correlate with the probability of perceiving
rivalrous visual stimuli. 13:1–11.
Polonsky A, Blake R, Braun J, Heeger DJ (2000) Neuronal activity in human primary visual cortex
correlates with perception during binocular rivalry. :1153–1159.
Ronconi L, Busch NA, Melcher D (2018) Alpha-band sensory entrainment alters the duration of temporal
windows in visual perception. Sci Rep 8:1–10 Available at: http://dx.doi.org/10.1038/s41598-018-
29671-5.
Rusiniak M, Wróbel A, Cieśla K, Pluta A, Lewandowska M, Wójcik J, Skarżyński PH, Wolak T (2018)
The relationship between alpha burst activity and the default mode network. Acta Neurobiol
Exp:92–106.
Samaha J, Iemi L, Postle BR (2017) Prestimulus alpha-band power biases visual discrimination
confidence, but not accuracy. Conscious Cogn 54:47–55 Available at:
http://dx.doi.org/10.1016/j.concog.2017.02.005.
Samaha J, Postle BR (2015) The Speed of Alpha-Band Oscillations Predicts the Temporal Resolution of
Visual Perception Report The Speed of Alpha-Band Oscillations Predicts the Temporal Resolution
of Visual Perception. Curr Biol 25:2985–2990 Available at:
http://dx.doi.org/10.1016/j.cub.2015.10.007.
Schmack K, Sekutowicz M, Rössler H, Brandl EJ, Müller DJ, Sterzer P (2013) The influence of
dopamine-related genes on perceptual stability. Eur J Neurosci.
Shpiro A, Moreno-Bote R, Rubin N (2009) Balance between noise and adaptation in competition models
of perceptual bistability. J Comput Neurosci 27:37–54.
Smulders FTY, ten Oever S, Donkers FCL, Quaedflieg CWEM, van de Ven V (2018) Single-trial log
transformation is optimal in frequency analysis of resting EEG alpha. Eur J Neurosci 48:2585–
2598.
Sobel K V., Blake R (2002) How context influences predominance during binocular rivalry. Perception.
Sterzer P, Kleinschmidt A, Rees G (2009) The neural bases of multistable perception. Trends Cogn Sci
13:310–318 Available at: https://doi.org/10.1016/j.tics.2009.04.006.
van Assen MALM, van Aert RCM, Wicherts JM (2014) Meta-Analysis Using Effect Size Distributions of
Only Statistically. Pychological Methods.
Vanrullen R (2016) How to evaluate phase differences between trial groups in ongoing
electrophysiological signals. :1–24.
Vanrullen R, Busch NA, Drewes J, Dubois J (2011) Ongoing EEG phase as a trial-by-trial predictor of
perceptual and attentional variability. Front Psychol 2:1–9.
Vanrullen R, Koch C (2003) Is perception discrete or continuous ? Trends Cogn Sci 7:207–213.
Zhang P, Jamison K, Engel S, He B, He S (2011) Binocular Rivalry Requires Visual Attention. Neuron
71:362–369 Available at: http://dx.doi.org/10.1016/j.neuron.2011.05.035.
727
728

Figure 1: A: Example of the Gabor patches used in the experimental protocol. B: Contrast of both Gabor patches was modulated simultaneously with different temporal patterns (low and high frequency rhythmic modulation, fast arrhythmic modulation). C: Participants continuously reported the conscious percept (red, green, mixed or none) while EEG was concurrently recorded. D: Image of our experimental setup. E: Target behavioural measure was individual characteristic duration of conscious percept (NAR). F: Behavioural measures were complemented with endogenous oscillatory activity recorded during the BR task.

Figure 2: A: Experimental procedure. All sessions started with a 5 minute recording during rest (eyes

closed) used to estimate IAF at rest. Next, the RGB level of the green stimulus was adjusted individually

and the stereoscope mirrors were calibrated. Prior to the experiment, two training blocks of a 2 minute

duration were run (0 Hz for Experiment 1, random conditions for Experiment 2 and 3). Finally,

participants ran the experimental blocks. B: In Experiment 1, participants ran 2 blocks of the 0 Hz

condition followed by 1 block of Replay (a condition neither used nor relevant for the current experiment)

until 9 blocks of 0 Hz were completed. C: In Experiment 2, participants performed 5 runs including a

block of each of the 4 experimental conditions: 3 Hz, IAF-2Hz, IAF and IAF+2Hz. The order was

assigned randomly, with the constraint that the same condition could not be presented in consecutive

blocks. D: In Experiment 3, the procedure was exactly the same as in Experiment 2 but the conditions

tested were 0 Hz, 3 Hz, IAF and arrIAF.

Figure 3: Natural alternation rate as a function of IAF during task for Experiments 1 (red) and 3 (green).

Dotted lines correspond to the linear fit of NAR as a function of IAF.

Figure 4: Entrainment in Experiment 2. A: Group average SSVEP. The frequencies are normalised with

respect to rest IAF (the one used during IAF entrainment) and 1/f background noise correction was

applied prior to the group averaging. Vertical dotted lines represent the target frequency at each condition

(only IAF is exactly equal to one in the normalised plot, shaded areas indicate the lower and upper bounds

for 3 Hz, IAF-2Hz and IAF+2Hz normalised values for each participant). B: Cross-coherence between the

EEG signal and the rhythmic entrainment at the frequencies corresponding to each entraining condition.

Dots represent individual values, black lines are the mean value for each condition and the dark shaded

area corresponds to the standard error of the mean. The red dotted line presents the value of the coherence

at a significance level 0.05. C: Cross-coherence spectrum for each of the experimental conditions.

768 **Figure 5:** Effect of entrainment on perceptual alternations. A: NAR for Experiment 2 at the different

769 experimental conditions. B: NAR for Experiment 3. C: NAR differences in the conditions of interest

770 (IAF-2 Hz compared to IAF +2 Hz, 0 Hz compared to 3 Hz and IAF compared to arrhythmic IAF). IAF-

771 2Hz was significantly slower than IAF+2 Hz, whereas 0 Hz and 3 Hz were equivalent (for a 10%

772 equivalence interval) and the IAF rhythmic condition was significantly faster than the arrhythmic

773 condition. Solid lines correspond to the group mean of each condition; dark shaded areas correspond to

774 the mean ± standard error of the mean. Light shaded areas correspond to the distribution of the individual

775 data and dots correspond to individual data points.

Figure 6: Entrainment in Experiment 3. A: Group average SSVEP. The frequencies are normalised with respect to rest IAF (the one used during IAF entrainment) and 1/f background noise correction was applied prior to the group averaging. Vertical dotted lines represent the target frequency at each condition (only IAF is exactly equal to one in the normalised plot, shaded area indicate the lower and upper bounds for 3 Hz, normalised values for each participant). B: Values of coherence between neural signal and rhythmic stimulation at the frequencies corresponding to each entraining condition. Dots represent individual values, black lines are the mean value for each condition and the dark shaded area corresponds to the standard error of the mean. The red dotted line presents the value of the cross-coherence at a significance level 0.05. C: Cross-coherence spectrum for each of the experimental conditions. D: Difference between normalised entropy at IAF and arrIAF conditions. Dots represent individual values, black thick line is the mean value and dark grey shaded area is the standard error of the mean.

Exp.	Cond. 1	Cond. 2	Tail	p-value	t-value	d.o.f.	d
2	IAF-2Hz	IAF	Left	0.08	-1.443	24	-0.29
2	IAF-2Hz	3 Hz	Both	$5 \cdot 10^{-8}$	7.801	24	1.56
2	IAF	3 Hz	Both	$7 \cdot 10^{-9}$	8.667	24	1.73
2	IAF	IAF+2Hz	Left	0.12	-1.223	24	-0.24
2	IAF+2Hz	3 Hz	Both	$1 \cdot 10^{-9}$	9.420	24	1.88
3	0 Hz	IAF	Both	$5 \cdot 10^{-10}$	9.108	29	1.66
3	0 Hz	arrIAF	Both	$6 \cdot 10^{-8}$	7.241	29	1.25
3	3 Hz	IAF	Both	$2 \cdot 10^{-10}$	9.420	29	171
3	3 Hz	arrIAF	Both	$5 \cdot 10^{-8}$	7.257	29	1.24

791   **Table 1:** Statistics for non-planned contrasts in Experiments 2 and 3. For all contrasts, we used paired t-

792   tests. In all contrasts but two, we used two tailed tests, as we did not have an a priori expectation about

793   the direction of the difference.

794   Figures