Effective connectivity and spatial selectivity-dependent fMRI changes elicited by microstimulation of pulvinar and LIP

Igor Kagan[1,2,3,*], Lydia Gibson[1,2,*], Elena Spanou[1], Melanie Wilke[1,2,3]

[1] *Decision and Awareness Group, Cognitive Neuroscience Laboratory, German Primate Center, Leibniz Institute for Primate Research, Kellnerweg 4, Goettingen, 37077, Germany*

[2] *Department of Cognitive Neurology, University of Goettingen, Robert-Koch-Str. 40, Goettingen, 37075, Germany*

[3] *Leibniz ScienceCampus Primate Cognition, Kellnerweg 4, Goettingen, 37077, Germany*

* Shared first authors

Running Head: Electrical microstimulation of pulvinar and LIP in behaving macaques

Corresponding author:
Igor Kagan ikagan@dpz.eu
German Primate Center, Leibniz Institute for Primate Research
Kellnerweg 4, Goettingen, 37077, Germany

Abstract

The thalamic pulvinar and the lateral intraparietal area (LIP) share reciprocal anatomical connections and are part of an extensive cortical and subcortical network involved in spatial attention and oculomotor processing. The goal of this study was to compare the effective connectivity of dorsal pulvinar (dPul) and LIP and to probe the dependency of microstimulation effects on task demands and spatial tuning properties of a given brain region. To this end, we applied unilateral electrical microstimulation in the dPul and LIP in combination with event-related BOLD fMRI in monkeys performing fixation and memory-guided saccade tasks. Microstimulation in both dPul and LIP enhanced task-related activity in monosynaptically-connected prefrontal cortex and along the superior temporal sulcus (STS) as well as in extrastriate cortex. Both dPul and LIP stimulation also elicited activity in several cortical areas in the opposite hemisphere, implying polysynaptic propagation of excitation. LIP microstimulation elicited strong activity in the opposite homotopic LIP while no homotopic activation was found during dPul stimulation. Despite extensive activation along the intraparietal sulcus evoked by LIP stimulation, there was a difference in frontal and occipital connectivity elicited by posterior and anterior LIP stimulation sites. Comparison of dPul stimulation with the adjacent but functionally distinct ventral pulvinar also showed distinct connectivity. On the level of single trial timecourses within a region, most microstimulation regions did not show task-dependence of stimulation-elicited response modulation. Across regions, however, there was an interaction between the task and the stimulation, and task-specific correlations between the initial spatial selectivity and the magnitude of stimulation effect were observed. Consequently, stimulation-elicited modulation of task-related activity was best fitted by an additive model scaled down by the initial response amplitude. In summary, we identified overlapping and distinct patterns of thalamocortical and corticocortical connectivity of the two key visuospatial areas, highlighting the dorsal bank and fundus of STS as a prominent node of shared circuitry. Spatial task-specific and partly polysynaptic modulations of cue and saccade planning delay period activity in both hemispheres exerted by unilateral pulvinar and parietal stimulation provide insight into the distributed interhemispheric processing underlying spatial behavior.

Keywords: memory saccades, eye movements, fixation, spatial selectivity, macaque monkeys

Highlights

- Electrical stimulation of pulvinar and LIP was used to study fMRI effective connectivity
- Both regions activated prefrontal cortex and the dorsal bank of superior temporal sulcus
- Activations within and across hemispheres suggest polysynaptic propagation
- Stimulation effects show interactions between task- and spatial selectivity
- Stimulation effects are best fitted by an additive model scaled by the initial response

1 Introduction

Spatial exploration and selection of visual stimuli for further analysis by goal-directed saccadic eye movements is an important feature of primate behavior. This behavior relies on a widely distributed subcortical-cortical brain circuitry including fronto-parietal cortices, superior colliculus (SC), basal ganglia and thalamus (Andersen and Cui, 2009; Glimcher, 2003; Gold and Shadlen, 2007; Wurtz et al., 2011; Wurtz and Hikosaka, 1986). While until recently most concepts of flexible visually-guided behavior emphasized interactions within cortical networks, direct connections between cortical regions are paralleled by indirect routes through higher-order thalamic nuclei such as pulvinar (dPul), raising the question what kind of information is contributed by the additional thalamic routes and how the processing compares between the functionally pertinent cortical and thalamic sites (Bridge et al., 2015; Grieve et al., 2000; Halassa and Kastner, 2017; Sherman and Guillery, 2011; Shipp, 2003). Two major nodes of the network that link vision with saccadic eye movements and where lesions lead to spatial neglect/extinction symptoms in monkeys and humans are the lateral intraparietal region and the dorsal pulvinar (Karnath et al., 2002; Mesulam, 1999; Storm et al., 2017; Wardak et al., 2002; M. Wilke et al., 2010). The causal influence of these two regions on interconnected brain circuitry is the focus of the current study.

In monkeys, the lateral intraparietal area (LIP) is located in the intraparietal sulcus (IPS) of the posterior parietal cortex (PPC), and its neurons encode the position and behavioral relevance of visual stimuli along with the direction of an upcoming saccade (Andersen et al., 1990; Barash et al., 1991; Bisley and Goldberg, 2010). Response fields in LIP are predominantly located in the contralateral hemifield (Ben Hamed et al., 2001; Blatt et al., 1990). Furthermore, LIP neurons exhibit delay period activity in memory-guided saccade tasks that reflects saccade preparation and the spatial allocation of attention (Andersen and Cui, 2009; Bisley and Goldberg, 2010; Colby and Goldberg, 1999). Area LIP is reciprocally interconnected with the pulvinar (Asanuma et al., 1985; Blatt et al., 1990; Hardy and Lynch, 1992; Romanski et al., 1997), which is a heterogeneous structure that has been subdivided in various manners based on histochemical, cytoarchitectonic or anatomical connectivity tracer techniques (Baldwin et al., 2017; Bridge et al., 2015; Gutierrez et al., 2000; Jones, 2007). Although not reflecting all detailed division schemes, several anatomical and functional studies have motivated a functional distinction between ventral and dorsal pulvinar portions (Arcaro et al., 2015; Kaas and Lyon, 2007; Komura et al., 2013; Stepniewska et al., 1994). The ventral pulvinar (vPul) encompasses the inferior pulvinar (PI) and the ventral part of the lateral pulvinar (PLvl), has strong reciprocal connections to visual cortices and contains several retinotopic maps with mainly contralateral visual receptive fields (Arcaro et al., 2015; Bender, 1981; Petersen et al., 1985). Ventral

pulvinar neurons are also modulated by visual attention (Bender and Youakim, 2001; Saalmann et al., 2012; Zhou et al., 2016) and show saccade-related activity that is mostly peri- or post-saccadic (Berman and Wurtz, 2011; Petersen et al., 1985; Robinson et al., 1986). The dorsal pulvinar (dPul) occupies the region above the level of the brachium of the superior colliculus and encompasses the medial pulvinar (PM) and the dorsal portion of the lateral pulvinar (PLdm) (Gutierrez et al., 2000; Kaas and Lyon, 2007). Similar to LIP and vPul, dPul neurons show enhancement for visual stimuli that are attended due to their behavioral relevance and/or indicate an upcoming saccade target (Bender and Youakim, 2001; Fiebelkorn et al., 2019) and discharge in cue and saccade execution phases with an overall contralateral preference (Benevento and Port, 1995; Dominguez-Vargas et al., 2017; Robinson et al., 1986; Schneider et al., 2020). But unlike vPul and to a certain extent LIP (Patel et al., 2010), dPul does not follow a clear retinotopic organization (Benevento and Miller, 1981; Benevento and Port, 1995; Petersen et al., 1987).

Providing causal evidence for involvement in spatial visuomotor behavior, reversible pharmacological inactivation in LIP or dPul leads to spatial neglect/extinction characterized by impaired responses to and exploration of the contralateral space (Christopoulos et al., 2018; Li et al., 1999; Liu et al., 2010; Petersen et al., 1987; Wardak et al., 2002, 2004; M. Wilke et al., 2010; Wilke et al., 2013). Particularly, employing visually-guided or memory-guided saccade tasks in monkeys, several studies have shown that local inactivation of LIP or the dorsal pulvinar leads to biased saccade choices towards the ipsilesional side while keeping saccade parameters largely intact, apart from a moderate increase of contralesional saccade latencies (Chen et al., 2016; Katz et al., 2016; Kubanek et al., 2015; Wardak et al., 2002; M. Wilke et al., 2010; Wilke et al., 2012, 2013). Similarly, electrical microstimulation in LIP or dorsal pulvinar induces a bias of perceptual and saccade choices towards contraversive (contralateral to the side of stimulation) spatial positions, and affects saccade latencies (Dai et al., 2014; Dominguez-Vargas et al., 2017; Hanks et al., 2006). Neglect-like deficits following stroke-induced parietal and pulvinar lesions in humans are also consistent with the critical contribution of both regions to visually-guided spatial behavior (Arend et al., 2008a, 2008b; Karnath, 2015; Karnath et al., 2002; Rafal et al., 2004; Van der Stigchel et al., 2010).

Area LIP and the dorsal pulvinar share a large number of afferent and efferent subcortical and cortical connections (Blatt et al., 1990; Lewis and Van Essen, 2000). Anatomical tracer studies have shown that both regions are reciprocally connected to other areas in the posterior parietal cortex (e.g. MIP, AIP, area 7) (Asanuma et al., 1985; Cappe et al., 2009, 2007; Grieve et al., 2000; Hardy and Lynch, 1992; Yeterian and Pandya, 1985), the lower and upper bank of the superior temporal sulcus (STS), eye movement-related areas in prefrontal cortex such as dorsolateral prefrontal cortex (dlPFC) and frontal eye fields (FEF, areas 8 and 45) (Homman-Ludiye et al., 2020; Lewis and Van Essen, 2000;

Romanski et al., 1997) and extrastriate visual cortex (Blatt et al., 1990). Similar functional connectivity patterns have been reported in human fMRI studies (Arcaro et al., 2018, 2015). While LIP projects to the intermediate layers of the superior colliculus (Andersen et al., 1990; Pare and Wurtz, 1997), dPul only receives afferent projections from the intermediate and deep layers of the superior colliculus without sending a projection back (Baldwin and Bourne, 2017; Bender and Butter, 1987; Benevento and Standage, 1983; Harting et al., 1980).

The combination of electrical microstimulation and fMRI (es-fMRI) has emerged as a powerful technique to probe the effective connectivity of a local cortical or subcortical stimulation site to distal brain regions *in vivo* (Field et al., 2008; Logothetis et al., 2010; Matsui et al., 2011; Moeller et al., 2008; Murayama et al., 2011; Petkov et al., 2015; Sultan et al., 2012), reviewed in Klink et al. (this issue). Es-fMRI allows the mapping of anatomical connections in a living animal with a precision that cannot be achieved with other in-vivo methods such as diffusion tensor imaging or resting state connectivity (Honey et al., 2009; Schilling et al., 2019). Furthermore, previous studies showed that es-fMRI provides functional information that goes beyond purely anatomical connections. Particularly, the strength of stimulation effects is modulated by the brain state (e.g. anaesthetized vs. awake) (Murris et al., 2020), the properties of the employed visual stimuli such as contrast or the visual responsiveness of activated voxels (Ekstrom et al., 2009, 2008) and task demands (Premereur et al., 2013). The first *goal* of this study was to identify the shared functional circuitry of dPul and LIP during oculomotor tasks. While es-fMRI effective connectivity of the pulvinar has not been investigated previously, es-fMRI effects of LIP have been studied as a control experiment in a work on grasp-related anterior intraparietal area AIP, but were not the main focus of this paper (Premereur et al., 2015). Second, we asked how the magnitude of stimulation effects of dPul and LIP on BOLD responses depends on task demands and/or spatial contingencies. A similar question has been addressed in a es-fMRI study on area FEF (Premereur et al., 2013), but not for parietal or thalamic regions. Based on our previous behavioral context-contingent effects of pulvinar microstimulation (Dominguez-Vargas et al., 2017) and es-fMRI studies in FEF showing visual stimulus-gated stimulation effects in early visual areas, and stronger effects in a saccade task compared to passive fixation (Ekstrom et al., 2009; Premereur et al., 2013), we hypothesized that microstimulation-elicited fMRI effects might be more pronounced in the saccade task compared to pure visual fixation. Third, we asked whether microstimulation in LIP and/or dPul interacts with the spatial tuning of a given brain region. For instance, we expected the largest microstimulation effect in the regions less optimally driven by the task (Ekstrom et al., 2008; Premereur et al., 2013), e.g. for contraversively tuned voxels in a ipsiversive saccade condition. To address these questions, we measured brain-wide fMRI BOLD activity while monkeys performed a sustained fixation or a memory saccade task towards ipsi- and contraversive spatial positions. This task, similarly to covert spatial attention tasks, elicits strong and

often contralaterally-tuned activation in the fronto-parietal and parieto-temporal network (Bogadhi et al., 2018; Caspari et al., 2015; Kagan et al., 2010; Wilke et al., 2012). We employed a slow time-resolved event-related design to isolate BOLD responses from different task epochs (Kagan et al., 2010), and compared interleaved trials with and without microstimulation, in an extensive dataset comprising microstimulation in several pulvinar and LIP locations.

2 Methods

Data and code availability statement

The data used in this study will be made available to the community via **PRIMatE Data Exchange (PRIME-DE)** repository (http://fcon_1000.projects.nitrc.org/indi/indiPRIME.html), and all custom source code will be uploaded to a publically accessible GitHub repository.

2.1 Procedures

All experimental procedures complied with the ARRIVE guidelines (https://arriveguidelines.org) and were conducted in accordance with the European Directive 2010/63/EU, the corresponding German law governing animal welfare, and German Primate Center institutional guidelines. The procedures were approved by the responsible government agency LAVES (Niedersaechsisches Landesamt fuer Verbraucherschutz und Lebensmittelsicherheit, Oldenburg, Germany).

2.2 Animal preparation

Two adult male rhesus monkeys (*Macaca mulatta*), weighing approximately 9 kg (monkey C) and 10.5 kg (monkey B), served as subjects. Surgical procedures were similar for both animals. In an initial surgery, each animal was implanted with an MRI-compatible polyetheretherketone (PEEK) head post embedded in a bone cement head cap (Palacos with gentamicin; BioMet) anchored by ceramic screws (Rogue Research) under general anesthesia and aseptic conditions. A separate surgery was performed in each animal to implant a PEEK MRI-compatible chamber (inside diameter 22 mm) allowing access to the right pulvinar and LIP. To aid the MR-guided pre-surgical planning of the chamber's location and angle in the stereotaxic space, performed in the Planner software package (https://github.com/shayo/Planner) (Ohayon and Tsao, 2012), several 1-2 mm indents that were filled with a MR-visible markers during scanning were embedded in the head cap. After confirming chamber positioning with post-surgical MRI scans (Monkey C, right hemisphere: center at 0.5A, 14.5R mm, tilted -11P, 27R degrees; Monkey B, right hemisphere: center at 3.5P, 17R mm, tilted 15P, 30R degrees; the coordinates are relative to stereotaxic zero, A – anterior, P – posterior, R – right; the tilts

are relative to the vertical axis normal to stereotaxic plane: P – posterior, top of the chamber tilted towards the back of the head, R – right, top of the chamber tilted towards the right ear). After confirming chamber positioning with a post-surgical MRI, a partial craniotomy was made inside the chamber.

2.3 Electrical microstimulation

An S88X dual output square pulse stimulator (Grass Products, Natus Neurology, USA) triggered by a custom MATLAB-based task controller software (https://github.com/dagdpz/monkeypsych) generated 200 ms trains of twin pulses at 300 Hz, which in turn triggered a battery-powered constant current stimulus isolator A365 (World Precision Instruments, USA) to produce 60 biphasic pulses. These devices were placed in the control room outside of the scanner Faraday cage. The current (100-250 µA) was delivered to the target structure using single monopolar electrodes (platinum-iridium, 100 mm length, 125 μm thick core, initial 2 cm glass coating with an exposed tip of 40 μm, total thickness of 230 μm including polyamide tubing coating, customer part ID: UEIK1, FHC Inc., USA). The electrode was connected to the lead wire using a small detachable stainless steel tubing connector made from 27G Spinocan Spinal needle (B. Braun). A return (reference) tungsten rod was placed in the recording chamber filled with saline. The two wires leading to the electrode and the reference were twisted together and fed into a shielded BNC cable that was connected, via a low pass filter (50 MHz cutoff) and the patch panel BNC or a waveguide of the scanner Faraday cage, to the output of the stimulus isolator. Voltage drop was monitored as the difference between voltage measured before and after a 10 kΩ resistor in series with the electrode using a 2 channel 2GS/s 200 MHz Tektronix TDS2022C oscilloscope. The manufacturer-specified impedance of the electrodes was 300-500 kΩ. The initial impedance measured in the lab at 1000 Hz before the experiment was 110-1400 kΩ. Since the impedance dropped dramatically after a few stimulation trains were applied, before each session 20-30 pulse trains were delivered to the electrode immersed in saline using 250 μA current, in order to bring the electrode impedance to a more stable regime. Following this procedure, the impedance ranged from 10 kΩ to 70 kΩ in monkey C and from 10 kΩ to 120 kΩ in monkey B. In each stimulation trial, 10 consecutive stimulation trains were applied at a frequency of 1 Hz (see section 2.6 for behavioral task).

2.4 MR imaging

Both animals were scanned in a 3 T MRI scanner (monkey C: Magnetom Tim Trio; Siemens; monkey B: Magnetom Prisma; Siemens). For planning of chamber and electrode positioning for each animal, high-resolution full-head T1-weighted (3D magnetization-prepared rapid gradient-echo, MPRAGE, 0.5 mm isometric) and additional T2-weighted (rapid acquisition with relaxation enhancement, RARE, 0.25 mm in plane, 1 mm slice thickness) images with the slice package aligned to the chamber vertical axis were acquired before and after chamber implantation in an awake state or under anesthesia using the built-in gradient body transmit coil and a custom single-loop receive coil (Windmiller Kolster Scientific). T1- and T2-weighted scans were coregistered and transformed into "chamber normal" orientation (aligned to the chamber vertical axis) and into anterior commissure-posterior commissure (AC–PC) space for electrode targeting and visualization. These images were acquired with the chamber and the grid filled with gadolinium (Magnevist; Bayer)/saline solution (proportion 1:200) with tungsten rods inserted in predefined grid locations for alignment purposes.

Functional BOLD data were acquired using a custom-made 4-channel receive coil and 1-channel transmit coil (Windmiller Kolster Scientific). In each experimental session, the transmit coil and each individual channel of the receive coil were tuned and matched to the resonance frequency of the scanner (~123.3 MHz) with manual knob adjustments, after being fixed in the final position around the monkey head, using portable probe tuning device (Morris Instruments Inc). In each session, advanced 3D shimming within a cuboid volume contained inside the brain followed by an interactive shimming was performed to maximize the T2* and minimize the spectral tuning curve full width at half maximum. Then, T2-weighted images with the slice package aligned to the chamber vertical axis (32 slices) were acquired. The first image was acquired with the electrode inserted into the brain but before lowering it to the targeted brain region. Based on that image, we measured the 2D-distance between the electrode tip and the center of the target structure and adjusted the electrode position accordingly. The second image was acquired after adjusting the electrode position to confirm proper placement within the target structure, allowing for the online control over electrode positioning. In addition, an in-plane (with the functional EPI slices) anatomical scan was acquired (T2-weighted, Turbo-Spin-Echo sequence, TR: 5 s, TE: 18 ms, flip angle: 180°, 0.75 mm x 0.75 mm in-plane resolution, slice thickness: 1.2 mm) for later coregistration with the functional images. Blood-oxygen-level dependent (BOLD) functional images were acquired with a T2*-weighted echo-planar imaging (EPI) single-shot sequence (TR: 2 s, TE: 27 ms, flip angle: 76°, bandwidth: 1302 Hz/pixel, 80x80 matrix, field of view: 96 mm, 1.2 mm x 1.2 mm in-plane resolution, 30 slices in monkey C and 32 slices in monkey B, slice thickness: 1.2 mm, inter slice time: 66.7 ms for monkey C and 61.8 ms for monkey B). Data were acquired in separate, consecutive runs of 454 volumes each. Before functional data

acquisition, the received signal was checked for RF artifacts and the level of background noise using the same echo-planar imaging sequence with the transmitter RF pulse amplitude set to zero, without and with the microstimulation circuit connections and the running microstimulation.

2.5 Pulvinar and LIP targeting

A custom-made MR-compatible polyetherimide (Ultem) grid (0.8 mm hole spacing, 0.43 mm hole diameter) and a custom-made plastic XYZ manipulator drive (design courtesy of Dr. Sebastian Moeller, (Moeller et al., 2008)) were used to position platinum-iridium electrodes (FHC, see section 2.3 for detailed specifications) in the corresponding grid hole. Grid hole planning and penetration trajectory estimation was based on anatomical MRI using Planner (Ohayon and Tsao, 2012) and BrainVoyager (Version 2.4.2.2070, 64-bit; Brain Innovation). During penetration, the electrode was protected by a custom-made MR compatible guide tube (polyimide coated fused silica, 430 μm outer diameter, 320 μm inner diameter; Polymicro Technologies). An MR compatible stopper (polyimide coated fused silica, 700 μm outer diameter, 530 μm inner diameter; Polymicro Technologies) ensured that the guide tube only penetrated the dura and minimally the cortex below. Before penetration, the electrode tip was aligned to the guide tube tip and was held in place by a drop of melted petroleum jelly. For each experimental session, the final electrode location was determined based on anatomical MRI (see section 2.4). In each animal, we stimulated several different sites, some located more anterior and the other located more posterior, both in the right dPul and the right dorsal LIP. For localization of the pulvinar stimulation sites we used the traditional segregation of the pulvinar nucleus into the medial, lateral and inferior pulvinar (PM, PL, and PI) as delineated in currently available online and downloadable atlases (Calabrese et al., 2015; Rohlfing et al., 2012). Some of the anatomical coronal templates were adapted from an earlier version of the NeuroMaps atlas (http://braininfo.rprc.washington.edu/PrimateBrainMaps/atlas/Mapindex.html) (Wu et al., 2000). **Figure 1A** shows the electrode positions in example sessions of dPul stimulation measured in T2-weighted MR images with the slice package aligned to the chamber vertical axis and reconstructed in T1-weighted MR images for each animal, and one coronal section through the center of the electrode tip probability map in AC-PC plane (see section 2.7 for a detailed description of the generation of the displayed probability maps). **Figure S1** (upper two rows in each monkey) shows the full extent of probability maps of the estimated electrode tip positions across dPul stimulation sessions. As can be seen, the dPul stimulation sites corresponded to PM and the dorsal part of PL (PLdm). The brachium of the SC (bsc) and other neighboring structures such as the reticular thalamic nucleus and the tail of the caudate nucleus were avoided. Slightly more anterior dPul(a) and slightly more posterior dPul sites were very close: nominally ~1.1 mm and ~0.8 mm apart from each other in monkey C and monkey B, respectively. Note that in several sessions in both monkeys the more

anterior dPul site was stimulated with currents lower than 250 μA (**Table 1**). Therefore, dPul(a) datasets serve as an additional controls, and main conclusions about the dorsal pulvinar stimulation effects are derived from the dPul datasets.

Additionally, in each animal we targeted one location in the ventral pulvinar (vPul) for direct comparison with the dorsal pulvinar stimulation. **Figure S2** shows the electrode position in an example session of vPul stimulation in a T2-weighted MR image with the slice package aligned to the chamber vertical axis and reconstructed in a T1-weighted MR image in monkey C, as well as the probability map of reconstructed electrode tip positions across all vPul sessions in comparison to dPul stimulation sessions, in the AC-PC space. The probability map indicates that the vPul stimulation sites mostly corresponded to the ventral part of lateral pulvinar (PL) known as PLvl (Kaas and Lyon, 2007). The local microstimulation-elicited BOLD fMRI activation maps for dPul and vPul, together with overlaid electrode probability maps demonstrate that the activation spread below the estimated electrode tip, hence the inferior pulvinar (PI) was also affected by vPul stimulation (**Figure S2**).

For localization of the LIP stimulation sites we used the segregation of LIP into a dorsal (LIPd) and a ventral (LIPv) zone (Saleem and Logothetis, 2006). **Figure 1A** show examples of measured and reconstructed electrode positions and one coronal section through the center of the electrode tip probability map in AC-PC plane. **Figure S1** (bottom two rows in each monkey) shows the full extent of probability maps of the estimated electrode tip positions across LIP stimulation sessions. In monkey C anterior and posterior LIPd sites were nominally ~3.4 mm apart from each other. In monkey B the two LIPd sites were nominally ~5 mm apart.

2.6 Behavioral paradigm

For training and scanning, monkeys sat or lay in custom-made horizontal MR compatible primate chairs (Rogue Research, Canada) in a sphinx position with their heads rigidly attached to the chair with a PEEK headholder. Visual stimuli (800x600 pixels) were back-projected onto a custom-made MR compatible tangential screen, placed in front of a monkey (**Figure 1B**). Visual cues and targets were displayed at one of three locations per hemifield (six locations in total) with an eccentricity of 12° of visual angle. Stimulus locations were arranged concentrically around the fixation spot at 0° (mid left), 30° (up left), 150° (up right), 180° (mid right), 210° (down right), and 330° (down left). Trials were presented in a pseudorandomized order to ensure a similar distribution of trial types throughout the duration of each fMRI run (15 min). Monocular eye position was monitored at 60 Hz with a MR-compatible infrared camera (Resonance Technology/Arrington Research, or MRC-Systems 60-M) and was recorded simultaneously with stimulus and timing parameters and digital triggers from the

scanner. Stimulus presentation, all behavioral control functions, and synchronization of the behavioral task with the scanner were programmed in MATLAB (R2014a, 64-bit; The MathWorks, Inc., USA) using the Psychophysics Toolbox (Brainard, 1997) and in-house developed software (https://github.com/dagdpz/monkeypsych).

To isolate the BOLD fMRI activity associated with the cue and the memory delay phases during stable fixation from the peri- and post-saccadic responses, we used a slow time-resolved event-related design with long trials. In one third of the trials, the animals performed a central eye fixation task. In the remaining two third of trials, they performed delayed memory-guided saccades. **Figure 1B** shows a schematic of the behavioral tasks and the timing of microstimulation. In training sessions the tasks were very similar except for a more variable timing to discourage that animals learn to anticipate the occurrence of trial events. Trials were initiated by fixating the central fixation spot (red dot, 0.25° diameter). Memory-guided saccade trials continued with an eye fixation period (monkey C: 10 s in experimental sessions, 12 - 14 s in training sessions; monkey B: 10 s in experimental sessions, 9.25 - 10.25 s in training sessions). Subsequently, a red filled circle (1° diameter) representing a visuospatial cue was presented for 200 ms either in the contraversive (left, contralateral to the side of stimulation, which was always in the right hemisphere in this study) or the ipsiversive (right, ipsilateral to the side of stimulation) hemifield while the animals were maintaining central eye fixation. The offset of the cue determined the beginning of the memory period (monkey C: 10 s in experimental sessions, 12 - 14 s in training sessions; monkey B: 10 s in experimental sessions and 9.5 - 10.5 s in training sessions) in which the animals were required to remember the cued spatial location and could plan a saccadic eye movement towards this location while maintaining central eye fixation. After the memory period, the fixation spot disappeared, signaling the time the animals were allowed to execute the saccadic eye movement towards the remembered location. If the animals performed a saccade towards the correct location within a radius of 5° to 7° around the center of the cued location within 500 ms, a target stimulus (red, filled circle, 1° diameter) appeared at the saccade location to signal to the animal that the saccade had been performed correctly. After fixating the saccade location for another few seconds (monkey C: 2 s in experimental sessions, 2 - 3 s in training sessions; monkey B: 1.5 s in experimental sessions, 1 - 1.5 s in training sessions) a fluid reward was delivered. In the fixation task, the animals were required to maintain central eye fixation until the end of the trial was signaled by the fixation spot offset (monkey C: 22.2 s in experimental sessions, 26 - 31 s in training sessions; monkey B: 21.7 s in experimental sessions, 20 - 22.5 s in training sessions) in order to get a fluid reward. In training sessions the fluid reward was preceded by a feedback sound. In both tasks, blink allowance time – a period when a fixation break was permitted - was 0.3 s. Trials with fixation breaks exceeding an allowance window of 4° to 5° radius around the fixation spot and trials with incorrect or too slow saccade execution were aborted and not rewarded. In addition to the online eye

tracking, a MR-compatible IR camera (MRC-Systems) connected to video-based motion-detection system (Pelco MD2001) were used to train the animals to minimize their jaw movements during the trials and to track jaw movements during scanning. Jaw movements were only allowed in the inter-trial interval (2 s after aborted trials with no reward, 5 s after correct trials with reward). Trials compromised by detected jaw motion were aborted.

In half of the trials, selected pseudo-randomly, the microstimulation (ten 200 ms trains separated by 800 ms) was delivered throughout the 10 s memory period in the memory-guided saccade task, starting at the time of the offset of the visual cue, or in the corresponding time window in the fixation task (**Figure 1B**). **Table 1** gives an overview of the number of sessions, EPI runs, and successful trials per stimulation site, animal, and sessions for all stimulation sites.

Figure 1. Microstimulation targeting, task and microstimulation timing. (A) Electrode localization in dPul and LIPd. Electrode positions measured in T2-weighted MR images (left column in each monkey) and reconstructed in T1-weighted images (middle column in each monkey), both aligned to the chamber vertical axis, in example sessions with microstimulation in dPul(a), dPul, LIPd-a, and LIPd-p. Outlines mark the respective target region. The right column for each monkey shows an example coronal section through the center of a probability map reconstructing the electrode tip position across all sessions, displayed on a T1-weighted MR image aligned to standard AC-PC space. R - right, y - distance from AC-PC origin along the anterior/posterior plane in mm. **(B)** Timing of one successful trial of the fixation task and the memory saccade task. Dashed circles illustrate the eye position fixation window but were not displayed on the screen. Correct trials were rewarded with a fluid reward. Jaw motion was only allowed during reward delivery and the inter-trial interval (ITI). In half of the trials stimulation was applied in the indicated time period. The timing and the properties of the microstimulation are depicted below the trial timing. On the right inset, the custom specifications used for the manufacturing the stimulating electrodes are depicted.

Table 1. Datasets. Number of EPI runs and successful trials per stimulation site, monkey, and session for stimulation in dorsal pulvinar (more anterior and more posterior sites), ventral pulvinar, anterior dorsal lateral intraparietal area (LIPd-a), and posterior dorsal lateral intraparietal area (LIPd-p).

Stimulation site	Session number	Monkey C			Monkey B		
		Runs	Current (µA)	Successful trials	Runs	Current (µA)	Successful trials
dPul (anterior)	1	3	200	69	7	150-250	133
	2	7	150	165	8	150-200	126
	3	6	150	132	7	150-250	124
	4	8	150	158	7	250	112
	5	7	150	140	7	200-250	124
	6	-		-	8	250	147
	7	-		-	6	250	106
	8	-		-	6	250	120
	9	-		-	7	250	136
dPul	1	3	250	62	7	250	142
	2	6	250	120	8	250	165
	3	6	250	137	7	250	132
	4	6	250	143	8	250	167
	5	5	250	90	6	250	128
	6	6	250	107	7	250	133
	7	9	250	163	9	250	168
	8	-	-	-	6	250	114
	9	-	-	-	8	250	120
dPul low current stimulation	1	7	100	129	-	-	-
	2	6	100	102	-	-	-
	3	6	100	138	-	-	-
	4	5	100	108	-	-	-
	5	6	100	134	-	-	-
	6	6	100	150	-	-	-
	7	10	100	240	-	-	-
vPul low current stimulation	1	3	100	52	6	100	122
	2	9	100	239	4	100	81
	3	5	100	117	7	100	123
	4	8	100	187	6	100	96
	5	6	100	137	8	100	168
	6	7	100	174	8	100	149
	7	6	100	147	6	100	129
	8	-	-	-	7	100	115
LIPd-a	1	8	100, 250	195	6	250	112
	2	6	250	157	8	250	129
	3	6	250	162	8	250	155
	4	8	250	211	11	250	216
	5	6	250	154	3	250	24
	6	-		-	10	250	181
	7	-		-	8	250	143
	8	-		-	9	250	131
	9	-		-	7	250	137
	10	-		-	7	250	128
	11	-		-	8	250	153
LIPd-p	1	6	250	157	6	250	123
	2	8	250	199	8	250	156
	3	5	250	131	7	250	122
	4	9	250	246	8	250	138
	5	6	250	155	7	250	128
	6	-		-	6	250	121
	7	-		-	6	250	123
	8	-		-	6	250	111
	9	-		-	6	250	123
	10	-		-	7	250	139

2.7 Data analysis

2.7.1 Behavioral analysis

For behavioral analysis, all trials of all experimental sessions were pooled for each stimulation site separately. In order to test the effects of microstimulation on task performance, first, the overall hit rate for all control trials without stimulation and for all stimulation trials was calculated, respectively, and a Chi-squared test was performed to determine the effect of microstimulation on the frequency of successful trials. If this Chi-squared test revealed a significant difference between the frequency of successful trials in control and stimulation conditions, additional Chi-squared tests were performed for each task (fixation, contraversive memory saccade, ipsiversive memory saccade) separately to test whether microstimulation affected task performance in a task-specific manner. Since trials in control and stimulation conditions were only different starting from stimulation onset, similar analyses were performed on the frequency of trials aborted during and after the stimulation period, in order to detect changes in task performance that were specific to the delivery of current pulse trains and to assess the effect of stimulation on subsequent saccade execution.

All eye movements with a minimum velocity of 15 °/s and a minimum duration of 10 ms were included in the analysis, including small saccades and eye blinks. The point in time when eye velocity passed the minimum velocity criterion determined eye movement onset. Movement offset was defined as the point in time when eye velocity dropped below 10 °/s. Using these detection criteria, the number of eye movements during the stimulation period was extracted for each trial and averaged across all trials of each experimental condition (control and stimulation for fixation, ipsiversive memory saccade, and contraversive memory saccade).

In order to test whether stimulation influenced fixation behavior, the number of eye movements during the stimulation period was then tested in a two-way ANOVA design with factors task (fixation, ipsiversive memory saccade, contraversive memory saccade) and stimulation (control, stimulation). Further post-hoc t-tests were performed if the ANOVA revealed significant effects of stimulation. To investigate the effects of microstimulation on saccade execution, saccade latencies were extracted from all saccade trials and again tested in a two-way ANOVA design with factors task (ipsiversive memory saccade, contraversive memory saccade) and stimulation (control, stimulation). Significant effects of stimulation on saccade latencies were further tested using post-hoc t-tests.

2.7.2 Estimation of electrode tip positions

For assessment of the variability in electrode tip positions across sessions, for each experimental session the location of the electrode tip was estimated based on the respective T2-weighted image acquired with the slice package aligned to the chamber vertical axis (see section 3.2.4) using BrainVoyager. For each stimulation site, a sphere (radius: 0.5 mm) was created around each session's estimated electrode tip position and a probability map was created based on the resulting volumes of interest showing the probability of overlapping voxels across all spherical volumes of interest. For better comparability, the resulting probability maps were then transformed into AC-PC space and overlaid onto the high-resolution, full-head T1-weighted anatomical image of each monkey.

2.7.3 Functional data processing

The first four EPI volumes were excluded from functional data analysis in order to eliminate transient effects of magnetic saturation. Preprocessing was performed using MATLAB (R2014a, 64-bit; The MathWorks, Inc., USA) and the NeuroElf toolbox (v1.0, http://neuroelf.net/). EPI data of each run was preprocessed using slice time correction and a high-pass temporal filter with a cut-off of three cycles per 15 min run. In addition, 3D motion correction was performed using the first functional volume included into the analysis as a reference. Coregistration and volume time course computation was done using BrainVoyager. First, the in-plane anatomical image of each session was coregistered to the high-resolution full-head T1-weighted anatomical scan in the AC-PC plane. Then, EPI runs were coregistered to the respective AC-PC registered in-plane anatomical image using rigid body transformations with automated initial alignment followed by careful manual fine-tuning of the resulting alignment based on anatomical sulcal landmarks. Finally, volume time courses were computed in AC-PC space using 1 mm x 1 mm x 1 mm voxel size, intersecting the volume time course data a with a mask only including voxels within the brain; this resulted in ~89,000 voxels in monkey C and ~110,000 voxels in monkey B as the total number of voxels considered for FDR correction. A 3D 1.5 mm x 1.5 mm x 1.5 mm Gaussian kernel was applied for spatial smoothing of the volume time courses.

2.7.4 GLM, ROI definition, and event-related averaging

General linear models (GLMs) were computed in MATLAB (R2014a, 64-bit; The MathWorks, Inc., USA) using the NeuroElf toolbox (v1.0, http://neuroelf.net/). For successful trials, all trial events except the initial fixation period – contraversive/ipsiversive cue and memory delay periods with and without stimulation, contraversive/ipsiversive saccades and subsequent peripheral target fixation, and the corresponding time windows in fixation trials, reward delivery and the subsequent inter-trial interval – were extracted and used as predictors for the GLM. In addition, there was one predictor for aborted

trials. The initial fixation served as GLM baseline condition. Based on these 11 task event predictors, design matrices were created using convolution with the macaque hemodynamic response function ((Kagan et al., 2010), time to positive peak: 3 s, time to negative peak: 10 s) in order to compute GLMs. For each run, 6 motion correction parameters were included as confound predictors. Volume time courses were z-score normalized. For each stimulation site, data from all sessions and runs were combined and analyzed using fixed-effects GLM, on volume data in individual subject AP-PC space.

For each animal and stimulation site, regions of interest (ROIs) were defined based on event-related statistical volume-based t-maps contrasting all three stimulation conditions (memory contraversive, memory ipsiversive, fixation) with the corresponding control (no stimulation) conditions across the tasks ("microstim > no microstim"), using an FDR-corrected threshold of q < 0.05 and a cluster threshold of k ≥ 20 1 mm^3 functional voxels. The resulting maps were converted to cluster tables using NeuroElf (https://neuroelf.net/wiki/doku.php?id=vmp.clustertable). Large contiguous clusters were further split to subclusters by searching for local maxima within each cluster (subclusters belonging to same cluster could be less than 20 voxels). The resulting (sub)clusters were exported as regions of interest (ROIs) and then revised using BrainVoyager, with a nomenclature based on rhesus monkey atlases (Calabrese et al., 2015; Saleem and Logothetis, 2006). Clusters covering more than one brain region were split accordingly by extracting a maximum volume of 2.5 x 2.5 x 2.5 mm^3 to 3.5 x 3.5 x 3.5 mm^3 around the center of activation belonging to a particular brain region avoiding spatial overlap between ROIs. Few clusters that could not unambiguously be assigned to gray matter were excluded from the analysis. For conjunction analysis, the highest t-value of the two contrast statistical maps corresponding to a FDR-corrected threshold q < 0.05 was used.

For each stimulation site dataset, BOLD time courses of each ROI were extracted using MATLAB and the NeuroElf toolbox. For event-related averaging, first, BOLD time course data were interpolated to the temporal resolution of 1 s. Then, for each trial even-related time course in %BOLD signal change was computed relative to the individual trial baseline, which was defined as the mean BOLD activity in the last 4 s of the initial fixation period before cue presentation in memory-saccade trials or the corresponding time window in fixation trials. Then, BOLD time courses were averaged across trials per each condition.

2.7.5 Region of interest analysis within and across ROIs

For ROI analysis, the mean response amplitude in %BOLD signal change between 2 and 9 s of the memory period or the corresponding time epoch in fixation trials was calculated for each ROI. The start of the analysis epoch was selected based on the observed fast responses to visual cues and the

stimulation onset (time to peak 3-4 s); the end of the epoch was selected so that the last second of the delay period, which can be potentially contaminated by the saccade response, was omitted.

Within each ROI, we calculated the main effect of task (memory saccade right - ipsiversive, memory saccade left – contraversive, fixation), the main effect of stimulation (no stimulation - control, stimulation) and the task × stimulation interaction, using a two-way ANOVA across single trial response amplitudes.

To investigate whether the magnitude of stimulation effect on BOLD responses depended on the task, we calculated the stimulation effect for each task and ROI as the difference between the mean BOLD response in the respective stimulation and control condition. Stimulation effects were then analyzed across ROIs using a two-way repeated measures ANOVA (rmANOVA) with the factors task (ipsiversive memory saccade, contraversive memory saccade, and fixation) and stimulation (no stimulation, stimulation) for each hemisphere separately. Significant effects were further investigated using paired-sample t-tests. Outlier ROIs with a mean BOLD response deviating more than 2.5 standard deviations from the mean BOLD response in control conditions across all ROIs for a given dataset were excluded from further across ROI analysis.

For each ROI we assessed the degree of contraversive selectivity, i.e., whether the BOLD response of an ROI was stronger to visual cues/saccade preparation presented in/towards spatial locations in the hemifield contralateral to the side of stimulation (contraversive memory saccade). To calculate contraversive selectivity index (CSI), we subtracted the mean BOLD response in ipsiversive trials from the mean BOLD response in contraversive trials for control and stimulation conditions for each ROI. Positive CSI values reflect contraversive tuning (contralateral for the stimulated hemisphere). Negative CSI values reflect ipsiversive tuning, i.e. preference towards the side of space that is ipsilateral to the side of stimulation (ipsiversive memory saccade condition). Note that the ipsiversive tuning is contralateral tuning for the non-stimulated (left) hemisphere.

To test whether stimulation effects on BOLD responses depended on the initial spatial tuning of ROIs, we categorized ROIs of both hemispheres according to their contraversive selectivity as indicated by CSI values for control trials (positive vs. negative values) and tested the differences between the mean BOLD response in the respective stimulation and control condition of each task in a two-way mixed ANOVA with between-subjects factor spatial tuning: contraversive selectivity, ipsiversive selectiivty – and within-subjects factor task condition: ipsiversive memory saccade, contraversive memory saccade, fixation (the "subject" here being each ROI). Interaction effects were further

investigated using paired-sample t-tests to compare stimulation effects between tasks for ROIs with contraversive and ipsiversive tuning, separately.

To test if microstimulation-elicited enhancement was best described by adding activity to the initial BOLD response in the control condition, or whether stimulation leads to multiplication of the BOLD response in the control condition, we fitted the relationship between BOLD response amplitudes (RA) with and without stimulation with an additive model: $RA_{stim} = RA_{cont} + a$, an additive model scaled by the control (no stimulation) response amplitude: $RA_{stim} = RA_{cont} + a*(1 - b*RA_{cont})$, and a multiplicative model $RA_{stim} = a*RA_{cont}$, for each task and hemisphere separately, using the *fit.m* function with *NonlinearLeastSquares* method in MATLAB. We then evaluated the resulting adjusted R-squared values as indicators of goodness of fit.

2.7.6 Surface maps

The individual monkeys' left and right hemisphere surfaces were reconstructed using standard BrainVoyager segmentation and surface reconstruction steps. The surface maps show the activity patterns extracted from the volume-based GLM contrasts (voxel-based map, "VMP" in BrainVoyager terminology), sampled along the gray matter/white matter boundary mesh vertex normal, from -0.5 mm to 1.5 mm, and mapped on the reconstructed surface boundary ("VMP"->"SMP" surface map conversion). Note that some volume activations above and below this 2 mm layer (e.g. in the arcuate sulcus and in the superior temporal sulcus) may not be visible on surface maps.

3 Results

3.1 Task performance and eye movements

During scanning, monkeys performed two oculomotor tasks: central fixation and delayed memory saccades (**Figure 1**). The memory trials were separated into left, i.e. contraversive (to the side of stimulation) and right (ipsiversive) saccades for further analysis. In a half of the trials, the microstimulation was applied for 10 s, corresponding to the memory delay period, or in the corresponding 10 s in fixation only trials. The stimulation of the different pulvinar and LIP sites (as well as control experiments in the ventral pulvinar) was conducted in separate sessions, generating 4 datasets per animal (dPul(a), dPul, LIPd-a, LIPd-p, see section 3.2). The main conclusions about the effect of the stimulation in the pulvinar were derived from the dPul dataset, and adjacent, slightly more anterior stimulation site dPul(a) served as a supplementary dataset.

Details of the behavioral analysis are shown in the **Supplementary Results (Figures S3-6)**. In short, both dPul and LIP stimulation did not lead to major impairments in the fixation or the memory saccade task. Only LIPd-p stimulation was associated with a slightly lower hit rate for memory saccades to the ipsiversive hemifield in monkey C. In monkey B, only dPul(a) stimulation led to a statistically significant but modest impairment in the execution of memory saccades to the contraversive hemifield (from 4% to 12%). Both dPul and LIP stimulation led to a modest increase in saccade latencies (~20-30 ms). More specifically, dPul and LIPd-p stimulation in monkey C were associated with longer latencies only for contraversive saccades, and a similar effect was found for LIPd-a stimulation in monkey B, while dPul(a) stimulation in monkey B significantly increased saccade latencies towards both the contraversive and the ipsiversive hemifield.

For the fMRI analysis, aborted trials were modeled in the GLM to account for the introduced variance, and hence might influence the statistics of activation patterns, but for the main statistical contrasts we only used successfully completed and rewarded trials (**Table 1**). The aborted trials were also excluded from event-related averaging of BOLD signal timecourses in the region-of-interest (ROI) analysis. This approach, the overall mild and not always consistent effects of the microstimulation on behavior, as well as asymmetrical (stronger in the stimulated hemisphere) patterns of microstimulation-elicited activation (see section 3.3) all support the notion that the fMRI results were not majorly affected by the changes in performance and eye movements.

3.2 BOLD activity induced by dPul and LIP microstimulation

3.2.1 Activation at and around stimulated sites

Microstimulation elicited strong local BOLD activity in the targeted dPul or LIP (**Figure 2**). The extent of activation around the electrode tip was substantially larger than expected from a passive current spread (Klink et al. this issue). Dorsal pulvinar stimulation affected mainly the medial and the lateral pulvinar (PM and PLdm) and also elicited activations in the adjacent ventral pulvinar (vPul) and the superior colliculus (SC) (**Figure 2A,B**). Stimulation of LIPd led to an extensive activation along the intraparietal sulcus, including dorsal and ventral LIP (LIPd and LIPv) and adjacent cortical regions (**Figure 2C**). Note that BOLD activity in the vicinity of the stimulating electrode (which was present both in stimulation and no stimulation trials) cannot be measured due to signal dropouts caused by the magnetic susceptibility artefacts introduced by metallic objects.

Figure 2. Local activation at the stimulation sites. (A) Coronal sections showing statistical t-maps of BOLD activation elicited by microstimulation in the two dPul sites (microstimulation > no microstimulation contrast, all tasks combined). In this and all other figures (unless stated otherwise), the activation maps were thresholded at q(FDR)<0.05 level and with a cluster threshold of 20 functional voxels, producing individual threshold t-value for each dataset and each monkey (see Materials and Methods): in each pair, left value is for monkey C, right value for monkey B. Cyan arrowheads indicate the location of the electrode tip, and an approximate angle. **(B)** Coronal sections showing activation (top image, red-yellow map, dPul dataset) through the center of the electrode tip reconstruction map (bottom image, cyan map). Schematic outlines of the pulvinar nuclei and neighboring structures were adapted from the NeuroMaps atlas (Rohlfing et al., 2012), shown in the middle. PM – medial pulvinar, PL – lateral pulvinar, PI – inferior pulvinar. L – left, R – right, S – superior, I – inferior, A – anterior, P – posterior, y - distance from AC-PC origin along the anterior/posterior plane, z - distance from AC-PC origin along the superior/inferior axis, in millimeters. **(C)** Coronal sections showing statistical t-maps of BOLD activation elicited by microstimulation in the two LIPd sites (microstimulation > no microstimulation contrast, all tasks combined).

3.2.2 Distal activation patterns

Both the pulvinar and parietal unilateral stimulation elicited widespread distal activations, in both hemispheres. **Figure 3A,B** shows statistical t-maps reflecting BOLD activity enhancement elicited by stimulation on the inflated brain surface based on the animal's individual anatomy. **Figure 3C** and **S7** show the same maps in the volume space on coronal sections. In the stimulated hemisphere, consistently across the two animals dPul stimulation activated primary and extrastriate visual cortex, an extensive swath encompassing dorsal and ventral superior temporal sulcus (STS) areas (dorsal/fundus: IPa, TAs, FST, PGa, TPO, MST; ventral: TEO, TEa, MT), parietal cortex (area 7a and many regions in the parieto-occipital and intraparietal sulcus: PO, LOP, MIP, LIP, VIP), and frontal and prefrontal cortex (45a/b, 8A/B, 46d/v) (**Figure 3A**). Subcortical regions such as insula, cingulate cortex, amygdala, caudate nucleus (monkey C), mediodorsal thalamus (MD, monkey B) and cerebellum were also activated (**Figure S7A**). Enhanced activity was also found in a subset of these cortical and subcortical regions in the opposite hemisphere, although to a lesser extent: visual cortex (both monkeys), STS, IPS and frontal cortex (monkey C), orbitofrontal cortex and caudate nucleus (monkey B). The activation in the opposite hemisphere suggests a polysynaptic transmission of microstimulation-elicited excitation, since to our knowledge there are no direct connections from the pulvinar to cortical regions in the contralateral hemisphere.

The above summary is derived from the main pulvinar datasets (dPul). Note that dPul(a) datasets, especially in monkey C, were more noisy and in some sessions lower currents (150 – 200 mA) were used; therefore we treat these datasets as supplementary (but present summary ROI statistics in subsequent analyses for completeness). **Figure S8** shows the comparison of dPul and dPul(a) stimulation sites. In both monkeys, the dPul and dPul(a) distal activation effects in the stimulated hemisphere coincided in the dorsal bank and the fundus of STS (MST, TPO, FST, IPa), ventral STS (MT, TEO), parietal area LIP, and prefrontal areas 8B and 45, while dlPFC areas 46d/v were only activated by the more posterior dPul stimulation.

Similar to dPul stimulation, in both animals LIPd stimulation activated a large number of cortical areas in the stimulated hemisphere: primary (V1) and extrastriate visual cortex (V2, V3, V4), dorsal STS (PGa, TPO, FST, MT, MST), other areas along and around the IPS (7, PO, LOP, LIP, MIP, VIP), prefrontal cortex (45, 8A/B, 46), and cingulate cortex (**Figure 3B**), as well as the cerebellum. Additionally, LIPd stimulation caused strong activation in the medial posterior parietal area 7m/PGM, and dorso-caudal area PEc. In the opposite hemisphere, homotopic IPS regions and several regions in the STS were activated, as well as motor and somatosensory cortex (area 5, 7op, 3a/b) in monkey C, and the dorsal premotor area PMd in monkey B. The homotopic activation of the opposite IPS

regions stands in a contrast to the dPul stimulation, which did not elicit the activation in the opposite pulvinar.

Figure 3. Cortical surface maps. Statistical t-maps showing BOLD activation elicited by microstimulation on the inflated brain cortical surface (lateral, medial) for **(A)** dPul stimulation and **(B)** LIPd stimulation (LIPd-a and LIPd-p maps overlaid). Major sulci and regions of interest are labeled. **(C)** dPul and LIPd stimulation activation maps in the volume space, coronal sections (2 mm spacing). L – left, R – right, S – superior, I – inferior, y - distance from AC-PC origin along the anterior/posterior plane in mm. See **Figure S7** for the same maps plotted separately for dPul and LIPd sites.

Figure 4. Overlap of thalamic and parietal activation. (A) Statistical t-maps showing BOLD activation on the inflated brain cortical lateral surface for dPul stimulation overlaid together with LIPd-a or dPul-LIPd-p stimulation. **(B)** Conjunction maps showing the overlap of activation elicited by the stimulation of dPul with LIPd-a or dPul-LIPd-p. Major overlap regions are labeled.

3.2.3 Overlap of pulvinar- and LIP-elicited effective connectivity

We next analyzed the overlap of dPul- and LIPd-elicited activations. **Figure 4** presents the overlaid dPul and LIPd activation maps (**4A**) and the conjunction maps (**4B**) in the stimulated hemisphere. The left (non-stimulated) hemisphere is not plotted since there was only a very minor overlap. Beyond LIP regions, the shared effective connectivity between dPul and either of the two LIPd sites was strongest in the dorsal bank of STS, in caudal, middle, and anterior parts of the sulcus (areas MST, FST, TPO, IPa) and, only for the LIPd-a, areas in pre(frontal) cortex (8A, 8B, 46d/v). These results implicate the dorsal bank and the fundus of STS as a main shared "node(s)" of the dorsal pulvinar-parietal circuitry.

3.2.4 Localized specificity of stimulation and distal connectivity

Despite a substantial local spread of BOLD activation, there are indications that the distal effects of stimulation were quite specific for the stimulation site. Firstly, the posterior and anterior LIP sites elicited partially distinct distal effects, especially in the prefrontal regions (please refer back to **Figure 3B**). Namely, LIPd-a stimulation caused extensive activity in prefrontal cortex (8A, 45, and 46), but activated fewer areas in visual cortex. Conversely, prefrontal activation during LIPd-p stimulation was mainly restricted to medial area 8B and area 46 whereas activation in visual cortex was more extensive. Secondly, we directly compared the activation patterns resulting from stimulation of dPul and adjacent, but anatomically and functionally distinct ventral pulvinar, vPul (encompassing the ventral division of lateral pulvinar, PLvl, and inferior pulvinar, PI). Since ventral pulvinar is smaller compared to dPul, and since we wanted to limit the spread of activation into the dorsal pulvinar, we stimulated vPul with a lower current strength (100 μA). Despite a partial overlap of local activations around the stimulation sites (but with clearly separated peaks of activation), the comparison of dPul and vPul distal stimulation effects showed divergent effective connectivity with a good correspondence to their respective anatomical connectivity, with vPul es-fMRI connectivity to early visual areas and the ventral bank of STS (**Figure 5**, **S9**). The stimulation of vPul also led to a distinct activation peak in SC, unlike the more diffuse dPul activation spread encompassing SC on the fringe but not exhibiting a separate SC peak. Further control experiments comparing a lower current strength (100 μA) and hence more local dPul stimulation with the vPul stimulation confirmed these findings (**Figure S2, S10**). Thus, even when the stimulation sites are only 2-3 mm apart, the distal activations show distinct patterns.

One unexpected finding was the activation of the frontal cortex following stimulation of vPul in area 8Ad/v and 45b in monkey C, and in areas 45a and 9/46v, with minor activity elicited also in 45b, in monkey B (**Figure S11**). To the best of our knowledge, direct anatomical connections from vPul to these regions were not reported, further suggesting polysynaptic propagation of microstimulation-

elicited fMRI activity. Likewise, the activation of SC both after dPul and vPul stimulation can also be polysynaptic via thalamo-cortical-collicular pathway, or via antidromic propagation.

Figure 5. Comparison of dPul and vPul stimulation effects. **(A)** Activation maps on the inflated brain surface, lateral view. **(B)** Zoomed in sections through the thalamus (1 mm spacing). **(C)** Same as (B), but with a higher statistic threshold (p<0.0001 uncorrected). L – left, R – right, S – superior, I – inferior, y - distance from AC-PC origin along the anterior/posterior plane in mm.

3.3 Effects of dPul and LIP microstimulation on BOLD responses during different task conditions

Up to this point, we looked at stimulation-elicited activations pooling across all task conditions. We next used such combined microstimulation > no microstimulation maps to extract a set of ROIs, individually for each dataset and each monkey (**Table S1**), and performed event-related averaging of single trial timecourses extracted from those ROIs (Materials and Methods). Example timecourses are shown in the **Figure 6**. The effect of microstimulation typically started 2 to 3 s after the stimulation onset (the first of 10 stimulation trains) and continued throughout the memory delay period or a corresponding fixation period. The stimulation modulated the task-related activity, which often showed a contralateral spatial selectivity of cue and sustained delay activity in control trials (i.e. stronger responses in the contraversive memory task condition for the stimulated hemisphere, and stronger responses in the ipsiversive condition for the opposite hemisphere). In many ROIs, the stimulation tended to enhance the initial task-related activity by a similar amount; in some other ROIs, the stimulation effect seemed to vary between task conditions.

To quantify these effects, single trial response amplitudes were extracted using the analysis window of 2 – 9 s after the onset of stimulation and submitted to a two-way ANOVA with factors task and stimulation, in each ROI. **Figure 7** shows a summary of such analysis for the dPul and LIPd-p stimulation sites, and **Figure S12** shows the analysis for dPul(a) and LIPd-a stimulation sites. Since ROIs were selected based on the positive stimulation effect in the statistical contrast map, main (positive) effect of the stimulation in many ROIs merely confirmed trial epoch-specific enhancement in BOLD signal timecourses. Here, the important question was whether the stimulation effect interacted with the task the animal was performing. At the level of single ROIs, however, there was very little evidence for task-dependence of the stimulation effect – even when the stimulation effect magnitude visually varied considerably between task conditions on the inspection of timecourses, the variability of single trial response amplitudes was too large to result in a statistically significant task × stimulation interaction. Consequently, only few ROIs, mostly in the visual cortex in monkey B, showed such interaction effect.

Figure 6. Example trial even-related average timecourses. Trial averages (and standard error) of %BOLD signal change relative to the last 4 s of the initial fixation are plotted for the last 5 s of the initial fixation, 10 s memory / fixation period coinciding with the stimulation epoch, and 5 s of saccadic response / fixation for the three task conditions, in control and stimulation trials. Timecourses were aligned to the onset of the stimulation that started immediately after the offset of 200 ms peripheral saccade cue, or an end of initial fixation period. The timing of trial events and the stimulation are indicated on the first column, second row panel (gray – fixation baseline, dark red – stimulation period). The analysis window (2 – 9 s) is indicated in the second row, second column panel. **(A)** dPul stimulation. **(B)** LIPd stimulation.

Figure 7. Stimulation effects derived from event-related average response amplitudes. dPul and LIPd-p datasets. For each ROI, the stimulation effect (calculated as a difference between average response amplitudes in %BOLD signal change in trials with and without stimulation) is shown for all trials (gray bar) and the three task conditions (green, red and blue bars) together with the results of two-way ANOVA across trials (asterisk –significant effect of stimulation, filled circle –significant effect of task, cross –significant interaction, p<0.05).

We next investigated the average stimulation effects for each task condition, *across ROIs*. **Figure 8** shows the average magnitude of the effect of microstimulation in all four stimulation sites for each task and hemisphere, and **Table 2** lists the corresponding statistics. We used repeated measures ANOVA across ROIs on the mean BOLD response amplitudes during stimulation and the corresponding periods in trials without stimulation to assess the main effect of task (contraversive saccades, ipsiversive saccades, fixation), the main effect of stimulation (no stimulation, stimulation), and the interaction between the task and the stimulation. Post-hoc paired t-tests contrasted stimulation effects between task conditions. In the right (stimulated) hemisphere both monkeys showed a significant main effect of task (and trivially, main effect of stimulation) for all stimulation sites except main effect of stimulation in dPul(a) in monkey C. More importantly, a significant interaction between the stimulation effect and the task was found (except in the dPul(a) and LIPd-p datasets in monkey B). In the left (non-stimulated) hemisphere the interaction was significant for dPul and LIPd-p stimulation sites in both monkeys. The post-hoc paired t-tests between task conditions showed many significant differences but the direction of the effect was not consistent between monkeys. In keeping with our initial prediction, in monkey C the contraversive saccade condition showed the strongest stimulation effect compared to fixation and ipsiversive saccade trials; while in monkey B, the fixation or ipsiversive task condition showed a strongest stimulation effect.

As expected, stimulation effects were stronger in the stimulated hemisphere compared to the opposite hemisphere, based on both the number of significantly activated ROIs (**Figure 7** and **S12**, **Table S1**) and on the statistical comparison of the stimulation effect strength combining task conditions. Across ROIs, 5/8 datasets showed higher stimulation enhancement in the stimulated hemisphere (two-sided t-test, p<0.001; the LIPd-a dataset in monkey B showed a trend with p=0.063, and dPul(a) datasets in both monkeys did not show this effect).

Figure 8. Effects of stimulation in dPul(a), dPul, LIPd-a, and LIPd-p on trial-averaged BOLD responses. Means and standard errors across ROIs for stimulation effects calculated as the difference in %BOLD signal change between trials with and without stimulation, for the contraversive memory saccade task (blue bars), the ipsiversive memory saccade task (red bars), and the fixation task (green bars). See **Table 2** for the corresponding statistics.

Table 2. Stimulation effects in different task conditions. Significant p-values (p<0.05) for the two-way repeated measures ANOVAs and t-tests are in bold font. p – p-value, F – F-statistics, t – t-value.

	Main effect task	Main effect stim.	Task × stim.	Contra vs ipsi	Contra vs fix	Ipsi vs fix	Main effect task	Main effect stim.	Task × stim.	Contra vs ipsi	Contra vs fix	Ipsi vs fix
Format	F p	F p	F p	t p	t p	t p	F p	F p	F p	t p	t p	t p
	Left hemisphere						Right hemisphere					
Monkey C												
dPul(a)	5.18	22.14	0.03	0.14	-0.12	-0.22	23.72	0.61	9.45	3.35	-0.39	-4.89
	0.010	**0.000**	0.969	0.888	0.909	0.825	**0.000**	0.438	**0.000**	**0.001**	0.699	**0.000**
dPul	7.34	51.15	13.91	4.55	2.36	-4.05	5.91	40.50	3.79	2.32	1.80	-0.85
	0.002	**0.000**	**0.000**	**0.000**	**0.030**	**0.001**	**0.004**	**0.000**	**0.027**	**0.026**	0.080	0.403
LIPd-a	0.89	17.34	1.49	0.38	1.82	1.33	29.76	48.12	3.13	1.81	-0.44	-2.65
	0.428	**0.002**	0.252	0.712	0.103	0.215	**0.000**	**0.000**	**0.049**	0.078	0.660	**0.012**
LIPd-p	38.51	71.90	8.57	3.29	-0.02	-3.95	9.51	23.85	11.91	4.23	0.96	-4.33
	0.000	**0.000**	**0.001**	**0.002**	0.988	**0.000**	**0.000**	**0.000**	**0.000**	**0.000**	0.343	**0.000**
Monkey B												
dPul(a)	0.19	35.49	1.44	-2.32	-0.76	0.79	13.04	63.16	1.77	1.10	1.72	0.87
	0.832	**0.009**	0.308	0.103	0.502	0.489	**0.000**	**0.000**	0.177	0.279	0.093	0.392
dPul	6.86	14.24	15.79	-2.12	4.43	4.37	32.87	80.64	23.11	-2.50	4.17	6.83
	0.005	**0.003**	**0.000**	0.058	**0.001**	**0.001**	**0.000**	**0.000**	**0.000**	**0.015**	**0.000**	**0.000**
LIPd-a	16.85	8.10	2.07	-0.73	-1.66	-1.39	37.04	28.77	14.17	-3.27	-4.38	-3.05
	0.000	**0.012**	0.144	0.477	0.118	0.184	**0.000**	**0.000**	**0.000**	**0.002**	**0.000**	**0.004**
LIPd-p	2.39	0.99	6.92	-0.27	-2.94	-4.40	17.09	66.45	2.41	-1.31	-2.16	-0.66
	0.128	0.354	**0.008**	0.794	**0.022**	**0.003**	**0.000**	**0.000**	0.098	0.200	**0.039**	0.515

3.4 Relationship between spatial selectivity and the magnitude of stimulation effects on BOLD responses

To further explore the dependence of stimulation effects on the task, and to understand the discrepancies between the two monkeys regarding the relative strength and direction of such effects, we next assessed whether the magnitude of stimulation effects on task-related activity depended on the initial spatial selectivity of the activated ROIs. To this end, we correlated contraversive selectivity index (CSI) derived from the control trials with the stimulation effect across all ROIs, separated by task condition. Note that for the ROIs in the stimulated right hemisphere the contraversive selectivity corresponds to the contralateral selectivity, while for the left hemisphere the contraversive selectivity is the ipsilateral selectivity. A significant correlation would suggest that the stimulation effect depends on initial spatial tuning, within a given task condition. Furthermore, if there is a difference in the above relationships between task conditions (i.e. different regression slopes), this would suggest a degree of task-dependence. To assess whether the relationship between the stimulation effect and CSI differed between task conditions, we grouped ROIs into two categories: with contraversive tuning (CSI > 0) or ipsiversive tuning (CSI < 0), and tested whether the magnitude of stimulation effects differed between tasks within each group using a two-way mixed ANOVA (see Materials and Methods). Further post-hoc t-tests were used to assess pairwise differences between tasks for each of the two ROI groups.

The resulting correlations for the contraversive and ipsiversive memory saccade task conditions are shown as scatter plots with least-square fits (**Figure 9**), and the corresponding statistics are shown in **Table 3**. The plots for all pair-wise task comparisons are shown in **Figure S13** and **S14**. To summarize the dependency of stimulation effects on spatial selectivity, for both dPul and LIP stimulation sites we consistently found positive correlations between stimulation-elicited enhancement and contraversive selectivity across ROIs in the ipsiversive memory saccade task (7 out of 8 datasets). In other words, stronger contraversive tuning of the respective ROIs was associated with a stronger microstimulation-elicited enhancement of BOLD responses to visual cues and motor preparation directed towards the ipsiversive hemifield. When significant, positive correlations were also found for the fixation task (3 out of 8 datasets), apart from one negative correlation in monkey B, LIPd-a site (**Figure S14**). Significant correlations for the contraversive task were less consistent: negative correlations for the dPul and dPul(a) sites in monkey C, but a positive correlation for the LIPd-a site in monkey B.

Figure 9. Microstimulation effect dependence on contraversive spatial selectivity and task. For each ROI, the task-specific effect of stimulation is plotted against ROI's initial (in trials without stimulation) contraversive selectivity index for contraversive (blue) and ipsiversive (red) memory saccade task conditions. Each dot represents one ROI (filled dots – right hemisphere, empty dots – left hemisphere); solid lines show linear fits of stimulation effects across ROIs. See **Table 3** for the corresponding statistics, and **Figure S14** and **S15** for data on all pairwise task condition comparisons.

Table 3. Relationship between stimulation effect, spatial selectivity and task condition. Significant p-values (p<0.05) for ANOVAs and t-tests are in bold font. For this analysis, ROIs from both hemispheres were combined. R – Pearson's linear correlation coefficient, N – number of ROIs before / after exclusion of outliers (see Materials and Methods), p – p-value, F – F-statistics (two-way mixed ANOVA), t – t-value.

	Corr. contra	Corr. ipsi	Corr. fix	Main effect tuning	Main effect task	Tuning × task	Contra vs ipsi CS<0	Contra vs ipsi CS>0	Contra vs fix CS<0	Contra vs fix CS>0	Ipsi vs fix CS<0	Ipsi vs fix CS>0
Format	R(N) p	R(N) p	R(N) p	F p	F p	F p	t p	t p	t p	t p	t p	t p
Monkey C												
dPul(a)	-0.53 (83/82) **0.000**	0.34 (83/82) **0.002**	-0.13 (83/82) 0.256	1.97 (1,80) 0.164	16.35 (2,160) **0.000**	24.36 (2,160) **0.000**	11.10 **0.000**	-1.03 0.308	4.59 **0.000**	-3.09 **0.003**	-4.63 **0.000**	-1.69 0.097
dPul	-0.47 (58/56) **0.000**	0.30 (58/56) **0.024**	0.27 (58/56) **0.047**	0.17 (1,54) 0.678	9.64 (2,108) **0.000**	16.53 (2,108) **0.000**	5.98 **0.000**	-0.60 0.556	4.84 **0.000**	-1.53 0.140	-2.10 **0.043**	-1.46 0.157
LIPd-a	-0.09 (52/50) 0.535	0.28 (52/50) 0.051	0.15 (52/50) 0.283	0.00 (1,48) 0.996	5.73 (2,96) **0.004**	3.99 (2,96) **0.022**	4.93 **0.001**	0.65 0.516	1.58 0.154	-0.37 0.713	-1.65 0.138	-1.11 0.273
LIPd-p	0.01 (86/84) 0.942	0.60 (86/84) **0.000**	0.39 (86/84) **0.000**	9.77 (1,82) **0.002**	6.93 (2,164) **0.001**	9.58 (2,164) **0.000**	6.66 **0.000**	-1.03 0.322	2.35 **0.022**	-2.54 **0.024**	-5.59 **0.000**	-1.79 0.098
Monkey B												
dPul(a)	0.02 (46/45) 0.885	0.34 (46/45) **0.021**	0.15 (46/45) 0.333	0.70 (1,43) 0.407	1.30 (2,86) 0.277	1.51 (2,86) 0.226	1.64 0.118	-0.74 0.468	1.25 0.227	0.80 0.434	-0.35 0.734	1.54 0.137
dPul	0.06 (90/86) 0.592	0.52 (90/86) **0.000**	0.27 (90/86) **0.012**	12.49 (1,84) **0.001**	32.96 (2,168) **0.000**	7.62 (2,168) **0.001**	0.81 0.422	-4.98 **0.000**	4.68 **0.000**	2.29 **0.027**	4.28 **0.000**	6.29 **0.000**
LIPd-a	0.51 (70/68) **0.000**	0.66 (70/68) **0.000**	-0.25 (70/68) **0.042**	8.62 (1,66) **0.005**	20.41 (2,132) **0.000**	12.19 (2,132) **0.000**	-1.28 0.207	-4.42 **0.000**	-4.90 **0.000**	-2.16 **0.039**	-5.12 **0.000**	0.99 0.331
LIPd-p	-0.15 (40/38) 0.381	0.41 (40/38) **0.012**	0.19 (40/38) 0.242	0.28 (1,36) 0.601	1.85 (2,72) 0.165	5.30 (2,72) **0.007**	4.20 **0.006**	-2.30 **0.029**	1.29 0.244	-3.91 **0.000**	-4.88 **0.003**	-1.60 0.120

Importantly, we consistently found interactions between the main effect of spatial tuning and the main effect of task condition on the stimulation effect magnitude: in 7 out of 8 sites, with the only exception being dPul(a) site in monkey B (**Table 3**). This indicates that across ROIs, the stimulation effects differed between ROIs with contraversive tuning and ROIs with ipsiversive tuning, depending on task condition. More specifically, in ROIs with ipsiversive tuning in monkey C, both dPul and LIP stimulation most strongly enhanced BOLD responses in the contraversive memory task and exhibited weakest effects in the ipsiversive memory task. Likewise, the ipsiversively-tuned ROIs in monkey B showed a consistent pattern with one exception: for dPul and LIPd-p datasets, the stimulation effect was strongest for the contraversive task (significant only in LIPd-p) but LIPd-a dataset showed the strongest effect in the fixation task. On the other hand, contraversively-tuned ROIs consistently showed a stronger stimulation effect in the ipsiversive task compared to the contraversive task and fixation (although these tendencies did not always reach significance) in both monkeys. Taken

together, our results indicate an inverse ("X-shaped") patterns of task-dependent effects for ROIs with contraversive tuning and ROIs with ipsiversive tuning. Furthermore, variations in the relative frequency of contraversively- and ipsiversively-tuned ROIs, as well as the slope and the intercept of the observed stimulation effect fits, could explain differences between average task-specific stimulation effects across monkeys and datasets.

3.5 Modeling effects of microstimulation on BOLD responses

Finally, we asked which model can account best for the observed stimulation effects on BOLD signal amplitudes. The two basic models represent an additive or a multiplicative mechanism, i.e. an enhancement of the initial (no stimulation) response by a constant added term, or a multiplication by a certain factor. Additionally, since we observed a dependence of the microstimulation effect on the initial tuning of a region (ROIs with a negative contraversive selectivity showed more enhancement in contraversive memory saccade trials, while ROIs with a positive contraversive selectivity showed more enhancement in ipsiversive memory saccade trials), we also fitted a variant of an additive model where the additive factor is scaled down by the initial response amplitude. Because of the stronger stimulation effect in the stimulated hemisphere, the fits were done separately for each hemisphere. The resulting adjusted R-squared values signifying the goodness of fit are shown in **Figure 10A**. In both monkeys, across most datasets and the two hemispheres, the best model was the additive scaled model, followed by the additive and then the multiplicative model (paired t-test, monkey C: additive (mean adjusted R-squared value 0.169) vs add. scaled (mean 0.363): p=0.016; additive vs multiplicative (mean -0.512) p=0.001; monkey B: additive (mean 0.321) vs add. scaled (mean 0.448) p=0.007; additive vs multiplicative (mean -0.365) p=0.003). This further confirms that while most individual ROIs did not show a significant task-dependence of the microstimulation effect across trials – as would otherwise be expected from a multiplicative model or an additive model with a strong difference between the additive factors across task conditions – the spatial selectivity, reflected in the initial response amplitude, plays a significant role in determining the strength of stimulation effect. The fitted additive coefficients matched the actual magnitude of stimulation effects on BOLD responses (compare **Figure 10B** with **Figure 8**). Furthermore, a simple qualitative simulation implementing the additive scaled stimulation effect could reproduce the "X-shaped" pattern of interaction between the task condition and the stimulation effect strength in relationship to contraversive spatial selectivity (**Figure S15**).

Figure 10. Stimulation effects are best approximated by additive scaled model. (A) Goodness of fit for modeling effects of microstimulation in different sites: dPul(a) (dots), dPul (squares), LIPd-a (diamonds), and LIPd-p (crosses). Adjusted R-squared values are shown separately for the right, stimulated hemisphere (solid lines) and the left hemisphere (dashed lines). **(B)** Fitted coefficients with 95% confidence intervals for the two additive models. Blue: contraversive memory saccade task, red: ipsiversive memory saccade task, green: fixation task.

4 Discussion

Unilateral microstimulation in both dPul and LIPd caused widespread fMRI BOLD activity enhancement, predominantly in prefrontal cortex, parietal cortex, the dorsal bank and fundus of superior temporal sulcus (STS) and in extrastriate visual cortex, in the stimulated hemisphere. Multiple regions along the dorsal bank of the STS stood out as a prominent shared "node" of es-fMRI connectivity for all dPul and LIPd stimulation sites. The es-fMRI activity was also present in the opposite hemisphere, suggesting polysynaptic propagation. The comparison of the more anterior to posterior LIPd stimulation locations showed extensive activation overlap along the intraparietal sulcus, while the two LIP sides evoked partially differing activation patterns in frontal and occipital regions. The comparison of dPul and vPul stimulation also showed distinct effective connectivity mainly consistent with known anatomical connections. For both pulvinar and parietal es-fMRI we observed task-specific correlations between the initial spatial selectivity and the magnitude of stimulation effects, across ROIs. The effects of microstimulation on BOLD signal amplitudes were best fitted by an additive model with an additive factor scaled down by response amplitude in control trials.

4.1 Dorsal pulvinar effective connectivity and comparison with anatomy

Microstimulation in the dPul elicited BOLD activity that was largely consistent with the reciprocal connections with visuomotor and polymodal cortical regions known from anatomical tracer studies in monkeys (Kaas and Lyon, 2007; Shipp, 2003). Thus, the extensive activity along the superior temporal sulcus (e.g. MT, MST, FST, TPO, PGa, TE, TEO), including anterior STS (IPa, TAa) (Yeterian and Pandya, 1991, 1989), posterior parietal cortex (e.g. LOP, LIP, MIP, VIP) (Asanuma et al., 1985; Cappe et al., 2009, 2007), prefrontal cortex (e.g. 8A/8B, 45, 46), insula (Romanski et al., 1997), cingulate and premotor cortices (Baleydier and Mauguiere, 1985) were expected. Those cortical regions are also largely consistent with human studies that investigated intracerebral-evoked responses from dorso-medial pulvinar in epileptic patients (Rosenberg et al., 2009) or used fMRI functional connectivity analyses (Arcaro et al., 2018). This dorsal pulvinar connectivity is conceptually in line with a number of lesion studies in monkeys and humans that highlight its critical contribution to higher-order visual, and oculomotor and skeletomotor functions involving spatial attention (Arend et al., 2008a; Karnath et al., 2002; Petersen et al., 1987; Rafal et al., 2004; Van der Stigchel et al., 2010), decision making and action selection (Komura et al., 2013; M. Wilke et al., 2010; Wilke et al., 2013)

and sensorimotor transformations for visually-guided reaching and grasping (Mundinano et al., 2018; M. Wilke et al., 2010; Wilke et al., 2017, 2018).

Some activations must have been of antidromic and/or polysynaptic origin. For example, the superior colliculus (SC) was activated, although it projects to pulvinar but does not receive a direct projection from it. Theoretically, this could due to antidromic activation or second-order activation through visual or parietal cortex (Baldwin and Bourne, 2017; Bender and Butter, 1987; Benevento and Standage, 1983; Harting et al., 1980; Murayama et al., 2011). Likewise, V1/V2 were activated by microstimulation in the dPul, but retrograde tracer studies typically do not report corresponding projections from the dorsal pulvinar (Adams et al., 2000). Since we applied a relatively high current strength (250 µA) (as typically used in es-fMRI studies (Matsui et al., 2011; Moeller et al., 2008; Tolias et al., 2005)), the early visual cortex activity could be due to co-activation of the ventral pulvinar or SC, and conversely, prefrontal activation by lower current vPul stimulation could have resulted from the co-activation of dPul. This view is supported by the absence of V1 activity in the control experiment where dPul was stimulated with lower (100 µA) currents, and which elicited very minor ventral pulvinar co-activation. On the other hand, the differences of distal activation patterns due to vPul or dPul stimulation, even with at higher currents, suggest that notwithstanding the apparent partial overlap of the local activation, the es-fMRI connectivity is strongly site-specific. Such specificity was also reported for the behavioral effects of dPul vs. vPul stimulation with comparable currents (Dominguez-Vargas et al., 2017). Therefore, our current hypothesis is that the intra-hemispheric activations not expected from the monosynaptic connectivity are due to polysynaptic propagation.

Since the pulvinar is known to project exclusively to ipsilateral cortical regions, the strongest evidence for polysynaptic propagation of es-fMRI effects via inter-hemispheric transcallosal connections is activity in cortical regions in the hemisphere opposite to the stimulation. Similar evidence for polysynaptic spread has been reported in previous es-fMRI studies with stimulation of somatosensory cortex, FEF, and LGN (Matsui et al., 2012, 2011; Murayama et al., 2011).

The activation in the cerebellum was not expected from anatomical studies in primates. Nonetheless, in cats, several anatomical tracer studies reported projections from the cerebellum to the contralateral LP nucleus, a structure partially homologous to the primate pulvinar (Itoh and Mizuno, 1979; Rodrigo-Angulo and Reinoso-Suarez, 1984), and a previous es-fMRI study in monkeys in the deep nuclei of the cerebellum found strong BOLD activity enhancement in the dorsal pulvinar(Sultan et al., 2012). Additionally, a patient with a selective (bilateral) lesion in the medial pulvinar exhibited a combination of parietal and cerebellar deficits (Wilke et al., 2018). Thus, the functional significance of pulvino-cerebellar pathway in monkeys/humans deserves further investigation.

4.2 LIPd effective connectivity and comparison with anatomy

The LIPd stimulation-elicited activation patterns were largely predicted by anatomical tracer studies (Blatt et al., 1990): both banks and the fundus of STS, various areas in the posterior parietal cortex (PO, LOP, MIP, 7a, area 5), prefrontal areas 45a/b, 8A/B and 46d/v, striate and extrastriate visual cortex, superior colliculus and cerebellum. Since anatomical tracer, electrophysiological and functional fMRI studies indicate that LIP is a heterogeneous area, even beyond the well-established ventral/dorsal subdivision (Blatt et al., 1990; Chen et al., 2016; Lewis and Van Essen, 2000; Liu et al., 2010; Patel et al., 2010), we had stimulated two different portions along the dorsal LIP. The anterior stimulation site LIPd-a was similar to a previous es-fMRI study in monkeys that stimulated an anterior portion of the LIP as control for stimulation in the grasp-related area AIP (Premereur et al., 2015). This study reported LIP-induced activation in MIP, FST, V3A and to a lesser extent in FEF. In comparison, the activation patterns in our study were more extensive and included more brain regions (e.g. cerebellum) and notably also visuomotor regions in the non-stimulated hemisphere. Generally, it is difficult to compare the stimulation results across studies due to many methodological variations. Since current strength was comparable and both studies used awake animals, this might is due to differences in other stimulation parameters such as longer stimulation durations in our study. While the macaque literature already makes an anatomical and functional division between ventral LIPd and LIPv subdivision (Lewis and Van Essen, 2000; Liu et al., 2010), the anterior-posterior axis is less well described. While the stimulation of the more posterior vs. anterior sides in LIPd yielded largely overlapping patterns, there was more extensive activation of FEF (area 8A) and dlPFC (area 46) with anterior LIPd-a stimulation. This anterior-posterior gradient should be followed up with systematic studies of electrophysiological properties (Premereur et al., 2015).

Unlike the dPul stimulation effects in the opposite hemisphere, the activation of heterotopic regions in the opposite parietal and parieto-temporal cortex by the LIP stimulation cannot be taken as a definitive proof of polysynaptic propagation, due to a presence of heterotopic transcallosal connections between these regions (Hedreen and Yin, 1981). It remains to be investigated if these connections are dense enough to support monosynaptic transcallosal activation, as an alternative to the polysynaptic transmission via homotopic LIP. Small activations in the premotor or cingulate cortex that we observed are unlikely to be mediated by the direct transcallosal connectivity (Lanz et al., 2017).

4.3 Similarities and differences of LIPd vs. dPul microstimulation effects

Given the considerable differences between thalamic and cortical local organization and projections, it was not *a priori* obvious how much similarity of microstimulation effects should be expected. The comparison of effective connectivity patterns of dPul and LIPd revealed many similarities across prefrontal, parietal, and most extensively, STS areas. This confirms that both regions belong to the same functional brain network supporting the visuospatial guidance of eye movements and attention (Bridge et al., 2015; Gold and Shadlen, 2007; Grieve et al., 2000; Halassa and Kastner, 2017). In support of this notion, inactivation of either region results in a saccade choice bias towards the ipsilesional space (Christopoulos et al., 2018; Wardak et al., 2002; Wilke et al., 2013, 2012). Nonetheless, dPul plays a role in reach as well as saccade choices, while LIP inactivation results mainly in an effector-specific bias for eye movements (Christopoulos et al., 2018; M. Wilke et al., 2010). While the prefrontal and parietal regions have been classically associated with spatial attention and decisions, one interesting location of overlap of the dPul and LIPd stimulation effect was the middle and the anterior portion of STS, encompassing areas TPO, FST, PGa and IPa. This finding is consistent with anatomy, since these areas receive strong input from both LIP (Blatt et al., 1990; Seltzer and Pandya, 1994) and the dorsal pulvinar (Gutierrez et al., 2000; Yeterian and Pandya, 1991, 1989). Importantly, it underlines the importance of these STS regions in the control of eye movements and visuospatial functions found in earlier electrophysiological and lesion studies (Bruce et al., 1981; Karnath, 2001; Luh et al., 1986; Scalaidhe et al., 1995; Watson et al., 1994). Confirming and extending these findings, these STS regions and the adjacent area called PITd have recently emerged as consistently being activated in monkey fMRI studies employing visual attention and delayed saccade tasks (Bogadhi et al., 2018; Caspari et al., 2015; Kagan et al., 2010; Patel et al., 2015). It is worth noting that inactivation-induced changes in these STS regions were also a common denominator of LIP, dPul and superior colliculus inactivation studies, accompanying contralesional saccade selection and attentional deficits (Bogadhi et al., 2019; Klink et al., this issue; Melanie Wilke et al., 2010; Wilke et al., 2013, 2012).

The major differences between the distal activation patterns following dPul vs. LIPd stimulation were the LIPd-only elicited activation of posterior IPS / PO sulci, the medial posterior parietal area 7m-PGM, related to eye movement generation (Thier and Andersen, 1998), as well as the homotopic LIP and adjacent IPS areas in the opposite hemisphere – whereas the pulvinar in the opposite hemisphere was not activated by dPul stimulation.

4.4 Task-dependent effects of microstimulation on BOLD activity

While structural connectivity is relatively fixed, dynamic synaptic connectivity can vary as a function of external inputs and behavioral demands. Thus, the combination of es-fMRI with a behavioral task can be used to investigate the contribution of a given brain region to underlying brain and behavioral processes. Only a few es-fMRI studies investigated microstimulation effects as a function of external inputs or behavioral task contingencies (Ekstrom et al., 2009, 2008; Premereur et al., 2013). A pioneering study using low current FEF microstimulation (26-58 µA) showed that higher order extrastriate areas directly connected to the FEF exhibited microstimulation effect irrespective of visual stimulation (mimicking attention-driven additive "baseline shifts"), while polysynaptically connected V1 showed microstimulation-induced activation only in the presence of visual stimulus, mostly in voxels not directly responsive to the visual stimulus (Ekstrom et al., 2008). Similarly, FEF stimulation induced strong fMRI signal increase in visual areas for the low contrast stimuli, and no or even negative effects for high contrast stimuli that already strongly drove fMRI activity (Ekstrom et al., 2009). Another study from the same group (Premereur et al., 2013) specifically addressed the task-dependence of FEF stimulation effects (83-316 µA) by comparing activity between fixation and saccade tasks with matched visual inputs. This study reported higher es-fMRI effects in visual areas in the more demanding saccade task as compared to passive fixation, but again in the voxels that were not strongly activated by the visual stimulus.

Based on these studies, and the predominately contralateral tuning of LIPd and dPul cue and delay period activity both on the level of single neurons and fMRI signals (Blatt et al., 1990; Caspari et al., 2015; Dominguez-Vargas et al., 2017; Fiebelkorn et al., 2019; Kagan et al., 2010; Patel et al., 2010; Schneider et al., 2020), we hypothesized that the effects of dPul and/or LIP stimulation might depend on the visual stimulation and the direction of the upcoming saccade. For instance, if microstimulation acts as a multiplicative gain mechanism we might expect the highest fMRI-elicited enhancement in the contraversive saccade task condition, at least in the stimulated hemisphere. On the other hand, it could be that the microstimulation effect is strongest when the initial (no microstimulation) task activity is weakest. These effects could be manifested both at the level of single ROIs, and/or across ROIs.

On the level of single trial BOLD response amplitudes within a single ROIs, there was very little indication of interaction between the stimulation and the task, in part due to a large variability of single trial timecourses. Across ROIs, however, a frequent interaction was observed in both dPul and LIP stimulation datasets. The direction of the effect was not consistent however between the two animals – while in monkey C, according to our initial prediction, the contraversive saccade task condition

produced the strongest microstimulation enhancement, it was not the case for monkey C (ipsiversive or fixation > contraversive).

We thus investigated how dPul and LIP microstimulation effects in the different tasks might be influenced by the initial spatial selectivity of stimulation-activated ROIs. Here we found a consistent relationship between the degree of contraversive selectivity and the strength of microstimulation-elicited activation, across ROIs. Specifically, the higher the contraversive selectivity of an ROI, the higher is the stimulation effect when monkeys are cued to perform a saccade into the ipsiversive hemifield. Conversely, such relationship was often inverted for the contraversive saccade task, during which ipsiversively-selective ROIs exhibited a higher stimulation effect. This "X-shape" interaction between the task condition and the initial spatial selectivity is reminiscent of FEF stimulation studies described above, showing that microstimulation effects in visual cortices are stronger with weaker visual stimuli and in task conditions not favored by the microstimulation-activated ROI (Ekstrom et al., 2009; Premereur et al., 2013). A plausible explanation is that strong initial task-related response leaves less "room" for modulation of neuronal activity (or a hemodynamically-coupled BOLD signal) by microstimulation before saturation is reached.

In line with these observations, the model that best described microstimulation effect amplitudes across ROIs was the additive mechanism, but with strength of the additive factor scaled down by the initial response amplitude. Across datasets, this model provided a better fit than a simple additive model, and was by far better than the multiplicative model. A simple qualitative simulation was able to reproduce the observed "X-shape" interaction pattern between the task condition and the initial spatial selectivity.

It remains to be elucidated how these microstimulation-elicited modulations of neural signals might be relevant for spatial behavior. Our simple single option instructed memory saccade task with a long delay, the prolonged microstimulation paradigm – with the last stimulation train ending 800 ms before the end of the delay period, and the analysis of only successful trials, were not designed to address stimulation-contingent changes of behavior. Faster trials with shorter stimulation periods and more demanding tasks involving attentional selection or a choice between different response options will be required to study behaviorally-relevant effects (Arsenault et al., 2014; Dominguez-Vargas et al., 2017). On the other hand, the time-resolved design of our experiment might afford the possibility to look more closely into the dynamics of the microstimulation effects, e.g. by analyzing transient cue response, the subsequent memory delay, and the saccade execution-related effects separately.

The ROIs in the present analysis were selected solely on the basis of positive activation by the microstimulation pooled over all task conditions, regardless of their task responsivity and other properties such as spatial tuning. One can argue that such selection, while most unbiased in a sense of identifying basic microstimulation-elicited enhancement, favors ROIs without strong task dependence – although the inspection of single ROI timecourses and response amplitudes indicated that in some ROIs a strong positive activation in only one of three task conditions was enough to have a significant effect. The strong overlap of stimulation-elicited activations and task-related activations in trials without stimulation, or even in no stimulation datasets (data not shown) also suggests that we did not miss any major relevant regions involved in the task-related processing. However, the microstimulation-driven mapping and ROI selection might be not most informative for deriving conclusions about global changes in spatial tuning due to microstimulation. For instance, the stimulation effect in the stimulated hemisphere was stronger and involved considerably more regions that in the opposite hemisphere. This could lead to changes of interhemispheric balance, and an overall stronger potentiation of contraversive spatial representations. Using a predefined, symmetrical set of ROIs across the two hemispheres might help addressing these crucial outstanding questions.

Authors contributions

IK: Conceptualization, Methodology, Software, Formal analysis, Resources, Data curation, Writing – Original Draft, Writing – Review & Editing, Visualization, Project administration, Supervision, Funding acquisition; **LG:** Conceptualization, Methodology, Software, Formal analysis, Investigation, Writing – Original Draft, Visualization; **ES:** Methodology, Software, Investigation, **MW:** Conceptualization, Methodology, Resources, Writing – Original Draft, Writing – Review & Editing, Supervision, Funding acquisition

Declarations of interest

Acknowledgements

We thank Dr. Vladislav Kozyrev for helping to collect a subset of the MRI data, Dr. Sebastian Moeller for sharing the design of the MR-compatible electrode drive and helping with the microstimulation setup, Dr. Peter Dechent and Dr. Juergen Baudewig for helping with setting up MRI data acquisition, and Dr. Jochen Weber for providing support for NeuroElf. We thank Ira Panolias, Dr. Daniela Lazzarini, Sina Plümer, Klaus Heisig, Leonore Burchardt, and Dirk Prüße for technical support. We also thank Stefan Treue, Alexander Gail, Hansjörg Scherberger, members of the Decision and Awareness Group, the Sensorimotor Group, and the Cognitive Neuroscience Laboratory for helpful discussions.

Funding

This research was supported by the Hermann and Lilly Schilling Foundation, German Research Foundation (DFG) grants WI 4046/1-1 and Research Unit GA1475-B4, KA 3726/2-1, and the Primate Platform of the DFG Center for Nanoscale Microscopy and Molecular Physiology of the Brain (CNMPB).

References

Adams, M.M., Hof, P.R., Gattass, R., Webster, M.J., Ungerleider, L.G., 2000. Visual cortical projections and chemoarchitecture of macaque monkey pulvinar. J Comp Neurol 419, 377–93.

Andersen, R.A., Asanuma, C., Essick, G., Siegel, R.M., 1990. Corticocortical connections of anatomically and physiologically defined subdivisions within the inferior parietal lobule. J Comp Neurol 296, 65–113.

Andersen, R.A., Cui, H., 2009. Intention, action planning, and decision making in parietal-frontal circuits. Neuron 63, 568–83.

Arcaro, M.J., Pinsk, M.A., Chen, J., Kastner, S., 2018. Organizing principles of pulvino-cortical functional coupling in humans. Nature Communications 9, 5382. https://doi.org/10.1038/s41467-018-07725-6

Arcaro, M.J., Pinsk, M.A., Kastner, S., 2015. The Anatomical and Functional Organization of the Human Visual Pulvinar. Journal of Neuroscience 35, 9848–9871. https://doi.org/10.1523/JNEUROSCI.1575-14.2015

Arend, I., Machado, L., Ward, R., McGrath, M., Ro, T., Rafal, R.D., 2008a. The role of the human pulvinar in visual attention and action: evidence from temporal-order judgment, saccade decision, and antisaccade tasks. Prog Brain Res 171, 475–83.

Arend, I., Rafal, R., Ward, R., 2008b. Spatial and temporal deficits are regionally dissociable in patients with pulvinar lesions. Brain 131, 2140–2152. https://doi.org/10.1093/brain/awn135

Arsenault, J.T., Rima, S., Stemmann, H., Vanduffel, W., 2014. Role of the Primate Ventral Tegmental Area in Reinforcement and Motivation. Current Biology 24, 1347–1353. https://doi.org/10.1016/j.cub.2014.04.044

Asanuma, C., Andersen, R.A., Cowan, W.M., 1985. The thalamic relations of the caudal inferior parietal lobule and the lateral prefrontal cortex in monkeys: divergent cortical projections from cell clusters in the medial pulvinar nucleus. J. Comp. Neurol. 241, 357–381. https://doi.org/10.1002/cne.902410309

Baldwin, M.K.L., Balaram, P., Kaas, J.H., 2017. The evolution and functions of nuclei of the visual pulvinar in primates. Journal of Comparative Neurology 525, 3207–3226. https://doi.org/10.1002/cne.24272

Baldwin, M.K.L., Bourne, J.A., 2017. The Evolution of Subcortical Pathways to the Extrastriate Cortex, in: Evolution of Nervous Systems. Elsevier, pp. 165–185. https://doi.org/10.1016/B978-0-12-804042-3.00081-6

Baleydier, C., Mauguiere, F., 1985. Anatomical evidence for medial pulvinar connections with the posterior cingulate cortex, the retrosplenial area, and the posterior parahippocampal gyrus in monkeys. J Comp Neurol 232, 219–28.

Barash, S., Bracewell, R.M., Fogassi, L., Gnadt, J.W., Andersen, R.A., 1991. Saccade-related activity in the lateral intraparietal area. II. Spatial properties. J Neurophysiol 66, 1109–24.

Ben Hamed, S., Duhamel, J.R., Bremmer, F., Graf, W., 2001. Representation of the visual field in the lateral intraparietal area of macaque monkeys: a quantitative receptive field analysis. Exp Brain Res 140, 127–144.

Bender, D.B., 1981. Retinotopic organization of macaque pulvinar. J Neurophysiol 46, 672–93.

Bender, D.B., Butter, C.M., 1987. Comparison of the effects of superior colliculus and pulvinar lesions on visual search and tachistoscopic pattern discrimination in monkeys. Exp Brain Res 69, 140–54.

Bender, D.B., Youakim, M., 2001. Effect of attentive fixation in macaque thalamus and cortex. J. Neurophysiol. 85, 219–234.

Benevento, L.A., Miller, J., 1981. Visual responses of single neurons in the caudal lateral pulvinar of the macaque monkey. The Journal of Neuroscience 1, 1268–1278.

Benevento, L.A., Port, J.D., 1995. Single neurons with both form/color differential responses and saccade-related responses in the nonretinotopic pulvinar of the behaving macaque monkey. Vis Neurosci 12, 523–44.

Benevento, L.A., Standage, G.P., 1983. The organization of projections of the retinorecipient and nonretinorecipient nuclei of the pretectal complex and layers of the superior colliculus to the lateral pulvinar and medial pulvinar in the macaque monkey. J. Comp. Neurol. 217, 307–336. https://doi.org/10.1002/cne.902170307

Berman, R.A., Wurtz, R.H., 2011. Signals Conveyed in the Pulvinar Pathway from Superior Colliculus to Cortical Area MT. Journal of Neuroscience 31, 373–384. https://doi.org/10.1523/JNEUROSCI.4738-10.2011

Bisley, J.W., Goldberg, M.E., 2010. Attention, intention, and priority in the parietal lobe. Annu. Rev. Neurosci. 33, 1–21. https://doi.org/10.1146/annurev-neuro-060909-152823

Blatt, G.J., Andersen, R.A., Stoner, G.R., 1990. Visual receptive field organization and cortico-cortical connections of the lateral intraparietal area (area LIP) in the macaque. J. Comp. Neurol. 299, 421–445. https://doi.org/10.1002/cne.902990404

Bogadhi, A.R., Bollimunta, A., Leopold, D.A., Krauzlis, R.J., 2019. Spatial Attention Deficits Are Causally Linked to an Area in Macaque Temporal Cortex. Curr. Biol. https://doi.org/10.1016/j.cub.2019.01.028

Bogadhi, A.R., Bollimunta, A., Leopold, D.A., Krauzlis, R.J., 2018. Brain regions modulated during covert visual attention in the macaque. Scientific Reports 8. https://doi.org/10.1038/s41598-018-33567-9

Brainard, D.H., 1997. The Psychophysics Toolbox. Spatial Vision 10, 433–436.

Bridge, H., Leopold, D.A., Bourne, J.A., 2015. Adaptive Pulvinar Circuitry Supports Visual Cognition. Trends Cogn Sci. https://doi.org/10.1016/j.tics.2015.10.003

Bruce, C., Desimone, R., Gross, C.G., 1981. Visual properties of neurons in a polysensory area in superior temporal sulcus of the macaque. Journal of neurophysiology 46, 369–84.

Calabrese, E., Badea, A., Coe, C.L., Lubach, G.R., Shi, Y., Styner, M.A., Johnson, G.A., 2015. A diffusion tensor MRI atlas of the postmortem rhesus macaque brain. NeuroImage 117, 408–416. https://doi.org/10.1016/j.neuroimage.2015.05.072

Cappe, C., Morel, A., Barone, P., Rouiller, E.M., 2009. The Thalamocortical Projection Systems in Primate: An Anatomical Support for Multisensory and Sensorimotor Interplay. Cerebral Cortex 19, 2025–2037. https://doi.org/10.1093/cercor/bhn228

Cappe, C., Morel, A., Rouiller, E.M., 2007. Thalamocortical and the dual pattern of corticothalamic projections of the posterior parietal cortex in macaque monkeys. Neuroscience 146, 1371–1387. https://doi.org/10.1016/j.neuroscience.2007.02.033

Caspari, N., Janssens, T., Mantini, D., Vandenberghe, R., Vanduffel, W., 2015. Covert shifts of spatial attention in the macaque monkey. The Journal of neuroscience : the official journal of the Society for Neuroscience 35, 7695–714. https://doi.org/10.1523/JNEUROSCI.4383-14.2015

Chen, M., Li, B., Guang, J., Wei, L., Wu, S., Liu, Y., Zhang, M., 2016. Two subdivisions of macaque LIP process visual-oculomotor information differently. Proc Natl Acad Sci U S A 113, E6263–E6270. https://doi.org/10.1073/pnas.1605879113

Christopoulos, V.N., Kagan, I., Andersen, R.A., 2018. Lateral intraparietal area (LIP) is largely effector-specific in free-choice decisions. Sci Rep 8, 8611. https://doi.org/10.1038/s41598-018-26366-9

Colby, C.L., Goldberg, M.E., 1999. Space and attention in parietal cortex. Annu Rev Neurosci 22, 319–49. https://doi.org/10.1146/annurev.neuro.22.1.319

Dai, J., Brooks, D.I., Sheinberg, D.L., 2014. Optogenetic and Electrical Microstimulation Systematically Bias Visuospatial Choice in Primates. Current Biology 24, 63–69. https://doi.org/10.1016/j.cub.2013.11.011

Dominguez-Vargas, A.-U., Schneider, L., Wilke, M., Kagan, I., 2017. Electrical Microstimulation of the Pulvinar Biases Saccade Choices and Reaction Times in a Time-Dependent Manner. J. Neurosci. 37, 2234–2257. https://doi.org/10.1523/JNEUROSCI.1984-16.2016

Ekstrom, L.B., Roelfsema, P.R., Arsenault, J.T., Bonmassar, G., Vanduffel, W., 2008. Bottom-up dependent gating of frontal signals in early visual cortex. Science 321, 414–7.

Ekstrom, L.B., Roelfsema, P.R., Arsenault, J.T., Kolster, H., Vanduffel, W., 2009. Modulation of the contrast response function by electrical microstimulation of the macaque frontal eye field. J Neurosci 29, 10683–94.

Fiebelkorn, I.C., Pinsk, M.A., Kastner, S., 2019. The mediodorsal pulvinar coordinates the macaque fronto-parietal network during rhythmic spatial attention. Nature Communications 10, 215. https://doi.org/10.1038/s41467-018-08151-4

Field, C.B., Johnston, K., Gati, J.S., Menon, R.S., Everling, S., 2008. Connectivity of the primate superior colliculus mapped by concurrent microstimulation and event-related FMRI. PLoS One 3, e3928. https://doi.org/10.1371/journal.pone.0003928

Glimcher, P.W., 2003. The neurobiology of visual-saccadic decision making. Annu Rev Neurosci 26, 133–79.

Gold, J.I., Shadlen, M.N., 2007. The Neural Basis of Decision Making. Annual Review of Neuroscience 30, 535–574. https://doi.org/10.1146/annurev.neuro.29.051605.113038

Grieve, K.L., Acuña, C., Cudeiro, J., 2000. The primate pulvinar nuclei: vision and action. Trends Neurosci. 23, 35–39.

Gutierrez, C., Cola, M.G., Seltzer, B., Cusick, C., 2000. Neurochemical and connectional organization of the dorsal pulvinar complex in monkeys. Journal of Comparative Neurology 419, 61–86.

Halassa, M.M., Kastner, S., 2017. Thalamic functions in distributed cognitive control. Nature Neuroscience 20, 1669–1679. https://doi.org/10.1038/s41593-017-0020-1

Hanks, T.D., Ditterich, J., Shadlen, M.N., 2006. Microstimulation of macaque area LIP affects decision-making in a motion discrimination task. Nat Neurosci 9, 682–9.

Hardy, S.P., Lynch, J.C., 1992. The spatial distribution of pulvinar neurons that project to two subregions of the inferior parietal lobule in the macaque. Cerebral Cortex 2, 217–230.

Harting, J.K., Huerta, M.F., Frankfurter, A.J., Strominger, N.L., Royce, G.J., 1980. Ascending pathways from the monkey superior colliculus: An autoradiographic analysis. J. Comp. Neurol. 192, 853–882. https://doi.org/10.1002/cne.901920414

Hedreen, J.C., Yin, T.C.T., 1981. Homotopic and heterotopic callosal afferents of caudal inferior parietal lobule in Macaca mulatta. Journal of Comparative Neurology 197, 605–621. https://doi.org/10.1002/cne.901970405

Homman-Ludiye, J., Mundinano, I.C., Kwan, W.C., Bourne, J.A., 2020. Extensive Connectivity Between the Medial Pulvinar and the Cortex Revealed in the Marmoset Monkey. Cereb. Cortex 30, 1797–1812. https://doi.org/10.1093/cercor/bhz203

Honey, C.J., Sporns, O., Cammoun, L., Gigandet, X., Thiran, J.P., Meuli, R., Hagmann, P., 2009. Predicting human resting-state functional connectivity from structural connectivity. PNAS 106, 2035–2040. https://doi.org/10.1073/pnas.0811168106

Itoh, K., Mizuno, N., 1979. A cerebello-pulvinar projection in the cat as visualized by the use of anterograde transport of horseradish peroxidase. Brain Res. 171, 131–134. https://doi.org/10.1016/0006-8993(79)90738-8

Jones, E.G., 2007. The lateral posterior and pulvinar nuclei., in: The Thalamus. Cambridge University Press, Cambridge, pp. 1009–1071.

Kaas, J.H., Lyon, D.C., 2007. Pulvinar contributions to the dorsal and ventral streams of visual processing in primates. Brain Research Reviews 55, 285–296. https://doi.org/10.1016/j.brainresrev.2007.02.008

Kagan, I., Iyer, A., Lindner, A., Andersen, R.A., 2010. Space representation for eye movements is more contralateral in monkeys than in humans. Proc Natl Acad Sci U S A 107, 7933–8. https://doi.org/10.1073/pnas.1002825107

Karnath, H.O., 2015. Spatial attention systems in spatial neglect. Neuropsychologia 75, 61–73. https://doi.org/10.1016/j.neuropsychologia.2015.05.019

Karnath, H.O., 2001. New insights into the functions of the superior temporal cortex. Nat Rev Neurosci 2, 568–76.

Karnath, H.O., Himmelbach, M., Rorden, C., 2002. The subcortical anatomy of human spatial neglect: putamen, caudate nucleus and pulvinar. Brain 125, 350–60.

Katz, L.N., Yates, J.L., Pillow, J.W., Huk, A.C., 2016. Dissociated functional significance of decision-related activity in the primate dorsal stream. Nature 535, 285–288. https://doi.org/10.1038/nature18617

Klink, P.C., et al. Combined Brain Perturbation and Neuroimaging in Non-human Primates. NeuroImage, this issue.

Komura, Y., Nikkuni, A., Hirashima, N., Uetake, T., Miyamoto, A., 2013. Responses of pulvinar neurons reflect a subject's confidence in visual categorization. Nature Neuroscience 16, 749–755. https://doi.org/10.1038/nn.3393

Kubanek, J., Li, J.M., Snyder, L.H., 2015. Motor role of parietal cortex in a monkey model of hemispatial neglect. Proceedings of the National Academy of Sciences of the United States of America. https://doi.org/10.1073/pnas.1418324112

Lanz, F., Moret, V., Ambett, R., Cappe, C., Rouiller, E.M., Loquet, G., 2017. Distant heterotopic callosal connections to premotor cortex in non-human primates. Neuroscience 344, 56–66. https://doi.org/10.1016/j.neuroscience.2016.12.035

Lewis, J.W., Van Essen, D.C., 2000. Mapping of architectonic subdivisions in the macaque monkey, with emphasis on parieto-occipital cortex. J Comp Neurol 428, 79–111. https://doi.org/10.1002/1096-9861(20001204)428:1<79::AID-CNE7>3.0.CO;2-Q

Li, C.S., Mazzoni, P., Andersen, R.A., 1999. Effect of reversible inactivation of macaque lateral intraparietal area on visual and memory saccades. J Neurophysiol 81, 1827–38.

Liu, Y., Yttri, E.A., Snyder, L.H., 2010. Intention and attention: different functional roles for LIPd and LIPv. Nature neuroscience 13, 495–500. https://doi.org/10.1038/nn.2496

Logothetis, N.K., Augath, M., Murayama, Y., Rauch, A., Sultan, F., Goense, J., Oeltermann, A., Merkle, H., 2010. The effects of electrical microstimulation on cortical signal propagation. Nat. Neurosci. 13, 1283–1291. https://doi.org/10.1038/nn.2631

Luh, K.E., Butter, C.M., Buchtel, H.A., 1986. Impairments in orienting to visual stimuli in monkeys following unilateral lesions of the superior sulcal polysensory cortex. Neuropsychologia 24, 461–70.

Matsui, T., Koyano, K.W., Tamura, K., Osada, T., Adachi, Y., Miyamoto, K., Chikazoe, J., Kamigaki, T., Miyashita, Y., 2012. FMRI activity in the macaque cerebellum evoked by intracortical microstimulation of the primary somatosensory cortex: evidence for polysynaptic propagation. PLoS One 7, e47515. https://doi.org/10.1371/journal.pone.0047515

Matsui, T., Tamura, K., Koyano, K.W., Takeuchi, D., Adachi, Y., Osada, T., Miyashita, Y., 2011. Direct comparison of spontaneous functional connectivity and effective connectivity measured by intracortical microstimulation: an fMRI study in macaque monkeys. Cereb Cortex 21, 2348–56. https://doi.org/10.1093/cercor/bhr019

Mesulam, M.M., 1999. Spatial attention and neglect: parietal, frontal and cingulate contributions to the mental representation and attentional targeting of salient extrapersonal events. Philos Trans R Soc Lond B Biol Sci 354, 1325–46.

Moeller, S., Freiwald, W.A., Tsao, D.Y., 2008. Patches with links: a unified system for processing faces in the macaque temporal lobe. Science 320, 1355–9.

Mundinano, I.-C., Fox, D.M., Kwan, W.C., Vidaurre, D., Teo, L., Homman-Ludiye, J., Goodale, M.A., Leopold, D.A., Bourne, J.A., 2018. Transient visual pathway critical for normal development of primate grasping behavior. Proc. Natl. Acad. Sci. U.S.A. 115, 1364–1369. https://doi.org/10.1073/pnas.1717016115

Murayama, Y., Augath, M., Logothetis, N.K., 2011. Activation of SC during electrical stimulation of LGN: retinal antidromic stimulation or corticocollicular activation? Magn Reson Imaging 29, 1351–7. https://doi.org/10.1016/j.mri.2011.08.002

Murris, S.R., Arsenault, J.T., Vanduffel, W., 2020. Frequency- and State-Dependent Network Effects of Electrical Stimulation Targeting the Ventral Tegmental Area in Macaques. Cereb Cortex 30, 4281–4296. https://doi.org/10.1093/cercor/bhaa007

Ohayon, S., Tsao, D.Y., 2012. MR-guided stereotactic navigation. J Neurosci Methods 204, 389–97. https://doi.org/10.1016/j.jneumeth.2011.11.031

Pare, M., Wurtz, R.H., 1997. Monkey posterior parietal cortex neurons antidromically activated from superior colliculus. J Neurophysiol 78, 3493–7.

Patel, G.H., Shulman, G.L., Baker, J.T., Akbudak, E., Snyder, A.Z., Snyder, L.H., Corbetta, M., 2010. Topographic organization of macaque area LIP. Proc Natl Acad Sci U S A 107, 4728–33. https://doi.org/10.1073/pnas.0908092107

Patel, G.H., Yang, D., Jamerson, E.C., Snyder, L.H., Corbetta, M., Ferrera, V.P., 2015. Functional evolution of new and expanded attention networks in humans. Proceedings of the National Academy of Sciences of the United States of America 112, 9454–9. https://doi.org/10.1073/pnas.1420395112

Petersen, S.E., Robinson, D.L., Keys, W., 1985. Pulvinar nuclei of the behaving rhesus monkey: visual responses and their modulation. Journal of neurophysiology 54, 867–886.

Petersen, S.E., Robinson, D.L., Morris, J.D., 1987. Contributions of the pulvinar to visual spatial attention. Neuropsychologia 25, 97–105.

Petkov, C.I., Kikuchi, Y., Milne, A.E., Mishkin, M., Rauschecker, J.P., Logothetis, N.K., 2015. Different forms of effective connectivity in primate frontotemporal pathways. Nat Commun 6, 6000. https://doi.org/10.1038/ncomms7000

Premereur, E., Janssen, P., Vanduffel, W., 2013. FEF-microstimulation causes task-dependent modulation of occipital fMRI activity. NeuroImage 67, 42–50. https://doi.org/10.1016/j.neuroimage.2012.11.017

Premereur, E., Van Dromme, I.C., Romero, M.C., Vanduffel, W., Janssen, P., 2015. Effective Connectivity of Depth-Structure–Selective Patches in the Lateral Bank of the Macaque Intraparietal Sulcus. PLoS biology 13, e1002072–e1002072.

Rafal, R., McGrath, M., Machado, L., Hindle, J., 2004. Effects of lesions of the human posterior thalamus on ocular fixation during voluntary and visually triggered saccades. J Neurol Neurosurg Psychiatry 75, 1602–6. https://doi.org/10.1136/jnnp.2003.017038

Robinson, D.L., Petersen, S.E., Keys, W., 1986. Saccade-related and visual activities in the pulvinar nuclei of the behaving rhesus monkey. Experimental Brain Research 62. https://doi.org/10.1007/BF00236042

Rodrigo-Angulo, M.L., Reinoso-Suarez, F., 1984. Cerebellar projections to the lateral posterior-pulvinar thalamic complex in the cat. Brain Res. 322, 172–176. https://doi.org/10.1016/0006-8993(84)91200-9

Rohlfing, T., Kroenke, C.D., Sullivan, E.V., Dubach, M.F., Bowden, D.M., Grant, K.A., Pfefferbaum, A., 2012. The INIA19 Template and NeuroMaps Atlas for Primate Brain Image Parcellation and Spatial Normalization. Front Neuroinform 6, 27. https://doi.org/10.3389/fninf.2012.00027

Romanski, L.M., Giguere, M., Bates, J.F., Goldman-Rakic, P.S., 1997. Topographic organization of medial pulvinar connections with the prefrontal cortex in the rhesus monkey. J. Comp. Neurol. 379, 313–332.

Rosenberg, D.S., Mauguiere, F., Catenoix, H., Faillenot, I., Magnin, M., 2009. Reciprocal Thalamocortical Connectivity of the Medial Pulvinar: A Depth Stimulation and Evoked Potential Study in Human Brain. Cerebral Cortex 19, 1462–1473. https://doi.org/10.1093/cercor/bhn185

Saalmann, Y.B., Pinsk, M.A., Wang, L., Li, X., Kastner, S., 2012. The Pulvinar Regulates Information Transmission Between Cortical Areas Based on Attention Demands. Science 337, 753–756. https://doi.org/10.1126/science.1223082

Saleem, K., Logothetis, N.K., 2006. A Combined MRI and Histology Atlas of the Rhesus Monkey Brain. Academic Press, New York.

Scalaidhe, S.P., Albright, T.D., Rodman, H.R., Gross, C.G., 1995. Effects of superior temporal polysensory area lesions on eye movements in the macaque monkey. J Neurophysiol 73, 1–19.

Schilling, K.G., Gao, Y., Stepniewska, I., Janve, V., Landman, B.A., Anderson, A.W., 2019. Histologically derived fiber response functions for diffusion MRI vary across white matter fibers-An ex vivo validation study in the squirrel monkey brain. NMR Biomed 32, e4090. https://doi.org/10.1002/nbm.4090

Schneider, L., Dominguez-Vargas, A.-U., Gibson, L., Kagan, I., Wilke, M., 2020. Eye position signals in the dorsal pulvinar during fixation and goal-directed saccades. J. Neurophysiol. 123, 367–391. https://doi.org/10.1152/jn.00432.2019

Seltzer, B., Pandya, D.N., 1994. Parietal, temporal, and occipital projections to cortex of the superior temporal sulcus in the rhesus monkey: a retrograde tracer study. J Comp Neurol 343, 445–63. https://doi.org/10.1002/cne.903430308

Sherman, S.M., Guillery, R.W., 2011. Distinct functions for direct and transthalamic corticocortical connections. Journal of Neurophysiology 106, 1068–1077. https://doi.org/10.1152/jn.00429.2011

Shipp, S., 2003. The functional logic of cortico-pulvinar connections. Philosophical Transactions of the Royal Society B: Biological Sciences 358, 1605–1624. https://doi.org/10.1098/rstb.2002.1213

Stepniewska, I., Preuss, T.M., Kaas, J.H., 1994. Thalamic connections of the primary motor cortex (M1) of owl monkeys. The Journal of Comparative Neurology 349, 558–582. https://doi.org/10.1002/cne.903490405

Storm, J.F., Boly, M., Casali, A.G., Massimini, M., Olcese, U., Pennartz, C.M.A., Wilke, M., 2017. Consciousness Regained: Disentangling Mechanisms, Brain Systems, and Behavioral Responses. The Journal of Neuroscience 37, 10882–10893. https://doi.org/10.1523/JNEUROSCI.1838-17.2017

Sultan, F., Augath, M., Hamodeh, S., Murayama, Y., Oeltermann, A., Rauch, A., Thier, P., 2012. Unravelling cerebellar pathways with high temporal precision targeting motor and extensive sensory and parietal networks. Nature Communications 3. https://doi.org/10.1038/ncomms1912

Thier, P., Andersen, R.A., 1998. Electrical microstimulation distinguishes distinct saccade-related areas in the posterior parietal cortex. Journal of Neurophysiology 80, 1713–1735.

Tolias, A.S., Sultan, F., Augath, M., Oeltermann, A., Tehovnik, E.J., Schiller, P.H., Logothetis, N.K., 2005. Mapping cortical activity elicited with electrical microstimulation using FMRI in the macaque. Neuron 48, 901–11.

Van der Stigchel, S., Arend, I., van Koningsbruggen, M.G., Rafal, R.D., 2010. Oculomotor integration in patients with a pulvinar lesion. Neuropsychologia 48, 3497–3504. https://doi.org/10.1016/j.neuropsychologia.2010.07.035

Wardak, C., Olivier, E., Duhamel, J.R., 2004. A deficit in covert attention after parietal cortex inactivation in the monkey. Neuron 42, 501–8.

Wardak, C., Olivier, E., Duhamel, J.-R., 2002. Saccadic target selection deficits after lateral intraparietal area inactivation in monkeys. J. Neurosci. 22, 9877–9884.

Watson, R.T., Valenstein, E., Day, A., Heilman, K.M., 1994. Posterior neocortical systems subserving awareness and neglect. Neglect associated with superior temporal sulcus but not area 7 lesions. Arch Neurol 51, 1014–21.

Wilke, M., Dechent, P., Bähr, M., 2017. Sarcoidosis Manifestion Centered on the Thalamic Pulvinar Leading to Persistent Astasia. Mov Disord Clin Pract 4, 898–900. https://doi.org/10.1002/mdc3.12544

Wilke, M., Kagan, I., Andersen, R.A., 2013. Effects of pulvinar inactivation on spatial decision-making between equal and asymmetric reward options. J Cogn Neurosci 25, 1270–1283. https://doi.org/10.1162/jocn_a_00399

Wilke, M., Kagan, I., Andersen, R.A., 2012. Functional imaging reveals rapid reorganization of cortical activity after parietal inactivation in monkeys. Proceedings of the National Academy of Sciences 109, 8274–8279. https://doi.org/10.1073/pnas.1204789109

Wilke, Melanie, Kagan, I., Andersen, R.A., 2010. Pulvinar inactivation alters cortical responses during spatial decision making, in: 280.11/SS19 Neuroscience Meeting Planner. Presented at the Society for Neuroscience Annual Meeting, San Diego, CA.

Wilke, M., Schneider, L., Dominguez-Vargas, A.-U., Schmidt-Samoa, C., Miloserdov, K., Nazzal, A., Dechent, P., Cabral-Calderin, Y., Scherberger, H., Kagan, I., Bähr, M., 2018. Reach and grasp deficits following damage to the dorsal pulvinar. Cortex 99, 135–149. https://doi.org/10.1016/j.cortex.2017.10.011

Wilke, M., Turchi, J., Smith, K., Mishkin, M., Leopold, D.A., 2010. Pulvinar Inactivation Disrupts Selection of Movement Plans. Journal of Neuroscience 30, 8650–8659. https://doi.org/10.1523/JNEUROSCI.0953-10.2010

Wu, J., Dubach, M.F., Robertson, J.E., 2000. Primate Brain Maps: Structure of the Macaque Brain: A Laboratory Guide with Original Brain Sections, Printed Atlas and Electronic Templates for Data and Schematics (including CD-ROM)., 1st Edition. ed. Elsevier Science, Amsterdam ; New York.

Wurtz, R.H., Hikosaka, O., 1986. Role of the basal ganglia in the initiation of saccadic eye movements. Prog Brain Res 64, 175–90. https://doi.org/10.1016/S0079-6123(08)63412-3

Wurtz, R.H., McAlonan, K., Cavanaugh, J., Berman, R.A., 2011. Thalamic pathways for active vision. Trends Cogn Sci 15, 177–84. https://doi.org/10.1016/j.tics.2011.02.004

Yeterian, E.H., Pandya, D.N., 1991. Corticothalamic connections of the superior temporal sulcus in rhesus monkeys. Exp Brain Res 83, 268–84.

Yeterian, E.H., Pandya, D.N., 1989. Thalamic connections of the cortex of the superior temporal sulcus in the rhesus monkey. J Comp Neurol 282, 80–97.

Yeterian, E.H., Pandya, D.N., 1985. Corticothalamic connections of the posterior parietal cortex in the rhesus monkey. J Comp Neurol 237, 408–26. https://doi.org/10.1002/cne.902370309

Zhou, H., Schafer, R.J., Desimone, R., 2016. Pulvinar-Cortex Interactions in Vision and Attention. Neuron 89, 209–220. https://doi.org/10.1016/j.neuron.2015.11.034