FGF8-mediated signaling regulates tooth developmental pace during odontogenesis

Chensheng Lin[1], Ningsheng Ruan[1], Linjun Li[1], Yibin Chen[1], Xiaoxiao Hu[1], YiPing Chen[2], Xuefeng Hu[1]*, Yanding Zhang[1]*

[1]Fujian Key Laboratory of Developmental and Neural Biology & Southern Center for Biomedical Research, College of Life Sciences, Fujian Normal University, Fuzhou, Fujian, 350117, P.R. China.

[2]Department of Cell and Molecular Biology, Tulane University, New Orleans, LA 70118, USA.

Running title: FGF8 regulates tooth developmental pace

*Correspondence to:

Yanding Zhang, Ph.D.

College of Life Sciences, Fujian Normal University (Qishan Campus), Fuzhou, Fujian, 350117, P.R. China.

Phone: +86-13706963408 (tel); E-mail: ydzhang@fjnu.edu.cn

or

Xuefeng Hu, Ph.D.

22   College of Life Sciences, Fujian Normal University (Qishan Campus), Fuzhou, Fujian,

23   350117, P.R. China.

24   Phone: +86-13055734898 (tel) ; E-mail: bioxfh@fjnu.edu.cn

Conflicts of interest

The authors declare no conflicts of interest.

Data Availability Statement

All the data are available within the article or its supplementary materials.

Abstract

The developing human and mouse teeth constitute an ideal model system to study the regulatory mechanism underlying organ growth control due to the fact that their teeth share highly conserved and well-characterized developmental processes and their developmental tempo vary notably. In the current study, we manipulated heterogenous recombination between human and mouse dental tissues and demonstrate that the dental mesenchyme dominates the tooth developmental tempo and FGF8 could be a critical player during this developmental process. Forced activation of FGF8 signaling in the dental mesenchyme of mice promoted cell proliferation, prevented cell apoptosis via p38 and perhaps PI3K-Akt intracellular signaling, and impelled the transition of the cell cycle from G1- to S-phase in the tooth germ, resulting in the slowdown of the tooth developmental pace. Our results provide compelling evidence that extrinsic signals can profoundly affect tooth developmental tempo and the dental mesenchymal FGF8 could be a pivotal factor in controlling developmental pace in a non-cell-autonomous manner during mammalian odontogenesis.

Key words: FGF8, tooth, developmental pace, non-cell-autonomous.

Introduction

Although mammals possess an almost identical body plan with morphologically and functionally similar organs and systems, their developmental programs and body sizes vary greatly among different species. The intrinsic developmental programs, together with extracellular signals, control the individual organ size and thus body size [1]. Generally, larger-bodied species own longer gestation lengths with slower developmental tempos to acquire larger body sizes [2,3]. Obviously, developmental tempo and organ size are two closely interrelated variables during individual growth (i.e. embryonic development and postnatal growth) and morphogenesis, both of which must be tightly harmonized to ensure the correct establishment of body plan and therefore are the result of a precisely controlled process.

The mechanisms that regulate organ size have been well investigated in non-mammalian models, particularly in *Drosophila melanogaster*, in which roles of TOR (target of rapamycin), insulin/IGF (insulin-like growth factor), Ras/Raf/MAPK, JNK (c-Jun N-terminal kinase), and Hippo signaling pathways are well defined [4]. In mammals, Hippo and IGF pathways have been evidenced to be involved in controlling organ size [5-10]. Additionally, several conserved families of secreted growth factors, including FGF (fibroblast growth factor), BMP (bone morphogenetic protein), and TGF-β (transforming growth factor-β), are deemed to be involved in the regulation of individual growth due to their ability to modulate cell proliferation and apoptosis during embryonic development [4,11,12]. However, the regulatory mechanisms of developmental pace in mammals are remain largely unknown.

Mammalian teeth have long served as a model organ for studying fundamental

questions in organ development on the molecular basis [13]. The main features of tooth

morphogenesis are conserved in the mammalian lineage [14-16], as they are all formed by

sequential and reciprocal interactions between epithelium and mesenchyme and share

similar processes during tooth development [15]. Despite these considerable homologies,

mammalian dentition undergoes remarkable morphological diversification, displaying

great variations not only in tooth shape, number, and size but also in developmental pace

[13,15,17-19]. Obviously, the period of tooth development in humans takes about 400 days

from the initiation stage at embryonic 6th week to tooth eruption at postnatal 6th month.

Whereas, this period in mice takes only about 20 days from the determination of tooth

forming sites at E10.5 to tooth eruption at postnatal 10th day [13]. Thus, the human and

mouse embryonic tooth provides an ideal model system for investigating the regulatory

mechanism underlying the developmental tempo, however, the molecular mechanism that

regulates the developmental pace of mammalian teeth remains elusive.

It is generally believed that diversified forms of organ morphogenesis (including the

developmental tempo) among species are achieved by tinkering conserved signaling

pathways instead of proposing novel ones during development [20-22]. The difference in

organismal growth rate and organ or body size between humans and mice primarily reflect

the difference in cell number rather than the cell size [1]. Many lines of evidence from

mouse models demonstrate that numerous growth factors of conserved families, such as

FGF, BMP, SHH, and Wnt, have been shown to play critical roles in tooth morphogenesis

by regulating cell proliferation, differentiation, and apoptosis during odontogenesis

[15,16,21,23-27], suggesting their potential roles in the regulation of the developmental tempo.

Here, we provide compelling evidence that it is the dental mesenchyme that dominates

tooth developmental tempo in a non-cell-autonomous manner and extrinsic FGF8 may

act as a critical factor in controlling this process during odontogenesis.

Materials and methods

Mouse models

Wnt1-Cre and *R26R^Fgf8* mice were described previously [28,29]. Dental mesenchyme-specifically overexpressed *Fgf8* mice were generated by crossing *Wnt1-Cre* mice with *R26R^Fgf8* mice. Mutant embryos were identified by PCR genotyping. All wild-type mice were on the CD-1 background. Mice were housed under the following identical conditions: 40-70% relative humidity, 12 hour light/dark cycle, and 20-25°C ambient temperature, in individually ventilated cages, with corncob granules for bedding and nesting, in groups of up to six. The mice were given free access to tap water and standard rodent chow (1010011; Jiangsu Xietong Pharmaceutical Bio-engineering Co., Ltd, Nanjing, China).

Tissue recombination, organ culture, and subrenal culture

Fresh lower jaws isolated from human fetuses of 14th-16th week gestation following medical termination of pregnancy were provided by Maternal and Child Health Hospital of Fujian Province, China. The precise embryonic age of fetuses was defined by the measurement of the crown-rump length (CRL) [30]. Human and E13.5 mouse embryonic tooth germs were dissected out in ice-cold PBS and then treated with Dispase II at 37°C for 20 min. The dental mesenchyme and dental epithelium were separated with the aid of fine forceps and incubated on ice for further tissue recombination experiments. Three types of tissue recombination were conducted and are summarized in Table 1: E13.5 mouse dental epithelium with bell-stage human dental mesenchyme (14-16 weeks of gestation) (referred to as mde+hdm); E13.5 mouse dental mesenchyme with bell-stage human dental epithelium (hde+mdm); and E13.5 mouse dental mesenchyme with mouse

dental epithelium from the same stage as control (mde+mdm). All recombinants were

cultured in the Trowell-type organ culture dish overnight, followed by subrenal grafting

in adult nude mice, as described previously [30,31]. The use of animals and human tissues

in this study was approved by the Ethics Committee of Fujian Normal University

(permission #2011005).

SU5402 injection

Two first mandibular premolar germs dissected out from the lower jaw of a human

fetus of 16[th] week gestation were cultured under the kidney capsules of two nude mice,

one germ each. SU5402 (Calbiochem, California, USA) and DMSO (Merck, Darmstadt,

Germany) was intraperitoneally injected into each recipient mouse at a dose of

25mg/Kg/d in 200 μL PBS twice a day for one week. The human tooth grafts were

harvested for histological analysis after another week in subrenal culture.

Lentivirus infection and cell re-aggregation

For efficient infection of lentivirus [32], isolated E13.5 mouse dental mesenchyme

was digested with 0.25% trypsin at 37 °C for 5 min and dispersed into a single-cell

suspension with the aid of a micropipette. To make a dental mesenchyme pellet, about

5×10^5 dental mesenchymal cells were infected with 20 μL concentrated lentivirus in 200

μL culture medium in an Eppendorf tube. The tube was incubated at 37°C in a 5% CO_2

environment for 2 hours with its cap open, centrifuged at 3000 rpm for 3 min, and then

incubated again in the same environment for 2 hours. Cell pellets were removed from the

tubes, recombined with E13.5 dental epithelium, and cultured in the Trowell-type organ

culture dish overnight. Recombinants were subsequently subjected to subrenal culture in

nude mice.

Histology, *in situ* hybridization, immunostaining, and Western blot

For histological analysis, harvested samples were fixed in 4% paraformaldehyde

(PFA)/PBS, dehydrated, embedded in paraffin, and sectioned at 7 μm. Standard

Hematoxylin/Eosin (H&E) and Azan dichromic staining were utilized for histological

analysis [33]. Ossified tissues or grafts were demineralized with EDTA/PBS for a week

prior to dehydration. For *in situ* hybridization, dental tissues from staged human embryos

were dissected out in ice-cold PBS and fixed in 4% RNase-free PFA/PBS at 4°C overnight.

Human cDNAs, including *FGF1* (MHS6278-202758608), *FGF2* (EHS1001-208062206),

FGF15/19 (MHS6278-202756089), *FGF18* (MHS6278-202830446), were purchased

from Open Biosystems (Alabama, USA) and were subcloned to pBluescript-KS plus

vector for riboprobe transcription. Whole-mount *in situ* hybridization and section *in situ*

hybridization were performed following standard protocols as described previously [34,35].

Immunostaining was performed as described previously [36]. The following primary

antibodies were used including the products of Santa Cruz Biotechnology (Texas, USA):

Ameloblastin, MMP20, DSP, K-Ras, FGF8, P21; ABCam: FGF4, FGF9, Caspase3,

Ccnd1; Cell Signaling Tech (Massachusetts, USA): pSmad1/5/8, pP38, pErk1/2, pJNK,

pAKT1, pPDK1; Merck Millipore: β-catenin; and Spring Bio: Ki67. Alexa Fluor 488 or

594 IgG (H+L) antibody (Thermo Fisher, Massachusetts, USA) and horseradish

peroxidase-coupled IgG antibody (Santa Cruz) were used as the secondary antibody. For

immunoblotting, pP38, pErk1/2, pJNK, pAkt, and Actin (Santa Cruz) were used as

primary antibodies. IRDye 800cw or 680cw IgG (Odyssey, Nebraska, USA) were used

as secondary antibodies. Immunoblotting was conducted as described previously [37].

Signals were observed using microscope (Olympus BX51, Tokyo, Japan), Laser scanning

confocal microscope (Zeiss LSM780, Jena, Germany), and Infrared Imaging Systems

(Odyssey Clx). Representative images were taken from a minimum of 3 replicate

experiments.

Micro-computed tomography imaging and analysis

Harvested mineralized teeth were scanned with X-ray micro-computed tomography

(microCT) equipment (μCT50, SCANCO Medical AG, Bruettisellen, Switzerland). The

microCT parameters were as follows: 19×19 mm-field of view, 15-μm voxel size, 70

kVp, 200 mA, 360° rotation, 1500ms exposure time. All datasets were exported in the

DICOM format with the isotropic 15-μm^3 voxel. Acquired height and width of the tooth

crown were analyzed with the attached software of μCT50.

Statistics analysis

Image J software (version 1.46r; National Institutes of Health, USA) was utilized for

Ki67-positive cell counting and to detect the densitometry of the western blot bands with

at least three biological replicates counted. Statistical analyses were performed with

GraphPad Prism 5 (Version 5.01; GraphPad Software Inc, California, USA). Data were

represented as mean ± SD. Paired *t*-test was used to evaluate differences between two

groups, and analysis of variance (ANOVA) was used to determine differences among

different groups. If there was a significance of ANOVA test, Bonferroni post-tests was

used. *p* values <0.05 is considered statistically significant difference.

Results

Tooth developmental pace is dominated by the dental mesenchyme in humans and mice.

Our previous studies have shown that epithelial sheets of human keratinocyte stem cells, when recombined with E13.5 mouse dental mesenchyme, can be induced to differentiate into enamel-secreting ameloblasts in only 10 days and apoptotically degraded in 4 weeks, whereas the mouse secondary branchial arch epithelium, when recombined with the human dental mesenchyme from the bell-stage, remains in an enamel-secreting status after 15-week *ex vivo* culture [30,38,39]. These results strongly imply that the duration of epithelial differentiation in tooth development could be dominated by the mesenchymal component in humans and mice. To validate our hypothesis, we comprehensively revisited these phenotypes by carrying out meticulously designed heterogenous recombination between human and mouse embryonic dental tissues (Table 1). As shown in Fig. 1A-C, it took more than 4 weeks for the mouse dental epithelium to deposit enamel when recombined with the human dental mesenchyme in mde+hdm chimeric tooth germs and the chimera generated a mineralized tooth after 16-week growth (Suppl. Fig. 1A), exhibiting a much longer developmental duration and slower tempo in comparison with the mouse dental epithelia that were combined with their own dental mesenchyme (Fig. 1D-F, Suppl. Fig. 1B). By contrast, it took only about 10 days for the human dental epithelium to deposit enamel when recombined with E13.5 mouse dental mesenchyme in hde+mdm chimeric tooth germs (Fig. 1G-I) and the chimera could achieve a mineralized tooth in less than 30 days as well as the mouse homogenous

(mde+mdm) tooth (Suppl. Fig. 1C), demonstrating a dramatically accelerated

developmental pace compared to the normal human tooth morphogenesis.

Immunostaining of two ameloblastic differentiation markers, ameloblastin and MMP20

(Matrix Metallopeptidase 20), further confirmed the development phase alteration of the

dental epithelium induced by the heterogenous dental mesenchyme. The expression of

these two markers in the mouse dental epithelium-derived pre-ameloblasts in mde+hdm

chimeric recombinants was delayed to 4^{th} week in subrenal culture (Fig. 1J-L), whereas

the expression of the two markers was antedated at 9-day in the human dental epithelium-

derived ameloblasts in hde+mdm chimeric recombinants in subrenal culture (Fig. 1P-R),

which is similar to the expression of them at 7-day in the ameloblasts derived from the

mouse dental epithelium in mde+mdm control recombinants (Fig. 1M-O). The possibility

of tissue contamination was ruled out using specific primary antibodies against mouse or

human MHC I antigens to distinguish between human and mouse tissues in the

heterogenous chimeras (Suppl. Fig. 2). Our results unambiguously indicate that the dental

mesenchyme is a crucial component that dominates the tooth developmental pace by

interaction with the dental epithelium via diffusible paracrine signals during

odontogenesis.

FGF signaling participates in the regulation of tooth developmental pace.

We next attempted to identify the paracrine signaling molecules that may be

involved in the regulation of tooth developmental tempo in humans and mice. Conserved

SHH, FGF, BMP, and Wnt pathways are all crucial signaling engaged in cell proliferation,

apoptosis and differentiation during odontogenesis, indicating that they may be key

players in orchestrating the fundamental mainspring of organ growth. Since the dental

mesenchyme is dominant in the regulation of tooth growth (Fig. 1), we exclude

epithelially expressed SHH [40]. To test whether mesenchymal BMP, FGF and WNT

signaling may be involved in the regulation of tooth developmental pace, we examined

the level of the active indicators of these pathway including pSmad1/5/8 for BMP, K-Ras

for FGF, and β-catenin for Wnt, respectively, in the chimeric tooth germs. The expression

level of K-Ras in both the dental epithelium and mesenchyme is much higher in the

mde+hdm chimeric tooth germ (Fig. 2E), where the development of mouse dental

epithelium-derived ameloblasts were dramatically delayed and remained immature after

4 weeks in subrenal culture, compared to that in the mouse homogenous recombinant

tooth germ (Fig. 2B). The expression levels of pSmad1/5/8 and β-catenin in mouse dental

epithelium-derived ameloblasts and adjacently interacted odontoblasts did not display

obvious   difference   as   compared   between   the   heterogenous   recombinants   and

homogenous ones (Fig. 2A,C,D,F). The enhanced epithelial FGF activity in the mde+hdm

chimeric tooth germ, evidenced by outstandingly enhanced activity of K-Ras, was

certainly induced by the human dental mesenchyme and may attribute to the protracted

differentiation of the mouse dental epithelium.

To explore whether the upregulated FGF signaling may be associated with the

prolonged tooth development phase in humans, we started with a search for candidate

FGFs that are persistently expressed in the human developing dental mesenchyme but

barely in the mouse one. Most expression patterns of 22 known FGF ligands in mammals

have been verified in mouse tooth germs at the cap and bell stages (summarized in Suppl.

Table 1)[41,42]. We have examined in our previous studies the expression patterns of *FGF3,*

FGF4, FGF7, FGF8, FGF9, and *FGF10* in human tooth germs at different stages using

in situ hybridization or immunohistochemistry [43]. Here, we screened the expression of

additional 12 *FGFs* that have not been identified in human tooth germs by RT-PCR. We

excluded *FGF11, FGF12, FGF13,* and *FGF14*, because they are not secreting ligands

and do not activate FGF receptors as well. We found that, among them, the transcripts of

FGF1, *FGF2*, *FGF15/19*, and *FGF18* were detectable at both the cap and bell stages

(data not shown). We further confirmed their expression in developing human teeth by

section *in situ* hybridization (Suppl. Fig. 3). When comparing the FGF expression data

between the human [43] and mouse [41,42], we noticed that the majority of FGF transcripts

are broadly distributed in the whole tooth germ including the dental epithelium and

mesenchyme throughout the early cap stage to the late bell stage in humans [43]. However,

their expression is restricted to the limited region of tooth germs, such as enamel knot,

and appeared in limited periods of time in mice (Suppl. Table 1)[41,42]. We believe that this

widely and persistently distributing pattern of FGF ligands in the human tooth germs is

closely related to their characteristics of tooth development. To further verify this notion,

we cultured human 1st premolar germs at the early bell stage under the kidney capsule of

nude mice with intraperitoneal injection of SU5402, a global FGF-signaling inhibitor (an

inhibitor of tyrosine phosphorylation of FGF receptor). The inhibition activity of SU5402

in human tooth germs was firstly evaluated after two days injection and showed that it

significantly inhibited FGF-mediated activation of FGFR downstream signal transduction,

pErk1/2 and pPDK1 (Fig. 2G). Then, we cultured the human molar germs with SU5402

for one week and found accelerated detinogenesis in the SU5402-treated molar germ with

the appearance of both predentin and dentin (Fig. 2I), as compared to DMSO-treated one

with appearance of only predentin (Fig. 2H), indicating pivotal roles of FGF-signaling in

regulation of tooth developmental tempo.

With further analyses of FGF expression data in humans and mice, we found that

FGF4, FGF8, and FGF9 ligands exhibit most distinctive patterns (Suppl. Table 1). These

three ligands are persistently expressed in the dental epithelium and mesenchyme at least

until the differentiation stage in humans [43]. However, in mice, the expression of FGF4 is

restricted to the enamel knot from the cap to bell stage and that of FGF9 to the enamel

knot at the cap stage and to the dental epithelium at the bell stage. Furthermore, FGF8 is

only expressed in the presumptive dental epithelium prior to and during the laminar stage

and is no longer expressed after the E12 bud stage [44]. Whereas, in humans, *FGF8*

transcripts are widely and intensively present in the whole tooth germ at least from the

early cap stage to the differentiation stage [43]. Importantly, we found retained expression

of FGF8, as well as FGF4 and FGF9, in the mouse dental epithelium derived pre-

ameloblasts in mde+hdm recombinants after 4 weeks in subrenal culture, where the

development of mouse ameloblasts was largely delayed (Fig. 2J-L), whereas FGF8 was

attenuated in the human dental epithelium derived pre-ameloblasts in hde+mdm

recombinants after 10 days in subrenal culture, where the development of human

ameloblasts was vastly accelerated (insert in Fig. 2K). Based on these observations, we

hypothesized that FGF signaling may play important roles and FGF8 could be a critical

factor in the regulation of tooth developmental pace.

Forced activation of FGF8 signaling in the mouse dental mesenchyme delays tooth development.

To test our hypothesis, we set to overexpress *Fgf8* in dental mesenchymal cells in

the E13.5 recombinant molar germ via lentivirus-mediated expression and subrenal

culture system. Histological examination revealed that it took up to 9 days for *Fgf8*-

overexpressed tooth grafts to start dentin secreting (Fig. 3B), whereas it only took 7 days

for control ones to do so (Fig. 3A). Moreover, Ki67 staining detected remarkably higher

level of proliferation rate in both the dental epithelium and mesenchyme of *Fgf8*-

overexpressed tooth grafts compared to the controls after 7 days in subrenal culture (Fig.

3C-E; *n*=6/group). These data indicate that activated mesenchymal FGF8 activity delays

tooth development and dramatically stimulates cell proliferation.

To corroborate the aforementioned *ex vivo* results, we further generated *Wnt1-Cre;*

R26R^{Fgf8} transgenic mice to verify whether forced activation of mesenchymal FGF8

signaling alters tooth development *in vivo*. The presence of FGF8 in the dental

mesenchyme of *Wnt1-Cre;R26R^{Fgf8}* mice was confirmed by anti-FGF8 immunostaining

(Suppl. Fig. 4). Most of mutant embryos die at around E12.5 and a few died around E14.5.

Although histological examination revealed severe craniofacial hypoplasia in 100% cases,

including exencephaly and the absence of identifiable tissues such as ears, auricles, eyes,

nose, as well as tongue (data not shown), the mutant mice still possessed a pair of intact

maxillary and mandibular molar germs (Fig. 4). More importantly, both the maxillary and

mandibular molar germ exhibited an obvious retardation of tooth development (Fig. 4).

Since most mutant embryos died at around E12.5, we grafted E11.5 mutant molar germs

under the kidney capsule for further development. Examination of histology and the deposition of dentin and enamel showed that the mutant maxillary molar developed with approximate one-day's delay when compared to the wild-type ones in nearly 100% (15/16, Fig. 5A-F), and the mutant mandibular molar developed with about two-day's delay in 2 out of 9 grafts with the rest 7 failing to develop (insert in Fig. 5D). Examination of the expression of odontoblast marker DSP and ameloblast marker ameloblastin in the grafted maxillary molars further confirmed this delay (Fig. 5I-L).

Interestingly, Azan staining showed that the mutant graft of maxillary molar germ formed a larger-sized tooth (Fig. 5G) than the wild-type one (Fig. 5H) after 18 days in subrenal culture. Further Micro-CT scanning revealed that the mutant graft indeed formed a larger-sized tooth with short and dumpy shape (Fig 5N) compared to the control (Fig. 5M) after 3 months in subrenal culture for complete mineralization, which exhibited no difference in the average height of the tooth crowns but about 20% wider in the average width in the mutants ($n = 11$) compared to the controls ($n = 7$, Fig. 5O). Apparently, higher rate of cell proliferation contributes greatly to the larger teeth, which was evidenced by more Ki67 positive cells presented in E13.5 mutant tooth germ and the grafted mutant teeth than controls (Fig. 6A-F, K-M, $n = 3$/each group). Meanwhile, while apoptotic marker Caspase3 staining was present in ameloblasts and odontoblasts in both mutant (Fig. 6I,J) and wild-type tooth germs (Fig. 6G,H), this apoptotic signal was non-detectable in the pulp cells of mutant tooth germs (Fig. 6I,J), whereas it was seen in that of wild-type ones (Fig. 6G,H) after 18 days in subrenal culture. These results further support the idea that ectopically activated FGF8 signaling in the mouse dental

mesenchyme stimulates cell proliferation and prevents cell apoptosis during

odontogenesis.

Ectopically activated dental mesenchymal FGF8 signaling is transduced via p38 and

PI3K-Akt intracellular pathway.

To determine which intracellular pathway is implicated in the signal transduction of

the ectopically activated dental mesenchymal FGF8 signals, we examined the levels of

phosphorylated Erk1/2, P38, JNK, and PI3K-Akt, the activity indicator of MAPK

pathway, in both E13.5 tooth germs and E11.5 ones after 9 days in subrenal culture. As

shown in Fig. 7, no signal intensity differences in pErk1/2 (Fig. 7A-D) and pJNK (Fig.

7E-H) staining were observed between wild-type and mutant teeth. Whereas, p38

signaling is dramatically elevated in *Wnt1-Cre;R26R^{Fgf8}* teeth with a considerably

increased amount of nuclear-localized pP38 cells both in the dental epithelium and dental

mesenchyme of E13.5 mutant tooth germs (Fig. 7J) and grafted teeth (Fig. 7L) as

compared to the controls (Fig. 7I,K). Meanwhile, the intensity of PI3K-Akt staining was

slightly enhanced and the amount of pAkt1$^+$ cells was increased in mutant E13.5 tooth

germs and grafted teeth (Fig. 7N,P) as compared to controls (Fig7 M,O). Western blot

analysis further confirmed the increased expression of pP38 and pAkt1 in E13.5 mutant

tooth germs (Fig. 7Q,R; *n* = 3). Our results indicate that the augmented dental

mesenchymal FGF8 activity enhances P38 and PI3K-Akt signaling in both the dental

mesenchyme and the dental epithelium, suggesting that they may function as intracellular

pathways for transducing FGF8 signal into nuclei.

Enhanced dental mesenchymal FGF8 signaling promotes transition of cell cycle

from G1- to S-stage during odontogenesis.

The rate of cell proliferation controlled by cell cycle progression is a key player in organ size control [4] and is also closely associated with tissue differentiation status, as higher cell proliferation always presents in developing tissues and lower one in differentiated ones. Since the elevation of *Fgf8* expression in the dental mesenchyme results in the significant promotion of cell proliferation and prevention of cell apoptosis (Fig. 6), we speculated that enhanced dental mesenchymal FGF8 signaling would alter cell cycling. To verify this hypothesis, we set forth to ascertain the distribution of cyclin D1 (Ccnd1) and its inhibitor $p21^{Cip1}$ (p21), a pair of critical regulators functioning stimulatory and repression effect, respectively, on inducing cell cycle entry and mediating the G1-S stage transition [45,46], in E13.5 tooth germs and grafted teeth. As shown in Fig. 8, although the same high density of $Ccnd1^+$ cells were present in the dental epithelium of E13.5 wild-type and mutant tooth germs, the dental mesenchyme in mutant tooth germs possessed a considerably increased amount of $Ccnd1^+$ cells in comparison to that in wild-type ones (Fig. 8A,E). On the other hand, $p21^+$ cells were abundantly present in the presumptive primary enamel knot of the dental epithelium and slightly in the dental mesenchyme in wild-type tooth germs but totally absent from the whole tooth germ in mutant ones (Fig. 8B,F). Thereout, more colocalization of Ccnd1 and p21 staining were found in the wild-type tooth germs than in the mutant ones (Fig. 8C-D, G-H). In grafted teeth, nuclear $Ccnd1^+$ cells were distributed not only in ameloblasts and odontoblasts (Fig. 8M,P-upper) but also all over the dental pulp in mutants (Fig. 8M,P-lower), rather than

being restricted to the dental mesenchymal cells adjacent to the odontoblasts in wild-types

(Fig. 8I,L). Although p21 was expressed in ameloblasts and dental mesenchymal cells in

both wild-type and mutant teeth, its expression level was relatively decreased in mutants

(Fig. 8J,L,N,P). In addition, the nuclear localization of p21 could be detected in dental

mesenchymal cells of wild-type teeth (Fig. 8L-lower), whereas it was found to be in the

cytoplasm of grafted mutant teeth (Fig. 8P-lower), which make it unable to cooperate

with nuclear Ccnd1 to regulate the cell cycle progression in the dental mesenchymal cells.

Taken together, these results provide evidence that enhanced mesenchymal FGF8 activity

observably elevates the activity of nuclear Ccnd1 while degrades the activity of nuclear

p21. Thus both stimulation of Ccnd1 and repression of p21 function to induce cell cycle

entry and promote the transition of cell cycle from G1- to S- stage, which facilitates cell

population to retain the high rate of proliferation.

Discussion

Tooth developmental tempo complies with the hereditary nature of dental mesenchyme.

A heterogenous recombination study carried out between the mouse and rat molar tooth germs has suggested that tooth size is determined by the dental mesenchyme [47,48]. Additionally, our recent studies showed that, in comparison with natural developmental pace, the differentiation duration of the mouse dental epithelium or non-dental epithelium into ameloblasts when induced by the human dental mesenchyme is significantly extended, whereas that of the human dental epithelium or epithelial stem cells into ameloblasts when induced by the mouse dental mesenchyme is significantly shortened [38,39]. These results are consistent with the difference of development progression speed between human and mouse teeth [13], suggesting a dominant role of the dental mesenchyme in regulation of tooth developmental tempo. This idea is further supported by our current heterogenous recombination study carried out between the mouse and human molar tooth germs (Fig. 1). We further provide unambiguous evidence that not only the tooth size but also the tooth developmental pace are obedient to the hereditary character of dental mesenchyme.

Dental mesenchymal FGF8 and synergistic effects of FGFs play critical roles in controlling the developmental tempo during mammalian odontogenesis

Conserved pathways, such as FGF, BMP, Hh, and WNT pathway, participate in the highly conserved molecular mechanism underlying organismal growth by corresponding alterations of cell growth and division [4,11,12]. FGFs, as the most common mitogens, are

positively correlated to the rate of cell growth and cell division and involved in the

induction and pattern formation during tooth development [49,50]. Fgf8 has been

demonstrated to play an important role in cell survival and proliferation of CNC cells [51],

and to exert a positive effect on proliferation of dental mesenchymal cells [44]. Our data

show that FGF signaling rather than BMP and WNT signaling is highly active in human-

mouse chimeric tooth germs where the differentiation duration of the mouse dental

epithelium is largely slowed down (Fig. 2), which is consistent with the continuously high

expression of FGFs, particularly FGF8, in human tooth germs, suggesting a potentially

positive role of FGF signaling in regulating the tooth developmental tempo during

odontogenesis in humans. Both our *ex vivo* lentivirus-mediated overexpression system

and *in vivo* transgenic mouse model support that forcibly activated dental mesenchymal

FGF8 signaling stimulates cell proliferation and inhibits cell apoptosis, thus prolonging

tooth developmental pace in mice (Fig. 3-6). FGF8 appears to be a critical factor in

regulating the tooth developmental pace during mammalian odontogenesis.

Functional redundancy complicates interpretation of FGF genetic studies.

Phenotypes in various *Fgfr* knockout mice have demonstrated that different FGF

receptors have both essential and redundant roles throughout development [49]. During

odontogenesis, FGF7 and FGF10 have been proven to act as surrogates of mesenchymal

FGF3 [52], and FGF4, FGF8, as well as FGF9 have also been suggested functional

redundancy and repetitive use in mouse tooth development [13]. Our previous and current

study show that several FGFs, especially FGF4, FGF8, and FGF9, have a broad and

persistent expression in the whole tooth germs in humans but are restricted to the limited

region of tooth germs in mice (Suppl. Table 1) [43]. These data suggest that tooth

developmental pace may be the product of synergistic effects of different FGFs in

mammals. Actually, in addition to the fact that FGF8 is a key factor to control tooth

developmental pace discovered in the current study, Fgf9 has also been reported to induce

cell proliferation in cultured mouse dental mesenchyme *in vitro* [44] as well as in cervical

loop development *in vivo* [53] and the *Fgf9* null mutant mice show a size-reduced labial

cervical loop [53], and FGF4 stimulates cell proliferation in both the dental epithelium and

mesenchyme as well [44,54], both suggesting their positive function on the regulation of

tooth growth. We speculate that, in humans, several broadly, intensively, and persistently

expressed FGF ligands may act in concert to confer growth capability (stimulation of

proliferation and prevention of apoptosis) on tooth tissues to achieve longer development

time-scale.

FGF pathway may also participate in the regulation of tooth size.

On the other hand, the developmental tempo and size of individual organs must be

tightly coordinated to match the body size and larger-bodied species always keep slower

developmental tempos to acquire larger organ sizes. Hippo signaling is the only admitted

pathway that plays its best-known physiological function on the control of organ size in

the mouse model for mammalian lineage [5-8], which mainly involves in the regulation of

cell proliferation and apoptosis and thereby cell number [9]. Activation or inhibition of

YAP/TAZ, a transcriptional co-activator of Hippo pathway, leads to the adjustment of cell

proliferation and apoptosis, thus affecting organ sizes and tissue homeostasis [5-8].

However, in tooth development, suppression of YAP *ex vivo* at the cap stage only leads

to ectopic cusp formation [55], while overexpression of YAP *in vivo* disrupts tooth

morphogenesis with abnormal enamel knot as well as greatly widened dental lamina;

teeth ultimately formed in YAP transgenic mice do not exhibit any growth delay or size

enlargement [6]. These data suggest that the renowned control function of Hippo pathway

in mammalian organ size is not applicable for the tooth organ. In support of this notion,

it has also been reported that size-control mechanisms differ across mammalian organs

[11] and enhanced Hippo signaling induces organ enlargements in the stomach, heart, liver,

and spleen, but not in lung, kidney, and intestine [56-58]. Besides, IGF signaling is the other

pathway that has been implied to have effects on organ size and *Igf2r* disrupted mice are

born 25-35% bigger than wild-types, however, it has also been revealed that IGF signaling

affects the size in some organs such as the cecum and colon but not the size of other

organs such as the skin and stomach [10]. In addition, it has indicated that tuning Bmp

activity via enhanced epithelial Noggin activity modulates the size and number of the

teeth in mice [24]. Herein, we uncovered that natural broadly and persistently distributing

pattern of FGF ligands, especially FGF8, in the human tooth germs may be also closely

related to its tooth size.

Dental mesenchymal FGF8 signal promotes cell cycle progression from G1- to S-

stage via enhancing p38 and PI3K-Akt signaling to stimulate proliferation and

prevent differentiation.

Transduction of FGF signaling is mediated by multiple intracellular pathways,

including P38, PI3K/Akt, Erk1/2, and JNK [49]. In our previous study, the similar

expression patterns of the indicators in Erk, P38, JNK, and PI3K-Akt pathways in mouse

and human tooth germs suggests their conservative regulatory functions during mammalian odontogenesis [43]. P38 has been reported to be involved in control of the cell cycle via the expression changes of P21, P27 and Ccnd3 in mouse skeletal muscle cells [59], Ccnd1 and c-Myc in human dental pulp stem cells [60], and Ccnd1 and P21 in rat mesenchymal stem cells [61], which is also significantly increased in our *Wnt1-Cre; R26R^Fgf8* mice (Fig. 7). The PI3K-Akt pathway is associated with a variety of cellular functions including cell growth, cell cycle regulation, and cell survival [62]. The increased PI3K-Akt level regulates the proliferation and differentiation of neural progenitor cells by the activation of the cell cycle through the elevation of the Ccnd1 and cyclin E, two cyclins essential for the G1-S transition during organogenesis [63]. Moreover, Akt protein is a central intracellular factor to transduce and integrate multiple extracellular growth factor signals, such as FGF, BMP, and GH/IGF, into a comprehensive signal network to supervise the processes of cell proliferation, growth, and survival [11]. The increased activity of PI3K-Akt pathway caused by augmented FGF8 in the cranial neural crest cell (Fig. 7) further supports its positive function in cell proliferation regulated by cell cycle during tooth development. Erk has been reported to provide a switch-like transition between proliferation and differentiation of muscle progenitors [64], while JNK pathway has been elucidated to play crucial role in negative growth regulation and likely performs a central regulator in both local growth coordination and global growth coordination [4]. These two pathways did not show any noticeable alteration in our *Fgf8* transgenic mice, which may attribute to their main function in control of negative growth or balancing cell proliferation and differentiation.

Cell growth and division are controlled by the cyclin depend kinases (CDKs) and

their binding partners (cyclins) that regulate cell cycle progression through the regulation

of the G1-S and G2-M transition. Many cell cycle regulators have been proved to control

organ growth, which appears to be generally conserved in *Drosophila* and mammals [4].

The G1-S phase transition is a rate-limiting step, a known checkpoint between cell growth

and cell division, requiring cell cycle regulators to monitor and regulate the progression

of the cell cycle [65]. The stimulatory effects of cyclins can be offset by CDKs inhibitors

(CDKIs) and the repression of CDKIs is also a prerequisite for G1-S transition. Thus

merely elevating the level of cyclin D1 is not sufficient to induce cell cycle entry [66].

Cyclin D1 (Ccnd1) is synthesized rapidly in the G1 phase and accumulates in the nucleus

and degraded as the cell enters the S phase [45]. P21, known as CDKI1, functions as a

regulator of cell cycle progression in G1 and S phase and inhibits all cyclin/CDK

complexes, which in turn results in cell cycle arrest and/or apoptosis [46]. Enhanced dental

mesenchymal FGF8 may promotes cell cycle progression through the regulation of the

G1-S transition, which further stimulates cell proliferation and prevents cell

differentiation and apoptosis, thereby resulting in prolonged tooth development. In line

with this notion, we found that the amount of nuclear $Ccnd1^+$ cells is dramatically

increased while that of nuclear $P21^+$ cells is markedly decreased in the dental

mesenchyme at both the early and differentiation stage in *Wnt1-Cre;R26R^{Fgf8}* mice as

compared to those in the control (Fig. 8), stating that forcedly enhanced FGF8 in the

dental mesenchyme could elevate the level of Ccnd1 and meanwhile suppress the level

of p21. Stimulation of Ccnd1 and repression of p21 both constitute the prerequisite for

inducing cell cycle entry and thereby promoting the G1-S transition.

FGF8 controls developmental pace in a non-cell-autonomous manner during early

odontogenesis.

Two latest studies have demonstrated that interspecies differences in developmental

tempo are caused by cell-autonomous differences in biochemical reaction speeds instead

of differences in signaling, the sequence of genes, or regulatory elements in *in vitro*

human and mouse platforms [67,68]. Moreover, in *Drosophila*, it is also reported that the

Minute mutation causes a prolongation of developmental pace and decreased cell division

rate in a cell-autonomous way via decreasing the expression of ribosomal proteins and

thereby reducing the rate of translation and protein synthesis [69,70]. Developmental

allochrony regulated in a cell-autonomous manner is highlighted in these studies.

However, our findings that the dental mesenchyme resorts to extrinsic signals such as

FGF to dominate the developmental tempo as well via tissue interaction in a non-cell-

autonomous manner during tooth development are inconsistent with these studies.

Additionally, the fact that the oscillation phase shift between Notch and Wnt signaling is

critical for the segmentation of presomitic mesoderm cells in multicellular organ

development in 2D assays *ex vivo* [71] also indicates the pivotal role of secretory signal

changes in the regulation of developmental pace. In *Drosophila*, organ growth are

orchestrated not only by the growth rate regulated via signaling pathways but also by the

growth duration regulated via hormonal signals and organ-autonomous processes

controlled by patterning genes and physical force [4]. Our heterogenous recombination

results provide compelling evidence to support the contribution of extrinsic cues in the

developmental tempo of teeth relying on tissue interaction during early development. Thus, we suggest that complicated regulatory networks involved in the developmental pace work both in a cell-autonomous and non-cell-autonomous way to orchestrate the organ morphogenesis, and, particularly, tissue interaction in a non-cell-autonomous way plays a major role during early organogenesis.

Tissue bioengineering based on the principle of developmental biology represents a promising approach for repair and regeneration of damaged tissues and organs in the future. However, unmatched development phase between humans and the model animals such as *Drosophila*, zebra fishes, and mice appear to be a major obstacle to acquire knowledge about organ growth control in humans. Uncovering of the molecular mechanism underlying organ growth control in humans is a prerequisite for the realization of replacement therapy in the future. Our study provides important FGF signaling information in tooth developmental process and enlightens our knowledge about the nature of FGF signaling in human tooth growth.

Acknowledgments

This study was supported by the National Natural Science Foundation of China (81870739, 82001002, 81271102, 81771034) and the Natural Science Foundation of Fujian Province (2019J01281, 2020J01180).

References

1.   Lloyd AC. The regulation of cell size. Cell. 2013;154(6):1194-205.

2.   Promislow DEL, Harvey PH. Living fast and dying young: A comparative analysis of life‐history
variation among mammals. J Zool Lond. 1990;220(3):417-37.

3.   Swanson EM, Dantzer B. Insulin-like growth factor-1 is associated with life-history variation across
Mammalia. Proc Biol Sci. 2014;281(1782):20132458.

4.   Gokhale RH, Shingleton AW. Size control: the developmental physiology of body and organ size
regulation. Wiley Interdiscip Rev Dev Biol. 2015;4(4):335-56.

5.   Barry ER, Morikawa T, Butler BL, Shrestha K, de la Rosa R, Yan KS, et al. Restriction of intestinal
stem cell expansion and the regenerative response by YAP. Nature. 2013;493(7430):106-10.

6.   Liu M, Zhao S, Wang XP. YAP overexpression affects tooth morphogenesis and enamel knot
patterning. J Dent Res. 2014;93(5):469-74.

7.   Yu FX, Zhao B, Guan KL. Hippo Pathway in Organ Size Control, Tissue Homeostasis, and Cancer.
Cell. 2015;163(4):811-28.

8.   Zhang P, Pei C, Wang X, Xiang J, Sun BF, Cheng Y, et al. A Balance of Yki/Sd Activator and E2F1/Sd
Repressor Complexes Controls Cell Survival and Affects Organ Size. Dev Cell. 2017;43(5):603-17
e5.

9.   Tumaneng K, Russell RC, Guan KL. Organ size control by Hippo and TOR pathways. Curr Biol.
2012;22(9):R368-79.

10.   Zaina S, Squire S. The soluble type 2 insulin-like growth factor (IGF-II) receptor reduces organ size
by IGF-II-mediated and IGF-II-independent mechanisms. J Biol Chem. 1998;273(44):28610-6.

11.   Penzo-Mendez AI, Stanger BZ. Organ-Size Regulation in Mammals. Cold Spring Harb Perspect Biol.
2015;7(9):a019240.

12.   Lui JC, Garrison P, Baron J. Regulation of body growth. Curr Opin Pediatr. 2015;27(4):502-10.

13.   Zhang YD, Chen Z, Song YQ, Liu C, Chen YP. Making a tooth: growth factors, transcription factors,
and stem cells. Cell Res. 2005;15(5):301-16.

14.   Polly PD. Gene networks, occlusal clocks, and functional patches: new understanding of pattern
and process in the evolution of the dentition. Odontology. 2015;103(2):117-25.

15.   Jheon AH, Seidel K, Biehs B, Klein OD. From molecules to mastication: the development and
evolution of teeth. Wiley Interdiscip Rev Dev Biol. 2013;2(2):165-82.

16.   Salazar-Ciudad I. Tooth patterning and evolution. Curr Opin Genet Dev. 2012;22(6):585-92.

17.   Schoenwolf GC BS, Brauer PR, Francis-West PH. Larsen's Human Embryology E-Book. 2014.

18.   Rodrigues HG, Renaud S, Charles C, Le Poul Y, Sole F, Aguilar JP, et al. Roles of dental development
and adaptation in rodent evolution. Nat Commun. 2013;4:2504.

19.   Line SR. Variation of tooth number in mammalian dentition: connecting genetics, development,
and evolution. Evol Dev. 2003;5(3):295-304.

20.   Edelman GM. Morphoregulation. Dev Dyn. 1992;193(1):2-10.

21.   Thesleff I, Sharpe P. Signalling networks regulating dental development. Mech Dev.
1997;67(2):111-23.

22.   Jernvall J, Thesleff I. Reiterative signaling and patterning during mammalian tooth morphogenesis.
Mech Dev. 2000;92(1):19-29.

23.   Tummers M, Thesleff I. The importance of signal pathway modulation in all aspects of tooth
development. J Exp Zool B Mol Dev Evol. 2009;312B(4):309-19.

24.   Plikus MV, Zeichner-David M, Mayer JA, Reyna J, Bringas P, Thewissen JG, et al. Morphoregulation
of teeth: modulating the number, size, shape and differentiation by tuning Bmp activity. Evol Dev.
2005;7(5):440-57.

25.   Tapaltsyan V, Charles C, Hu J, Mindell D, Ahituv N, Wilson GM, et al. Identification of novel Fgf
enhancers and their role in dental evolution. Evol Dev. 2016;18(1):31-40.

26.   Thesleff I, Mikkola M. The role of growth factors in tooth development. Int Rev Cytol.
2002;217:93-135.

27.   Tucker A, Sharpe P. The cutting-edge of mammalian development; how the embryo makes teeth.
Nat Rev Genet. 2004;5(7):499-508.

28.   Danielian PS, Muccino D, Rowitch DH, Michael SK, McMahon AP. Modification of gene activity in
mouse embryos in utero by a tamoxifen-inducible form of Cre recombinase. Curr Biol.
1998;8(24):1323-6.

29.   Lin C, Yin Y, Bell SM, Veith GM, Chen H, Huh SH, et al. Delineating a conserved genetic cassette
promoting outgrowth of body appendages. Plos Genet. 2013;9(1):e1003231.

30.   Hu X, Lin C, Shen B, Ruan N, Guan Z, Chen Y, et al. Conserved odontogenic potential in embryonic
dental tissues. J Dent Res. 2014;93(5):490-5.

31.   Zhang Y, Wang S, Song Y, Han J, Chai Y, Chen Y. Timing of odontogenic neural crest cell migration
and tooth-forming capability in mice. Dev Dyn. 2003;226(4):713-8.

32.   Song Y, Zhang Z, Yu X, Yan M, Zhang X, Gu S, et al. Application of lentivirus-mediated RNAi in
studying gene function in mammalian tooth development. Dev Dyn. 2006;235(5):1334-44.

33.   Hu X, Xu S, Lin C, Zhang L, Chen Y, Zhang Y. Precise chronology of differentiation of developing
human primary dentition. Histochem Cell Biol. 2014;141(2):221-7.

34.   St Amand TR, Zhang Y, Semina EV, Zhao X, Hu Y, Nguyen L, et al. Antagonistic signals between
BMP4 and FGF8 define the expression of Pitx1 and Pitx2 in mouse tooth-forming anlage. Dev Biol.
2000;217(2):323-32.

35.   Dong X, Shen B, Ruan N, Guan Z, Zhang Y, Chen Y, et al. Expression patterns of genes critical for
BMP signaling pathway in developing human primary tooth germs. Histochem Cell Biol.
2014;142(6):657-65.

36.   Xiong W, He F, Morikawa Y, Yu X, Zhang Z, Lan Y, et al. Hand2 is required in the epithelium for
palatogenesis in mice. Dev Biol. 2009;330(1):131-41.

37.   Yang G, Yuan G, Ye W, Cho KW, Chen Y. An atypical canonical bone morphogenetic protein (BMP)
signaling pathway regulates Msh homeobox 1 (Msx1) expression during odontogenesis. J Biol
Chem. 2014;289(45):31492-502.

38.   Wang B, Li L, Du S, Liu C, Lin X, Chen Y, et al. Induction of human keratinocytes into enamel-
secreting ameloblasts. Dev Biol. 2010;344(2):795-9.

39.   Hu X, Lee JW, Zheng X, Zhang J, Lin X, Song Y, et al. Efficient induction of functional ameloblasts
from human keratinocyte stem cells. Stem Cell Res Ther. 2018;9(1):126.

40.   Hu X, Zhang S, Chen G, Lin C, Huang Z, Chen Y, et al. Expression of SHH signaling molecules in the
developing human primary dentition. BMC Dev Biol. 2013;13:11.

41.   Li CY, Prochazka J, Goodwin AF, Klein OD. Fibroblast growth factor signaling in mammalian tooth
development. Odontology. 2014;102(1):1-13.

42.   Cam Y, Neumann MR, Oliver L, Raulais D, Janet T, Ruch JV. Immunolocalization of acidic and basic
fibroblast growth factors during mouse odontogenesis. Int J Dev Biol. 1992;36(3):381-9.

43.   Huang F, Hu X, Fang C, Liu H, Lin C, Zhang Y, et al. Expression profile of critical genes involved in

FGF signaling pathway in the developing human primary dentition. Histochem Cell Biol.
2015;144(5):457-69.

44. Kettunen P, Thesleff I. Expression and function of FGFs-4, -8, and -9 suggest functional
655 redundancy and repetitive use as epithelial signals during tooth morphogenesis. Dev Dyn.
656 1998;211(3):256-68.

45. Baldin V, Lukas J, Marcote MJ, Pagano M, Draetta G. Cyclin D1 is a nuclear protein required for
658 cell cycle progression in G1. Genes Dev. 1993;7(5):812-21.

46. Gartel AL, Radhakrishnan SK. Lost in transcription: p21 repression, mechanisms, and
660 consequences. Cancer Res. 2005;65(10):3980-5.

47. Cai J, Cho SW, Kim JY, Lee MJ, Cha YG, Jung HS. Patterning the size and number of tooth and its
662 cusps. Dev Biol. 2007;304(2):499-507.

48. Jiang TX, Jung HS, Widelitz RB, Chuong CM. Self-organization of periodic patterns by dissociated
664 feather mesenchymal cells and the regulation of size, number and spacing of primordia.
665 Development. 1999;126(22):4997-5009.

49. Brewer JR, Mazot P, Soriano P. Genetic insights into the mechanisms of Fgf signaling. Genes Dev.
667 2016;30(7):751-71.

50. Lan Y, Jia S, Jiang R. Molecular patterning of the mammalian dentition. Semin Cell Dev Biol.
669 2014;25-26:61-70.

51. Abzhanov A, Tzahor E, Lassar AB, Tabin CJ. Dissimilar regulation of cell differentiation in
671 mesencephalic (cranial) and sacral (trunk) neural crest cells in vitro. Development.
672 2003;130(19):4567-79.

52. Kratochwil K, Galceran J, Tontsch S, Roth W, Grosschedl R. FGF4, a direct target of LEF1 and Wnt
674 signaling, can rescue the arrest of tooth organogenesis in Lef1(-/-) mice. Genes Dev.
675 2002;16(24):3173-85.

53. Kurosaka H, Islam MN, Kuremoto K, Hayano S, Nakamura M, Kawanabe N, et al. Core binding
677 factor beta functions in the maintenance of stem cells and orchestrates continuous proliferation
678 and differentiation in mouse incisors. Stem Cells. 2011;29(11):1792-803.

54. Jernvall J, Kettunen P, Karavanova I, Martin LB, Thesleff I. Evidence for the role of the enamel knot
680 as a control center in mammalian tooth cusp formation: non-dividing cells express growth
681 stimulating Fgf-4 gene. Int J Dev Biol. 1994;38(3):463-9.

55. Kwon HJ, Li L, Jung HS. Hippo pathway/Yap regulates primary enamel knot and dental cusp
683 patterning in tooth morphogenesis. Cell Tissue Res. 2015;362(2):447-51.

56. Song H, Mak KK, Topol L, Yun K, Hu J, Garrett L, et al. Mammalian Mst1 and Mst2 kinases play
685 essential roles in organ size control and tumor suppression. Proc Natl Acad Sci U S A.
686 2010;107(4):1431-6.

57. Heallen T, Zhang M, Wang J, Bonilla-Claudio M, Klysik E, Johnson RL, et al. Hippo pathway inhibits
688 Wnt signaling to restrain cardiomyocyte proliferation and heart size. Science.
689 2011;332(6028):458-61.

58. Dong J, Feldmann G, Huang J, Wu S, Zhang N, Comerford SA, et al. Elucidation of a universal size-
691 control mechanism in Drosophila and mammals. Cell. 2007;130(6):1120-33.

59. Irazoqui AP, Boland RL, Buitrago CG. Actions of 1,25(OH)2-vitamin D3 on the cellular cycle depend
693 on VDR and p38 MAPK in skeletal muscle cells. J Mol Endocrinol. 2014;53(3):331-43.

60. Vandomme J, Touil Y, Ostyn P, Olejnik C, Flamenco P, El Machhour R, et al. Insulin-like growth
factor 1 receptor and p38 mitogen-activated protein kinase signals inversely regulate signal

transducer and activator of transcription 3 activity to control human dental pulp stem cell
quiescence, propagation, and differentiation. Stem Cells Dev. 2014;23(8):839-51.

61. Stockl S, Bauer RJ, Bosserhoff AK, Gottl C, Grifka J, Grassel S. Sox9 modulates cell survival and
adipogenic differentiation of multipotent adult rat mesenchymal stem cells. J Cell Sci.
2013;126(Pt 13):2890-902.

62. Castellano E, Downward J. RAS Interaction with PI3K: More Than Just Another Effector Pathway.
Genes Cancer. 2011;2(3):261-74.

63. Xue XJ, Yuan XB. Nestin is essential for mitogen-stimulated proliferation of neural progenitor cells.
Mol Cell Neurosci. 2010;45(1):26-36.

64. Michailovici I, Harrington HA, Azogui HH, Yahalom-Ronen Y, Plotnikov A, Ching S, et al. Nuclear to
cytoplasmic shuttling of ERK promotes differentiation of muscle stem/progenitor cells.
Development. 2014;141(13):2611-20.

65. Elledge SJ. Cell cycle checkpoints: preventing an identity crisis. Science. 1996;274(5293):1664-72.

66. Chambard JC, Lefloch R, Pouyssegur J, Lenormand P. ERK implication in cell cycle regulation.
Biochim Biophys Acta. 2007;1773(8):1299-310.

67. Rayon T, Stamataki D, Perez-Carrasco R, Garcia-Perez L, Barrington C, Melchionda M, et al.
Species-specific pace of development is associated with differences in protein stability. Science.
2020;369(6510).

68. Matsuda M, Hayashi H, Garcia-Ojalvo J, Yoshioka-Kobayashi K, Kageyama R, Yamanaka Y, et al.
Species-specific segmentation clock periods are due to differential biochemical reaction speeds.
Science. 2020;369(6510):1450-5.

69. Saebøe-Larssen S, Lyamouri M, Merriam J, Oksvold MP, Lambertsson A. Ribosomal protein
insufficiency and the minute syndrome in Drosophila: a dose-response relationship. Genetics.
1998;148(3):1215-24.

70. Morata G, Ripoll P. Minutes: mutants of drosophila autonomously affecting cell division rate. Dev
Biol. 1975;42(2):211-21.

71. Sonnen KF, Lauschke VM, Uraji J, Falk HJ, Petersen Y, Funk MC, et al. Modulation of Phase Shift
between Wnt and Notch Signaling Oscillations Controls Mesoderm Segmentation. Cell.
2018;172(5):1079-90 e12.

Tables

Table. 1 Ratio of tooth formation in different types of tissue recombinants.

Type of Recombinant		No. of	No. of Tooth	Ratio (%)
Epithelium	Mesenchyme	Recombinant	Formation	
mde	mdm	28	28	100
mde	hdm	78	76	97.4
hde	mdm	27	24	88.9

727 Abbreviations: mdm, E13.5 mouse dental mesenchyme; hdm, bell-stage human dental mesenchyme

728 (14-16 weeks of gestation); mde, E13.5 mouse dental epithelium; hde, bell-stage human dental

729 epithelium (14-16 weeks of gestation).

Figure legends

Fig. 1. Heterogenous recombination of human and mouse embryonic dental tissues reveals the dominant role of the dental mesenchyme in the control of tooth developmental pace. A-C: Azan dichromic staining (stains dentin blue and enamel red) of recombinants of E13.5 mouse dental epithelium and bell-stage human dental mesenchyme cultured under the kidney capsule for 3 weeks (**A**), 4 weeks (**B**), 8 weeks (**C**). Note that no deposition of dentin and enamel was seen in the chimeric tooth germs until 4 weeks in subrenal culture. **D-F**: Azan dichromic staining of recombinants of E13.5 mouse dental epithelium and E13.5 mouse dental mesenchyme cultured under the kidney capsule for 6 days (**D**), 7 days (**E**), and 8 days (**F**). Note that the deposition of dentin and enamel was not seen until 7 and 8 days, respectively, in subrenal culture. **G-I**: Azan dichromic staining of recombinants of bell-stage human dental epithelium and E13.5 mouse dental mesenchyme cultured under the kidney capsule for 7 days (**G**), 9 days (**H**), and 10 days (**I**). Note that the deposition of dentin and enamel was not seen until 9 and 10 days, respectively, in subrenal culture. **J-R**: The expression of ameloblastin and MMP20 in recombinants of E13.5 mouse dental epithelium and bell-stage human dental mesenchyme cultured under the kidney capsule for 3 weeks (**J**), 4 weeks (**K** and **L**), in recombinants of E13.5 mouse dental epithelium and mouse dental mesenchyme for 6 days (**M**), 7 days (**N** and **O**), and in recombinants of bell-stage human dental epithelium and mouse dental mesenchyme for 7 days (**P**) and 9 days (**Q** and **R**). hdm, human dental mesenchyme; hde, human dental epithelium; mdm, mouse dental mesenchyme; mde, mouse dental epithelium; hod, human odontoblasts; mod, mouse odontoblasts; mam,

mouse ameloblasts; d, dentin; e, enamel. Scale bar = 50 μm.

Fig. 2. FGF signaling is involved in the regulation of tooth developmental pace. A-C:

Immunostaining showing the expression of pSmad1/5/8 (**A**), K-Ras (**B**), and β-catenin

(**C**) in recombinants of mouse dental epithelium and mouse dental mesenchyme cultured

under the kidney capsule for 7 days. **D-F**: Immunostaining showing the expression of

pSmad1/5/8 (**D**), K-Ras (**E**), and β-catenin (**F**) in recombinants of mouse dental

epithelium and human dental mesenchyme cultured under the kidney capsule for 4 weeks.

G: Human 1st premolar germs of 16th week gestation were transplanted under the kidney

capsule and cultured in the absence or presence of SU5402 for 2 days. Note the decreased

expression of pErk1/2 and pPDK1 in the presence of SU5402. **H-I**: Human 1st premolar

germs of 16th week gestation were transplanted under the kidney capsule and cultured in

the absence (**H**) or presence (**I**) of SU5402 for 1 week. Note the accelerated detinogenesis

with the appearance of dentin (**I**, red arrow) in the presence of SU5402. **J-L**:

Immunohistochemical staining showing the expression of FGF4 (**J**), FGF8 (**K**), and

FGF9 (**L**) in recombinants of mouse dental epithelium and human dental mesenchyme

cultured under the kidney capsule for 4 weeks and the expression of FGF8 (insert in **K**)

in recombinants of human dental epithelium and mouse dental mesenchyme cultured

under the kidney capsule for 10 days. mam, mouse ameloblasts; mod, mouse odontoblasts;

hod, human odontoblasts; am, ameloblasts; od, odontoblasts; iee, inner enamel epithelium;

sr, stellate reticulum; dm, dental mesenchyme; d, dentin; pre-d, predentin. Scale bar = 25

μm.

Fig. 3. Augmented FGF8 signaling in the mouse dental mesenchyme delays tooth development and increases cell proliferation. A, **B**: Azan dichromic staining of recombinants of E13.5 mouse dental epithelium and re-aggregate of dental mesenchymal cells infected with control LV-*eGFP* (**A**) or LV-*Fgf8-eGFP* (**B**) cultured under the kidney capsule for 6 days (insert in **A**), 7 days, and 9 days (insert in **B**). **C**, **D**: Immunostaining showing the expression of Ki67 in recombinants of E13.5 mouse dental epithelium and re-aggregate of dental mesenchymal cells infected with control LV-*eGFP* (**C**) or LV-*Fgf8-eGFP* (**D**) cultured under the kidney capsule for 7 days. **E**: Comparison of Ki67-positive cells in the dental mesenchyme and dental epithelium of the recombinants. Standard deviation values were shown as error bars, and ** indicates $p < 0.01$, $n = 6$/group. **F**: GFP staining of recombinants of E13.5 mouse dental epithelium and re-aggregate of dental mesenchymal cells infected with control LV-*eGFP* cultured under the kidney capsule for 11 days. dm, dental mesenchyme; dp, dental pulp; am, ameloblasts; sr, stellate reticulum; od, odontoblasts; d, dentin; Epi, epithelium; Mes, mesenchyme. Scale bar = 25 μm.

Fig. 4. Delayed tooth development in early molar germs of *Wnt1-Cre;R26R^{Fgf8}* mice. A-H: H&E staining showing the morphology of maxillary and mandibular molar in E13.5 wild-type (**A,C**) and mutant (**B,D**) embryos and in E14.5 wild-type (**E,G**) and mutant (**F,H**) embryos. de, dental epithelium; dm, dental mesenchyme. Scale bar = 50 μm.

Fig. 5. Grafts of *Wnt1-Cre;R26R^{Fgf8}* molar germs exhibit delayed tooth development

and enlarged tooth size. A-H: Azan dichromic staining of E11.5 wild-type and mutant

maxillary molar germs cultured under the kidney capsule for 8 days (**A,B**), 9 days (**C,D**),

10 days (**E,F**), 12 days (insert in **F**), and 18 days (**G,H**). Azan dichromic staining of E11.5

mutant mandibular molar germs cultured under the kidney capsule for 9 days (insert in

D). **I-L**: Immunostaining showing the expression of DSP (**I,J**) and Ameloblastin (**K,L**) in

E11.5 wild-type and mutant molar grafts cultured under the kidney capsule for 9 days. **M**,

N: Representative Micro-CT images of E11.5 wild-type (**M**) and mutant molar germs (**N**)

cultured under the kidney capsule for 3 months. **O**: Comparison of the height and width

of the tooth crown between wild-type and mutant mineralized teeth cultured under the

kidney capsule for 3 months. Standard deviation values were shown as error bars. ns

indicates $p > 0.05$, and ** indicates $p < 0.01$, $n = 7$ and 11/group in **O**. de, dental

epithelium; dm, dental mesenchyme; am, ameloblasts; od, odontoblasts; d, dentin; e,

enamel. Scale bar: A-H, I-L = 50 μm; M and N = 500 μm.

Fig. 6. Ectopically activated dental mesenchymal FGF8 signaling stimulates cell

proliferation and prevents cell apoptosis during odontogenesis. A-F: Immunostaining

showing the Ki67 positive cells in E13.5 wild-type (**A**) and mutant (**D**) molar germs and

in the grafts of E11.5 wild-type and mutant molar germ cultured under the kidney

capsule for 9 days (**B,E**) and 18 days (**C,F**). **G-J**: Immunostaining showing the expression

of Caspase3 in the grafts of E11.5 wild-type (**G**, boxed area is magnified in **H**) and mutant

(**I**, boxed area is magnified in **J**) molar germ cultured under the kidney capsule for 18

days. Note the absence of Caspase3 expression in the dental mesenchyme (red asterisk)

excepting odontoblasts in mutant grafts. **K-M**: The percentage of Ki67 positive cells in

the dental mesenchyme of wild-type and mutant E13.5 tooth germs (**K**) and tooth grafts

cultured under the kidney capsule for 9 days (**I**) and 18 days (**M**), $n = 3$/group, * indicates

$p < 0.05$. Standard deviation values were shown as error bars. am, ameloblasts; od,

odontoblasts; d, dentin; e, enamel. Scale bar = 100 μm.

Fig. 7. Ectopically activated dental mesenchymal FGF8 signaling is transduced via

p38 and PI3K-Akt intracellular pathways. A-D: Immunostaining showing the

expression of pErk1/2 in E13.5 wild-type (**A**) and mutant (**B**) molar germs and in E11.5

wild-type (**C**) and mutant (**D**) molar germs cultured under the kidney capsule for 9 days.

E-H: Immunostaining showing the expression of pJNK in E13.5 wild-types (**E**) and

mutants (**F**) molar germs and in E11.5 wild-type (**G**) and mutant (**H**) molar germs cultured

under the kidney capsule for 9 days. **I-L**: Immunostaining showing the expression of pP38

in E13.5 wild-type (**I**) and mutant (**J**) molar germs and in E11.5 wild-type (**K**) and mutant

(**L**) cultured under the kidney capsule for 9 days. **M-P**: Immunostaining showing the

expression of pAkt1 in E13.5 wild-types (**M**) and mutants (**N**) molar germs and in E11.5

wild-type (**O**) and mutant (**P**) molar germs cultured under the kidney capsule for 9 days.

Q: Western-blotting of pErk1/2, pP38, pJNK, and pAKT1 expressed in E13.5 wild-type

and mutant molar germs. **R**: Quantification of immunoblotting data in **Q**. $n = 3$/group. *

indicates $p < 0.05$. Standard deviation values were shown as error bars. de, dental

epithelium; dm, dental mesenchyme; am, ameloblasts; od, odontoblasts. Scale bar = 25

μm.

Fig. 8. Enhanced dental mesenchymal FGF8 signaling increases the level of nuclear Ccnd1 and decreases the level of nuclear p21. Immunostaining showing the colocalization of Ccnd1 and P21 in E13.5 wild-type (**A-D**) and mutant (**E-H**) molar germs and E11.5 wild-type (**I-L**) and mutant (**M-P**) molar germs cultured under the kidney capsule for 9 days. Boxed areas in (**C**), (**G**), (**K**), and (**O**) are magnified in (**D**), (**H**), (**L**), and (**P**), respectively. Note the significantly increased proportion of Ccnd1-positive cells (white arrow) and the shift of P21 from the nuclei to the cytoplasm (yellow arrow) in the mutant dental mesenchyme (**P**) as compared with wild-type ones (**L**). de, dental epithelium; dm, dental mesenchyme; am, ameloblasts; od, odontoblasts. Scale bar: A-N, P-S = 50 μm; O, T = 25 μm.