FGF8-mediated signaling regulates tooth developmental pace and size during odontogenesis

Chensheng Lin[1], Ningsheng Ruan[1], Linjun Li[1], Yibin Chen[1], Xiaoxiao Hu[1], YiPing Chen[2], Xuefeng Hu[1]*, Yanding Zhang[1]*

[1]Fujian Key Laboratory of Developmental and Neural Biology & Southern Center for Biomedical Research, College of Life Sciences, Fujian Normal University, Fuzhou, Fujian, 350117, P.R. China.

[2]Department of Cell and Molecular Biology, Tulane University, New Orleans, LA 70118, USA.

Running title: FGF8 regulates tooth developmental pace and size

*Correspondence to:

Yanding Zhang, Ph.D.

College of Life Sciences, Fujian Normal University (Qishan Campus), Fuzhou, Fujian, 350117, P.R. China.

Phone: +86-13706963408 (tel); E-mail: ydzhang@fjnu.edu.cn

or

Xuefeng Hu, Ph.D.

College of Life Sciences, Fujian Normal University (Qishan Campus), Fuzhou, Fujian, 350117, P.R. China.

Phone: +86-13055734898 (tel) ; E-mail: bioxfh@fjnu.edu.cn

Disclosure page

Conflicts of interest

The authors declare no conflicts of interest.

Data Availability Statement

All the data are available within the article or its supplementary materials.

Abstract

The developing human and mouse teeth constitute an ideal model system to study the regulatory mechanism underlying organ growth control due to the fact that their teeth share highly conserved and well-characterized developmental processes and their developmental tempo and size vary notably. In the current study, we manipulated heterogenous recombination between human and mouse dental tissues and demonstrate that the dental mesenchyme dominates the tooth developmental tempo and size and FGF8 could be a critical player during this developmental process. Forced activation of FGF8 signaling in the dental mesenchyme of mice promoted cell proliferation, prevented cell apoptosis via p38 and perhaps PI3K-Akt intracellular signaling, and impelled the transition of the cell cycle from G1- to S-phase in the tooth germ, resulting in the slowdown of the tooth developmental pace and the enlargement of the tooth size. Our results provide compelling evidence that extrinsic signals can profoundly affect tooth developmental tempo and size and the dental mesenchymal FGF8 could be a pivotal factor in controlling developmental pace and size in a non-cell-autonomous manner during mammalian odontogenesis.

Key words: FGF8, tooth, developmental pace, size, non-cell-autonomous.

Introduction

Although mammals possess an almost identical body plan with morphologically and functionally similar organs and systems, their developmental tempo and body sizes vary greatly among different species. Generally, larger-bodied species own longer gestation lengths with slower developmental tempos to acquire larger body sizes (Promislow and Harvey, 1990, Swanson and Dantzer, 2014). Obviously, developmental tempo and organ size must be tightly harmonized to ensure the correct establishment of body plan and therefore both of them are the result of a precisely controlled process. The mechanisms that regulate organ growth (i.e. embryonic development and postnatal growth) and size have been well investigated in non-mammalian models, particularly in *Drosophila melanogaster*, in which roles of TOR (target of rapamycin), insulin/IGF (insulin-like growth factor), Ras/Raf/MAPK, JNK (c-Jun N-terminal kinase), and Hippo signaling pathways are well defined (Gokhale and Shingleton, 2015).

In mammals, several conserved families of secreted growth factors, including FGF (fibroblast growth factor), BMP (bone morphogenetic protein), TGF-β (transforming growth factor-β), GH (growth hormone)/IGF, and Hippo/YAP, are deemed to be involved in the regulation of organ growth and organ size due to their ability to modulate cell proliferation and apoptosis during embryonic development (Penzo-Mendez and Stanger, 2015, Lui et al., 2015, Gokhale and Shingleton, 2015). However, no direct evidence can explain roles of pathways abovementioned in controlling developmental pace and organ size in mammals, except for the Hippo and IGF pathway in size control (Barry et al., 2013, Liu et al., 2014, Yu et al., 2015, Zhang et al., 2017, Tumaneng et al., 2012, Zaina and Squire, 1998).

Mammalian teeth have long served as a model organ for studying fundamental questions in organ development on the molecular basis (Zhang et al., 2005). The main features of tooth morphogenesis are conserved in the mammalian lineage (Polly, 2015, Jheon et al., 2013, Salazar-Ciudad, 2012), as they are all formed by sequential and reciprocal interactions between epithelium and mesenchyme and share similar processes during tooth development (Jheon et al., 2013). Despite these considerable homologies,

mammalian dentition undergoes remarkable morphological diversification, displaying great variations not only in tooth shape and number but also tooth size and growth rate (Schoenwolf GC, 2014, Rodrigues et al., 2013, Jheon et al., 2013, Zhang et al., 2005, Line, 2003). Obviously, the human dentition appears much larger size than the mouse one. In addition, the period of tooth development in humans takes about 400 days from the initiation stage at embryonic 6^{th} week to tooth eruption at postnatal 6^{th} month. Whereas, this period in mice takes only about 20 days from the determination of tooth forming sites at E10.5 to tooth eruption at postnatal 10^{th} day (Zhang et al., 2005). Thus, the human and mouse embryonic tooth provides an ideal model system for investigating the regulatory mechanism underlying the developmental tempo and size.

It is generally believed that diversified forms of organ morphogenesis (including growth rate and size) among species are achieved by tinkering conserved signaling pathways instead of proposing novel ones during development (Edelman, 1992, Thesleff and Sharpe, 1997, Jernvall and Thesleff, 2000). The difference in organismal growth rate and organ or body size between humans and mice primarily reflect the difference in cell number rather than the cell size (Lloyd, 2013). Many lines of evidence from mouse models demonstrate that numerous growth factors of conserved families, such as FGF, BMP, SHH, and Wnt, have been shown to play critical roles in tooth morphogenesis by regulating cell proliferation, differentiation, and apoptosis during odontogenesis (Tummers and Thesleff, 2009, Plikus et al., 2005, Tapaltsyan et al., 2016, Jheon et al., 2013, Thesleff and Sharpe, 1997, Thesleff and Mikkola, 2002, Tucker and Sharpe, 2004, Salazar-Ciudad, 2012). Studies in mice demonstrated that the Hippo pathway, a signaling pathway that is well-known to control organ size in mammals, does not involve in the regulation of tooth developmental pace and size during odontogenesis (Liu et al., 2014, Kwon et al., 2015) , but tuning down the BMP signaling by overexpressing secreted *Noggin* in the dental epithelium could reduce the tooth size. Here, we took advantage of the differences in developmental tempo and size of teeth in mice and humans, and provide compelling evidence that it is the dental mesenchyme that dominates tooth developmental tempo and size in a non-cell-autonomous manner and extrinsic FGF8 may act as a critical factor in controlling these processes during odontogenesis.

Materials and methods

Mouse models

Wnt1-Cre and *R26R^{Fgf8}* mice were described previously [Danielian et al., 1998, Lin et al., 2013]. Dental mesenchyme-specifically overexpressed *Fgf8* mice were generated by crossing *Wnt1-Cre* mice with *R26R^{Fgf8}* mice. Mutant embryos were identified by PCR genotyping. All wild-type mice were on the CD-1 background. Mice were housed under the following identical conditions: 40-70% relative humidity, 12 hour light/dark cycle, and 20-25°C ambient temperature, in individually ventilated cages, with corncob granules for bedding and nesting, in groups of up to six. The mice were given free access to tap water and standard rodent chow (1010011; Jiangsu Xietong Pharmaceutical Bio-engineering Co., Ltd, Nanjing, China).

Tissue recombination, organ culture, and subrenal culture

Fresh lower jaws isolated from human fetuses of 14th-16th week gestation following medical termination of pregnancy were provided by Maternal and Child Health Hospital of Fujian Province, China. The precise embryonic age of fetuses was defined by the measurement of the crown-rump length (CRL) (Hu et al., 2014a). Human and E13.5 mouse embryonic tooth germs were dissected out in ice-cold PBS and then treated with Dispase II at 37℃ for 20 min. The dental mesenchyme and dental epithelium were separated with the aid of fine forceps and incubated on ice for further tissue recombination experiments. Three types of tissue recombination were conducted and are summarized in Table 1: E13.5 mouse dental epithelium with bell-stage human dental mesenchyme (14-16 weeks of gestation) (referred to as mde+hdm); E13.5 mouse dental mesenchyme with bell-stage human dental epithelium (hde+mdm); and E13.5 mouse dental mesenchyme with mouse dental epithelium from the same stage as control (mde+mdm). All recombinants were cultured in the Trowell-type organ culture dish overnight, followed by subrenal grafting in adult nude mice, as described previously (Zhang et al., 2003, Hu et al., 2014a). The use of animals and human tissues in this study was approved by the Ethics Committee of Fujian Normal University (permission #2011005).

SU5402 injection

Two first mandibular premolar germs dissected out from the lower jaw of a human

fetus of 16th week gestation were cultured under the kidney capsules of two nude mice,

one germ each. SU5402 (Calbiochem, California, USA) and DMSO (Merck, Darmstadt,

Germany) was intraperitoneally injected into each recipient mouse at a dose of

25mg/Kg/d in 200 μL PBS twice a day for one week. The human tooth grafts were

harvested for histological analysis after another week in subrenal culture.

Lentivirus infection and cell re-aggregation

For efficient infection of lentivirus (Song et al., 2006), isolated E13.5 mouse dental

mesenchyme was digested with 0.25% trypsin at 37 °C for 5 min and dispersed into a

single-cell suspension with the aid of a micropipette. To make a dental mesenchyme pellet,

about 5×10^5 dental mesenchymal cells were infected with 20 μL concentrated lentivirus

in 200 μL culture medium in an Eppendorf tube. The tube was incubated at 37℃ in a 5%

CO_2 environment for 2 hours with its cap open, centrifuged at 3000 rpm for 3 min, and

then incubated again in the same environment for 2 hours. Cell pellets were removed from

the tubes, recombined with E13.5 dental epithelium, and cultured in the Trowell-type

organ culture dish overnight. Recombinants were subsequently subjected to subrenal

culture in nude mice.

Histology, *in situ* hybridization, immunostaining, and Western blot

For histological analysis, harvested samples were fixed in 4% paraformaldehyde

(PFA)/PBS, dehydrated, embedded in paraffin, and sectioned at 7 μm. Standard

Hematoxylin/Eosin (H&E) and Azan dichromic staining were utilized for histological

analysis (Hu et al., 2014b). Ossified tissues or grafts were demineralized with EDTA/PBS

for a week prior to dehydration. For *in situ* hybridization, dental tissues from staged

human embryos were dissected out in ice-cold PBS and fixed in 4% RNase-free PFA/PBS

at 4°C overnight. Human cDNAs, including *FGF1* (MHS6278-202758608), *FGF2*

(EHS1001-208062206), *FGF15/19* (MHS6278-202756089), *FGF18* (MHS6278-

202830446), were purchased from Open Biosystems (Alabama, USA) and were

subcloned to pBluescript-KS plus vector for riboprobe transcription. Whole-mount *in situ*

hybridization and section *in situ* hybridization were performed following standard

protocols as described previously (St Amand et al., 2000, Dong et al., 2014).

Immunostaining was performed as described previously (Xiong et al., 2009). The following primary antibodies were used including the products of Santa Cruz Biotechnology (Texas, USA): Ameloblastin, MMP20, DSP, K-Ras, FGF8, P21; ABCam: FGF4, FGF9, Caspase3, Ccnd1; Cell Signaling Tech (Massachusetts, USA): pSmad1/5/8, pP38, pErk1/2, pJNK, pAKT1; Merck Millipore: β-catenin; and Spring Bio: Ki67. Alexa Fluor 488 or 594 IgG (H+L) antibody (Thermo Fisher, Massachusetts, USA) and horseradish peroxidase-coupled IgG antibody (Santa Cruz) were used as the secondary antibody. For immunoblotting, pP38, pErk1/2, pJNK, pAkt, and Actin (Santa Cruz) were used as primary antibodies. IRDye 800cw or 680cw IgG (Odyssey, Nebraska, USA) were used as secondary antibodies. Immunoblotting was conducted as described previously (Yang et al., 2014). Signals were observed using microscope (Olympus BX51, Tokyo, Japan), Laser scanning confocal microscope (Zeiss LSM780, Jena, Germany), and Infrared Imaging Systems (Odyssey Clx). Representative images were taken from a minimum of 3 replicate experiments.

Micro-computed tomography imaging and analysis

Harvested mineralized teeth were scanned with X-ray micro-computed tomography (microCT) equipment (μCT50, SCANCO Medical AG, Bruettisellen, Switzerland). The microCT parameters were as follows: 19×19 mm-field of view, 15-μm voxel size, 70 kVp, 200 mA, 360° rotation, 1500ms exposure time. All datasets were exported in the DICOM format with the isotropic 15-μm^3 voxel. Acquired height and width of the tooth crown were analyzed with the attached software of μCT50.

Statistics analysis

Image J software (version 1.46r; National Institutes of Health, USA) was utilized for Ki67-positive cell counting and to detect the densitometry of the western blot bands with at least three biological replicates counted. Statistical analyses were performed with GraphPad Prism 5 (Version 5.01; GraphPad Software Inc, California, USA). Data were represented as mean ± SD. Paired *t*-test was used to evaluate differences between two groups, and analysis of variance (ANOVA) was used to determine differences among different groups. If there was a significance of ANOVA test, Bonferroni post-tests was used. *p* values <0.05 is considered statistically significant difference.

Results

Tooth developmental tempo and size is dominated by the dental mesenchyme in humans and mice.

Our previous studies have showed that epithelial sheets of human keratinocyte stem cells, when recombined with E13.5 mouse dental mesenchyme, can be induced to differentiate into enamel-secreting ameloblasts in only 10 days and apoptotically degraded in 4 weeks to achieve small-sized chimeric teeth with similar size to mouse ones, whereas the mouse secondary branchial arch epithelium, when recombined with the human dental mesenchyme from the bell-stage, remains in an enamel-secreting status after 15-week *ex vivo* culture and acquired large-sized teeth in the end, a size similar to human deciduous teeth (Hu et al., 2014a, Wang et al., 2010, Hu et al., 2018). These results strongly imply that the duration of epithelial differentiation and size control in tooth development could be dominated by the mesenchymal component in humans and mice. To validate our hypothesis, we comprehensively revisited these phenotypes by carrying out meticulously designed heterogenous recombination between human and mouse embryonic dental tissues (Table 1). As shown in Fig. 1A-C, it took more than 4 weeks for the mouse dental epithelium to deposit enamel when recombined with the human dental mesenchyme in mde+hdm chimeric tooth germs, exhibiting a much longer developmental duration and slower tempo in comparison with the mouse dental epithelia that were combined with their own dental mesenchyme (Fig. 1E-G). By contrast, it took only about 10 days for the human dental epithelium to deposit enamel when recombined with E13.5 mouse dental mesenchyme in hde+mdm chimeric tooth germs (Fig. 1I-K), demonstrating a dramatically accelerated developmental pace compared to the normal human tooth morphogenesis. When the mouse dental epithelia were recombined with the human dental mesenchyme, in addition to the compliance of developmental pace with that of the human dental mesenchyme, the chimeric tooth germ generated an enlarged tooth after 16-week growth for sufficient mineralization (Fig. 1D), whereas when the human dental epithelium was recombined with the mouse dental mesenchyme, the developmental pace of chimeric teeth did not obeys that of the human dental epithelium but that of the mouse dental

mesenchyme, and thereby achieved a compact-size tooth equivalent to the size of mouse homogenous (mde+mdm) teeth after grown under the kidney capsule for 30 days (Fig. 1H,L). Immunostaining of two ameloblastic differentiation markers, ameloblastin and MMP20 (Matrix Metallopeptidase 20), further confirmed the development phase alteration of the dental epithelium induced by the heterogenous dental mesenchyme. The expression of these two markers in the mouse dental epithelium-derived pre-ameloblasts in mde+hdm chimeric recombinants was delayed to 4^{th} week in subrenal culture (Fig. 1M-O), whereas the expression of the two markers was antedated at 9-day in the human dental epithelium-derived ameloblasts in hde+mdm chimeric recombinants in subrenal culture (Fig. 1S-U), which is similar to the expression of them at 7-day in the ameloblasts derived from the mouse dental epithelium in mde+mdm control recombinants (Fig. 1P-R). The possibility of tissue contamination was ruled out using specific primary antibodies against mouse or human MHC I antigens to distinguish between human and mouse tissues in the heterogenous chimeras (Fig. S1). Our results unambiguously indicate that the dental mesenchyme is a crucial component that dominates the tooth developmental pace and size by interaction with the dental epithelium via diffusible paracrine signals during odontogenesis.

FGF signaling participates in the regulation of tooth developmental pace and size.

We next attempted to identify the paracrine signaling molecules that may be involved in the regulation of tooth developmental tempo in humans and mice. Despite the fact that Hippo pathway that is present during tooth development as well is currently the only well-recognized signaling pathway in the control of the size and growth rate for several organs such as liver, knockout or overexpression experiments do not demonstrate its function in the supervision of developmental pace and size during tooth development (Liu et al., 2014, Kwon et al., 2015). Conserved SHH, FGF, BMP, and Wnt pathways are all crucial signaling engaged in cell proliferation, apoptosis and differentiation during odontogenesis, indicating that they may be key players in orchestrating the fundamental mainspring of organ growth. Since the dental mesenchyme is dominant in the regulation of tooth growth (Fig. 1), we exclude epithelially expressed SHH (Hu et al., 2013). To test

whether mesenchymal BMP, FGF and WNT signaling may be involved in the regulation of tooth developmental pace and size, we examined the level of the active indicators of these pathway including pSmad1/5/8 for BMP, K-Ras for FGF, and β-catenin for Wnt, respectively, in the chimeric tooth germs. The expression level of K-Ras in both the dental epithelium and mesenchyme is much higher in the mde+hdm chimeric tooth germ (Fig. 2E), where the development of mouse dental epithelium-derived ameloblasts were dramatically delayed and remained immature after 4 weeks in subrenal culture, compared to that in the mouse homogenous recombinant tooth germ (Fig. 2B). By contrast, the expression levels of pSmad1/5/8 and β-catenin did not display obvious difference as compared between the heterogenous recombinants and homogenous ones (Fig. 2A,C,D,F). The enhanced epithelial FGF activity in the mde+hdm chimeric tooth germ, evidenced by outstandingly enhanced activity of K-Ras, was certainly induced by the human dental mesenchyme and may attribute to the protracted differentiation of the mouse dental epithelium.

To explore whether the upregulated FGF signaling may be associated with the prolonged tooth development phase and large tooth size in humans, we started with a search for candidate FGFs that are persistently expressed in the human developing dental mesenchyme but barely in the mouse one. Most expression patterns of 22 known FGF ligands in mammals have been verified in mouse tooth germs at the cap and bell stages (summarized in Table S1)(Li et al., 2014, Cam et al., 1992). We have examined in our previous studies the expression patterns of *FGF3, FGF4, FGF7, FGF8, FGF9,* and *FGF10* in human tooth germs at different stages using *in situ* hybridization or immunohistochemistry (Huang et al., 2015). Here, we screened the expression of additional 12 *FGFs* that have not been identified in human tooth germs by RT-PCR. We excluded *FGF11, FGF12, FGF13,* and *FGF14*, because they are not secreting ligands and do not activate FGF receptors as well. We found that, among them, the transcripts of *FGF1*, *FGF2*, *FGF15/19*, and *FGF18* were detectable at both the cap and bell stages (data not shown). We further confirmed their expression in developing human teeth by section *in situ* hybridization (Fig. S2). Comparison of FGF expression data between the human (Huang et al., 2015) and mouse (Li et al., 2014, Cam et al., 1992), we noticed that

the majority of FGF transcripts are broadly distributed in the whole tooth germ including the dental epithelium and mesenchyme throughout the early cap stage to the late bell stage in humans (Huang et al., 2015). However, their expression is restricted to the limited region of tooth germs, such as enamel knot, and appeared in limited periods of time in mice (Table S1)(Li et al., 2014, Cam et al., 1992). We believe that this widely and persistently distributing pattern of FGF ligands in the human tooth germs is closely related to their characteristics of tooth development. To further verify this notion, we cultured human 1st premolar germs at the early bell stage under the kidney capsule of nude mice with intraperitoneal injection of SU5402, a global FGF-signaling inhibitor, for one week and found accelerated detinogenesis in the SU5402-treated molar germ with the appearance of both predentin and dentin (Fig. 2G), as compared to DMSO-treated one with appearance of only predentin (Fig. 2H), indicating pivotal roles of FGF-signaling in regulation of tooth developmental tempo.

With further analyses of FGF expression data in humans and mice, we found that FGF4, FGF8, and FGF9 ligands exhibit most distinctive patterns (Table S1). These three ligands are persistently expressed in the dental epithelium and mesenchyme at least until the differentiation stage in humans (Huang et al., 2015). However, in mice, the expression of FGF4 is restricted to the enamel knot from the cap to bell stage and that of FGF9 to the enamel knot at the cap stage and to the dental epithelium at the bell stage. Furthermore, FGF8 is only expressed in the presumptive dental epithelium prior to and during the laminar stage and is no longer expressed after the E12 bud stage (Kettunen and Thesleff, 1998). Whereas, in humans, *FGF8* transcripts are widely and intensively present in the whole tooth germ at least from the early cap stage to the differentiation stage (Huang et al., 2015). Importantly, we found retained expression of FGF8, as well as FGF4 and FGF9, in the mouse dental epithelium derived pre-ameloblasts in mde+hdm recombinants after 4 week in subrenal culture, where the development of mouse ameloblasts was largely delayed (Fig. 2I-K). Based on these observations, we hypothesized that FGF signaling may play important roles and FGF8 could be a critical factor in the regulation of tooth developmental pace and size.

Forced activation of FGF8 signaling in the mouse dental mesenchyme delays tooth development and enlarges tooth size.

To test our hypothesis, we set to overexpress *Fgf8* in dental mesenchymal cells in the E13.5 recombinant molar germ via lentivirus-mediated expression and subrenal culture system. Histological examination revealed that it took up to 9 days for *Fgf8*-overexpressed tooth grafts to start dentin secreting (Fig. 3B), whereas it only took 7 days for control ones to do so (Fig. 3A). Moreover, Ki67 staining detected remarkably higher level of proliferation rate in both the dental epithelium and mesenchyme of *Fgf8*-overexpressed tooth grafts compared to the controls after 7 days in subrenal culture (Fig. 3C-E; *n*=6/group). These data indicate that activated mesenchymal FGF8 activity delays tooth development and dramatically stimulates cell proliferation.

To corroborate the aforementioned *ex vivo* results, we further generated *Wnt1-Cre; R26R^{Fgf8}* transgenic mice to verify whether forced activation of mesenchymal FGF8 signaling alters tooth development *in vivo*. The presence of FGF8 in the dental mesenchyme of *Wnt1-Cre;R26R^{Fgf8}* mice was confirmed by anti-FGF8 immunostaining (Fig. S3). Most of mutant embryos die at around E12.5 and a few died around E14.5. Although histological examination revealed severe craniofacial hypoplasia in 100% cases, including exencephaly and the absence of identifiable tissues such as ears, auricles, eyes, nose, as well as tongue (data not shown), the mutant mice still possessed a pair of intact maxillary and mandibular molar germs (Fig. 4). More importantly, both the maxillary and mandibular molar germ exhibited an obvious retardation of tooth development (Fig. 4). Since most mutant embryos died at around E12.5, we grafted E11.5 mutant molar germs under the kidney capsule for further development. Examination of histology and the deposition of dentin and enamel showed the mutant teeth developed with approximate one-day's delay when compared to the wild-type ones (Fig. 5A-F). Examination of the expression of odontoblast marker DSP and ameloblast marker ameloblastin in the grafted teeth further confirmed this delay (Fig. 5I-L).

Interestingly, Azan staining showed that the mutant graft formed a larger-sized tooth (Fig. 5G) than the wild-type one (Fig. 5H) after 18 days in subrenal culture. To precisely evaluate the effect of mesenchymal *Fgf8* on tooth size, the tooth grafts were cultured

under the kidney capsule for 3 months for complete mineralization. Micro-CT scanning revealed that the mutant graft indeed formed a larger-sized tooth with short and dumpy shape (Fig 5N) compared to the control (Fig. 5M). Measurement of the height and width of the tooth crowns exhibited no difference in the average height of the tooth crowns between the mutants ($n = 11$) and wild-types ($n = 7$) but about 20% wider in the average width in the mutants compared to the controls (Fig. 5O). Apparently, higher rate of cell proliferation contributes greatly to the larger teeth, which was evidenced by more Ki67 positive cells presented in E13.5 mutant tooth germ and the grafted mutant teeth than controls (Fig. 6A-F, K-M, $n = 3$/each group). Meanwhile, while apoptotic marker Caspase3 staining was present in ameloblasts and odontoblasts in both mutant (Fig. I,J) and wild-type tooth germs (Fig. 6G,H), this apoptotic signal was non-detectable in the pulp cells of mutant tooth germs (Fig. 6I,J), whereas it was seen in that of wild-type ones (Fig. 6G,H) after 18 days in subrenal culture. These results further support the idea that ectopically activated FGF8 signaling in the mouse dental mesenchyme stimulates cell proliferation and prevents cell apoptosis during odontogenesis.

Ectopically activated dental mesenchymal FGF8 signaling is transduced via p38 and PI3K-Akt intracellular pathway.

To determine which intracellular pathway is implicated in the signal transduction of the ectopically activated dental mesenchymal FGF8 signals, we examined the levels of phosphorylated Erk1/2, P38, JNK, and PI3K-Akt, the activity indicator of MAPK pathway, in both E13.5 tooth germs and E11.5 ones after 9 days in subrenal culture. As shown in Fig. 7, no signal intensity differences in pErk1/2 (Fig. 7A-D) and pJNK (Fig. 7E-H) staining were observed between wild-type and mutant teeth. Whereas, p38 signaling is dramatically elevated in *Wnt1-Cre;R26R^{Fg/8}* teeth with a considerably increased amount of nuclear-localized pP38 cells both in the dental epithelium and dental mesenchyme of E13.5 mutant tooth germs (Fig. 7J) and grafted teeth (Fig. 7L) as compared to the controls (Fig. 7I,K). Meanwhile, the intensity of PI3K-Akt staining was slightly enhanced and the amount of pAkt1$^+$ cells was increased in mutant E13.5 tooth germs and grafted teeth (Fig. 7N,P) as compared to controls (Fig7 M,O). Western blot

analysis further confirmed the increased expression of pP38 and pAkt1 in E13.5 mutant

tooth germs (Fig. 7Q,R; n = 3). Our results indicate that the augmented dental

mesenchymal FGF8 activity enhances P38 and PI3K-Akt signaling in both the dental

mesenchyme and the dental epithelium, suggesting that they may function as intracellular

pathways for transducing FGF8 signal into nuclei.

Enhanced dental mesenchymal FGF8 signaling promotes transition of cell cycle

from G1- to S-stage during odontogenesis.

The rate of cell proliferation controlled by cell cycle progression is a key player in

organ size control (Gokhale and Shingleton, 2015) and is also closely associated with

tissue differentiation status, as higher cell proliferation always presents in developing

tissues and lower one in differentiated ones. Since the elevation of *Fgf8* expression in the

dental mesenchyme results in the significant promotion of cell proliferation and

prevention of cell apoptosis (Fig. 6), we speculated that enhanced dental mesenchymal

FGF8 signaling would alter cell cycling. To verify this hypothesis, we set forth to

ascertain the distribution of cyclin D1 (Ccnd1) and its inhibitor p21^{Cip1} (p21), a pair of

critical regulators functioning stimulatory and repression effect, respectively, on inducing

cell cycle entry and mediating the G1-S stage transition (Baldin et al., 1993, Gartel and

Radhakrishnan, 2005), in E13.5 tooth germs and grafted teeth. As shown in Fig. 8,

although the same high density of Ccnd1$^+$ cells were present in the dental epithelium of

E13.5 wild-type and mutant tooth germs, the dental mesenchyme in mutant tooth germs

possessed an considerably increased amount of Ccnd1$^+$ cells in comparison to that in

wild-type ones (Fig. 8A,E). On the other hand, p21$^+$ cells were abundantly present in the

presumptive primary enamel knot of the dental epithelium and slightly in the dental

mesenchyme in wild-type tooth germs but totally absent from the whole tooth germ in

mutant ones (Fig. 8B,F). Thereout, more nuclear colocalization of Ccnd1 and p21 staining

were found in the presumptive primary enamel knot of the wild-type tooth germs than in

the mutant ones (Fig. 8C-D, G-H), which makes p21 functioning with Ccnd1 to regulate

the cell cycle progression and inhibits cell proliferation in the epithelial signal center in

the wild-type tooth germ. In grafted teeth, nuclear Ccnd1$^+$ cells were distributed not only

in ameloblasts and odontoblasts (Fig. 8M,P-up) but also all over the dental pulp in mutants (Fig. 8M,P-down), rather than being restricted to the dental mesenchymal cells adjacent to the odontoblasts in wild-types (Fig. 8I,L). Although p21 was expressed in ameloblasts and dental mesenchymal cells in both wild-type and mutant teeth, its expression level was relatively decreased in mutants (Fig. 8J,L,N,P). In addition, the nuclear localization of p21 could be detected in dental mesenchymal cells of wild-type teeth (Fig. 8L-down), whereas it was found to be in the cytoplasm of grafted mutant teeth (Fig. 8P-down), which make it unable to cooperate with nuclear Ccnd1 to regulate the cell cycle progression in the dental mesenchymal cells. Taken together, these results provide evidence that enhanced mesenchymal FGF8 activity observably elevates the activity of nuclear Ccnd1 while degrades the activity of nuclear p21. Thus both stimulation of Ccnd1 and repression of p21 function to induce cell cycle entry and promote the transition of cell cycle from G1- to S- stage, which facilitates cell population to retain the high rate of proliferation.

Discussion

Hippo signaling is the only admitted pathway that plays its best-known physiological function on the control of organ size in the mouse model for mammalian lineage (Barry et al., 2013, Liu et al., 2014, Yu et al., 2015, Zhang et al., 2017), which mainly involves in the regulation of cell proliferation and apoptosis and thereby cell number (Tumaneng et al., 2012). Activation or inhibition of YAP/TAZ, a transcriptional co-activator of Hippo pathway, leads to the adjustment of cell proliferation and apoptosis, thus affecting organ sizes and tissue homeostasis (Barry et al., 2013, Liu et al., 2014, Yu et al., 2015, Zhang et al., 2017). However, in tooth development, suppression of YAP *ex vivo* at the cap stage only leads to ectopic cusp formation (Kwon et al., 2015), while overexpression of YAP *in vivo* disrupts tooth morphogenesis with abnormal enamel knot as well as greatly widened dental lamina; teeth ultimately formed in YAP transgenic mice do not exhibit any growth delay or size enlargement (Liu et al., 2014). These data suggest that the renowned control function of Hippo pathway in mammalian organ size is not applicable for the tooth organ. In support of this notion, it has also been reported that size-control mechanisms differ across mammalian organs (Penzo-Mendez and Stanger, 2015) and enhanced Hippo signaling induces organ enlargements in the stomach, heart, liver, and spleen, but not in lung, kidney, and intestine (Song et al., 2010, Heallen et al., 2011, Dong et al., 2007). Besides, IGF signaling is the other pathway that has been implied to have effects on organ size since *Igf2r* disrupted mice are born 25-35% bigger than wild-types, however, it has also been revealed that IGF signaling affects the size in some organs such as the cecum and colon but not the size of other organs such as the skin and stomach (Zaina and Squire, 1998), which also support the notion.

A heterogenous recombination study carried out between the mouse and rat molar tooth germs has suggested that tooth size is determined by the dental mesenchyme (Cai et al., 2007, Jiang et al., 1999). Additionally, our recent studies showed that, in comparison with natural developmental pace, the differentiation duration of the mouse dental epithelium or non-dental epithelium into ameloblasts when induced by the human dental mesenchyme is significantly extended, whereas that of the human dental

epithelium or epithelial stem cells into ameloblasts when induced by the mouse dental

mesenchyme is significantly shortened (Wang et al., 2010, Hu et al., 2018). These results

are consistent with the difference of development progression speed between human and

mouse teeth (Zhang et al., 2005), suggesting a dominant role of the dental mesenchyme

in regulation of tooth developmental tempo. This idea is further supported by our current

heterogenous recombination study carried out between the mouse and human molar tooth

germs (Fig. 1). We further provide unambiguous evidence that not only the tooth size but

also the tooth developmental pace are obedient to hereditary character of the dental

mesenchyme.

Conserved pathways, such as FGF, BMP, Hh, and WNT pathway, participate in the

highly conserved molecular mechanism underlying organismal growth by corresponding

alterations of cell growth and division (Penzo-Mendez and Stanger, 2015, Lui et al., 2015,

Gokhale and Shingleton, 2015). FGFs, as the most common mitogens, are positively

correlated to the rate of cell growth and cell division and involved in the induction and

pattern formation during tooth development (Brewer et al., 2016, Lan et al., 2014). Fgf8

has been demonstrated to play an important role in cell survival and proliferation of CNC

cells (Abzhanov et al., 2003), and to exert a positive effect on proliferation of dental

mesenchymal cells (Kettunen and Thesleff, 1998). Our data show that FGF signaling

rather than BMP and WNT signaling is highly active in human-mouse chimeric tooth

germs where the differentiation duration of the mouse dental epithelium is largely slowed

down (Fig. 2), which is consistent with the continuously high expression of FGFs,

particularly FGF8, in human tooth germs, suggesting a potentially positive role of FGF

signaling in regulating the tooth developmental tempo and size during odontogenesis in

humans. Both our *ex vivo* lentivirus-mediated overexpression system and *in vivo*

transgenic mouse model support that forcibly activated dental mesenchymal FGF8

signaling stimulates cell proliferation and inhibits cell apoptosis, thus prolonging tooth

developmental pace and enlarges tooth size in mice (Fig. 3-6). FGF8 appears to be a

critical factor in regulating the tooth developmental pace and size during mammalian

odontogenesis.

Functional redundancy complicates interpretation of FGF genetic studies.

Phenotypes in various *Fgfr* knockout mice have demonstrated that different FGF receptors have both essential and redundant roles throughout development (Brewer et al., 2016). During odontogenesis, FGF7 and FGF10 have been proven to act as surrogates of mesenchymal FGF3 (Kratochwil et al., 2002), and FGF4, FGF8, as well as FGF9 have also been suggested functional redundancy and repetitive use in mouse tooth development (Zhang et al., 2005). Our previous and current study show that several FGFs, especially FGF4, FGF8, and FGF9, have a broad and persistent expression in the whole tooth germs in humans but are restricted to the limited region of tooth germs in mice (Huang et al., 2015) (Table S1). These data suggest that tooth developmental pace and size may be the product of synergistic effects of different FGFs in mammals. Actually, in addition to the fact that FGF8 is a key factor to control tooth developmental pace and size discovered in the current study, Fgf9 has also been reported to induce cell proliferation in cultured mouse dental mesenchyme *in vitro* (Kettunen and Thesleff, 1998) as well as in cervical loop development *in vivo* (Kurosaka et al., 2011) and the *Fgf9* null mutant mice show a size-reduced labial cervical loop (Kurosaka et al., 2011), and FGF4 stimulates cell proliferation in both the dental epithelium and mesenchyme as well (Kettunen and Thesleff, 1998, Jernvall et al., 1994), both suggesting their positive function on the regulation of tooth growth and size. We speculate that, in humans, several broadly, intensively, and persistently expressed FGF ligands may act in concert to confer growth capability (stimulation of proliferation and prevention of apoptosis) on tooth tissues to achieve longer development time-scale and larger size.

Transduction of FGF signaling is mediated by multiple intracellular pathways, including P38, PI3K/Akt, Erk1/2, and JNK (Brewer et al., 2016). In our previous study, the similar expression patterns of the indicators in Erk, P38, JNK, and PI3K-Akt pathways in mouse and human tooth germs suggests their conservative regulatory functions during mammalian odontogenesis (Huang et al., 2015). P38 has been reported to be involved in control of the cell cycle via the expression changes of P21, P27 and Ccnd3 in mouse skeletal muscle cells (Irazoqui et al., 2014), Ccnd1 and c-Myc in human dental pulp stem cells (Vandomme et al., 2014), and Ccnd1 and P21 in rat mesenchymal stem cells (Stockl et al., 2013), which is also significantly increased in our *Wnt1-Cre; R26R^{Fgf8}* mice (Fig.

7). The PI3K-Akt pathway is associated with a variety of cellular functions including cell

growth, cell cycle regulation, and cell survival (Castellano and Downward, 2011). The

increased PI3K-Akt level regulates the proliferation and differentiation of neural

progenitor cells by the activation of the cell cycle through the elevation of the Ccnd1 and

cyclin E, two cyclins essential for the G1-S transition during organogenesis (Xue and

Yuan, 2010). Moreover, Akt protein is a central intracellular factor to transduce and

integrate multiple extracellular growth factor signals, such as FGF, BMP, and GH/IGF,

into a comprehensive signal network to supervise the processes of cell proliferation,

growth, and survival (Penzo-Mendez and Stanger, 2015). The increased activity of PI3K-

Akt pathway caused by augmented FGF8 in the cranial neural crest cell (Fig. 7) further

supports its positive function in cell proliferation regulated by cell cycle during tooth

development. Erk has been reported to provide a switch-like transition between

proliferation and differentiation of muscle progenitors (Michailovici et al., 2014), while

JNK pathway has been elucidated to play crucial role in negative growth regulation and

likely performs a central regulator in both local growth coordination and global growth

coordination (Gokhale and Shingleton, 2015). These two pathways did not show any

noticeable alteration in our *Fgf8* transgenic mice, which may attribute to their main

function in control of negative growth or balancing cell proliferation and differentiation.

Cell growth and division are controlled by the cyclin depend kinases (CDKs) and

their binding partners (cyclins) that regulate cell cycle progression through the regulation

of the G1-S and G2-M transition. Many cell cycle regulators have been proved to control

organ growth and size, which appears to be generally conserved in *Drosophila* and

mammals (Gokhale and Shingleton, 2015). The G1-S phase transition is a rate-limiting

step, a known checkpoint between cell growth and cell division, requiring cell cycle

regulators to monitor and regulate the progression of the cell cycle (Elledge, 1996). The

stimulatory effects of cyclins can be offset by CDKs inhibitors (CDKIs) and the

repression of CDKIs is also a prerequisite for G1-S transition. Thus merely elevating the

level of cyclin D1 is not sufficient to induce cell cycle entry (Chambard et al., 2007).

Cyclin D1 (Ccnd1) is synthesized rapidly in the G1 phase and accumulates in the nucleus

and degraded as the cell enters the S phase (Baldin et al., 1993). P21, known as CDKI1,

functions as a regulator of cell cycle progression in G1 and S phase and inhibits all cyclin/CDK complexes, which in turn results in cell cycle arrest and/or apoptosis (Gartel and Radhakrishnan, 2005). Enhanced dental mesenchymal FGF8 may promotes cell cycle progression through the regulation of the G1-S transition, which further stimulates cell proliferation and prevents cell differentiation and apoptosis, thereby resulting in prolonged tooth development and larger tooth size. In line with this notion, we found that the amount of nuclear Ccnd1$^+$ cells is dramatically increased while that of nuclear P21$^+$ cells is markedly decreased in the dental mesenchyme at both the early and differentiation stage in *Wnt1-Cre;R26R^{Fgf8}* mice as compared to those in the control (Fig. 8), stating that forcedly enhanced FGF8 in the dental mesenchyme could elevate the level of Ccnd1 and meanwhile suppress the level of p21. Stimulation of Ccnd1 and repression of p21 both constitute the prerequisite for inducing cell cycle entry and thereby promoting the G1-S transition.

Two latest studies have demonstrated that interspecies differences in developmental tempo are caused by cell-autonomous differences in biochemical reaction speeds instead of differences in signaling, the sequence of genes, or regulatory elements in *in vitro* human and mouse platforms (Rayon et al., 2020, Matsuda et al., 2020). Moreover, in *Drosophila*, it is also reported that the *Minute* mutation causes a prolongation of developmental pace and decreased cell division rate in a cell-autonomous way via decreasing the expression of ribosomal proteins and thereby reducing the rate of translation and protein synthesis (Saebøe-Larssen et al., 1998, Morata and Ripoll, 1975). Developmental allochrony regulated in a cell-autonomous manner is highlighted in these studies. However, our findings that the dental mesenchyme resorts to extrinsic signals such as FGF to dominate the developmental tempo and size as well via tissue interaction in a non-cell-autonomous manner during tooth development are inconsistent with these studies. Additionally, the fact that the oscillation phase shift between Notch and Wnt signaling is critical for the segmentation of presomitic mesoderm cells in multicellular organ development in 2D assays *ex vivo* (Sonnen et al., 2018) also indicates the pivotal role of secretory signal changes in the regulation of developmental pace. In *Drosophila*, organ growth and size are orchestrated not only by the growth rate regulated via signaling

pathways but also by the growth duration regulated via hormonal signals and organ-
autonomous processes controlled by patterning genes and physical force (Gokhale and
Shingleton, 2015). The developmental tempo and size of individual organs must be tightly
coordinated to match the body size and larger-bodied species always keep slower
developmental tempos to acquire larger organ sizes. Our heterogenous recombination
results provide compelling evidence to support the contribution of extrinsic cues in the
developmental tempo and size of teeth relying on tissue interaction during early
development. Thus, we suggest that complicated regulatory networks involved in the
developmental pace and size work both in a cell-autonomous and non-cell-autonomous
way to orchestrate the organ morphogenesis, and, particularly, tissue interaction in a non-
cell-autonomous way plays a major role during early organogenesis.

Tissue bioengineering based on the principle of developmental biology represents a
promising approach for repair and regeneration of damaged tissues and organs in the
future. However, unmatched development phase and organ size between humans and the
model animals such as *Drosophila*, zebra fishes, and mice appear to be a major obstacle
to acquire knowledge about organ growth control in humans. Uncovering of the molecular
mechanism underlying organ growth control in humans is a prerequisite for the realization
of replacement therapy in the future. Our study provides important FGF signaling
information in tooth developmental process and enlightens our knowledge about the
nature of FGF signaling in human tooth growth.

Acknowledgments

This study was supported by the National Natural Science Foundation of China (81870739, 82001002, 81271102, 81771034) and the Natural Science Foundation of Fujian Province (2019J01281, 2020J01180).

References

Abzhanov, A., Tzahor, E., Lassar, A. B. & Tabin, C. J. 2003. Dissimilar regulation of cell differentiation in mesencephalic (cranial) and sacral (trunk) neural crest cells in vitro. *Development,* 130**,** 4567-79.

Baldin, V., Lukas, J., Marcote, M. J., Pagano, M. & Draetta, G. 1993. Cyclin D1 is a nuclear protein required for cell cycle progression in G1. *Genes Dev,* 7**,** 812-21.

Barry, E. R., Morikawa, T., Butler, B. L., Shrestha, K., de la Rosa, R., Yan, K. S., Fuchs, C. S., Magness, S. T., Smits, R., Ogino, S., Kuo, C. J. & Camargo, F. D. 2013. Restriction of intestinal stem cell expansion and the regenerative response by YAP. *Nature,* 493**,** 106-10.

Brewer, J. R., Mazot, P. & Soriano, P. 2016. Genetic insights into the mechanisms of Fgf signaling. *Genes Dev,* 30**,** 751-71.

Cai, J., Cho, S. W., Kim, J. Y., Lee, M. J., Cha, Y. G. & Jung, H. S. 2007. Patterning the size and number of tooth and its cusps. *Dev Biol,* 304**,** 499-507.

Cam, Y., Neumann, M. R., Oliver, L., Raulais, D., Janet, T. & Ruch, J. V. 1992. Immunolocalization of acidic and basic fibroblast growth factors during mouse odontogenesis. *Int J Dev Biol,* 36**,** 381-9.

Castellano, E. & Downward, J. 2011. RAS Interaction with PI3K: More Than Just Another Effector Pathway. *Genes Cancer,* 2**,** 261-74.

Chambard, J. C., Lefloch, R., Pouyssegur, J. & Lenormand, P. 2007. ERK implication in cell cycle regulation. *Biochim Biophys Acta,* 1773**,** 1299-310.

Danielian, P. S., Muccino, D., Rowitch, D. H., Michael, S. K. & McMahon, A. P. 1998. Modification of gene activity in mouse embryos in utero by a tamoxifen-inducible form of Cre recombinase. *Curr Biol,* 8**,** 1323-6.

Dong, J., Feldmann, G., Huang, J., Wu, S., Zhang, N., Comerford, S. A., Gayyed, M. F., Anders, R. A., Maitra, A. & Pan, D. 2007. Elucidation of a universal size-control mechanism in Drosophila and mammals. *Cell,* 130**,** 1120-33.

Dong, X., Shen, B., Ruan, N., Guan, Z., Zhang, Y., Chen, Y. & Hu, X. 2014. Expression patterns of genes critical for BMP signaling pathway in developing human primary tooth germs. *Histochem Cell Biol,* 142**,** 657-65.

Edelman, G. M. 1992. Morphoregulation. *Dev Dyn,* 193**,** 2-10.

Elledge, S. J. 1996. Cell cycle checkpoints: preventing an identity crisis. *Science,* 274**,** 1664-72.

Gartel, A. L. & Radhakrishnan, S. K. 2005. Lost in transcription: p21 repression, mechanisms, and consequences. *Cancer Res,* 65**,** 3980-5.

Gokhale, R. H. & Shingleton, A. W. 2015. Size control: the developmental physiology of body and organ size regulation. *Wiley Interdiscip Rev Dev Biol,* 4**,** 335-56.

Heallen, T., Zhang, M., Wang, J., Bonilla-Claudio, M., Klysik, E., Johnson, R. L. & Martin, J. F. 2011. Hippo pathway inhibits Wnt signaling to restrain cardiomyocyte proliferation and heart size. *Science,* 332**,** 458-61.

Hu, X., Lee, J. W., Zheng, X., Zhang, J., Lin, X., Song, Y., Wang, B., Hu, X., Chang, H. H., Chen, Y., Lin, C. P. & Zhang, Y. 2018. Efficient induction of functional ameloblasts from human keratinocyte stem cells. *Stem Cell Res Ther,* 9**,** 126.

Hu, X., Lin, C., Shen, B., Ruan, N., Guan, Z., Chen, Y. & Zhang, Y. 2014a. Conserved odontogenic potential in embryonic dental tissues. *J Dent Res,* 93**,** 490-5.

Hu, X., Xu, S., Lin, C., Zhang, L., Chen, Y. & Zhang, Y. 2014b. Precise chronology of differentiation
of developing human primary dentition. *Histochem Cell Biol,* 141**,** 221-7.

Hu, X., Zhang, S., Chen, G., Lin, C., Huang, Z., Chen, Y. & Zhang, Y. 2013. Expression of SHH
signaling molecules in the developing human primary dentition. *BMC Dev Biol,* 13**,** 11.

Huang, F., Hu, X., Fang, C., Liu, H., Lin, C., Zhang, Y. & Hu, X. 2015. Expression profile of critical
genes involved in FGF signaling pathway in the developing human primary dentition.
Histochem Cell Biol, 144**,** 457-69.

Irazoqui, A. P., Boland, R. L. & Buitrago, C. G. 2014. Actions of 1,25(OH)2-vitamin D3 on the cellular
cycle depend on VDR and p38 MAPK in skeletal muscle cells. *J Mol Endocrinol,* 53**,** 331-
43.

Jernvall, J., Kettunen, P., Karavanova, I., Martin, L. B. & Thesleff, I. 1994. Evidence for the role of
the enamel knot as a control center in mammalian tooth cusp formation: non-dividing cells
express growth stimulating Fgf-4 gene. *Int J Dev Biol,* 38**,** 463-9.

Jernvall, J. & Thesleff, I. 2000. Reiterative signaling and patterning during mammalian tooth
morphogenesis. *Mech Dev,* 92**,** 19-29.

Jheon, A. H., Seidel, K., Biehs, B. & Klein, O. D. 2013. From molecules to mastication: the
development and evolution of teeth. *Wiley Interdiscip Rev Dev Biol,* 2**,** 165-82.

Jiang, T. X., Jung, H. S., Widelitz, R. B. & Chuong, C. M. 1999. Self-organization of periodic patterns
by dissociated feather mesenchymal cells and the regulation of size, number and spacing of
primordia. *Development,* 126**,** 4997-5009.

Kettunen, P. & Thesleff, I. 1998. Expression and function of FGFs-4, -8, and -9 suggest functional
redundancy and repetitive use as epithelial signals during tooth morphogenesis. *Dev Dyn,* 211**,**
256-68.

Kratochwil, K., Galceran, J., Tontsch, S., Roth, W. & Grosschedl, R. 2002. FGF4, a direct target of
LEF1 and Wnt signaling, can rescue the arrest of tooth organogenesis in Lef1(-/-) mice. *Genes
Dev,* 16**,** 3173-85.

Kurosaka, H., Islam, M. N., Kuremoto, K., Hayano, S., Nakamura, M., Kawanabe, N., Yanagita, T.,
Rice, D. P., Harada, H., Taniuchi, I. & Yamashiro, T. 2011. Core binding factor beta functions
in the maintenance of stem cells and orchestrates continuous proliferation and differentiation
in mouse incisors. *Stem Cells,* 29**,** 1792-803.

Kwon, H. J., Li, L. & Jung, H. S. 2015. Hippo pathway/Yap regulates primary enamel knot and dental
cusp patterning in tooth morphogenesis. *Cell Tissue Res,* 362**,** 447-51.

Lan, Y., Jia, S. & Jiang, R. 2014. Molecular patterning of the mammalian dentition. *Semin Cell Dev
Biol,* 25-26**,** 61-70.

Li, C. Y., Prochazka, J., Goodwin, A. F. & Klein, O. D. 2014. Fibroblast growth factor signaling in
mammalian tooth development. *Odontology,* 102**,** 1-13.

Lin, C., Yin, Y., Bell, S. M., Veith, G. M., Chen, H., Huh, S. H., Ornitz, D. M. & Ma, L. 2013.
Delineating a conserved genetic cassette promoting outgrowth of body appendages. *PLoS
Genet,* 9**,** e1003231.

Line, S. R. 2003. Variation of tooth number in mammalian dentition: connecting genetics,
development, and evolution. *Evol Dev,* 5**,** 295-304.

Liu, M., Zhao, S. & Wang, X. P. 2014. YAP overexpression affects tooth morphogenesis and enamel
knot patterning. *J Dent Res,* 93**,** 469-74.

Lloyd, A. C. 2013. The regulation of cell size. *Cell,* 154**,** 1194-205.

Lui, J. C., Garrison, P. & Baron, J. 2015. Regulation of body growth. *Curr Opin Pediatr,* 27**,** 502-10.

Matsuda, M., Hayashi, H., Garcia-Ojalvo, J., Yoshioka-Kobayashi, K., Kageyama, R., Yamanaka, Y.,
Ikeya, M., Toguchida, J., Alev, C. & Ebisuya, M. 2020. Species-specific segmentation clock
periods are due to differential biochemical reaction speeds. *Science,* 369**,** 1450-1455.

Michailovici, I., Harrington, H. A., Azogui, H. H., Yahalom-Ronen, Y., Plotnikov, A., Ching, S.,
Stumpf, M. P., Klein, O. D., Seger, R. & Tzahor, E. 2014. Nuclear to cytoplasmic shuttling of
ERK promotes differentiation of muscle stem/progenitor cells. *Development,* 141**,** 2611-20.

Morata, G. & Ripoll, P. 1975. Minutes: mutants of drosophila autonomously affecting cell division
rate. *Dev Biol,* 42**,** 211-21.

Penzo-Mendez, A. I. & Stanger, B. Z. 2015. Organ-Size Regulation in Mammals. *Cold Spring Harb*
Perspect Biol, 7**,** a019240.

Plikus, M. V., Zeichner-David, M., Mayer, J. A., Reyna, J., Bringas, P., Thewissen, J. G., Snead, M.
L., Chai, Y. & Chuong, C. M. 2005. Morphoregulation of teeth: modulating the number, size,
shape and differentiation by tuning Bmp activity. *Evol Dev,* 7**,** 440-57.

Polly, P. D. 2015. Gene networks, occlusal clocks, and functional patches: new understanding of
pattern and process in the evolution of the dentition. *Odontology,* 103**,** 117-25.

Promislow, D. E. L. & Harvey, P. H. 1990. Living fast and dying young: A comparative analysis of
life-history variation among mammals. *Proceedings of the Zoological Society of London,* 220**,**
417-437.

Rayon, T., Stamataki, D., Perez-Carrasco, R., Garcia-Perez, L., Barrington, C., Melchionda, M.,
Exelby, K., Lazaro, J., Tybulewicz, V. L. J., Fisher, E. M. C. & Briscoe, J. 2020. Species-
specific pace of development is associated with differences in protein stability. *Science,* 369.

Rodrigues, H. G., Renaud, S., Charles, C., Le Poul, Y., Sole, F., Aguilar, J. P., Michaux, J., Tafforeau,
P., Headon, D., Jernvall, J. & Viriot, L. 2013. Roles of dental development and adaptation in
rodent evolution. *Nat Commun,* 4**,** 2504.

Saebøe-Larssen, S., Lyamouri, M., Merriam, J., Oksvold, M. P. & Lambertsson, A. 1998. Ribosomal
protein insufficiency and the minute syndrome in Drosophila: a dose-response relationship.
Genetics, 148**,** 1215-24.

Salazar-Ciudad, I. 2012. Tooth patterning and evolution. *Curr Opin Genet Dev,* 22**,** 585-92.

Schoenwolf GC, B. S., Brauer PR, Francis-West PH 2014. Larsen's Human Embryology E-Book.

Song, H., Mak, K. K., Topol, L., Yun, K., Hu, J., Garrett, L., Chen, Y., Park, O., Chang, J., Simpson,
R. M., Wang, C. Y., Gao, B., Jiang, J. & Yang, Y. 2010. Mammalian Mst1 and Mst2 kinases
play essential roles in organ size control and tumor suppression. *Proc Natl Acad Sci U S A,*
107**,** 1431-6.

Song, Y., Zhang, Z., Yu, X., Yan, M., Zhang, X., Gu, S., Stuart, T., Liu, C., Reiser, J., Zhang, Y. &
Chen, Y. 2006. Application of lentivirus-mediated RNAi in studying gene function in
mammalian tooth development. *Dev Dyn,* 235**,** 1334-44.

Sonnen, K. F., Lauschke, V. M., Uraji, J., Falk, H. J., Petersen, Y., Funk, M. C., Beaupeux, M., François,
P., Merten, C. A. & Aulehla, A. 2018. Modulation of Phase Shift between Wnt and Notch
Signaling Oscillations Controls Mesoderm Segmentation. *Cell,* 172**,** 1079-1090 e12.

St Amand, T. R., Zhang, Y., Semina, E. V., Zhao, X., Hu, Y., Nguyen, L., Murray, J. C. & Chen, Y.
2000. Antagonistic signals between BMP4 and FGF8 define the expression of Pitx1 and Pitx2
in mouse tooth-forming anlage. *Dev Biol,* 217**,** 323-32.

Stockl, S., Bauer, R. J., Bosserhoff, A. K., Gottl, C., Grifka, J. & Grassel, S. 2013. Sox9 modulates

cell survival and adipogenic differentiation of multipotent adult rat mesenchymal stem cells. *J Cell Sci,* 126**,** 2890-902.

Swanson, E. M. & Dantzer, B. 2014. Insulin-like growth factor-1 is associated with life-history variation across Mammalia. *Proc Biol Sci,* 281**,** 20132458.

Tapaltsyan, V., Charles, C., Hu, J., Mindell, D., Ahituv, N., Wilson, G. M., Black, B. L., Viriot, L. & Klein, O. D. 2016. Identification of novel Fgf enhancers and their role in dental evolution. *Evol Dev,* 18**,** 31-40.

Thesleff, I. & Mikkola, M. 2002. The role of growth factors in tooth development. *Int Rev Cytol,* 217**,** 93-135.

Thesleff, I. & Sharpe, P. 1997. Signalling networks regulating dental development. *Mech Dev,* 67**,** 111-23.

Tucker, A. & Sharpe, P. 2004. The cutting-edge of mammalian development; how the embryo makes teeth. *Nat Rev Genet,* 5**,** 499-508.

Tumaneng, K., Russell, R. C. & Guan, K. L. 2012. Organ size control by Hippo and TOR pathways. *Curr Biol,* 22**,** R368-79.

Tummers, M. & Thesleff, I. 2009. The importance of signal pathway modulation in all aspects of tooth development. *J Exp Zool B Mol Dev Evol,* 312B**,** 309-19.

Vandomme, J., Touil, Y., Ostyn, P., Olejnik, C., Flamenco, P., El Machhour, R., Segard, P., Masselot, B., Bailliez, Y., Formstecher, P. & Polakowska, R. 2014. Insulin-like growth factor 1 receptor and p38 mitogen-activated protein kinase signals inversely regulate signal transducer and activator of transcription 3 activity to control human dental pulp stem cell quiescence, propagation, and differentiation. *Stem Cells Dev,* 23**,** 839-51.

Wang, B., Li, L., Du, S., Liu, C., Lin, X., Chen, Y. & Zhang, Y. 2010. Induction of human keratinocytes into enamel-secreting ameloblasts. *Dev. Biol.,* 344**,** 795-9.

Xiong, W., He, F., Morikawa, Y., Yu, X., Zhang, Z., Lan, Y., Jiang, R., Cserjesi, P. & Chen, Y. 2009. Hand2 is required in the epithelium for palatogenesis in mice. *Dev Biol,* 330**,** 131-41.

Xue, X. J. & Yuan, X. B. 2010. Nestin is essential for mitogen-stimulated proliferation of neural progenitor cells. *Mol Cell Neurosci,* 45**,** 26-36.

Yang, G., Yuan, G., Ye, W., Cho, K. W. & Chen, Y. 2014. An atypical canonical bone morphogenetic protein (BMP) signaling pathway regulates Msh homeobox 1 (Msx1) expression during odontogenesis. *J Biol Chem,* 289**,** 31492-502.

Yu, F. X., Zhao, B. & Guan, K. L. 2015. Hippo Pathway in Organ Size Control, Tissue Homeostasis, and Cancer. *Cell,* 163**,** 811-28.

Zaina, S. & Squire, S. 1998. The soluble type 2 insulin-like growth factor (IGF-II) receptor reduces organ size by IGF-II-mediated and IGF-II-independent mechanisms. *J Biol Chem,* 273**,** 28610-6.

Zhang, P., Pei, C., Wang, X., Xiang, J., Sun, B. F., Cheng, Y., Qi, X., Marchetti, M., Xu, J. W., Sun, Y. P., Edgar, B. A. & Yuan, Z. 2017. A Balance of Yki/Sd Activator and E2F1/Sd Repressor Complexes Controls Cell Survival and Affects Organ Size. *Dev Cell,* 43**,** 603-617 e5.

Zhang, Y., Wang, S., Song, Y., Han, J., Chai, Y. & Chen, Y. 2003. Timing of odontogenic neural crest cell migration and tooth-forming capability in mice. *Dev Dyn,* 226**,** 713-8.

Zhang, Y. D., Chen, Z., Song, Y. Q., Liu, C. & Chen, Y. P. 2005. Making a tooth: growth factors, transcription factors, and stem cells. *Cell Res,* 15**,** 301-16.

767 **Tables**

768 **Table. 1 Ratio of tooth formation in different types of tissue recombinants.**

Type of Recombinant		No. of Recombinant	No. of Tooth Formation	Ratio (%)
Epithelium	Mesenchyme			
mde	mdm	28	28	100
mde	hdm	78	76	97.4
hde	mdm	27	24	88.9

769 Abbreviations: mdm, E13.5 mouse dental mesenchyme; hdm, bell-stage human dental mesenchyme

770 (14-16 weeks of gestation); mde, E13.5 mouse dental epithelium; hde, bell-stage human dental

771 epithelium (14-16 weeks of gestation).

Figure legends

Fig. 1. Heterogenous recombination of human and mouse embryonic dental tissues reveals the dominant role of the dental mesenchyme in the control of tooth developmental pace and size. A-C: Azan dichromic staining (stains dentin blue and enamel red) of recombinants of E13.5 mouse dental epithelium and bell-stage human dental mesenchyme cultured under the kidney capsule for 3 weeks (**A**), 4 weeks (**B**), 8 weeks (**C**). Note that no deposition of dentin and enamel was seen in the chimeric tooth germs until 4 weeks in subrenal culture. **D**: A lateral view of mde+hdm chimeric tooth crowns after 16 weeks in subrenal culture. **E-G**: Azan dichromic staining of recombinants of E13.5 mouse dental epithelium and E13.5 mouse dental mesenchyme cultured under the kidney capsule for 6 days (**E**), 7 days (**F**), and 8 days (**G**). Note that the deposition of dentin and enamel was not seen until 7 and 8 days, respectively, in subrenal culture. **H**: A lateral view of mde+mdm tooth crowns after 30 days in subrenal culture. **I-K**: Azan dichromic staining of recombinants of bell-stage human dental epithelium and E13.5 mouse dental mesenchyme cultured under the kidney capsule for 7 days (**I**), 9 days (**J**), and 10 days (**K**). Note that the deposition of dentin and enamel was not seen until 9 and 10 days, respectively, in subrenal culture. **L**: A lateral view of hde+mdm chimeric tooth crowns after 30 weeks in subrenal culture. **M-U**: The expression of ameloblastin and MMP20 in recombinants of E13.5 mouse dental epithelium and bell-stage human dental mesenchyme cultured under the kidney capsule for 3 weeks (**M**), 4 weeks (**N** and **O**), in recombinants of E13.5 mouse dental epithelium and mouse dental mesenchyme for 6 days (**P**), 7 days (**Q** and **R**), and in recombinants of bell-stage human dental epithelium and mouse dental mesenchyme for 7 days (**S**) and 9 days (**T** and **U**). hdm, human dental mesenchyme; hde, human dental epithelium; mdm, mouse dental mesenchyme; mde, mouse dental epithelium; hod, human odontoblasts; mod, mouse odontoblasts; mam, mouse ameloblasts; d, dentin; e, enamel. Scale bar: A-C, E-G, I-K, and M-U = 50 μm, D, H, and L = 500 μm.

Fig. 2. FGF signaling is involved in the regulation of tooth developmental pace and size. A-C: Immunostaining showing the expression of pSmad1/5/8 (**A**), K-Ras (**B**), and β-catenin (**C**) in recombinants of mouse dental epithelium and mouse dental mesenchyme cultured under the kidney capsule for 7 days. **D-F**: Immunostaining showing the expression of pSmad1/5/8 (**D**), K-Ras (**E**), and β-catenin (**F**) in recombinants of mouse dental epithelium and human dental mesenchyme cultured under the kidney capsule for 4 weeks. **G-H**: Human 1^{st} premolar germs of 16^{th} week gestation were transplanted under the kidney capsule and cultured in the absence (**G**) or presence (**H**) of SU5402 for 1 week. Note the accelerated detinogenesis with the appearance of dentin (**H**, red arrow) in the presence of SU5402. **I-K**: Immunohistochemical staining showing the expression of FGF4 (**I**), FGF8 (**J**), and FGF9 (**K**) in recombinants of mouse dental epithelium and human dental mesenchyme cultured under the kidney capsule for 4 weeks. mam, mouse ameloblasts; mod, mouse odontoblasts; hod, human odontoblasts; am, ameloblasts; od, odontoblasts; sr, stellate reticulum; d, dentin; pre-d, predentin. Scale bar = 25 μm.

Fig. 3. Augmented FGF8 signaling in the mouse dental mesenchyme delays tooth development and increases cell proliferation. A, **B**: Azan dichromic staining of recombinants of E13.5 mouse dental epithelium and re-aggregate of dental mesenchymal cells infected with control LV-*eGFP* (**A**) or LV-*Fgf8-eGFP* (**B**) cultured under the kidney capsule for 6 days (insert in **A**), 7 days, and 9 days (insert in **B**). **C**, **D**: Immunostaining showing the expression of Ki67 in recombinants of E13.5 mouse dental epithelium and re-aggregate of dental mesenchymal cells infected with control LV-*eGFP* (**C**) or LV-*Fgf8-eGFP* (**D**) cultured under the kidney capsule for 7 days. **E**: Comparison of Ki67-positive cells in the dental mesenchyme and dental epithelium of the recombinants. Standard deviation values were shown as error bars, and ** indicates $p < 0.01$, $n = 6$/group. **F**: GFP staining of recombinants of E13.5 mouse dental epithelium and re-aggregate of dental mesenchymal cells infected with control LV-*eGFP* cultured under the kidney capsule for 11 days. dm, dental mesenchyme; dp, dental pulp; am, ameloblasts; sr, stellate reticulum; od, odontoblasts; d, dentin; Epi, epithelium; Mes, mesenchyme. Scale bar = 50 μm.

Fig. 4. Delayed tooth development in early molar germs of *Wnt1-Cre;R26R^{Fgf8}* mice.
A-H: H&E staining showing the morphology of maxillary and mandibular molar in E13.5 wild-type (**A,C**) and mutant (**B,D**) embryos and in E14.5 wild-type (**E,G**) and mutant (**F,H**) embryos. de, dental epithelium; dm, dental mesenchyme. Scale bar = 50 μm.

Fig. 5. Grafts of *Wnt1-Cre;R26R^{Fgf8}* molar germs exhibit delayed tooth development and enlarged tooth size. A-H: Azan dichromic staining of E11.5 wild-type and mutant molar germs cultured under the kidney capsule for 8 days (**A,B**), 9 days (**C,D**), 10 days (**E,F**), and 18 days (**G,H**). **I-L**: Immunostaining showing the expression of DSP (**I,J**) and Ameloblastin (**K,L**) in E11.5 wild-type and mutant molar grafts cultured under the kidney capsule for 9 days. **M**, **N**: Representative Micro-CT images of E11.5 wild-type (**M**) and mutant molar germs (**N**) cultured under the kidney capsule for 3 months. **O**: Comparison of the height and width of the tooth crown between wild-type and mutant mineralized teeth cultured under the kidney capsule for 3 months. Standard deviation values were shown as error bars. ns indicates $p > 0.05$, and ** indicates $p < 0.01$, $n = 7$ and 11/group in **O**. de, dental epithelium; dm, dental mesenchyme; am, ameloblasts; od, odontoblasts; d, dentin; e, enamel. Scale bar: A-H, I-L = 50 μm; M and N = 500 μm.

Fig. 6. Ectopically activated dental mesenchymal FGF8 signaling stimulates cell proliferation and prevents cell apoptosis during odontogenesis. A-F: Immunostaining showing the Ki67 positive cells in E13.5 wild-type (**A**) and mutant (**D**) molar germs and in the grafts of E11.5 wild-type and mutant molar germ cultured under the kidney capsule for 9 days (**B,E**) and 18 days (**C,F**). **G-J**: Immunostaining showing the expression of Caspase3 in the grafts of E11.5 wild-type (**G**, boxed area is magnified in **H**) and mutant (**I**, boxed area is magnified in **J**) molar germ cultured under the kidney capsule for 18 days. Note the absence of Caspase3 expression in the dental mesenchyme (red asterisk) excepting odontoblasts in mutant grafts. **K-M**: The percentage of Ki67 positive cells in the dental mesenchyme of wild-type and mutant E13.5 tooth germs (**K**) and tooth grafts cultured under the kidney capsule for 9 days (**I**) and 18 days (**M**), $n = 3$/group, * indicates $p < 0.05$. Standard deviation values were shown as error bars. am, ameloblasts; od,

odontoblasts; d, dentin; e, enamel. Scale bar = 100 μm.

Fig. 7. Ectopically activated dental mesenchymal FGF8 signaling is transduced via

p38 and PI3K-Akt intracellular pathways. A-D: Immunostaining showing the

expression of pErk1/2 in E13.5 wild-type (**A**) and mutant (**B**) molar germs and in E11.5

wild-type (**C**) and mutant (**D**) molar germs cultured under the kidney capsule for 9 days.

E-H: Immunostaining showing the expression of pJNK in E13.5 wild-types (**E**) and

mutants (**F**) molar germs and in E11.5 wild-type (**G**) and mutant (**H**) molar germs cultured

under the kidney capsule for 9 days. **I-L**: Immunostaining showing the expression of pP38

in E13.5 wild-type (**I**) and mutant (**J**) molar germs and in E11.5 wild-type (**K**) and mutant

(**L**) cultured under the kidney capsule for 9 days. **M-P**: Immunostaining showing the

expression of pAkt1 in E13.5 wild-types (**M**) and mutants (**N**) molar germs and in E11.5

wild-type (**O**) and mutant (**P**) molar germs cultured under the kidney capsule for 9 days.

Q: Western-blotting of pErk1/2, pP38, pJNK, and pAKT1 expressed in E13.5 wild-type

and mutant molar germs. **R**: Quantification of immunoblotting data in **Q**. $n = 3$/group. *

indicates $p < 0.05$. Standard deviation values were shown as error bars. de, dental

epithelium; dm, dental mesenchyme; am, ameloblasts; od, odontoblasts. Scale bar = 25

μm.

Fig. 8. Enhanced dental mesenchymal FGF8 signaling increases the level of nuclear

Ccnd1 and decreases the level of nuclear p21. Immunostaining showing the

colocalization of Ccnd1 and P21 in E13.5 wild-type (**A-D**) and mutant (**E-H**) molar germs

and E11.5 wild-type (**I-L**) and mutant (**M-P**) molar germs cultured under the kidney

capsule for 9 days. Boxed areas in (**C**), (**G**), (**K**), and (**O**) are magnified in (**D**), (**H**), (**L**),

and (**P**), respectively. Note the significantly increased proportion of Ccnd1-positive cells

(white arrow) and the shift of P21 from the nuclei to the cytoplasm (yellow arrow) in the

mutant dental mesenchyme (**P**) as compared with wild-type ones (**L**). de, dental

epithelium; dm, dental mesenchyme; am, ameloblasts; od, odontoblasts. Scale bar: A-N,

P-S = 50 μm; O, T = 25 μm.

Figure 1

Figure 2

Figure 3

897 Figure 4

900 Figure 5

Figure 6

904   Figure 7

906 Figure 8