Transcriptional activity differentiates families of Marine Group II *Euryarchaeota* in the

coastal ocean

Running title: Euryarchaeal transcriptomes in the coastal ocean

Julian Damashek[1,3], Aimee Oyinlade Okotie-Oyekan[1,4], Scott Michael Gifford[2], Alexey

Vorobev[1,5], Mary Ann Moran[1], James Timothy Hollibaugh[1]

[1]Department of Marine Sciences, University of Georgia, Athens, GA, USA

[2]Department of Marine Sciences, University of North Carolina, Chapel Hill, NC, USA

[3]Present address: Department of Biology, Utica College, Utica, NY, USA

[4]Present address: Environmental Studies Program, University of Oregon, Eugene, OR, USA

[5]Present address: INSERM U932, PSL University, Institut Curie, Paris, France

Corresponding author:

Julian Damashek

Utica College

1600 Burrstone Road, Utica, NY 13502

(315) 223-2326; judamash@utica.edu

Version 2 for biorXiv

ABSTRACT

Marine Group II *Euryarchaeota* (*Candidatus* Poseidoniales), abundant but yet-

uncultivated members of marine microbial communities, are thought to be (photo)heterotrophs

that metabolize dissolved organic matter (DOM) such as lipids and peptides. However, little is

known about their transcriptional activity. We mapped reads from a metatranscriptomic time

series collected at Sapelo Island (GA, USA) to metagenome-assembled genomes to determine

the diversity of transcriptionally-active *Ca.* Poseidoniales. Summer metatranscriptomes had the

highest abundance of *Ca.* Poseidoniales transcripts, mostly from the O1 and O3 genera within

Ca. Thalassarchaeaceae (MGIIb). In contrast, transcripts from fall and winter samples were

predominantly from *Ca.* Poseidoniaceae (MGIIa). Genes encoding proteorhodopsin, membrane-

bound pyrophosphatase, peptidase/proteases, and part of the ß-oxidation pathway were highly

transcribed across abundant genera. Highly transcribed genes specific to *Ca.* Thalassarchaeaceae

included xanthine/uracil permease and receptors for amino acid transporters. Enrichment of *Ca.*

Thalassarchaeaceae transcript reads related to protein/peptide, nucleic acid, and amino acid

transport and metabolism, as well as transcript depletion during dark incubations, provided

further evidence of heterotrophic metabolism. Quantitative PCR analysis of South Atlantic Bight

samples indicated consistently abundant *Ca.* Poseidoniales in nearshore and inshore waters.

Together, our data suggest *Ca.* Thalassarchaeaceae are important photoheterotrophs potentially

linking DOM and nitrogen cycling in coastal waters.

INTRODUCTION

Since the initial discovery of Marine Group II (MGII) *Euryarchaeota* [1,2], definitive determination of their physiology and ecological roles has remained challenging due to the lack of a cultivated isolate. Nonetheless, as data describing MGII distributions throughout the oceans have increased, several patterns have emerged: MGII are often highly abundant in the euphotic zone and in coastal waters, can reach high abundance following phytoplankton blooms, and are largely comprised of two subclades, MGIIa and MGIIb [3,4]. Early metagenomic studies provided the first evidence that MGII may be aerobic (photo)heterotrophs [5-7], a hypothesis supported by incubation experiments [8-10] and by the gene content of diverse metagenome-assembled genomes (MAGs) [11-15]. Two recent studies deepened our understanding of the phylogenomics and metabolic potential of MGII by analyzing hundreds of MAGs, highlighting clade-specific differences in genomic potential for transport and degradation of organic molecules, light harvesting proteorhodopsins, and motility [16,17]. Here, we refer to MGII as the putative order "*Candidatus* Poseidoniales," MGIIa and MGIIb as the putative families "*Ca.* Poseidoniaceae" and "*Ca.* Thalassarchaeaceae," respectively, and putative genera as specified by Rinke et al. [16]. We occasionally use "MGIIa" and "MGIIb" for consistency with previous literature.

Metatranscriptomics is one strategy for gleaning information about microbial activity in the environment. *Ca.* Poseidoniales transcripts can be abundant in marine metatranscriptomes, suggesting transiently high transcriptional activity [18,19]. When metatranscriptome reads from the Gulf of Aqaba were mapped to metagenomic contigs from the Mediterranean Sea, genes involved in amino acid transport, carbon metabolism, and cofactor synthesis were highly transcribed in the aggregate euryarchaeal community [20,21]. In another study, mapping deep-

sea metatranscriptome reads to novel *Ca.* Poseidoniales MAGs indicated transcription of genes

related to protein, fatty acid, and carbohydrate transport and metabolism, likely fueling aerobic

heterotrophy [15]. Finally, a metaproteomics study found abundant euryarchaeal transport

proteins for L-amino acids, branched-chain amino acids, and peptides throughout the Atlantic

Ocean [22]. Despite these advances, little is known about similarities or differences in gene

transcription between *Ca.* Poseidoniales and Thalassarchaeaceae.

We report MAG-resolved metatranscriptomic analyses of *Ca.* Poseidoniales in coastal

waters near Sapelo Island (GA, USA). Prior work suggested *Ca.* Poseidoniales are sporadically

active at Sapelo Island [23] and may comprise the majority of archaea in mid-shelf surface

waters of the South Atlantic Bight (SAB) [24]. Since other studies thoroughly described the

genomic content of *Ca.* Poseidoniales MAGs, our focus instead was determining which clades

were transcriptionally active and identifying highly or differentially transcribed genes. We used

two Sapelo Island MAGs [25] combined with recent *Ca.* Poseidoniales MAG collections [16,17]

to competitively recruit reads from a metatranscriptomic time series [26] and an incubation

experiment [23] to determine which clades were active over time. We then used representative

MAGs from highly active genera to determine which *Ca.* Poseidoniales genes were transcribed.

Finally, we used quantitative PCR (qPCR) to measure the abundance of *Ca.* Poseidoniales 16S

rRNA genes in DNA samples throughout the SAB to assess the prevalence *Ca.* Poseidoniales in

this region.

MATERIALS AND METHODS

PHYLOGENOMICS

Phylogenomic analyses compared SIMO Bins 19-2 and 31-1 (ref. 25) to previously-

reported *Ca.* Poseidoniales MAGs, including those binned from TARA Oceans, Mediterranean

Sea, Red Sea, Gulf of Mexico, Guaymas basin, and Puget Sound metagenomes [11,13,15-17,27-

32] (Table S1). Average nucleotide identity (ANI) was calculated using fastANI [33] to compare

non-redundant MAGs from Tully [17] to 15 Port Hacking MAGs [16] and the two SIMO MAGs;

any with ANI <98.5% were added to the non-redundant set. Phylogenomic analysis used sixteen

ribosomal proteins [34] within anvi'o v4 (ref. 35). All genomes were converted to contig

databases and proteins were identified using HMMER [36], concatenated, aligned using

MUSCLE [37], and used to build a phylogenomic tree using FastTree [38] within anvi'o.

COMPETITIVE READ MAPPING

We used competitive read mapping to determine which *Ca.* Poseidoniales genera were

transcriptionally active in Sapelo Island metatranscriptomes. Surface water for all

metatranscriptomes was collected seasonally in 2008, 2009, and 2014 from the dock at Marsh

Landing, Sapelo Island (31°25′4.08 N, 81°17′43.26 W) as previously described [23,26]. Briefly,

3-8 L of surface water was pumped through a 3-µm pore size prefilter followed by a 0.2-µm pore

size collection filter, which was frozen on liquid nitrogen. RNA from the free was extracted from

the collection filter as previously described [23,26], including the addition of internal RNA

standards to calculate volumetric transcript abundances [39]. Analyses of "field" communities

included Gifford et al. metatranscriptomes (iMicrobe Accession CAM_P_0000917) [26] and the

T$_0$ metatranscriptomes from Vorobev et al. [23], while "dark incubation" analyses included only

Vorobev et al. samples (T$_0$, which were processed immediately after collection, and T$_{24}$, which

were processed following 24 hours of *in situ* incubation in dark containers; NCBI BioProject

PRJNA419903). Temperature, salinity, dissolved oxygen, pH, and turbidity data corresponding

to metatranscriptome sampling times were downloaded from the NOAA National Estuarine

Research Reserve System website (http://cdmo.baruch.sc.edu; last accessed 16 July 2020).

Contigs from all MAGs used for phylogenomics (Table S1) were used as a read mapping

database using Bowtie2 v.2.2.9 (ref. 40) with the "very-sensitive" flag. Samtools v.1.3.1 (ref. 41)

was used to index BAM files, which were profiled and summarized in anvi'o. Contig genus

identity was imported to the anvi'o contig database as an external collection. The number of

transcripts L^{-1} was calculated by scaling the number of mapped reads by the volume of water

filtered and the recovery of internal standards (reported in [23,26]) as previously described [39].

Seasonal transcript abundances were compared using a one-way ANOVA test in R [42] with data

log-transformed as necessary to improve normality. When ANOVA results were significant,

groupings were defined post-hoc with Tukey's Honest Significant Difference (HSD) test using

the agricolae R package [43].

Non-metric multidimensional scaling (NMDS) analysis of metatranscriptome hits was

conducted using the vegan R package [44]. NMDS input was a distance matrix constructed by

Hellinger-transforming the table of transcript hits and calculating Euclidean distance between

samples [45]. Genus vectors were calculated using the *envfit* command. Significance of

groupings were tested by permutational multivariate analysis of variance (*adonis* command) with

999 permutations.

MAG-SPECIFIC ANNOTATION AND TRANSCRIPT ANALYSIS

Gene-specific analyses focused on three MAGs: two from the SIMO collection (SIMO

Bin 19-2, Genbank: VMDE00000000; SIMO Bin 31-1, VMBU00000000; [25]) and one

previously binned from Red Sea metagenomes (RS440, PBUZ00000000; [27]). These MAGs

represented genera O1, O3, and M, respectively, which were highly abundant in

metatranscriptomes (see Fig. 1). RS440 was selected due to a high number of transcripts

recruited when genus M was abundant (data not shown).

MAGs were annotated using the archaeal database in Prokka v.1.13 (ref. 46), using

DIAMOND [47] to search against all orthologous groups in eggNOG-mapper v.1 (refs. 45,46),

and using the BlastKOALA portal (https://www.kegg.jp/blastkoala/, last accessed 6 March 2019)

[48]. Putative genes for carbohydrate-active enzymes, peptidases, and membrane transport

proteins were identified using HMMER searches of dbCAN2 (HMMdb v.7) [49,50], MEROPS

v.12.0 (ref. 51), and the Transporter Classification Database [52], respectively.

Transcript reads were mapped to MAGs (combined into a single database such that each

read mapped to only one MAG) to identify *Ca.* Poseidoniales genes that were highly or

differentially transcribed. Coverage was calculated by profiling BAM files in anvi'o and

normalized to coverage per million reads (CPM) by dividing by the total number of reads per

sample. For each MAG, "highly transcribed" genes were the 5% of putative genes with the

highest median CPM across metatranscriptomes (SIMO Bin 19-2: 63 genes, SIMO Bin 31-1: 70

genes, RS440: 77 genes).

DESeq2 v.3.11 (ref. 53) was used to identify genes from each MAG that were

differentially transcribed when each genus was highly transcriptionally active. For each MAG,

"treatment" samples in DESeq2 were those where the respective genus recruited ≥50% of *Ca.*

Poseidoniales reads from the metatranscriptome. Thus, positive fold-change values are genes

transcribed at higher levels when the genus is highly transcriptionally active (compared to other

metatranscriptomes). DESeq2 was also used to identify differentially transcribed genes for each

MAG between T_0 and T_{24} samples in high tide (HT) dark incubations [23]. Since T_{24} samples

were the "treatment" condition in DESeq2, positive fold-change values here are genes

transcribed at higher levels in T_{24} compared to T_0 samples. In all DESeq2 analyses, genes with

Benjamini-Hochberg adjusted $p<0.1$ were counted as having significantly different transcription.

QUANTITATIVE PCR

DNA samples from SAB field campaigns in 2014 and 2017 [24,54] were used as

templates for qPCR reactions targeting the *Ca.* Poseidoniales 16S rRNA gene. Samples included

the variety of shelf habitats (inshore, nearshore, mid-shelf, shelf-break, and oceanic as previously

defined [24]; Fig. S1) and multiple depths when possible. DNA from the entire microbial

community (no prefiltration) was extracted as previously described [54]. Primers were GII-554-f

[55] and Eury806-r [56] with cycling conditions as previously reported [57] (Table S2).

Reactions (25 µL, triplicate) used iTaq Universal Green SYBR Mix (Bio-Rad, Hercules, CA) in

a C1000 Touch Thermal Cycler/CFX96 Real-Time System (Bio-Rad, Hercules, CA). Each plate

included a no-template control and a standard curve (serial dilutions of a linearized plasmid

containing a previously-sequenced, cloned amplicon). Abundance of *Ca.* Poseidoniales 16S

rRNA genes was compared to published bacterial 16S rRNA gene abundance from the same

samples [24,54]. Regional variability of gene abundance was assessed using a one-way ANOVA

and a post-hoc HSD test as described above. Model II regressions of log-transformed qPCR data

were estimated using the lmodel2 R package [58] as previously described [54]. All plots were

constructed with anvi'o or the ggplot2 R package [59].

RESULTS

EURYARCHAEOTAL MAGs

SIMO Bins 19-2 and 31-1 were estimated as 82.5-92.3% and 77.5-96.2% complete,

respectively, with redundancy <0.6% [25]. Phylogenomics placed both in the putative family *Ca.*

Thalassarchaeaceae (MGIIb) and genera O1 (SIMO Bin 19-2) and O3 (SIMO Bin 31-1; Fig. S2).

Phylogenomic groupings were generally consistent with previous findings [16,17].

Both SIMO MAGs contained a proteorhodopsin gene. Presence of a methionine residue

at position 315 suggested absorption of green light [60], and both proteorhodopsin genes grouped

in "Archaea Clade B" [11,17,61] (Fig. S3). Both MAGs included partial or complete pathways

indicating aerobic heterotrophic growth, such as glycolysis, the TCA cycle, and electron

transport chain components (Table S3). Pathways for metabolism of compounds such as fatty

acids, peptides, and proteins were also present, as were transport systems and metabolic

pathways for amino acids and nucleotides.

DOMINANT GENERA IN FIELD METATRANSCRIPTOMES

There were significant seasonal differences in transcript recruitment by the combined set

of *Ca.* Poseidoniales MAGs ($F_{3,28}$=4.9, p=0.007): most summer samples had >10^{10} *Ca.*

Poseidoniales transcripts L^{-1}, significantly more than in other seasons (Fig. 1A,C; Table S4). The

diversity of transcriptionally-active *Ca.* Poseidoniales also changed seasonally. Genera O1 and

O3 accounted for most reads mapped from summer samples (typically 89.5-99.5% of *Ca.*

Poseidoniales reads), with most mapping to O1 (Fig. 1A,B; Fig. S4; Table S4). HT (and not low

tide; LT) metatranscriptomes from July 2014 also had a moderate fraction of reads (37.5-39.6%)

mapped to *Ca.* Poseidoniaceae. In contrast to summer samples, November 2008 and May 2009

transcripts were predominantly O3, while those from February 2009 and October 2014 were

mostly *Ca.* Poseidoniaceae genera M, L1, or L2 (Fig. 1A,B, Table S4).

HIGHLY TRANSCRIBED *CA.* POSEIDONIALES GENES

Many highly transcribed *Ca.* Thalassarchaeaceae (MGIIb) genes were involved in

translation, transcription, replication/repair, or post-translation protein modification (Fig. 2), as

expected by their fundamental role in cellular processes. Genes encoding proteins involved in

energy production or conservation (ATPases, a pyrophosphate-energized proton pump, and

proteorhodopsin) were also highly transcribed. Notably, the *aapJ* and *livK* genes, encoding

ligand-binding receptors for L-amino acid and branch-chain amino acid transporters,

respectively, were among the most highly transcribed genes in both *Ca.* Thalassarchaeaceae

MAGs (Fig. 2, Table S3).

Many of the highly transcribed genes mapping to the *Ca.* Poseidoniaceae (MGIIa) MAG

were not highly transcribed by *Ca.* Thalassarchaeaceae, including genes encoding a carbamoyl

phosphate synthetase subunit (*carA*), a family 2 glycosyl transferase, chromosomal protein

MC1b, and a ftsX-like permease. The *carA* gene had the highest median coverage of *Ca.*

Poseidoniaceae genes across coastal metatranscriptomes (Fig. 2, Table S3). While both *Ca.*

Thalassarchaeaceae MAGs also contained the carbamoyl phosphate synthetase genes, neither

transcribed *carA* at high levels (Table S3).

Twelve genes were highly transcribed by *Ca.* Thalassarchaeaceae and not by *Ca.*

Poseidoniaceae, including genes encoding ATP synthase, transcription initiation factor IIB,

halocyanin, phytoene desaturase, protein translocase, xanthine/uracil permease, and receptors for

amino acid transporters (Fig. 2, Table S3). Other than those encoding ribosomal proteins, only

six genes were highly transcribed in all three MAGs: a chaperone protein, a ribonucleoside-

diphosphate reductase, translation elongation factor 1A, 3-hydroxyacyl-CoA dehydrogenase, a

membrane-bound pyrophosphatase (*hppA*), and proteorhodopsin (Fig. 2).

DIFFERENTIAL GENE TRANSCRIPTION

We were interested in identifying genes with variable transcription levels when genera

O1, O3, and M were highly transcriptionally active in the ocean. Twenty-three genes were

differentially transcribed in O1-active samples (Fig. 3, Table S5), sixteen of which had higher

abundance in O1-active metatranscriptomes compared to other metatranscriptomes. These highly

transcribed genes encoded proteorhodopsin, two copper-containing redox proteins (halocyanin

and plastocyanin), and proteins involved in lipid metabolism (3-hydroxyacyl-CoA

dehydrogenase and oligosaccharyltransferase), nucleotide transport/metabolism (ribonucleotide-

diphosphate reductase and xanthine/uracil permease), and amino acid transport (ligand-binding

receptor for a L-amino acid transporter, *aapJ*). Differentially transcribed genes mapping to the

O3 MAG were mostly depleted in O3-active metatranscriptomes and largely encoded proteins of

unknown function; only the gene encoding ribosomal protein L12 was enriched in O3-active

samples (Fig. S5, Table S5). Only four genes mapping to the M MAG were differentially

transcribed in M-active metatranscriptomes. Annotated genes encoded chromosomal protein

MC1b, an ATPase subunit, and a glycosyl transferase, which all had significantly fewer

transcripts in M-rich samples (Fig. S6, Table S5). One gene of unknown function was enriched

compared to other samples.

DARK INCUBATION METATRANSCRIPTOMES

248 Incubation had little effect on transcription by the dominant genera in LT samples (Fig. 4,

249 Table S4). In contrast, there were distinct shifts in transcriptionally active populations during

250 incubations of all HT samples. July HT metatranscriptomes initially contained 60.4-62.5% *Ca.*

251 Thalassarchaeaceae (MGIIb) while hits from the corresponding T_{24} samples were 98.1-99.3%

252 *Ca.* Thalassarchaeaceae, due to increased transcript hits to genus O1 (Fig. 4, Table S4).

253 Likewise, October 2014 HT samples initially contained 65.0-66.6% hits to *Ca.* Poseidoniaceae

254 (MGIIa) but changed to 78.3-98.8% hits to *Ca.* Thalassarchaeaceae at T_{24} due to an increase in

255 hits to O1 (Fig. 4).

256 DESeq2 identified 40 differentially transcribed genes mapping to the O1 MAG between

257 HT T_0 and T_{24} metatranscriptomes. Four O1 genes had higher transcription at T_{24}, including

258 xanthine/uracil permease (*pbuG*) and a ligand-binding receptor for a general amino acid

259 transporter (*aapJ*; Fig. 5, Table S6). The 36 O1 genes transcribed at lower levels encoded

260 proteins involved in repair of UV-damaged DNA, amino acid or nucleotide metabolism,

261 coenzyme synthesis, peptidases or proteases, transcription, DNA replication, and lipid

262 biosynthesis, as well as phytoene desaturase (*crtD*) and multiple subunits of pyruvate

263 dehydrogenase (*pdhC*, *pdhA*). None of the genes mapping to the O3 or M MAGs were

264 transcribed at significantly different levels between HT incubation timepoints ($p>0.1$ for all

265 genes; Table S6).

267 16S rRNA GENE ABUNDANCE

268 *Ca.* Poseidoniales 16S rRNA genes were detected in all SAB DNA samples (*n*=208),

269 with a range from 1.6×10^4 to 7.6×10^8 genes L^{-1} (Table S7). Standard curves for the *Ca.*

270 Poseidoniales assay always had $r^2>0.99$ (mean ± standard deviation: 0.99±0.001) and the mean

efficiency was 93.4% (±2.0%; Table S2). When data from all cruises were combined, *Ca.*

Poseidoniales genes were most abundant throughout inshore and nearshore waters and least

abundant in shelf-break and oceanic samples ($F_{4,204}$=18.5, $p<0.001$; Fig. 6A). There was a strong

linear relationship between log-transformed bacterial and *Ca.* Poseidoniales 16S rRNA gene

abundances, with bacterial abundances 2-3 orders of magnitude higher (Fig. 6B).

DISCUSSION

ABUNDANCE OF *CA*. POSEIDONIALES GENES IN THE SOUTH ATLANTIC BIGHT

Given typical *Ca.* Poseidoniales abundance of 10^6-10^7 genes or cells L^{-1} in oligotrophic

waters [62-64] and 10^7-10^8 genes or cells L^{-1} in coastal waters [9,55,57,65-67], the abundance of

their 16S rRNA genes in the coastal SAB (nearly 10^9 genes L^{-1} in some samples) is among the

highest measured in the ocean. Greater gene abundance in inshore, nearshore, and mid-shelf

waters indicates *Ca.* Poseidoniales are more abundant over the shallow shelf than further

offshore (Fig. 6A), matching clone library data from the SAB [24] and data from the Central

California Current and the Black Sea [9,66]. The DOM in productive, turbid SAB coastal waters

supports highly active heterotrophic microbial populations [68,69]. Our data suggest large

populations of *Ca.* Poseidoniales are part of this heterotrophic community, and the correlation

between *Ca.* Poseidoniales and bacterial abundance (Fig. 6B) suggests common factors influence

the abundance of these two communities.

DIVERSITY OF TRANSCRIPTIONALLY ACTIVE *CA*. POSEIDONIALES

While numerous studies have demonstrated high abundance of *Ca.* Poseidoniales in the

coastal ocean (e.g., [57,67]), little is known about which clades are transcriptionally active in

these regions. At Sapelo Island, the striking dominance of the *Ca.* Thalassarchaeaceae (MGIIb)

genera O1 and O3 in summer samples, which also contained the highest levels of aggregate *Ca.*

Poseidoniales transcripts (Fig. 1), indicates that *Ca.* Thalassarchaeaceae are most active during

the summertime. Outliers to this pattern were July 2014 HT samples, which contained abundant

Ca. Poseidoniaceae (MGIIa) transcripts (though *Ca.* Thalassarchaeaceae still comprised the

majority of their *Ca.* Poseidoniales reads). A previous study found relatively high salinity and

DOM enriched in marine-origin molecules over the mid-shelf SAB during July 2014 [70]. Our

data suggest *Ca.* Poseidoniaceae were relatively active over the shelf during this time and were

transported inshore during flood tides, leading to shifts in transcriptional diversity between LT

waters (dominated by *Ca.* Thalassarchaeaceae) and HT waters (which included a higher number

of *Ca*. Poseidoniaceae).

Ca. Poseidoniaceae (MGIIa) are the predominant euryarchaeal family in many coastal

ecosystems, particularly in summer (e.g., [9,14,57,71,72]), but it is unclear what general patterns

govern *Ca.* Poseidoniales distributions in coastal waters worldwide. Studies of *Ca.* Poseidoniales

ecology often focus on distributions with depth, typically finding abundant *Ca.*

Thalassarchaeaceae (MGIIb) in deeper waters and *Ca.* Poseidoniaceae (MGIIa) more prevalent

in euphotic waters (e.g., [12,55,73-75]). A recent mapping of global ocean metagenome reads

showed that coastal populations of *Ca.* Poseidoniales were primarily *Ca.* Poseidoniaceae, though

Ca. Thalassarchaeaceae MAGs recruited a substantial number of reads from some coastal

metagenomes [17]. Our data match this latter pattern, with *Ca.* Poseidoniales populations in

surface waters off Sapelo Island dominated by highly active *Ca.* Thalassarchaeaceae (Fig. 1).

The higher abundance of MAGs from genera O1 and O3 (also referred to as MGIIb.12 and

MGIIb.14) in some mesopelagic and coastal samples with relatively high temperature (~23−

30°C; [17]) may explain the unusual pattern found in SAB waters: these genera peaked in

summer at Sapelo Island, when water temperatures were 29–30°C (Table S4), suggesting they

may be adapted to growth at relatively low light and high temperature.

CA. POSEIDONIALES GENE TRANSCRIPTION PATTERNS

Sapelo Island metatranscriptome reads that mapped to *Ca.* Poseidoniales were analyzed

in three ways:

(1) We determined sets of "highly transcribed" genes mapping to MAGs of

transcriptionally-active *Ca* Poseidoniales genera (5% of MAG genes with the highest

median transcript coverage);

(2) We identified genes mapping to *Ca* Poseidoniales MAGs that were differentially

transcribed when its genus was highly active (≥50% of *Ca.* Poseidoniales transcripts

in a sample; Fig. 1, Table S4);

(3) We identified genes mapping to *Ca* Poseidoniales MAGs that were differentially

transcribed at the beginning versus end of dark incubations. Since dark incubation

separates indigenous microbes from light and sources of short-lived substrates,

transcription of corresponding transporters and metabolic genes ceases during

incubations as substrates are consumed. We therefore assume transcript depletion in

T_{24} compared to T_0 metatranscriptomes indicates genes that were transcriptionally

active in the field [23]. This interpretation was bolstered by significant transcript

depletions for genes involved in repairing UV damage to DNA (*uvrA*, *uvrC*, and

cofH; Fig. 5), an expected result given alleviation of UV stress in dark incubations.

In the following sections, we synthesize these analyses to discuss *Ca.* Poseidoniales

transcriptional activity related to DOM metabolism, transport/metabolism of amino acids and

nucleotides, and basic energetic processes. Though lack of a cultivated representative limits the

analysis to computationally-inferred functions, these data provide hypotheses regarding the

activity of *Ca.* Poseidoniales families in the coastal ocean.

PROTON GRADIENTS AND ELECTRON TRANSPORT

Our analysis revealed that *Ca* Poseidoniales genes involved in establishing

transmembrane proton gradients were highly transcribed in our samples. Genes encoding

proteorhodopsin were among the most highly transcribed by both *Ca.* Thalassarchaeaceae MAGs

and were highly transcribed in O1-active samples (Figs. 2,3). Proteorhodopsins consist of a

retinal chromophore linked to a transmembrane protein and use light energy to pump protons

across the cell membrane [76]. The resulting energy can be coupled to ATP production or other

chemiosmotic processes and often supports photoheterotrophy, though its function varies widely

[61]. Proteorhodopsin genes are highly transcribed in the photic zone of both open ocean and

coastal waters (e.g., [77-79]) and our data indicate coastal *Ca.* Poseidoniales conform to this

pattern, consistent with recent evidence from other regions [21,80,81]. High transcription of

proteorhodopsin supports the photoheterotrophic lifestyle hypothesized for *Ca.* Poseidoniales

(e.g., [9,11,12,16,17]).

Since O1 proteorhodopsin transcript abundance did not differ between the beginning and

end of dark incubations (Fig. 5), light may not regulate *Ca.* Thalassarchaeaceae proteorhodopsin

transcription. However, depletion of O1 *crtD* (carotenoid 3,4-desaturase) transcripts during dark

incubation (Fig. 5) suggests light may regulate retinal synthesis. Whether proteorhodopsin

transcription responds to light varies among marine bacteria (e.g., [82,83]), and the function of

constitutive transcription is not straightforward: while some bacteria use proteorhodopsin to

produce ATP when carbon-limited [84], high amounts of proteorhodopsin in other bacteria can

physically stabilize membranes even when inactive [85]. The high proteorhodopsin transcription

in our data emphasizes, but provides little mechanistic clarification of, the physiological role for

proteorhodopsin in *Ca.* Poseidoniales (Table 1).

Numerous genes encoding putative electron transport proteins were highly transcribed by

at least one *Ca.* Poseidoniales family (Fig. 2). Like proteorhodopsin, a halocyanin gene was

among the most highly transcribed *Ca.* Thalassarchaeaceae genes and was enriched when genus

O1 was transcriptionally active (Fig. 3). Halocyanins are involved in the electron transport chain

and have been posited to increase the energy yield of aerobic respiration in *Ca.*

Thalassarchaeaceae to stimulate rapid growth [12]. The similar transcription patterns of

proteorhodopsin and halocyanin suggests proteorhodopsin activity in coastal *Ca.*

Thalassarchaeaceae may function to increase growth rates during respiration, aiding rapid

population growth when conditions permit.

The *hppA* gene from all *Ca* Poseidoniales MAGs was highly transcribed (Fig. 2; Table 1).

This gene (which is found across many domains of life [86], including a diverse array of marine

microbes [87]) putatively encodes a membrane-bound pyrophosphatase, which generates a

proton gradient via hydrolysis of pyrophosphate, a byproduct of numerous cellular processes

[86]. In metatranscriptomes from a phytoplankton bloom, enrichment of *hppA* transcripts

suggested high pyrophosphate-based energy conservation in oligotrophic waters [87], and *hppA*

transcripts were similarly abundant at night in an oligotrophic lake [88]. Although widespread in

MAGs from *Ca.* Poseidoniales [16], *hppA* has not been previously recognized as a potentially

important part of their metabolism. Our data suggest *Ca.* Poseidoniales may be capable of using

pyrophosphatase (along with proteorhodopsin) to generate a protonmotive force (Table 1).

POTENTIAL IMPORTANCE OF MARINE DOM IN *CA.* POSEIDONIALES METABOLISM

Although the T_0 metatranscriptomes from summer versus fall were dominated by

transcripts from different *Ca.* Poseidoniales families, a 24-hour dark incubation consistently

favored transcription by *Ca.* Thalassarchaeaceae when samples were collected at HT (Fig. 4).

This tidal stage-linked increase in *Ca.* Thalassarchaeaceae transcription could relate to

differences in DOM availability between HT and LT, consistent with numerous studies

implicating DOM in shaping *Ca.* Poseidoniales populations [9,10,89]. The HT Sapelo Island

DOM pool is primarily of marine origin while LT DOM is more riverine- and marsh-derived

[70], which may select for growth of different *Ca.* Poseidoniales families in the water masses

present at different tidal stages. Furthermore, the depletion of transcripts encoding two pyruvate

dehydrogenase subunits (*pdhA* and *pdhC*) during incubations (Fig. 5) suggests *Ca.* Poseidoniales

were metabolizing phytoplankton photosynthate *in situ*. Alternatively, these tidal stage-driven

transcriptional patterns may relate to differential light adaptation in populations originating in

offshore versus nearshore waters. Inshore populations, potentially adapted to life in turbid

waters, may increase transcription upon dark enclosure, whereas offshore populations

(transported shoreward during flood tide) may be adapted to clearer waters and reduce

transcription in dark conditions.

Multiple lines of evidence indicate coastal *Ca.* Poseidoniales may have been

metabolizing proteins and fatty acids. High transcription of genes encoding proteases or

peptidases from all *Ca.* Poseidoniales MAGs (Fig. 2) suggests likely metabolism of proteins or

peptides by both families (Table 1). Furthermore, decreased transcription of protease genes

mapping to the O1 MAG during dark incubations (Fig. 5) suggests protein metabolism by *Ca.*

Thalassarchaeaceae may have been active *in situ* prior to incubation. While some of these genes

could be involved in intracellular recycling (particularly lon protease and cytosol

aminopeptidase), active protein metabolism is consistent with previous experiments

demonstrating protein assimilation [10] and high transcription of *Ca.* Poseidoniales peptidase

genes in other marine regions [15,21]. In addition to genes encoding protein catabolism, high

transcription of the 3-hydroxyacyl-CoA dehydrogenase gene from all three MAGs (Fig. 2), and

its enrichment in O1-active field samples (Fig. 3), suggests a likely importance of fatty acid

metabolism for both *Ca.* Poseidoniales families (Table 1), matching widespread ß-oxidation

genes in *Ca.* Poseidoniales MAGs [16,17] and transcriptional data from the deep ocean [15].

DISTINCT PATTERNS OF AMINO ACID AND NUCLEOTIDE UPTAKE AND

METABOLISM

Transcription of *livK* and *aapJ* appears to differentiate *Ca.* Poseidoniales families in the

coastal ocean (Table 1). These genes putatively encode ligand-binding receptors for ABC

transporters: *aapJ* for a general L-amino acid transporter and *livK* for a branched-chain amino

acid transporter [90]. Both are commonly present in *Ca.* Thalassarchaeaceae (MGIIb) but not *Ca.*

Poseidoniaceae (MGIIa) [17] and were among the most highly transcribed *Ca.*

Thalassarchaeaceae genes (Fig. 2). Previous studies noted high transcription of euryarchaeal *livK*

and *aapJ* genes in the water column of the Red Sea [20,21], at the Mid-Cayman Rise [15], and

throughout the Atlantic Ocean [22]. Our data suggest this activity was probably associated with

Ca. Thalassarchaeaceae.

The *aapJ* and *livK* genes were collocated with genes putatively encoding the full

transporters in the *Ca.* Thalassarchaeaceae MAGs (Table S3). Unfortunately, it is difficult to

guess their substrates from sequence data alone: AAP transporters are typically capable of

transporting a range of L-amino acids [90] while LIV transporters can be highly specific for

leucine, specific for branched-chain amino acids, or transport diverse amino acids [90-92]. In soil

bacteria grown under inorganic nitrogen limitation, elevated transcription of *aapJ* is linked to

organic nitrogen use [93], but it is unclear whether this mechanism translates to *Ca.*

Thalassarchaeaceae since amino acids could be used for numerous anabolic or catabolic

processes. In addition to these binding proteins, the depletion of transcripts from O1 genes

putatively involved in the shikimate pathway of aromatic amino acid synthesis (3-

phosphoshikimate 1-carboxyvinyltransferase and anthranilate synthase) during incubations (Fig.

5) suggests *Ca.* Thalassarchaeaceae in this genus may have been synthesizing aromatic amino

acids *in situ*.

The combination of high *pbuG* transcription by *Ca.* Thalassarchaeaceae (MGIIb) with the

high numbers of O1 *pbuG* transcripts in O1-active samples and dark incubation endpoints (Figs.

2,3,5) suggests an important role for xanthine/uracil permease (the putative product of *pbuG*) in

Ca. Thalassarchaeaceae metabolism. In some phytoplankton, *pbuG* is transcribed during

nitrogen-stressed growth [94-96], potentially allowing access to DON. However, *pbuG* and

xanthine dehydrogenase (*xdh*) are also transcribed when xanthine is catabolized by marine

bacteria [97]. Both *Ca.* Thalassarchaeaceae MAGs contain putative xanthine dehydrogenase

genes (*xdhC* and *yagS*; Table S3), suggesting the ability to catabolize xanthine (Table 1).

Transcription levels of *carA*, putatively encoding part of carbamoyl phosphate synthetase,

appears to be a distinct trait of *Ca.* Poseidoniaceae (MGIIa): while all three MAGs contained this

gene (Table S3), only *Ca.* Poseidoniaceae *carA* transcription was high. Since carbamoyl

phosphate synthetase is a key enzyme for arginine and pyrimidine synthesis from bicarbonate

[98], high *carA* transcription suggests these pathways may be important for *Ca.* Poseidoniaceae

growth or survival, though it is not clear whether this transcription is related to synthesis of

amino acids, nucleotides, or both.

CONCLUSIONS

Our metatranscriptomic data and associated experiments provide a novel window into the

activity of *Ca.* Poseidoniales families (formerly "MGIIa" and "MGIIb"). They indicate an

important role for *Ca.* Thalassarchaeaceae (MGIIb) as coastal photoheterotrophs, particularly in

warm waters. High transcription of proteorhodopsin and membrane-bound pyrophosphatase

genes suggested common methods for establishing proton gradients. Furthermore, high

transcription of genes involved in protein/peptide metabolism and ß-oxidation of fatty acids

confirmed peptide and lipid metabolism as a common trait. However, high transcription of *Ca.*

Thalassarchaeaceae genes encoding amino acid binding proteins and nucleotide transporters

suggests uptake of these substrates may distinguish the two families. These data confirm the

importance of DOM metabolism by *Ca.* Poseidoniales and suggest a potential role for organic

nitrogen in *Ca.* Thalassarchaeaceae metabolism.

ACKNOWLEDGMENTS

We thank Qian Liu, Bradley Tolar, the Georgia Coastal Ecosystems LTER field team, the

University of Georgia Marine Institute at Sapelo Island staff, and the captain and crew of the

R/V *Savannah* for assistance with original field sampling. This work was funded in part by NSF

OCE grant 1538677 and OPP grant 1643466 (to JTH), NSF Grant IOS1656311 (to MAM), NSF

OCE grant 1832178 (to the Georgia Coastal Ecosystems LTER), as well as by resources and

expertise from the Georgia Advanced Computing Resource Center, a partnership between the

University of Georgia's Office of the Vice President for Research and Office of the Vice

President for Information Technology.

CONFLICT OF INTEREST

The authors declare no competing financial interests.

REFERENCES
1.   Fuhrman JA, McCallum K, Davis AA. Novel major archaebacterial group from marine plankton. Nature
1992;356:148–149.

2.   DeLong EF. Archaea in coastal marine environments. Proceedings of the National Academy of Sciences
1992;89:5685–5689.

3.   Zhang CL, Xie W, Martín-Cuadrado A-B, Rodriguez-Valera F. Marine Group II Archaea, potentially
important players in the global ocean carbon cycle. Frontiers in Microbiology 2015;6:1108.

4.   Santoro AE, Richter RA, Dupont CL. Planktonic marine archaea. Annual Review of Marine Science
2019;11:131–158.

5.   Béjà O, Suzuki MT, Koonin EV, Aravind L, Hadd A, Nguyen LP, et al. Construction and analysis of
bacterial artificial chromosome libraries from a marine microbial assemblage. Environmental
Microbiology 2000;2:516–529.

6.   Moreira D, Rodriguez-Valera F, López-García P. Analysis of a genome fragment of a deep-sea
uncultivated Group II euryarchaeote containing 16S rDNA, a spectinomycin-like operon and several
energy metabolism genes. Environmental Microbiology 2004;6:959–969.

7.   Frigaard N-U, Martinez A, Mincer TJ, DeLong EF. Proteorhodopsin lateral gene transfer between marine
planktonic Bacteria and Archaea. Nature 2006;439:847–850.

8.   Alonso C, Pernthaler J. Concentration-dependent patterns of leucine incorporation by coastal
picoplankton. Applied and Environmental Microbiology 2006;72:2141–2147.

9.   Orsi WD, Smith JM, Wilcox HM, Swalwell JE, Carini P, Worden AZ, et al. Ecophysiology of
uncultivated marine euryarchaea is linked to particulate organic matter. The ISME Journal 2015;9:1747–
1763.

10.   Orsi WD, Smith JM, Liu S, Liu Z, Sakamoto CM, Wilken S, et al. Diverse, uncultivated bacteria and
archaea underlying the cycling of dissolved protein in the ocean. The ISME Journal 2016;10:2158–2173.

11.   Iverson V, Morris RM, Frazar CD, Berthiaume CT, Morales RL, Armbrust EV. Untangling genomes from
metagenomes: revealing an uncultured class of marine *Euryarchaeota*. Science 2012;335:587–590.

12.   Martín-Cuadrado A-B, Garcia-Heredia I, Gonzaga Moltó A, López-Ubeda R, Kimes N, López-Garcia P,
et al. A new class of marine Euryarchaeota group II from the mediterranean deep chlorophyll maximum.
The ISME Journal 2015;9:1619–1634.

13.   Thrash JC, Seitz KW, Baker BJ, Temperton B, Gillies LE, Rabalais NN, et al. Metabolic roles of
uncultivated bacterioplankton lineages in the northern Gulf of Mexico "dead zone." mBio 2017;8:e01017–
7.

14.   Orellana LH, Francis TB, Krüger K, Teeling H, Müller M-C, Fuchs BM, et al. Niche differentiation
among annually recurrent coastal Marine Group II Euryarchaeota. The ISME Journal 2019;13:3024–3036.

15.   Li M, Baker BJ, Anantharaman K, Jain S, Breier JA, Dick GJ. Genomic and transcriptomic evidence for
scavenging of diverse organic compounds by widespread deep-sea archaea. Nature Communications
2015;6:8933.

16.   Rinke C, Rubino F, Messer LF, Youssef N, Parks DH, Chuvochina M, et al. A phylogenomic and
ecological analysis of the globally abundant Marine Group II archaea (*Ca.* Poseidoniales ord. nov.). The
ISME Journal 2019;13:663–675.

17.   Tully BJ. Metabolic diversity within the globally abundant Marine Group II Euryarchaea offers insight
into ecological patterns. Nature Communications 2019;10:271.

18.   Baker BJ, Sheik CS, Taylor CA, Jain S, Bhasi A, Cavalcoli JD, et al. Community transcriptomic assembly
reveals microbes that contribute to deep-sea carbon and nitrogen cycling. The ISME Journal 2013;7:1962–
1973.

19.   Ottesen EA, Young CR, Gifford SM, Eppley JM, Marin R, Schuster SC, et al. Multispecies diel
transcriptional oscillations in open ocean heterotrophic bacterial assemblages. Science 2014;345:207–212.

20.   Miller D, Pfreundt U, Hou S, Lott SC, Hess WR, Berman-Frank I. Winter mixing impacts gene expression
in marine microbial populations in the Gulf of Aqaba. Aquatic Microbial Ecology 2017;80:223–242.

21.   Hou S, Pfreundt U, Miller D, Berman-Frank I, Hess WR. mdRNA-Seq analysis of marine microbial
communities from the northern Red Sea. Scientific Reports 2016;6:35470.

22.   Bergauer K, Fernandez-Guerra A, Garcia JAL, Sprenger RR, Stepanauskas R, Pachiadaki MG, et al.
Organic matter processing by microbial communities throughout the Atlantic water column as revealed by
metaproteomics. Proceedings of the National Academy of Sciences 2018;115:E400–E408.

23.   Vorobev A, Sharma S, Yu M, Lee J, Washington BJ, Whitman WB, et al. Identifying labile DOM
components in a coastal ocean through depleted bacterial transcripts and chemical signals. Environmental
Microbiology 2018;20:3012–3030.

24.   Liu Q, Tolar BB, Ross MJ, Cheek JB, Sweeney CM, Wallsgrove NJ, et al. Light and temperature control
the seasonal distribution of thaumarchaeota in the South Atlantic bight. The ISME Journal 2018;12:1473–
1485.

25.   Damashek J, Edwardson CF, Tolar BB, Gifford SM, Moran MA, Hollibaugh JT. Coastal ocean
metagenomes and curated metagenome-assembled genomes from Marsh Landing, Sapelo Island (Georgia,
USA). Microbiology Resource Announcements 2019;8:e00934–19.

26.   Gifford SM, Sharma S, Moran MA. Linking activity and function to ecosystem dynamics in a coastal
bacterioplankton community. Frontiers in Microbiology 2014;5:185.

27.   Tully BJ, Graham ED, Heidelberg JF. The reconstruction of 2,631 draft metagenome-assembled genomes
from the global oceans. Scientific Data 2018;5:170103.

28.   Tully BJ, Sachdeva R, Graham ED, Heidelberg JF. 290 metagenome-assembled genomes from the
Mediterranean Sea: a resource for marine microbiology. PeerJ 2017;5:e3558.

29.   Delmont TO, Quince C, Shaiber A, Esen ÖC, Lee ST, Rappé MS, et al. Nitrogen-fixing populations of
Planctomycetes and Proteobacteria are abundant in surface ocean metagenomes. Nature Microbiology
2018;3:804–813.

30.   Haroon MF, Thompson LR, Parks DH, Hugenholtz P, Stingl U. A catalogue of 136 microbial draft
genomes from Red Sea metagenomes. Scientific Data 2016;3:160050.

31.   Parks DH, Rinke C, Chuvochina M, Chaumeil P-A, Ben J Woodcroft, Evans PN, et al. Recovery of nearly
8,000 metagenome-assembled genomes substantially expands the tree of life. Nature Microbiology
2017;2:1533–1542.

32. Haro-Moreno JM, López Pérez M, la Torre de JR, Picazo A, Camacho A, Rodriguez-Valera F. Fine
metagenomic profile of the Mediterranean stratified and mixed water columns revealed by assembly and
recruitment. Microbiome 2018;6:128.

33. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI analysis of
90K prokaryotic genomes reveals clear species boundaries. Nature Communications 2018;9:5114.

34. Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, et al. A new view of the tree of
life. Nature Microbiology 2016;1:16048.

35. Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, et al. Anvi'o: an advanced analysis
and visualization platform for 'omics data. PeerJ 2015;3:e1319.

36. Eddy SR. Accelerated profile HMM searches. PLoS Computational Biology 2011;7:e1002195.

37. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic
Acids Research 2004;32:1792–1797.

38. Price MN, Dehal PS, Arkin AP. FastTree 2 – approximately maximum-likelihood trees for large
alignments. PLoS ONE 2010;5:e9490.

39. Satinsky BM, Gifford SM, Crump BC, Moran MA. Use of Internal Standards for Quantitative
Metatranscriptome and Metagenome Analysis. In: DeLong EF, editor. *Methods in Enzymology Microbial
Metagenomics, Metatranscriptomics, and Metaproteomics*. Elsevier, Inc.; 2013. pages 237–250.

40. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature Methods 2012;9:357–359.

41. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format
and SAMtools. Bioinformatics 2009;25:2078–2079.

42. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria. https://www.R-project.org/. 2019.

43. de Mendiburu F. agricolae: Statistical procedures for agricultural research. R package version 1.3-1.
https://CRAN.R-project.org/package=agricolae. 2019.

44. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan: Community ecology
package. R package version 2.5-5. https://CRAN.R-project.org/package=vegan. 2019.

45. Legendre P, Legendre L. Numerical Ecology. 3rd ed. Elsevier; 2012.

46. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics 2014;30:2068–2069.

47. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. Nature Methods
2014;12:59–60.

48. Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG Tools for Functional
Characterization of Genome and Metagenome Sequences. Journal of Molecular Biology 2016;428:726–
731.

49. Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. The carbohydrate-active enzymes
database (CAZy) in 2013. Nucleic Acids Research 2014;42:D490–D495.

50. Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, et al. dbCAN2: a meta server for automated
carbohydrate-active enzyme annotation. Nucleic Acids Research 2018;46:W95–W101.

51. Rawlings ND, Barrett AJ, Thomas PD, Huang X, Bateman A, Finn RD. The MEROPS database of
proteolytic enzymes, their substrates and inhibitors in 2017 and a comparison with peptidases in the
PANTHER database. Nucleic Acids Research 2017;46:D624–D632.

52. Saier MH Jr, Reddy VS, Tsu BV, Ahmed MS, Li C, Moreno-Hagelsieb G. The Transporter Classification
Database (TCDB): recent advances. Nucleic Acids Research 2016;44:D372–D379.

53. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with
DESeq2. Genome Biology 2014;15:550.

54. Damashek J, Tolar BB, Liu Q, Okotie Oyekan AO, Wallsgrove NJ, Popp BN, et al. Microbial oxidation of
nitrogen supplied as selected organic nitrogen compounds in the South Atlantic Bight. Limnology and
Oceanography 2019;64:982–995.

55. Massana R, Murray AE, Preston CM, DeLong EF. Vertical distribution and phylogenetic characterization
of marine planktonic *Archaea* in the Santa Barbara Channel. Applied and Environmental Microbiology
1997;63:50–56.

56. Teira E, Reinthaler T, Pernthaler A, Pernthaler J, Herndl GJ. Combining catalyzed reporter deposition-
fluorescence in situ hybridization and microautoradiography to detect substrate utilization by bacteria and
archaea in the deep ocean. Applied and Environmental Microbiology 2004;70:4411–4414.

57. Galand PE, Gutiérrez-Provecho C, Massana R, Gasol JM, Casamayor EO. Inter-annual recurrence of
archaeal assemblages in the coastal NW Mediterranean Sea (Blanes Bay Microbial Observatory).
Limnology and Oceanography 2010;55:2117–2125.

58. Legendre P. lmodel2: Model II regression. R package version 1.7-3. https://CRAN.R-
project.org/package=lmodel2. 2018.

59. Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York: Springer; 2016.

60. Man D, Wang W, Sabehi G, Aravind L, Post AF, Massana R, et al. Diversification and spectral tuning in
marine proteorhodopsins. The EMBO Journal 2003;22:1725–1731.

61. Pinhassi J, DeLong EF, Beja O, González JM, Pedrós-Alio C. Marine bacterial and archaeal ion-pumping
rhodopsins: genetic diversity, physiology, and ecology. Microbiology and Molecular Biology Reviews
2016;80:929–954.

62. Herndl GJ, Reinthaler T, Teira E, van Aken H, Veth C, Pernthaler A, et al. Contribution of *Archaea* to
total prokaryotic production in the deep Atlantic Ocean. Applied and Environmental Microbiology
2005;71:2303–2309.

63. Liu H, Zhang CL, Yang C, Chen S, Cao Z, Zhang Z, et al. Marine Group II dominates planktonic archaea
in water column of the northeastern South China Sea. Frontiers in Microbiology 2017;8:1098.

64. Church MJ, DeLong EF, Ducklow HW, Karner MB, Preston CM, Karl DM. Abundance and distribution
of planktonic Archaea and Bacteria in the waters west of the Antarctic Peninsula. Limnology and
Oceanography 2003;48:1893–1902.

65. Mincer TJ, Church MJ, Taylor LT, Preston C, Karl DM, DeLong EF. Quantitative distribution of
presumptive archaeal and bacterial nitrifiers in Monterey Bay and the North Pacific Subtropical Gyre.
Environmental Microbiology 2007;9:1162–1175.

66. Stoica E, Herndl GJ. Contribution of *Crenarchaeota* and *Euryarchaeota* to the prokaryotic plankton in the
coastal northwestern Black Sea. Journal of Plankton Research 2007;29:699–706.

67. Xie W, Luo H, Murugapiran SK, Dodsworth JA, Chen S, Sun Y, et al. Localized high abundance of
Marine Group II archaea in the subtropical Pearl River Estuary: implications for their niche adaptation.
Environmental Microbiology 2018;20:734–754.

68. Wang ZA, Cai W-J, Wang Y, Ji H. The southeastern continental shelf of the United States as an
atmospheric CO_2 source and an exporter of inorganic carbon to the ocean. Continental Shelf Research
2005;25:1917–1941.

69. Letourneau ML, Medeiros PM. Dissolved organic matter composition in a marsh-dominated estuary:
response to seasonal forcing and to the passage of a hurricane. Journal of Geophysical Research:
Biogeosciences 2019;124:1545–1559.

70. Medeiros PM, Babcock Adams L, Seidel M, Castelao RM, Di Iorio D, Hollibaugh JT, et al. Export of
terrigenous dissolved organic matter in a broad continental shelf. Limnology and Oceanography
2017;62:1718–1731.

71. Herfort L, Schouten S, Abbas B, Veldhuis MJW, Coolen MJL, Wuchter C, et al. Variations in spatial and
temporal distribution of Archaea in the North Sea in relation to environmental variables. FEMS
Microbiology Ecology 2007;62:242–257.

72. Hugoni M, Taib N, Debroas D, Domaizon I, Jouan Dufournel I, Bronner G, et al. Structure of the rare
archaeal biosphere and seasonal dynamics of active ecotypes in surface coastal waters. Proceedings of the
National Academy of Sciences 2013;110:6004–6009.

73. Bano N, Ruffin S, Ransom B, Hollibaugh JT. Phylogenetic composition of Arctic Ocean archaeal assemblages and comparison with Antarctic assemblages. Applied and Environmental Microbiology 2004;70:781–789.

74. Parada AE, Fuhrman JA. Marine archaeal dynamics and interactions with the microbial community over 5 years from surface to seafloor. The ISME Journal 2017;11:2510–2525.

75. Pereira O, Hochart C, Auguet J-C, Debroas D, Galand PE. Genomic ecology of Marine Group II, the most common marine planktonic Archaea across the surface ocean. MicrobiologyOpen 2019;8:e852.

76. Béjà O, Aravind L, Koonin EV, Suzuki MT, Hadd A, Nguyen LP, et al. Bacterial rhodopsin: evidence for a new type of phototrophy in the sea. Science 2000;289:1902–1906.

77. Frias-Lopez J, Shi Y, Tyson GW, Coleman ML, Schuster SC, Chisholm SW, et al. Microbial community gene expression in ocean surface waters. Proceedings of the National Academy of Sciences 2008;105:3805–3810.

78. Poretsky RS, Hewson I, Sun S, Allen AE, Zehr JP, Moran MA. Comparative day/night metatranscriptomic analysis of microbial communities in the North Pacific subtropical gyre. Environmental Microbiology 2009;11:1358–1375.

79. Satinsky BM, Crump BC, Smith CB, Sharma S, Zielinski BL, Doherty M, et al. Microspatial gene expression patterns in the Amazon River Plume. Proceedings of the National Academy of Sciences 2014;111:11085–11090.

80. Alonso-Sáez L, Morán XAG, González JM. Transcriptional patterns of biogeochemically relevant marker genes by temperate marine bacteria. Frontiers in Microbiology 2020;11:465.

81. Arandia-Gorostidi N, González JM, Huete-Stauffer T, Ansari MI, Morán XAG, Alonso-Sáez L. Light supports cell-integrity and growth rates of taxonomically diverse coastal photoheterotrophs. Environmental Microbiology 2020;22:3823-3837.

82. Giovannoni SJ, Bibbs L, Cho J-C, Stapels MD, Desiderio R, Vergin KL, et al. Proteorhodopsin in the ubiquitous marine bacterium SAR11. Nature 2005;438:82–85.

83. Gómez-Consarnau L, González JM, Coll-Lladó M, Gourdon P, Pascher T, Neutze R, et al. Light stimulates growth of proteorhodopsin-containing marine Flavobacteria. Nature 2007;445:210–213.

84. Steindler L, Schwalbach MS, Smith DP, Chan F, Giovannoni SJ. Energy starved *Candidatus* Pelagibacter ubique substitutes light-mediated ATP production for endogenous carbon respiration. PLoS ONE 2011;6:e19725.

85. Song Y, Cartron ML, Jackson PJ, Davison PA, Dickman MJ, Zhu D, et al. Proteorhodopsin overproduction enhances the long-term viability of *Escherichia coli*. Applied and Environmental Microbiology 2020;86:e02087–19.

86. Baykov AA, Malinen AM, Luoto HH, Lahti R. Pyrophosphate-fueled Na^+ and H^+ transport in prokaryotes. Microbiology and Molecular Biology Reviews 2013;77:267–276.

87. Rinta-Kanto JM, Sun S, Sharma S, Kiene RP, Moran MA. Bacterial community transcription patterns during a marine phytoplankton bloom. Environmental Microbiology 2012;14:228–239.

88. Vila-Costa M, Sharma S, Moran MA, Casamayor EO. Diel gene expression profiles of a phosphorus limited mountain lake using metatranscriptomics. Environmental Microbiology 2013;15:1190–1203.

89. Needham DM, Fuhrman JA. Pronounced daily succession of phytoplankton, archaea and bacteria following a spring bloom. Nature Microbiology 2016;1:16005.

90. Hosie AHF, Poole PS. Bacterial ABC transporters of amino acids. Research in Microbiology 2001;152:259–270.

91. Anderson JJ, Oxender DL. *Escherichia coli* transport mutants lacking binding protein and other components of the branched-chain amino acid transport systems. Journal of Bacteriology 1977;130:384–392.

92. Montesinos ML, Herrero A, Flores E. Amino acid transport in taxonomically diverse cyanobacteria and
identification of two genes encoding elements of a neutral amino acid permease putatively involved in
recapture of leaked hydrophobic amino acids. Journal of Bacteriology 1997;179:853–862.

93. Walshaw DL, Reid CJ, Poole PS. The general amino acid permease of *Rhizobium leguminosarum* strain
3841 is negatively regulated by the Ntr system. FEMS Microbiology Letters 1997;152:57–64.

94. Berg GM, Shrager J, Glöckner G, Arrigo KR, Grossman AR. Understanding nitrogen limitation in
Aureococcus anophagefferens (Pelagophyceae) through cDNA and qRT-PCR analysis. Journal of
Phycology 2008;44:1235–1249.

95. Wurch LL, Gobler CJ, Dyhrman ST. Expression of a xanthine permease and phosphate transporter in
cultures and field populations of the harmful alga *Aureococcus anophagefferens*: tracking nutritional
deficiency during brown tides. Environmental Microbiology 2014;16:2444–2457.

96. Cooper JT, Sinclair GA, Wawrik B. Transcriptome analysis of *Scrippsiella trochoidea* CCMP 3099
reveals physiological changes related to nitrate depletion. Frontiers in Microbiology 2016;7:639.

97. Cunliffe M. Purine catabolic pathway revealed by transcriptomics in the model marine bacterium
Ruegeria pomeroyi DSS-3. FEMS Microbiology Ecology 2016;92:fiv150.

98. Charlier D, Le Minh PN, Roovers M. Regulation of carbamoylphosphate synthesis in *Escherichia coli*: an
amazing metabolite at the crossroad of arginine and pyrimidine biosynthesis. Amino Acids 2018;50:1647–
1661.

TABLES AND FIGURES

Table 1 Transcriptional traits shared or distinct among *Ca.* Poseidoniales families.[1]

Table 1 Transcriptional traits shared or distinct among euryarchaeal families[1]

Putative function	Relevant gene(s)	Distribution	Evidence
Proteorhodopsin	Proteorhodopsin gene	Both families	Highly transcribed (Fig. 2); enriched when O1 active (Fig. 3)
Pyrophosphatase	*hppA*	Both families	Highly transcribed (Fig. 2)
Protease/peptidase	*pepF*, *lonB*, *pepA*, *pip*, carboxypeptidase A, subtilase	Both families	Highly transcribed (Fig. 2); depletion during dark incubation (Fig. 5)
ß-oxidation	3-hydroxyacyl-CoA dehydrogenase	Both families	Highly transcribed (Fig. 2); enriched when O1 active (Fig. 3)
Proteorhodopsin retinal synthesis	*crtI*, *crtD*	*Ca.* Thalassarchaeaceae only	Highly transcribed in SIMO MAGs (Fig. 2); depletion during dark incubation (Fig. 5)
Electron transport	Halocyanin gene	*Ca.* Thalassarchaeaceae only	Highly transcribed in SIMO MAGs (Fig. 2); enriched when O1 active (Fig. 3)
Amino acid transport/metabolism	*aapJ*, *livK*, *aroA*, *trpE*	*Ca.* Thalassarchaeaceae only	Highly transcribed (Fig. 2); depletion during dark incubation (Fig. 5)
Xanthine/uracil permease	*pbuG*	*Ca.* Thalassarchaeaceae only	Highly transcribed (Fig. 2); enriched when O1 active (Fig. 3); depletion during incubations (Fig. 5)
Amino acid synthesis	*carA*	*Ca.* Poseidoniaceae only	Highly transcribed (Fig. 2)
Carbohydrate synthesis	Family 2 glycosyl transferase	*Ca.* Poseidoniaceae only	Highly transcribed (Fig. 2)

[1]Putative *Ca.* Poseidoniales families are *Ca.* Poseidoniaceae (MGIIa; MAG RS440) and *Ca.* Thalassarchaeaceae (MGIIb; SIMO Bins 19-2, 31-1).

Fig. 1 A) Relative abundance of *Ca.* Poseidoniales genera in Sapelo Island metatranscriptomes (*n*=24). The dendrogram (top) shows grouping by similarity. Season is indicated by color beneath sample names. The bar chart shows the abundance of *Ca.* Poseidoniales transcripts L^{-1} and the stacked bar charts show the relative abundance of genera (% total *Ca.* Poseidoniales transcripts), colored by genus. Dominant genera are indicated below the stacked bar chart. "Highly active" samples for each genus are marked and were used for analysis of differential transcription. **B)** NMDS of *Ca.* Poseidoniales metatranscriptome reads using a Hellinger distance matrix of genera relative abundances. Each point represents a sample, with color and shape denoting season and day/night, respectively. Shading indicates sampling year and original study. Arrows represent vectors for each genus. NMDS stress and the results of a PERMANOVA analyzing sums of squares by season are indicated. **C)** Boxplots of *Ca.* Poseidoniales reads L^{-1}, grouped by season (winter, *n*=4; spring, *n*=3; summer, *n*=10; fall, *n*=7). Values from individual metatranscriptomes are overlain. Results of an ANOVA are indicated; letters at the top indicate post-hoc groups according to Tukey's HSD test.

741
Fig. 2 Boxplots of highly-transcribed MAG genes (top 5%) in Sapelo Island field metatranscriptomes. Overlain points
show CPM for individual metatranscriptomes (n=24). Shading indicates COG functional category assigned by
eggNOG-mapper (genes assigned to group S were similar to proteins of unknown function in the COG database, while
genes with no COG assignment did not match proteins in the COG database). Diamonds indicate genes highly
transcribed in all MAGs (pink), in *Ca.* Thalassarchaeaceae (MGIIb) MAGs only (gray), or in the *Ca.* Poseidoniaceae
(MGIIa) MAG and one *Ca.* Thalassarchaeaceae MAG (green).

Fig. 3 Log₂-fold change of SIMO Bin 19-2 (genus O1) genes differentially transcribed in field metatranscriptomes
where transcriptional activity of *Ca.* Poseidoniales was dominated by genus O1 (see Fig. 1, Table S4), calculated with
DESeq2. Error bars show estimated standard error. Only genes with adjusted *p*-values <0.1 are shown. Color indicates
COG functional category (see Fig. 2). Bold indicates genes in the top 5% median transcript coverage across field
metatranscriptomes (Fig. 2).

Fig. 4 Comparisons of competitive read mapping to *Ca.* Poseidoniales genera at the beginning and end of 24-hour
incubations of Sapelo Island water conducted by Vorobev et al. [23]. The dendrogram (top) shows grouping by
similarity. The bar chart below the dendrogram is the abundance of *Ca.* Poseidoniales transcripts L⁻¹. Stacked bar
charts show the relative abundance of genera (% total *Ca.* Poseidoniales transcript reads), colored by genus.

Fig. 5 Log$_2$-fold change of SIMO Bin 19-2 (genus O1) genes differentially abundant in T$_{24}$ versus T$_0$
metatranscriptomes from Sapelo Island high tide waters. Error bars show estimated standard error. Only genes with
adjusted *p*-values<0.1 are shown. Color indicates COG functional category (see Fig. 2). Bold indicates genes in the
top 5% median transcript coverage across field metatranscriptomes (Fig. 2).

Fig. 6 A) Violin plots of log-transformed *Ca.* Poseidoniales 16S rRNA gene abundances across regions in the SAB,
with overlain boxplots. Width of the violin plot corresponds to data probability density. Color denotes sampling region.
Letters above boxes denote post-hoc grouping according to Tukey's HSD test. **B)** Scatterplot of bacterial and *Ca.*
Poseidoniales 16S rRNA gene abundances in the SAB. The solid line shows the best fit of a model II (major axis)
linear regression, with dashed lines showing a 95% confidence interval of the slope. Regression parameters are shown
on the plot.