Running head:

OASTLs confer Se resistance and degrade L-SeCys in *Arabidopsis*

Corresponding author:

Prof. Moshe Sagi

French Associates Institute for Agriculture and Biotechnology of Drylands, Blaustein Institutes for Desert
Research, Sede Boqer campus, Ben-Gurion University, P.O.B. 653, Beer Sheva 84105, Israel

fax +972 8 6472984; e-mail gizi@bgu.ac.il

Research category:

Ecophysiology and Sustainability, Associate Editor Graham Farquhar
11

12 **Title:**

13   Impairment in O-acetylserine-(thiol) lyase A and B, but not C, confers higher selenate
14 sensitivity and uncovers role for A, B and C as L-Cys and L-SeCys desulfhydrases in
15 *Arabidopsis*
16
17

Authors:

Assylay Kurmanbayeva[1]
Aizat Bekturova[1]
Aigerim Soltabayeva[1]
Sudhakar Srivastava[1]
Dinara Oshanova[1]
Zhadyrassyn Nurbekova[1]
Moshe Sagi[1]*
Author addresses:

[1] Plant Stress laboratory, French Associates Institute for Agriculture and Biotechnology of Drylands,
Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev, Sede Boqer Campus
84990, Israel (A.K., A.B., A.S., S.S.,D.O., Z.N., M.S.)
30

31 **Summary:**

32 The cytosolic OASTLA and chloroplastic OASTLB have significantly higher desulfhydrase
33 activity rates than the cytosolic DES1 and are able to degrade L-Cys and L-SeCys to sulfide and
34 selenide, respectively in *Arabidopsis*.
35

Footnotes:

List of author contributions:

A.K. designed and performed experiments and analyzed the data. A.B. involved in SO kinetic assay and

selenate detection. A.S. involved in RNA extraction. D.O. performed Western-blot, S.S and Z.N.

participated in native SDS-PAGE assays. M.S. conceived the original idea, designed the research plan,

and supervised the research work; the article was jointly written by A.K. and M.S.

This research was supported by the Israel Science Foundation grant no. 212/13

* Corresponding author; e-mail gizi@bgu.ac.il

ABSTRACT

The role of the cytosolic O-acetylserine-(thiol) lyase A (OASTLA), chloroplastic OASTLB and mitochondrion OASTLC in plant resistance/sensitivity to selenate was studied in *Arabidopsis* plants. Impairment in OASTLA and B resulted in reduced biomass, chlorophyll and soluble protein levels compared with impaired OASTL C and Wild-Type treated with selenate. The lower organic-Se and protein-Se levels followed by decreased organic-S, S in proteins and total glutathione in *oastlA* and *oastlB* compared to Wild-Type and *oastlC* are indicative that Se accumulation is not the main cause for the stress symptoms, but rather the interference of Se with the S-reduction pathway. The increase in sulfite oxidase, adenosine 5′-phosphosulfate reductase, sulfite reductase and OASTL activity levels, followed by enhanced sulfite and sulfide, indicate a futile anabolic S-starvation response to selenate-induced organic-S catabolism in *oastlA* and *oastlB* compared to Wild-Type and *oastlC*.

Additionally, the catabolic pathway of L-cysteine degradation was enhanced by selenate, and similar to L-cysteine producing activity, *oastlA* and *B* exhibited a significant decrease in L-cysteine desulfhydrase (DES) activity, compared with WT, indicating a major role of OASTLs in L-cysteine degradation. This notion was further evidenced by sulfide dependent DES in-gel activity, immunoblotting, immunoprecipitation with specific antibodies and identification of unique peptides in activity bands generated by OASTLA, B and C. Similar responses of the OASTLs in Seleno-Cysteine degradation was demonstrated in selenate stressed plants. Notably, no L-cysteine and L-Seleno-Cysteine DES activity bands but those related to OASTLs were evident. These results indicate the significance of OASTLs in degrading L-cysteine and L-SelenoCysteine in *Arabidopsis*.

Keywords: Sulfur and selenium metabolism**,** L-Cys desulfhydrase, OASTLs, SeCys desulfhydrase, *Arabidopsis*

INTRODUCTION

The amino acid selenocysteine, a component of selenoproteins in humans, contains Se (Papp et al., 2007). Plants serve as a source of Se for mammalians; yet Se is not essential for plants. The chemical similarity of Se to S, enables plants to readily uptake Se via root sulfate transporters (Pilon-Smits and Quinn, 2010; Sors et al., 2005) and incorporate it into organic Se compounds via the S assimilation pathway components (Grant et al., 2011). The pathway (see Supplemental Fig. S1) is initiated with the adenylation of sulfate/selenate by ATP sulfurylase (ATPS, EC 2.7.7.4) to generate adenosine 5′-phosphosulfate (APS)/adenosine 5′-phosphoselenate (APSe). APS/APSe is then reduced to sulfite/selenite by the plastidic APS reductase (APR, EC 1.8.99.2) (Mroczek-Zdyrska and Wójcik, 2012; Schiavon et al., 2015; White, 2015). The toxic sulfite/selenite is further reduced to sulfide by the chloroplast-localized sulfite reductase (SiR, EC 1.8.7.1) (Khan et al., 2010; Yarmolinsky et al., 2013), whereas selenite can be reduced enzymatically to selenide by SiR and non-enzymatically by reduced glutathione [GSH (Fisher et al., 2016; Seko et al., 1989; White, 2015)]. Further, the generated sulfide/selenide together with O-acetylserine (OAS), the latter catalyzed by serine acetyltransferase (SAT, EC 2.3.1.30), are incorporated into cysteine (Cys) or selenocysteine (SeCys) in a reaction catalyzed by the O-acetylserine-(thiol) lyase (OASTL, EC 2.5.1.47) in cytosol (OASTLA), plastid (OASTLB), and mitochondria (OASTLC) [(Pilon-Smits and Womg, 2012; White, 2015; Wirtz and Hell, 2006) (see in Supplemental Fig. S1)]. The replacement of Cys by SeCys in proteins is thought to be a cause for selenate toxicity in *Arabidopsis* (Sabbagh and Van Hoewyk, 2012; Van Hoewyk, 2018).

Three major Se toxicity mechanisms were recently concluded in plants: competition between Se and S in primary and secondary metabolism, incorporation of SeCys and SeMet into proteins, and disruption caused by oxidative stress of metabolism and cellular structures (Lima et al., 2018; White, 2018). Amongst others, the role of Cys catabolic enzymes in L-SeCys degradation was not shown. Moreover, while the cytosolic L-cysteine desulfhydrase1 (DES1, EC 4.4.1.1) was shown to catalyze the degradation of cysteine to sulfide, pyruvate and ammonia (Alvarez et al., 2012b; Gotor et al., 2013; Romero et al., 2013), DES1 or other L-Cys catabolic enzymes were not shown to increase L-Cys and/or L-SeCys degradation in response to selenate.

In the current study we show that *oastlA and oastlB Atabidopsis* mutants are more sensitive to the presence of selenate in the growth medium than *oastlC* and WT. The

significantly lower selenium level in the protein fraction and biomass of *oastlA* and *oastlB*

compared to WT and *oastlC* mutant led us to explore the interference of selenate in the sulfate

reduction pathway as the cause for the lower growth rate in the two mutants. The results

demonstrate that in the absence of active OASTLA and OASTLB, the selenate interference in

the S assimilation pathway leads to a decrease in organic S, S in protein fraction and total

glutathione and has a more negative effect on the resulting biomass accumulation than the

expected positive effect of lower selenium content in the organic fraction. Increase in the activity

of core enzymes of the sulfate reduction pathway, the APR and SiR, followed by enhanced

sulfite and sulfide levels uncovers a futile anabolic S-starvation response to the selenate induced

organic-S catabolism in *oastlA* and *oastlB* compared to WT and *oastlC*. A major role for the

OASTL proteins in acting as L-Cys and L-SeCys desulfhydrases in control and Selenate stressed

plants was demonstrated as well.

RESULTS

oastlA **and** *oastlB* **mutants are more sensitive to selenate than** *WT* **and** *oastlC*

Replacement of cysteine by selenocysteine in proteins is considered to be toxic to plants

(Lima et al., 2018; White, 2018); whereas minimizing the incorporation of selenocysteine into

proteins has been shown to be an effective strategy to increase Se tolerance in plants (Van

Hoewyk et al., 2005). *Arabidopsis* mutants impaired in the OASTLs: the cytosolic (*oastlA*),

plastidal (*oastlB*) and mitochondrial (*oastlC*), were employed to examine whether these

impairments will reduce selenium in the protein fraction and thus enhance plant tolerance to

selenium toxicity.

Figure 1. The effect of 40 µM selenate treatment on growth parameters and Se in protein fraction in WT, *oastlA, oastlB* and *oastlC KO* mutants of *Arabidopsis thaliana*. A. Phenotype of plants grown for 14 days under control (left panel) and 40µM selenate treated (right panel) conditions. The weight of plant biomass accumulation (B) and relative weight to their own controls (C) supplemented with or without selenate for 14 days. The values are means ± SE (n = 15). Remaining chlorophyll level (D) and Se in protein fraction (E). The values are means ± SE (n = 4). Values denoted with different letters are significantly different according to the T- test analyses, P < 0.05 (JMP 8.0 software, http://www.jmp.com/). Different uppercase letters indicate significant differences between control and Se treatment of each genotype. Asterisks indicate significant differences between *WT* and *oas-tlA, oas-tlB, oastlC* KO plants subjected to the same treatment.

The mutation of the various OASTL mutant plants was verified by immunoblot analysis, using an OASTL C polyclonal antiserum which also cross-reacts with OASTL A and OASTL B (kindly provided by Prof. Dr. S. Kopriva, University of Cologne, Köln). The three bands detected in WT fractionated proteins corresponded to OASTL A, B, and C from bottom to top respectively, whereas the lower, middle or upper bands were missing in the fractionated proteins of *oastlA, oastlB* or *oastlC* mutants respectively (Supplemental Fig. S2A).

oastlA, oastlB or *oastlC* mutants grown with *WT* plants on 0.5 MS agar medium in the presence and absence (control) of 40 µM selenate for 2 weeks exhibited similar growth as *WT* plant in the absence of selenate and showed a significant reduction in growth rate and chlorophyll level in response to selenate. Yet, the reduction in biomass and chlorophyll content

shown in *oastlA* and *oastlB* mutants was significantly higher than in *WT* and *oastlC* (Fig. 1A-D),

in spite of the lower Se level in the protein fraction of the *oastlA* and *oastlB* (Fig. 1E).

Since at high levels selenate and Se can act as pro-oxidants and cause oxidative stress

(Grant et al., 2011; Mroczek-Zdyrska and Wójcik, 2012), Se and selenate levels were determined

in the leaves. Yet, no higher total Se and selenate were detected in *oastlA* and *oastlB* as

compared to *WT* and *oastlC* (Supplemental Fig. S2B and C) which could be attributed to the

decreased biomass accumulation and higher chlorophyll degradation rates in *oastlA* and *oastlB*

mutants. These results indicate that the levels of Se in the protein fraction, total Se and selenate

in leaf tissue are not the only cause for the higher sensitivity to Selenate noticed in *oastlA* and

oastlB compared to WT and *oastlC* mutant.

The interference in sulfate assimilation by selenate is stronger in *oastA* and *B* than in WT

and *oastlC* mutant

Since higher sensitivity in the *oastlA* and *oastlB* mutants was not related to Se content in

the leaves, the sulfate reduction pathway was further studied, estimating the level of selenate

interference. Cysteine biosynthesis, the last step of the sulfate reduction pathway catalyzed by

OASTLs was studied in the three mutants in comparison to WT. *OASTLA, OASTLB* or *OASTLC*

transcripts were not expressed in *oastlA, oastlB* or *oastlC* mutants respectively, indicating the

complete knockout of these genes (Supplemental Fig. S2D-F). Importantly, under control

conditions *OASTLB* transcript was enhanced in *AtoastA* and *C* mutants, *OASTLA* in *AtoastB*,

whereas *OASTC* was enhanced in *AtoastB,* indicating a level of complementation to compensate

for the mutation among the various transcripts. The presence of selenate in the growth medium

resulted in a significant enhancement of the transcripts, exhibiting the increase of the three

transcripts in *WT*, as well as *OASTLA* and *B* transcripts in *oastlC*, whereas in *oastlA* the transcript

of *OASTLB* and in *oastlB OASTLA,* both were significantly increased (Supplemental Fig. S2D-

F). Similarly, the absence of either OASTLA or C in *oastlA* or *oastlC* mutants respectively,

resulted in the enhancement of the two other functioning OASTLs in each of the mutants as

compared to the proteins expression in WT leaves grown under unstressed conditions, whereas in

oastlB only OASTLA exhibited such response (Supplemental Fig. S2A). Supplementation of

selenate to the growth medium resulted in increased expression of OASTLB and C proteins in

WT leaves, and the enhancement of OASTLB in *oastlA*, OASTLA in *oastlB* and OASTLB in

AtoastlC as compared to their control unstressed plants (Supplemental Fig. S2A).

In-spite of the enhanced protein expression of OASTLB and C in the absence of selenate

and of OASTLB in the presence of selenate, the impairment in OASTLA resulted in a strongly

decreased cysteine production activity of OASTL in *AtoastlA* mutant, exhibiting a reduction in

activity by 87% and 72% as compared to WT under control and selenate treated conditions,

respectively. The *AtoastlB* mutant showed a significant decline of 28% in both conditions, while

in *oastlC* OASTL activity was similar to the activity in WT plants. Interestingly, the activity in

all the examined genotypes was significantly increased with selenate treatment, likely as the

result of the enhanced expression of at least one of the active OASTL proteins (Fig. 2A). The

comparable activity rates to WT found in *oastlC* mutant might be the result of the enhanced

expression of the two major proteins, the OASTLA and B as compared to WT in control as well

as under stressed conditions (Supplemental Fig. S2A). Taken together, the enhanced OASTL

activity in leaves of selenate stressed WT and the OASTL mutants may indicate S-limitation type

response shown before in *Arabidopsis* leaves in response to limited S supply (Barraso et al.,

1995; Hesse et al., 1999).

To examine this assumption the Cys, total glutathione and total organic S levels, as well

as the S level in protein fraction were detected. No differences were noticed in cysteine level

between WT and mutants in control and selenate treated plants, whereas in selenate treated

plants a higher cysteine level was evident with WT and *oastlC* compared to these plants grown

under control conditions (Fig. 2B). Determination of total glutathione revealed a decrease only in

oastlC leaves whereas the other mutants exhibited comparable levels to WT leaves in plants

grown in control conditions. In contrast, *oastlA* and *oastB* showed decreased total glutathione

level compared to WT and *oastlC* leaves in plants exposed to selenate. Significantly, all

genotypes exposed to selenate exhibited significantly lower glutathione levels than control plants

(Fig. 2C). Determination of the organic S level revealed a significant reduction in all the

genotypes at 40 μM selenate compared to untreated controls. Under the selenate stress, *oastlA*

and *oastlB* exhibited a significant decrease in organic S as compared to WT and *oastlC* (Fig.2D).

Figure 2. Cys biosynthesis and S containing compounds in WT, *oastlA, oastlB* and *oastlC* KO mutants supplemented with or without selenate for 14 days. The effect of 40 µM selenate treatment on OAS-TL activity (A), Cysteine level (B), Total glutathione (C), Total organic S (D) and S in protein fraction (E). The values are means ± SE (n = 4). Values denoted with different letters are significantly different according to T-test analyses, $P < 0.05$ (JMP 8.0 software, http://www.jmp.com/). Different uppercase letters indicate significant differences between control and Se treatment of each genotype. Asterisks indicate significant differences between WT and the KO plants *oastlA, oastlB and oastlC* subjected to the same treatment.

The level of S in the protein fraction, under control conditions, was higher in *oastlA* than in WT, whereas in the other mutants it was similar to *WT*. Selenate supplementation resulted in a significantly lower S content in the protein fraction in *oastlA* and *oastlB* as compared to their controls and selenate treated WT and *oastlC* (Fig.2E). Taken together, the lower total glutathione, organic S and S in protein fraction in the leaf tissue of mutants impaired in OASTLA or B proteins fed with selenate (Fig. 2C, D and E), indicates possible interference in the S reduction pathway by selenate, resulting in retarded plant growth as compared to WT and *oastlC* mutant.

The presence of selenate in the growth medium induces catabolic processes in *Arabidopsis*

Cysteine can be found in plants as free cysteine, being among the amino acids that build proteins, as well as an intermediate of Met biosynthesis, GSH and many other thiol species (Hesse and Hoefgen, 2003; Ravanel et al., 1998). Under the long day (14/10 light/dark) growth conditions free cysteine levels in leaves of plants growing in plates containing 0.5 MS medium,

were similar in the mutants compared to WT either in control or in selenate treated plants (Fig.

2B). Considering the lowest cysteine generating activity level of OASTL was evident in *oastlA*

and *B* (Fig. 2B), this result indicates a possible catabolic protein degrading activity which can act

as an additional source of free L-cysteine in *oastlA* and *B*. In support of this notion is the

significantly lower soluble protein content evident in *oastlA* and *B* compared to WT and *oastlC*

mutant grown with selenate in the growth medium (Fig. 3A).

Recently, a lower biomass accumulation rate, the result of reduced organic S content was

shown to be the consequence of higher L-cysteine degradation by desulfhydrase (DES) activity

of OASTL in the perennial halophyte *Sarcocornia*, but not in the annual halophyte *Salicornia*,

both fed with high sulfate (Kurmanbayeva et al., 2017). To examine whether OASTLs play a

role in cysteine degradation also in non-halophyte plants, we studied first the expression of the

cytosolic *DES1* claimed previously to play an important role in cytosolic L-Cys degradation

(Álvarez et al., 2010). Unlike the *OASTLs* whose transcript levels were enhanced in WT plants in

response to selenate (Supplemental Fig. S2 D-F), a significant decrease in *DES1* transcript

abundance was evident in WT, *oastlB* and *C,* whereas in *oastlA* mutant a non-significant

decrease was noticed. Under control and selenate stress conditions, *DES1* transcript abundance

was the highest in *oastlB* (Fig. 3B).

Unlike the *DES1* transcript abundance in *WT* and *OASTLs* mutants, but similar to the L-

Cys generation activities of the OASTLs (Fig. 2A), the total L-Cys DES kinetic activity was

enhanced in all genotypes grown with selenate compared to unstressed control plants (Fig. 3C).

DES activity rate was significantly lower in *oastlA* and *oastlB* mutants compared to *WT* grown

under unstressed conditions, while when exposed to selenate *oastlA* exhibited a significant lower

activity rate than *WT* and the other genotypes (Fig.3C). These results indicate a possible role of

these OASTL isoforms in L-Cys degradation and not only in L-Cys biosynthesis (Figs.2A, 3C).

Figure 3. Soluble protein level and S metabolism components in WT, *oastlA, oastlB* and *oastlC* KO mutants supplemented with or without selenate for 14 days. The effect of 40 µM selenate treatment on soluble protein content (A), transcript expression of DES1 (B), L-Cysteine desulfhydrase (DES) activity (C), sulfide (D) and sulfite levels (E), Sulfite oxidase (SO) activity (F), Adenosine 5′-phosphosulfate reductase (APR) activity (G) and Sulfite reductase (SiR) activity (H). Transcript level of *DES1* was detected by quantitative reverse transcription-PCR using *ACTIN2* as the housekeeping transcript for normalization. The relative expression in the various genotypes was analyzed using the normalized WT control as reference. The values are means ± SE (n = 4). Values denoted with different letters are significantly different according to T-test analyses, P < 0.05 (JMP 8.0 software, http://www.jmp.com/). Different uppercase letters indicate significant differences between control and Se treatment of each genotype. Asterisks indicate significant differences between WT and *oastlA, oastlB, oastlC* plants subjected to the same treatment.

Notably, in spite of the lower or similar DES activity rate, *oastlA* and *oastlB* mutants exhibited higher sensitivity to selenate stress, accumulating lower biomass. This can be attributed to the lowest capacity of L-Cys biosynthesis by OASTL in *oastlA* and *oastlB* mutants grown

with selenate that resulted in low levels of organic sulfur, total glutathione as well as low S in the

protein fraction (Fig. 2A,C,D,E). This could result in S-starvation type responses such as the

higher sulfide level evident in these mutants compared to WT and *oastlC* when stressed by

selenate, exhibiting, excluding *oastlC,* higher sulfide than the control unstressed plants, (Fig.

3D). The higher sulfite level detected in these mutants compared to WT and *oastlC* grown with

selenate (Fig. 3E), shown before when detected on fresh weight basis to be toxic in *Arabidopsis*

and tomato plants (Brychkova et al., 2012; Brychkova et al., 2013; Yarmolinsky et al., 2014), is

likely another S-starvation type response and an additional cause for the mutants' sensitivity to

selenate. Interestingly, while under control unstressed conditions SO activity was similar in

mutants and in WT treated with selenate, the mutants exhibited a drastically higher SO activity

rate than WT (Fig. 3F). This enhanced SO activity in the mutants compared to WT is a typical

response of SO to sulfite increase as was shown before (Brychkova et al., 2012; 2013;

Yarmolinsky et al., 2013; 2014). The selenate induced sulfite increase in *oastlA* and *oastlB*

mutants is most likely the result of the increased APR activity rate in these mutants compared to

WT and *oastlC* grown with selenate (Fig. 3G). The increase in APR activity rate in *oastlA* and

oastlB leaves is an expected response to S-starvation (Lee et al., 2011 and references therein), as

featured in the current study by lower glutathione, organic S and S in the protein fraction of these

mutants compared to WT and *oastlC* (Fig. 2C,D,E). Further reinforcement for this notion is the

significant increase in SiR activity resulting in the enhanced sulfide in *oastlA* and *oastlB* treated

with selenate, whereas no differences between the genotypes were noticed in the unstressed

plants (Fig. 3H, D). Taken together, the increase in SO, APR and SiR activity, followed by the

enhanced sulfite and sulfide levels (Fig. 3D-H), indicates a futile anabolic S-starvation response

to the selenate induced organic-S catabolism (Fig. 2C, D, E) in *oastlA* and *oastlB* compared to

Wild-Type and *oastlC*.

OAS-TLs degrades L-Cys and L-SeCys to H$_2$S and H$_2$Se respectively

The decrease in L-Cys desulfydrase kinetic activity in the *oastlA* and *oastlB* mutants

compared to WT (Fig. 3C) raised the possibility that OASTLs may act as L-Cys and L-SeCys

desulfhydrases. This was examined by employing crude protein extracted from leaves of 14 day

old selenate stressed and unstressed WT, *oastlA*, *B* and *C* plants, fractionated by native SDS-

PAGE (Sagi and Fluhr, 2001; Srivastava et al., 2017; Yesbergenova et al., 2005) to distinguish

between the activity bands of the various OASTLs. The gels were then subjected to DES activity

reaction solution containing L-Cys or L-SeCys as a substrate under reducing conditions,

generated in the presence of β-Mercaptoethanol. The reaction of lead (as lead acetate) with

sulfide or selenide (the consequence of desulfhydration activity that degrades L-Cys or L-SeCys,

respectively), results in a brown-dark precipitate exhibiting the activity band at the position of

the fractionated active enzyme. When the substrates (L-Cys or L-SeCys) or reducing conditions

(β-Mercaptoethanol) were omitted from the reaction solution, the activity bands, the precipitated

product of the generated lead sulfide or lead selenide, were not seen (Supplemental Fig. S3A). In

contrast, when the substrate was present under reducing condition, three distinguishable activity

bands were present in the location of the fractionated WT proteins (Fig.4A). Significantly, the

fractionated proteins extracted from *oastlA*, *oastlB* and *oastlC* mutants exhibited the absence of

the lowest, middle and most upper activity band respectively, compared to WT, with L-Cys as

the substrate (Fig.4A). When L-SeCys was present in the reaction solution as the substrate, the

two lowest activity bands with the same mobility as with the L-Cys desulfhydrase activity were

present in the fractionated WT proteins, whereas the fractionated proteins of *oastlA* and *oastlB*

mutants were missing the lowest and middle bands, respectively (Fig. 4B). The absence of the

most upper activity band of OASTLC in WT and *oastlA*, *oastlB* mutants, is likely the result of

the relatively low L-SeCys desulfhydrase activity rate by OASTLC, being below the detection

level when employing L-SeCys as a substrate. Importantly, the presence of selenate in the

growth medium resulted in the general enhancement of L-Cys and L-SeCys desulfhydrase

activities of OAS-TL enzymes compared to the control conditions (Fig.4A and B). These results

indicate that OASTLA, OASTLB and OASTLC play a role not only in Cys/SeCys biosynthesis,

but also in degrading L-Cys and L-SeCys.

Interestingly, while the homozygous *Atdes1* KO T-DNA mutant SALK_103855 was

shown to express ca 45 to 60% of *DES1* (AT5G28030) compared to WT root transcript (see

Figure 7C in Mei et al., 2019), the homozygous KO mutant SALK_103855 exhibited almost the

absence and ca 29% transcript expression compared to WT employing oligonucleotides upstream

and downstream along the transcript length, respectively. SALK_205358, shown before to be a

des1 null mutant (Jin et al., 2018) exhibited ca 40% decrease in *DES1* compared to the

expression in WT leaves, employing the oligonucleotides described above (Supplemental Fig.

S3B and E). Yet, in-spite of the transcript decrease the *Atdes1* KO (SALK_205358) exhibited

kinetic activities of cysteine generation by OASTLs and total desulfhydration activity in control

and selenate stressed plants comparable to WT (Fig. S3C and D, respectively).

Since SALK_103855, KO mutant of *Atdes1* grown on MS medium was also

demonstrated to be responsible for only a minor portion of the Cys-desulfudrase activity

compared to WT [14% of the total activity in leaves of plants grown in plates containing agar

with MS medium (Alvarez et al., 2010)], an in-gel desulfydrase activity was employed to further

examine the significance of DES1 and the OASTLs in desulfhydrase activity in Arabidopsis

plants. NATIVE-SDS was employed to fractionate the proteins (Sagi and Fluhr, 2001;

Yesbergenova et al., 2005) for desulfhydrase in-gel activities in WT and homozygous

SALK_205358 *Atdes1* mutant employing L-Cys or L-SeCys as substrates. The existence of

OASTLA, OASTLB and OASTLC activity bands with relative intensity in *Atdes1* mutant as

with WT was evident (Supplemental Fig. S3E). We further examined L-Cys desulfhydrase

activity in SALK_103855 leaves and further noticed similar activity bands in WT and the *Atdes1*

homozygous mutant (Supplemental Fig. S3F, left insert). These results demonstrate that the

OASTL enzymes may play a significant role in degrading L-cys and L-SeCys to sulfide and

selenide respectively in Arabidopsis leaves.

To further support the identification of the enzymes which generated the activity bands, a

Western blot analysis employing the lower middle and upper sliced activity bands in WT (Fig.

4A and B) was performed after the fractionation of the proteins in SDS-PAGE, using OASTL C

antibody which cross reacts also with OAS-TL A and B proteins (kindly provided by Prof. R.

Hell, Heidelberg University, Germany). Different mobility of the single bands was demonstrated

(Fig.4C left side), supporting the notion of OASTL A, B and C in L-Cys and OASTL A, B

isoforms playing a role in L-SeCys degradation (Fig.4C right side). Additional support for this

notion was achieved by immunoprecipitation analysis. The proteins that generated L-Cys and

SeCys DES activity bands were fully pulled down after the proteins were immunoprecipitated by

Arabidopsis OAS-TL B specific antibody (kindly provided by Prof. Dr. S. Kopriva, University

of Cologne, Germany) that also cross reacts with OASTLA and C isoforms (Fig. 4D left and

right side respectively).

A Native-SDS with L-Cys DES activity

Col-0 | oastlA | oastlB | oastlC
Con Se 40 | Con Se 40 | Con Se 40 | Con Se 40

Upper → ; middle → ; lower →

	Col-0 Con	Col-0 Se 40	oastlA Con	oastlA Se 40	oastlB Con	oastlB Se 40	oastlC Con	oastlC Se 40
Upper	1.00	1.94	1.74	1.68	1.35	1.93		
middle	1.00	1.39	1.16	1.37			1.27	1.92
lower	1.00	1.12			1.24	1.54	1.02	1.19

B Native-SDS with SeCys DES activity

	Col-0 Con	Col-0 Se 40	oastlA Con	oastlA Se 40	oastlB Con	oastlB Se 40	oastlC Con	oastlC Se 40
	1	2.00	1.65	2.24			1.12	1.56
	1	1.44			1.34	1.90	2.04	2.15

C Immunoblotting in SDS — Bands taken from Native SDS L-Cys DES activity (Low., Mid., Up.); Crude protein Col-0. Bands taken from Native SDS SeCys DES activity (Lower, Upper); Crude protein Col-0. kDa 34→ . OASTL C, OASTL B, OASTL A.

D Immunoprecipitation — Native-SDS with L-Cys | Native-SDS with SeCys. Crude protein Col-0 | OAS-TL B antibody (+, -). Crude protein Col-0 | OAS-TL B antibody (+, -).

Figure 4. NATIVE-SDS on L-Cys and SeCys desulfhydrase activities in WT, oastlA, oastlB and oastlC KO mutants supplemented with or without selenate for 14 days. Effect of 40 µM selenate treatment on L-Cysteine (A) and SeCys (B) degradation activities. C. Immunodetection of the sliced WT activity bands from 4A are in left insert and the sliced WT activity bands sliced from 4B are in the right insert. D. Desulfhydrase activity of WT crude protein extract without immunoprecipitation (left lane) with immunoprecipitation in the presence (+) or absence (-) of specific antibodies. L-cys and L-Secys desulfhydrase activities are left and right inserts respectively. Red arrows indicate activity band.

Finally, the identification of tripsinized unique peptides in the three sliced activity bands after the fractionation of WT crude protein extract (Fig.4A and B) was performed. The lowest sliced activity bands of L-Cys and L-SeCys desulfhydrase activities revealed 23 and 20 unique peptides overlapping 66% and 65% of the Arabidopsis OAS-TL A protein sequence, respectively (Supplemental Table S1A-C). Together with the OAS-TL A peptides, a lower amount of 14 and 9 peptides of OASTLB were identified in the lowest sliced L-Cys and SeCys desulfhydrase activity bands, respectively (Supplemental Table S1A-C). In the middle activity band, sliced from L-Cys and SeCys desulfhydrase activity, 23 and 18 unique peptides were identified showing 58% and 60.5% overlap of OAS-TL B protein sequence, respectively (Supplemental Table S1A-C). Additionally, 18 and 15 peptides of OASTLA proteins in the L-Cys and SeCys desulfhydrase activity bands and 6 unique peptides of OASTLC protein were identified in L-Cys desulfhydrase activity. The upper band sliced from L-Cys desulfhydrase activity revealed 13 specific unique peptides with 45% coverage of OASTLC protein. Six and 11 unique peptides of OASTLA and B proteins respectively were identified as well (Supplemental Table S1A-C). The

lack of a full separation between the 3 OASTLs activity bands is likely the result of the relatively high identity between the three proteins [ca 70 to 80% identity (Supplemental Fig.S4)] that prevented the clear complete separation in-spite of the 16 cm fractionation length by SDS-PAGE. Yet, based on the identification by the use of the three mutants exposed to in-gel activity, immunoblot and immunoprecipitation by the specific antibodies employed (Fig. 4), as well as the majority of the overlapping unique peptides (Supplemental Table S1), we can conclude that OASTLA, B and C are localized to the highest, second-highest and the lowest activity bands, respectively of L-Cys and SeCys desulfhydrase activity.

Significantly, only two unique peptides of DES1 were identified in the lower and middle bands and none in the most upper band of L-Cys desulfhydrase activity bands (Supplemental Table S1). Considering that the number of theoretic tryptic peptides in DES1 is similar to that of OASTLs (estimated by the use of Expasy: https://web.expasy.org/peptide_mass/), these results indicate that DES1 is a less abundant protein or it undergoes ionization. Yet, since the semi quantitation was made by calculating the peak area of each peptide, whereas to overcome differential ionization, the area of the protein was calculated as the average of the three most intense peptides from each protein; the data indicate that OASTL A, B and C are more abundant than DES1 protein (Supplemental Table S1D).

The less abundant DES1 protein in the activity bands does not necessarily indicate decreased L-Cys and L-SeCys DES activity by the WT DES1 compared to the OASTLs. Yet, firstly, the kinetic activity revealed a major significant decrease of 80 and 65% in the Cys-desulfhydrase activity of the cytosolic *oastlA* leaves in plants grown in control or selenate stressed plants, respectively compared to WT (Fig.3C). Additionally, activity bands, additive to OASTL, A, B and C, that could be attributed to DES1, were not noticed, neither above, nor below and none were noticed at the position of the migrated OASTL A, B and C when the activity bands were absent in the related impaired mutants under control and selenate stressed conditions (Fig. 4A and B). Taken together these results (Supplemental Table S1A-D, Figs 3C, 4A, B) strongly support the notion that OAS-TL A, B and C proteins play a major role in L-Cys and L-SeCys degradation in *Arabidopsis*. In the presence of selenate, plants impaired in OASTL A and B exhibit higher catabolic processing rates of chlorophyll and protein degradation, the result of enhanced sulfur metabolism interference in the mutants compared with WT.

DISCUSSION

Impairment in OASTLA and OASTLB leads to higher sensitivity to selenate

Non-specific incorporation of seleno-amino acids SeCys and SeMet into proteins instead of cysteine and methionine is thought to be a main cause for Se induced toxicity in many plants (Sors et al., 2005; Terry et al., 2000). Yet, lower Se in protein fraction was evident in *apr2* KO mutants exhibiting stronger sensitivity than WT to the Se treatment (Grant et al., 2011). Similarly, determination of the total Se and Se content in protein fraction revealed a significantly lower rate in *oastlA* and *oastlB* mutants, compared with WT and *oastlC* (Fig. 1E and Supplemental Fig. S2B). This indicates that the accumulated Se in *oastlA* and *oastlB* mutant proteins is not the main cause for their stronger sensitivity to Se compared with WT.

The uptake and assimilation of Se has similar effects on gene expression as with the case of S-starvation, resulting in greater S uptake and assimilation rates by plants [(White, 2018) and references therein]. Plants exposed to high Se level exhibited symptoms such as growth inhibition and chlorosis (Grant et al., 2011; Sabbagh and Van Hoewyk, 2012; Van Hoewyk, 2018). Interestingly, the overexpression of OASTL in Arabidopsis and tobacco increased tolerance to the toxicity of Cd stress by increasing thiol availability (Domínguez-Solís et al., 2004). Among the three *Arabidopsis* OASTL mutants grown for 2 weeks with 40μM selenate, a decrease in biomass accumulation and chlorophyll content compared to WT was noticed in *oastlA* and *oastlB* but not in *oastlC* mutant (Fig.1). This is likely since the cytosolic and chloroplastic OASTL A and B respectively, exhibit significantly higher activity rates compared with the mitochondrion localized OASTL C, the latter being responsible for only 5-12% of OASTL activity in WT plants (Heeg et al., 2008; Kuske et al., 1996). The results demonstrate that dissimilar to WT and impairment in OASTLC, in the absence of active OASTLA and OASTLB, the selenate interference in S assimilation pathway leads to a stronger decrease in organic S, S in protein fraction and total glutathione, and has a more negative effect on the remaining chlorophyll level and the resulting biomass accumulation than the expected positive effect of lower selenium content in the organic fraction (Figs 1 and 2).

The free cysteine levels were similar in WT and the mutants exposed to selenate (Fig.2B), whereas the Cys generation activity rate differed as shown here in response to selenate (Fig.2A). This was also shown previously in an O-acetylserine (thiol) lyase reduced isoform by RNA interference in potato, as well as in *Arabidopsis* KO mutants (Heeg et al., 2008;

Riemenschneider et al., 2005), indicating the feasibility of an additional Cys source affecting free

Cys level. The source can be protein degradation, indicated by the lower soluble protein level in

oastlA and *oastlB* KO mutants compared to WT and *oastlC* treated with selenate (Fig. 3A). The

protein degradation is the result of selenate induced stress, being stronger in *oastlA* and *B* as

indicated by the lower remaining chlorophyll (Fig. 1D), organic-S, and S in protein fraction (Fig.

2D and E).

Sulfite is a strong nucleophile that should be tightly regulated to avoid its deleterious

reaction with a wide variety of cellular components (Brychkova et al., 2007; Lang et al., 2007;

Brychkova et al., 2013; Yarmolinsky et al., 2013; Brychkova et al., 2015). The increased sulfite

level in *oastlA* and *oastlB* mutants compared to WT (Fig. 3D) is a deleterious effect of a futile S-

starvation response, the result of the low organic sulfur, total glutathione as well as low S in the

protein fraction (Fig. 2A,C,D,E) in *oastlA* and *oastlB* mutants treated with selenate. APR activity

increase in *oastlA* and *oastlB* leaves that caused the enhanced sulfite level in these 2 mutants is a

response to S-starvation (Lee et al., 2011) and the followed sulfide level increase, the result of

higher SiR activity in these 2 mutants, further supports the notion of selenate induced S-

starvation. The futileness of such response is firstly because of the waste of reducing agents and

ATP (see in Supplementary Fig. S1) in the process ending with excess sulfide, the result of the

inability of *oastlA* and *oastlB* plants to assimilate sulfide to L-Cys generation. Additionally, the

sulfite level increase, shown before to induce oxidative stress (Brychkova et al., 2012;

Yarmolinsky et al., 2014), indeed led to oxidative stress (Supplementary Fig. S5) that had an

additive deleterious effect on the growth of *oastlA* and *oastlB* mutant plants. The oxidative stress

could further induce APR to generate more sulfite as demonstrated before (Bick et al., 2001;

Koprivova et al., 2008). Notably, the levels of hydrogen peroxide (H_2O_2), the carbonyl toxic

malondialdehyde (MDA), the product of lipid peroxidation by prolonged oxidative stress (Tian

et al., 2017) and the antioxidant anthocyanins (Gould et al., 2002; Xu et al., 2017), were

enhanced (on FW, and except H_2O_2, on soluble protein basis as well) in the selenate treated

plants compared to control plants. MDA and anthocyanin levels were significantly higher in

oastlA and *oastlB* compared to the selenate treated WT and *oastlC* mutant, whereas, H_2O_2 was

significantly higher in the 3 mutants compared to WT treated with selenate. The enhancement in

H_2O_2, MDA and anthocyanins levels in *oastlA* and *oastlB* mutants (Supplemental Fig. S5) is

indicative of the higher oxidative stress level in these plants as shown before (Gould et al., 2002;

Srivastava et al., 2017; Xu et al., 2017; Yarmolinsky et al., 2014). Taken together the results
indicate that the decrease in *oastlA* and *oastlB* biomass accumulation is likely the result of
selenate induced S-starvation responses causing waste of reducing agents and ATP, as well as
sulfite induced oxidative stress.

OASTL A and B act as major L-Cys and L-SeCys desulfhydrases in *Arabidopsis*
The similar free cysteine level in WT and *oastlA* and *B* mutants, compared with the lower
cysteine generating activity rate of OASTL and the lower soluble protein level evident in the
mutants exposed to selenate (Figs 2A, B, Supplemental S3A), led us to examine a role for
OASTL proteins in degrading L-Cys and SeCys to sulfide and selenide respectively.
Additionally, albeit in a small rate comparing to the biosynthesis by OASTLs, the degrading
activity of the active enzymes is likely beneficial to plants, decreasing the levels of L-SeCys
(compare the rate in Fig. 2A to the rate in Fig.3C) and its incorporation into proteins.
L-Cys desulfhydrase activity in WT showed 3 activity bands, where the upper band intensity
was weaker than that of the other two bands. L-SeCys desulfhydrase exhibited only 2 activity
bands, with the absence of the upper band likely the result of detection limits (Fig. 4A and B).
Importantly, fractionated proteins extracted from *oastlA* and *oastlB* KO mutants exhibited in-gel
activity absent of the lower and middle activity band, respectively with both activity types; the L-
Cys and SeCys DES, whereas the most upper band was seen only when L-Cys was used as the
substrate and only in WT, *oastlA* and *oastlB* but not in *oastlC* (Fig. 4A and B). These results
indicate that the activity bands with the highest, second-highest and the lowest mobility are of
OASTLA, B and C, respectively. Additionally, the results show that the three OASTLs have the
capacity of L-SeCys and L-Cys DES activities. This was further supported by the use of specific
antibodies to identify the sliced activity bands by Western blot after further SDS fractionation
(Fig. 4C), as well as by immunoprecipitation assay (Fig. 4D). Finally it was supported by the
identification of the tripsinized unique peptides (Supplemental Table S1) in the three sliced WT
activity bands (Fig.4A and B).
Importantly, DES1 was shown to degrade L-Cys, estimated by the use of the *des1* null mutants
to be responsible for up to 14% of the total L-Cys desulfhydrase activity compared to WT plants
grown in plated on MS medium (Álvarez et al., 2010). This suggests the existence of additional
source/s that can contribute to the majority (ca 86%) of H_2S production by degrading L-Cys.

Notably, the impairment of *oastlA* and *oastlB* mutants led to a significantly decreased sulfide

production activity rate by L-Cys desulfhydrase [ca 76% and 18% reduction respectively

(Fig3C)] compared to WT, indicating the important role of the OASTLs in degrading L-Cys.

Intriguingly, only very few unique peptides of DES1 were identified in the major activity bands,

the highest and second-highest mobility bands, of L-Cys desulfhydrase activity and none in the

SeCys desulfhydrase activity bands, indicating that the OASTLs are more abundant proteins than

DES1 (Supplemental Table S1). Taking together i) the lower abundancy of DES1 compared to

the OASTLs and ii) the significant decrease in the Cys-desulfhydrase kinetic activity of the

oastlA leaves in plants grown in control or selenate stressed plants compared to WT (Fig.3C) as

well as iii) the absence of activity bands, additive to OASTL A, B and C activity bands that

could be attributed to DES1 in control and selenate stressed condition (Fig. 4A and B), the result

strongly support the notion that OASTL A, B and C play a major role in L-Cys and L-SeCys

degradation in *Arabidopsis*.

CONCLUSIONS

Using *Arabidopsis* O-acetylserine-(thiol) lyases mutants, impaired in the biosynthesis of cysteine

(Cys) from the substrates sulfide and O-acetylserine, it was shown that the absence of active

OASTLA and OASTLB, confers higher reduced growth, lower remaining chlorophyll, lower

sulfur in protein fraction and lower total glutathione than in selenate stressed WT and *oastlC*

mutant as the result of the interference of selenate with the sulfate reduction pathway.

Additionally it was shown that the absence of active OASTLA or OASTLB confers higher

sensitivity to selenate compared to WT or the absence of active OASTLC. Further, it was shown

here that the cytosolic OASTLA, chloroplat localized OASTLB and the mitochondrion localized

OASTLC has an important role in desulfhydrase activity degrading not only L-Cys but also

SeCys to sulfide and selenide, respectively.

MATERIALS AND METHODS

Plant Material, Growth Conditions and Selenium treatment

Arabidopsis thaliana var. Columbia wild type (*WT*), *oastlA (Salk_072213), oastlB (Salk*

_021183), oastlC (Salk_000860) and des1 (SALK_205358) lines were used for Selenium

treatment. SALK_103855 (*des1*) was employed for in-gel DES activity and transcript

expression. All the mutants were derived from the Columbia (*Col*) ecotype. Experiments were

carried out in the growth room at Sde Boqer Campus, Ben-Gurion University of the Negev,

under 14 h light/10 h darkness, 22˚C and 75–85% relative humidity under photosynthetically

active radiation of 150 µmol m^{-2} sec^{-1} as described in Brychkova et al., 2007.

Seeds were germinated and grown in standard 90 mm Petri dishes on solid ½ MS

medium for 9 days (2 days in dark and 7 days in growth room). The seedlings were transferred to

large Petri dishes of 155x30 mm diameter and height respectively, supplied with ½ MS medium

(Murashige and Skoog, 1962) supplemented with or without 40µM of Na_2SeO_4. All treatments

were performed in six replicates. The weight of shoot biomass accumulation was determined 14

days after the treatment and results were expressed as average plant growth rate in mg plant^{-15}

RNA Isolation, cDNA Preparation, and transcript analyses by Real-time PCR

To quantify the transcripts using quantitative reverse transcriptase-polymerase chain reaction

(qRT-PCR), total RNA was prepared using the Aurum^{TM} total RNA Mini Kit (Bio-Rad,

Hercules, CA) according to the manufacturer's instructions. The cDNA was prepared in a 10 µl

volume containing 350 ng of plant total RNA that was reverse-transcribed with an iScript^{TM}

cDNA Synthesis Kit using modified MMLV-derived reverse transcriptase (Bio-Rad), a blend of

oligo-d(T) and random hexamer primers, according to the manufacturer's instructions. The

generated cDNA was diluted 10 times and the quantitative analysis of transcripts was performed

employing the sets of primers from both sites of insertions shown in Supplemental Table S2 as

previously described (Brychkova et al., 2007). The transcripts level detected by Real-time PCR

were normalized using *ACTIN2* (At3g18780) and *ELONGATION FACTOR1-α* (At5g60390) and

compared to WT grown under the control condition. The data are presented as relative

expression (means ± SE, *n* = 4). All PCR fragments were sequenced for verification.

Chlorophyll, MDA, H_2O_2 and Anthocyanin determination

For detection of remaining content of chlorophyll (%) the leaf discs (7 mm diameter)

were immersed in 90% ethanol and incubated at 4°C for 2 d in the dark. Absorbance of the

extracted chlorophyll was measured at 652 nm, and total chlorophyll was estimated (Ritchie,

2006). MDA levels were determined as described by Srivastava et.al., 2017.

For detecting H_2O_2, frozen leaves were extracted in 50 mM P buffer (pH 7.5) at a ratio of

1:8 (w/v) and centrifuged (Eppendorf 5417R) twice at 18,000g for 20 min. The reaction mixture

for detecting H_2O_2 consisted of 0.85 mm 4-aminoantipyrine, 3.4 mm 3,5-dichloro-2-

hydroxobenzene sulfonate, 4.5 U mL^{-1} HRP in 2 mL of 50 mM P buffer (pH 7.5) in the presence

or absence of 2 mM tungstic acid and 100 μM DPI as described in Yesbergenova et al. 2005.

Absorbance was measured after 5 min at 515 nm as described above.

The anthocyanin content was determined based on a modification of protocols described

by Laby et al. (2000) and Kant et al. (2006). Approximately 100 mg of fresh plants grown in ½

MS medium were crushed in 600 μL methanol acidified with 1% HCl. The extract was

centrifuged for 10 sec at 4000g. Five hundred μL of double distilled water were added to the

collected sand, mixture was gently vortexed and then 700 μL chloroform was added and mixed

for 20 sec followed by centrifugation at 4000 g for 2 min. The total anthocyanin in the aqueous

phase was determined by detecting the optical density (OD) at A530 and A657 nm. The amount

of anthocyanin was calculated by subtracting the A657 from the A530 (Laby et al., 2000).

Enzyme activity measurements

Protein extraction, desalting, concentration determination and kinetic assays for APR, SO, SiR,

OAS-TL and DES activities were assayed and expressed as described before; OAS-TL and SiR

activity in nmol Cys mg^{-1} protein min^{-1}, APR and SO in nmol sulfite mg^{-1} protein min^{-1} and

DES activity in nmol H_2S mg^{-1} protein min^{-1} (Brychkova et al., 2012a; Brychkova et al., 2012b;

Kurmanbayeva et.al., 2017). In brief, APR activity employing APS as substrate was detected

using the sulfite-specific fuchsine colorimetric detection method. SO activity was measured as

the disappearance of sulfite. The desalted protein extracts were treated with 1 mM tungstate for

30 min at 4°C to inhibit SO activity. SiR activity was estimated by the coupled SiR/OASTL

assay with the addition of NADPH and tungstic acid. The resultant generated Cys was detected

as described before (Gaitonde, 1967; Brychkova et al., 2012b). OAS-TL activity was initiated by

incubating 75 μL Reaction mixture (RM) containing 25 mM OAS, 1 mM DTT, 1.42 mM

Pyridoxal-phosphate (PLP), 60 mM Na_2S and 200 mM P-buffer (pH 7.5) with 25 μL of protein

extract for 10 min at 30°C. Generated Cysteine was detected at 560 nm by a spectrophotometer

(Gaitonde, 1967).

DES activity was detected as H_2S formation in the presence of L-cysteine. The assay solution contained 0.1 M Tris-HCl, pH 9.0, 2.5 mM dithiothreitol, 0.8 mM L-cysteine and 10 μg of desalted protein in a total volume of 0.2 mL. After incubation at 37 °C for 30 min, H_2S was detected according to Bloem et al., 2004 with 30 mM $FeCl_3$ dissolved in 1.2 N HCl and 20 mM N,N-dimethyl-p-phenylenediamine dihydrochloride dissolved in 7.2 N HCl. The formation of methylene blue was measured at 670 nm. Na_2S*9H_2O was used as a standard.

Protein extraction, fractionation for in gel DES activity

Whole protein from Arabidopsis seedlings was extracted as described by Sagi and Fluhr, 2001. Concentrations of total soluble protein in the resulting supernatant were determined according to Bradford, 1976. Native-SDS PAGE was carried out as follows: samples were subjected to a Bio-Rad Protean II xi slab cell (Bio-Rad, Richmond, CA, USA), with a discontinuous buffer system (Laemmli, 1970) in 7.5% (w/v) polyacrylamide separating gels and 4% (w/v) stacking gels. Native-SDS PAGE was carried out using 1.5 mm thick slabs loaded with 200 μg of proteins. Regeneration of the active proteins after native-SDS PAGE was carried out by removal of the SDS as described by Sagi and Fluhr, 2001 with slight modification: by shaking the gel for 1 h in 10 mM Tris-HCl buffer (pH 7.8) solution (65 ml buffer per ml of gel) containing 2 mM EDTA.

DES activity was detected using a modification of an in-gel visualization protocol for H_2S (Manchenko, 2002). Lead acetate was employed to detect the generated H_2S and possibly H_2Se, producing dark brown lead sulfide/selenide bands. The reaction solution contained 0.15 M Tris–HCl, pH 8.5, 1 mM Dithiothreitol (DTT), 50 mM β-mercaptoethanol, 5 mM L-cysteine or 2.5 mM SelenoCystine, 0.1 mM pyridoxal 5-phosphate (PLP), 0.4 mM lead acetate. The reaction was stopped by immersion of the gel in double-distilled water.

Western Blot and Immunoprecipitation Analysis

Western blot proteins were separated by SDS–PAGE carried out in 12.5% polyacrylamide gels and were blotted to polyvinylidene difluoride membranes (Immun-Blot membranes, Bio-Rad). The blotted proteins were subjected to immuno-detection with antibodies raised against Arabidopsis chloroplastic OAS-TL C (diluted1:1000) (kindly provided by Prof.

R.Hell, University of Heidelberg, Heidelberg) and followed by secondary antibodies as described

by Heeg et al., 2008.

For the immunoprecipitation assay 200 µg of protein from *WT* were incubated with 80 µl

of the OAS-TL B (diluted1:5000) (kindly provided by Prof. Dr. S. Kopriva, University of

Cologne, Köln) antibodies in Tris-buffer saline (TBS) for 30 min at room temperature and then

kept at 4°C for overnight. Protein extract that had not been mixed with antibody was employed

as control. The mixture and the control solutions were incubated with 120 µl of Protein G

Agarose at 4°C for 2h with continuous shaking and then centrifuged at 10,000g for 5 min,

followed by removal of supernatant for analysis by the Native-SDS DES assay.

Determination of selenium and sulfur-containing metabolites

To determine the total selenium content, 100 mg of the dried and powdered leaves were

placed in glass tubes, digested with 70% $HN0_3$, heated at 220^OC for 4 hours and quantified by

inductively coupled plasma emission spectrometry (ICP-AES) (Kalra, 1997).

Se in protein fraction: 400 g fresh weight was extracted in 2 mL buffer containing 100

mM NaCl, 50 mM Tris HCl (pH 7.5), 0.5% (v/v) Triton X-100, 1 mM DTT, and 0.1 mM PMSF.

The homogenate was cleared by centrifugation (7,500g for 10 min). A small sample was taken

for protein determination, and the volume of the extract was measured. The proteins in the

extract were precipitated by adding TCA to a final concentration of 15% (w/v). The mixture was

incubated on ice for 30 min and then centrifuged for 20 min at 7,000g at 4°C. The pellet was

washed with ice-cold acetone, dried, and dissolved in 1 mL of concentrated nitric acid. After acid

digestion, the Se was determined by ICP (Pilon-Smits et al., 1999).

Selenate detection was done according to (Schiavon et al., 2012) with slight

modifications. The frozen tissues (200 mg) were ground in liquid nitrogen, and then 5 mL of

distilled water was added. The obtained extracts were filtered (0.22µm, Millipore) and analyzed

for sulfate and selenate concentrations by Ion chromatography using a Dionex IonPac AS19

4x250mm column.

Organic S and Se were calculated by subtracting the sum of sulfate/selenate, sulfite and

sulfide from the total sulfur/selenium content.

Determination of sulfite, free Cys and total glutathione was performed as described

(Brychkova et al., 2013; Yarmolinsky et al., 2014). Sulfide was extracted (1:4, w/v) in 0.1 M

Tris–HCl, pH 8.0 buffer in the presence of 0.5 M sodium sulfate to minimize sulfite self-
oxidation to sulfate. After centrifugation at 1800 g for 15 min, the supernatant was further de-
proteinized employing Sephadex G-25 Column (Pd-10, GE Healthcare) loaded with the same
extraction solution. Immediately after the separation through the column 600 µl of 2% cadmium
acetate were added to the 600 µl of the sample and kept on ice. The levels of sulfide were
detected by 40 µl NN-Dimethyl-1,4-phenylen-diammonium-dichloride dissolved in 7.2 N HCl
and 4 µl FeCl3 dissolved in 1.2 N HCl from 240 µl of preincubated sample with 2% cadmium
acetate (Siegel, 1965). Sulfide was detected at 625 nm.

Protein sequencing

To identify the proteins participating in the cysteine desulfhydrase activity, the activity bands
from the in-gel activity of DES were sliced from the native gel, and fractionated with 12.5%
SDS-PAGE (Fig. 4). Thereafter the proteins were stained by Coomassie Brilliant Blue, and the
stained bands were excised from the gel, trypsinized and the resulting peptides were separated by
HPLC and analyzed by LC-MS/MS on Q-Exactive (Thermo) at The Smoler Protein Research
Center (Technion University, Haifa, Israel).

All the identified peptides were filtered with high confidence, top rank, mass accuracy, and a
minimum of 2 peptides. High confidence peptides passed the 1% FDR threshold. (*FDR =false
discovery rate, is the estimated fraction of false positives in a list of peptides). Semi quantitation
was done by calculating the peak area of each peptide. The area of the protein is the average of
the three most intense peptides from each protein. Analysis of peptide sequences was performed
by employing Proteome Discoverer™ Software ver. 1.4.1.14 (Thermo Fisher Scientific Inc.
https://www.thermofisher.com/order/catalog/product/ IQLAAEGABSFAKJMAUH).

Accession Numbers

Sequence data for this article can be found in the Arabidopsis Genome Initiative or
GenBank/EMBL databases under the following accession numbers: At4g14880 *(OASTL A)*,
At2g43750 (*OASTL B*), At3g59760 (*OASTL C*), AT5G28030 (*DES1*), AT3G18780 (*ACTIN 2*),
AT5G60390 (*EF 1-α*)

Supplemental Figures

Supplemental Figure S1. Schematic model of Sulfur (S) and Selenium (Se) metabolism in Arabidopsis plants [after White (2015)].

Supplemental Figure S2. OASTLs protein and transcript expressions and Se containing compounds in WT, *oastlA, oastlB* and *oastlC KO* mutants supplemented with or without selenate for 14 days.

Supplemental Figure S3. S/Se metabolism activities and DES1 transcript in WT and *Atdes1* KO mutants supplemented with or without selenate for 14 days.

Supplemental Figure S4. Multiple sequence alignment of OASTL A, B and C proteins of in *Arabidopsis thaliana*.

Supplemental Figure S5. H_2O_2 MDA and anthocyanin level in WT, *oastlA, oastlB* and *oastlC* KO mutants supplemented with or without selenate for 14 days.

Supplemental Tables

Supplemental Table S1. Identified and overlapped unique peptides from L-Cys and L-SeCys desulfhydrase activity. **Supplemental Table S2.** List of gene primers used for quantitative real-time PCR.

ACKNOWLEDGMENT:

We thank Prof. Dr. S. Kopriva (University of Cologne, Köln) and Prof. R.Hell, (Universität Heidelberg, Heidelberg) for providing OAS-TL A and B antibodies.

FIGURE LEGENDS

Figure 1. The effect of 40 μM selenate treatment on growth parameters and Se in protein fraction in WT, *oastlA, oastlB* and *oastlC KO* mutants of *Arabidopsis thaliana*. A. Phenotype of plants grown for 14 days under control (left panel) and 40μM selenate treated (right panel) conditions. The weight of plant biomass accumulation (B) and relative weight to their own controls (C) supplemented with or without selenate for 14 days. The values are means ± SE (n = 15). Remaining chlorophyll level (D) and Se in protein fraction (E). The values are means ± SE (n = 4). Values denoted with different letters are significantly different according to the T- test analyses, $P < 0.05$ (JMP 8.0 software, http://www.jmp.com/). Different uppercase

letters indicate significant differences between control and Se treatment of each genotype. Asterisks indicate significant differences between *WT* and *oas-tlA, oas-tlB, oastlC* KO plants subjected to the same treatment.

Figure 2. Cys biosynthesis and S containing compounds in WT, *oastlA, oastlB* and *oastlC* KO mutants supplemented with or without selenate for 14 days. The effect of 40 μM selenate treatment on OAS-TL activity (A), Cysteine level (B), Total glutathione (C), Total organic S (D) and S in protein fraction (E). The values are means ± SE (n = 4). Values denoted with different letters are significantly different according to T-test analyses, P < 0.05 (JMP 8.0 software, http://www.jmp.com/). Different uppercase letters indicate significant differences between control and Se treatment of each genotype. Asterisks indicate significant differences between WT and the KO plants *oastlA, oastlB and oastlC* subjected to the same treatment.

Figure 3. Soluble protein level and S metabolism components in WT, *oastlA, oastlB* and *oastlC* KO mutants supplemented with or without selenate for 14 days. The effect of 40 μM selenate treatment on soluble protein content (A), transcript expression of DES1 (B), L-Cysteine desulfhydrase (DES) activity (C), sulfide (D) and sulfite levels (E), Sulfite oxidase (SO) activity (F), Adenosine 5′-phosphosulfate reductase (APR) activity (G) and Sulfite reductase (SiR) activity (H). Transcript level of DES1 was detected by quantitative reverse transcription-PCR using *ACTIN2* as the housekeeping transcript for normalization. The relative expression in the various genotypes was analyzed using the normalized WT control as reference. The values are means ± SE (n = 4). Values denoted with different letters are significantly different according to T-test analyses, P < 0.05 (JMP 8.0 software, http://www.jmp.com/). Different uppercase letters indicate significant differences between control and Se treatment of each genotype. Asterisks indicate significant differences between WT and *oastlA, oastlB, oastlC* plants subjected to the same treatment.

Figure 4. NATIVE-SDS on L-Cys and SeCys desulfhydrase activities in WT, oastlA, oastlB and oastlC KO mutants supplemented with or without selenate for 14 days. Effect of 40 μM selenate treatment on L-Cysteine (A) and SeCys (B) degradation activities. C. Immunodetection of the sliced WT activity bands from 4A are in left insert and the sliced WT activity bands sliced from 4B are in the right insert. D. Desulfhydrase activity of WT crude protein extract without immunoprecipitation (left lane) with immunoprecipitation in the presence (+) or absence (-) of specific antibodies. L-cys and L-Secys desulfhydrase activities are left and right inserts respectively. Red arrows indicate activity band.

Supplemental Figures

Supplemental Figure S1. Schematic model of Sulfur (S) and Selenium (Se) metabolism in Arabidopsis plants [after White (2015)]. ATP sufurylase (ATPS) catalyzes the adenylation of sulfate/selenate to adenosine 5′-phosphosulfate (APS)/adenosine phosphoselenate (APSe) using ATP as an electron donor. Then, APS/APSe is reduced by the plastidic enzyme APS reductase (APR) to sulfite/selenite in the presence of two molecules of reduced glutathione, which acts as an electron donor. The generated sulfite/selenite can be reduced to sulfide/selenide by the Sulfite Reductase (SiR) employing 3 molecules of reduced ferredoxin. Alternatively, selenite can also be reduced non-enzymatically by glutathione (GSH) to selenide. The sulfide/selenide together with O-acetyl-L-Serine (OAS) are the substrates for Cysteine (Cys)/Selenocysteine (SeCys) biosynthesis catalyzed by O-acetylserine-(thiol) lyase (OAS-TL).

Supplemental Figure S2. OASTLs protein and transcripts expression, as well as Se containing compounds in *WT, AtoastlA, AtoastlB* and *AtoastlC* KO mutants supplemented with or without selenate for 14 days. Effect of 40μM Selenate treatment on OASTLA, OASTLB and OASTLC level in WT and the various mutants analysed by Western blot employing specific antibody (A), total Se (B), selenate (C) and transcripts analyses of *OASTLA* (D), *OASTLB* (E) and *OASTLC* (F). Transcript levels of *OASTLA, OASTLB* and *OASTLC* were detected in the various genotypes by quantitative reverse transcription-PCR using *ACTIN2* as housekeeping transcript for normalization. The relative expression of each normalized *OASTL* in the various genotypes was analyzed using WT control as the reference. The values are means ± SE (n = 4). Values denoted with different letters are significantly different according to T-test analyses, P < 0.05 (JMP 8.0 software, http://www.jmp.com/). Different uppercase letters indicate significant differences between control and Se treatment of each genotype. Asterisks indicate significant differences between WT and *oas-tlA, oastlB, oastlC* KO plants subjected to the same treatment. The specific antibody kindly provided by Prof. Dr. S. Kopriva, University of Cologne, Germany.

Supplemental Figure S3. S/Se metabolism activities and DES1 transcript in WT and *Atdes1* KO mutants supplemented with or without selenate for 14 days. L-Cys/SeCys NATIVE-SDS degradation activities in the presence and absence of L-Cys/SeCys or BME (A), *DES1* transcripts expression with primers constructed upstream and downstream along the transcript length (B), OAS-TL (C) and DES kinetic activities (D), as well as, L-Cysteine (left side) and SeCys (right side) in-gel desulfhydrase activities in WT and *Atdes1* SALK_205358 leaves (E). DES1 transcript expression and in-gel L-Cysteine desulfhydrase activity in WT and *Atdes1* SALK_103855 (F). Transcript level of *Des1* was detected by quantitative reverse transcription-PCR using *ACTIN2* as the housekeeping transcript for normalization.

The relative expression in the *Atdes1* with the 2 coupled primers was analyzed using the normalized WT

control as reference. In-gel L-Cysteine desulfhydrase activities were employed after protein fractionation

in SDS gel performed in Protean® II xi (www.bio-rad.com) in A and E and mini Protean3 (www.bio-

rad.com) in F. The values are means ± SE (n = 4). Values denoted with different letters are significantly

different according to T-test analyses, $P < 0.05$ (JMP 8.0 software, http://www.jmp.com/). Different

uppercase letters indicate significant differences between control and Se treatment of each genotype.

Different lower-case letters indicate differences between genotypes within the same primer. Asterisks

indicate significant differences between WT and *des1* KO plants subjected to the same treatment

Supplemental Figure S4. Multiple sequence alignment of OASTL A, B and C proteins in *Arabidopsis*

thaliana. The percentage of sequence identity between the proteins is presented below.

Supplemental Figure S5. Hydrogen peroxide (H_2O_2), malondialdehyde (MDA) and anthocyanin level in

WT, *AtoastlA, AtoastlB* and *AtoastlC* KO mutants supplemented with or without selenate for 14 days. The

effect of 40µM Selenate treatment on H_2O_2 (A), MDA (B) and anthocyanin (C) levels calculated on FW

(left) and soluble protein basis (right). The values are means ± SE (n = 4). Values denoted with different

letters are significantly different according to T-test analyses, $P < 0.05$ (JMP 8.0 software,

http://www.jmp.com/). Different uppercase letters indicate significant differences between control and Se

treatment of each genotype.

Supplemental Tables

Supplemental Table S1. Identified and overlapped unique peptides from L-Cys and L-SeCys

desulfhydrase activity. **A**. List of the number coverage percentage **B**. Identified and overlapped unique

peptides from L-Cys desulfhydrase activity. **C**. Identified and overlapped unique peptides from L-SeCys

desulfhydrase activity. D. Abundance of OASTL A, B and C as compared to DES1 protein at the lower,

middle and upper activity bands. The semi quantitation was made by calculating the peak area of

each peptide, whereas to overcome differential ionization, the area of the protein was calculated

as the average of three most intense peptides from each protein. The calculated area of the DES

protein is marked in red. The activity bands (as shown in Fig. 4A and B) were sliced and fractionated in

12.5 % SDS-PAGE. Further the bands were digested by trypsin and analyzed by LC-MS/MS as described

in Materials and Methods. Blue color – overlapped areas. The presented data are representative of 3

independent experiments with similar results. *(see below).

Supplemental Table S2. List of gene primers used for quantitative real-time PCR.

REFERENCES

Álvarez C, Calo L, Romero LC, García I, Gotor C (2010) An O-acetylserine (thiol) lyase
homolog with L-cysteine desulfhydrase activity regulates cysteine homeostasis in
Arabidopsis. *Plant Physiology* **152**: 656-669

Barroso C, Vega J, Gotor C (1995) A new member of the cytosolic O-acetylserine(thiol)lyase
gene family in Arabidopsis thaliana. FEBS Letters **363**: 1-5

Bick J-A, Setterdahl AT, Knaff DB, Chen Y, Pitcher LH, Zilinskas BA, Leustek T (2001)
Regulation of the plant-type 5 '-adenylyl sulfate reductase by oxidative stress.
Biochemistry **40**: 9040-9048

Bloem E, Riemenschneider A, Volker J, Papenbrock J, Schmidt A, Salac I, Haneklaus S,
Schnug E (2004) Sulphur supply and infection with Pyrenopeziza brassicae influence L-
cysteine desulphydrase activity in Brassica napus L. *Journal of Experimental Botany* **55**:
2305-2312

Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram
quantities of protein utilizing the principle of protein-dye binding. *Analytical*
biochemistry **72**: 248-254

Brychkova G, Xia Z, Yang G, Yesbergenova Z, Zhang Z, Davydov O, Fluhr R, Sagi M
(2007) Sulfite oxidase protects plants against sulfur dioxide toxicity. *The Plant journal:*
for cell and molecular biology **50**: 696-709

Brychkova G, Yarmolinsky D, Fluhr R, Sagi M (2012a) The determination of sulfite levels
and its oxidation in plant leaves. Plant Sci **190**: 123-130

Brychkova G, Yarmolinsky D, Ventura Y, Sagi M (2012b) A novel in-gel assay and an
improved kinetic assay for determining in vitro sulfite reductase activity in plants. Plant
and Cell Physiology **53**: 1507-1516

Brychkova G, Grishkevich V, Fluhr R, Sagi M (2013) An essential role for tomato sulfite
oxidase and enzymes of the sulfite network in maintaining leaf sulfite homeostasis. *Plant*
physiology **161**: 148-164

Brychkova G, Yarmolinsky D, Batushansky A, Grishkevich V, Khozin-Goldberg I, Fait A,
Amir R, Fluhr R, Sagi M (2015) Sulfite oxidase activity is essential for normal sulfur,
nitrogen and carbon metabolism in tomato leaves. *Plants* **4**: 573-605

Domínguez-Solís JR, López-Martín MC, Ager FJ, Ynsa MD, Romero LC, Gotor C (2004)
Increased cysteine availability is essential for cadmium tolerance and accumulation in
Arabidopsis thaliana. *Plant Biotechnology Journal* **2**: 469-476

Fisher B, Yarmolinsky D, Abdel-Ghany S, Pilon M, Pilon-Smits EA, Sagi M, Van Hoewyk
D (2016) Superoxide generated from the glutathione-mediated reduction of selenite
damages the iron-sulfur cluster of chloroplastic ferredoxin. *Plant physiology and*
biochemistry **106**: 228-235

Gaitonde M (1967) A spectrophotometric method for the direct determination of cysteine in the
presence of other naturally occurring amino acids. *Biochemical Journal* **104**: 627

Grant K, Carey NM, Mendoza M, Schulze J, Pilon M, Pilon-Smits EA, Van Hoewyk D
(2011) Adenosine 5′-phosphosulfate reductase (APR2) mutation in Arabidopsis
implicates glutathione deficiency in selenate toxicity. *Biochemical Journal* **438**: 325-335

Gould KS, Mckelvie J, Markham KR (2002) Do anthocyanins function as antioxidants in leaves? Imaging of H_2O_2 in red and green leaves after mechanical injury. *Plant Cell and Environment* **25**: 1261 – 1269

Gupta M, Gupta S (2017) An overview of selenium uptake, metabolism, and toxicity in plants. *Frontiers in Plant Science* **7**: 2074

Heeg C, Kruse C, Jost R, Gutensohn M, Ruppert T, Wirtz M, Hell R (2008) Analysis of the Arabidopsis O-acetylserine (thiol) lyase gene family demonstrates compartment-specific differences in the regulation of cysteine synthesis. *The Plant Cell* **20**: 168-185

Hesse H, Lipke J, Altmann T, Höfgen R (1999) Molecular cloning and expression analyses of mitochondrial and plastidic isoforms of cysteine synthase (O-acetylserine (thiol) lyase) from Arabidopsis thaliana. Amino Acids **16**: 113-131

Hesse H, Hoefgen R (2003) Molecular aspects of methionine biosynthesis. *Trends in plant science* **8**: 259-262

Holdorf MM, Owen HA, Lieber SR, Yuan L, Adams N, Dabney-Smith C, Makaroff CA (2012) Arabidopsis ETHE1 encodes a sulfur dioxygenase that is essential for embryo and endosperm development. *Plant physiology* **160**: 226-236

Jin Z, Sun L, Yang G, Pei Y (2018) Hydrogen sulfide regulates energy production to delay leaf senescence induced by drought stress in Arabidopsis. Frontiers in Plant Science **9**: 1722

Kalra Y (1997). "Handbook of reference methods for plant analysis," CRC press.

Kant S, Kant P, Raveh E, Barak S (2006) Evidence that differential gene expression between the halophyte, Thellungiella halophila, and Arabidopsis thaliana is responsible for higher levels of the compatible osmolyte proline and tight control of Na+ uptake in T. halophila. Plant, Cell Environ **29**: 1220–1234

Khan MS, Haas FH, Samami AA, Gholami AM, Bauer A, Fellenberg K, Reichelt M, Hänsch R, Mendel RR, Meyer AJ (2010) Sulfite reductase defines a newly discovered bottleneck for assimilatory sulfate reduction and is essential for growth and development in Arabidopsis thaliana. *The Plant Cell* **22**: 1216-1231

Koprivova A, North KA, Kopriva S (2008) Complex signaling network in regulation of adenosine 5′-phosphosulfate reductase by salt stress in Arabidopsis roots. Plant Physiology **146**: 1408-1420

Krüßel L, Junemann J, Wirtz M, Birke H, Thornton JD, Browning LW, Poschet G, Hell R, Balk J, Braun H-P (2014) The mitochondrial sulfur dioxygenase ETHYLMALONIC ENCEPHALOPATHY PROTEIN1 is required for amino acid catabolism during carbohydrate starvation and embryo development in Arabidopsis. *Plant physiology* **165**: 92-104

Kurmanbayeva A, Bekturova A, Srivastava S, Soltabayeva A, Asatryan A, Ventura Y, Khan MS, Salazar O, Fedoroff N, Sagi M (2017) Higher novel L-Cys degradation activity results in lower organic-S and biomass in Sarcocornia than the related saltwort, Salicornia. *Plant physiology* **175**: 272-289

Kuske CR, Hill KK, Guzman E, Jackson PJ (1996) Subcellular location of O-acetylserine sulfhydrylase isoenzymes in cell cultures and plant tissues of Datura innoxia Mill. *Plant physiology* **112**: 659-667

Laby RJ, Kincaid MS, Kim D, Gibson SI (2000) The Arabidopsis sugar-insensitive mutants sis4 and sis5 are defective in abscisic acid synthesis and response. Plant J **23**: 587–596

Lang C, Popko J, Wirtz M, Hell R, Herschbach C, Kreuzwieser J, Rennenberg H, Mendel RR, Haensch R (2007) Sulphite oxidase as key enzyme for protecting plants against sulphur dioxide. Plant, Cell & Environment **30**: 447-455

Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *nature* **227**: 680

Lee BR, Koprivova A, Kopriva S (2011) The key enzyme of sulfate assimilation, adenosine 5′-phosphosulfate reductase, is regulated by HY5 in Arabidopsis. The Plant Journal **67**: 1042-1054

Lima LW, Pilon-Smits EA, Schiavon M (2018) Mechanisms of selenium hyperaccumulation in plants: A survey of molecular, biochemical and ecological cues. *Biochimica et Biophysica Acta (BBA)-General Subjects* **1862**: 2343-2353

Manchenko GP (2002). "Handbook of detection of enzymes on electrophoretic gels," CRC press.

Mroczek-Zdyrska M, Wójcik M (2012) The influence of selenium on root growth and oxidative stress induced by lead in Vicia faba L. minor plants. *Biological trace element research* **147**: 320-328

Murashige T, Skoog F (1962) A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiologia plantarum* **15**: 473-497

Papp LV, Lu J, Holmgren A, Khanna KK (2007) From selenium to selenoproteins: synthesis, identity, and their role in human health. *Antioxidants & redox signaling* **9**: 775-806

Pilon-Smits EA, Hwang S, Lytle CM, Zhu Y, Tai JC, Bravo RC, Chen Y, Leustek T, Terry N (1999) Overexpression of ATP sulfurylase in Indian mustard leads to increased selenate uptake, reduction, and tolerance. *Plant Physiology* **119**: 123-132

Pilon-Smits EA, Quinn CF (2010). Selenium metabolism in plants. *In* "Cell biology of metals and nutrients", pp. 225-241. Springer.

Pilon-Smits EA, Womg M (2012). Plant selenium metabolism–genetic manipulation, phytotechnological applications, and ecological implications. *In* "Environmental contamination: Health risks and ecological restoration", pp. 293-311. CRC Press Boca Raton, FL.

Ravanel S, Gakière B, Job D, Douce R (1998) The specific features of methionine biosynthesis and metabolism in plants. *Proceedings of the National Academy of Sciences* **95**: 7805-7812

Riemenschneider A, Riedel K, Hoefgen R, Papenbrock J, Hesse H (2005) Impact of reduced O-acetylserine (thiol) lyase isoform contents on potato plant metabolism. *Plant physiology* **137**: 892-900

Ritchie RJ (2006) Consistent sets of spectrophotometric chlorophyll equations for acetone, methanol and ethanol solvents. Photosynth Res **89**: 27–41

Sabbagh M, Van Hoewyk D (2012) Malformed selenoproteins are removed by the ubiquitin–proteasome pathway in Stanleya pinnata. *Plant and Cell Physiology* **53**: 555-564

Sagi M, Fluhr R (2001) Superoxide production by plant homologues of the gp91phox NADPH oxidase. Modulation of activity by calcium and by tobacco mosaic virus infection. *Plant Physiology* **126**: 1281-1290

Schiavon M, Moro I, Pilon-Smits EA, Matozzo V, Malagoli M, Dalla Vecchia F (2012) Accumulation of selenium in Ulva sp. and effects on morphology, ultrastructure and antioxidant enzymes and metabolites. *Aquatic toxicology* **122**: 222-231

Schiavon M, Pilon M, Malagoli M, Pilon-Smits EA (2015) Exploring the importance of sulfate transporters and ATP sulphurylases for selenium hyperaccumulation—a comparison of Stanleya pinnata and Brassica juncea (Brassicaceae). *Frontiers in plant science* **6**: 2

Seko Y, Saito Y, Kitahara J, Imura N (1989). Active oxygen generation by the reaction of selenite with reduced glutathione in vitro. *In* "Selenium in biology and medicine", pp. 70-73. Springer.

Siegel LM (1965) A direct microdetermination for sulfide. *Analytical biochemistry* **11**: 126-132

Sors TG, Ellis DR, Na GN, Lahner B, Lee S, Leustek T, Pickering IJ, Salt DE (2005) Analysis of sulfur and selenium assimilation in Astragalus plants with varying capacities to accumulate selenium. *The Plant Journal* **42**: 785-797

Srivastava S, Brychkova G, Yarmolinsky D, Soltabayeva A, Samani T, Sagi M (2017) Aldehyde Oxidase 4 plays a critical role in delaying silique senescence by catalyzing aldehyde detoxification. *Plant physiology* **173**: 1977-1997

Terry N, Zayed A, De Souza M, Tarun A (2000) Selenium in higher plants. *Annual review of plant biology* **51**: 401-432

Tian M, Hui M, Thannhauser TW, Pan S, Li L (2017) Selenium-induced toxicity is counteracted by sulfur in broccoli (Brassica oleracea L. var. italica). *Frontiers in plant science* **8**: 1425

Van Hoewyk D (2018) Defects in endoplasmic reticulum-associated degradation (ERAD) increase selenate sensitivity in Arabidopsis. *Plant signaling & behavior* **13**: e1171451

Van Hoewyk D, Garifullina GF, Ackley AR, Abdel-Ghany SE, Marcus MA, Fakra S, Ishiyama K, Inoue E, Pilon M, Takahashi H (2005) Overexpression of AtCpNifS enhances selenium tolerance and accumulation in Arabidopsis. *Plant Physiology* **139**: 1518-1528

White PJ (2015) Selenium accumulation by plants. *Annals of botany* **117**: 217-235

White PJ (2018) Selenium metabolism in plants. *Biochimica et Biophysica Acta (BBA)-General Subjects* **1862**: 2333-2342

White PJ, Bowen HC, Parmaguru P, Fritz M, Spracklen W, Spiby R, Meacham M, Mead A, Harriman M, Trueman L (2004) Interactions between selenium and sulphur nutrition in Arabidopsis thaliana. *Journal of Experimental Botany* **55**: 1927-1937

Wirtz M, Hell R (2006) Functional analysis of the cysteine synthase protein complex from plants: structural, biochemical and regulatory properties. *Journal of plant physiology* **163**: 273-286

Xu Z, Mahmood K, Rothstein SJ (2017) ROS induces anthocyanin production via late biosynthetic genes and anthocyanin deficiency confers the hypersensitivity to ROS-generating stresses in Arabidopsis. *Plant and Cell Physiology* **58**: 1364–1377

Yarmolinsky D, Brychkova G, Kurmanbayeva A, Bekturova A, Ventura Y, Khozin-Goldberg I, Eppel A, Fluhr R, Sagi M (2014) Impairment in Sulfite Reductase Leads to Early Leaf Senescence in Tomato Plants. *Plant Physiology* **165**: 1505–1520

Yarmolinsky D, Brychkova G, Fluhr R, Sagi M (2013) Sulfite reductase protects plants against sulfite toxicity. *Plant physiology* **161**: 725-743

Yesbergenova Z, Yang G, Oron E, Soffer D, Fluhr R, Sagi M (2005) The plant Mo-hydroxylases aldehyde oxidase and xanthine dehydrogenase have distinct reactive oxygen species signatures and are induced by drought and abscisic acid. *The Plant Journal* **42**: 862-876

Figure 1. The effect of 40 μM selenate treatment on growth parameters and Se in protein fraction in WT, *oastlA, oastlB* and *oastlC KO* mutants of *Arabidopsis thaliana*. A. Phenotype of plants grown for 14 days under control (left panel) and 40μM selenate treated (right panel) conditions. The weight of plant biomass accumulation (B) and relative weight to their own controls (C) supplemented with or without selenate for 14 days. The values are means ± SE (n = 15). Remaining chlorophyll level (D) and Se in protein fraction (E). The values are means ± SE (n = 4). Values denoted with different letters are significantly different according to the T- test analyses, P < 0.05 (JMP 8.0 software, http://www.jmp.com/). Different uppercase letters indicate significant

Figure 2. Cys biosynthesis and S containing compounds in WT, *oastlA, oastlB* and *oastlC* KO mutants supplemented with or without selenate for 14 days. The effect of 40 μM selenate treatment on OAS-TL activity (A), Cysteine level (B), Total glutathione (C), Total organic S (D) and S in protein fraction (E). The values are means ± SE (n = 4). Values denoted with different letters are significantly different according to T-test analyses, P < 0.05 (JMP 8.0 software, http://www.jmp.com/). Different uppercase letters indicate significant differences between control and Se treatment of each genotype. Asterisks indicate significant differences between WT and the KO plants *oastlA, oastlB and oastlC* subjected to the same treatment.

Figure 3. Soluble protein level and S metabolism components in WT, *oastlA, oastlB* and *oastlC* KO mutants supplemented with or without selenate for 14 days. The effect of 40 μM selenate treatment on soluble protein content (A), transcript expression of DES1 (B), L-Cysteine desulfhydrase (DES) activity (C), sulfide (D) and sulfite levels (E), Sulfite oxidase (SO) activity (F), Adenosine 5′-phosphosulfate reductase (APR) activity (G) and Sulfite reductase (SiR) activity (H). Transcript level of *DES1* was detected by quantitative reverse transcription-PCR using *ACTIN2* as the housekeeping transcript for normalization. The relative expression in the various genotypes was analyzed using the normalized WT control as reference. The values are means ± SE (n = 4). Values denoted with different letters are significantly different according to T-test analyses, P < 0.05 (JMP 8.0 software, http://www.jmp.com/). Different uppercase letters indicate significant differences between control and Se treatment of each genotype. Asterisks indicate significant differences between WT and *oastlA, oastlB, oastlC* plants subjected to the same

Figure 4. NATIVE-SDS on L-Cys and SeCys desulfhydrase activities in WT, oastlA, oastlB and oastlC KO mutants supplemented with or without selenate for 14 days. Effect of 40 μM selenate treatment on L-Cysteine (A) and SeCys (B) degradation activities. C. Immunodetection of the sliced WT activity bands from 4A are in left insert and the sliced WT activity bands sliced from 4B are in the right insert. D. Desulfhydrase activity of WT crude protein extract without immunoprecipitation (left lane) with immunoprecipitation in the presence (+) or absence (-) of specific antibodies. L-cys and L-Secys desulfhydrase activities are left and right inserts respectively. Red arrows indicate activity band.

Parsed Citations

Álvarez C, Calo L, Romero LC, García I, Gotor C (2010) An O-acetylserine (thiol) lyase homolog with L-cysteine desulfhydrase activity regulates cysteine homeostasis in Arabidopsis. Plant Physiology 152: 656-669
Google Scholar: Author Only Title Only Author and Title

Barroso C, Vega J, Gotor C (1995) A new member of the cytosolic O-acetylserine(thiol)lyase gene family in Arabidopsis thaliana. FEBS Letters 363: 1-5
Google Scholar: Author Only Title Only Author and Title

Bick J-A, Setterdahl AT, Knaff DB, Chen Y, Pitcher LH, Zilinskas BA, Leustek T (2001) Regulation of the plant-type 5 '-adenylyl sulfate reductase by oxidative stress. Biochemistry 40: 9040-9048
Google Scholar: Author Only Title Only Author and Title

Bloem E, Riemenschneider A, Volker J, Papenbrock J, Schmidt A, Salac I, Haneklaus S, Schnug E (2004) Sulphur supply and infection with Pyrenopeziza brassicae influence L-cysteine desulphydrase activity in Brassica napus L. Journal of Experimental Botany 55: 2305-2312
Google Scholar: Author Only Title Only Author and Title

Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Analytical biochemistry 72: 248-254
Google Scholar: Author Only Title Only Author and Title

Brychkova G, Xia Z, Yang G, Yesbergenova Z, Zhang Z, Davydov O, Fluhr R, Sagi M (2007) Sulfite oxidase protects plants against sulfur dioxide toxicity. The Plant journal: for cell and molecular biology 50: 696-709
Google Scholar: Author Only Title Only Author and Title

Brychkova G, Yarmolinsky D, Fluhr R, Sagi M (2012a) The determination of sulfite levels and its oxidation in plant leaves. Plant Sci 190: 123-130
Google Scholar: Author Only Title Only Author and Title

Brychkova G, Yarmolinsky D, Ventura Y, Sagi M (2012b) A novel in-gel assay and an improved kinetic assay for determining in vitro sulfite reductase activity in plants. Plant and Cell Physiology 53: 1507-1516
Google Scholar: Author Only Title Only Author and Title

Brychkova G, Grishkevich V, Fluhr R, Sagi M (2013) An essential role for tomato sulfite oxidase and enzymes of the sulfite network in maintaining leaf sulfite homeostasis. Plant physiology 161: 148-164
Google Scholar: Author Only Title Only Author and Title

Brychkova G, Yarmolinsky D, Batushansky A, Grishkevich V, Khozin-Goldberg I, Fait A, Amir R, Fluhr R, Sagi M (2015) Sulfite oxidase activity is essential for normal sulfur, nitrogen and carbon metabolism in tomato leaves. Plants 4: 573-605
Google Scholar: Author Only Title Only Author and Title

Domínguez-Solís JR, López-Martín MC, Ager FJ, Ynsa MD, Romero LC, Gotor C (2004) Increased cysteine availability is essential for cadmium tolerance and accumulation in Arabidopsis thaliana. Plant Biotechnology Journal 2: 469-476
Google Scholar: Author Only Title Only Author and Title

Fisher B, Yarmolinsky D, Abdel-Ghany S, Pilon M, Pilon-Smits EA, Sagi M, Van Hoewyk D (2016) Superoxide generated from the glutathione-mediated reduction of selenite damages the iron-sulfur cluster of chloroplastic ferredoxin. Plant physiology and biochemistry 106: 228-235
Google Scholar: Author Only Title Only Author and Title

Gaitonde M (1967) A spectrophotometric method for the direct determination of cysteine in the presence of other naturally occurring amino acids. Biochemical Journal 104: 627
Google Scholar: Author Only Title Only Author and Title

Grant K, Carey NM, Mendoza M, Schulze J, Pilon M, Pilon-Smits EA, Van Hoewyk D (2011) Adenosine 5′-phosphosulfate reductase (APR2) mutation in Arabidopsis implicates glutathione deficiency in selenate toxicity. Biochemical Journal 438: 325-335
Google Scholar: Author Only Title Only Author and Title

Gould KS, Mckelvie J, Markham KR (2002) Do anthocyanins function as antioxidants in leaves? Imaging of H2O2 in red and green leaves after mechanical injury. Plant Cell and Environment 25: 1261 – 1269
Google Scholar: Author Only Title Only Author and Title

Gupta M, Gupta S (2017) An overview of selenium uptake, metabolism, and toxicity in plants. Frontiers in Plant Science 7: 2074
Google Scholar: Author Only Title Only Author and Title

Heeg C, Kruse C, Jost R, Gutensohn M, Ruppert T, Wirtz M, Hell R (2008) Analysis of the Arabidopsis O-acetylserine (thiol) lyase gene family demonstrates compartment-specific differences in the regulation of cysteine synthesis. The Plant Cell 20: 168-185
Google Scholar: Author Only Title Only Author and Title

Hesse H, Lipke J, Altmann T, Höfgen R (1999) Molecular cloning and expression analyses of mitochondrial and plastidic isoforms of cysteine synthase (O-acetylserine (thiol) lyase) from Arabidopsis thaliana. Amino Acids 16: 113-131

Google Scholar: Author Only Title Only Author and Title

Hesse H, Hoefgen R (2003) Molecular aspects of methionine biosynthesis. Trends in plant science 8: 259-262
  Google Scholar: Author Only Title Only Author and Title

Holdorf MM, Owen HA, Lieber SR, Yuan L, Adams N, Dabney-Smith C, Makaroff CA (2012) Arabidopsis ETHE1 encodes a sulfur dioxygenase that is essential for embryo and endosperm development. Plant physiology 160: 226-236
  Google Scholar: Author Only Title Only Author and Title

Jin Z, Sun L, Yang G, Pei Y (2018) Hydrogen sulfide regulates energy production to delay leaf senescence induced by drought stress in Arabidopsis. Frontiers in Plant Science 9: 1722
  Google Scholar: Author Only Title Only Author and Title

Kalra Y (1997). "Handbook of reference methods for plant analysis," CRC press.

Kant S, Kant P, Raveh E, Barak S (2006) Evidence that differential gene expression between the halophyte, Thellungiella halophila, and Arabidopsis thaliana is responsible for higher levels of the compatible osmolyte proline and tight control of Na+ uptake in T. halophila. Plant, Cell Environ 29: 1220–1234
  Google Scholar: Author Only Title Only Author and Title

Khan MS, Haas FH, Samami AA, Gholami AM, Bauer A, Fellenberg K, Reichelt M, Hänsch R, Mendel RR, Meyer AJ (2010) Sulfite reductase defines a newly discovered bottleneck for assimilatory sulfate reduction and is essential for growth and development in Arabidopsis thaliana. The Plant Cell 22: 1216-1231
  Google Scholar: Author Only Title Only Author and Title

Koprivova A, North KA, Kopriva S (2008) Complex signaling network in regulation of adenosine 5′-phosphosulfate reductase by salt stress in Arabidopsis roots. Plant Physiology 146: 1408-1420
  Google Scholar: Author Only Title Only Author and Title

Krüßel L, Junemann J, Wirtz M, Birke H, Thornton JD, Browning LW, Poschet G, Hell R, Balk J, Braun H-P (2014) The mitochondrial sulfur dioxygenase ETHYLMALONIC ENCEPHALOPATHY PROTEIN1 is required for amino acid catabolism during carbohydrate starvation and embryo development in Arabidopsis. Plant physiology 165: 92-104
  Google Scholar: Author Only Title Only Author and Title

Kurmanbayeva A, Bekturova A, Srivastava S, Soltabayeva A, Asatryan A, Ventura Y, Khan MS, Salazar O, Fedoroff N, Sagi M (2017) Higher novel L-Cys degradation activity results in lower organic-S and biomass in Sarcocornia than the related saltwort, Salicornia. Plant physiology 175: 272-289
  Google Scholar: Author Only Title Only Author and Title

Kuske CR, Hill KK, Guzman E, Jackson PJ (1996) Subcellular location of O-acetylserine sulfhydrylase isoenzymes in cell cultures and plant tissues of Datura innoxia Mill. Plant physiology 112: 659-667
  Google Scholar: Author Only Title Only Author and Title

Laby RJ, Kincaid MS, Kim D, Gibson SI (2000) The Arabidopsis sugar-insensitive mutants sis4 and sis5 are defective in abscisic acid synthesis and response. Plant J 23: 587–596
  Google Scholar: Author Only Title Only Author and Title

Lang C, Popko J, Wirtz M, Hell R, Herschbach C, Kreuzwieser J, Rennenberg H, Mendel RR, Haensch R (2007) Sulphite oxidase as key enzyme for protecting plants against sulphur dioxide. Plant, Cell & Environment 30: 447-455
  Google Scholar: Author Only Title Only Author and Title

Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. nature 227: 680
  Google Scholar: Author Only Title Only Author and Title

Lee BR, Koprivova A, Kopriva S (2011) The key enzyme of sulfate assimilation, adenosine 5′-phosphosulfate reductase, is regulated by HY5 in Arabidopsis. The Plant Journal 67: 1042-1054
  Google Scholar: Author Only Title Only Author and Title

Lima LW, Pilon-Smits EA, Schiavon M (2018) Mechanisms of selenium hyperaccumulation in plants: A survey of molecular, biochemical and ecological cues. Biochimica et Biophysica Acta (BBA)-General Subjects 1862: 2343-2353
  Google Scholar: Author Only Title Only Author and Title

Manchenko GP (2002). "Handbook of detection of enzymes on electrophoretic gels," CRC press.

Mroczek-Zdyrska M, Wójcik M (2012) The influence of selenium on root growth and oxidative stress induced by lead in Vicia faba L. minor plants. Biological trace element research 147: 320-328
  Google Scholar: Author Only Title Only Author and Title

Murashige T, Skoog F (1962) A revised medium for rapid growth and bio assays with tobacco tissue cultures. Physiologia plantarum 15: 473-497
  Google Scholar: Author Only Title Only Author and Title

Papp LV, Lu J, Holmgren A, Khanna KK (2007) From selenium to selenoproteins: synthesis, identity, and their role in human health. Antioxidants & redox signaling 9: 775-806

Google Scholar: Author Only Title Only Author and Title

Pilon-Smits EA, Hwang S, Lytle CM, Zhu Y, Tai JC, Bravo RC, Chen Y, Leustek T, Terry N (1999) Overexpression of ATP sulfurylase in Indian mustard leads to increased selenate uptake, reduction, and tolerance. Plant Physiology 119: 123-132

Pilon-Smits EA, Quinn CF (2010). Selenium metabolism in plants. In "Cell biology of metals and nutrients", pp. 225-241. Springer.

Pilon-Smits EA, Womg M (2012). Plant selenium metabolism–genetic manipulation, phytotechnological applications, and ecological implications. In "Environmental contamination: Health risks and ecological restoration", pp. 293-311. CRC Press Boca Raton, FL.

Ravanel S, Gakière B, Job D, Douce R (1998) The specific features of methionine biosynthesis and metabolism in plants. Proceedings of the National Academy of Sciences 95: 7805-7812

Riemenschneider A, Riedel K, Hoefgen R, Papenbrock J, Hesse H (2005) Impact of reduced O-acetylserine (thiol) lyase isoform contents on potato plant metabolism. Plant physiology 137: 892-900

Ritchie RJ (2006) Consistent sets of spectrophotometric chlorophyll equations for acetone, methanol and ethanol solvents. Photosynth Res 89: 27–41

Sabbagh M, Van Hoewyk D (2012) Malformed selenoproteins are removed by the ubiquitin–proteasome pathway in Stanleya pinnata. Plant and Cell Physiology 53: 555-564

Sagi M, Fluhr R (2001) Superoxide production by plant homologues of the gp91phox NADPH oxidase. Modulation of activity by calcium and by tobacco mosaic virus infection. Plant Physiology 126: 1281-1290

Schiavon M, Moro I, Pilon-Smits EA, Matozzo V, Malagoli M, Dalla Vecchia F (2012) Accumulation of selenium in Ulva sp. and effects on morphology, ultrastructure and antioxidant enzymes and metabolites. Aquatic toxicology 122: 222-231

Schiavon M, Pilon M, Malagoli M, Pilon-Smits EA (2015) Exploring the importance of sulfate transporters and ATP sulphurylases for selenium hyperaccumulation-a comparison of Stanleya pinnata and Brassica juncea (Brassicaceae). Frontiers in plant science 6: 2

Seko Y, Saito Y, Kitahara J, Imura N (1989). Active oxygen generation by the reaction of selenite with reduced glutathione in vitro. In "Selenium in biology and medicine", pp. 70-73. Springer.

Siegel LM (1965) A direct microdetermination for sulfide. Analytical biochemistry 11: 126-132

Sors TG, Ellis DR, Na GN, Lahner B, Lee S, Leustek T, Pickering IJ, Salt DE (2005) Analysis of sulfur and selenium assimilation in Astragalus plants with varying capacities to accumulate selenium. The Plant Journal 42: 785-797

Srivastava S, Brychkova G, Yarmolinsky D, Soltabayeva A, Samani T, Sagi M (2017) Aldehyde Oxidase 4 plays a critical role in delaying silique senescence by catalyzing aldehyde detoxification. Plant physiology 173: 1977-1997

Terry N, Zayed A, De Souza M, Tarun A (2000) Selenium in higher plants. Annual review of plant biology 51: 401-432

Tian M, Hui M, Thannhauser TW, Pan S, Li L (2017) Selenium-induced toxicity is counteracted by sulfur in broccoli (Brassica oleracea L. var. italica). Frontiers in plant science 8: 1425

Van Hoewyk D (2018) Defects in endoplasmic reticulum-associated degradation (ERAD) increase selenate sensitivity in Arabidopsis. Plant signaling & behavior 13: e1171451

Van Hoewyk D, Garifullina GF, Ackley AR, Abdel-Ghany SE, Marcus MA, Fakra S, Ishiyama K, Inoue E, Pilon M, Takahashi H (2005) Overexpression of AtCpNifS enhances selenium tolerance and accumulation in Arabidopsis. Plant Physiology 139: 1518-1528

White PJ (2015) Selenium accumulation by plants. Annals of botany 117: 217-235

White PJ (2018) Selenium metabolism in plants. Biochimica et Biophysica Acta (BBA)-General Subjects 1862: 2333-2342
Google Scholar: Author Only Title Only Author and Title

White PJ, Bowen HC, Parmaguru P, Fritz M, Spracklen W, Spiby R, Meacham M, Mead A, Harriman M, Trueman L (2004) Interactions between selenium and sulphur nutrition in Arabidopsis thaliana. Journal of Experimental Botany 55: 1927-1937
Google Scholar: Author Only Title Only Author and Title

Wirtz M, Hell R (2006) Functional analysis of the cysteine synthase protein complex from plants: structural, biochemical and regulatory properties. Journal of plant physiology 163: 273-286
Google Scholar: Author Only Title Only Author and Title

Xu Z, Mahmood K, Rothstein SJ (2017) ROS induces anthocyanin production via late biosynthetic genes and anthocyanin deficiency confers the hypersensitivity to ROS-generating stresses in Arabidopsis. Plant and Cell Physiology 58: 1364–1377
Google Scholar: Author Only Title Only Author and Title

Yarmolinsky D, Brychkova G, Kurmanbayeva A, Bekturova A, Ventura Y, Khozin-Goldberg I, Eppel A, Fluhr R, Sagi M (2014) Impairment in Sulfite Reductase Leads to Early Leaf Senescence in Tomato Plants. Plant Physiology 165: 1505–1520
Google Scholar: Author Only Title Only Author and Title

Yarmolinsky D, Brychkova G, Fluhr R, Sagi M (2013) Sulfite reductase protects plants against sulfite toxicity. Plant physiology 161: 725-743
Google Scholar: Author Only Title Only Author and Title

Yesbergenova Z, Yang G, Oron E, Soffer D, Fluhr R, Sagi M (2005) The plant Mo-hydroxylases aldehyde oxidase and xanthine dehydrogenase have distinct reactive oxygen species signatures and are induced by drought and abscisic acid. The Plant Journal 42: 862-876
Google Scholar: Author Only Title Only Author and Title