Slipped strand mispairing in the gene encoding sialidase NanH3 in

***Gardnerella* spp.**

Shakya P. Kurukulasuriya, Mo H. Patterson, Janet E. Hill[#]

Department of Veterinary Microbiology, University of Saskatchewan, Saskatoon, Canada

Running head: Slipped-strand mispairing in *Gardnerella* sialidase

#Address correspondence to Janet E. Hill, Janet.Hill@usask.ca

Abstract

Cell wall proteins with sialidase activity are involved in carbohydrate assimilation, adhesion to mucosal surfaces, and biofilm formation. *Gardnerella* spp. inhabit the human vaginal microbiome and encode up to three sialidase enzymes, two of which are suspected to be cell wall associated. Here we demonstrate that the gene encoding extracellular sialidase NanH3 is found almost exclusively in *G. piotii* and closely related *Gardnerella* genome sp. 3, and its presence correlates with sialidase positive phenotype in a collection of 112 *Gardnerella* isolates. The *nanH3* gene sequence includes a homopolymeric repeat of cytosines that varies in length within cell populations, indicating that this gene is subject to slipped-strand mispairing, a mechanisms of phase variation in bacteria. Variation in the length of the homopolymer sequence results in encoding of either the full length sialidase protein or truncated peptides lacking the sialidase domain due to introduction of reading-frame shifts and premature stop codons. Phase variation in NanH3 may be involved in immune evasion or modulation of adhesion to host epithelial cells, and formation of biofilms characteristic of the vaginal dysbiosis known as bacterial vaginosis.

Introduction

Bacterial vaginosis (BV) is a condition that is characterized by altered composition of the vaginal microbiota and occurs when the healthy microbiota is replaced by an overgrowth of mixed aerobic and anaerobic species, including *Gardnerella* spp. (1, 2). An abundance of *Gardnerella* spp. is often found in cases of symptomatic BV, although they are also found in healthy women with no clinical signs or symptoms of BV (3). *Gardnerella* spp. can be resolved into four subgroups based on cpn60 barcodes sequencing (4) or whole genome sequencing (5). Recently, the description of *Gardnerella vaginalis* was amended and three new species were defined within the genus *Gardnerella*: *G. leopoldii*, *G. swidsinskii*, and *G. piotii* (6)*. G. piotii* and *G. vaginalis* correspond to cpn60 subgroup B and C, respectively. *G. leopoldii* and G. *swidsinskii* were previously grouped together as subgroup A based on cpn60 sequences of the isolates available at that time. Isolates belonging to cpn60 subgroup D corresponded to three distinct genome species: genome species 8, 9 and 10 (7).

In addition to the characteristic change in microbiota, and elevated pH, sialidase activity in vaginal fluid is a diagnostic marker of BV (8, 9). Sialidase enzymes cleave the glycosidic linkages of sialic acids from terminal glycans including vaginal mucins, immunoglobulin A molecules, and epithelial cell surface glycoproteins (10, 11). Activity on the latter may be related to the adhesion of bacteria to epithelial cells; the initial step in biofilm formation (12). Previously, we have demonstrated that sialidase activity is almost exclusively confined to *G. piotii* and *Gardnerella* genome species 3 (cpn60 subgroup B) (13).

A putative sialidase gene, *nanH1* (sialidase A) was identified in *Gardnerella* spp. and was initially thought to be the gene responsible for sialidase activity (14). Although *nanH1* appears to be present in all sialidase positive strains, it is also found in sialidase activity negative strains (13).

This observation combined with the lack of signal peptide on the NanH1 proteins suggests that

this protein is an intracellular enzyme, likely involved in manipulation of nutritional substrates as

is the case in many bacteria (15). The discrepancy between *nanH1* presence and enzyme activity

led to the recognition of two additional sialidase genes (*nanH2* and *nanH3*) in some *Gardnerella*

isolates (16), which are unevenly distributed among *Gardnerella* spp. (17). Based on biochemical

studies, Robinson et al. (16) concluded that NanH2 and NanH3 are the proteins responsible for

sialidase activity detected in vaginal fluid. The presence of signal peptides and/or C-terminal

transmembrane alpha-helices in some of the identified sequences further suggested that they are

secreted and tethered to the cell, although some proportion may subsequently be released into the

extracellular environment due to proteolytic activity (16). Cell wall anchored proteins with

sialidase activity in other species have been found to be involved in adhesion to mucosal surfaces

and biofilm initiation as well as carbohydrate assimilation (18–21).

Interestingly, in our examination of genome sequences of our collection of *Gardnerella*

isolates, we noted that *nanH3* contains a homopolymeric tract of cytosine residues. Genomic

regions that contain short, homogenous or heterogenous repeats are susceptible to slipped-strand

mispairing in which the length of the repeat region can change with each replication (22). The

result of this modulation is phase variation: a reversible process in which the expression of the

encoded protein can be rapidly switched on and off (23). Phase variation in cell surface proteins

can result in immune evasion and alteration of biofilm phenotypes (24, 25).

Here we determined the distribution of genes encoding sialidases NanH2 and NanH3 in

the context of newly reclassified *Gardnerella* spp. and demonstrated that *nanH3* is subject to

slipped-strand mispairing.

Methods

Protein domain identification and sequence alignment

DNA and protein sequence alignments were performed with Clustal Omega (EMBL_EBI) and NCBI BLAST (basic local alignment search tool). InterproScan (https://www.ebi.ac.uk/interpro/) and SignalP were used to predict the location of functional domains and signal peptides (26).

Bacterial strains and culture conditions

Gardnerella strains (n = 112) from a previously described culture collection were used in the study (13). Sialidase activity for all isolates had been determined previously (13) using a quantitative assay and fluorogenic substrate 2'-(4-methylumbelliferyl)-α-D-N-acetylneuraminic acid sodium salt hydrate (11), and whole genome sequences for 36 of these isolates were previously determined (27). Complete strain information and sequence accessions are provided in Table S1.

Gardnerella isolates were grown on Columbia Sheep Blood Agar plates (BBL, Becton, Dickinson and Company, Sparks, MD, USA) at 37°C for 48 hours with anaerobic BD GasPak EZ (Becton, Dickinson and Company, Sparks, MD, USA). For broth cultures, a few colonies from the plate were collected with a 10 μl inoculation loop and used to inoculate NYC III (ATCC 1685 medium; per litre: 2.4g HEPES, 15 g Proteose peptone, 3.8 g Yeast extract, 5 g NaCl, 5 g Glucose). Broth cultures were incubated at 37°C for 48 hours in anaerobic conditions.

PCR screen for *nanH3*

Genomic DNA was purified from broth cultures using a modified salting-out procedure (28). All DNA extracts were initially tested by PCR for the universal cpn60 barcode to confirm the quality of the DNA.

To screen isolates for the presence of *nanH3*, PCR primers were designed based on

multiple sequence alignments of 15 *nanH3* sequences obtained from the Integrated Microbial

Genomes database (https://img.jgi.doe.gov/). Degenerate primers were designed to account for

sequence variability within the gene sequence and to amplify a product of 375 bp in length

(JH0684: 5'-GTT GTA GAR CTT TCT GAT GG-3', JH0685: 5'-YRY TAT TAT CGC CCT CAT

ATA-3'). PCR reactions contained 1 × PCR Buffer (0.2 M Tris-HCl at pH 8.4, 0.5 M KCl), 2.5

μM $MgCl_2$, 0.40 μM dNTP, 0.20 μM forward primer, 0.20 μM reverse primer, 2 U Taq DNA

Polymerase, ultrapure water and 2 μl of template DNA in a final volume of 50 μl. PCR reactions

were conducted using the following thermocycling parameters in a Mastercycler Pro 6321

(Eppendorf AG, Hamburg, Germany): 94 °C for 3 minutes, 40 cycles of (94 °C for 30 seconds, 55

°C for 30 seconds, 72 °C for 30 seconds), 72 °C for 1 minute, hold at 20 °C. PCR products were

visualized under UV light on a 1.0 % (w/v) agarose gel containing ethidium bromide.

Homopolymer PCR, cloning and sequencing

To determine the length of the homopolymer region of *nanH3,* four strains (W11, VN014,

VN015, NR032) were cultured on Columbia Agar plates with 5% (v/v) sheep blood. Primers

JH0780 (5'-ATG ATT GGA ACA GCG CAT AAA G-3') and JH0781 (5'-GAT TTC TCC ACC

TAC AGT TAC C-3') were designed to PCR amplify a region including basepairs 2-310 of the

open reading frame of *nanH3*.

The DNA sequence (308 bp) was amplified by PCR in a Mastercycler Pro 6321 (Eppendorf

AG, Hamburg, Germany). The components of the PCR reaction mix (50 μl per reaction) were

added to achieve final concentrations of 1×High Fidelity PCR buffer (60 mM $Tris-SO_4$ (pH 8.9),

18 mM $(NH_4)_2SO_4$), 0.2 mM dNTP mix, 2 mM $MgSO_4$, 1U Platinum High Fidelity (Hi-Fi) proof

reading Taq polymerase (Invitrogen, Carlsbad, CA, USA). Two colonies of each *G. piotii* or

genome sp. 3 strain were randomly picked and added to separate PCR reactions using sterile

toothpicks. Thermocycling conditions included 35 cycles of denaturation at 94°C for 15 seconds,

annealing at 55°C for 30 seconds, extension at 68°C for 1 minute, and final extension at 68°C for

5 minutes. PCR products were visualized on a 1% (w/v) agarose gel. PCR products were purified

using QiaQuick PCR purification kit (Qiagen, Hilden, Germany) and purified PCR products were

sequenced using the amplification primers (JH0780, JH0781).

To clarify the exact length of the homopolymer region, amplicons generated from the two

colonies of each of the four strains were A-tailed in 10 μl reactions containing 1× Platinum PCR

buffer (20 mM Tris HCl (pH 8.4), 50 mM KCl), 2 mM $MgCl_2$, 0.5 mM dATP, 5 U Platinum Taq

polymerase (Invitrogen, Carlsbad, CA, USA) and less than 500 ng of the purified PCR product.

The reaction mixture was incubated at 72 °C for 20 minutes in Mastercycler Pro 6321 (Eppendorf

AG, Hamburg, Germany). End-modified PCR products were ligated into pGEM-T Easy vector

(Promega, Madison, WI, USA). The vector-insert construct was used to transform chemically

competent DH5α cells or OneShot Top 10 *E. coli* (Invitrogen, Carlsbad, CA, USA) and plated on

LB/AMP/X-gal agar media. Ten white colonies were randomly selected from the transformants of

each *Gardnerella* strain and transferred into LB + ampicillin broth. Cultures were grown overnight

at 37 °C. Plasmid DNA was isolated using QiaPrep Spin Miniprep kit (Qiagen, Hilden, Germany).

Plasmids were sequenced using vector primers T7 and SP6.

Results

Distribution of extracellular sialidases in *Gardnerella* spp. isolates.

Robinson et al. (16) had previously reported that sialidase activity was associated with the

presence of either *nanH2* or *nanH3* in a collection of 34 *Gardnerella* isolates but did not report

the species or subgroup affiliation. In order to confirm this observation and to reconcile the

distribution of genes and activities with the new *Gardnerella* taxonomic framework (6), we

queried an unrelated set of 36 isolates in our culture collection for which whole genome sequence

data and sialidase activity data were available (Table S1). The presence of the genes was assessed

by aligning *nanH2* of *G. piotii* JCP8151B (ATJH01000056) and *nanH3* of *Gardnerella* genome

sp. 3 strain W11 to the genome sequences using BLASTn. BLASTn results were dichotomous,

with either one clear match per genome (E = 0.0, bitscore 1275-5049, percent identity 95-99%) or

no significant hits reported. Sialidase activity was reported in 13/36 isolates (Table 1) including

12 *G. piotii* or genome sp. 3 (cpn60 subgroup B) isolates and one *G. vaginalis* isolate. The *nanH3*

gene was found in 12/13 sialidase activity positive isolates while *nanH2* was present in only 7/13

isolates. Six isolates possessed both *nanH2* and *nanH3* genes in their genomes. VN002 was

sialidase activity positive but only *nanH2* was identified in the genome sequence. This isolate

subsequently screen PCR positive for *nanH3* (see next section), suggesting that the gene was not

included in the shotgun assembly of the genome, which consisted of multiple gap-containing

scaffolds. None of the *G. swidsinskii, G. leopoldii* or subgroup D isolates were sialidase positive

and none contained *nanH2* or *nanH3*.

Interestingly, the NanH3 of *G. piotii* strain JCP8151B investigated by Robinson et al. was

reported to lack a signal peptide (16). When we examined the sequence upstream of *nanH3* in the

JCP8151B sequence (Genbank accession ATJH01000033), we found that there is an alternative

start codon, and a cytosine homopolymer. The frame-shift that resulted in the annotation of the

gene without the signal peptide-encoding N-terminus is caused by the homopolymer. This was

also the case for isolates GH020, GH022, N170, and NR026 where the homopolymer created a

frame-shift that resulted in a protein annotation lacking the signal peptide encoded by sequence

immediately upstream of the homopolymer. In N144, W11, N101, N153 and N95, the

homopolymer length did not create a frame-shift and the annotated NanH3 protein sequence

includes a signal peptide. In GH007, GH019, and GH015 we were unable to determine if a signal-

peptide encoding sequence was present due to assembly gaps immediately upstream of the

homopolymer in these genomes (Table S1).

Signal peptides were detected in 4/7 predicted NanH2 sequences (GH019, GH020, N101

and N153). In the other three cases where a *nanH2* sequence was identified, we could not conclude

if a signal peptide was included in the protein sequence provided in the Genbank annotation since

assembly gaps in those regions of the genomes affected the annotation of these proteins.

175   Correlation of *nanH3* with sialidase activity

In order to examine further the relationship of *nanH3* to sialidase activity, 112 *Gardnerella*

spp. isolates for which sialidase activity data was available (13) were screened for the presence of

nanH3 using PCR primers JH0684/JH0685 (Figure 1). Sialidase activity was detected in 32/33 *G.*

piotii/genome sp. 3 isolates and 3/35 *G. vaginalis* isolates. All sialidase activity positive isolates

were positive for *nanH3* by PCR with the exception of one *G. piotii* isolate (WP027) that was PCR

negative. Unfortunately, genome sequence data was not available for WP027.

182   Characterization of a poly-C homopolymer in *nanH3*

The protein sequences encoded by all nine *nanH3* sequences without assembly gaps were

predicted to encode a signal peptide (amino acids 1-32), a sialidase domain (amino acids 183-547

in W11) and a C-terminal transmembrane domain (amino acids 789-807 in W11) (Figure 2A). We

observed a poly-cytosine homopolymer (8-14 cytosines) in all *nanH3* genes identified in the whole

genome sequences, approximately 100 bases from the start of the open reading frame. The

homopolymer occurred immediately following the region of the sequence predicted to encode the

signal peptide (amino acids 1-32) (Figure 2B). When Sanger sequencing was performed on PCR

products corresponding to nucleotides 2-310 of *nanH3* amplified from genomic DNA extracted

from broth cultures of strain W11, the results suggested variable lengths of the homopolymer.

Specifically, clean data was obtained upstream of the poly-C tract but sequence 3' to that region

was indicative of a mixed template (Figure S1). Since it has been demonstrated that simple repeats

such as homopolymers can be subject to slipped strand mispairing, we set out to determine if the

homopolymer region of *nanH3* varied in length within and between strains of *G. piotii*/genome

species 3.

PCR product libraries were made from two colonies each of isolates W11, VN014, VN015

and NR032 (8 clone libraries total). Plasmids were purified from 10 colonies from each of the 8

clone libraries and sequenced. High quality sequence data was obtained from 15, 14, 18 and 16

clones from the W11, VN014, VN015 and NR032 libraries, respectively. Example results are

shown in Figure S2. The length of the homopolymer varied from 8 to 14 among all strains (Table

2), and within each strain the sequence flanking the homopolymer was identical. *In silico*

translation of the encoded polypeptides showed that most homopolymer lengths resulted in a

truncated peptide (37-44 amino acids), while full-length protein (812 aa) would result when there

were nine or twelve cytosines in the homopolymer region. Truncated peptides would include the

signal peptide (amino acids 1-32) but no part of the predicted sialidase domain (amino acids 183-

547) (Figure 2). Interestingly, the most common length of the poly-C region varied among strains.

C_{10} was most common in VN014 and VN015, while C_9 and C_{12} were most frequently observed in

NR032 and W11, respectively. VN015 had the highest number of homopolymer variants (Figure

3). A second open reading frame within the *nanH3* open reading frame was also identified,

corresponding to amino acids Met_{397}-Tyr_{812} of the NanH3 protein. These 415 aa polypeptides

encompass part of the predicted sialidase domain, and the C-terminal transmembrane domain but

lacks a signal peptide.

Discussion

The genus *Gardnerella* is known as a hallmark of BV and their abundance is used as a

criterion for the laboratory diagnosis of the condition (29). In addition to participating in biofilms

that coat epithelial cells in BV, *Gardnerella* spp. produce enzymes and a cholesterol dependent

cytolysin (vaginolysin) that contribute to degrading the protective barriers of the vaginal mucosa

(30). Sialidase activity can provide nutrients by releasing sialic acid moieties from vaginal

sialylated mucins (11), altering the physical properties of vaginal mucus. Removal of sialic acid

residues from epithelial cell surface glycans can facilitate bacterial adhesion and initiation of

biofilm formation (12, 18, 20, 21).

The two putative sialidase genes (*nanH2* and *nanH3*) in *Gardnerella* spp. have been

demonstrated to encode proteins with high enzymatic potency (16). Taken together, the result of

this study and our current study show that *nanH3* is more common and that *nanH2* is virtually

never found without *nanH3*. It is now clear that extracellular sialidase activity is a property of *G.*

piotii and the closely related *Gardnerella* genome sp. 3. We found three occurrences of sialidase

positive, *nanH3* positive *G. vaginalis* isolates (3/35 isolates screened), and Vaneechoutte et al. (6)

reported sialidase activity in 1/4 *G. vaginalis* isolates used in the amendment of the genus. This

infrequent prevalence of *nanH3* in *G. vaginalis* could be the result of lateral gene transfer from *G.*

piotii since these species co-exist in the same microbiome and women are usually colonized by

more than one species (7, 31, 32).

The *nanH3* sequences we examined were predicted to encode a signal peptide, a sialidase

domain and a C-terminal membrane domain, suggesting a cell-wall tethered protein, similar to

SiaBb2 of *Bifidobacterium bifidum*, which enhances adhesion to intestinal mucosal surfaces and

contributes to carbohydrate assimilation (21). The presence of the homopolymer immediately

downstream of the signal peptide encoding region presents a challenge to automated annotation of

this sequence. Sequencing and assembly methods commonly used in bacterial whole genome

sequencing struggle with homopolymeric sequences, which can result in frame-shifts such as we

observed in the *nanH3* sequence of JCP8151B.

Genomic regions that contain homogenous or heterogenous repeats are prone to changes

in length of the repeat at each replication due to slipped strand mispairing (22). This can lead to

consequent changes in the transcription or translation product of a gene depending on whether the

slipped strand mispairing occurs within the open reading frame or in extragenic regions such as

promoter sequences. Phase variation results when slipped strand mispairing creates "on" and "off"

states of expression of a protein. Phase variation of cell surface proteins has been documented in

many bacterial species including *Neisseria* spp. (33–35), *Salmonella* spp. (36–38), *Treponema*

pallidum (39, 40) and *Helicobacter pylori* (25). It was first described in the *opa* genes that encode

opacity surface proteins of *Neisseria* spp. (41). The *opa* genes contain CTCTT pentamer repeats

within the signal peptide encoding region and the expression of the Opa protein is regulated by

slipped-strand mispairing. With six, nine or twelve CTCTT repeats the initiation codon is in frame

with the remaining *opa* gene translating the Opa protein. Four or eight coding repeats makes the

initiation codon out of frame with rest of the *opa* codons. The phase "on" state allows them to

adhere to specific surface receptors on host cells (42).

Our results clearly show variation of the homopolymer length among *G. piotii* and

Gardnerella genome sp. 3 within colonies growing on agar, with anywhere from 8 to 14 C's

observed. The effect of this change would be a mixture of cells with translation of NanH3 on or

off. Why would *G. piotii* have a sialidase subject to phase variation? Two possible reasons are

immune evasion and cell adhesion (biofilm initiation and dispersal).

Although BV is often referred to as a non-inflammatory condition because of the lack of

typical clinical signs of inflammation (leukocyte infiltration, pain, redness, swelling), there is

evidence of a host immune response to bacteria associated with BV, including *Gardnerella* spp..

IgA specific for *Gardnerella* produced vaginolysin has been detected in women with clinical signs

of BV (43), and IgA levels correlate with IL-8 expression in vaginal secretions (44). *Gardnerella*

has also been shown to stimulate pro-inflammatory cytokines *in vitro* (45). Balancing this

inflammatory response are the actions of bacterial enzymes like prolidases and sialidases that can

degrade the effectors of inflammation. Whether there is a specific host response to NanH3 remains

to be determined.

The initial step in biofilm formation is adhesion, which in the case of BV associated

biofilms is to epithelial cells (2). *Streptococcus pneumoniae* surface proteins with sialidase activity

have been shown to reveal carbohydrate ligands for bacterial adhesion to host cells (18). Similarly,

Soong et al. (19) showed that a cell wall associated sialidase of *Pseudomonas aeruginosa* plays a

critical role in facilitating respiratory mucosa infection through biofilm formation. Specific

antibodies against *B. bifidum* SiaBb2, a cell wall tethered sialidase, inhibit adhesion to cells (21).

The involvement of *Gardnerella* sialidases in adhesion has not been demonstrated directly,

although it has been reported that inhibition of sialidase activity in *Gardnerella* strain JCP8066 by

an antiviral drug (Zanamavir) resulted in reduction of adherence to vaginal epithelial cells *in vitro*

(12). *Gardnerella* spp. can form multi-species biofilms *in vitro* (46) and although "*Gardnerella*

vaginalis" has been shown to participate in multispecies biofilms with other BV associated

bacteria *in vivo* (47, 48) and *in vitro* (49)*, it is not yet known how each of the *Gardnerella* spp.

participate in this process. Interestingly, *G. piotii* and *Gardnerella* genome sp. 3 (cpn60 subgroup

B) have been previously associated with "intermediate" grades of vaginal dysbiosis (7, 31, 50, 51)

and may contribute to transition from eubyosis to dysbiosis and enhance colonization by other BV

associated anaerobes. Phase variation in a cell surface associated sialidase enzyme might be critical

for turning on or off the initial adhesion of *Gardnerella* to epithelial cells and thus the cascade of

events following that result in the biofilm covered "clue cells" typical of BV (52).

It seems likely that some of the NanH3 protein produced by *G. piotii* does not remain

tethered to the cell surface, as is the case with *S. pneumoniae* (53), since the sequence upstream of

the C-terminal transmembrane domain may be susceptible to proteolytic cleavage, and

heterologously expressed NanH3 lacking the transmembrane domain is active (16). Taken

together, the activity of NanH3 while anchored to the cell surface or released into the extracellular

environment, and the likelihood that it is subject to phase variation through slipped-strand

mispairing, make this protein an intriguing puzzle for future study. Development of genetic tools

to create mutants where NanH3 production is locked "on" or "off" would be a significant step

toward understanding the role of this protein in the initiation and maintenance of vaginal dysbiosis.

Acknowledgements

This research was funded by a Natural Sciences and Engineering Council of Canada Discovery

Grant to JEH. The authors are grateful to Champika Fernando for excellent technical assistance,

and to all our colleagues in the Hill Lab for helpful discussions and feedback.

References

1.   Nugent RP, Krohn MA, Hillier SL. 1991. Reliability of diagnosing bacterial vaginosis is improved

by a standardized method of gram stain interpretation. J Clin Microbiol 29:297–301.

2.   Rosca AS, Castro J, Sousa LGV, Cerca N. 2020. *Gardnerella* and vaginal health: the truth is out

there. FEMS Microbiol Rev 44:73–105.

3.   van de Wijgert JH, Borgdorff H, Verhelst R, Crucitti T, Francis S, Verstraelen H, Jespers V. 2014.

The vaginal microbiota: what have we learned after a decade of molecular characterization? PLoS

One 9:e105998.

4.   Paramel Jayaprakash T, Schellenberg JJ, Hill JE. 2012. Resolution and characterization of distinct

cpn60-based subgroups of *Gardnerella vaginalis* in the vaginal microbiota. PLoS ONE 7:e43009.

5.   Ahmed A, Earl J, Retchless A, Hillier SL, Rabe LK, Cherpes TL, Powell E, Janto B, Eutsey R,

Hiller NL, Boissy R, Dahlgren ME, Hall BG, Costerton JW, Post JC, Hu FZ, Ehrlich GD. 2012.

Comparative genomic analyses of 17 clinical isolates of *Gardnerella vaginalis* provide evidence of

multiple genetically isolated clades consistent with subspeciation into genovars. J Bacteriol

194:3922–37.

6.   Vaneechoutte M, Guschin A, Van Simaey L, Gansemans Y, Van Nieuwerburgh F, Cools P. 2019.

Emended description of *Gardnerella vaginalis* and description of *Gardnerella leopoldii* sp. nov.,

Gardnerella piotii sp. nov. and *Gardnerella swidsinskii* sp. nov., with delineation of 13 genomic

species within the genus *Gardnerella*. Int J Syst Evol Microbiol 69:679–687.

7.   Hill JE, Albert AYK. 2019. Resolution and co-occurrence patterns of *Gardnerella leopoldii,*

Gardnerella swidsinskii, Gardnerella piotii and *Gardnerella vaginalis* within the vaginal

microbiome. Infect Immun 87:e00532-19.

8.   McGregor JA, French JI, Jones W, Milligan K, McKinney PJ, Patterson E, Parker R. 1994.

Bacterial vaginosis is associated with prematurity and vaginal fluid mucinase and sialidase: results

of a controlled trial of topical clindamycin cream. Am J Obstet Gynecol 170:1048–1059.

9.   Myziuk L, Romanowski B, Johnson SC. 2003. BVBlue test for diagnosis of bacterial vaginosis. J

Clin Microbiol 41:1925–8.

10.   Briselden AM, Moncla BJ, Stevens CE, Hillier SL. 1992. Sialidases (neuraminidases) in bacterial

vaginosis and bacterial vaginosis-associated microflora. J Clin Microbiol 30:663–6.

11.   Lewis WG, Robinson LS, Gilbert NM, Perry JC, Lewis AL. 2013. Degradation, foraging, and

depletion of mucus sialoglycans by the vagina-adapted Actinobacterium *Gardnerella vaginalis*. J

Biol Chem 288:12067–79.

12.   Govinden G, Parker JL, Naylor KL, Frey AM, Anumba DOC, Stafford GP. 2018. Inhibition of

sialidase activity and cellular invasion by the bacterial vaginosis pathogen *Gardnerella vaginalis*.

Arch Microbiol 200:1129–1133.

13.   Schellenberg JJ, Paramel Jayaprakash T, Withana Gamage N, Patterson MH, Vaneechoutte M, Hill

JE. 2016. *Gardnerella vaginalis* subgroups defined by cpn60 sequencing and sialidase activity in

isolates from Canada, Belgium and Kenya. PLoS One 11:e0146510.

14.   Santiago GL, Deschaght P, El Aila N, Kiama TN, Verstraelen H, Jefferson KK, Temmerman M,

Vaneechoutte M. 2011. *Gardnerella vaginalis* comprises three distinct genotypes of which only two

produce sialidase. Am J Obstet Gynecol 204:450.e1–7.

15.   Corfield T. 1992. Bacterial sialidases—roles in pathogenicity and nutrition. Glycobiology 2:509–

521.

16.   Robinson LS, Schwebke J, Lewis WG, Lewis AL. 2019. Identification and characterization of

NanH2 and NanH3, enzymes responsible for sialidase activity in the vaginal bacterium *Gardnerella*

vaginalis. J Biol Chem 294:5230–5245.

17.   Potter RF, Burnham C-AD, Dantas G. 2019. In silico analysis of *Gardnerella* genomospecies

detected in the setting of bacterial vaginosis. Clin Chem 65:1375–1387.

18.   Navarre WW, Schneewind O. 1999. Surface proteins of Gram-positive bacteria and mechanisms of

their targeting to the cell wall envelope. Microbiol Mol Biol Rev MMBR 63:174–229.

19.   Soong G, Muir A, Gomez MI, Waks J, Reddy B, Planet P, Singh PK, Kanetko Y, Wolfgang MC,

Hsiao Y-S, Tong L, Prince A. 2006. Bacterial neuraminidase facilitates mucosal infection by

participating in biofilm production. J Clin Invest 116:2297–2305.

20.   Parker D, Soong G, Planet P, Brower J, Ratner AJ, Prince A. 2009. The NanA neuraminidase of

Streptococcus pneumoniae is involved in biofilm formation. Infect Immun 77:3722–30.

21.   Nishiyama K, Yamamoto Y, Sugiyama M, Takaki T, Urashima T, Fukiya S, Yokota A, Okada N,

Mukai T. 2017. *Bifidobacterium bifidum* extracellular sialidase enhances adhesion to the mucosal

surface and supports carbohydrate assimilation. MBio 8.

22.   Woude MW van der, Bäumler AJ. 2004. Phase and antigenic variation in bacteria. Clin Microbiol

Rev 17:581–611.

23.   Phillips ZN, Tram G, Seib KL, Atack JM. 2019. Phase-variable bacterial loci: how bacteria gamble

to maximise fitness in changing environments. Biochem Soc Trans 47:1131–1141.

24.   Seib KL, Jen FE-C, Scott AL, Tan A, Jennings MP. 2017. Phase variation of DNA

methyltransferases and the regulation of virulence and immune evasion in the pathogenic *Neisseria*.

Pathog Dis 75.

25.   Bergman M, Del Prete G, van Kooyk Y, Appelmelk B. 2006. Helicobacter pylori phase variation,

immune modulation and gastric autoimmunity. 2. Nat Rev Microbiol 4:151–159.

26.   Nielsen H, Tsirigos KD, Brunak S, von Heijne G. 2019. A brief history of protein sorting prediction.

Protein J 38:200–216.

27.   Khan S, Vancuren SJ, Hill JE. 2020. A generalist lifestyle allows rare *Gardnerella* spp. to persist at

low levels in the vaginal microbiome. bioRxiv 2020.08.25.267146.

28.   Martin-Platero AM, Valdivia E, Maqueda M, Martinez-Bueno M. 2007. Fast, convenient, and

economical method for isolating genomic DNA from lactic acid bacteria using a modification of the

protein "salting-out" procedure. Anal Biochem 366:102–4.

29.   Nugent RP, Krohn MA, Hillier SL. 1991. Reliability of diagnosing bacterial vaginosis is improved

by a standardized method of gram stain interpretation. J Clin Microbiol 29:297–301.

30.   Schellenberg JJ, Patterson MH, Hill JE. 2017. *Gardnerella vaginalis* diversity and ecology in

relation to vaginal symptoms. Res Microbiol doi:10.1016/j.resmic.2017.02.011.

31.   Balashov SV, Mordechai E, Adelson ME, Gygax SE. 2014. Identification, quantification and

subtyping of *Gardnerella vaginalis* in noncultured clinical vaginal samples by quantitative PCR. J

Med Microbiol 63:162–75.

32.   Albert AY, Chaban B, Wagner EC, Schellenberg JJ, Links MG, van Schalkwyk J, Reid G,

Hemmingsen SM, Hill JE, Money D. 2015. A study of the vaginal microbiome in healthy Canadian

women utilizing cpn60-based molecular profiling reveals distinct *Gardnerella* subgroup community

state types. PLoS One 10:e0135620.

33.   Bayliss CD, Hoe JC, Makepeace K, Martin P, Hood DW, Moxon ER. 2008. *Neisseria meningitidis*

escape from the bactericidal activity of a monoclonal antibody is mediated by phase variation of

lgtG and enhanced by a mutator phenotype. Infect Immun 76:5038–5048.

34.   Apicella MA, Shero M, Jarvis GA, Griffiss JM, Mandrell RE, Schneider H. 1987. Phenotypic

variation in epitope expression of the *Neisseria gonorrhoeae* lipooligosaccharide. Infect Immun

55:1755–1761.

35.   van Putten JP. 1993. Phase variation of lipopolysaccharide directs interconversion of invasive and

immuno-resistant phenotypes of *Neisseria gonorrhoeae*. EMBO J 12:4043–4051.

36.   Lederberg J, Iino T. 1956. Phase variation in *Salmonella*. Genetics 41:743–757.

37.   Silverman M, Zieg J, Hilmen M, Simon M. 1979. Phase variation in *Salmonella*: genetic analysis of

a recombinational switch. Proc Natl Acad Sci 76:391–395.

38.   Bonifield HR, Hughes KT. 2003. Flagellar phase variation in *Salmonella enterica* is mediated by a

posttranscriptional control mechanism. J Bacteriol 185:3567–3574.

39.   Fraser CM, Norris SJ, Weinstock GM, White O, Sutton GG, Dodson R, Gwinn M, Hickey EK,

Clayton R, Ketchum KA, Sodergren E, Hardham JM, McLeod MP, Salzberg S, Peterson J, Khalak

H, Richardson D, Howell JK, Chidambaram M, Utterback T, McDonald L, Artiach P, Bowman C,

Cotton MD, Fujii C, Garland S, Hatch B, Horst K, Roberts K, Sandusky M, Weidman J, Smith HO,

Venter JC. 1998. Complete genome sequence of *Treponema pallidum*, the syphilis spirochete.

Science 281:375–388.

40.   Pinto M, Borges V, Antelo M, Pinheiro M, Nunes A, Azevedo J, Borrego MJ, Mendonça J,

Carpinteiro D, Vieira L, Gomes JP. 2016. Genome-scale analysis of the non-cultivable *Treponema*

pallidum reveals extensive within-patient genetic variation. 1. Nat Microbiol 2:1–11.

41.   Stern A, Brown M, Nickel P, Meyer TF. 1986. Opacity genes in *Neisseria gonorrhoeae*: Control of

phase and antigenic variation. Cell 47:61–71.

42.   Stern A, Nickel P, Meyer TF, So M. 1984. Opacity determinants of *Neisseria gonorrhoeae*: Gene

expression and chromosomal linkage to the gonococcal pilus gene. Cell 37:447–456.

43.   Cauci S, Scrimin F, Driussi S, Ceccone S, Monte R, Fant L, Quadrifoglio F. 1996. Specific immune

response against *Gardnerella vaginalis* hemolysin in patients with bacterial vaginosis. Am J Obstet

Gynecol 175:1601–5.

44.   Cauci S, Driussi S, Guaschino S, Isola M, Quadrifoglio F. 2002. Correlation of local Interleukin-

1beta levels with specific IgA response against *Gardnerella vaginalis* cytolysin in women with

bacterial vaginosis. Am J Reprod Immunol 47:257–264.

45.   Eade CR, Diaz C, Wood MP, Anastos K, Patterson BK, Gupta P, Cole AL, Cole AM. 2012.

Identification and characterization of bacterial vaginosis-associated pathogens using a

comprehensive cervical-vaginal epithelial coculture assay. PLoS One 7:e50106.

46.   Khan S, Voordouw MJ, Hill JE. 2019. Competition among *Gardnerella* subgroups from the human

vaginal microbiome. Front Cell Infect Microbiol 9:374.

47.   Machado A, Cerca N. 2015. Influence of biofilm formation by *Gardnerella vaginalis* and other

anaerobes on bacterial vaginosis. J Infect Dis 212:1856–61.

48.   Machado D, Castro J, Palmeira-de-Oliveira A, Martinez-de-Oliveira J, Cerca N. 2015. Bacterial

vaginosis biofilms: Challenges to current therapies and emerging solutions. Front Microbiol 6:1528.

49.   Castro J, Machado D, Cerca N. 2019. Unveiling the role of Gardnerella vaginalis in polymicrobial

Bacterial Vaginosis biofilms: the impact of other vaginal pathogens living as neighbors. ISME J

13:1306–1317.

50.   Hilbert DW, Schuyler JA, Adelson ME, Mordechai E, Sobel JD, Gygax SE. 2017. *Gardnerella*

433   *vaginalis* population dynamics in bacterial vaginosis. Eur J Clin Microbiol Infect Dis

434   https://doi.org/10.1007/s10096-017-2933-8.

435   51.   Shipitsyna E, Krysanova A, Khayrullina G, Shalepo K, Savicheva A, Guschin A, Unemo M. 2019.

436   Quantitation of all four *Gardnerella vaginalis* clades detects abnormal vaginal microbiota

437   characteristic of bacterial vaginosis more accurately than putative *G. vaginalis* sialidase A gene

438   count. Mol Diagn Ther 23:139–147.

439   52.   Simoes JA, Discacciati MG, Brolazo EM, Portugal PM, Dini DV, Dantas MC. 2006. Clinical

440   diagnosis of bacterial vaginosis. Int J Gynaecol Obstet 94:28–32.

441   53.   Lock RA, Paton JC, Hansman D. 1988. Purification and immunological characterization of

442   neuraminidase produced by *Streptococcus pneumoniae*. Microb Pathog 4:33–43.

445   **Table 1.** Distribution of *nanH2* and *nanH3* genes in *Gardnerella* spp. whole genome sequences

Isolate	cpn60 Subgroup	Species	Sialidase activity[1]	*nanH2*	*nanH3*
GH007	B	*G. piotii*	+	+	+
GH019		*G. piotii*	+	+	+
GH020		*G. piotii*	+	+	+
GH022		*G. piotii*	+	+	+
N170		Genome species 3	+	-	+
NR026		Genome species 3	+	-	+
VN002		*G. piotii*	+	+	-[2]
N144		Genome species 3	+	-	+
W11		Genome species 3	+	-	+
N101		Genome species 3	+	+	+
N153		Genome species 3	+	+	+
N95		Genome species 3	+	-	+
GH005	A	*G. leopoldii*	-	-	-
NR015		Unknown	-	-	-
NR016		*G. swidsinskii*	-	-	-
NR017		*G. leopoldii*	-	-	-
NR019		*G. leopoldii*	-	-	-
NR020		*G. swidsinskii*	-	-	-
NR021		*G. swidsinskii*	-	-	-
VN003		*G. leopoldii*	-	-	-
WP021		*G. swidsinskii*	-	-	-
WP022		*G. swidsinskii*	-	-	-
NR010		*G. leopoldii*	-	-	-
N072		*G. swidsinskii*	-	-	-
GH021	C	*G. vaginalis*	-	-	-
NR001		*G. vaginalis*	-	-	-
NR037		*G. vaginalis*	-	-	-
NR038		*G. vaginalis*	-	-	-
NR039		*G. vaginalis*	-	-	-
WP023		*G. vaginalis*	-	-	-
N165		*G. vaginalis*	-	-	-
GH015		*G. vaginalis*	+	-	+
NR003	D	Genome sp. 8	-	-	-
NR047		Unknown	-	-	-
WP012		Genome sp. 9	-	-	-
N160		Genome sp. 10	-	-	-

446   [1]Previously determined (13)
447   [2]PCR positive for *nanH3*

Table 2. Homopolymer length variants in *Gardnerella piotii* and genome sp. 3 isolates.

Isolate	Homopolymer sequence	Predicted peptide length (amino acids)[1]
W11	CAACTA- C_{11}-ATGAACAAA	43
	CAACTA- C_{12}-ATGAACAAA	812
	CAACTA- C_{13}-ATGAACAAA	38
	CAACTA- C_{14}-ATGAACAAA	44
VN014	CAACTA- C_9-TCGAACAAA	812
	CAACTA- C_{10}-TCGAACAAA	37
	CAACTA- C_{11}-TCGAACAAA	43
VN015	CAACTA- C_9-TCGAACAAA	812
	CAACTA- C_{10}-TCGAACAAA	37
	CAACTA- C_{11}-TCGAACAAA	43
	CAACTA- C_{12}-TCGAACAAA	812
	CAACTA- C_{13}- - -GAACAAA	43
NR032	CAACTA- C_8-ATGAACAAA	42
	CAACTA- C_9-ATGAACAAA	812

[1]Full-length NanH3 is 812 amino acids

Figure 1. Correlation of *nanH3* and sialidase activity in 112 *Gardnerella* isolates. Numbers of isolates positive or negative for sialidases activity and *nanH3* PCR. cpn60 subgroup affiliation and species are indicated below each crosstab. Sialidase results were determined by Schellenberg *et al*. (13).

467 **Figure 2.** A. Predicted locations of signal peptide, sialidase and transmembrane domains of

468 NanH3 from strain W11 based on InterProScan analysis. B. Location of predicted signal peptide,

469 cleavage site and adjacent homopolymer in the *nanH3* gene of strain W11. Analysis was

470 conducted with SignalP-5.0.

Figure 3. Frequencies of the homopolymer length variants in isolates W11, VN014, VN015 and

NR032.

Supplemental Materials

Table S1. *Gardnerella* strains used in the study, cpn60 subgroup affiliations, whole genome

sequence accession numbers, sialidase activity phenotypes, presence/absence of *nanH2, nanH3,*

signal peptide encoding sequences and homopolymers.

Figure S1. Sequencing of PCR products amplified directly from colonies of isolate W11.

PCR was performed with primers flanking the homopolymer region as described in the methods.

In each electropherogram, after eleven peaks of cytosine, the sequencing signal shows multiple

overlapping peaks suggesting the presence of multiple template sequences with different

homopolymer lengths.

Figure S2. Electropherograms of the homopolymer regions in two W11 colonies. Colony 6 had

individuals with 11, 12 and 13 C's in the homopolymeric region. Five out of eight colonies had

12C's in the homopolymer region. Colony 8 had individuals with 11, 12 and 14 C's in the

homopolymeric region. Five out of eight colonies had 12 C's in the homopolymer region. Having

12 C's makes the coding region in-frame. When there is 11,13 or 14 C's, a premature STOP codon

is generated possibly making the protein truncated.