Fixation and effective size in a haploid-diploid population with asexual reproduction

*Kazuhiro Bessho, **Sarah P. Otto

* Saitama Medical University, 38 Morohongo Moroyama-machi, Iruma-gun, Saitama 350-0495,

Japan

**Department of Zoology, The University of British Columbia, Vancouver, BC, V6T 1Z4,

Canada

* besshokazuhiro.reserch@gmail.com; (+81) 049-276-2030

** otto@zoology.ubc.ca; (+1) 604-822-2778

*Keywords***:** haploid-diploid life cycle, Wright-Fisher model, fixation probability, effective

population size, heteromorphic and isomorphic life cycle

Abstract

The majority of population genetic theory assumes fully haploid or diploid organisms

with obligate sexuality, despite complex life cycles with alternating generations being commonly

observed. To reveal how natural selection and genetic drift shape the evolution of haploid-diploid

populations, we analyze a stochastic genetic model for populations that consist of a mixture of

haploid and diploid individuals, allowing for asexual reproduction and niche separation between

haploid and diploid stages. Applying a diffusion approximation, we derive the fixation

probability and describe its dependence on the reproductive values of haploid and diploid stages,

which depend strongly on the extent of asexual reproduction in each phase and on the ecological

differences between them.

1. Introduction

Sexual reproduction in eukaryotes generally consists of an alternation of generations, where meiosis halves the number of chromosomes to produce haploids and syngamy brings together haploid gametes to produce diploids. The extent of development in each ploidy phase varies substantially (Bell 1982; 1994). In diplontic organisms, at one extreme, development and growth occur only in the diploid phase, as is observed in most animals. Haplontic organisms, at the other extreme, undergo mitotic growth only in the haploid stage, as is seen in some green algae. In between these extremes, many terrestrial plants, macroalgae, and fungi exhibit both haploid and diploid growth (haploid-diploid life cycles). These stages are typically free living in macroalgae, with either macroscopically similar (isomorphic) or distinct (heteromorphic) forms in the haploid and diploid stage (Raper and Flexer 1970; Wilson 1981; Mable and Otto 1998; Coelho 2007).

To explain variation in life cycles, several theoretical models have analyzed the deterministic dynamics of a modifier allele that alters the time spent in haploid and diploid phases (e.g., Perrot et al. 1991; Otto and Goldstein 1992; Goldstein 1992; Otto 1994; Orr and Otto 1994; Jenkins and Kirkpatrick 1995; Otto and Marks 1996; Scott and Rescan 2017). However, there are some gaps between these models and the complexities seen in many haploid-diploid species. For example, these models often treat haploid and diploid individuals as ecologically equivalent, despite the frequent observation of niche differences and seasonal shifts in prevalence (e.g., Drew 1949; Slocum 1980; Dethier 1981). Most of these models also assume obligate sexuality (but see Otto and Marks 1996), despite asexuality being frequently observed among haploid-diploid species ("asexual looping"). Furthermore, while several models have explored how haploid-diploid life cycles might evolve, the impact of haploid-diploid life cycles

on evolutionary processes remains underexplored (see, e.g., Bessho and Otto 2017 on the impact

on fixation probabilities and Immler et al. 2012 on the maintenance of variation).

Here we contribute to evolutionary theory for haploid-diploid populations by

calculating the fixation probability of mutations using a stochastic genetic model. This builds

upon our previous work (Bessho and Otto 2017) by accounting for asexual looping and niche

differences between ploidy phases, both of which are common in macroalgae (Bell 1982; de

Wreede and Klinger 1988; Hawkes 1990). Haploid and diploid phases often differ

physiologically, and even isomorphic haploids and diploids may differ ecologically (Hannach

and Santelices 1985; Destombe et al. 1993; Dyck and de Wreede 2006; Thornber et al., 2006;

Vieira et al. 2018). We therefore explore different forms of density dependence, acting either

globally on the total population size (as in Bessho and Otto 2017) or locally on the population

size of haploids and diploids separately (Figure 1). We show that the fate of a mutation depends

strongly on the reproductive values of haploids and diploids, which in turn depend on the extent

of asexual reproduction and ecological differences between the phases.

2. Model

In Bessho and Otto (2017), we calculated the fixation probabilities by tracking the

dynamics of a resident allele (R) and a mutant allele (M) in haploid and diploid individuals, using

both a Wright-Fisher and a Moran model. In that model, reproduction was obligately sexual,

individuals were ecologically equivalent, and the total population size was held constant (global

density dependence). Below, we calculate the fixation probability by first considering asexual

reproduction in each phase, assuming that haploids and diploids are ecologically equivalent

(global population regulation), and then determine how these results are affected by niche

differences (local population regulation that is ploidy specific).

2.1. Haploid-diploid Wright-Fisher model with global regulation and asexual looping

Let $X_{(GT)}(t)$ be a random variable that represents the number of individuals with "genotype" (GT) at time t with resident (R) and mutant alleles (M), and let $x_{(GT)}(t)$ represent a particular outcome of this random variable. In the global regulation model, we assume a constant population, $x_R + x_M + x_{RR} + x_{RM} + x_{MM} = N_{tot}$, that is strictly regulated regardless of the ploidy of the individuals.

The reproductive output and the degree of asexuality are characterized by $w_{(GT)}$ and a_H for haploids [(GT) = R or M] and $w_{(GT)}$ and a_D for diploids [(GT) = RR, RM, and MM]. Specifically, diploid individuals produce $(1 - a_D)w_{(GT)}$ haploid spores (sexual reproduction) and $a_D w_{(GT)}$ diploid offspring (asexual loop). Similarly, haploids produce $(1 - a_H)w_{(GT)}/2$ female gametes (sexual reproduction) and $a_H w_{(GT)}$ haploid offspring (asexual loop), where we assume that the species is monoecious and invests equal resources in male and female gametes. During syngamy, we assume that male gametes are not limiting, that mating is random, and that female gametes are successfully fertilized with male gametes, at a rate $f_{(GT)}$ [(GT) = R or M], becoming diploid zygotes. For clarity, we describe the model with non-overlapping generations, although we note that overlapping generations can be considered by including surviving adults in the counts of asexual offspring ($a_D w_{(GT)}$ and $a_H w_{(GT)}$).

We define the selection coefficient ($s_{(GT)}$) and the degree of dominance (h) acting upon the mutant allele such that: $f_M = f_R\left(1 - s_M^f\right)$, $w_M = w_R(1 - s_M^w)$, $w_{RM} = w_{RR}(1 - s_{RM}^w)$, $w_{MM} = w_{RR}(1 - s_{MM}^w)$, and $h = s_{RM}^w/s_{MM}^w$. To perform the diffusion approximation, we assume that selection is weak, $s_M^f = \epsilon \tilde{s}_M^f$ and $s_{(GT)}^w = \epsilon \tilde{s}_{(GT)}^w$, where ϵ is a small parameter.

2.2. Haploid-diploid Wright-Fisher model with local regulation and asexual looping

We then consider the case where density dependence regulates haploid and diploid

populations separately, which may occur if they have different resource needs or utilize different

habitats or microhabitats (for short-hand, we refer to this case as "local regulation"). More

specifically, we assume that the population size of haploids and diploids is separately regulated

and remains constant N_H and N_D ($x_R + x_M = N_H$ and $x_{RR} + x_{RM} + x_{MM} = N_D$), respectively.

We set $N_H + N_D = N_{tot}$, $\hat{\rho}_H^L = N_H/N_{tot}$, and $\hat{\rho}_D^L = N_D/N_{tot}$, which will then allow us to

compare the results of local and global regulation. Holding population sizes constant is assumed

strictly for mathematical convenience but may be reasonable for populations whose sizes are

strongly regulated by the availability of appropriate habitat.

3. Fixation probability in a haploid-diploid population

3.1. Fixation probability in the global regulation model

The fixation probability in a haploid-diploid population can be derived using a diffusion

approximation (Bessho and Otto 2017), but doing so requires that we approximate the dynamics

to reduce the dimensionality from four variables ($x_R, x_M, x_{RR}, x_{RM}, x_{MM}$, which sum to N_{tot}) down

to one. We do so by using a separation of time scales, deriving the first and second moments of the

mutant allele frequency. Specifically, we transform the number of individuals of each genotype,

$x_{(GT)}$, into new variables that allow us to separate the slower evolutionary dynamics and the faster

ecological dynamics (Appendix A):

$$p_{ave} = c_H\, p_H + c_D\, p_D, \tag{1a}$$

$$\delta_p = p_D - p_H, \tag{1b}$$

$$\eta_{HW} = 1 - \frac{1}{2p_D(1-p_D)}\frac{x_{RM}}{x_{RR} + x_{RM} + x_{MM}}, \tag{1c}$$

$$\rho_H = \frac{x_R + x_M}{N_{tot}}, \tag{1d}$$

where p_{ave} indicates the average allele frequency of haploids and diploids weighted by the class

reproductive values (c_H and c_D, where $c_H + c_D = 1$, see next paragraph), δ_p indicates the

difference in allele frequencies between haploids and diploids, η_{HW} indicates the departure from

the Hardy-Weinberg equilibrium in diploids, and ρ_H indicates the frequency of haploids in the

population. Within these equations, the frequencies of mutant alleles in haploids and diploids are

$p_H = x_M/(x_R + x_M)$ and $p_D = \left(\frac{x_{RM}}{2} + x_{MM}\right)/(x_{RR} + x_{RM} + x_{MM})$. As similar variables are

used in the model with local population regulation, we use superscripts to indicate the form of

population regulation ("*Model*" is *G* for global or *L* for local regulation).

The class reproductive values of haploids and diploids are defined as follows. In linear

models, "reproductive value" is a measure of the expected fraction of the population in the long-

term future that descends from an individual of a particular type (e.g., age or stage class). Class

reproductive values, as defined by Taylor (1990) and Rousset (2004, p.153), scale these individual

reproductive values up to the whole population of each class (i.e., the product of the individual

reproductive values times the class size). In the models considered here, the dynamics are non-

linear because of competition for resources (N_{tot}). Nevertheless, we can approximate reproductive

values by assuming that the population is near equilibrium with only resident alleles and by holding

the strength of competition constant (see Supplementary *Mathematica* file for all calculations).

Doing so, we find that the class reproductive values of haploids and diploids, expressed as

proportions that sum to one, are:

$$c_H^G = \frac{(1 - a_H)\frac{f_R}{2}w_R(\hat{\rho}_H^G)^2}{(1 - a_H)\frac{f_R}{2}w_R(\hat{\rho}_H^G)^2 + (1 - a_D)w_{RR}(\hat{\rho}_D^G)^2}, \tag{2a}$$

$$c_D^G = \frac{(1 - a_D)w_{RR}(\hat{\rho}_D^G)^2}{(1 - a_H)\frac{f_R}{2}w_R(\hat{\rho}_H^G)^2 + (1 - a_D)w_{RR}(\hat{\rho}_D^G)^2}. \tag{2b}$$

where $\hat{\rho}_H^G = 1 - \hat{\rho}_D^G$ is the equilibrium frequency of haploids in the global model (Eq. A.4). As a

special case of interest, when populations are purely sexual ($a_H = a_D = 0$), we can plug the

equilibrium for $\hat{\rho}_H^G$ from Eq. (A.4) into (2) and show that $c_H^G = c_D^G = 1/2$.

As discussed in Appendix A (see also Bessho and Otto 2017), equations (2) provide the

only weights that allow ecological and evolutionary time scales to be separated when calculating

the average allele frequency in equation (1a), which is why we take that to be the evolutionarily

relevant average. Although one might initially think that diploids should count twice as much

because they contain two allele copies and that the evolutionarily relevant average allele frequency

would depend on the population sizes of haploids and diploids, a strict alternation of generations

($a_H = a_D = 0$) ensures that haploids and diploids contribute equally to long-term future

generations, so that their reproductive values are equal and the evolutionarily relevant average

allele frequency is $p_{ave} = (1/2)p_H + (1/2)p_D$ (Bessho and Otto 2017).

As with our previous model, we can track the slow evolutionary dynamics for the

expected change in average allele frequency p_{ave} under weak selection, once the fast ecological

dynamics have stabilized, as which point we can show that there are similar allele frequencies in

haploids and diploids ($\delta_p \approx 0$), diploids are approximately at Hardy-Weinberg equilibrium

($\eta_{HW} \approx 0$), and the ratio of haploids is similar among mutant and resident genotypes ($\rho_H \approx \hat{\rho}_H^G$)

(Appendix A, File S1). Furthermore, to leading order, the second moment of change in allele

frequency is equal to the neutral case and can be derived in the diffusion limit ($N_{tot} \to \infty$).

Given a single variable, p_{ave}, changing slowly over evolutionary time, we can then use

standard diffusion methods to calculate the fixation probability of a mutation in a haploid-diploid

population, $u(p_0^{Model})$, where "*Model*" is G for global and L for local (considered in the next

section). The diffusion is a function of the first and second moments of change in the mutant allele

frequency, $m^{Model}(p_{ave})$ and $v^{Model}(p_{ave})$, both measured in time units of N_{tot} generations:

$$u\left(p_0^{Model}\right) = \frac{\int_0^{p_0^{Model}} \exp[-2Q^{Model}(p')]\, dp'}{\int_0^1 \exp[-2Q^{Model}(p')]\, dp'}, \tag{3a}$$

$$m^{Model}(p_{ave}) = \frac{N_{tot}p_{ave}(1-p_{ave})}{2}\left[2s_{ave}^{Model} + 2c_D^{Model}p_{ave}(1-2h)s_{MM}^w\right], \tag{3b}$$

$$v^{Model}(p_{ave}) = \frac{p_{ave}(1-p_{ave})\left[\left(c_D^{Model}\right)^2 \hat{\rho}_H^{Model} + \left(c_H^{Model}\right)^2 \left(2\hat{\rho}_D^{Model}\right)\right]}{\hat{\rho}_H^{Model}\left(2\hat{\rho}_D^{Model}\right)}. \tag{3c}$$

162   where p_0^{Model} is the initial allele frequency of mutants (we focus on the case with a single initial

163   mutant   allele, $p_0^{Model} = 1/\left[N_{tot}\left(\hat{\rho}_H^{Model} + 2\hat{\rho}_D^{Model}\right)\right]$)   and $Q^{Model}(p) =$

164     $\int (m^{Model}(p)/v^{Model}(p))dp$. For the global regulation model, the average selection acting upon

165   rare mutant alleles across haploid and diploid stages, s_{ave}^G, can be calculated from the first moment

166   equation (Appendix 1) and equals:

$$s_{ave}^{Model} = c_H^{Model}s_M^w + c_D^{Model}s_{RM}^w + \phi^{Model}c_D^{Model}\frac{s_M^f}{2}, \tag{4a}$$

$$\phi^{Model} = \frac{(1-a_H)\widetilde{w}_R\hat{\rho}_H^{Model}}{(1-a_H)\widetilde{w}_R\hat{\rho}_H^{Model} + a_D w_{RR}\hat{\rho}_D^{Model}}. \tag{4b}$$

where $\widetilde{w}_R = f_R w_R/2$ is the fitness of haploids considering the cost of sex. As we will see later,

this equation is valid for local regulation model. The term ϕ^{Model} indicates the fraction of the

diploids in the next generation that come from the union of gametes rather than diploid asexual

reproduction. With obligately sexual haploid-diploids ($a_H = a_D = 0$, where $c_H^{Model} = c_D^{Model} =$

$1/2$ and $\phi^{Model} = 1$), these results coincide with those of Bessho and Otto (2017).

3.2. Fixation probability in the local regulation model

We next derive the fixation probability in a haploid-diploid population when density

dependence regulates haploid and diploid populations separately (Figure 1b), by again applying a

transformation of variables and separation of time scales. For the local regulation model, the

appropriate weights for the average allele frequency are similar to the global regulation model,

where now the class reproductive values, expressed as proportions, are:

$$c_H^L = \frac{1 + \dfrac{a_H w_R \hat{\rho}_H^L}{(1 - a_D) w_{RR} \hat{\rho}_D^L}}{2 + \dfrac{a_H w_R \hat{\rho}_H^L}{(1 - a_D) w_{RR} \hat{\rho}_D^L} + \dfrac{2}{f_R}\dfrac{a_D w_{RR} \hat{\rho}_D^L}{(1 - a_H) w_R \hat{\rho}_H^L}}, \tag{5a}$$

$$c_D^L = \frac{1 + \dfrac{2}{f_R}\dfrac{a_D w_{RR} \hat{\rho}_D^L}{(1 - a_H) w_R \hat{\rho}_H^L}}{2 + \dfrac{a_H w_R \hat{\rho}_H^L}{(1 - a_D) w_{RR} \hat{\rho}_D^L} + \dfrac{2}{f_R}\dfrac{a_D w_{RR} \hat{\rho}_D^L}{(1 - a_H) w_R \hat{\rho}_H^L}}. \tag{5b}$$

After applying a separation of time scales and conducting a diffusion approximation, we conclude

that the solution for the fixation probability in a haploid-diploid population, Eqs. (3), remains valid

for the local regulation model (Supplementary *Mathematica* file), with the average selection

coefficient now being given by Eqs. (5).

3.3. Effective genetic parameters

Using the first and second moments of change in allele frequency, we derive effective

genetic parameters to compare our results to the dynamics found in the classical model for fully

haploid or fully diploid organisms (Bessho and Otto 2017). More specifically, we define the

effective selection coefficient (s_e), dominance coefficient (h_e), and effective population size (N_e)

that would result in the same expected change in allele frequency and variance as in the classical

diploid model of selection.

For selection, the diploid model is: $\Delta p_{ave} = s_e p_{ave}(1 - p_{ave})[h_e + (1 - 2h_e)p_{ave}]$

(Crow and Kimura 1970; Bessho and Otto 2017). Because this equation depends on the allele

frequency in the same way as Eq. (3b), we can find the effective and dominance selection

coefficient from $\Delta p_{ave}/[p_{ave}(1-p_{ave})] = s_e h_e$ when $p_{ave} = 0$ and

$\Delta p_{ave}/[p_{ave}(1-p_{ave})] = s_e(1-h_e)$ when $p_{ave} = 1$, yielding:

$$s_e^{Model} = 2s_{ave}^{Model} + 2\hat{c}_D^{Model}\frac{(1-2h)s_{MM}^w}{2}. \tag{6a}$$

$$h_e^{Model} = \frac{2s_{ave}^{Model}}{4s_{ave}^{Model} + 2\hat{c}_D^{Model}(1-2h)s_{MM}^w}. \tag{6b}$$

When the mutation is additive ($h = 1/2$), these effective parameters are $s_e^{Model} = 2s_{ave}^{Model}$ and

$h_e^{Model} = 1/2$.

We next derive the variance effective population size by equating the one generation

change in variance (Eq. 3c divided by the time scale, N_{tot}) to the variance in allele frequency

expected in the classical Wright-Fisher model, $p_{ave}(1-p_{ave})/2N_e$ with N_e diploid individuals,

obtaining:

$$N_e^{Model} = \frac{p_{ave}(1-p_{ave})}{2(v^{Model}/N_{tot})} = \frac{N_{tot}\,\hat{\rho}_H^{Model}\hat{\rho}_D^{Model}}{\left(c_D^{Model}\right)^2\hat{\rho}_H^{Model} + 2\left(c_H^{Model}\right)^2\hat{\rho}_D^{Model}}. \tag{7}$$

203   Plugging these effective parameters into the formula from the fixation probability in the

204 classical diploid Wright-Fisher model (Kimura 1957; 1962; Crow and Kimura 1970, p. 427), the

205 fixation probability in a haploid-diploid population given by Eq. 3a can be expressed as:

$$u(p_0^{Model}) = \frac{\int_0^{p_0^{Model}} \exp[-2N_e^{Model}s_e^{Model}\{(2h_e^{Model}-1)p'(1-p')+p'\}]\,dp'}{\int_0^1 \exp[-2N_e^{Model}s_e^{Model}\{(2h_e^{Model}-1)p'(1-p')+p'\}]\,dp'}. \tag{8}$$

206 Assuming an initially rare and additive mutation ($h = 1/2$) with weak positive selection in a large

207 population ($s_e^{Model}N_e^{Model}p_0^{Model} \approx 0$ and $s_e^{Model}N_e^{Model} \gg 1$), we obtain the classic

208 approximation, $u(p_0^{Model}) \approx 2s_e^{Model}N_e^{Model}p_0^{Model}$, which upon substituting from Eq. 7 yields:

$$u(p_0^{Model}) \approx \frac{2\hat{\rho}_H^{Model}\hat{\rho}_D^{Model}}{(\hat{\rho}_H^{Model}+2\hat{\rho}_D^{Model})\left[\left(c_D^{Model}\right)^2\hat{\rho}_H^{Model} + 2\left(c_H^{Model}\right)^2\hat{\rho}_D^{Model}\right]}2s_{ave}^{Model}. \tag{9a}$$

209 For example, because haploids and diploids have the same reproductive values in the obligately

210 sexual case ($\hat{c}_H^{Model} = \hat{c}_D^{Model} = 1/2$), we obtain:

$$u(p_0^{Model}) \approx \frac{8\hat{\rho}_H^{Model}\hat{\rho}_D^{Model}}{\left(\hat{\rho}_H^{Model}+2\hat{\rho}_D^{Model}\right)^2}\,2s_{ave}^{Model} \tag{9b}$$

(Eq. 13a in Bessho and Otto 2017), or simply $u\left(p_0^{Model}\right) \approx 2s_{ave}^{Model}$ if haploid and diploid

population sizes are equal in terms of number of chromosomes ($\hat{\rho}_H^{Model} = 2/3$).

In the next three sections, we explore the implications of these results for the evolution

of haploid-diploid populations.

3.4. Effective selection in a haploid-diploid population

The strength of selection averaged across haploids and diploids, s_{ave}^{Model}, plays a key

role in the evolution of haploid-diploid populations. When a mutation is rare, both the rate of

change in allele frequency (Eq. 3b) and the approximate fixation probability (Eq. 10a) are

proportional to s_{ave}^{Model}. We thus begin by exploring how s_{ave}^{Model} varies as we alter the amount of

asexual reproduction in haploid and diploid phases. We focus on the case where the mutation does

not affect fertilization success ($s_M^f = 0$), so that the average selection becomes:

$$s_{ave}^{Model} = c_H^{Model} s_M^w + c_D^{Model} s_{RM}^w, \tag{10}$$

in both global and local regulation models (Eqs. (4) and (6)).

The relative evolutionary importance of selection in the haploid and diploid phases is

thus determined by the class reproductive values, c_H^{Model} and c_D^{Model} (where $c_H^{Model} +$

$c_D^{Model} = 1$). Figures 2 (global regulation) and 3 (local regulation) illustrate the proportional

reproductive value of haploids, c_H^{Model}, as a function of the degree of asexual reproduction in

haploids (x-axis) and diploids (ranging from 0.05 in red to 0.95 in blue). With global regulation,

the frequency of haploidy within the population, $\hat{\rho}_H^G$ (given by Eq. A.4), varies with the

parameters (see inset graphs in Figure 2), rising with the frequency of haploid asexuality (x-axis

in inset) but declining with more asexuality in diploids (from red to blue). By contrast, with local

regulation, the frequency of haploidy is held fixed by the strict density dependent competition that

we have assumed ($\hat{\rho}_H^L = 0.8$ in Figure 3(a)(b) and 0.3 in 3(c)(d)).

In the left panels, haploids have a higher fertility ($w_R/w_{RR} = 5$), leading to a higher

haploid reproductive value, c_H^{Model}, especially with local regulation when haploids are also more

common ($\hat{\rho}_H^L = 0.8$ in Figure 3a). In the right panels, diploids have a higher fertility ($w_{RR}/w_R =$

5), leading to a lower haploid reproductive value, especially when haploids are rare ($\hat{\rho}_H^L = 0.3$ in

Figure 3d).

When haploids are primarily sexual ($a_H \approx 0$), increasing asexuality of the haploid stage

typically causes the reproductive value of haploids to rise, unless diploids are fitter and more

frequent (Figure 3d and blue curves in Figure 2b). At the other extreme, the reproductive value of

haploids typically plummets to zero as haploid reproduction becomes primarily asexual ($a_H \approx 1$)

while diploids remain sexual, particularly with local regulation (Figure 3), because haploids then

act as a genetic "sink" contributing little to the diploid sub-population. This downward trend when

haploids are predominantly asexual is also seen with global regulation if diploids are more fit

(Figure 2b), except when the diploid population does not sustain itself and goes extinct, which

occurs when $a_D < 0.2$ and $a_H = 1$. The net result can thus be non-monotonic (purple curves

with $0.2 < a_D < 0.4$ in Figure 2b and Figure 3(a)(b)(c)).

3.4. Effective population size in a haploid-diploid population

We next consider the effective size of haploid-diploid populations with varying degrees

of asexuality. Figure 4 plots the effective population size (Eq. 8) relative to the total population

size, N_e^{Model}/N_{tot}, as a function of the frequency of haploids, $\hat{\rho}_H^{Model}$ (x-axis), and the class

reproductive values (with c_H^{Model} ranging from 0.05 in blue to 0.95 in red). As noted by Bessho

and Otto (2017), the effective population size is highest – and drift weakest – at intermediate

frequencies of haploids and diploids, which ensures the least sampling error as organisms alternate

generations.

When haploids and diploids have equal reproductive values, as in the fully sexual case

($c_H^{Model} = c_D^{Model} = 1/2$), the effective population size is maximized at $\hat{\rho}_H^{Model} \approx 0.586$. With

asexual reproduction, the peak shifts towards whichever ploidy level has the higher reproductive

value. For example, if haploids have a high reproductive value (red) then the effective population

size is maximized at a higher frequency of haploids, reducing the amount of genetic drift in that

phase. Although not illustrated, the peak shifts to $N_e^{Model} = \hat{\rho}_D^{Model} N_{tot}$ when future populations

descend only from diploids ($c_H^{Model} = 0$) and to $N_e^{Model} = \left(\hat{\rho}_H^{Model}/2\right)N_{tot}$ when future

populations descend only from haploids ($c_H^{Model} = 1$), effectively becoming diplontic or haplontic,

respectively (with the $1/2$ arising because haploids have half the number of chromosomes).

Of course, the reproductive values, as well as the frequency of haploids with global

population regulation ($\hat{\rho}_H^G$), depend in turn on the fitness parameters and the extent of asexuality,

as explored in the previous section. Figures 5 (global regulation) and 6 (local regulation) illustrate

the effective population size as a function of the frequency of haploid asexuality, a_H (x-axis), and

the frequency of diploid asexuality (a_D rising from red to blue), using the parameters in Figures 2

and 3, respectively. The trends are often non-monotonic, with N_e^{Model}/N_{tot} values varying

around $1/2$ when the parameter values are intermediate. The effective population size is often

higher when diploids rarely reproduce asexually (red) rather than when they frequently do (blue),

although there are exceptions (particularly when the fitness and frequency of diploids is high).

3.6. Fixation probability in a haploid-diploid population

We next compare the above results with numerical simulations estimating the fixation

probability of a newly arisen mutation in a haploid-diploid population. When simulating the global

regulation model, we assumed that the population has reached the demographic equilibrium,

$\hat{\rho}_H^G N_{tot}$ haploids and $\hat{\rho}_D^G N_{tot}$ diploids (see Appendix A). We then chose one resident allele R at

random and replaced it with a mutant allele M. After mutation, offspring were sampled from the

parental generation according to a multinomial distribution with expected frequencies given by

$x_{(GT)}$, repeating until the mutant allele fixed or was lost from the population. We estimated the

fixation probability as the fraction of 10,000 replicate simulations leading to fixation.

We here consider the additive case ($h = 1/2$ and $h_e = 1/2$), where the fixation

probability (Eq. 8) simplifies to:

$$u\left(p_0^{Model}\right) = \frac{\exp[-2p_0^{Model}N_e^{Model}s_e^{Model}] - 1}{\exp[-2N_e^{Model}s_e^{Model}] - 1} \qquad (11)$$

and where $s_e^{Model} = 2\,s_{ave}^{Model}$ (Eq. 6). Figure 7 plots the fixation probability as a function of the

average selection pressure, s_{ave}^{Model}, when the reproductive values and chromosome numbers in

haploids and diploids are equal ($c_H^{Model} = c_D^{Model}$ and $\hat{\rho}_H^{Model} = 2/3$) and $s_{ave}^{Model} = [(1/2)s_M^w +$

$(1/2)s_{RM}^w]$. The diffusion Eq. (11) provides an excellent fit, as does the approximation Eq. (9b)

for selection coefficients that are positive and not too weak. In this case, the results are the same

with global and local population regulation (Fig. 7a and 7b, respectively) and are insensitive to

how much selection occurs in the haploid or diploid phases (s_M^{Model} and s_{RM}^{Model}, respectively), as

long as s_{ave}^{Model} is held constant (see additional simulations in Supplementary *Mathematica* file).

As expected, the extent of selection in the haploid versus diploid phase matters more when the

mutation is not additive ($h \neq 1/2$ and $h_e \neq 1/2$) (supplementary *Mathematica* file).

Next, we illustrate the approximate fixation probability, Eq. (9a), as a function of the

degree of asexuality (a_H and a_D) when the population size is globally (Fig. 8) or locally (Fig. 9)

regulated, assuming only selection in haploids or only in diploids. For example, with additive

mutations, the fixation probability can be approximated as $u \approx 4c_H^{Model}N_e^{Model}p_0^{Model}s_M^w$ when

selection occurs only in the haploid phase or $u \approx 4c_D^{Model}N_e^{Model}p_0^{Model}s_{RM}^w$ with selection only

in the diploid phase, indicating that the fate of mutations depends as much on the strength of

selection as on the reproductive value of the ploidy phase in which selection acts (as illustrated in

Fig. 2 and 3). Figures 8 (global) and 9 (local) illustrate how the fixation probability depends on the

various parameters in the model, particularly the amount of asexual reproduction in haploids (x-

axis) and diploids (a_D rising from red to blue). The trends can be understood by the combined

effects of the parameters on the reproductive value and the effective population size (e.g., Fig. 9(a)

is proportional to the product of Fig. 3(a) and Fig. 6(c)).

4. Discussion

Across the phylogenetic tree of life, organisms have diverse and complex reproductive

strategies (Bell 1982). Classical population genetic theory has, however, focused most on fully

haploid or diploid life cycles with obligate sexuality. In this article we develop a stochastic model

for the population genetics of haploid-diploid organisms considering demography, asexuality, and

habitat differentiation between haploid and diploid stages. Using a separation of time scales, we

derive a diffusion approximation for the change in allele frequency, allowing us to estimate the

fixation probability of new mutations, the effective strengths of selection and dominance, as well

as the effective population size of haploid-diploid populations.

5.1. Natural selection in a haploid-diploid population

Our results indicate that the strength of natural selection and the extent of genetic drift

depend strongly on the reproductive value of haploid versus diploid phases. In the simplest case,

when the effect of a mutation is weak, additive, positive, and absent in the gamete stage ($s_M^f = 0$),

the fixation probability is proportional to the effective strength of selection (Eq. 10), $s_e^{Model} = $

$2s_{ave}^{Model}$, which in turn is proportional to the amount of selection in and the reproductive value of

haploids and diploids (Eqs. 2, and 5).

These analytical results reveal some evolutionary principles for populations that undergo

an alternation of generations. One consequence is that the balance of opposing selection pressures

in haploids and diploids (Eqs. 4 and 6) depends not only on the selection coefficients, but also on

the relative reproductive values of haploids (c_H^{Model}) versus diploids (c_D^{Model}). Thus, the very

direction of evolution depends on the extent of asexuality in the two phases and the relative survival

and fertility of haploids versus diploids when there is "ploidally antagonistic selection" (Immler et

al. 2012).

335   The efficacy of selection to fine tune traits in haploids and diploids also depends on the

class reproductive values. For example, when the population is regulated by local density

dependence (i.e., the haploid and diploid phases are spatially or temporally distinct), higher

reproductive success in haploids increases the efficiency of haploid selection (compare Figure 3a

to 3b). However, when there is extremely rare sexuality in haploids (a_H near one), diploid

selection tends to be more effective because of increasing competition between offspring from

haploids. By contrast, the trends differ with global density dependence (e.g., species that are more

isomorphic with small ecological differences between stages). For example, the reproductive value

of haploids remains high even when they reproduce primarily asexually in the global regulation

model (see Figure 2 when a_H approaches one), because haploids then make up a larger proportion

of the total population size (see inset figures). Thus, whether selection is effective in the haploid

phase when that phase mainly reproduces asexually is quite sensitive to the nature of competition.

347   Our work can also be useful in the design of field studies and the interpretation of data

for species that alternate generations. To understand the efficiency of selection on haploid and

diploid phases, we not only need data about the fraction of haploids and diploids and their fertility

and mortality (e.g., Thornber and Gains 2004; Vieira et al., 2018a; Vieira et al. 2018b), but we also

need to know about the extent of asexuality in each phase and whether they compete for common

or different resources.

5.2. Genetic drift and effective size

The impact of random genetic drift on the genetic diversity of haploid-diploid population

depends on the effective population size (Eq. 7). As we had found previously in a haploid-diploid

model with obligate sexuality (Bessho and Otto 2017, pp. 431), the effective population size with

asexuality is generally smaller than the total number of individuals and again depends strongly on

the reproductive value of each phase (Figures 4-6). With obligate sexuality, the reproductive values

of haploids and diploids are equal, and the effective population size is maximized (drift minimized)

when haploids comprise 2/3 of the population, making the number of chromosomes equal between

haploids and diploids. Asexual reproduction, however, causes the reproductive value of haploids

and diploids to differ (Eqs. 2 and 5). Consequently, drift is lessened if the phase with the higher

reproductive value is more common (see shifts in peaks in Figure 4).

5.3. "Ploidally-structured" population

The key role that reproductive values play in this work is analogous to the role that patch

dynamics play in two-patch models of evolution. In a spatially structured population, subdivided

local populations are genetically connected by migration. A haploid-diploid system can be seen as

being ploidally structured, where gene flow describes the movement of alleles through sexual

reproduction, with meiosis causing flow to haploidy and syngamy flow to diploidy. We note that

our research reveals that all qualitative results are equally accurate for evolution in a two-patch

system (see Supplementary *Mathematica* File). For example, fixation probability strongly depends

on class reproductive values of each patch.

This analogy suggests an interesting idea: complex reproductive systems can be

considered and analyzed using the tools of metapopulation theory. For example, many eukaryotes

including terrestrial plants, insects, and fishes, often exhibit ploidy variation, including polyploid

members (Otto and Whitton 2000; Comai 2005). In such species, individuals characterized by

different numbers of chromosomes coexist, with complex reproductive relationships causing gene

flow between them (Ramsey and Schemske 1998). Similarly, social insects often exhibit complex

sex determination systems linked with ploidy levels (haplodiploidy).

Our research suggests that these ploidally-structured populations can be fruitfully treated

as metapopulations. Selection and drift in populations with diploids, triploids, and tetraploids can,

for example, be considered as a three-patch model. In this system, we conjecture that the average

strength of selection that is evolutionarily relevant would be the mean selection coefficient in each

ploidy class, weighted by its class reproductive value, with additional terms coming from

reproductive interactions (akin to the term of s_M^f in Eqs. 4).

Many evolutionary aspects of haploid-diploid populations remain to be investigated. One

avenue that we are exploring is how model parameters can be estimated from field data. For

example, the analogy between spatially and ploidally structured population suggests that genetic

differences between haploids and diploid can be used to estimate gene flow between them (i.e.,

rates of sex), akin to using Fst to inform estimates of migration (e.g., Slatkin 1987). Another fruitful

avenue for further work is to determine how fluctuations in population size affect the effective

population size of species that alternate generations. In classical population genetics theory, such

fluctuations can be captured by using the harmonic mean population in place of the total population

size (Karlin 1968). It is unclear, however, whether the same is true in haploid-diploid populations.

Can the harmonic total population size simply replace N_{tot} in the global model of population

regulation? Similarly, can the harmonic population sizes of haploids and diploids replace N_H and

N_D with local regulation? The answer is unclear because population size fluctuations perturb the

fast ecological dynamics away from the steady state (especially $\hat{\rho}_H^{Model}$), and the impact of these

perturbations on selection and drift is unknown. Further research is needed to clarify evolutionary

processes in the wide variety of species that alternate generations.

Acknowledgements

We here thank Alireza Ghaseminejad for help in developing our manuscript. This project was

funded by a Grand-in-Aid from a Japan Society for the Promotion of Science (JSPS) to KB

(16J05204) and by a Discovery grant from the Natural Sciences and Engineering Research Council

of Canada (NSERCRGPIN-2016-03711) to S.P.O.

Appendix A. Fixation probability in a haploid-diploid Wright-Fisher model using a

diffusion approximation

A.1. Equilibrium with global regulation

We derive the fixation probability in a haploid-diploid population using a diffusion

approximation (e.g., Bessho and Otto 2017). We first derive the stable equilibrium in the global

regulation model, allowing for asexual reproduction in each phase. In the Wright-Fisher model,

all individuals reproduce and then the parents die (non-overlapping generations). Let $b_{(GT)}$

represent the number of reproductive cells of each type in the next generation:

$$b_R = (1 - a_D)w_{RR}x_{RR} + (1 - a_D)\frac{w_{RM}x_{RM}}{2} + a_H w_R x_R, \tag{A.1a}$$

$$b_M = (1 - a_D)\frac{w_{RM}x_{RM}}{2} + (1 - a_D)w_{MM}x_{MM} + a_H w_M x_M, \tag{A.1b}$$

$$b_{RR} = (1 - a_H)\frac{f_R}{2}\frac{w_R^2 x_R^2}{w_R x_R + w_M x_M} + a_D w_{RR}x_{RR}, \tag{A.1c}$$

$$b_{RM} = (1 - a_H)\frac{f_R + f_M}{2}\frac{w_R w_M x_R x_M}{w_R x_R + w_M x_M} + a_D w_{RM}x_{RM}, \tag{A.1d}$$

$$b_{MM} = (1 - a_H)\frac{f_M}{2}\frac{w_M^2 x_M^2}{w_R x_R + w_M x_M} + a_D w_{MM}x_{MM}. \tag{A.1e}$$

418   The probability that a reproductive cell of genotype (GT) is sampled from the offspring produced

419   by the previous generation of adults is

$$q_{(GT)} = \frac{b_{(GT)}}{b_R + b_M + b_{RR} + b_{RM} + b_{MM}}.$$
    (A.2)

420   Therefore, the composition of offspring in the next generation is given by the multinomial

421   distribution, sampling N_{tot} individuals in proportion to Eq. (A2). Using Eq. (A1) and (A2), we

422   describe the conditional expectation of change in the number of individuals of genotype (GT),

423    $\Delta X_{(GT)}(t) = X_{(GT)}(t+1) - X_{(GT)}(t)$, as

$$E[\Delta X_{(GT)}(t)|\vec{X}(t) = \vec{x}] = N_{tot}q_{(GT)} - x_{(GT)},$$
    (A.3)

424   where $E\left[\Delta F\left(X_{(GT)}(t)\right)|\vec{X}(t) = \vec{x}\right]$ is the conditional expected value for change in the function

425   F of the random variable given that $\vec{X}(t) = (X_R(t) \quad X_M(t) \quad X_{RR}(t) \quad X_{RM}(t) \quad X_{MM}(t))^T$

426   equals $\vec{x} = (x_R \quad x_M \quad x_{RR} \quad x_{RM} \quad x_{MM})^T$.

427   To simplify this fully stochastic system, we assume that the resident population is large

428   and treat demographic changes deterministically prior to the appearance of the mutation.

429   Considering the dynamics of the resident population, we then find the equilibrium of these

430   dynamical equations by solving $N_{tot}q_R - \hat{x}_R = 0$ and $N_{tot}q_{RR} - \hat{x}_{RR} = 0$ ($\hat{x}_R + \hat{x}_{RR} = N_{tot}$).

431   Setting $\hat{x}_R = \hat{\rho}_H^G N_{tot}$ and $\hat{x}_{RR} = \hat{\rho}_D^G N_{tot}$, the fraction of haploids $\hat{\rho}_H^G$ (and diploids $\hat{\rho}_D^G = 1 -$

432    $\hat{\rho}_H^G$) at equilibrium becomes,

$$\hat{\rho}_H^G = \frac{a_H w_R + a_D w_{RR} - 2w_{RR} + \sqrt{4(1 - a_H)(1 - a_D)\widetilde{w}_R w_{RR} + (a_H w_R - a_D w_{RR})^2}}{2[a_H w_R + (1 - a_H)\widetilde{w}_R - w_{RR}]},$$
    (A.4)

where $\widetilde{w}_R = f_R w_R/2$ is the fitness of haploids considering the cost of sex (see Supplementary

Mathematica file for the step-by-step derivation). We note that, when the fertility of haploids is

much greater than that of diploids ($w_R \gg w_{RR}$), the frequency of haploids in a population

approaches $a_H/\{a_H + [(1 - a_H)f_R/2]\}$, which is less than one because sexual reproduction of

the haploids produces diploids (the $(1 - a_H)f_R/2$ term). Conversely, when the fertility of

diploids is much greater than haploids ($w_R \ll w_{RR}$), the frequency of haploids approaches $1 -$

a_D, the rate at which diploids undergo meiosis.

A.2. First moment of change in allele frequency

442   To derive the first moment of change in allele frequency, $m^{Model}(p_{ave})$, we apply a

443   separation of time scales (e.g., Nagylaki 1976; Otto and Day 2007; Bessho and Otto 2017).

444   Details of the calculation are represented in the Supplementary *Mathematica* file. We first

445   transform the expected change in the number of individuals of each type (five variables that sum

446   to N_{tot}) into the expected change in a new set of four variables, $\Theta \in \{p_{ave}, \delta_p, \eta_{HW}, \rho_H\}$,

447   described by the functions:

$$\mathrm{E}[\Delta\Theta|\vec{X}(t) = \vec{x}] = f_\Theta^{Model}(\epsilon, p_{ave}, \vec{\theta}), \tag{A.5}$$

448   where $\vec{\theta} = (\delta_p, \eta_{HW}, \rho_H)$ and ϵ is proportional to the selection coefficients and assumed small

449   (the functions f are given explicitly in the Supplementary *Mathematica* file). With local

450   regulation, ρ_H is assumed fixed at N_H/N_{tot} and dropped from the variable set, Θ.

451   To constant order (setting the small changes due to selection to zero, $\epsilon \to 0$), the fast

ecological dynamics of the system are described by: $f_\Theta^{Model}(0, p_{ave}, \vec{\theta})$. This system of

equations rapidly approaches a steady state found by solving $f_\Theta^{Model}(0, p_{ave}, \vec{\theta}) = 0$, which

gives $\bar{\delta}_p = \bar{\eta}_{HW} = 0$, and $\bar{\rho}_H = \hat{\rho}_H^G$ (Eq. A.4). To this order, the steady state change in allele

frequency is zero, $f_{p_{ave}}^{Model}(0, p_{ave}, \vec{\theta}) = 0$. We then describe slower changes, including changes

in allele frequency due to selection, by describing the deviations that occur around this steady

state. Specifically, to order ϵ, the variables are allowed to deviate from the steady state by $\delta_p =$

$\tilde{\delta}_p\epsilon$, $\eta_{HW} = \tilde{\eta}_{HW}\epsilon$, and $\rho_H = \hat{\rho}_H^G + \tilde{\rho}_H\epsilon$, and the dynamics $f_\Theta^{Model}(\epsilon, p_{ave}, \vec{\theta})$ are then

approximated using a Taylor series expansion. Defining the average allele frequency by

combining haploid and diploid populations using an arbitrary weighting, $p_{ave} = \omega p_H +$

$(1 - \omega)p_D$, we show in the Supplementary *Mathematica* file that setting the weights proportional

to the class reproductive values (given by Eq. (2) with global regulation and Eq. (5) with local

regulation) is the only choice that separates evolutionary change in p_{ave} from changes in the

other variables to order ϵ. Defining the average allele frequency in this way (Eq. 1a) and taking

the Taylor series, the change in allele frequency becomes:

$$E[\Delta p_{ave}|\vec{X}(t) = \vec{x}] \approx M^{Model}(p_{ave}) \tag{A.6}$$

$$= \frac{p_{ave}(1 - p_{ave})}{2}\left[2s_{ave}^{Model} + 2c_D^{Model}p_{ave}(1 - 2h)s_{MM}^w\right].$$

A.3. Second moment of change in average allele frequency

We next derive the second moment of change in average allele frequency in a haploid-

diploid population with asexuality. We again assume that the population size is very large, that

selection is very weak, and that the system has approached the steady state in $(\delta_p, \eta_{HW}, \rho_H)$,

ignoring deviations that are of $O(\epsilon)$. Because selection is assumed weak, the second moment is

well approximated by that of the neutral model (to constant order, $\epsilon \to 0$).

Under these assumptions, the fraction of haploids in a population is relatively fixed in

both the global and local regulation models, and we can sample the haploid offspring according

to a binomial distribution, with expectation and variance: $E[X_M|\vec{X}(t) = \vec{x}] = q_M N_H$ and

$Var[X_M|\vec{X}(t) = \vec{x}] = q_M(1 - q_M)N_H$ where $q_M = b_M/(b_R + b_M)$. To simplify the equation,

we set $E[X_{(GT)}|\vec{X}(t) = \vec{x}] = m_{(GT)}$ and $Var[X_{(GT)}|\vec{X}(t) = \vec{x}] = v_{(GT)}$, finding that:

$$E[\Delta X_M|\vec{X}(t) = \vec{x}] = m_M - x_M, \tag{A.7a}$$

$$E[(\Delta X_M)^2|\vec{X}(t) = \vec{x}] = v_M + m_M^2 - 2m_M x_M + x_M^2. \tag{A.7b}$$

In terms of allele frequencies (rather than numbers), we have the first and second moments for

the haploid offspring population, $E\left[\frac{\Delta X_M}{N_H}|\vec{X}(t) = \vec{x}\right] = \frac{1}{N_H}E\left[\Delta X_M|\vec{X}(t) = \vec{x}\right]$ and

$E\left[\left(\frac{\Delta X_M}{N_H}\right)^2|\vec{X}(t) = \vec{x}\right] = \frac{1}{N_H^2}E\left[(\Delta X_M)^2|\vec{X}(t) = \vec{x}\right]$.

Similarly, the diploid offspring are sampled according to a trinomial distribution, with

expectation, variance, and covariance: $m_{(GT)} = q_{(GT)}N_D$, $v_{(GT)} = q_{(GT)}\left(1 - q_{(GT)}\right)N_D$, and

$Cov\left[X_{RM}, X_{MM}|\vec{X}(t) = \vec{x}\right] = q_{RM}q_{MM}N_D$, where $q_{(GT)} = b_{(GT)}/(b_{RR} + b_{RM} + b_{MM})$. To derive

the moments of the allele frequency in diploids, we define, $y_M = (x_{RM}/2) + x_{MM}$ and $Y_M =$

$(X_{RM}/2) + X_{MM}$. The moments of random variable Y are then:

$$E\left[Y_M|\vec{X}(t) = \vec{x}\right] = \frac{m_{RM}}{2} + m_{MM}, \tag{A.8a}$$

$$E\left[Y_M^2|\vec{X}(t) = \vec{x}\right] = \frac{v_{RM} + m_{RM}^2}{4} + \left(Cov\left[X_{RM}, X_{MM}|\vec{X}(t) = \vec{x}\right] + m_{RM}m_{MM}\right) + (v_{MM} + m_{MM}^2), \tag{A.8b}$$

$$E\left[\Delta Y_M|\vec{X}(t) = \vec{x}\right] = \frac{E\left[\Delta X_{RM}|\vec{X}(t) = \vec{x}\right]}{2} + E\left[\Delta X_{MM}|\vec{X}(t) = \vec{x}\right] - y_M, \tag{A.8c}$$

$$E\left[(\Delta Y_M)^2|\vec{X}(t) = \vec{x}\right] = E\left[Y_M^2|\vec{X}(t) = \vec{x}\right] - 2E\left[Y_M|\vec{X}(t) = \vec{x}\right]y_M + y_M^2. \tag{A.8d}$$

486   To consider the change in average allele frequency across the entire population, we

487   define $Z_M = c_H^{Model}\frac{X_M}{N_H} + c_D^{Model}\frac{Y_M}{N_D}$ and consider the expectation of change in this random

488   variable. Plugging in Eqs. (7a), (7b), (8c), and (8d), we have

$$E\left[(\Delta Z_M)^2|\vec{X}(t) = \vec{x}\right] = \frac{p_{ave}(1 - p_{ave})\left[\left(c_D^{Model}\right)^2\hat{\rho}_H^{Model} + \left(c_H^{Model}\right)^2\left(2\hat{\rho}_D^{Model}\right)\right]}{\hat{\rho}_H^{Model}\left(2\hat{\rho}_D^{Model}\right)N_{tot}}. \tag{A.9}$$

489   After transforming time scales using the variable $\tau = t/N_{tot}$ and defining $P(\tau) = Z(N_{tot}\tau)$,

490   we have the diffusion coefficient $v^{Model}(p_{ave}) = \lim_{N_{tot}\to\infty} E\left[\frac{(P(\tau+\Delta\tau)-P(\tau))^2}{\Delta\tau}\right]$ by taking the limit

491    $N_{tot} \to \infty$, giving Eq. (3c). Similarly, we derive the drift coefficient using Eq. (A6) ($m^{Model} =$

492    $M^{Model}N_{tot}$), giving Eq. (3b).

References

Bell, G. 1982. The Masterpiece of Nature: The Genetics and Evolution of Sexuality. University of California Press. Berkeley, CA.

Bell, G., 1994. The comparative biology of the alternation of generations. Lectures on mathematics in the life sciences 25, 1–26.

Bessho, K., Otto, S. P. 2017. Fixation probability in a haploid-diploid population. Genetics 205, 421-440.

Comai, L. 2005. The advantages and disadvantages of being polyploid. Nature Reviews Genetics 6, 836-846.

Coelho, S. M., Peters, A. F., Charrier, B., Roze, D., Destombe, C., Valero, M., Cock, J. M., 2007. Complex life cycles of multicellular eukaryotes: new approaches based on the use of model organisms. Gene 406, 152–170.

Crow, J. F., Kimura, M. (1970). An introduction to population genetics theory. The Blackburn Press, New Jersey.

Deithier, M. N. (1981). Heteromorphic algal life histories: The seasonal pattern and response to herbivory of the brown crust, *Ralfsia californica*. Oecologia 49, 333-339.

Destombe, C., Godin, J., Nocher, M., Richerd, S., Valero, M. (1993). Differences in response between haploid and diploid isomorphic phases of *Gracilaria verrucosa* (Rhodophyta: Gigartinales) exposed to artificial environmental conditions. Hydrobiologia 260/261, 131-137.

De Wreede, R. E., Klinger, T. 1988. Reproductive strategies in algae. Plant reproductive ecology: Patterns and strategies, 267-284.

Drew, K. M. (1949). Conchocelis-phase in the life-history of *Porphyra umbilicalis* (L.) Kütz. Nature 164, 748-749.

Dyck, L. J., De Wreede, R. E. (2006). Seasonal and spatial patterns of population density in the marine macroalga *Mazzaella splendens* (Gigartinales, Rhodophyta). Phycological Research, 54(1), 21-31.

Goldstein, D. B. 1992. Heterozygote advantage and the evolution of dominant diploid phase. Genetics 132, 1195–1198.

Haghes, J. S., Otto, S. P. 1999. Ecology and the evolution of biphasic life cycles. The American Naturalist 154, 306–330.

Hannach, G., Santelicesm, B. 1985. Ecological differences between the isomorphic reproductive phases of two species of *Iridaea* (Rhodophyta: Gigartinales). Marine Ecology Progress Series 22, 291–303.

Hawkes, M. W. 1990. Reproductive strategies. Biology of the red algae, 455-476.

529 Immler, S., G. Arnqvist, and S. P. Otto. 2012. Ploidally antagonistic selection maintains stable
530   genetic polymorphism. 66:55–65.

531 Jenkins, C. D., Kirkpatrick, M., 1995. Deleterious mutation and the evolution of genetic life
532   cycles. Evolution, 512–520.

533 Karlin, S. 1968. Rates of approach to homozygosity for finite stochastic models with variable
534   population size. The American Natualist 102, 443-455.

535 Kimura, M., 1957. Some problems of stochastic processes in genetics. The Annals of
536   Mathematical Statistics, 882-901.

537 Kimura, M., 1962. On the probability of fixation of mutant genes in a population. Genetics 47,
538   713.

539 Mable, K. B., Otto, P. S., 1998. The evolution of life cycles with haploid and diploid phases.
540   BioEssays 20, 453–462.

541 Nagylaki, T., 1976. The evolution of one-and two-locus systems. Genetics 83, 583-600.

542 Orr, H. A., Otto, S. P. 1994. Does diploidy increase the rate of adaptation? Genetics 136, 1475–
543   1480.

544 Otto, S., Goldstein, D., 1992. Recombination and the evolution of diploidy. Genetics 131, 745–
545   751.

546 Otto, S., 1994. The role of deleterious and beneficial mutations in the evolution of ploidy levels.
547   Lectures on Mathematica in the Life Sciences 25, 69–98.

548 Otto, S. P., Marks, J. C. 1996. Mating system and the evolutionary transition between haploidy
549   and diploidy. Biological Journal of the Linnean Society 57, 197-218.

550 Otto, S. P., Day, T., 2007. A biologist's guide to mathematical modeling in ecology and
551   evolution. Princeton University Press.

552 Otto, S. P., Whitton, J. 2000. Polyploid incidence and evolution. Annu. Rev. Genet. 34, 401-437.

553 Perrot, V., Richerd, S., Valéro, M., 1991. Transition from haploidy to diploidy. Nature 351, 315–
554   317.

555 Ramsey, J., Schemske, D. W. 1998. Pathways, mechanisms, and rates of polyploid formation in
556   flowering plants. Annual reviews of Ecology and Systematics 29: 467-501.

557 Raper, J. R., Flexer, A. S. (1970). The road to diploidy with emphasis on a detour. In *Symp. Soc.*
558   *Gen. Microbiol* (Vol. 20, pp. 401-432).

559 Rousset, F. (2004). Genetic structure and selection in subdivided populations. Princeton
560   University Press, New Jersey.

561 Scott, M. F., Rescan, M. 2016. Evolution of haploid-diploid life cycles when haploid and diploid
562   fitnesses are not equal. Evolution 71, 215–226.

563 Slatkin, M. 1987. Gene flow and the geographic struture of natural populations. Science New
564   Series 236: 787-792.

565 Slocum C. J. 1980. Differential susceptibility to grazers in two phases of an intertidal alga:

566   Advantages of heteromorphic generations. Journal of Experimental Marine Biology and

567   Ecology 46, 99-110.

568 Taylor, P. D. 1980. Allele-Frequency Change in a Class-Structured Population. The American

569   Naturalist 135, 95-106.

570 Thornber, C. S., Gains, S. D. 2004. Population demographics in species with biphasic life cycle.

571   Ecology 85, 1661-1674.

572 Thornber, C., Stachowicz, J. J., Gains, S. 2006. Tissue type matters: selective herbivory on

573   different life history stages of an isomorphic alga. Ecology 87, 2255-2263.

574 Vieira, V. M. N. C. S., Engelen, A. H., Huanel, O. R., Guillemin, M. 2018a. Haploid females in

575   the isomorphic biphasic life-cycle of *Gracilaria chilensis* excel in survival. BMC

576   Evolutionary Biology 18:174.

577 Vieira, V. M. N. C. S., Engelen, A. H., Huanel, O. R., Guillemin, M. 2018b. Differentiation of

578   haploid and diploid fertilities in *Gracilaria chilensis* affect ploidy ratio. BMC

579   Evolutionary Biology 18:183.

580 Willson, M. F. 1981. On the evolution of complex life cycles in plants: a review and an

581   ecological perspective. Annals of the Missouri Botanical Garden 68, 275-300.

Figure Captions

Fig. 1. An illustration of the haploid-diploid models. (a) In the global regulation model, both

haploids and diploids occupy the same habitat and density dependence holds the total population

size N_{tot} constant. (b) In the local regulation model, each ploidy stage occupies a different

habit, therefore density dependence regulates the population size of haploids (N_H) and diploids

(N_D) separately.

Fig. 2. Class reproductive value of haploids in the global regulation model, c_H^G. Curves show c_H^G

as a function of the degree of haploid asexuality, a_H (x-axis), with the degree of diploid

asexuality ranging in color from $a_D = 0.05$ (red) to 0.95 (blue) in increments of 0.05. Other

parameters are set as: (a) $f_R = 0.5$, $w_R = 5000$, $w_{RR} = 1000$, (b) $f_R = 0.5$, $w_R = 1000$,

$w_{RR} = 5000$. The resulting frequency of haploids, $\hat{\rho}_H^G$ (Eq. A.4), is shown in the inset plots.

Fig. 3. Proportional reproductive value of haploids in the local regulation model, c_H^L. Parameters

are the same as Fig. 2, except that haploids are held fixed at a frequency of (a)(b) $\hat{\rho}_H^L = 0.8$ or

(c)(d) $\hat{\rho}_H^L = 0.3$. We consider the case when (a)(c) haploid fitness parameter is larger than

diploid $w_R = 5000$, $w_{RR} = 1000$ ($w_H > w_D$), and when (b)(d) diploid fitness parameter is

larger than haploid $w_R = 1000$, $w_{RR} = 5000$ ($w_H < w_D$).

Fig. 4. The effective population size of a haploid-diploid population. The relative effective

population size over the total population size (N_e^{Model}/N_{tot}, Eq. 7) is shown as a function of the

frequency of haploids ($\hat{\rho}_H^{Model}$, x-axis), when the haploid reproductive value (c_H^{Model}) varies from

0.05 (blue) to 0.95 (red) in increments of 0.05. This figure applies to both global and local

regulation models.

Fig. 5.   The effective population size of a haploid-diploid population in the global regulation

model as a function of the degree of asexuality, a_H and a_D. Parameters are the same as in Fig. 2

and determine the relative class reproductive values (c_H^G) and fraction of haploids ($\hat{\rho}_H^G$) according

to Eqs. (2a) and (A.4).

Fig. 6. The effective population size of a haploid-diploid population in the local regulation

model. Parameters are the same as in Fig. 3 and determine the relative class reproductive values

(c_H^L) by Eqs. (5). The frequencies of haploids are held fixed at a frequency of (a)(b) $\hat{\rho}_H^L = 0.9$,

(c)(d) $\hat{\rho}_H^L = 0.8$, (e)(f) $\hat{\rho}_H^L = 0.6$, or (g)(h) $\hat{\rho}_H^L = 0.3$.

Fig. 7. Fixation probability given the average strength of selection, s_{ave}^{Model}, for global (a) and

local (b) regulation models. The solid curve gives the analytical result from the diffusion

approximation (Eq. (11)) and the dashed curve gives the linear approximation (Eq. (9b)). Black

dots indicate the fixation probability estimated from 10000 numerical simulations with 95% CI

(Wilson score interval for binomial). Parameters: $N_{tot} = 90$, $N_H = 60$, $N_D = 30$, $f_R = 0.5$,

$w_R = w_{RR} = 1000$, $a_H = a_D = 0.1$, $h = 0.5$, $s_M^f = 0$, $s_M^w = s_{RM}^w = s_{ave}^{Model}$, such that the

fraction of haploids in the resident population is $\hat{\rho}_H^{Model} = 2/3$ and class reproductive values

are equal $c_H^{Model} = c_D^{Model} = 1/2$. Holding $s_{ave}^{Model} = 1/2(s_M^w + s_{RM}^w)$ constant, similar results

are obtained for a range of different choices of s_M^w and s_{RM}^w (see supplementary *Mathematica*

file).

Fig. 8. The fixation probability in a haploid-diploid population in the global regulation model.

Curves gives the linear approximation for the fixation probability (Eq. (9a)). Parameters are the

same as in Fig. 2. Selection acts only in the haploid or diploid phase, with selection coefficients

set as $s_M^w = 0.02$ for haploid selection (a)(b) and $s_{RM}^w = 0.02$ for diploid selection (c)(d).

Fig. 9. The fixation probability in a haploid-diploid population in the local regulation model.

Curves gives the linear approximation for the fixation probability (Eq. (9a)). Parameters are the

same as in Fig. 3 and Fig. 8. The frequency of haploids is held fixed at (a)(b)(e)(f) $\hat{\rho}_H^L = 0.8$,

(c)(d)(g)(h) $\hat{\rho}_H^L = 0.3$. Fitness of haploids is higher than diploids in panels (a)(c)(e)(g) ($w_H >$

w_D), and the opposite condition is considered in panels (b)(d)(f)(h) ($w_H < w_D$). Selection only

occurs in the haploid (a)(b)(c)(d) or diploid stage (e)(f)(g)(h).

Supporting information

S1. Supplementary *Mathematica* file.

Figure1

(a)   (b)

Global regulation   **Local regulation**

Total population size: N_{tot} N_{H} N_{D}

Bessho and Otto

Figure 2

Bessho and Otto

Figure 3

Bessho and Otto

Figure 4

Bessho and Otto

Figure 5

Bessho and Otto

Figure 6

Figure 7

(a) global regulation

(b) local regulation

Bessho and Otto

Figure 8

Bessho and Otto

Figure 9