Fixation and effective size in a haploid-diploid population with asexual reproduction

*Kazuhiro Bessho, **Sarah P. Otto

* Saitama Medical University, 38 Morohongo Moroyama-machi, Iruma-gun, Saitama 350-0495,

Japan

**Department of Zoology, The University of British Columbia, Vancouver, BC, V6T 1Z4,

Canada

* besshokazuhiro.reserch@gmail.com; (+81) 049-276-2030

** otto@zoology.ubc.ca; (+1) 604-822-2778

*Corresponding author

besshokazuhiro.reserch@gmail.com (KB)

*Co-corresponding author

otto@zoology.ubc.ca (SPO)

*Keywords***:** haploid-diploid life cycle, Wright-Fisher model, fixation probability, variance

effective population size, class reproductive value

Highlight

- Classical models consider fully haploid or diploid populations

- We model haploid-diploid life cycles allowing for asexual reproduction

- We obtain the fixation probability of alleles subject to selection and drift

- Reproductive values of haploid and diploid stages shape their evolution

Abstract

The majority of population genetic theory assumes fully haploid or diploid organisms with obligate sexuality, despite complex life cycles with alternating generations being commonly observed. To reveal how natural selection and genetic drift shape the evolution of haploid-diploid populations, we analyze a stochastic genetic model for populations that consist of a mixture of haploid and diploid individuals, allowing for asexual reproduction and niche separation between haploid and diploid stages. Applying a diffusion approximation, we derive the fixation probability and describe its dependence on the reproductive values of haploid and diploid stages, which depend strongly on the extent of asexual reproduction in each phase and on the ecological differences between them.

1. Introduction

Sexual reproduction in eukaryotes generally consists of an alternation of generations, where meiosis halves the number of chromosomes to produce haploids and syngamy brings together haploid gametes to produce diploids. The extent of development in each ploidy phase varies substantially (Bell 1982; 1994). In diplontic organisms, at one extreme, development and growth occur only in the diploid phase, as is observed in most animals. Haplontic organisms, at the other extreme, undergo mitotic growth only in the haploid stage, as is seen in some green algae. In between these extremes, many terrestrial plants, macroalgae, and fungi exhibit both haploid and diploid growth (haploid-diploid life cycles). These stages are typically free living in macroalgae, with either macroscopically similar (isomorphic) or distinct (heteromorphic) forms in the haploid and diploid stage (Raper and Flexer 1970; Wilson 1981; Mable and Otto 1998; Coelho 2007).

To explain variation in life cycles, several theoretical models have analyzed the deterministic dynamics of a modifier allele that alters the time spent in haploid and diploid phases (e.g., Perrot et al. 1991; Otto and Goldstein 1992; Goldstein 1992; Otto 1994; Orr and Otto 1994; Jenkins and Kirkpatrick 1995; Otto and Marks 1996; Scott and Rescan 2017). However, there are some gaps between these models and the complexities seen in many haploid-diploid species. For example, these models often treat haploid and diploid individuals as ecologically equivalent, despite the frequent observation of niche differences and seasonal shifts in prevalence (e.g., Slocum 1980; Dethier 1981). Most of these models also assume obligate sexuality (but see Otto and Marks 1996), despite asexuality being frequently observed among haploid-diploid species ("asexual looping"). Furthermore, while several models have explored how haploid-diploid life cycles might evolve, the impact of haploid-diploid life cycles on

evolutionary processes remains underexplored (see, e.g., Bessho and Otto 2017 on the impact on

fixation probabilities and Immler et al. 2012 on the maintenance of variation).

Here we contribute to evolutionary theory for haploid-diploid populations by

calculating the fixation probability of mutations using a stochastic genetic model. This builds

upon our previous work (Bessho and Otto 2017) by accounting for asexual looping and niche

differences between ploidy phases, both of which are common in macroalgae (Bell 1982; de

Wreede and Klinger 1988; Hawkes 1990). Haploid and diploid phases often differ

physiologically, and even isomorphic haploids and diploids may differ ecologically (Hannach

and Santelices 1985; Destombe et al. 1993; Dyck and de Wreede 2006; Thornber et al., 2006;

Vieira et al. 2018). We therefore explore different forms of density dependence, acting either

globally on the total population size (as in Bessho and Otto 2017) or locally on the population

size of haploids and diploids separately (Figure 1). We show that the fate of a mutation depends

strongly on the reproductive values of haploids and diploids, which in turn depend on the extent

of asexual reproduction and ecological differences between the phases.

2. Model

In Bessho and Otto (2017), we calculated the fixation probabilities by tracking the

dynamics of a resident allele (*R*) and a mutant allele (*M*) in haploid and diploid individuals, using

both a Wright-Fisher and a Moran model. In that model, reproduction was obligately sexual,

individuals were ecologically equivalent, and the total population size was held constant (global

density dependence). Below, we calculate the fixation probability by first considering asexual

reproduction in each phase, assuming that haploids and diploids are ecologically equivalent

(global population regulation), and then determine how these results are affected by niche

differences (local population regulation that is ploidy specific).

2.1. Haploid-diploid Wright-Fisher model with global regulation and asexual looping

Let $X_{(GT)}(t)$ be a random variable that represents the number of individuals of a

particular genotype ("*GT*") at time t, involving the resident (*R*) and mutant alleles (*M*) (e.g., *GT*

= *RM* for a heterozygous diploid and *GT* = *M* for a mutant haploid). Let $x_{(GT)}(t)$ represent a

particular outcome of this random variable. In the global regulation model, we assume a constant

population, $x_R + x_M + x_{RR} + x_{RM} + x_{MM} = N_{tot}$, that is strictly regulated regardless of the

ploidy of the individuals.

The reproductive output and the degree of asexuality are characterized by $w_{(GT)}$ and

a_H for haploids [$(GT) = R$ or M] and $w_{(GT)}$ and a_D for diploids [$(GT) = RR$, *RM*, and *MM*].

Specifically, diploid individuals produce $(1 - a_D)w_{(GT)}$ haploid spores (sexual reproduction)

and $a_D w_{(GT)}$ diploid offspring (asexual loop). Similarly, haploids produce $(1 - a_H)w_{(GT)}/2$

female gametes (sexual reproduction) and $a_H w_{(GT)}$ haploid offspring (asexual loop), where we

assume that the species is monoecious and invests equal resources in male and female gametes.

During syngamy, we assume that male gametes are not limiting, that mating is random, and that

female gametes are successfully fertilized with male gametes, at a rate $f_{(GT)}$ [$(GT) = R$ or M],

becoming diploid zygotes. For clarity, we describe the model with non-overlapping generations,

although we note that overlapping generations can be considered by including surviving adults in

the counts of asexual offspring ($a_D w_{(GT)}$ and $a_H w_{(GT)}$).

We define the selection coefficient ($s_{(GT)}$) and the degree of dominance (*h*) acting upon

the mutant allele such that: $f_M = f_R\left(1 - s_M^f\right)$, $w_M = w_R(1 - s_M^w)$, $w_{RM} = w_{RR}(1 - s_{RM}^w)$,

$w_{MM} = w_{RR}(1 - s_{MM}^w)$, and $h = s_{RM}^w/s_{MM}^w$. To perform the diffusion approximation, we assume

that selection is weak, $s_M^f = \epsilon \tilde{s}_M^f$ and $s_{(GT)}^w = \epsilon \tilde{s}_{(GT)}^w$, where ϵ is a small parameter (parameter

ϵ is order $O(1/N_{tot})$).

2.2. Haploid-diploid Wright-Fisher model with local regulation and asexual looping

116    We then consider the case where density dependence regulates haploid and diploid

populations separately, which may occur if they have different resource needs or utilize different

habitats or microhabitats (for short-hand, we refer to this case as "local regulation"). More

specifically, we assume that the population size of haploids and diploids is separately regulated

and remains constant N_H and N_D ($x_R + x_M = N_H$ and $x_{RR} + x_{RM} + x_{MM} = N_D$), respectively.

We set $N_H + N_D = N_{tot}$, $\hat{\rho}_H^L = N_H/N_{tot}$, and $\hat{\rho}_D^L = N_D/N_{tot}$, which will then allow us to

compare the results of local and global regulation. Holding population sizes constant is assumed

strictly for mathematical convenience but may be reasonable for populations whose sizes are

strongly regulated by the availability of appropriate habitat.

3. Fixation probability in a haploid-diploid population

3.1. Fixation probability in the global regulation model

128    The fixation probability in a haploid-diploid population can be derived using a

diffusion approximation (Bessho and Otto 2017), but doing so requires that we approximate the

dynamics to reduce the dimensionality from four variables ($x_R, x_M, x_{RR}, x_{RM}, x_{MM}$, which sum to

N_{tot}) down to one. We do so by using a separation of time scales, deriving the first and second

moments of the mutant allele frequency. Specifically, we transform the number of individuals of

each genotype, $x_{(GT)}$, into new variables that allow us to separate the slower evolutionary

dynamics and the faster ecological dynamics (Appendix A):

$$p_{ave} = c_H\, p_H + c_D\, p_D, \tag{1a}$$

$$\delta_p = p_D - p_H, \tag{1b}$$

$$\eta_{HW} = 1 - \frac{1}{2p_D(1-p_D)} \frac{x_{RM}}{x_{RR} + x_{RM} + x_{MM}}, \tag{1c}$$

$$\rho_H = \frac{x_R + x_M}{N_{tot}}, \tag{1d}$$

where p_{ave} indicates the average allele frequency of haploids and diploids weighted by the class

reproductive values (c_H and c_D, where $c_H + c_D = 1$, see next paragraph), δ_p indicates the

difference in allele frequencies between haploids and diploids, η_{HW} indicates the departure from

the Hardy-Weinberg equilibrium in diploids, and ρ_H indicates the frequency of haploids in the

population. Within these equations, the frequencies of mutant alleles in haploids and diploids are

$p_H = x_M/(x_R + x_M)$ and $p_D = \left(\frac{x_{RM}}{2} + x_{MM}\right)/(x_{RR} + x_{RM} + x_{MM})$. As similar variables are

used in the model with local population regulation, we use superscripts to indicate the form of

population regulation ("*Model*" is *G* for global or *L* for local regulation).

The class reproductive values of haploids and diploids are defined as follows. In linear

models, "reproductive value" is a measure of the expected fraction of the population in the

long-term future that descends from an individual of a particular type (e.g., age or stage class).

Class reproductive values, as defined by Taylor (1990) and Rousset (2004, p.153), scale these

individual reproductive values up to the whole population of each class (i.e., the product of the

individual reproductive values times the class size). In the models considered here, the dynamics

are non-linear because of competition for resources (N_{tot}). Nevertheless, we can approximate

reproductive values by assuming that the population is near equilibrium with only resident alleles

and by holding the strength of competition constant (Appendix A, Supplementary *Mathematica*

file; Supplementary parts). Doing so, we find that the class reproductive values of haploids and

diploids, expressed as proportions that sum to one, are:

$$c_H^G = \frac{(1-a_H)\frac{f_R}{2}w_R(\hat{\rho}_H^G)^2}{(1-a_H)\frac{f_R}{2}w_R(\hat{\rho}_H^G)^2 + (1-a_D)w_{RR}(\hat{\rho}_D^G)^2}, \tag{2a}$$

$$c_D^G = \frac{(1 - a_D)w_{RR}(\hat{\rho}_D^G)^2}{(1 - a_H)\frac{f_R}{2}w_R(\hat{\rho}_H^G)^2 + (1 - a_D)w_{RR}(\hat{\rho}_D^G)^2}. \tag{2b}$$

where $\hat{\rho}_H^G = 1 - \hat{\rho}_D^G$ is the equilibrium frequency of haploids in the global model (Eq. A.4). As

a special case of interest, when populations are purely sexual ($a_H = a_D = 0$), we can plug the

equilibrium for $\hat{\rho}_H^G$ from Eq. (A.4) into (2) and show that $c_H^G = c_D^G = 1/2$. As discussed in

section 3.4 below, increasing the degree of asexual reproduction of haploids tends to increase

their reproductive value, but there are exceptions when diploids are more fertile and

predominantly asexual. Similarly, increasing the extent of asexual reproduction among diploids

tends to increase the reproductive value of diploids, except when haploids are more fertile and

predominantly asexual. These exceptions occur because passage through the more productive

ploidy level (via sexual reproduction) then becomes essential for individuals of the less

productive ploidy to contribute substantially to future generations.

As discussed in Appendix A (see also Bessho and Otto 2017), equations (2) provide the

only weights that allow ecological and evolutionary time scales to be separated when calculating

the average allele frequency in equation (1a), which is why we take that to be the evolutionarily

relevant average. Although one might initially think that diploids should count twice as much

because they contain two allele copies and that the evolutionarily relevant average allele

frequency would depend on the population sizes of haploids and diploids, having a strict

alternation of generations in a fully sexual population ($a_H = a_D = 0$) ensures that haploids and

diploids contribute equally to long-term future generations, so that their reproductive values are

equal and the evolutionarily relevant average allele frequency is $p_{ave} = (1/2)p_H + (1/2)p_D$

(Bessho and Otto 2017).

As with our previous model, we can track the slow evolutionary dynamics for the

expected change in average allele frequency p_{ave} under weak selection, once the fast ecological

dynamics have stabilized, as which point we can show that there are similar allele frequencies in

haploids and diploids ($\delta_p \approx 0$), diploids are approximately at Hardy-Weinberg equilibrium

($\eta_{HW} \approx 0$), and the ratio of haploids is similar among mutant and resident genotypes ($\rho_H \approx \hat{\rho}_H^G$)

(Appendix A, Supplementary *Mathematica* file; Diffusion approximation for global (local)

regulation model). Furthermore, to leading order, the second moment of change in allele

frequency is equal to the neutral case and can be derived in the diffusion limit ($N_{tot} \to \infty$).

Given a single variable, p_{ave}, changing slowly over evolutionary time, we can then

use standard diffusion methods to calculate the fixation probability of a mutation in a

haploid-diploid population, $u(p_0^{Model})$, where "*Model*" is *G* for global and *L* for local (considered

in the next section). The diffusion is a function of the first and second moments of change in the

mutant allele frequency, $m^{Model}(p_{ave})$ and $v^{Model}(p_{ave})$, both measured in time units of N_{tot}

generations:

$$u\left(p_0^{Model}\right) = \frac{\int_0^{p_0^{Model}} \exp[-2Q^{Model}(p')]\, dp'}{\int_0^1 \exp[-2Q^{Model}(p')]\, dp'}, \tag{3a}$$

$$m^{Model}(p_{ave}) = \frac{N_{tot}\, p_{ave}(1 - p_{ave})}{2}\left[2s_{ave}^{Model} + 2c_D^{Model} p_{ave}(1 - 2h)s_{MM}^w\right], \tag{3b}$$

$$v^{Model}(p_{ave}) = \frac{p_{ave}(1 - p_{ave})\left[\left(c_D^{Model}\right)^2 \hat{\rho}_H^{Model} + \left(c_H^{Model}\right)^2 \left(2\hat{\rho}_D^{Model}\right)\right]}{\hat{\rho}_H^{Model}\left(2\hat{\rho}_D^{Model}\right)}. \tag{3c}$$

where p_0^{Model} is the initial allele frequency of mutants (we focus on the case with a single initial

mutant   allele, $p_0^{Model} = 1/\left[N_{tot}\left(\hat{\rho}_H^{Model} + 2\hat{\rho}_D^{Model}\right)\right]$)   and

$Q^{Model}(p) = \int (m^{Model}(p)/v^{Model}(p))\, dp$. We note that Eq (3a) is obtained from the backward

equation for the fixation probability (Bessho and Otto 2017) and Eqs. (3b) and (3c) are derived in

Appendix A. For the global regulation model, the average selection acting upon rare mutant

alleles across haploid and diploid stages, s_{ave}^G, can be calculated from the first moment equation

(see Supplementary *Mathematica* file; Diffusion approximation for the global (local) regulation

model) and equals:

$$s_{ave}^{Model} = c_H^{Model} s_M^w + c_D^{Model} s_{RM}^w + \phi^{Model} c_D^{Model} \frac{s_M^f}{2}, \tag{4a}$$

$$\phi^{Model} = \frac{(1 - a_H)\widetilde{w}_R \hat{\rho}_H^{Model}}{(1 - a_H)\widetilde{w}_R \hat{\rho}_H^{Model} + a_D w_{RR} \hat{\rho}_D^{Model}}. \tag{4b}$$

where $\widetilde{w}_R = f_R w_R/2$ is the fertility of haploids considering the cost of sex. As we will see later,

this equation is valid for local regulation model. The term ϕ^{Model} indicates the fraction of the

diploids in the next generation that are sexually produced (from the union of haploid-produced

gametes) rather than asexually produced by diploid parents. With obligately sexual

haploid-diploids ($a_H = a_D = 0$, where $c_H^{Model} = c_D^{Model} = 1/2$ and $\phi^{Model} = 1$), these results

coincide with those of Bessho and Otto (2017).

3.2. Fixation probability in the local regulation model

We next derive the fixation probability in a haploid-diploid population when density

dependence regulates haploid and diploid populations separately (Figure 1b), by again applying a

transformation of variables and separation of time scales. For the local regulation model, the

appropriate weights for the average allele frequency are similar to the global regulation model,

where now the class reproductive values, expressed as proportions, are:

$$c_H^L = \frac{1 + \dfrac{a_H w_R \hat{\rho}_H^L}{(1 - a_D)w_{RR}\hat{\rho}_D^L}}{2 + \dfrac{a_H w_R \hat{\rho}_H^L}{(1 - a_D)w_{RR}\hat{\rho}_D^L} + \dfrac{2}{f_R}\dfrac{a_D w_{RR}\hat{\rho}_D^L}{(1 - a_H)w_R\hat{\rho}_H^L}}, \tag{5a}$$

$$c_D^L = \frac{1 + \dfrac{2}{f_R}\dfrac{a_D w_{RR}\hat{\rho}_D^L}{(1 - a_H)w_R\hat{\rho}_H^L}}{2 + \dfrac{a_H w_R \hat{\rho}_H^L}{(1 - a_D)w_{RR}\hat{\rho}_D^L} + \dfrac{2}{f_R}\dfrac{a_D w_{RR}\hat{\rho}_D^L}{(1 - a_H)w_R\hat{\rho}_H^L}}. \tag{5b}$$

After applying a separation of time scales and conducting a diffusion approximation, we

conclude that the solution for the fixation probability in a haploid-diploid population, Eqs. (3),

remains valid for the local regulation model (Supplementary *Mathematica* file; Diffusion

approximation for the local regulation model), with the average selection coefficient now being

given by Eqs. (5).

3.3. Effective genetic parameters

Using the first and second moments of change in allele frequency, we derive effective

genetic parameters to compare our results to the dynamics found in the classical model for fully

haploid or fully diploid organisms (Bessho and Otto 2017). More specifically, we define the

effective selection coefficient (s_e), dominance coefficient (h_e), and effective population size (N_e)

that would result in the same expected change in allele frequency and variance as in the classical

diploid model of selection.

For selection, the diploid model is: $\Delta p_{ave} = s_e p_{ave}(1 - p_{ave})[h_e + (1 - 2h_e)p_{ave}]$

(Crow and Kimura 1970; Bessho and Otto 2017). Because this equation depends on the allele

frequency in the same way as Eq. (3b), we can find the effective and dominance selection

coefficient   from $\Delta p_{ave}/[p_{ave}(1 - p_{ave})] = s_e h_e$ when $p_{ave} = 0$ and

$\Delta p_{ave}/[p_{ave}(1 - p_{ave})] = s_e(1 - h_e)$ when $p_{ave} = 1$, yielding:

$$s_e^{Model} = 2s_{ave}^{Model} + 2\hat{c}_D^{Model} \frac{(1 - 2h)s_{MM}^w}{2}. \tag{6a}$$

$$h_e^{Model} = \frac{2s_{ave}^{Model}}{4s_{ave}^{Model} + 2\hat{c}_D^{Model}(1 - 2h)s_{MM}^w}. \tag{6b}$$

When the mutation is additive ($h = 1/2$), these effective parameters are $s_e^{Model} = 2s_{ave}^{Model}$ and

$h_e^{Model} = 1/2$.

We next derive the variance effective population size (Crow and Kimura 1970) by

equating the one generation change in variance (Eq. 3c divided by the time scale, N_{tot}) to the

variance in allele frequency expected in the classical Wright-Fisher model, $p_{ave}(1 - p_{ave})/2N_e$

with N_e diploid individuals, obtaining:

$$N_e^{Model} = \frac{p_{ave}(1 - p_{ave})}{2(v^{Model}/N_{tot})} = \frac{N_{tot}\,\hat{\rho}_H^{Model}\hat{\rho}_D^{Model}}{\left(c_D^{Model}\right)^2\hat{\rho}_H^{Model} + 2\left(c_H^{Model}\right)^2\hat{\rho}_D^{Model}}. \qquad (7)$$

233   Plugging these effective parameters into the formula from the fixation probability in

234   the classical diploid Wright-Fisher model (Kimura 1957; 1962; Crow and Kimura 1970, p. 427),

235   the fixation probability in a haploid-diploid population given by Eq. 3a can be expressed as:

$$u(p_0^{Model}) = \frac{\int_0^{p_0^{Model}} \exp[-2N_e^{Model}s_e^{Model}\{(2h_e^{Model}-1)p'(1-p') + p'\}]\,dp'}{\int_0^1 \exp[-2N_e^{Model}s_e^{Model}\{(2h_e^{Model}-1)p'(1-p') + p'\}]\,dp'}. \qquad (8)$$

236   Assuming an initially rare and additive mutation ($h = 1/2$) with weak positive selection in a

237   large population ($s_e^{Model}N_e^{Model}p_0^{Model} \approx 0$ and $s_e^{Model}N_e^{Model} \gg 1$), we obtain the classic

238   approximation, $u\left(p_0^{Model}\right) \approx 2s_e^{Model}N_e^{Model}p_0^{Model}$, which upon substituting from Eq. 7 yields:

$$u(p_0^{Model}) \approx \frac{2\hat{\rho}_H^{Model}\hat{\rho}_D^{Model}}{(\hat{\rho}_H^{Model} + 2\hat{\rho}_D^{Model})\left[\left(c_D^{Model}\right)^2\hat{\rho}_H^{Model} + 2\left(c_H^{Model}\right)^2\hat{\rho}_D^{Model}\right]}\,2s_{ave}^{Model}. \qquad (9a)$$

239   For example, because haploids and diploids have the same reproductive values in the obligately

240   sexual case ($\hat{c}_H^{Model} = \hat{c}_D^{Model} = 1/2$), we obtain:

$$u\left(p_0^{Model}\right) \approx \frac{8\hat{\rho}_H^{Model}\hat{\rho}_D^{Model}}{\left(\hat{\rho}_H^{Model} + 2\hat{\rho}_D^{Model}\right)^2}\,2s_{ave}^{Model} \qquad (9b)$$

(Eq. 13a in Bessho and Otto 2017), or simply $u\left(p_0^{Model}\right) \approx 2s_{ave}^{Model}$ if haploid and diploid

population sizes are equal in terms of number of chromosomes ($\hat{\rho}_H^{Model} = 2/3$).

In the next three sections, we explore the implications of these results for the evolution

of haploid-diploid populations.

3.4. Effective selection in a haploid-diploid population

The strength of selection averaged across haploids and diploids, s_{ave}^{Model}, plays a key

role in the evolution of haploid-diploid populations. When a mutation is rare, both the rate of

change in allele frequency (Eq. 3b) and the approximate fixation probability (Eq. 10a) are

proportional to s_{ave}^{Model}. We thus begin by exploring how s_{ave}^{Model} varies as we alter the amount

of asexual reproduction in haploid and diploid phases. We focus on the case where the mutation

does not affect fertilization success ($s_M^f = 0$), so that the average selection becomes:

$$s_{ave}^{Model} = c_H^{Model} s_M^w + c_D^{Model} s_{RM}^w, \qquad (10)$$

in both global and local regulation models (Eqs. (4) and (6)).

The relative evolutionary importance of selection in the haploid and diploid phases is

thus determined by the class reproductive values, c_H^{Model} and c_D^{Model} (where $c_H^{Model} +$

$c_D^{Model}=1$). Figures 2 (global regulation) and 3 (local regulation) illustrate the proportional

reproductive value of haploids, c_H^{Model}, as a function of the degree of asexual reproduction in

haploids (x-axis) and diploids (ranging from 0.05 in red to 0.95 in blue). With global regulation,

the frequency of haploidy within the population, $\hat{\rho}_H^G$ (given by Eq. A.4), varies with the

parameters (see inset graphs in Figure 2), rising with the frequency of haploid asexuality (x-axis

in inset) but declining with more asexuality in diploids (from red to blue).

By contrast, with local regulation, the frequency of haploidy is held fixed by the strict

density dependent competition that we have assumed ($\hat{\rho}_H^L = 0.8$ in Figure 3(a)(b) and 0.3 in

3(c)(d)). In the left panels, haploids have a higher fertility ($w_R/w_{RR} = 5$), leading to a higher

haploid reproductive value, c_H^{Model}, especially with local regulation when haploids are also more

common ($\hat{\rho}_H^L = 0.8$ in Figure 3a). In the right panels, diploids have a higher fertility ($w_{RR}/w_R =$

5), leading to a lower haploid reproductive value, especially when haploids are rare ($\hat{\rho}_H^L = 0.3$ in

Figure 3d).

When haploids are primarily sexual ($a_H \approx 0$), increasing asexuality of the haploid

stage typically causes the reproductive value of haploids to rise, unless diploids are fitter and

more frequent (Figure 3d and blue curves in Figure 2b). At the other extreme, the reproductive

value of haploids typically plummets to zero as haploid reproduction becomes primarily asexual

($a_H \approx 1$) while diploids remain sexual, particularly with local regulation (Figure 3), because

haploids then act as a genetic "sink" contributing little to the diploid sub-population. This

downward trend when haploids are predominantly asexual is also seen with global regulation if

diploids are more fit (Figure 2b), except when the diploid population does not sustain itself and

goes extinct, which occurs when $a_D < 0.2$ and $a_H = 1$. The net result can thus be

non-monotonic (purple curves with $0.2 < a_D < 0.4$ in Figure 2b and Figure 3(a)(b)(c)).

3.5. Effective population size in a haploid-diploid population

We next consider the effective size of haploid-diploid populations with varying degrees

of asexuality. Figure 4 plots the effective population size (Eq. 8) relative to the total population

size, N_e^{Model}/N_{tot}, as a function of the frequency of haploids, $\hat{\rho}_H^{Model}$ (x-axis), and the class

reproductive values (with c_H^{Model} ranging from 0.05 in blue to 0.95 in red). As noted by Bessho

and Otto (2017), the effective population size is highest – and drift weakest – at intermediate

frequencies of haploids and diploids, which ensures the least sampling error as organisms

alternate generations.

When haploids and diploids have equal reproductive values, as in the fully sexual case

($c_H^{Model} = c_D^{Model} = 1/2$), the effective population size is maximized at $\hat{\rho}_H^{Model} \approx 0.586$. With

asexual reproduction, the peak shifts towards whichever ploidy level has the higher reproductive

value. For example, if haploids have a high reproductive value (red) then the effective population

size is maximized at a higher frequency of haploids, reducing the amount of genetic drift in that

phase. Although not illustrated, the peak shifts to $N_e^{Model} = \hat{\rho}_D^{Model} N_{tot}$ when future

populations descend only from diploids ($c_H^{Model} = 0$) and to $N_e^{Model} = (\hat{\rho}_H^{Model}/2)N_{tot}$ when

future populations descend only from haploids ($c_H^{Model} = 1$), effectively becoming diplontic or

haplontic, respectively (with the $1/2$ arising because haploids have half the number of

chromosomes).

Of course, the reproductive values, as well as the frequency of haploids with global

population regulation ($\hat{\rho}_H^G$), depend in turn on the fertility parameters and the extent of asexuality,

as explored in the previous section. Figures 5 (global regulation) and 6 (local regulation)

illustrate the effective population size as a function of the frequency of haploid asexuality, a_H

(x-axis), and the frequency of diploid asexuality (a_D rising from red to blue), using the

parameters in Figures 2 and 3, respectively. The trends are often non-monotonic, with N_e^{Model}/

N_{tot} values varying around $1/2$ when the parameter values are intermediate. The effective

population size is often higher when diploids rarely reproduce asexually (red) rather than when

they frequently do (blue), although there are exceptions.

To check whether our analytical result is correct, we simulated a haploid-diploid

population for a neutral allele (Figure 7 for the global regulation and Figure 8 for the local

regulation). In the classical WF model, the expected heterozygosity within a simulation is

$E[H(t)] = H(0)\left(1 - \frac{1}{2N}\right)^t \approx H(0) \exp\left[-\frac{t}{2N}\right]$. Our simulations reveal that the effective

population size predicts the average dynamics of $E[H_e(t)]$ (yellow line) where $H_e(t) =$

$2p_{ave}(1 - p_{ave})$. A slight discrepancy is observed when diploids reproduce primarily asexually

(Fig. 7d and 8h), because the assumption that departures from Hardy-Weinberg remain small

becomes accurate with drift; nevertheless, our formula for the effective population size continues

to perform reasonably well even in this case (especially for the first few hundred generations).

Here, we have tracked the "effective" heterozygosity calculated as $H_e(t) =$

$2p_{ave}(1 - p_{ave})$ where p_{ave} is the average allele frequency based on the (typically unknown)

class reproductive values. We also find, however, that the effective population size equally well

predicts loss of heterozygosity using the naive allele frequency that one might observe by

randomly sampling alleles from a population:

$p_{naive} = (2x_{MM} + x_{RM} + x_M)/(2x_{MM} + 2x_{RM} + 2x_{RR} + x_M + x_R)$ (see Supplementary

Mathematica file; Figure 7 and 8). Even the loss of heterozygotes in the diploid population,

$x_{RM}/(x_{MM} + x_{RM} + x_{RR})$, is well described by the effective population size derived here.

3.6. Fixation probability in a haploid-diploid population

We next compare the above results with numerical simulations estimating the fixation

probability of a newly arisen mutation in a haploid-diploid population. When simulating the

global regulation model, we assumed that the population has reached the demographic

equilibrium, $\hat{\rho}_H^G N_{tot}$ haploids and $\hat{\rho}_D^G N_{tot}$ diploids (see Appendix A). We then chose one

resident allele R at random and replaced it with a mutant allele M. After mutation, offspring were

sampled from the parental generation according to a multinomial distribution with expected

frequencies given by $x_{(GT)}$, repeating until the mutant allele fixed or was lost from the

population. We estimated the fixation probability as the fraction of 100,000 replicate simulations

leading to fixation.

We here consider the additive case ($h = 1/2$ and $h_e = 1/2$), where the fixation

probability (Eq. 8) simplifies to:

$$u\left(p_0^{Model}\right) = \frac{\exp[-2p_0^{Model} N_e^{Model} s_e^{Model}] - 1}{\exp[-2N_e^{Model} s_e^{Model}] - 1} \tag{11}$$

and where $s_e^{Model} = 2\, s_{ave}^{Model}$ (Eq. 6). Figure 9 plots the fixation probability as a function of the

average selection pressure, s_{ave}^{Model}, when the reproductive values and chromosome numbers in

haploids and diploids are equal ($c_H^{Model} = c_D^{Model}$ and $\hat{\rho}_H^{Model} = 2/3$) and $s_{ave}^{Model} = [(1/$

$2)s_M^w + (1/2)s_{RM}^w]$. The diffusion Eq. (11) provides an excellent fit, as does the approximation

Eq. (9b) for selection coefficients that are positive and not too weak. In this case, the results are

the same with global and local population regulation (Fig. 9a and 9b, respectively) and are

insensitive to how much selection occurs in the haploid or diploid phases (s_M^{Model} and s_{RM}^{Model},

respectively), as long as s_{ave}^{Model} is held constant (see additional simulations in Supplementary

Mathematica file; Figure 9). As expected, the extent of selection in the haploid versus diploid

phase matters more when the mutation is not additive ($h \neq 1/2$ and $h_e \neq 1/2$) (supplementary

Mathematica file; non additive mutation). Additional simulations are presented in the

supplementary *Mathematica* file (Supplementary: Fixation probability for global (or local)

regulation model (simulation)).

These confirm that the analytical results remain valid across a broad parameter space

(combinations of higher haploid fertility vs diploid fertility, higher haploid asexuality vs diploid

asexuality, and higher frequency of haploids vs diploids).

Next, we illustrate the approximate fixation probability, Eq. (9a), as a function of the

degree of asexuality (a_H and a_D) when the population size is globally (Fig. 10) or locally (Fig.

11) regulated, assuming only selection in haploids or only in diploids. For example, with additive

mutations, the fixation probability can be approximated as $u \approx 4c_H^{Model} N_e^{Model} p_0^{Model} s_M^w$ when

selection occurs only in the haploid phase or $u \approx 4c_D^{Model} N_e^{Model} p_0^{Model} s_{RM}^w$ with selection only

in the diploid phase, indicating that the fate of mutations depends as much on the strength of

selection as on the reproductive value of the ploidy phase in which selection acts (as illustrated in

Fig. 2 and 3). Figures 10 (global) and 11 (local) illustrate how the fixation probability depends on

the various parameters in the model, particularly the amount of asexual reproduction in haploids

(x-axis) and diploids (a_D rising from red to blue). The trends can be understood by the combined

effects of the parameters on the reproductive value and the effective population size (e.g., Fig.

11(a) is proportional to the product of Fig. 3(a) and Fig. 6(c)).

4. Discussion

Across the phylogenetic tree of life, organisms have diverse and complex reproductive

strategies (Bell 1982). Classical population genetic theory has, however, focused most on fully

haploid or diploid life cycles with obligate sexuality. In this article we develop a stochastic

model for the population genetics of haploid-diploid organisms considering demography,

asexuality, and habitat differentiation between haploid and diploid stages. Using a separation of

time scales, we derive a diffusion approximation for the change in allele frequency, allowing us

to estimate the fixation probability of new mutations, the effective strengths of selection and

dominance, as well as the effective population size of haploid-diploid populations.

5.1. Natural selection in a haploid-diploid population

Our results indicate that the strength of natural selection and the extent of genetic drift

depend strongly on the reproductive value of haploid versus diploid phases. In the simplest case,

when the effect of a mutation is weak, additive, positive, and absent in the gamete stage ($s_M^f = 0$),

the fixation probability is proportional to the effective strength of selection (Eq. 10), $s_e^{Model} =$

$2s_{ave}^{Model}$, which in turn is proportional to the amount of selection in and the reproductive value of

haploids and diploids (Eqs. 2, and 5).

These analytical results reveal some evolutionary principles for populations that

undergo an alternation of generations. One consequence is that the balance of opposing selection

pressures in haploids and diploids (Eqs. 4 and 6) depends not only on the selection coefficients,

but also on the relative reproductive values of haploids (c_H^{Model}) versus diploids (c_D^{Model}). Thus,

the very direction of evolution depends on the extent of asexuality in the two phases and the

relative survival and fertility of haploids versus diploids when there is "ploidally antagonistic

selection" (Immler et al. 2012).

The efficacy of selection to fine tune traits in haploids and diploids also depends on the

class reproductive values. For example, when the population is regulated by local density

dependence (i.e., the haploid and diploid phases are spatially or temporally distinct), higher

reproductive success in haploids increases the efficiency of haploid selection (compare Figure 3a

to 3b). However, when there is extremely rare sexuality in haploids (a_H near one), diploid

selection tends to be more effective because of increasing competition between offspring from

haploids. By contrast, the trends differ with global density dependence (e.g., species that are

more isomorphic with small ecological differences between stages). For example, the

reproductive value of haploids remains high even when they reproduce primarily asexually in the

global regulation model (see Figure 2 when a_H approaches one), because haploids then make up

a larger proportion of the total population size (see inset figures). Thus, whether selection is

effective in the haploid phase when that phase mainly reproduces asexually is quite sensitive to

the nature of competition.

Our work can also be useful in the design of field studies and the interpretation of data

for species that alternate generations. To understand the efficiency of selection on haploid and

diploid phases, we not only need data about the fraction of haploids and diploids and their

fertility and mortality (e.g., Thornber and Gains 2004; Vieira et al., 2018a; Vieira et al. 2018b),

but we also need to know about the extent of asexuality in each phase and whether they compete

for common or different resources.

5.2. Genetic drift and effective size

The impact of random genetic drift on the genetic diversity of haploid-diploid

population depends on the effective population size (Eq. 7). As we had found previously in a

haploid-diploid model with obligate sexuality (Bessho and Otto 2017, pp. 431), the effective

population size with asexuality is generally smaller than the total number of individuals and

again depends strongly on the reproductive value of each phase (Figures 4-6). With obligate

sexuality, the reproductive values of haploids and diploids are equal, drift is minimized for a new

allele (the effective population size times p_0 is maximized) when haploids comprise 2/3 of the

population, making the number of chromosomes equal between haploids and diploids. Asexual

reproduction, however, causes the reproductive value of haploids and diploids to differ (Eqs. 2

and 5). Consequently, drift is lessened if the phase with the higher reproductive value is more

common (see shifts in peaks in Figure 4).

5.3. Evolution in haploid-diploid populations with asexuality

We find that evolution in haploid-diploid populations is substantially different than

either fully haploid or fully diploid populations and also substantially different than fully sexual

haploid-diploid populations. The model considered in this paper allows selection in both haploid

and diploids phases but does not require a strict alternation of generations, allowing individuals

to produce offspring of either the same ploidy type (asexually) or the opposite ploidy type

(sexually).

Here we consider the implications for evolution, focusing on the simplest case of

additive selection with weak selection, where the fixation probability is given by Eq. (9a). This

formula indicates that evolution in a haploid-diploid population is dominated by (i) the average

selection across haploid and diploid stages, (ii) the ploidy ratio in the population, and (iii) the

class reproductive values of haploids and diploids. Comparing this probability with the classical

result 2s reveals that evolution in a haploid-diploid population differs substantially in three ways

from evolution in a fully haploid (or diploid) population.

First, and most obvious, evolutionary changes are shaped by selection in both haploid

and diploid stages. As we see in subsection 5.1, because the reproductive values of haploids and

diploids determine how selection in each phase contributes to the effective selection coefficient

(6a), selection is not a simple average across ploidy stages but depends on the details of how the

species reproduces. Second, the ploidy ratio of haploids versus diploids strongly affects evolution.

In particular, because genetic drift is strongest when either haploids or diploids are rare (Figure

4), the effective size dramatically decreases when the ploidy ratio is strongly skewed (i.e., most

individuals are haploids or diploids; Figure 4), reducing the efficacy of selection. Third, the

reproductive values of haploid and diploid stages depend in a complex way on selection in the

two phases and their mode of reproduction (Figures 2-4), affecting the course of evolution

through both selection and genetic drift.

Evolutionary dynamics in completely sexual haploid-diploid populations is

substantially simpler. In particular, with a strict alternation of generations (no asexuality), the

class reproductive values of haploids and diploids are equal (Bessho and Otto 2017). As a

consequence, selection across the life cycle, is half the strength of selection in haploids and half

in diploids, regardless of the frequency of the two types within the population.

With both asexual and sexual reproduction, however, the average selection, the ploidy

ratio, and the reproductive values of haploids and diploids all depend on the fertility parameters

and the degree of asexuality. The resulting interplay alters the fixation probability of new

mutations, as illustrated in Figures 10 and 11. While the interplay is complex and often

non-monotonic, some patterns emerge.

An important take-home result is that the form of competition acting to regulate

population size strongly affects the reproductive values of haploids and diploids (Eqs. (2) and

(5)) and thus dramatically alters the chance that a new beneficial mutation fixes (equation 8;

Figures 10 and 11). We contrast two forms of population regulation: global, where haploids and

diploids are competitively equivalent, and local, where only individuals of a given ploidy level

compete with one another.

With global regulation, the fixation probability for an allele that benefits the haploid

phase is generally higher when haploids are more asexual and diploids more sexual (Figure 10),

increasing the fraction of haploids in the population (Figure 2), but there are exceptions when

diploids have higher fertility and reproduce predominantly asexually (bluer curves in Figure 10b).

Similarly, the fixation probability for an allele that benefits the diploid phase is generally higher

when diploids are more asexual and haploids more sexual (Figure 10), increasing the fraction of

diploids experiencing selection, with exceptions when haploids have higher fertility and are

predominantly asexual (right hand side of Figure 10c). In summary, selection in a particular

ploidy phase is typically more effective when that phase is more common and tends to breed true

(asexual reproduction), with exceptions when the other ploidy phase is a greater evolutionary

"source" of future generations, with a higher reproductive value (higher fertility and more

asexual). These exceptions occur because passage through the other ploidy phase then becomes

essential for an allele to avoid loss while rare, even when selection does not act in the other

phase.

The patterns with local regulation of population size differ substantially. Because local

population regulation by ploidy level determines the equilibrium fraction of haploids and

diploids, the fixation probability tends to be less dependent on the exact degree of asexuality, as

long as there is some sexual and some asexual reproduction (flatter curves with more overlapping

red/blue in Figure 11). In this case, the fixation probability of an allele favoured in the haploid

phase is higher if haploids are more fertile (Figure 11 panels a and c versus b and d) and if

haploids are more common (panels a and b versus c and d), with the reverse holding for alleles

favoured in the diploid phase. When populations are predominantly sexual or predominantly

asexual, however, there can be rapid changes in the fixation probability (Figure 11), resulting

from rapid changes in the reproductive value of each phase (Figure 3).

5.4. "Ploidally-structured" populations

The key role that reproductive values play in this work is analogous to the role that

patch dynamics play in two-patch models of evolution. In a spatially structured population,

subdivided local populations are genetically connected by migration. A haploid-diploid system

can be seen as being ploidally structured, where gene flow describes the movement of alleles

through sexual reproduction, with meiosis causing flow to haploidy and syngamy flow to

diploidy. We note that our research reveals that all qualitative results are equally accurate for

evolution in a two-patch system (see Supplementary *Mathematica* File; Comparison with

two-patch system). For example, fixation probability strongly depends on class reproductive

values of each patch.

This analogy suggests an interesting idea: complex reproductive systems can be

considered and analyzed using the tools of metapopulation theory. For example, many

eukaryotes including terrestrial plants, insects, and fishes, often exhibit ploidy variation,

including polyploid members (Otto and Whitton 2000; Comai 2005). In such species, individuals

characterized by different numbers of chromosomes coexist, with complex reproductive

relationships causing gene flow between them (Ramsey and Schemske 1998). Similarly, social

insects often exhibit complex sex determination systems linked with ploidy levels

(haplodiploidy).

Our research suggests that these ploidally-structured populations can be fruitfully

treated as metapopulations. Selection and drift in populations with diploids, triploids, and

tetraploids can, for example, be considered as a three-patch model. In this system, we conjecture

that the average strength of selection that is evolutionarily relevant would be the mean selection

coefficient in each ploidy class, weighted by its class reproductive value, with additional terms

coming from reproductive interactions (akin to the term of s_M^f in Eqs. 4).

Many evolutionary aspects of haploid-diploid populations remain to be investigated.

One avenue that we are exploring is how model parameters can be estimated from field data. For

example, the analogy between spatially and ploidally structured population suggests that genetic

differences between haploids and diploid can be used to estimate gene flow between them (i.e.,

rates of sex), akin to using Fst to inform estimates of migration (e.g., Slatkin 1987). Another

fruitful avenue for further work is to determine how fluctuations in population size affect the

effective population size of species that alternate generations. In classical population genetics

theory, such fluctuations can be captured by using the harmonic mean population in place of the

total population size (Karlin 1968). It is unclear, however, whether the same is true in

haploid-diploid populations. Can the harmonic total population size simply replace N_{tot} in the

global model of population regulation? Similarly, can the harmonic population sizes of haploids

and diploids replace N_H and N_D with local regulation? The answer is unclear because

population size fluctuations perturb the fast ecological dynamics away from the steady state

(especially $\hat{\rho}_H^{Model}$), and the impact of these perturbations on selection and drift is unknown.

Further research is needed to clarify evolutionary processes in the wide variety of species that

alternate generations.

Acknowledgements

We here thank Alireza Ghaseminejad for help in developing our manuscript. We also thank H.

Ohtsuki and Y. Uchiumi for the helpful comment.

Funding sources

This project was funded by a Grand-in-Aid from a Japan Society for the Promotion of Science

(JSPS) to KB (16J05204) and by a Discovery grant from the Natural Sciences and Engineering

539   Appendix A. Fixation probability in a haploid-diploid Wright-Fisher model using a

540   diffusion approximation

A.1. Equilibrium with global regulation

We derive the fixation probability in a haploid-diploid population using a diffusion

approximation (e.g., Bessho and Otto 2017). We first derive the stable equilibrium in the global

regulation model, allowing for asexual reproduction in each phase. In the Wright-Fisher model,

all individuals reproduce and then the parents die (non-overlapping generations). Let $b_{(GT)}$

represent the number of reproductive cells of each type in the next generation:

$$b_R = (1 - a_D)w_{RR}x_{RR} + (1 - a_D)\frac{w_{RM}x_{RM}}{2} + a_H w_R x_R, \tag{A.1a}$$

$$b_M = (1 - a_D)\frac{w_{RM}x_{RM}}{2} + (1 - a_D)w_{MM}x_{MM} + a_H w_M x_M, \tag{A.1b}$$

$$b_{RR} = (1 - a_H)\frac{f_R}{2}\frac{w_R^2 x_R^2}{w_R x_R + w_M x_M} + a_D w_{RR}x_{RR}, \tag{A.1c}$$

$$b_{RM} = (1 - a_H)\frac{f_R + f_M}{2}\frac{w_R w_M x_R x_M}{w_R x_R + w_M x_M} + a_D w_{RM}x_{RM}, \tag{A.1d}$$

$$b_{MM} = (1 - a_H)\frac{f_M}{2}\frac{w_M^2 x_M^2}{w_R x_R + w_M x_M} + a_D w_{MM}x_{MM}. \tag{A.1e}$$

547   The probability that a reproductive cell of genotype (*GT*) is sampled from the offspring produced

548   by the previous generation of adults is

$$q_{(GT)} = \frac{b_{(GT)}}{b_R + b_M + b_{RR} + b_{RM} + b_{MM}}. \tag{A.2}$$

549   Therefore, the composition of offspring in the next generation is given by the multinomial

550   distribution, sampling N_{tot} individuals in proportion to Eq. (A2). Using Eq. (A1) and (A2), we

551   describe the conditional expectation of change in the number of individuals of genotype (*GT*),

552    $\Delta X_{(GT)}(t) = X_{(GT)}(t + 1) - X_{(GT)}(t)$, as

$$\text{E}\big[\Delta X_{(GT)}(t)|\vec{X}(t) = \vec{x}\big] = N_{tot} q_{(GT)} - x_{(GT)}, \tag{A.3}$$

where $\text{E}\left[\Delta F\left(X_{(GT)}(t)\right)|\vec{X}(t) = \vec{x}\right]$ is the conditional expected value for change in the function

F of the random variable given that $\vec{X}(t) = (X_R(t) \quad X_M(t) \quad X_{RR}(t) \quad X_{RM}(t) \quad X_{MM}(t))^T$

equals $\vec{x} = (x_R \quad x_M \quad x_{RR} \quad x_{RM} \quad x_{MM})^T$.

To simplify this fully stochastic system, we assume that the resident population is large

and treat demographic changes deterministically prior to the appearance of the mutation.

Considering the dynamics of the resident population, we then find the equilibrium of these

dynamical equations by solving $N_{tot} q_R - \hat{x}_R = 0$ and $N_{tot} q_{RR} - \hat{x}_{RR} = 0$ $(\hat{x}_R + \hat{x}_{RR} = N_{tot})$.

Setting $\hat{x}_R = \hat{\rho}_H^G N_{tot}$ and $\hat{x}_{RR} = \hat{\rho}_D^G N_{tot}$, the fraction of haploids $\hat{\rho}_H^G$ (and diploids $\hat{\rho}_D^G = 1 -$

$\hat{\rho}_H^G$) at equilibrium becomes,

$$\hat{\rho}_H^G = \frac{a_H w_R + a_D w_{RR} - 2w_{RR} + \sqrt{4(1 - a_H)(1 - a_D)\tilde{w}_R w_{RR} + (a_H w_R - a_D w_{RR})^2}}{2[a_H w_R + (1 - a_H)\tilde{w}_R - w_{RR}]}, \tag{A.4}$$

where $\tilde{w}_R = f_R w_R/2$ is the fertility of haploids considering the cost of sex (see Supplementary

Mathematica file for the step-by-step derivation; Supplementary parts). We note that, when the

fertility of haploids is much greater than that of diploids ($w_R \gg w_{RR}$), the frequency of haploids

in a population approaches $a_H/\{a_H + [(1 - a_H)f_R/2]\}$, which is less than one because sexual

reproduction of the haploids produces diploids (the $(1 - a_H)f_R/2$ term). Conversely, when the

fertility of diploids is much greater than haploids ($w_R \ll w_{RR}$), the frequency of haploids

approaches $1 - a_D$, the rate at which diploids undergo meiosis.

A.2. First moment of change in allele frequency

To derive the first moment of change in allele frequency, $m^{Model}(p_{ave})$, we apply a

separation of time scales (e.g., Nagylaki 1976; Otto and Day 2007; Bessho and Otto 2017).

Details of the calculation are represented in the Supplementary *Mathematica* file (Diffusion

approximation for the global (local) regulation model). We first transform the expected change in

the number of individuals of each type (five variables that sum to N_{tot}) into the expected change

in a new set of four variables, $\Theta \in \{p_{ave}, \delta_p, \eta_{HW}, \rho_H\}$, described by the functions:

$$\mathrm{E}\left[\Delta\Theta \,|\, \vec{X}(t) = \vec{x}\right] = f_\Theta^{Model}\left(\epsilon, p_{ave}, \vec{\theta}\right), \tag{A.5}$$

where $\vec{\theta} = \left(\delta_p, \eta_{HW}, \rho_H\right)$ and ϵ is proportional to the selection coefficients and assumed small

(the functions f are given explicitly in the Supplementary *Mathematica* file). With local

regulation, ρ_H is assumed fixed at N_H/N_{tot} and dropped from the variable set, Θ.

To constant order (setting the small changes due to selection to zero, $\epsilon \to 0$), the fast

ecological dynamics of the system are described by: $f_\Theta^{Model}\left(0, p_{ave}, \vec{\theta}\right)$. This system of

equations rapidly approaches a steady state found by solving $f_\Theta^{Model}\left(0, p_{ave}, \vec{\theta}\right) = 0$, which

gives $\bar{\delta}_p = \bar{\eta}_{HW} = 0$, and $\bar{\rho}_H = \hat{\rho}_H^G$ (Eq. A.4). To this order, the steady state change in allele

frequency is zero, $f_{p_{ave}}^{Model}\left(0, p_{ave}, \vec{\theta}\right) = 0$. We then describe slower changes, including changes

in allele frequency due to selection, by describing the deviations that occur around this steady

state. Specifically, to order ϵ, the variables are allowed to deviate from the steady state by

$\delta_p = \tilde{\delta}_p\epsilon$, $\eta_{HW} = \tilde{\eta}_{HW}\epsilon$, and $\rho_H = \hat{\rho}_H^G + \tilde{\rho}_H\epsilon$, and the dynamics $f_\Theta^{Model}\left(\epsilon, p_{ave}, \vec{\theta}\right)$ are then

approximated using a Taylor series expansion. Defining the average allele frequency by

combining haploid and diploid populations using an arbitrary weighting, $p_{ave} = \omega p_H +$

$(1 - \omega)p_D$, we show in the Supplementary *Mathematica* file (Class reproductive value in

Supplementary parts) that setting the weights proportional to the class reproductive values (given

by Eq. (2) with global regulation and Eq. (5) with local regulation) is the only choice that

separates evolutionary change in p_{ave} from changes in the other variables to order ϵ. Defining

the average allele frequency in this way (Eq. 1a), taking the Taylor series, and reporting the

results in the initial parameters (e.g., $s_{(GT)}^w$ rather than $\tilde{s}_{(GT)}^w$, where $s_{(GT)}^w = \epsilon \tilde{s}_{(GT)}^w$) the change

in allele frequency becomes:

$$E[\Delta p_{ave} | \vec{X}(t) = \vec{x}] \approx M^{Model}(p_{ave}) \tag{A.6}$$

$$= \frac{p_{ave}(1 - p_{ave})}{2} \left[2s_{ave}^{Model} + 2c_D^{Model} p_{ave}(1 - 2h)s_{MM}^w \right].$$

In the supplementary Mathematica file, we represent an example of the simulation of

4-dimensional dynamics calculating the fraction of times that the mutation fixed in the

population (see section "Example of 4-dimensional dynamics")

A.3. Second moment of change in average allele frequency

We next derive the second moment of change in average allele frequency in a

haploid-diploid population with asexuality. We again assume that the population size is very

large, that selection is very weak, and that the system has approached the steady state in

$(\delta_p, \eta_{HW}, \rho_H)$, ignoring deviations that are of $O(\epsilon)$. Because selection is assumed weak, the

second moment is well approximated by that of the neutral model (to constant order, $\epsilon \to 0$).

Under these assumptions, the fraction of haploids in a population is relatively fixed in

both the global and local regulation models, and we can sample the haploid offspring according

to a binomial distribution, with expectation and variance: $E[X_M | \vec{X}(t) = \vec{x}] = q_M N_H$ and

$Var[X_M | \vec{X}(t) = \vec{x}] = q_M(1 - q_M)N_H$ where $q_M = b_M / (b_R + b_M)$. To simplify the equation, we

set $E[X_{(GT)} | \vec{X}(t) = \vec{x}] = m_{(GT)}$ and $Var[X_{(GT)} | \vec{X}(t) = \vec{x}] = v_{(GT)}$, finding that:

$$E[\Delta X_M | \vec{X}(t) = \vec{x}] = m_M - x_M, \tag{A.7a}$$

$$E[(\Delta X_M)^2 | \vec{X}(t) = \vec{x}] = v_M + m_M^2 - 2m_M x_M + x_M^2. \tag{A.7b}$$

In terms of allele frequencies (rather than numbers), we have the first and second moments for

the haploid offspring population, $E\left[\frac{\Delta X_M}{N_H} | \vec{X}(t) = \vec{x}\right] = \frac{1}{N_H} E[\Delta X_M | \vec{X}(t) = \vec{x}]$ and

$E\left[\left(\frac{\Delta X_M}{N_H}\right)^2 | \vec{X}(t) = \vec{x}\right] = \frac{1}{N_H^2} E[(\Delta X_M)^2 | \vec{X}(t) = \vec{x}].$

Similarly, the diploid offspring are sampled according to a trinomial distribution, with

expectation, variance, and covariance: $m_{(GT)} = q_{(GT)}N_D$, $v_{(GT)} = q_{(GT)}(1 - q_{(GT)})N_D$, and

$\text{Cov}[X_{RM}, X_{MM} | \vec{X}(t) = \vec{x}] = q_{RM}q_{MM}N_D$, where $q_{(GT)} = b_{(GT)}/(b_{RR} + b_{RM} + b_{MM})$. To derive

the moments of the allele frequency in diploids, we define, $y_M = (x_{RM}/2) + x_{MM}$ and

$Y_M = (X_{RM}/2) + X_{MM}$. The moments of random variable Y are then:

$$E[Y_M | \vec{X}(t) = \vec{x}] = \frac{m_{RM}}{2} + m_{MM}, \tag{A.8a}$$

$$E[Y_M^2 | \vec{X}(t) = \vec{x}] = \frac{v_{RM} + m_{RM}^2}{4} + \left(\text{Cov}[X_{RM}, X_{MM} | \vec{X}(t) = \vec{x}] + m_{RM}m_{MM}\right) + (v_{MM} + m_{MM}^2), \tag{A.8b}$$

$$E[\Delta Y_M | \vec{X}(t) = \vec{x}] = \frac{E[\Delta X_{RM} | \vec{X}(t) = \vec{x}]}{2} + E[\Delta X_{MM} | \vec{X}(t) = \vec{x}] - y_M, \tag{A.8c}$$

$$E[(\Delta Y_M)^2 | \vec{X}(t) = \vec{x}] = E[Y_M^2 | \vec{X}(t) = \vec{x}] - 2E[Y_M | \vec{X}(t) = \vec{x}]y_M + y_M^2. \tag{A.8d}$$

620   To consider the change in average allele frequency across the entire population, we

621   define $Z_M = c_H^{Model} \frac{X_M}{N_H} + c_D^{Model} \frac{Y_M}{N_D}$ and consider the expectation of change in this random

622   variable. Plugging in Eqs. (7a), (7b), (8c), and (8d), we have

$$E[(\Delta Z_M)^2 | \vec{X}(t) = \vec{x}] = \frac{p_{ave}(1 - p_{ave})\left[\left(c_D^{Model}\right)^2 \hat{\rho}_H^{Model} + \left(c_H^{Model}\right)^2 \left(2\hat{\rho}_D^{Model}\right)\right]}{\hat{\rho}_H^{Model}\left(2\hat{\rho}_D^{Model}\right)N_{tot}}. \tag{A.9}$$

After transforming time scales using the variable $\tau = t/N_{tot}$ and defining $P(\tau) = Z(N_{tot}\tau)$,

we have the diffusion coefficient $v^{Model}(p_{ave}) = \lim_{N_{tot} \to \infty} E\left[\frac{(P(\tau + \Delta\tau) - P(\tau))^2}{\Delta\tau}\right]$ by taking the

limit $N_{tot} \to \infty$, giving Eq. (3c). Similarly, we derive the drift coefficient using Eq. (A6)

($m^{Model} = M^{Model}N_{tot}$), giving Eq. (3b).

A.4. Class reproductive value of global regulation model

We now calculate the reproductive value of being haploid or diploid, assuming only

resident individuals. Specifically, we are interested in knowing what the long-term contribution

of an individual is to the future population if it is sampled from either haploids or diploids. Here,

we describe the population dynamics in a haploid-diploid population of resident alleles,

$$x'_R = N_{tot} \frac{(1 - a_D)w_{RR}x_{RR} + a_H w_R x_R}{[(1 - a_D)w_{RR}x_{RR} + a_H w_R x_R] + \left[(1 - a_H)\frac{f_R}{2} w_R x_R + a_D w_{RR}x_{RR}\right]}, \qquad \text{(A.10a)}$$

$$x'_{RR} = N_{tot} \frac{(1 - a_H)\frac{f_R}{2} w_R x_R + a_D w_{RR}x_{RR}}{[(1 - a_D)w_{RR}x_{RR} + a_H w_R x_R] + \left[(1 - a_H)\frac{f_R}{2} w_R x_R + a_D w_{RR}x_{RR}\right]}. \qquad \text{(A.10b)}$$

where, x'_R and x'_{RR} indicate the average number of haploids and diploid at the next time step.

The reproductive value of an individual type is generally calculated for linear models

without density dependence. Because the dynamics of x_R and x_{RR} converge to equilibrium, we

modify the recursions so that the total offspring pool size (denominator) is held constant at its

steady state value. This gives us a linear model that allows us to focus on the effects of individual

fitness and movement without the density-dependent feedback:

$$x'_R = \frac{(1 - a_D)w_{RR}x_{RR} + a_H w_R x_R}{[(1 - a_D)w_{RR}\hat{\rho}_D^G + a_H w_R \hat{\rho}_H^G] + \left[(1 - a_H)\frac{f_R}{2} w_R \hat{\rho}_H^G + a_D w_{RR}\hat{\rho}_D^G\right]}, \qquad \text{(A.11a)}$$

$$x'_{RR} = \frac{(1 - a_H)\frac{f_R}{2} w_R x_R + a_D w_{RR}x_{RR}}{[(1 - a_D)w_{RR}\hat{\rho}_D^G + a_H w_R \hat{\rho}_H^G] + \left[(1 - a_H)\frac{f_R}{2} w_R \hat{\rho}_H^G + a_D w_{RR}\hat{\rho}_D^G\right]}, \qquad \text{(A.11b)}$$

643   Eqs. (A.11) can be described by the matrix, $\begin{pmatrix} x'_R \\ x'_{RR} \end{pmatrix} = \mathbf{W} \begin{pmatrix} x_R \\ x_{RR} \end{pmatrix}$. At demographic

644   equilibrium, the leading eigenvalue of matrix $\mathbf{W}$ should be one (see details in Supplementary

645   *Mathematica* file; Supplementary parts in "Diffusion approximation for the global regulation

646   model"). The left eigenvector associated with this eigenvalue is:

$$\left(\frac{(1 - a_H)f_R(1 - a_D - \hat{\rho}_H^G)}{(1 - a_D)\left(f_R\hat{\rho}_H^G - a_H(2 + (-2 + f_R)\hat{\rho}_H^G)\right)} \quad 1 \right)^T \tag{A.12}$$

647   which gives the relative individual reproductive value for a single individual haploid or diploid.

648   The class reproductive values scale this up the contribution of each class,

$$\left(\hat{\rho}_H^G \frac{(1 - a_H)f_R(1 - a_D - \hat{\rho}_H^G)}{(1 - a_D)\left(f_R\hat{\rho}_H^G - a_H(2 + (-2 + f_R)\hat{\rho}_H^G)\right)} \quad \hat{\rho}_D^G \right)^T . \tag{A.13}$$

649   These are proportional to the class reproductive values, Eqs. (2).

651   A.4. Class reproductive value of local regulation model

652   We next calculate the reproductive value of being haploid or diploid with local

653   regulation. We describe the population dynamics in a haploid-diploid population of resident

654   alleles,

$$x_R' = N_H \frac{(1 - a_D)w_{RR}x_{RR} + a_H w_R x_R}{(1 - a_D)w_{RR}x_{RR} + a_H w_R x_R}, \tag{A.14a}$$

$$x_{RR}' = N_D \frac{(1 - a_H)\frac{f_R}{2}w_R x_R + a_D w_{RR}x_{RR}}{(1 - a_H)\frac{f_R}{2}w_R x_R + a_D w_{RR}x_{RR}}. \tag{A.14b}$$

655   For the local regulation model, because both the haploid and diploid population sizes are fixed,

656   we have the trivial dynamics $x_R' = N_H$ and $x_{RR}' = N_D$.

657   To assess the impact of slight changes around this state, we again hold the size of the

658   offspring pool constant at its steady-state value, obtaining the linear set of equations:

$$x_R' = \hat{\rho}_H^L \frac{(1 - a_D)w_{RR}x_{RR} + a_H w_R x_R}{(1 - a_D)w_{RR}\hat{\rho}_D^L + a_H w_R \hat{\rho}_H^L}, \tag{A.15a}$$

$$x_{RR}' = \hat{\rho}_D^L \frac{(1 - a_H)\frac{f_R}{2}w_R x_R + a_D w_{RR}x_{RR}}{(1 - a_H)\frac{f_R}{2}w_R \hat{\rho}_H^L + a_D w_{RR}\hat{\rho}_D^L}. \tag{A.15b}$$

661   Eqs. (A.15) can be described by matrix, $\begin{pmatrix} x'_R \\ x'_{RR} \end{pmatrix} = \mathbf{W} \begin{pmatrix} x_R \\ x_{RR} \end{pmatrix}$. The leading eigenvalue of

662 $\mathbf{W}$ is again one because of the assumption that the population is at its steady-state size (see

663 details in Supplementary *Mathematica* file; Supplementary parts in "Diffusion approximation for

664 the local regulation model"), and the associated left eigenvector becomes,

$$\left(\frac{(1-a_H)f_R w_R (w_{RR} - a_D w_{RR} + a_H w_R \hat{\rho}_H^L - w_{RR}\hat{\rho}_H^L + a_D w_{RR}\hat{\rho}_H^L)}{(1-a_D)w_{RR}(2a_D w_{RR} + f_R w_R \hat{\rho}_H^G - a_H f_R w_R \hat{\rho}_H^L - 2a_D w_{RR}\hat{\rho}_H^L)} \quad 1 \right)^T. \qquad (A.16)$$

This vector gives the relative reproductive value for a single individual that is either haploid or

diploid. The class reproductive values scale this up, with the contribution of each class being

proportional to the class reproductive values, Eqs. (5).

References

Bell, G. 1982. The Masterpiece of Nature: The Genetics and Evolution of Sexuality. University
of California Press. Berkeley, CA.
Bell, G., 1994. The comparative biology of the alternation of generations. Lectures on
mathematics in the life sciences 25, 1–26.
Bessho, K., Otto, S. P. 2017. Fixation probability in a haploid-diploid population. Genetics 205,
421-440.
Comai, L. 2005. The advantages and disadvantages of being polyploid. Nature Reviews Genetics
6, 836-846.
Coelho, S. M., Peters, A. F., Charrier, B., Roze, D., Destombe, C., Valero, M., Cock, J. M., 2007.
Complex life cycles of multicellular eukaryotes: new approaches based on the use of
model organisms. Gene 406, 152–170.
Crow, J. F., Kimura, M. (1970). An introduction to population genetics theory. The Blackburn
Press, New Jersey.

Deithier, M. N. (1981). Heteromorphic algal life histories: The seasonal pattern and response to
herbivory of the brown crust, *Ralfsia californica*. Oecologia 49, 333-339.

Destombe, C., Godin, J., Nocher, M., Richerd, S., Valero, M. (1993). Differences in response
between haploid and diploid isomorphic phases of *Gracilaria verrucosa* (Rhodophyta:
Gigartinales) exposed to artificial environmental conditions. Hydrobiologia 260/261,
131-137.

De Wreede, R. E., Klinger, T. 1988. Reproductive strategies in algae. Plant reproductive ecology:
Patterns and strategies, 267-284.

Dyck, L. J., De Wreede, R. E. (2006). Seasonal and spatial patterns of population density in the
marine macroalga *Mazzaella splendens* (Gigartinales, Rhodophyta). Phycological
Research, 54(1), 21-31.

Goldstein, D. B. 1992. Heterozygote advantage and the evolution of dominant diploid phase.
Genetics 132, 1195–1198.

Haghes, J. S., Otto, S. P. 1999. Ecology and the evolution of biphasic life cycles. The American
Naturalist 154, 306–330.

Hannach, G., Santelicesm, B. 1985. Ecological differences between the isomorphic reproductive
phases of two species of *Iridaea* (Rhodophyta: Gigartinales). Marine Ecology Progress
Series 22, 291–303.

Hawkes, M. W. 1990. Reproductive strategies. Biology of the red algae, 455-476.

Immler, S., G. Arnqvist, and S. P. Otto. 2012. Ploidally antagonistic selection maintains stable
genetic polymorphism. 66:55–65.

Jenkins, C. D., Kirkpatrick, M., 1995. Deleterious mutation and the evolution of genetic life
cycles. Evolution, 512–520.

Karlin, S. 1968. Rates of approach to homozygosity for finite stochastic models with variable
population size. The American Natualist 102, 443-455.

Kimura, M., 1957. Some problems of stochastic processes in genetics. The Annals of
Mathematical Statistics, 882-901.

Kimura, M., 1962. On the probability of fixation of mutant genes in a population. Genetics 47,
713.

Mable, K. B., Otto, P. S., 1998. The evolution of life cycles with haploid and diploid phases.
BioEssays 20, 453–462.

Nagylaki, T., 1976. The evolution of one-and two-locus systems. Genetics 83, 583-600.

Orr, H. A., Otto, S. P. 1994. Does diploidy increase the rate of adaptation? Genetics 136,
1475–1480.

Otto, S., Goldstein, D., 1992. Recombination and the evolution of diploidy. Genetics 131,
745–751.

Otto, S., 1994. The role of deleterious and beneficial mutations in the evolution of ploidy levels. Lectures on Mathematica in the Life Sciences 25, 69–98.

Otto, S. P., Marks, J. C. 1996. Mating system and the evolutionary transition between haploidy and diploidy. Biological Journal of the Linnean Society 57, 197-218.

Otto, S. P., Day, T., 2007. A biologist's guide to mathematical modeling in ecology and evolution. Princeton University Press.

Otto, S. P., Whitton, J. 2000. Polyploid incidence and evolution. Annu. Rev. Genet. 34, 401-437.

Perrot, V., Richerd, S., Valéro, M., 1991. Transition from haploidy to diploidy. Nature 351, 315–317.

Ramsey, J., Schemske, D. W. 1998. Pathways, mechanisms, and rates of polyploid formation in flowering plants. Annual reviews of Ecology and Systematics 29: 467-501.

Raper, J. R., Flexer, A. S. (1970). The road to diploidy with emphasis on a detour. In *Symp. Soc. Gen. Microbiol* (Vol. 20, pp. 401-432).

Rousset, F. (2004). Genetic structure and selection in subdivided populations. Princeton University Press, New Jersey.

Scott, M. F., Rescan, M. 2016. Evolution of haploid-diploid life cycles when haploid and diploid fitnesses are not equal. Evolution 71, 215–226.

Slatkin, M. 1987. Gene flow and the geographic struture of natural populations. Science New Series 236: 787-792.

Slocum C. J. 1980. Differential susceptibility to grazers in two phases of an intertidal alga: Advantages of heteromorphic generations. Journal of Experimental Marine Biology and Ecology 46, 99-110.

Taylor, P. D. 1980. Allele-Frequency Change in a Class-Structured Population. The American Naturalist 135, 95-106.

Thornber, C. S., Gains, S. D. 2004. Population demographics in species with biphasic life cycle. Ecology 85, 1661-1674.

Thornber, C., Stachowicz, J. J., Gains, S. 2006. Tissue type matters: selective herbivory on different life history stages of an isomorphic alga. Ecology 87, 2255-2263.

Vieira, V. M. N. C. S., Engelen, A. H., Huanel, O. R., Guillemin, M. 2018a. Haploid females in the isomorphic biphasic life-cycle of *Gracilaria chilensis* excel in survival. BMC Evolutionary Biology 18:174.

Vieira, V. M. N. C. S., Engelen, A. H., Huanel, O. R., Guillemin, M. 2018b. Differentiation of haploid and diploid fertilities in *Gracilaria chilensis* affect ploidy ratio. BMC Evolutionary Biology 18:183.

Willson, M. F. 1981. On the evolution of complex life cycles in plants: a review and an ecological perspective. Annals of the Missouri Botanical Garden 68, 275-300.

Figure Captions

Fig. 1. An illustration of the haploid-diploid models. (a) In the global regulation model, both

haploids and diploids occupy the same habitat and density dependence holds the total population

size N_{tot} constant. (b) In the local regulation model, each ploidy stage occupies a different habit,

therefore density dependence regulates the population size of haploids (N_H) and diploids (N_D)

separately.

Fig. 2. Class reproductive value of haploids in the global regulation model, c_H^G. Curves show c_H^G

as a function of the degree of haploid asexuality, a_H (x-axis), with the degree of diploid

asexuality ranging in color from $a_D = 0.05$ (red) to 0.95 (blue) in increments of 0.05. Other

parameters are set as: (a) $f_R = 0.5$, $w_R = 5000$, $w_{RR} = 1000$, (b) $f_R = 0.5$, $w_R = 1000$,

$w_{RR} = 5000$. The resulting frequency of haploids, $\hat{\rho}_H^G$ (Eq. A.4), is shown in the inset plots.

Note that if haploids are strictly asexual ($a_H = 1$), diploids persist only when $a_D w_{RR} > w_R$ (i.e.,

when $a_D > 0.2$ in panel b), which accounts for the switch in behavior observed in this panel.

Fig. 3. Proportional reproductive value of haploids in the local regulation model, c_H^L. Parameters

are the same as Fig. 2, except that haploids are held fixed at a frequency of (a)(b) $\hat{\rho}_H^L = 0.8$ or

(c)(d) $\hat{\rho}_H^L = 0.3$. We consider the case when (a)(c) haploid fertility is larger than diploid

$w_R = 5000$, $w_{RR} = 1000$, and when (b)(d) diploid fertility is larger than haploid $w_R = 1000$,

$w_{RR} = 5000$.

Fig. 4. The effective population size of a haploid-diploid population. The relative effective

population size over the total population size (N_e^{Model}/N_{tot}, Eq. 7) is shown as a function of the

frequency of haploids ($\hat{\rho}_H^{Model}$, x-axis), when the haploid reproductive value (c_H^{Model}) varies from

0.05 (blue) to 0.95 (red) in increments of 0.05. This figure applies to both global and local

regulation models.

Fig. 5. The effective population size of a haploid-diploid population in the global regulation

model as a function of the degree of asexuality, a_H and a_D. Parameters are the same as in Fig. 2

and determine the relative class reproductive values (c_H^G) and fraction of haploids ($\hat{\rho}_H^G$) according

to Eqs. (2a) and (A.4).

Fig. 6. The effective population size of a haploid-diploid population in the local regulation model.

Parameters are the same as in Fig. 3 and determine the relative class reproductive values (c_H^L) by

Eqs. (5). The frequencies of haploids are held fixed at a frequency of (a)(b) $\hat{\rho}_H^L = 0.9$, (c)(d)

$\hat{\rho}_H^L = 0.8$, (e)(f) $\hat{\rho}_H^L = 0.6$, or (g)(h) $\hat{\rho}_H^L = 0.3$.

Fig. 7. The estimated heterozygosity over time within a globally regulated haploid-diploid

population, as measured from 1000 replicate simulations of a neutral model (black dots). The

solid yellow curve gives the analytical result for the expected heterozygosity with the effective

population size, $\mathrm{E}[H_e(t)] = H(0)\left(1 - \frac{1}{2N_e^{Model}}\right)^t$. Parameters: $N_{tot} = 90$, $f_R = 0.5$. We

consider the case when (a)(c) haploids are more fertile than diploids ($w_R = 5000$, $w_{RR} = 1000$)

and when (b)(d) diploids are more fertile than haploids ($w_R = 1000$, $w_{RR} = 5000$), considering

asexual reproduction at a high rate among (a)(b) haploids ($a_H = 0.8$, $a_D = 0$) or (c)(d) diploids

($a_H = 0$, $a_D = 0.8$).

Fig. 8. The estimated heterozygosity over time within a locally regulated haploid-diploid

population, as measured from 1000 replicate simulations of a neutral model (black dots). The

solid yellow curve gives the analytical result for the expected heterozygosity. Parameters are the

same as Fig. 7. We consider the case when (a)(c)(e)(g) haploids are more fertile ($w_R = 5000$,

$w_{RR} = 1000$), when (b)(d)(f)(h) diploids are more fertile ($w_R = 1000$, $w_{RR} = 5000$), when

(a)(b)(c)(d) haploids are often asexual ($a_H = 0.8$, $a_D = 0$), when (e)(f)(g)(h) diploids are often

asexual ($a_H = 0$, $a_D = 0.8$), holding haploids fixed at frequency of $\hat{\rho}_H^L = 0.8$ in (a)(b)(e)(f) and

at frequency of $\hat{\rho}_H^L = 0$ in (c)(d)(g)(h).

Fig. 9. Fixation probability given the average strength of selection, s_{ave}^{Model}, for global (a) and

local (b) regulation models. The solid curve gives the analytical result from the diffusion

approximation (Eq. (11)) and the dashed curve gives the approximation with weak, additive, and

positive selection (Eq. (9b)). Black dots indicate the fixation probability estimated from 100000

numerical simulations with 95% CI (Wilson score interval for binomial). Parameters: $N_{tot} = 90$,

$N_H = 60$, $N_D = 30$, $f_R = 0.5$, $w_R = w_{RR} = 1000$, $a_H = a_D = 0.1$, $h = 0.5$, $s_M^f = 0$,

$s_M^w = s_{RM}^w = s_{ave}^{Model}$, such that the fraction of haploids in the resident population is $\hat{\rho}_H^{Model} =$

$2/3$ and class reproductive values are equal $c_H^{Model} = c_D^{Model} = 1/2$. Holding $s_{ave}^{Model} =$

$1/2(s_M^w + s_{RM}^w)$ constant, similar results are obtained for a range of different choices of s_M^w and

s_{RM}^w (see supplementary *Mathematica* file).

Fig. 10. The fixation probability in a haploid-diploid population in the global regulation model.

Curves gives the linear approximation for the fixation probability (Eq. (9a)). Parameters are the

same as in Fig. 2. Selection acts only in the haploid or diploid phase, with selection coefficients

set as $s_M^w = 0.02$ for haploid selection (a)(b) and $s_{RM}^w = 0.02$ for diploid selection (c)(d).

Fig. 11. The fixation probability in a haploid-diploid population in the local regulation model.

Curves gives the linear approximation for the fixation probability (Eq. (9a)). Parameters are the

same as in Fig. 3 and Fig. 10. The frequency of haploids is held fixed at (a)(b)(e)(f) $\hat{\rho}_H^L = 0.8$,

(c)(d)(g)(h) $\hat{\rho}_H^L = 0.3$. Fertility of haploids is higher than diploids in panels (a)(c)(e)(g), and the

opposite condition is considered in panels (b)(d)(f)(h). Selection only occurs in the haploid

(a)(b)(c)(d) or diploid stage (e)(f)(g)(h).

Supporting information

S1. Supplementary *Mathematica* file.

Figure1

(a)

Global regulation

(b)

Local regulation

Total population size: N_{tot}

N_H N_D

Bessho and Otto

Figure 2

(a) Higher haploid fertility

high a_D

c_H^G

low a_D

$\hat{\rho}_H^G$

a_H

(b) Higher diploid fertility

low a_D

$a_D = 0.2$

high a_D

a_H

Bessho and Otto

Figure 3

(a) Higher haploid fertility   (b) Higher diploid fertility

high a_D

low a_D

$\hat{\rho}_H^L = 0.8$ c_H^L

(c)

(d)

$\hat{\rho}_H^L = 0.3$ c_H^L

a_H

a_H

Bessho and Otto

Figure 4

$$\frac{N_e^{Model}}{N_{tot}}$$

small $\hat{c}_H^{Model}$

large $\hat{c}_H^{Model}$

$\hat{\rho}_H^{Model}$

Bessho and Otto

Figure 5

$$\frac{N_e^{Model}}{N_{tot}}$$

(a) Higher haploid fertility **(b)** Higher diploid fertility

small a_D

large a_D

a_H a_H

Bessho and Otto

Figure 6

Figure 7

Bessho and Otto

Figure 8

Figure 9

(a) global regulation

(b) local regulation

Bessho and Otto

Figure 10

Bessho and Otto

Figure 11