High-sensitivity vision restoration via ectopic expression of chimeric rhodopsin in mice

Yusaku Katada[1,2], Kazuho Yoshida[3], Kenta Kobayashi[4], Hideyuki Okano[5], Hideki Kandori[3],

Kazuo Tsubota[2*], Toshihide Kurihara[1,2*]

[1]Laboratory of Photobiology, [2]Department of Ophthalmology, [5]Department of Physiology,

Keio University School of Medicine, Shinanomachi, Shinjuku-ku, Tokyo, Japan

[3]Department of Life Science and Applied Chemistry, Nagoya Institute of Technology,

Nagoya, Aichi, Japan

[4]Section of Viral Vector Development, Center for Genetic Analysis of Behavior, National

Institute for Physiological Sciences, National Institutes of Natural Sciences, Okazaki, Aichi,

Japan

*Corresponding authors:

Toshihide Kurihara, MD, PhD

Laboratory of Photobiology, Department of Ophthalmology, Keio University School of

Medicine; 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan

Tel.: +81-3-5363-3204, Fax: +81-3- 5363-3274

E-mail: kurihara@z8.keio.jp

20   Kazuo Tsubota, MD, PhD

21   Professor and Department Chair, Department of Ophthalmology, Keio University School

22   of Medicine; 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan

23   Tel.: +81-3-5363-3269, Fax: +81-3- 3358-5961

24   E-mail: tsubota@z3.keio.jp

Abstract

Photoreception requires amplification by mammalian rhodopsin through G protein

activation, which requires a visual cycle. To achieve this in retinal gene therapy, we

incorporated human rhodopsin cytoplasmic loops into *Gloeobacter* rhodopsin, thereby

generating *Gloeobacter* and human chimeric rhodopsin (GHCR). In a murine model of

inherited retinal degeneration, we induced retinal GHCR expression by intravitreal

injection of a recombinant adeno-associated virus vector. Retinal explant and visual

thalamus electrophysiological recordings, behavioral tests, and histological analysis

showed that GHCR restored dim-environment vision and prevented the progression of

retinal degeneration. Thus, GHCR may be a potent clinical tool for the treatment of retinal

disorders.

Introduction

  Inherited retinal degeneration (IRD) is a major cause of vision loss. More than 2 million

people worldwide are blind due to IRD[1], and few effective treatments exist. For retinitis

pigmentosa (RP), one of the most common forms of IRD, previous studies have reported

vision restoration in animal models using various molecules as optogenetic actuators[2–9]. In

addition, clinical trials are under way to investigate the effects of introducing

channelrhodopsin 2 (RST-001, ClinicalTrials.gov Identifier: NCT01648452) and

ChrimsonR (GS-030, ClinicalTrials.gov Identifier: NCT03326336) into retinal ganglion

cells (RGCs) via gene transduction achieved by intravitreal injection of recombinant

adeno-associated virus (rAAV). However, microbial opsins, such as channelrhodopsin 2,

require high light intensity, such as outdoor light intensity levels, to function[10–12]. They

cannot restore vision in dimly lit environments, such as indoors or at night, and strong

light irradiation can promote retinal degeneration[13,14]. Physiological photoreception

mediated by mammalian rhodopsin, however, relies on amplification through G protein

activation. Although the introduction of human rhodopsin improved photosensitivity in

mice, it is unclear how the chromophore retinal is metabolized in the retina where the

visual cycle is broken. Animal rhodopsin also causes toxicity if all-trans retinal is not

properly metabolized[15,16], and is, thus, hampered by safety and stability concerns in terms

of clinical application.

In this study, to achieve light sensitivity, stability, and safety, we attempted to restore

vision in mice using *Gloeobacter* and human chimeric rhodopsin (GHCR)[17,18].

Results

Design of GHCR

Although there is no sequence identity between microbial and animal opsin, both possess

similar chromophore (retinal) and protein (seven-transmembrane helix) structures. As we

previously reported[18], to generate GHCR, we replaced the second and third intracellular

loops of *Gloeobacter* rhodopsin with human sequences and introduced the E132Q

mutation (**Figure S1**). Previous work has shown that GHCR induces G protein activation

in vitro[18].

Restoring light-evoked activity in the retina with GHCR

We injected a viral vector (rAAV-DJ or rAAV-2) containing the GHCR coding sequence

under the control of the hybrid promoter comprising the CMV immediate-early enhancer,

CBA promoter, and CBA intron 1/exon 1, known as the CAGGS promoter,

(CAGGS-GHCR; **Figure 1a**) into the vitreous humor of 10-week-old *rd1* mice. We

adopted the rAAV-DJ vector to achieve more efficient, widespread gene transfer[19,20], and

used rAAV-2 as a benchmark, as it has already been used in the clinic[21]. The retinas were

harvested 2–4 months later. Enhanced green fluorescent protein (EGFP) reporter gene

expression was observed in the retina and in both the ganglion cell layer and the inner

nuclear layer (**Figure S2a, b**). To evaluate the function of ectopically expressed GHCR in

the mouse retina, we performed multi-electrode array (MEA) recording to record the

extracellular potential of RGCs (**Figure S2c**). As a result of photoreceptor degeneration,

the untreated control retina showed no RGC response as detected by MEA (**Figure 1b**). In

contrast, the treated retinas showed obvious light-induced responses down to 10^{14}

photons/cm^2/s of white light-emitting diode (LED) irradiation (**Figure 1c**).

Next, to create a stable vector for human gene therapy, we designed a codon-optimized

version of GHCR (coGHCR) and fused the ER2 endoplasmic reticulum (ER) export signal

to its C-terminus to increase gene expression levels. Immunolabelling revealed expression

across the whole retina, including in the bipolar cells, of treated *rd1* mice (**Figures 1d**). As

a result, the firing rate increased significantly, and a photoresponse was confirmed down

to 10^{13} photons/cm^2/s, which had not observed before optimization (**Figure 1e, f**).

Moreover, the number of firing cells per unit area also increased significantly (**Figure 1g**).

Since rhodopsin shows selectivity for Gi/o class G proteins upon heterologous

expression[22–25], we measured Gi/o activation with a homogeneous time-resolved

fluorescence (HTRF) cyclic adenosine monophosphate (cAMP) assay. We observed a 5-

fold increase in activation in coGHCR-treated compared with GHCR-treated mice (**Figure**

1h). The maximum spectral sensitivity of retinas treated with coGHCR was around 500

nm, and a photoresponse was obtained even upon stimulation with light with a wavelength

>600 nm (**Figure 1i**).

Restoration of visual cortex responses by GHCR

To investigate whether retinal light responses were transmitted to the visual cortex, we

then examined visual evoked potentials (VEPs) generated by the visual cortex (**Figure**

2a). The output from the RGCs is sent through their axons (optic nerve) to the lateral

geniculate nucleus (LGN) of the thalamus, which is a region of the diencephalon, then

from the LGN to the primary visual cortex in the occipital lobe of the cerebral cortex. For

these experiments, we used *rd1* mice in which both eyes had been treated with the AAV-

DJ-CAGGS-GHCR, AAV-DJ-CAGGS-coGHCR, or control EGFP (AAV-DJ-CAGGS-

EGFP) vectors. Significant VEPs were not detected in the control or GHCR-treated mice.

In contrast, VEPs were observed in coGHCR-treated mice (**Figure 2b**). In response to 3

cd·s/m^2 light stimulation, the average VEP amplitude in coGHCR-treated mice was

significantly higher (56.4 µV; n = 6) than those in GHCR-treated mice (22.1 µV; n = 8)

and control mice (17.9 µV; n = 6) (**Figure 2c**). Based on this result, all subsequent

experiments were performed using coGHCR.

Characterization of the in vivo responses restored by GHCR transduction

Next, light-dark transition (LDT) testing was performed to investigate whether ectopic

expression of coGHCR in degenerating retinas led to behavioral changes due to vision

restoration (**Figure 3a**). Rodents with intact vision tend to stay in dark places as they are

nocturnal and feel uneasy in bright environments, whereas blind rodents spend roughly

half of their time in bright places. The coGHCR-treated mice spent significantly less time

in the bright area compared with the untreated *rd1* mutant mice (**Figure 3b**), thereby

confirming vision restoration via behavioral analysis. Furthermore, in order to directly

compare the effects of coGHCR with genes in clinical trials, we treated *rd1* mice with

chimeric rhodopsin (AAV-6-CAGGS-coGHCR), microbial opsin (AAV-6-CAGGS-

ChrimsonR[26]), animal rhodopsin (AAV-6-CAGGS-human rhodopsin), or the control EGFP

(AAV-6-CAGGS-EGFP) vector. At an illuminance of 3,000 lux, a significant reduction in

the time spent in the bright half of the observation area was noted for coGHCR-treated

mice (0.32; n = 6) compared with control mice (0.50; n = 8) (**Figure 3c**). A similar

tendency was observed in ChrimsonR-treated mice (0.36; n = 6). However, no obvious

change was observed in human rhodopsin-treated mice (0.48; n = 6). When the experiment

was carried out at an illumination of 10 lux, human rhodopsin-treated mice showed a

significant change in the time spent in the bright area (0.40; n = 6), whereas ChrimsonR-

treated mice did not show an obvious change (0.55; n = 6) (**Figure 3d**). The coGHCR-

treated mice again spent significantly less time in the bright area illuminated at 10 lux

(0.40; n = 6).

Restored object recognition function upon GHCR gene therapy

LDT testing measures only light and dark discrimination. Visual recognition testing

(VRT) was performed to evaluate whether the mice could recognize an object with the

restored level of vision. Mice use vision for their cognitive functions, and are attracted to

fighting videos[27-29]. We examined mice in a place preference apparatus with a tablet

showing a fighting video (**Figure 3e**). The ratio of the time spent in the area with the

fighting compared with the time spent in the control area (showing a video of an empty

cage with the same illuminance) over 15 minutes was measured. The coGHCR-treated

(AAV-DJ-CAGGS-coGHCR) mice spent significantly more time in the fighting video half

of the apparatus (0.55, n = 33) than the untreated *rd1* mice (0.50, n = 30). On the other

hand, microbial opsin-treated (AAV-DJ-CAGGS-C1V1[30]) mice spent roughly equivalent

time in each half (0.49, n = 20) (**Figure 3f**).

GHCR protective effects against retinal degeneration

We employed another mouse model of retinal degeneration using mice with the P23H

RHO mutation, referred to as P23H mice[31]. P23H mice were selected to evaluate the

protective effect because they have slower retinal degeneration than *rd1* mice. We

subretinally delivered AAV DJ-CAGGS-coGHCR and the control (AAV DJ-CAGGS-

EGFP) vector into postnatal day (PND) 0–1 mouse retinas, targeting the outer retina, and

quantified the protective effects of the vector via morphological and electrophysiological

examination. Subretinal injection of AAV-DJ efficiently induced gene expression in the

murine outer retina (**Figure 4a**). Optical coherence tomography (OCT) showed that the

outer retinal thickness (ORT), which is the thickness from the outer nuclear layer (ONL)

to the rod outer segment (ROS), of coGHCR-treated mice (50.0 μm; n = 13) was

significantly greater than that of the control mice (42.7 μm; n = 10) at PND 30 (**Figure 4b,**

c). The ORT of the treated mice remained significantly greater than that of control mice

until PND 50 (**Figure S3**).

Electroretinography (ERG) revealed that the treated mice had larger rod, mixed, and cone

162 response amplitudes (141.2 µV, 271.4 µV, and 159.0 µV, respectively; n = 9) than the

163 control mice (70.4 µV, 158.7 µV, and 99.1 µV, respectively; n = 14) at PND 30 (**Figure**

164 **4d, e**). All amplitudes in the control mice gradually decreased, whereas all amplitudes in

165 the coGHCR-treated mice continued to increase until PND 42 (**Figure S4a, b c**).

166 Thereafter, the amplitudes in the treated mice also gradually decreased, although they

167 remained significantly higher than those in the control mice until PND 66.

168  We also performed terminal deoxynucleotidyl transferase dUTP nick end labeling

169 (TUNEL) to detect apoptosis in the retinas. The number of TUNEL-positive cells in the

170 coGHCR-treated mouse ONL (289.7 cells; n = 3) was significantly lower than that in the

171 control mouse ONL (67.3 cells; n = 3) at PND 31 (**Figure 5a–c**).

172  To expand these observations, we obtained transmission electron microscopy (TEM)

173 images of transverse sections from PND 31 mice. Consistent with the OCT results, the

174 ONL (**Figure 5d**) and ROS (**Figure 5e**) of coGHCR-treated mice were relatively intact

175 compared with those of controls, and the ROS structure was less disorganized (**Figure 5f**).

176 In addition, coGHCR-treated mice had less swelling of their ER, a feature that is

177 indicative of ER stress (**Figure 5g**).

178  We also performed western blotting to investigate ER stress. Expression of the ER stress

179 marker ATF4 was significantly lower, and expression of BiP, PERK, ATF6, and pIRE1

tended to be lower in treated mice than in control mice at PND 14 (**Figure S5a, b**).

Discussion

In this study, we demonstrated that ectopic expression of coGHCR is an effective

method of vision restoration in mice with retinal degeneration. MEA revealed that

photoresponses were maintained for retinal irradiance levels as low as 10^{13} photons/cm^2/s.

This is consistent with the response of the treated mice to 10 lux illumination in the

behavioral test, and represents a significant improvement in sensitivity compared with that

observed in previous studies of vision restoration with microbial opsins (threshold: 10^{14} to

10^{17} photons/cm^2/s)[2–7], LiGluR/MAG photoswitches (threshold: 10^{15}–10^{16}

photons/cm^2/s)[32,33], or photoactivated ligands (AAQ threshold: 10^{15} photons/cm^2/s[34] and

DENAQ threshold: 4×10^{13} photons/cm^2/s[35]). Although some vectors restored greater

sensitivity, such as human rhodopsin and Opto-mgluR6 (10^{12} photons/cm^2/s)[8], our LDT

results at 3,000 lux (similar to a cloudy outdoor environment) suggest that photobleaching

of rhodopsin like these does not work in bright environments. coGHCR is adaptable to a

light environment ranging from at least 10 lux (similar to a night light levels with

streetlights) to 3,000 lux, and is, thus, a suitable single-opsin vision restoration tool.

Furthermore, the typical channelrhodopsins have a spectrum limited to blue light[36],

which limits their use as a visual restoration tool. On the other hand, GHCR has a

spectrum peak around 500 nm and facilitates responses to red light. In addition, behavioral

tests showed that coGHCR enabled responses to both sustained and transient stimulation

lasting 10 ms. These findings suggested that coGHCR gene therapy can restore sensitivity

to multiple light environments encountered in daily life.

The ERG amplitudes in coGHCR-treated mice continued to increase until PND 42, likely

because the coGHCR-mediated signal was additive with the innate amplitude. This is

consistent with the fact that gene expression of the AAV-DJ vector peaks at approximately

1.5 months after administration[19]. We observed no apparent changes in the shapes of the

ERG waveforms in the coGHCR-treated mice.

coGHCR is known to activate Gt and Gi/o pathways; when expressed ectopically in ON

bipolar cells, it is expected to inhibit responses. However, the restored responses observed

by MEA were all ON responses. In addition, the electrophysiological and behavioral

results were similar to physiological responses, and no reversal reaction observed. In *rd1*

mice, photoreceptors are mostly lost by 4 weeks after birth and no optical response is

obtained after 7 weeks at the latest[37,38]. Therefore, responses from residual photoreceptors

are unlikely in this study. A similar phenomenon has been confirmed in previous studies;

the excitatory response is hypothesized to result from disinhibition of inhibitory amacrine

cells[6,8,9].

The safety of ectopic expression of opsins, such as channelrhodopsin 2, has been

previously reported[3,39,40]. To our knowledge, this is the first report of their protective

effects against retinal degeneration. In vitro studies have shown that the P23H opsin is

misfolded and retained in the ER[41]. ER retention of P23H opsin can induce the unfolded

protein response, leading to apoptosis[42–45]. Our results suggest that expression of coGHCR

in the retinal outer layer suppressed ER stress and photoreceptor apoptosis, which led to

protection against degeneration. The lack of 11-cis-retinal induces cytotoxicity during the

development of ROS in P23H mice[46]. Since coGHCR uses all-trans retinal as a

chromophore, like microbial opsin, it does not consume cis-retinal and is free from

photobleaching. Therefore, the expressed coGHCR may suppress cis-retinal consumption

via photoreceptor substitution. If this hypothesis is correct, the protection effect of

coGHCR may not be applicable to patients with all IRD genotypes. However, there are

more than 140 known RP-linked rhodopsin mutations, and those that result in protein

misfolding and retention in the ER are the most prevalent[47,48].

In summary, the coGHCR vector has the advantages of both animal and microbial opsin

as a vision regeneration tool. It restores sensitivity and an action spectrum that enables

vision in lighting ranging from levels found outdoors to those in dimly lit indoor

environments via G protein stimulation without the risk of bleaching; it can also be

expected to protect against the progression of retinal degeneration in the majority of IRD

patients. These results suggest that coGHCR is worthy of consideration for clinical

application as a gene therapy for IRD.

Methods

Study approval: All of the animal experiments were conducted in accordance with

protocols approved by Institutional Animal Care and Use Committee of Keio University

School of Medicine (#2808).

Please see supplemental information for detail.

Author Contribution

Y.K. and T.K. designed the research, wrote the manuscript. Y.K. performed the retinal

histology, MEA, ERG and VEP recordings, and WB, TEM, LDT and VRT experiments.

K.Y. performed plasmid construction. K.K. performed AAV production. Y.K. performed

data processing and analysis. H.K., H.O and T.K. made critical revisions of the

manuscript. T.K. supervised the research.

Acknowledgements

Y.K. was supported by grants from the Keio University Doctorate Student Grant-in-Aid

Program and Grant-in-Aid for Research Activity Start-up. T.K. was supported by Grants-

in-Aid from Takeda Science Foundation and the Keio University Medical Science Fund.

We would like to thank Editage (www.editage.com) for English language editing.

Declaration of Interests

Y.K., H.K., K.T., and T.K. are inventors on pending patents (PCT/JP2017/031579、

PCT/JP2019/ 1565) related to this work. Y.K., K.T., and T.K. are equity holders in Restore

Vision, Inc.

References

1.   Sahel, J.-A., Marazova, K. & Audo, I. Clinical characteristics and current

therapies for inherited retinal degenerations. *Cold Spring Harb. Perspect. Med.* **5**,

a017111 (2014).

2.   Bi, A. *et al.* Ectopic Expression of a Microbial-Type Rhodopsin Restores Visual

Responses in Mice with Photoreceptor Degeneration. *Neuron* **50**, 23–33 (2006).

3.   Doroudchi, M. M. *et al.* Virally delivered Channelrhodopsin-2 Safely and

Effectively Restores Visual Function in Multiple Mouse Models of Blindness.

Mol. Ther. **19**, 1220–1229 (2011).

4.   Lagali, P. S. *et al.* Light-activated channels targeted to ON bipolar cells restore

visual function in retinal degeneration. *Nat. Neurosci.* **11**, 667–675 (2008).

5.   Cronin, T. *et al.* Efficient transduction and optogenetic stimulation of retinal

bipolar cells by a synthetic adeno-associated virus capsid and promoter. *EMBO*

Mol. Med. **6**, 1–16 (2014).

6.   Macé, E. *et al.* Targeting channelrhodopsin-2 to ON-bipolar cells with vitreally

administered AAV restores ON and OFF visual responses in blind mice. *Mol.*

Ther. **23**, 7–16 (2014).

7.   Busskamp, V. *et al.* Genetic Reactivation of Cone Photoreceptors Restores

Visual Responses in Retinitis Pigmentosa. *Science (80-.).* **329**, 413–417 (2010).

8.   Van Wyk, M., Pielecka-Fortuna, J., Löwel, S. & Kleinlogel, S. Restoring the ON

Switch in Blind Retinas: Opto-mGluR6, a Next-Generation, Cell- Tailored

Optogenetic Tool. *PLoS Biol* **13**, (2015).

9.   Pienaar, A., Bedford, R., Davis, K. & Bishop, P. N. Restoration of vision with

ectopic expression of human rod opsin. *Curr. Biol.* **25**, 1–35 (2015).

10.   Kleinlogel, S. *et al.* Ultra light-sensitive and fast neuronal activation with the Ca

2+-permeable channelrhodopsin CatCh. *Nat. Neurosci.* **14**, 513–518 (2011).

11.   Kleinlogel, S. *et al.* A gene-fusion strategy for stoichiometric and co-localized

expression of light-gated membrane proteins. *Nat. Methods* **8**, 1083–1091 (2011).

12.   Pan, Z. H., Ganjawala, T. H., Lu, Q., Ivanova, E. & Zhang, Z. ChR2 mutants at

L132 and T159 with improved operational light sensitivity for vision restoration.

PLoS One **9**, (2014).

13.   Grimm, C. *et al.* Protection of Rpe65-deficient mice identifies rhodopsin as a

mediator of light-induced retinal degeneration. *Nat. Genet.* **25**, 63–66 (2000).

14.   Vaughan, D. K., Coulibaly, S. F., Darrow, R. M. & Organisciak, D. T. A

morphometric study of light-induced damage in transgenic rat models of retinitis

pigmentosa. *Investig. Ophthalmol. Vis. Sci.* **44**, 848–855 (2003).

15.   Różanowska, M. & Sarna, T. Light-induced Damage to the Retina: Role of

299   Rhodopsin Chromophore Revisited. *Photochem. Photobiol.* **81**, 1305 (2005).

300   16.   Maeda, A. *et al.* Involvement of all-trans-retinal in acute light-induced

301   retinopathy of mice. *J. Biol. Chem.* **284**, 15173–15183 (2009).

302   17.   Nakatsuma, A. *et al.* Chimeric microbial rhodopsins containing the third

303   cytoplasmic loop of bovine rhodopsin. *Biophys. J.* **100**, 1874–1882 (2011).

304   18.   Sasaki, K. *et al.* Chimeric Proton-Pumping Rhodopsins Containing the

305   Cytoplasmic Loop of Bovine Rhodopsin. *PLoS One* **9**, e91323 (2014).

306   19.   Katada, Y., Kobayashi, K., Tsubota, K. & Kurihara, T. Evaluation of AAV-DJ

307   vector for retinal gene therapy. *PeerJ*

308   20.   Grimm, D. *et al.* In vitro and in vivo gene therapy vector evolution via

309   multispecies interbreeding and retargeting of adeno-associated viruses. *J. Virol.*

310   **82**, 5887–5911 (2008).

311   21.   Keeler, A. M. & Flotte, T. R. Recombinant Adeno-Associated Virus Gene

312   Therapy in Light of Luxturna (and Zolgensma and Glybera): Where Are We, and

313   How Did We Get Here? *Annu. Rev. Virol.* **6**, 601–621 (2019).

314   22.   Bailes, H. J. & Lucas, R. J. Human melanopsin forms a pigment maximally

315   sensitive to blue light (λmax $\approx$ 479 nm) supporting activation of Gq/11 and Gi/o

signalling cascades. *Proc. R. Soc. B Biol. Sci.* **280**, (2013).

23. Li, X. *et al.* Fast noninvasive activation and inhibition of neural and network

activity by vertebrate rhodopsin and green algae channelrhodopsin. *Proc. Natl.*

Acad. Sci. U. S. A. **102**, 17816–17821 (2005).

24. Gutierrez, D. V. *et al.* Optogenetic control of motor coordination by G i/o

protein-coupled vertebrate rhodopsin in cerebellar purkinje cells. *J. Biol. Chem.*

286, 25848–25858 (2011).

25. Cao, P. *et al.*   Light-sensitive coupling of rhodopsin and melanopsin to G i/o

and G q signal transduction in Caenorhabditis elegans . *FASEB J.* **26**, 480–491

(2012).

26. Klapoetke, N. C. *et al.* Independent optical excitation of distinct neural

populations. *Nat. Methods* **11**, 338–346 (2014).

27. Watanabe, S., Shinozuka, K. & Kikusui, T. Preference for and discrimination of

videos of conspecific social behavior in mice. *Anim. Cogn.* **19**, 523–531 (2016).

28. Watanabe, S. Preference for and Discrimination of Paintings by Mice. *PLoS One*

8, (2013).

29. Yakura, T. *et al.* Visual recognition of mirror, video-recorded, and still images in

rats. *PLoS One* **13**, 1–11 (2018).

30.   Erbguth, K., Prigge, M., Schneider, F., Hegemann, P. & Gottschalk, A. Bimodal

335   Activation of Different Neuron Classes with the Spectrally Red-Shifted

336   Channelrhodopsin Chimera C1V1 in Caenorhabditis elegans. *PLoS One* **7**,

337   (2012).

338   31.   Sakami, S., Kolesnikov, A. V., Kefalov, V. J. & Palczewski, K. P23H opsin

339   knock-in mice reveal a novel step in retinal rod disc morphogenesis. *Hum. Mol.*

340   *Genet.* **23**, 1723–1741 (2014).

341   32.   Caporale, N. *et al.* LiGluR restores visual responses in rodent models of inherited

342   blindness. *Mol. Ther.* **19**, 1212–1219 (2011).

343   33.   Gaub, B. M. *et al.* Restoration of visual function by expression of a light-gated

344   mammalian ion channel in retinal ganglion cells or ON-bipolar cells. (2014).

345   doi:10.1073/pnas.1414162111

346   34.   Polosukhina, A. *et al.* Photochemical Restoration of Visual Responses in Blind

347   Mice. *Neuron* **75**, 271–282 (2012).

348   35.   Tochitsky, I. *et al.* Restoring visual function to blind mice with a photoswitch

349   that exploits electrophysiological remodeling of retinal ganglion cells. *Neuron*

350   **81**, 800–813 (2014).

36.   Nagel, G. *et al.* Channelrhodopsin-2, a directly light-gated cation-selective

membrane channel. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 13940–5 (2003).

37. Fujii, M., Sunagawa, G. A., Kondo, M., Takahashi, M. & Mandai, M. Evaluation

of micro Electroretinograms Recorded with Multiple Electrode Array to Assess

Focal Retinal Function. *Sci. Rep.* 1–12 (2016). doi:10.1038/srep30719

38. Chang, B. *et al.* Two mouse retinal degenerations caused by missense mutations

in the beta-subunit of rod cGMP phosphodiesterase gene. *Vision Res.* **47**, 624–33

(2007).

39. Sugano, E. *et al.* Immune responses to adeno-associated virus type 2 encoding

channelrhodopsin-2 in a genetically blind rat model for gene therapy. *Gene Ther.*

18, 266–274 (2011).

40. Ivanova, E. & Pan, Z. H. Evaluation of the adeno-associated virus mediated long-

term expression of channelrhodopsin-2 in the mouse retina. *Mol. Vis.* **15**, 1680–

1689 (2009).

41. Chapple, J. P. *et al.* Unfolding retinal dystrophies: A role for molecular

chaperones? *Trends Mol. Med.* **7**, 414–421 (2001).

42. Liu, X., Garriga, P. & Khorana, H. G. Structure and function in rhodopsin:

Correct folding and misfolding in two point mutants in the intradiscal domain of

rhodopsin identified in retinitis pigmentosa. *Proc. Natl. Acad. Sci. U. S. A.* **93**,

4554–4559 (1996).

43.   Frederick, J. M. *et al.* Mutant rhodopsin transgene expression on a null

372   background. *Investig. Ophthalmol. Vis. Sci.* **42**, 826–833 (2001).

373   44.   Leonard, K. C. *et al.* XIAP protection of photoreceptors in animal models of

374   retinitis pigmentosa. *PLoS One* **2**, (2007).

375   45.   Lin, J. H. *et al.* IRE1 signaling affects cell fate during the unfolded protein

376   response. *Science (80-.).* **318**, 944–949 (2007).

377   46.   Sakami, S. *et al.* Probing mechanisms of photoreceptor degeneration in a new

378   mouse model of the common form of autosomal dominant retinitis pigmentosa

379   due to P23H opsin mutations. *J. Biol. Chem.* **286**, 10551–10567 (2011).

380   47.   Mendes, H. F., Van Der Spuy, J., Chapple, J. P. & Cheetham, M. E. Mechanisms

381   of cell death in rhodopsin retinitis pigmentosa: Implications for therapy. *Trends*

382   *Mol. Med.* **11**, 177–185 (2005).

383   48.   Yoshida, T. *et al.* The use of induced pluripotent stem cells to reveal pathogenic

384   gene mutations and explore treatments for retinitis pigmentosa. *Mol. Brain* **7**, 1–

385   11 (2014).

388   **Figure legends**

Figure 1. Ectopic GHCR expression restores light responses in the *rd1* mouse retina

(a) DNA expression cassette schematic. The GHCR coding sequence is driven by the

CAGGS promoter, flanked by inverted terminal repeats (ITR), and stabilized by a

polyadenylation signal sequence (pA) and a woodchuck hepatitis posttranscriptional

regulatory element (WPRE). (b, c, e) Raster plots and peri-stimulus time histograms for

light stimulation of control (AAV-DJ-CAGGS-EGFP) (b), GHCR-treated (AAV-DJ-

CAGGS-GHCR) (c), and coGHCR-treated (AAV-DJ-CAGGS-coGHCR) mice (e).

Responses to exposure to a white LED with varying light intensity for 1.0 s. Gray shading

around the averaged traces represents the standard error of the mean (SEM). (d) Confocal

image of a transverse *rd1* mouse retina section 2 months after AAV-DJ-CAGGS-coGHCR

intravitreal injection. Green, FLAG tag antibody signal (vector); red, PKCα signal

(bipolar cells); blue, 4′,6-diamidino-2-phenylindolenuclear (DAPI) counterstaining.

Scale bar, 50 μm. (f) Quantitation of the firing rates of RGCs transduced with GHCR or

coGHCR at the indicated light intensity. (g) Histogram showing the number of RGCs that

responded to light per unit area (2.6 mm^2) of the retinas of GHCR- or coGHCR-treated

mice (n = 3 each). (g) Changes in cAMP consumption in response to Gi/o-coupled G-

protein-coupled receptor activation in HEK293T cells transfected with GHCR and

coGHCR (n = 3 each). (h) Spectral sensitivity induced by coGHCR. Error bars represent

the SEM. Data were analyzed with Student's two-tailed t-test; * represents $p < 0.05$, **

represents $p < 0.01$, and *** represents $p < 0.001$.

Figure 2. coGHCR restored vision in *rd1* mice through the primary visual cortex

(a) Schematic view of the VEP recording strategy. (b) Representative VEP traces from

GHCR-treated, coGHCR-treated, and control mice. (c) The average amplitude of the

VEPs in the control (AAV-DJ-CAGGS-EGFP), GHCR-treated (AAV-DJ-CAGGS-

GHCR), and coGHCR-treated (AAV-DJ-CAGGS-coGHCR) mice. The stimulus was a

white LED flash (3 cd·s/m^2). Signals were low-pass filtered at 300 Hz and averaged over

60 trials. Error bars represent the SEM. Data were analyzed with one-way analysis of

variance (ANOVA) and Tukey's multiple comparison test; * represents $p < 0.05$. V1,

visual cortex.

Figure 3. coGHCR-treated mouse behavior indicated vision restoration

(a) LDT testing schematic. Mice were tested in a $30 \times 45 \times 30$-cm box with equally sized

bright and dark chambers connected by a 5×5-cm opening, across which the mice could

move freely. (b) Percentage of time spent in the bright area (total, 10 min) by wild type (n

= 4), and control (AAV-DJ-CAGGS-EGFP) (n = 7) and coGHCR-treated (AAV-DJ-

CAGGS-coGHCR) *rd1* mice (n = 6). LDT test, 10 lux illumination. (c, d) The percentage

of time spent in the bright area (total, 10 min) by wild type (n = 6), and control (AAV-6-

CAGGS-EGFP) (n = 8), coGHCR-treated (AAV-6-CAGGS-coGHCR) (n = 6),

ChrimsonR-treated (AAV-6-CAGGS-ChrimsonR) (n = 6), and human rhodopsin-treated

(AAV-6-CAGGS-human-rhodopsin) *rd1* mice (n = 6). LDT test with 3,000 lux (c) and 10

lux (d) illumination. (e) VRT setup. Time spent in areas showing a video of mice fighting

(object half, blue) or an empty cage (control half, red) was measured. (f) Distribution of

time spent in the object half by wild type (n = 6), and control (no treatment) (n = 8), AAV-

2-coGHCR-treated (AAV-2-CAGGS-coGHCR) (n = 6), AAV-DJ-coGHCR-treated

(AAV-DJ-CAGGS- coGHCR) (n = 6), and AAV-DJ-C1V1-treated (AAV-DJ-CAGGS-

C1V1) *rd1* mice (n = 6). LDT test with 10 lux (c) and 3,000 lux (d) illumination. Black

line, average value. Error bars represent the SEM. Data were analyzed with one-way

ANOVA and Tukey's multiple comparison test; * represents $p < 0.05$, ** represents $p <$

0.01, and *** represents $p < 0.001$.

Figure 4. Ectopic coGHCR expression protects against photoreceptor degeneration

(a) Confocal image of a transverse section through the P23H retina 2 months after AAV-

DJ-CAGGS-coGHCR subretinal injection. Green, FLAG tag fused to the C-terminus of

coGHCR; blue, DAPI nuclear counterstaining. Scale bar, 100 μm. (b) OCT retinal image

sections from coGHCR-treated and control (AAV-DJ-CAGGS-EGFP subretinally

injected) mice at PND 30. The white arrow indicates the measured ORT (from ONL to

cone outer segment). Scale bar, 20 μm. (c) Histogram of the measured ORT of the

coGHCR-treated and control mice at PND 30. (d, e) Representative ERG waveforms (rod

response, mixed response, and cone response) of coGHCR-treated and control mice (d).

Histograms of the average ERG amplitudes from panel d at PND 30 (e). Error bars

represent SEM. Data were analyzed with the unpaired t-test; *** represents $p < 0.001$.

GCL, ganglion cell layer; INL, inner nuclear layer.

Figure 5. coGHCR treatment suppressed retinal apoptosis and ER stress

(a, b) TUNEL-stained transverse sections (a) and enlarged images of the white squares (b)

of coGHCR-treated and control (AAV-DJ-CAGGS-EGFP subretinally injected) mouse

retinas at PND 31. Red, TUNEL-positive cells; blue, DAPI nuclear counterstaining. Scale

bar, 1,000 μm in (a) and 100 μm in (b). (c) Histogram of the number of TUNEL-positive

cells in the ONLs of coGHCR-treated and control mice at PND 31. Error bars represent

SEM. Data were analyzed with the unpaired t-test; ** represents $p < 0.01$. (d) TEM

images of transverse sections from coGHCR-treated and control mice at PND 31, showing

the outer retinal layer (d), the outer segment at low magnification (e) and high

magnification (f), and the inner segment (g). The arrowhead indicates swollen ER. Scale

bar, 20 μm in (d), 5 μm in (e), 1 μm in (f), and 500 nm in (g).