Title:

Persistence and invasiveness of high-level heteroplasmy through biparental transmission of a selfish mitochondrion in *Drosophila*

Running title: Biparental transmission of a selfish mitochondrion in *Drosophila*

Authors: Guilherme C. Baião[1,§], Anton Strunov[2,§], Eleanor Heyworth[1,§], Daniela I. Schneider[2,*], Julia Thoma[2], Lisa Klasson[1,#], Wolfgang J. Miller[2,#]

Author affiliations: [1] Molecular Evolution, Department of Cell and Molecular Biology, Science for Life Laboratory, Uppsala University, Husargatan 3, 751 24, Uppsala, Sweden; [2] Lab Genome Dynamics, Department Cell & Developmental Biology, Center for Anatomy and Cell Biology, Medical University of Vienna, Schwarzspanierstraße 17, 1090, Vienna, Austria.

§ The authors contributed equally

\# Corresponding authors, and equal contribution

* Current address: Moffitt Cancer Center, 12902 Magnolia Drive, Tampa, FL 33612, USA.

Emails:

guilherme.baiao@icm.uu.se
anton.strunov@meduniwien.ac.at
heyworth.eleanor@gmail.com
daniela.schneider@moffitt.org
thoma.julia@yahoo.de
lisa.klasson@icm.uu.se
wolfgang.miller@meduniwien.ac.at

Keywords: Heteroplasmy, mitochondria, biparental & paternal transmission, selfish genetic elements, Drosophila, evolution, genomics, introgression, development.

ABSTRACT

Heteroplasmy is the coexistence of more than one type of mitochondria in an organism. Although widespread sequencing has identified several cases of transient or low-level heteroplasmy that primarily occur through mutation or paternal leakage, stable, high-titer heteroplasmy remains rare in animals. In this study we present a unique, stable and high-level heteroplasmy in male and female flies belonging to the neotropical *Drosophila paulistorum* species complex. We show that mitochondria of *D. paulistorum* are polyphyletic and form two clades, α and β, with two subclades each. Mitochondria of the $\alpha2$ subclade appear functional based on their genomic integrity but are exclusively found in heteroplasmic flies and never in homoplasmy, suggesting that they are a secondary mitotype with distinct functionality from the primary mitochondria. Using qPCR, we show that $\alpha2$ titer do not respond to energetic demands of the cell and are generally higher in males than females. By crossing hetero- and homoplasmic flies, we find that $\alpha2$ can be transmitted to their offspring via both parents and that levels are dependent on nuclear background. Following $\alpha2$ mitotype levels during embryogenesis, we demonstrate that this secondary mitotype replicates rapidly just after fertilization of the egg in a period when primary mitochondria are dormant. This so-called "Replication precox" mitochondrial phenotype likely prevents the $\alpha2$ mitotype from being outcompeted by the primary mitotype – and thereby secures its persistence and further spread as a selfish mitochondrion, we hereby designate "Spartacus". Finally, we reconstruct the evolutionary history of mitochondria in the willistoni subgroup uncovering signs of multiple mitochondrial losses and introgressions. Our data indicate an α-like mitochondrial ancestor in the willistoni subgroup, with the β mitotype likely acquired via introgression from an unidentified donor. We hypothesize that the selfish characteristics of $\alpha2$ might have emerged as a response to competition for inheritance with the introgressed β mitotype.

INTRODUCTION

It is generally assumed that individuals of most animal species carry a single, maternally inherited mitochondrial haplotype (a mitotype) [1, 2]. However, recent advances in sequencing technology and genomic analyses show that mitochondrial heteroplasmy, the presence of multiple mitotypes in one individual, is found in a number of vertebrate and invertebrate species [3, 4].

Most often, heteroplasmy is the result of either mitochondrial mutations that accumulate over an individual's lifetime, or paternal leakage. In both cases, the additional mitotype is usually transient and persists at low levels within an individual [5]. Most host-mitochondria relationships are ancient, with both parties coevolving over extended periods of time, and disturbances caused by the introduction and persistence of new mitotypes may reduce fitness or cause disease [6, 7]. Stable, long-term heteroplasmy is therefore relatively uncommon and may be evolutionarily disadvantageous (Reviewed in [8]). Accordingly, selection against heteroplasmy has been proposed as an explanation for the evolution of uniparental inheritance in mitochondria [9]. Mechanistically, transmission of male mitochondria in animals is prevented by both pre- and postmating defense and destruction mechanisms of the host ([5, 10-15] and references therein), and if these fail, the rare paternal mitotype is usually outcompeted by maternal mitochondria during early development or within subsequent generations [16].

The best-known case of animal heteroplasmy is that of freshwater mussels of the family *Unionidae*, which display double uniparental mitochondrial inheritance (DUI). In these species, two different mitochondrial haplotypes, M and F, are transmitted exclusively by males and females, respectively [17, 18]. Stable heteroplasmy is also found in several isopods in which mitochondria encode different tRNA variants [19-21]. Among insects, prevalence (the frequency of heteroplasmic individuals in a population) at various levels (the proportions of different mitotypes within an individual) has been described in natural populations of the bed bug *Cimex lectularius* L. [22], the neotropical ants of the *Ectatomma ruidum* complex [23], the brown dog tick *Rhipicephalus* spp. [24], and the leaf beetles *Gonioctena intermedia* [25]. For the genetic model system *Drosophila*, natural heteroplasmy has been detected in *D. mauritiana* [26], *D. simulans* [27-29], *D. melanogaster* [30], and *D. subobscura* [31, 32]. Slightly higher prevalence and levels have been found in interspecies F_1 hybrids of different closely related *Drosophila* species ([33-35] and references therein). In spite of a growing number of reports on heteroplasmy in animals including humans (recently reviewed in [36]), little is known as to how heteroplasmic systems have evolved and what the functional differences between mitotypes could be (recently reviewed in [37]).

However, when discussing causes and consequences of mitochondrial heteroplasmy, it might be useful to consider mitotypes as primary or secondary endosymbionts, similarly to

what is done for bacterial endosymbionts of insects where multiple infections and obligate mutualism are common phenomena. Hence, primary bacterial endosymbionts as well as mitochondria are normally maternally inherited, long-term co-evolved with the host and have a well-defined function that make them obligate for the eukaryotic host cell, while secondary bacterial endosymbionts and recently acquired or evolved mitotypes are non-fixed, can have variable, often condition-specific and sometimes harmful effects on their hosts [38-40]. While primary endosymbionts might evolve from secondary endosymbionts as a way for the host to secure higher fitness over long evolutionary time, secondary symbionts can be maintained either because they provide a benefit under specific conditions or because they have the ability to cheat the host and be transmitted selfishly without any positive effect [40]. Similarly, secondary mitochondria might thus be maintained in the host for the same reason and result in heteroplasmy (reviewed in [37]). Additionally, obligate dependencies on multiple endosymbionts evolved in several insect. In such cases, one symbiont has often lost genes so that the genome of the other contain complementary genes involved in a function that was previously performed by only one of them [41, 42]. Hence, similarly, stable heteroplasmy might also be maintained if mitotypes with complementary gene sets are present in the same host.

The Neotropical *Drosophila willistoni* group has been a model for evolutionary studies since the 1950s [43]. Within it, species belonging to the willistoni subgroup have received the most attention, with *Drosophila paulistorum,* in particular, being the focus of many studies on speciation and hybridization [44, 45]. *D. paulistorum* is a complex of six semispecies under incipient speciation, namely Amazonian (AM), Andean Brazilian (AB), Centro-American (CA), Interior (IN), Orinocan (OR) and Transitional (TR) [46], each displaying partially overlapping geographical distribution and varying degrees of pre- and postmating reproductive incompatibilities with the others ([44], and references therein). All semispecies also carry the bacterial endosymbiont *Wolbachia*, which provides mutualistic functions and causes premating isolation between members of different semispecies pairs [47-51]. The whole willistoni subgroup is a recent radiation and phylogenetic incongruences between *D. paulistorum* nuclear and mitochondrial evolutionary histories [52-54] suggest that hybridization and introgression events between semispecies may have occurred in the past [53].

In the present study, we discovered and investigated the unusual biparental inheritance of mitochondria resulting in stable and high-level heteroplasmy in *Drosophila paulistorum*. We assembled the mitochondrial genomes of six species of the willistoni subgroup and three representatives of the closely related saltans species group. We discovered that *D. paulistorum* lines carry mitochondria from two non-sister clades - α and β - and identified several lines exhibiting stable heteroplasmy. We further determined that both mitotypes can be divided into two subclades, called 1 and 2, where the $\alpha2$ mitotype is so far observed exclusively in heteroplasmic lines. This finding implies that $\alpha2$ is unable to provide full mitochondrial functionality, and might therefore be regarded as a secondary mitotype, in

contrast to the fully functional primary mitotypes (α1, β1 and β2), which can exist in

homoplasmy.

In addition, the α2 mitotype was found in 41% of the tested *D. paulistorum* lines, where

it is stably maintained at variable but significant levels with up to 66% depending on host

background. Unexpectedly, this secondary mitotype, unlike the other primary mitochondria we

investigated, was not distributed within the fly in a way that correlated with the expected

OXPHOS-mediated ATP demands of different tissues. Yet, the α2 genome showed no signs

of deterioration, suggesting it may perform some unknown function for the host. Additionally,

we demonstrated that α2 mitochondria are biparentally transmitted and start replicating rapidly

after egg fertilization, while the primary mitochondria are still quiescent. This replication

advantage, which we name the "Replication precox" phenotype, allows the α2 to reach high

levels during early embryogenesis and thus to avoid being outcompeted by the much larger

number of maternally inherited primary mitochondria. Hence, we propose that α2 is a newly

discovered selfish mitotype that can persist and spread in and between populations, and

possibly even across species by introgression, via the combination of biparental transmission

and the Replication precox phenotype.

Using phylogenetic reconstruction, we observed a complex evolutionary history of

mitochondria in *D. paulistorum*, with both ancient and recent introgressions. We suggest that

an α-like mitochondrion was the ancestral mitotype of the species and that heteroplasmy might

have originated through introgression of the β mitotype from an unsampled or extinct donor,

a concept known as 'ghost' introgression [55-57]. We speculate that the unusual biparental

transmission mode of α2 might have developed through competition with the β mitochondria

for maternal transmission.

RESULTS

High level heteroplasmy is common in the *D. paulistorum* species complex

In the course of our phylogenetic survey we sequenced PCR products of the mitochondrial

Cytochrome Oxidase 1 (*CO1*) gene from several *D. paulistorum* lines for typing and found

multiple robust double peaks in the resulting chromatograms derived from single flies of the

Orinocan O11 line. The double peaks indicated the presence of two bases within a diagnostic

*Hin*dIII restriction site (**Fig 1A**). Thus, to independently confirm heteroplasmy, we performed

diagnostic RFLP/*Hin*dIII Southern blots on single fly DNA probed with *CO1*. We found that the

O11 line is truly heteroplasmic, with individual flies carrying two mitotypes, hereafter called α

and β, at approximately similar levels (**Fig 1B**). In contrast, single flies of the Centro-American

C2 and Amazonian A28 lines were homoplasmic and carried either the α or β mitotype,

respectively (**Fig 1B**). By using *CO1*-specific RFLP/*Hin*dIII-PCR we further showed that all

tested O11 flies (20 females, 20 males) exhibited both the α and β specific bands (**Fig 1C**),
which suggested fixation of the high-level heteroplasmic state in this *D. paulistorum* long-term
lab line.

167
Fig 1. High-level heteroplasmy in the *D. paulistorum* Orinocan line O11. **(A)** Chromatogram
derived from *CO1* direct Sanger sequencing of one O11 female covering the α-diagnostic
*Hin*dIII restriction site (underlined). The asterisk highlights the double peak with Cytosine (C)
for α and Thymine (T) for β-mitotype. **(B)** Single-fly *CO1* Southern blot digested with *Hin*dIII.
While the Amazonian females of the A28 line (n=3) are homoplasmic for the β-mitotype (4.8kb)
and the ones of the Centro-American line C2 (n=2) are homoplasmic for α (2.4kb), all randomly
picked Orinocan O11 females (n=3) show high-level heteroplasmy of the α- and β-mitotypes
(4.8 and 2.4kb). **(C)** α-diagnostic *CO1* RFLP/*Hin*dIII-PCR screen on O11 single flies (n=20
for each sex). Undigested (U) and digested (D) *CO1*-PCR fragments per fly are shown (bottom
right in **C**). As all tested single flies (40/40) exhibited clear double bands after *Hin*dIII digestion,
hence fixation of the α/β heteroplasmy in the O11 line was assumed.

Whole-genome sequencing shows widespread heteroplasmy forming two distinct
mitochondrial clades and four subclades in D. paulistorum

Next, we used Illumina sequencing to screen for heteroplasmy on a genome-wide level in *D.*
paulistorum and several related species. We sequenced a total of 23 *Drosophila* lines
representing six species of the willistoni group and three species of the closely related saltans
group (**Fig 2A, Table S1**). Thirteen *D. paulistorum* lines were sequenced, including
representatives of the six classical semispecies and seven recently collected isofemale lines

from Brazil and French Guiana (**Table S1**). Five of the sequenced *D. paulistorum* lines (38.5%)
were found to be heteroplasmic, i.e., the Orinocan FG103, FG111 and O11 lines, the
Amazonian FG572 line and the Andean-Brazilian MS line (**Fig 2B**). Importantly, heteroplasmy
was not detected in any other tested neotropical species apart from *D. paulistorum*.
190

191
192
Fig 2. Phylogenies of the sequenced *Drosophila* lines. Trees based on (**A**) the nuclear
genes *28S*, *Adh*, *CAD/r*, *Ddc*, *Hb* and *Per*; and (**B**) whole mitochondrial genomes. Fly lines in
bold are heteroplasmic. *D. paulistorum* mitochondria split into two distinct non-sister clades: α
(green) and β (blue). Support for the branch defining α as sister clade to other species of the
willistoni subgroup is low for whole genome (WG) phylogeny, but high when only first and
second codon positions of protein coding genes are analyzed (1+2). The same topologies are
obtained by maximum likelihood (GTRGAMMA model) and Bayesian analyses. Colored
circles next to the nodes indicate ML bootstrap support (left half) and Bayesian posterior
probability (right half). Black corresponds to 99-100% bootstrap support or 0.99-1.0 posterior
probability, grey represents 70-98% bootstrap support or 0.7-0.98 posterior probability, white
indicates support below 70%. Branches leading to the outgroups and marked by two
transverse lines were shortened and don't follow the presented scale.
205
Among all sequenced lines of willistoni and saltans group species, a total of 28 mitochondrial
genomes were assembled, one for each homoplasmic *Drosophila* line and two for each of the
heteroplasmic *D. paulistorum* lines (see Methods). The genomes are approximately 16.000
bp long, have a GC content around 21% (**Table S2**), and encode the same 13 proteins, 22
tRNAs and two rRNAs as the mitochondria of *D. melanogaster* and most other animals.
Importantly, no loss-of-function mutations were found, indicating that all genomes are likely to
encode fully functional mitochondria. The assembled mitochondrial genomes also show
considerably higher read coverage than nuclear contigs from the same lines, which

distinguishes them from potential nuclear mitochondrial DNA sequences (NUMTs) (**Table S3**).
All of the assembled mitochondrial genomes have short control regions (CR), between 1.1
kbp in the willistoni group and 1.3 kbp in the saltans group, compared to the 4.6 kbp CR in *D.*
melanogaster. A few substitutions occur between *D. paulistorum* lines in or near the putative
origins of replication, but no major differences in length are observed between CRs (**Table S2**,
Fig S1).
Phylogenetic analysis of the assembled genomes revealed that the mitochondria of *D.*
paulistorum are polyphyletic and split into two major clades (**Fig 2B**), the α and β mitotypes
that were previously detected in single flies of O11 (**Fig 1**). The mitochondrial genomes of the
α and β clades are relatively distantly related to each other with an average nucleotide
divergence of 2.94%. A majority of the substitutions that occur between them in protein coding
genes are synonymous (**Tables S4, S5**), indicating that purifying selection is acting on the
genomes of both clades. Based on the phylogeny, the α and β clades can be further divided
into two subclades, named 1 and 2 **(Fig 2B)**. Interestingly, while the $\alpha 1$, $\beta 1$ and $\beta 2$ mitotypes
occur both in homoplasmic and heteroplasmic lines, the $\alpha 2$ mitotype is never found in
homoplasmy and is present in all heteroplasmic lines, where it is partnered with a mitotype
from another subclade.
231
Level and prevalence of heteroplasmy in D. paulistorum lines
To further investigate the heteroplasmy of *D. paulistorum*, we assessed its prevalence (the
percentage of heteroplasmic flies within a line) and levels (percentage of each mitotype within
a sample) using mitotype-specific RFLP-PCR. Seventeen lines belonging to the six *D.*
paulistorum semispecies were tested and seven (41%) displayed heteroplasmy (**Table 1**).
Apart from the lines already identified as heteroplasmic through genome sequencing (**Fig 2B)**,
we further identified heteroplasmy in the two Amazonian lines FG16 and FG572 as well as in
the Andean-Brazilian Yellow lines. Interestingly, the Orinocan FG103 line was clearly
heteroplasmic at the time of sequencing, shortly after its field collection and the establishment
of the isofemale line in 2014/2015 (**Table 1**), but thereafter lost the $\alpha 2$ mitotype in consecutive
generations. Heteroplasmic lines were thus observed in the Amazonian, Andean-Brazilian and
Orinocan semispecies and originate from both long-term lab lines and recently collected field
samples (**Table S1**).

Table 1. Levels of α2 and prevalence of heteroplasmy in *D. paulistorum* lines based on Illumina sequencing of ovary samples (α2 level Illumina) and RFLP-PCR from adult flies (α2 level RFLP). A dash (-) in the α2 level from Illumina sequencing means that we are unable to determine that the α2 mitotype existed. Levels of α2 obtained through RFLP-PCR are presented separately for females (left) and males (right). Prevalence of heteroplasmy among tested individuals (Het. prevalence) is presented both as a percentage and in absolute numbers (in parenthesis). Abbreviations: AB: Andean-Brazilian, AM: Amazonian, CA: Centro-American, Coll: Collection year, Het: heteroplasmy, IN: Interior, ND: Not determined, OR: Orinocan, Spp: Semispecies, TR: Transitional. Further details about the fly lines are provided in **Table S1**.

Line	Coll.	Spp.	Mitotype	α2 level Illumina	α2 level RFLP	Het. prevalence
O11	1957	OR	α2/ß2	19.4%	30.7±3.7/36.8±3.6	100% (39/39)
FG103	2014	OR	α2/ß2	6.5%	ND§	0% (0/40)§
FG111	2014	OR	α2/ß2	2.3%	13.4±1.1/18.3±2.9	25% (10/40)
FG295	2014	OR	ß2	-	0	0% (0/40)
TP37	2009	OR	ß2	-	0	0% (0/20)
POA1	2003	OR	ß2	-	0	0% (0/20)
RP	1995	OR	ß2	-	0	0% (0/20)
A28	1952	AM	ß1	-	0	0% (0/40)
FG16	2018	AM	α2/ß1	ND	57.2±8.6/66.5±7.2	55% (22/40)
FG18	2018	AM	ß1	ND	0	0% (0/40)
FG572	2015	AM	α2/ß1	49.5%	45.3±8.1/63.7±7.0	90% (36/40)
MS	1962	AB	α1/α2	19.1%	21.8±3.1/36.0±4.1	100% (40/40)
Yellow	1960	AB	α1/α2	ND	20.4±4.7/27.4±4.3	100% (40/40)
C2	1954	CA	α1	-	0	0% (0/20)
White	1960	CA	α1	ND	0	0% (0/20)
L1	1958	IN	ß2	-	0	0% (0/20)
SM	1956	TR	ß2	-	0	0% (0/20)

§The line FG103 was heteroplasmic in 2014/15 (determined by Sanger and Illumina sequencing), but has lost the α2 mitotype afterwards (verified by RFLP-PCR and Sanger sequencing in 2018/19).

Whereas heteroplasmy is fixed in the three independent long-term lab lines O11, MS and Yellow, α2 prevalence ranges between 25% and 90% in the recently collected isofemale lines FG111, FG16 and FG572 (**Table 1**). Levels of α2 were high within individual flies of most lines (**Table 1, Fig S2)** and appear to be associated with the nuclear background of the host. The three heteroplasmic lines of the Orinocan semispecies, FG103, FG111 and O11 displayed different α2 levels (**Table 1**). However, the two lab lines of the Andean-Brazilian semispecies (MS, Yellow) showed levels of α2 that were similar to each other at 20%. Finally, the highest α2 levels (45-60%) were detected in the heteroplasmic lines of the Amazonian semispecies, FG16 and FG572 (**Table 1, Fig S2**). In addition, we found that levels of α2 were significantly higher in male flies compared to females in all tested lines except for FG111 ($p<0.001$ for FG572, MS, $p<0.01$ for Yellow and FG16, $p<0.05$ for O11, Tukey HSD test, **Fig S2; Supplemental data file**). However, since α2 levels were measured as a ratio between mitotypes, it is not clear if the higher level in males results from fluctuations in copy numbers of α2, of the primary mitotype (α1, β1 or β2), or of both.

Titers of α2 and β2 mitotypes in *D. paulistorum* tissues

To explore the dynamics and potential function of α2 and β2 mitochondria, we measured the genome copy numbers of each mitotype relative to nuclear genome copies (from here on referred to as titer) in the heteroplasmic O11 line of the Orinocan semispecies. Since mitochondrial copy numbers are variable in different tissues and under different conditions, their location and titer can provide clues as to their function. Hence, besides whole flies we also investigated α2 and β2 titers in the head, thorax, abdominal carcasses without gonads and gonads of male and female O11 flies using discriminant qPCR. We also explored the potential influence of sex, age, and infection with the endosymbiont *Wolbachia* on titers of both mitotypes. We found that titers of the primary β2 mitotype (**Fig 3A**) were significantly influenced by tissue ($F_{4,80}$=25.44, p<0.001). As expected, β2 is found in higher titers in energy-demanding tissues where ATP is provided by oxidative phosphorylation (OXPHOS) such as in the flight muscle of the thorax ([5], and references therein). Further, we found a weak significant interaction between age and sex ($F_{1,80}$=6.12, p<0.05) potentially driven by younger male flies having slightly more β2 mitochondria than older males, in combination with different tissue location patterns between the sexes. Females had overall lower numbers of the primary β2 mitotype per nuclear genome copy than males ($F_{1,80}$=28.68, p<0.001).

In contrast to β2, we found no significant effect of fly tissue ($F_{4,80}$=1.426, p=0.23) or age ($F_{1,80}$=0, p=0.99) on the titer of α2. Additionally, there was no significant effect of *Wolbachia* infection on the titer of either the β2 ($F_{1,80}$=1.41, p=0.24) or the α2 ($F_{1,80}$=0, p=0.97) mitotypes. The copy number of α2 is, however, significantly affected by fly sex (**Fig 3B**, $F_{1,80}$=51.34, p<0.001), with males carrying approximately twice as many α2 copies as females per nuclear genome. Significant higher titers of α2 in O11 males corroborate our finding of generally higher α2 levels in heteroplasmic males compared to heteroplasmic female flies (**Table 1; Fig S2** and **Supplemental Data file**), confirming that the significant differences between the sexes in the percentage data are due to more copies of α2, and not for example, lower β2 numbers.

The fact that only the titers of the β2 mitotype varied depending on fly tissue, but not the ones of α2, supports our conclusion that β2 is the functional and hence the primary mitochondrion and that α2 might be dysfunctional and/or not able to generate sufficient energy for its host.

Fig 3. Absolute copies of β2 and α2 mitochondria in O11 *D. paulistorum* flies, relative to fly nuclear genome copy. Bar charts present **(A)** β2 and **(B)** α2 mitochondria in male and female flies, dissected tissues and across two time points (1 and 3 weeks) as well as whether the flies had been treated with antibiotics leading to a knockdown of *Wolbachia*. Abbreviations: WT: wild-type flies. KD: *Wolbachia* knock-down flies. Three biological repeats are plotted with error bars of 1 standard error.

Biparental transmission and levels of the α2 mitotype in intra- and inter-semispecies hybrids of *D. paulistorum*

We used *CO1* RFLP-PCR to investigate if paternal transmission plays a role in the maintenance of high level heteroplasmy in *D. paulistorum*. Relative levels of α2 in F_1 offspring were measured in various crosses involving homoplasmic females and heteroplasmic males in intra- and intersemispecies combinations. Semispecies designation of the tester lines was determined by compatibility crosses (**Table S6**). By this, we uncovered unusual high levels of paternal transmission of α2. In intra-semispecies crosses, all tested F_1 flies in the FG295 (β2) x O11 (β2 and α2) cross (40/40, **Fig 4A**) and 57.5% flies in the A28 x FG572 cross (23/40, **Fig 4B**) were heteroplasmic (Supplemental data for Fig 4). F_1 female hybrids from FG295 x O11 cross had significantly lower α2 levels than O11 females (parental line) (*p*<0.001, Tukey HSD test), whereas the F_1 male hybrids from the same cross as well as both F_1 sexes from the A28 x FG572 cross did not significantly differ in levels from their parental ones of the same sex (*p*=0.99 for former and *p*=0.06 and *p*=0.46 for the latter two, respectively). As

heteroplasmy is not fixed in the FG572 line (**Table 1**) paternal introgression in A28 was also incomplete (**Fig 4B).**

In inter-semispecies crosses between Amazonian and Orinocan flies all tested F_1 flies of A28 (β1) x O11 (40/40, **Fig 4C**) and 45% flies of FG18 (β1) x O11 (18/40, **Fig 4D**) were heteroplasmic. In both combinations, the α2 mitotype levels of F_1 flies of both sexes were significantly lower than in the paternal O11 line (*p*<0.001 for both crosses, Tukey HSD test). Combining data from all four crosses (**Fig 4 A-D**), paternal transmission of α2 was found in 76% of F_1 hybrids (121/160). This finding clearly demonstrates that the α2 mitochondria of *D. paulistorum* are not only sporadically leaking but are highly transmissible via males. To independently confirm these results, we tested a subset of the heteroplasmic F_1 hybrids for their α2 levels using a second mitochondrial locus (*CO2*) and observed patterns similar to *CO1* (**Fig S3**).

Additionally, we investigated whether inheritance of paternal mitochondria is α2-specific. For this, we used mitotype-specific RFLP-PCR or Sanger sequencing on F_1 hybrid flies generated by eight different cross combinations aimed at tracing paternal transmission of either α1 (number of F_1 flies assayed = 59), β1 (31), or the β2 (76) mitochondria. Paternal transmission was not detected in any of the 166 tested F_1 flies (**Table S7**). Hence, a high level of paternal transmission is unique for α2 mitochondria.

We verified our findings using highly sensitive discriminant qPCR on A28 x O11 interspecies crosses. We assessed quantities of the β1, β2 and α2 mitotypes in O11 males (α2 and β2), A28 females (β1) and F_1 hybrid offspring. As expected, the pure parental lines possessed only the expected mitotypes (**Fig 4E, F**). Despite robust paternal inheritance of α2 in all tested F_1 hybrids (16/16), the hybrid offspring do not show any significant differences in titers for any mitotype when compared to their homoplasmic mothers, which is likely due to low α2 titers in the hybrids and relatively few biological replicates. However, significantly different titers for all mitotypes were seen when F_1 hybrids are compared to their heteroplasmic fathers **(Fig 4E, F, G)**, i.e. higher β1, lower β2 and lower α2 titers are seen in F_1 hybrids compared to the paternal line. In contrast to the less sensitive RFLP-PCR method, our discriminant qPCR method allowed us to detect very low levels (~0.1-0.3%) of paternal β2 leakage in 11 out of 16 hybrid offspring (Supplementary data file).

As expected for an inter-semispecies cross [44, 45], all F_1 hybrids were sterile and did not produce any live larvae when crossed either to siblings or backcrossed to parental lines (**Table S8**). This prevented exploration of the mitotype dynamics in further generations from this combination of parental lines. We were also unable to explore the dynamics of the reverse cross (female O11 with male A28), and thereby test if the β1 mitotype could be paternally transmitted, as none of the crosses we set up (0/200 attempts) produced larvae.

Fig. 4. Levels of the paternal $\alpha2$ mitotype in intra- and inter-semispecies hybrids of *D. paulistorum*. Paternal inheritance of $\alpha2$ mitochondria in different *D. paulistorum* hybrids determined by RFLP-PCR **(A-D)**. Scatter plots present relative $\alpha2$ levels in single fly F_1 hybrids derived from homoplasmic females (circle) and heteroplasmic males (square) of **(A, B)** intra-semispecies (FG295 x O11; A28 x FG572) and **(C, D)** inter-semispecies crosses (A28 x O11; FG18 x O11). Relative levels in F_1 hybrids (F_1 circle, F_1 square) were calculated based on *Hind*III-RFLP using *CO1* locus, and compared to the maternal (G_0m fem, G_0m mal) and paternal (G_0p fem, G_0p mal) lines. Single values are plotted including median (red crossbar) of all individuals and mean (red rhombus) of heteroplasmic flies (n=20 per sex, flies were 7-10 days old). Significance of difference between $\alpha2$ levels was estimated by ANOVA ($p<0.05*$, $p<0.001***$). Paternal $\alpha2$-inheritance determined by discriminant qPCR **(E-G)**. Number of copies of $\beta1$ **(E)**, $\beta2$ **(F)** and $\alpha2$ **(G)** mitochondrial genomes in 3-week old O11, A28 parents and F_1 hybrids relative to fly nuclear genome copy. Data was generated from crosses from single virgin males and single virgin female flies. N=8 in all cases and error bars are 1 standard error. Post-hoc significances are shown as *** = $p<0.001$, ** = $p<0.01$, * = $p<0.05$.

High invasiveness and stable persistence of the α2 mitochondria over multiple

generations

After identifying that α2 was the only mitotype paternally transmitted in high frequency to F_1 flies, we investigated persistence of heteroplasmy over further generations. We used discriminant qPCR to follow the inheritance patterns of the α2 mitotype during four generations of fertile intra-semispecies FG295 (homoplasmic β2) and O11 (heteroplasmic α2 and β2) hybrid crosses. We followed four single-fly G_0 crosses in each direction (FG295 x O11 and O11 x FG295), and mated a randomly picked pair of virgin siblings to each other in each consecutive generation (F_1-F_4) to explore single-fly inheritance patterns of the α2 mitotype (**Fig 5**). The results showed both maternal (**Fig 5A**) and paternal (**Fig 5B**) transmission of the α2 mitotype, as well as high α2 invasiveness (ability to invade a homoplasmic line) and persistence (ability to be maintained over generations) in F_4 flies (13/16).

In both crossing directions, all F_1 hybrids (16/16) were heteroplasmic, hence demonstrating not only paternal but biparental transmission of α2. However, in the following generations (F_2-F_4), transmission success was more varied. All tested offspring from three of the four crosses starting with a heteroplasmic mother carried both α2 and β2, and one cross lost the heteroplasmy at F_3 (**Fig 5A**), whereas the crosses starting with a heteroplasmic father showed more stochasticity in transmission, with the α2 mitotype not necessarily transferred to both tested offspring in these later generations (**Fig 5B**). However, more individuals need to be tested in order to confirm this pattern.

In agreement with earlier results, we found that there were differences in mitochondrial titers between heteroplasmic and homoplasmic flies, which are dependent on sex. While titers of the β2 mitotype were not significantly different between hybrid males and females across the experiment (Wilcoxon rank sum test W=866, p=0.29), male hybrids had higher titers of the α2 mitotype compared to hybrid females (Wilcoxon rank sum test W=1208, p<0.001). Overall, heteroplasmic flies also carried a higher total mitochondrial load than homoplasmic flies (Wilcoxon rank sum test W=183, p<0.001). After separating the data by the direction of the cross and the sex of the fly, we found that heteroplasmic sons had significantly higher total mitochondrial titers than their homoplasmic parents (Wilcoxon rank sum test W=5, p<0.05 for female parents; W=2, p<0.01 for male parents), while heteroplasmic daughters do not (Wilcoxon rank sum test W=10, p=0.138; W=12, p=0.13 for male parents), demonstrating the noticeably higher titers in male heteroplasmic flies specifically. In contrast, heteroplasmic fathers have significantly higher total mitochondrial titers than their heteroplasmic sons and daughters (sons W=47, p<0.05; daughters W=43, p<0.05), whereas the difference in titer between heteroplasmic mothers and their heteroplasmic offspring is non-significant for either sex (sons W=19, p=0.382; daughters W= 38, p=0.202).

Given that we only observe significant variability in the α2 titer between sexes as well

as between parents and their offspring (no significant difference was observed for any test on

β2), the higher titer of mitochondria in heteroplasmic flies overall suggest that the presence of

α2 increases the number of total mitochondria per cell and is added on top of the primary

mitochondria, i.e., β2. The reduction in total mitochondrial titer between heteroplasmic fathers

of the O11 line and their heteroplasmic offspring of both sexes cannot be explained by the

significantly different α2 titers observed between male and female flies. Instead, this result

indicates an interaction between mitotype titer and host nuclear background, which is more

pronounced in males.

Fig 5. **Presence and titer of α2 and β2 in (A) O11 x FG295 and (B) FG295 x O11 hybrids.**
Relative copies of α2 to number of fly nuclear copies is shown in green. Relative copies of β2
to number of fly nuclear copies is shown in blue. Values were rounded up if the decimal was
over 0.5 and rounded down if under 0.5. The figure depicts four biological repeats of each
direction of the hybrid cross, each representing a single-pair mating of a male and female fly
for each line. A mated pair derived from randomly picked virgin flies of each generation was
tested for four generations after the initial mating and copy numbers of α2 and β2 mitochondria
were quantified through discriminant qPCR. ∗ means that α2 was below the cutoff threshold
on qPCR and the fly is considered homoplasmic.

To further investigate whether fly nuclear background affects the persistence of the α2

mitotype we backcrossed virgin F_1 hybrids from O11 x FG295 to the FG295 line (**Fig 6A**),

thereby increasing the nuclear background that coevolved with the homoplasmic β2 mitotype

in each generation.   As before, all F_1 flies acquired the α2 mitotype when transmitted

maternally (**Fig 6A**, left column), but their respective levels were lower than in the G_0 parent

flies ($p<0.001$, Tukey HSD test). Backcrossing these F_1 flies to the homoplasmic FG295 line

(BC1, maternal introgression) showed a decrease in α2 prevalence in the offspring (**Table S9**).

As in F_1 flies, the levels of α2 in offspring from the hybrid parents were significantly lower than

in G_0 flies ($p<0.001$, Tukey HSD test). A second backcross of the BC1 offspring to the
homoplasmic FG295 line revealed that $\alpha2$ prevalence decreased in the BC2 offspring from
the BC1 hybrid mothers, whereas their paternal counterparts kept a similar prevalence as that
observed in BC1. Overall $\alpha2$ levels in BC2 flies were significantly lower in both sexes in all
backcrossing directions ($p<0.001$, Tukey HSD test) compared to their parental levels (G_0
females - Mean=44.5, SD=4.5; G_0 males - Mean=52.4, SD=4.8).

Again, all F_1 flies also inherited the $\alpha2$ mitotype when initially transmitted paternally
(**Fig 6A**, right column). Compared to G_0 flies of the same sex, the levels of $\alpha2$ were significantly
lower in F_1 females but not significantly different in F_1 males ($p<0.01$ and $p=0.56$, respectively,
Tukey HSD test). Backcrossing of these F_1 flies to the FG295 line (BC1, paternal introgression)
showed a decrease in $\alpha2$ prevalence in the offspring when transmitted both maternally and
paternally (**Table S9**). Compared to the F_1 flies of the same sex, the levels of $\alpha2$ in all BC
offspring from both hybrid parents decreased to approximately 20% in females and 30% in
males ($p<0.001$, Tukey HSD test).

A second backcross of the BC1 offspring to the FG295 line revealed a pattern similar
to the one of the reciprocal O11 x FG295 cross. The prevalence of $\alpha2$ was low in the
backcrossed offspring derived from hybrid mothers whereas the flies from hybrid fathers had
a prevalence similar to that observed in BC1. The only exception was the offspring obtained
from mothers derived from the F_1 hybrid fathers (40% for females and 40% for males). By
increasing the FG295 nuclear background, all BC2 offspring showed significantly lower $\alpha2$
levels ($p<0.001$, Tukey HSD test; see **Table S9**) compared to the levels in their heteroplasmic
O11 parents of the same sex. In summary, we observed that backcrossing heteroplasmic F_1
hybrid flies from both crossing directions to a homoplasmic $\beta2$ line not only lowers relative $\alpha2$
levels but also results in a decrease in heteroplasmy prevalence in further generations. These
data indicate that the $\alpha2$ mitotype can successfully invade previously homoplasmic lines,
although its transmission success is affected by the host nuclear background.

Additionally, we investigated the capacity of $\alpha2$ to invade other species by performing
crosses between females of *D. equinoxialis* and *D. willistoni* and males of *D. paulistorum* O11
line (**Fig 6B, 6C**). Since hybrid flies from these interspecies crosses are fully sterile, we could
not assess $\alpha2$ prevalence, persistence and levels beyond F_1 flies. Both cross combinations
showed 100% prevalence of the $\alpha2$ mitotype in F_1 offspring, although due to the high
embryonic lethality of hybrids we were only able to analyze eight F_1 flies in the *D. equinoxialis*
x O11 cross (**Fig 6B**) and two F_1 flies in the *D. willistoni* x O11 cross (**Fig 6C**). In agreement
with earlier experiments [58], no F_1 males were obtained in either interspecies crosses, thus,
only females were analyzed for mitochondria levels. Similar to results derived from *D.*
paulistorum inter-semispecies crosses (**Fig 4**) F_1 females of *D. equinoxialis* x O11 cross (**Fig**
6B) had significantly lower $\alpha2$ levels in comparison to G_0 flies (11%, $p<0.001$, Tukey HSD

test). α2 levels in the two F$_1$ females of *D. willistoni* x O11 cross were also decreased, but due to the extremely low number of viable F$_1$ hybrid flies (**Fig 6C**) we were not able to analyze significance of this result.

Figure 6. Prevalence and levels of the α2 mitotype introgressed in different genetic backgrounds. (A) Dynamics of α2 prevalence (blue) and levels (green) determined by α-diagnostic *CO1*/*Hin*dIII RFLP in females and males by backcrossing F$_1$ flies derived from O11 x FG295 (left: maternal introgression; white x black) and FG295 x O11 (right: paternal introgression; black x white) parents to FG295 flies. Arrows in this figure represent a backcross to homoplasmic (β2) FG295 flies. **(B, C)** Dynamics of α2 prevalence (blue) and levels (green) determined by the α2-diagnostic CO2/*Msp*I RFLP in F$_1$ flies from interspecies crosses of **(B)** *D. equinoxialis* (orange) and **(C)** *D. willistoni* (red) females to heteroplasmic *D. paulistorum* O11 males (white). The skull symbolizes complete absence of F$_1$ males in both crosses. Lines inside the green bars show standard deviation. Values below sex symbols indicate the number of single flies used in RFLP-PCR analysis and listed in **Table S9**.

α2 dynamics during embryogenesis in heteroplasmic lines and F$_1$ hybrids uncover their biparental transmission mode

As demonstrated by our crossing experiments, the biparentally inherited α2 mitotype of *D. paulistorum* is highly invasive and maintained at consistently high levels in the adult fly. Since the number of mitochondrial DNA copies per sperm cell is much smaller than in a mature egg [59], and paternally leaked mitochondria are usually both passively diluted and actively destroyed [15], we asked how paternally inherited α2 mitochondria avoid depletion. By using RFLP-PCR, we analyzed α2 levels in unfertilized eggs and embryos from the O11, FG572 and MS lines (**Fig 7A-C**) as well as F$_1$ embryos from reciprocal crosses between O11 and FG295 flies (**Fig 7D, E**). The α2 levels at different stages of embryogenesis were compared to those in unfertilized eggs (E0) and parental flies (G$_0$f - females, G$_0$m - males). Strikingly, none of the five lines analyzed showed traces of α2 in unfertilized eggs. The finding is not

surprising for the F_1 flies from FG295 x O11 cross (**Fig 7E**), where the eggs come from

homoplasmic FG295 mothers with only the β2 mitotype, but is unexpected for mothers of the

four heteroplasmic lines (**Fig 7A-D**) that carry high α2 levels (**Table 1**). We infer that α2 levels

are very low in unfertilized eggs of heteroplasmic mothers and below the detection level of our

RFLP-PCR method. Immediately after fertilization (E1; 0-30 minutes after egg laying - AEL),

however, α2 reached relatively high levels (similar or close to parental) in F_1 embryos of all of

the lines except for the ones obtained from the maternal O11 x FG295 cross (**Fig 7D**). For

O11 and FG572 E1 embryos (**Fig 7A, B**) median α2 levels were not significantly different from

the levels of either parent ($p>0.05$; here and further on ANOVA with post-hoc Tukey HSD test),

except for FG572 males ($p<0.01$; Supplemental data of **Fig 7**). Median α2 levels of E1

embryos from MS, FG295 x O11 and O11 x FG295 (**Fig 7C-E**) were significantly lower than

in parents ($p<0.05$). In mid-embryogenesis (E2; 3-4 hours AEL) α2 levels of O11 were only

significantly decreased compared to E1 ($p<0.001$; **Fig 7A**), whereas in the four other lines (**Fig**

7B-E) α2 levels continued to increase (statistically non-significant in FG572, MS embryos and

F_1 of O11 x FG295, $p>0.05$; but significant in F_1 of FG295 x O11 crosses, $p<0.001$). In late

embryogenesis (E3; 18-20 hours AEL) all lines showed a significant increase in α2 levels

compared to E2 ($p<0.001$) except for FG572 and F_1 embryos from the maternal O11 x FG295

cross shown in **Fig 7B** and **D** ($p>0.05$). Only in F_1 embryos from the maternal O11 x FG295

cross α2 levels at E3 were significantly lower in comparison to parents (9.7%; $p<0.001$). In all

other cases the level of α2 in E3 embryos reached similar or even significantly higher levels

compared to their respective heteroplasmic parents ($p<0.001$ for G_0f in O11 and MS as well

as G_0m in MS; $p<0.05$ for FG295 x O11; $p>0.05$ for G_0f in FG572, FG295 x O11 and G_0m in

O11 and FG572).

Additionally, we compared α2 levels of embryos to the levels in emerged flies (F_1f and

F_1m, **Fig 7**). There were no significant differences in α2 levels between G_0 and F_1 flies of the

same sex except for F_1 flies from the intra-semispecies crosses O11 x FG295 and FG295 x

O11 ($p<0.01$). In comparison between F_1 flies and embryos of heteroplasmic lines we found

similar statistically significant differences as between embryos and G_0 flies (see Supplemental

data of **Fig 7**). F_1 flies from both O11 x FG295 and FG295 x O11 crosses (**Fig 7D, E**) had

significantly higher α2 levels than at any embryonic stages ($p<0.05$), except in late embryos

(E3) from the paternal FG295 x O11 cross with no significant difference to F_1 males ($p=0.99$),

but significantly higher α2 levels than F_1 females ($p=0.002$). Strikingly, a comparison of the

maternal (**Fig 7D**) and paternal (**Fig 7E**) α2 dynamics between embryonic stages (E1-E3) and

F_1 adult flies (**Fig 7D, E**) uncovered significant differences in α2 levels depending on the mode

of transmission ($p<0.001$ for all embryonic stages and F_1 flies). Hence during *D. paulistorum*

embryogenesis, paternally introgressed α2 mitochondria are replicating faster and possibly

earlier than α2 that were transmitted by the female.

Fig 7. Relative α2 levels during *D. paulistorum* embryogenesis determined by *CO1*/*Hind*III (A, B, D, E) and CO2/*Msp*I (C) RFLP-PCR. Scatter dot plots combined with box plots show levels of α2 (%) in the heteroplasmic lines O11 **(A)**, FG572 **(B)**, and MS **(C)** and α2 levels in F_1 hybrids from maternal **(D)** and paternal **(E)** introgressions obtained from reciprocal crosses between parental heteroplasmic O11 and homoplasmic FG295 flies. Dynamics of α2 levels are presented in adult parental females (G_0f) and males (G_0m), unfertilized eggs (E0), embryos of 0-30 min (E1), 3-4 hrs (E2), and 18-20 hrs (E3) of development at 25°C and F_1 adult females (F_1f) and males (F_1m). Each dot represents the α2 level in percent determined from a pool of 20 staged embryos collected for each sample. Females are shown as circles, males as squares and embryos as triangles. Mean values for every stage and sex are depicted with red rhombus.

Collectively, these data strongly suggest that α2 arrives in very low copy numbers from both sexes to the mature egg and, in order to avoid out-dilution by the primary mitotype, starts replicating immediately after fertilization. This strategy works regardless of its transmission mode and thereby ensures that α2 already reaches high levels in mature embryos (E3), equivalent to levels of the adult fly. As the strict maternally-inherited primary mitochondria of *D. melanogaster* and other animals are quiescent and do not replicate during early embryogenesis [13, 60], this early replication behavior is unique for α2 and we hence name it "Replication precox" (see Discussion).

Genomic features unique to the α2 mitotype

The unusual properties of the α2 mitochondria prompted us to search for unique genomic

features that distinguish this secondary mitotype from the other *D. paulistorum* mitochondria

(**Table 2, 3**). However, only three non-synonymous substitutions were found to be exclusive

to the α2 mitotype (**Table 2**).

Table 2. Mutations unique to the α2 clade among protein-coding genes of the sampled

***D. paulistorum* lines**. Three non-synonymous substitutions are unique to α2. Among these,

only the H -> N substitution in COIII results in a change in amino acid property. Amino acid

and nucleotide changes are presented from other mitotypes to α2. Nr = number.

Gene	Nr non-synonymous	Nr synonymous	Amino-acid change	Chemical properties
ATP6	0	3		
ATP8	0	1		
COI	0	5		
COII	0	1		
COIII	1	0	H -> N	Positive charge to neutral
CYTB	0	2		
ND1	0	2		
NADH2	0	3		
ND3	1	0	V -> M	No change
ND4	1	3	V -> I	No change
ND4L	0	0		
ND5	0	4		
ND6	0	4		

One of the substitutions is found in *ND3* and causes a replacement of Valine with Methionine

(aa position 18 in *D. paulistorum*, 21 in humans) and another substitution in *ND4* causes a

replacement of Valine with Isoleucine (aa position 379 in *D. paulistorum*, 391 in humans). Both

of these replacements are between chemically equivalent amino acids and likely have little

functional impact on the proteins. However, the substitution in *CO3* causes the replacement

of a positively charged Histidine by a neutral Asparagine in α2 (aa position 7 in *D. paulistorum*,

6 in humans) and is predicted to have medium functional impact according to the

MutationAssessor server (http://mutationassessor.org/r3/). An inspection of multi-species

alignments in Uniprot and Pfam revealed that Asparagine is rarely found at this position of the

CO3 protein in other animals. The only organisms in the database with Asparagine in the

same position are a few species of mites, representatives of some invertebrate marine phyla,

and a wide variety of mollusks. Interestingly, among the mollusk sequences, many are from

the F mitotype of heteroplasmic unionid mussels with doubly uniparental inheritance (DUI).

Additionally, the selfish α2 mitotype differs from the primary mitotypes of *D. paulistorum* by

four A to G transitions in the large rRNA gene and a T to A transversion in tRNA-Ala and a 3

bp insertion in tRNA-Leu2 (**Table 3**), both of which are located in the T-loop.

Table 3. Mutations unique to the α2 subclade among non-protein-coding genes in the sampled *D. paulistorum* lines. Substitutions are found in the large rRNA and tRNA-Ala while an insertion occurs in the tRNA-Leu2. In both tRNAs, the mutations are located in the T-loop.

Gene	Number of mutations	Nucleotide changes
lrRNA	4	A -> G (4x)
tRNA-Ala	1	T -> A
tRNA-Leu2	1	- (β) / AT (α1) -> ATATT

Since the nuclear-encoded protein Elongation factor Tu (EFTu) is known to interact specifically with the tRNA T-loop during translation [61, 62], we took a closer look at this protein in *D. paulistorum*. Two divergent copies of EFTu are encoded in the nuclear genome of *D. melanogaster*, *mEFTu-1* and *mEFTu-2*. The two genes have largely sex-specific expression in adults [63] and the proteins differ somewhat in binding specificities to tRNAs with and without T- and D-arms [64]. Using our previously assembled transcriptomes of A28, C2 and O11 [50] we identified orthologs of the two *D. melanogaster mEFTu* genes in *D. paulistorum*. An alignment of the two proteins together with orthologous proteins from all *Drosophila* species found in Flybase (release 2019_04) showed that mEFTu-2, the gene with testis-specific expression, has unique amino acids in two positions in *D. paulistorum*. Interestingly, both positions are close to the C-terminal end of the protein, which is the region that interacts with the T-stem of the tRNA [64], suggesting a potential mito-nuclear coadaptation. However, since only one of the three sequenced *D. paulistorum* lines currently carries the α2 mitotype, and we have no data to say whether the other two, A28 and C2, can carry or have ever carried the α2 mitotype, no firm conclusions can be drawn.

The evolutionary history of mitochondria in *D. paulistorum* and the *willistoni* subgroup

Well-supported phylogenies can provide useful insights into the evolutionary history of genetic elements. Although our whole-genome mitochondrial phylogeny showed very high support in almost all internal nodes, the node defining the relationship between α, β and the clade containing *D. willistoni*, *D. insularis* and *D. tropicalis* (hereafter referred to as the willistoni clade) has low support (**Fig 2B**). Hence, in order to confirm the topology of our mitochondrial phylogeny, we investigated the cause of the poorly supported node by testing whether recombination between clades or bias in nucleotide composition could be affecting the phylogenetic signal.

Nucleotide composition creates conflicting phylogenetic signals between the α, β and the willistoni mitochondrial clades

We used several methods to search for signs of recombination between the clades. First, we created single gene trees for the 13 mitochondrial protein coding genes and tested for conflicts

between the resulting topologies. For eleven of the 13 trees a definite topology could not be
determined as the bootstrap support was less than 80% on the node defining the relationships
of the α, β and willistoni clades. The remaining two trees had different topologies, as the *ND2*
gene featured the α clade as sister either to the willistoni clade (BS = 81%) and the *ND5* gene
placed α as sister to the β clade (*ND5* gene, BS = 90%). Second, we ran tests for intragenic
recombination on all protein coding genes. Only two genes, *CO2* and *ND3*, were detected with
marginally significant signals ($p<0.05$) using Phipack [65]. Finally, we estimated the
recombination over mutation rate along the whole mitochondrial genome to be low (0.01).
These results indicate that the recombination rate is low and that recombination signals are
weak and inconsistent. Thus, even if some recombination occurred between mitochondria, it
is unlikely to be the cause of the low support on the internal node of the whole-genome
mitochondrial tree.

Nucleotide composition bias and codon usage can also influence phylogenies and are
particularly likely to affect the more neutral third codon position. Hence, we built separate trees
using either only the third or a combination of the first and second codon positions of the
protein coding genes. Interestingly, we observed significantly increased support for α as sister
to the willistoni clade when the third codon position was removed (**Fig 2B**). In contrast, the
tree based only on the third codon position showed high (Bayesian posterior probability 0.99)
or fairly good (ML bootstrap 88%) support for the α and β clades being sisters (**Fig S4**). The
conflicting phylogenetic signal between the two first codon positions and the third codon
position suggests that nucleotide composition bias and perhaps codon usage are indeed the
cause of low support on the node defining the relationship between the α, β and willistoni
clades in the whole genome mitochondrial tree.

Since the third codon position is more likely to be affected by such biases, we conclude
that the topology supported by the first and second codon positions of protein coding genes
(Fig 2B) is most likely the accurate representation of the evolutionary relationships between
mitochondrial clades. Thus, *D. paulistorum* mitochondria are polyphyletic, and α is sister to
the willistoni rather than to the β clade. The resulting phylogeny **(Fig 2B)** implies an α-like
mitochondrial ancestor in *D. paulistorum,* and suggests that β was most likely acquired from
a so far unidentified lineage within the *willistoni* subgroup.

Mitochondrial and nuclear phylogenies show a complex evolutionary history of mitochondria
in the willistoni subgroup

A comparison between nuclear and mitochondrial phylogenies shows that neither mitotype
relationship nor presence of heteroplasmy follow the topology of the nuclear phylogeny.
Mitotypes are shared by relatively distantly related lines belonging to different semispecies,
as in the case of α1 in C2 (CA) and MS (AB) and β2 in O11 (OR), L1 (IN) and SM (TR),

whereas the primary β1 mitotype is exclusively found in AM semispecies (**Fig 2B**).
Heteroplasmic lines are also spread across the nuclear phylogeny, and the mitotype that
accompanies α2 in heteroplasmy varies according to what semispecies the line belongs. This
pattern suggests that the mitochondria of *D. paulistorum* were not only vertically transmitted
but have also been introgressed and lost multiple times during evolution. Further introgression
events are indicated by the positioning of *D. paulistorum* and *D. equinoxialis* in relation to other
species of the willistoni subgroup on either tree. In the nuclear phylogeny, *D. paulistorum* and
D. equinoxialis are the most derived species in the willistoni group, but the mitochondrial
phylogeny places the α and β clades in a basal position with regards to the willistoni clade
(**Fig 2A, B)**.

NUMTs provide insights into the evolutionary history of mitochondria in the willistoni subgroup
Insertions of mitochondrial DNA in the nuclear genome (NUMTs) can provide valuable
information about mitochondrial evolution. Using Illumina reads we generated whole genome
assemblies for all willistoni subgroup lines (**Fig 2A**) and searched for contigs containing
sequences with at least 90% identity to our mitochondrial genomes. Potential NUMTs were
identified based on (*i*) coverage similar to that of nuclear genes, (*ii*) low identity towards the α
and β genomes, and (*iii*) presence of mixed nuclear-mitochondrial sequences (see Methods).
Sequences classified as NUMTs were found in assemblies of all fly lines (**Tables S3, S8**).
Interestingly, eleven of the 13 *D. paulistorum* lines, including all lines carrying only β
mitochondria, shared a putative NUMT that spans a genomic region containing the *ND2* and
CO1 genes. Contigs containing this sequence had similar coverage to nuclear genes in all
lines except FG295 and L1, where the NUMT had lower coverage than the nuclear average,
and FG572, in which the NUMT coverage was slightly higher than the nuclear average. In the
L1 and SM lines, the contigs also included non-mitochondrial sequences which confirm their
nuclear location (**Table S3**, **S10**). A phylogenetic analysis of this genomic region suggested
that this putative NUMT forms a clade together with the α mitotype and the mitochondria of
other willistoni subgroup species (**Fig 8**).
699

Fig 8. Phylogeny of the mitochondrial region spanning the _ND2-CO1_ genes in the willistoni subgroup. NUMT sequences recovered from genomic assemblies of _D. paulistorum_ lines branch together with the α and the willistoni clades. Fly lines in bold are heteroplasmic. The same topologies are obtained by maximum likelihood (GTRGAMMA model) and Bayesian analyses. Colored circles next to the nodes indicate ML bootstrap support (left half) and Bayesian posterior probability (right half). Black corresponds to 99-100% bootstrap support or 0.99-1.0 posterior probability, grey represents 70-98% bootstrap support or 0.7-0.98 posterior probability, white indicates support below 70%. Branches leading to the outgroups and marked by two transverse lines were shortened and don't follow the presented scale.

The presence of the sequence in nearly all _D. paulistorum_ lines together with the nuclear-like

topology suggest that this NUMT stems from a single nuclear insertion that occurred prior to

the diversification of _D. paulistorum_ semispecies. The ancient origin together with the

phylogenetic position of this NUMT close to α and the willistoni clades further support the

hypothesis of an α-like ancestral mitotype of _D. paulistorum_.

Finally, we observed that the genome assembly of _D. equinoxialis_ contains several

β−related contigs, which were not found in species of the willistoni clade (**Tables S3, S8**). The

coverage of these contigs is lower than the nuclear average for the line and they do not contain

matches to non-mitochondrial sequences. Hence, while these _D. equinoxialis_ sequences

might stem from NUMTs, we cannot fully exclude that they represent low-level contamination.

Nevertheless, it is interesting to note that the presence of β sequences in _D. equinoxialis_ might

suggest that the last common ancestor of *D. paulistorum* and *D. equinoxialis* carried a β-like mitochondrion.

DISCUSSION

We report on a unique system of stable mitochondrial heteroplasmy in the neotropical *D. paulistorum* species complex that might exist due to biparental transmission. Using sequencing, RFLP-PCR and bioinformatic analyses we discovered high-level heteroplasmy in 41% of the tested fly lines including representatives from three of the six *Drosophila paulistorum* semispecies (Andean-Brazilian, Amazonian and Orinocan). However, given the limited number of lines currently available from the Centro-American, Interior and Transitional semispecies, it is possible that heteroplasmy may be even more broadly distributed among the species. We also found that heteroplasmy is fixed with balanced levels of each mitotype in individuals of three long-term laboratory lines (O11, MS and Yellow), but not in the more recently collected lines, in which α2 levels are also tightly balanced but not fixed (**Table 1**). These results suggest that a line has to be genetically homogeneous for heteroplasmy to be fixed and that levels are controlled by the nuclear genetic background. In the course of our current study we observed that the Orinocan FG103 line collected in 2014 lost the secondary mitotype within the first two years in the lab, suggesting that long-term maintenance of fly lines under lab conditions might also contribute to an underestimation of prevalence. Further sampling will be necessary to determine the prevalence of heteroplasmy in natural populations.

Function of the α2 mitotype in *D. paulistorum*

Using phylogenetic analysis, we found that the mitochondria of *D. paulistorum* are polyphyletic and form two phylogenetically distinct clades - α and β - each further divided into two subclades, 1 and 2. Heteroplasmy is always associated with the α2 mitotype, and this mitotype is in turn never observed in homoplasmy. Such a pattern may occur because α2 alone does not perform all of the functions required for sustaining a fly, or perhaps because homoplasmic α2 lines are rare and difficult to sample in the wild, or both. Although systematic field sampling would be necessary to determine which hypothesis is correct, we found additional evidence suggesting that α2 might not provide full mitochondrial functionality. Mitochondria are usually enriched in energy-demanding tissues [66, 67], and by studying the heteroplasmic O11 line we found that only the β2 mitotype follows this pattern. The α2 mitotype, on the other hand, has unchanging titers across fly tissues. This suggests that β2 is the mitotype that provides energy via oxidative phosphorylation, while α2 does not respond to the energy demands of the host. Despite this, the genes encoded on the α2 genome appear to be evolving under purifying selection, as indicated by low dN/dS ratios and the absence of pseudogenes. In

addition, very few genomic differences are seen between α2 and the closely related α1

mitotype (0,67% divergence), which since it is the sole mitotype found in two Centro-American

D. paulistorum lines (C1 and Yellow) must be fully functional. The low sequence divergence

between α1 and α2 strongly suggests that the transition from a primary α-like ancestor to a

selfish mitotype took place relatively recently. Considering these facts, we surmise that α2 is

likely functional based on its genome but that its steady titer throughout tissues and absence

in homoplasmy suggest that it does not provide sufficient energy to sustain a fly, at least under

the standard laboratory conditions of our experiments.

Biparental inheritance of the α2 mitotype in *D. paulistorum*

The question remains as to how and why α2 is maintained in *D. paulistorum* flies. It may serve

a yet undefined essential or condition-specific function for the fly, or alternatively it may be

cheating the system and perpetuate selfishly while providing no or few benefits to the host.

One possible way of accomplishing the latter is for the mitotype to increase its own vertical

transmission, for example through biparental inheritance.

Using RFLP-PCR and discriminant qPCR, we detected high level paternal

transmission of the α2 mitotype in offspring from both intra- and inter-semispecies crosses of

D. paulistorum (**Fig 4, 5 and 6A, B**), and even in offspring from crosses between *D.*

paulistorum and two closely related species of the willistoni subgroup (**Fig 6C**). None of the

other mitotypes found in *D. paulistorum* were paternally transmitted in high levels. Generally,

active pre- and post-fertilization recognition and elimination mechanisms prevent paternal

transmission of mitochondria in animals (recently reviewed in [15]). In *D. melanogaster,* the

mechanism is not yet fully understood, but two genes, *EndoG* and *tamas,* are known to be

involved in eliminating mitochondrial DNA from developing spermatids [13, 14, 68]. After

fertilization, paternally leaked mitochondria are eliminated by endolytic and autophagic

degradation systems [69] while the rare paternal mitochondria that escape are passively

outcompeted by the larger number of maternally transmitted mitochondria already present in

the egg [70]. In order for a mitochondrion to be transmitted paternally and – even more

importantly – to stably persist in adult flies in large numbers, at least one of these mechanisms

need to be bypassed. We did not specifically determine if the α2 mitotype evades the active

mechanisms that eliminate paternal mitochondrial DNA (reviewed in [15]), but the observation

that none of the other mitotypes of *D. paulistorum* were paternally transmitted in large numbers

(**Table S7**) implies that there is no general loss of such active defense functions in *D.*

paulistorum. The spurious low level (~0.1-0.3%) paternal leakage of β2 in some hybrid crosses

could be a result of a partial breakdown of such mechanisms due to the mixed nuclear hybrid

background ([23] and references therein). However, low levels of paternally transmitted β2 is

also in line with previous reports of paternal leakage within *Drosophila* species [26, 29, 30],

indicating that similar levels of leakage can occur even without interspecies hybridization and breakdown of defenses. Thus, the mechanism for paternal transmission is specific to the invasive α2 mitotype.

In *D. melanogaster* and *D. simulans,* the spermatozoon is estimated to carry approximately 20 copies of mitochondrial DNA [59], which is five to six orders of magnitude less than the number of copies found in an egg [59]. We expected that similarly small numbers of the α2 mitochondrion are present in *D. paulistorum* sperm and at the onset of embryo development when α2 is transmitted by the male. However, using relative RFLP-PCR, we found that rapid and premature replication in early embryos resulted in α2 levels that were similar to those of adult flies just a few hours after fecundation (**Fig 7**). Previous studies of *D. melanogaster* showed that maternally transmitted mitochondria are quiescent and do not replicate for up to 10 hours after fertilization [13, 60], which seems to be an evolutionary conserved state (reviewed in [71]). Hence, the premature and unique temporal replication of the α2 mitotype, that we name "Replication precox", might allow this mitotype to escape passive dilution by the still dormant primary mitotype and thereby guarantee its presence in high levels in adult flies. We also found that the α2 mitotype level increased earlier and faster in hybrid embryos when it was transmitted paternally rather than maternally (**Fig 7D, E**). This could be due to more α2 mitochondria being deposited by the sperm than by the egg, or to a higher rate or earlier onset of α2 replication when it is paternally inherited. Since we found that O11 males have significantly higher levels of α2 than females, our data suggest that the former hypothesis is possibly more likely. One tempting mechanistic hypothesis for the Replication precox phenotype is that the nuclear encoded mitochondrial replication machinery is preferentially and prematurely recruited to α2 in the early embryo. The elucidation of the mechanism for diagnostic recognition and selective replication of α2 is currently underway in our laboratory.

Variability in mitochondrial titer and levels between sexes

We found that α2 titers were consistently and significantly higher in males than in females. Since α2 does not seem to be a fully functional mitotype, the higher α2 titer in males cannot be interpreted purely based on different energy needs between sexes. So why would males have a higher titer of α2 mitochondria than females? One possibility is that high titer in males has evolved specifically to facilitate paternal transmission of the invasive α2 mitotype, since a higher titer increases the chance of transmission. Alternatively, higher male titer and the Replication precox phenotype of α2 could have co-evolved in order to compensate for a reduced functionality of the maternally transmitted primary mitotype in males. The mother's curse hypothesis suggests that DNA of strictly maternally inherited mitochondria co-evolve

with females and might therefore accumulate mutations that are harmful to male functions
[72]. If such mutations have accumulated in the maternally transmitted primary mitochondria
of *D. paulistorum*, the biparentally transmitted α2 mitotype might have co-evolved with males
and express a still undetermined function that is more important to males than females. In the
O11 line, higher relative titer of α2 compared to the primary mitotype β2 are found in heads
and gonads (**Fig 3B**), suggesting that perhaps behavior and fertility might be important further
areas of investigation for α2 specific functions in males.

In addition, our data suggest that the total mitochondrial titer is significantly affected by
the presence of α2, as this secondary mitotype appears in addition to the primary
mitochondrial copies in a cell. A similar quantitative effect on mitochondrial copy numbers was
previously observed in heteroplasmic *D. subobscura* flies carrying a selfish mitotype with a
large deletion that results in a replicative advantage [73].

Our crossing data also suggests that the nuclear genetic background influences the
total mitochondrial titer, particularly in males. This is indicated by a significant reduction in
overall mitochondrial titer between pure O11 heteroplasmic males and heteroplasmic hybrid
offspring of both intra- and interspecific crosses, but not between pure O11 heteroplasmic
females and hybrid offspring. This observation is in line with long-term co-evolution between
the heteroplasmic cytoplasm and the nuclear genome in males. In this context it is interesting
to note that two of the genomic differences that we found in the α2 mitochrial genome are
located in the T-loops of tRNAs and the nuclear encoded gene *mEFTU-2* that is specifically
expressed in testis of *D. melanogaster* has mutations in a region where the protein is predicted
to interacts with the tRNA T-loop [63, 64]. Hence, if assuming a similar male-specific
expression of *mEFTU-2* in *D. paulistorum*, this observation might strengthen the idea of a co-
adaptation between the α2 mitotype and male functions.

Evolutionary history of the α and β mitotypes in the neotropical willistoni group

The ancestrality of α-like mitochondria in *D. paulistorum* is strongly supported by our
phylogenetic analysis after we accounted for the nucleotide composition bias in our
mitochondrial phylogeny, since the α clade is clearly sister to the other willistoni group species.
Furthermore, the presence of an α-related NUMT, which is shared by most sequenced *D.*
paulistorum lines (**Table S10**) and shows a topology that is similar to our nuclear tree (**Fig 8**),
indicates that this sequence was transferred from an α-like ancestral mitochondrion to the
nuclear genome before semispecies diversification.

Since the β mitotype does not share an immediate common ancestor with α or any
other lineage in our phylogeny, its presence in *D. paulistorum* is likely the result of a "ghost"
introgression from an unsampled or possibly even extinct species [55-57]. The phylogenetic
position of β suggests an origin within the willistoni group but outside the willistoni subgroup.

Thus, possible donors include species from the other two subgroups within the willistoni group, the

bocainensis and the alagitans. Unfortunately, sequence data is not available for all species in

the *bocainensis* subgroup, but *D. nebulosa,* which is included in our phylogeny, branches

basally to β. A similar position was observed for three other representatives of the bocainensis

subgroup, *D. capricorni, D. fumipennis* and *D. sucinea* in previously published single gene

trees [54]. No sequence data is available for any alagitans species. Alternatively, the β

mitotype may be the genomic remnant of an extinct species of the willistoni group which once

hybridized with *D. paulistorum*. Cases of introgressions followed by extinction of the donor

species have been reported in a variety of organisms and are often associated with deep

mitochondrial divergence in clades with otherwise high nuclear similarity [57, 74]. To our

knowledge, the *D. paulistorum* species complex is the first reported case of stable

heteroplasmy originating from mitochondrial introgression.

Our analyses allow us to further infer that the introgression of β likely happened before

the diversification of the semispecies in *D. paulistorum*. The presence of the β mitotype in 11

of 17 sampled lines from four different semispecies of *D. paulistorum* makes it more

parsimonious that the mitotype was inherited once from a common ancestor rather than

acquired multiple times. The ancestral introgression of β is further suggested by the observed

convergence in third codon position between the α and β mitotypes (**Fig S4**). Regardless of

whether the underlying mechanism of convergence is exposure of the β genome to the

mutational bias (amelioration) or the selective pressure (codon usage) of *D. paulistorum*, it

likely required significant time to act. A third indication that β introgression predates *D.*

paulistorum diversification is the presence of potential β-NUMTs in *D. equinoxialis*. Although

we could not conclusively determine the nuclear origin of these sequences, their presence

might indicate that *D. equinoxialis* at some point carried a β-like mitochondrion. If so, β could

have been introgressed already in the common ancestor of *D. paulistorum* and *D. equinoxialis*.

A comparison between the nuclear and mitochondrial trees suggests that several other

introgression events might have occurred in the willistoni subgroup apart from the acquisition

of β. We observed that the α clade branches basally to the willistoni clade even though nuclear

data positions *D. paulistorum* as one of the most derived species in the willistoni subgroup.

This suggests that perhaps the extant α mitotype does not directly descend from the ancestral

mitochondria of *D. paulistorum*, even if the ancestral mitochondrion was also an α-like

mitotype. Instead, the phylogenetic placement of all extant α mitochondria indicates that an

introgression in the ancestor of *D. paulistorum* and *D. equinoxialis* from an unsampled deep

diverging α-carrying lineage within the willlistoni group might have given rise to the current α

mitotypes in a similar mode to what we propose for β. One possibility is that the two extant

mitotypes were introgressed at the same time, either together or separately.

Incongruences between the nuclear and mitochondrial trees also indicate more recent
introgressions between *D. paulistorum* semispecies and lines, as mitotype distribution does
not follow the nuclear phylogeny. Since *D. paulistorum* is a recent radiation, i.e., a species
cluster *in statu nascendi* ([44] and references therein), such pattern could be explained by
hybridization in a more recent past, when the pre- and postmating incompatibilities that exist
today between semispecies (reviewed in [75], [44], and this study) were still incomplete.
Although both F_1 hybrid sexes from our inter-semispecies crosses were sterile, which would
have prevented mitochondrial introgression, possibly some rare fertile F_1 female flies could be
found with extended tests and/or other line combinations. As discussed in early literature [76],
it is possible that sporadic hybridization occurs between less discriminating and partially
compatible populations or semispecies, and those could perhaps lead to the serendipitous
events of mitochondrial introgression. As shown earlier, the Transitional semispecies is
particularly permissive to inter-semispecies crossings and has the potential to bridge borders
[77]. Specific lines of other semispecies have also been shown to be less isolated from others,
potentially allowing at least some rare cases of genetic leakage and exchange [78]. For the
one Transitional line included in our study, SM, we see a strong discrepancy between the
nuclear and mitochondrial trees, with the mitochondrial genome nested within the Orinocan
mitochondrial clade and the nuclear tree positioning it with Centro-American. Both our
phylogenetic analysis and experimental data from crosses suggest that $\alpha2$ mitochondria are
particularly likely to be transferred in crosses between semispecies, and even when crossed
with sister species.

Another possible driver of mitochondrial introgression after a hybridization event is the
maternally inherited endosymbiont *Wolbachia*. *Wolbachia* strains with strong reproductive
manipulation phenotypes or which provide enhanced host fitness can spread fast in the
population and are known to cause sweeps of the mitotype associated with infected females
[79, 80]. *D. paulistorum* is known to carry obligate *Wolbachia* symbionts, but it is not yet known
if these cause any reproductive manipulation besides premating isolation [47, 49]. Thus, we
currently have no indication that *Wolbachia* has any part in the introgression, past or on-going.

Introgression and heteroplasmy as drivers of the evolution of selfish mitochondrial
traits

Our inferences on mitochondrial evolution within *D. paulistorum* provide us with a hypothesis
about how features specific to $\alpha2$ may have evolved. We speculate that the introgression of β
into an α-carrying *D. paulistorum* likely created competition between mitotypes for a chance
to be transferred to the next host generations. If the ancestrally introgressed β mitotype was
more efficient at maternal transmission or perhaps just a better energy producer than the
ancestral α, selection could have made it the more dominant mitotype in the *D. paulistorum*

species complex. Over time, this could have led to a reduction in prevalence and possibly
resulted in extinction of the α mitotype from the species. In this scenario, the evolution of
biparental transmission and the Replication precox phenotype may have allowed the α2 lineage
to persist selfishly next to the primary mitotype. Such traits would perhaps guarantee the
transmission of α2 to the next host generation independently of its ability to meet the energetic
needs of the host or to compete with the maternal mitotype for presence in the egg.

Can we be sure that α2 is cytoplasmic?

Finally, we are aware of a potential alternative hypothesis which would suggest that α2 is not
a cytoplasmic mitotype but might stem from a so-called "mega-NUMT". It was recently shown
that mega-NUMTs can mimic heteroplasmy and extremely rare cases of paternal
mitochondrial DNA leakage in humans and other vertebrates (recently reviewed in [81]). In
this case, mitochondrial DNA-concatemers produced by extrachromosomal rolling circle
replication were found as large-sized chromosomal insertions in human autosomes and
wrongly assumed to be unusual biparental transmission of an additional mitotype [82].
However, in *D. paulistorum*, we reject this alternative hypothesis based on the following
reasons: Contrary to mega-NUMTs, which rapidly accumulate degenerative mutations and
structural variation, the genome of the α2 mitotype resembles a fully functional mitochondrial
DNA in size and content. Furthermore, the α2 genome is highly conserved within and between
heteroplasmic *D. paulistorum* semispecies as well as between long-term lab lines and recently
collected lines, which suggests that it evolves under purifying and not neutral selection. In
addition, mega-NUMTs are cryptic and if detected they are transmitted at very low frequencies
and thereby do not follow strict Mendelian inheritance. This is in sharp contrast to α2 that
shows almost perfect biparental transmission within a line and upon introgression this selfish
mitotype has the capacity for invasive spreading. Finally, our backcross experiments by
introducing α2 in a homoplasmic nuclear backgrounds clearly demonstrate the persistance
and spreading capacity of this selfish mitotype, which is self-sufficient and hence not
associated with any mega-NUMTs.

CONCLUSION

In this paper we present *D. paulistorum* as the first example of an animal with high level, stable
heteroplasmy, which is derived from introgression and is robustly maintained in both sexes.
The system is remarkable for involving a secondary mitotype with a seemingly functional
genome, which appears to not respond to the energy demands of the host and instead relies
on biparental inheritance and rapid replication in the early embryo as potentially selfish

strategies for spreading and persisting. Additionally, its higher titer and tighter co-evolution
with the nuclear background in males might hint at a specific male related function.

Given its intact putatively functional genome and recently evolved selfish phenotype,
the $\alpha2$ mitotype can potentially be referred to as a sexually transmitted, parasitic infection
rather than a primary maternally inherited mitochondrion. Hence, since $\alpha2$ appears to have
assumed a lifestyle with less host control than typical primary mitochondria, we propose to
name it "*Spartacus*" in reference to the famous ancient Roman slave, gladiator and then
commander who escaped from imperial oppression by rebellion, for at least a short period of
his lifetime.

Our results contribute to the increasing realization, driven by recent large-scale
sequencing efforts, that animal mitochondria often do not follow the dominant model of
homoplasmic, maternally inherited organelles, neither on ecological or evolutionary times
scales. Sometimes even organisms, which have been studied for decades, such as *D.*
paulistorum, harbor unexpected complexity.

MATERIALS & METHODS

Fly strains, semispecies typing, and *Wolbachia* knockdown
Information regarding the *Drosophila* lines used in this study is summarized in **Table S1**. All
flies were kept at 25±1°C on Formula 4-24 *Drosophila* instant food (Carolina, USA) with a 12
hr light-dark cycle. Typing of the biological semispecies of the *D. paulistorum* isofemale lines,
which were recently collected from French Guiana (FG16, FG103, FG111, FG295 and FG572),
was done by analyzing their reproductive compatibility towards reference lab strains from each
of the six semispecies (A28 (AM), O11 (OR), C2 (CA), L1 (IN), SM (TR) and MS (AB)**).** Each
new line was assigned to the semispecies with which it produced fertile F1 offspring. Crosses
were performed in 2-3 biological and three technical replicas (in total 30 crosses in 2-3
biological replicas). Each biological replicate consisted of 3-6 females and 3-6 males, which
were kept in the same vial for 3-5 days before transfer into a new vial for generating one
technical replicate. Egg-to-adult development was monitored and information on all crosses
including the number of F_1 flies hatched is summarized in **Table S6**.

1003

Wolbachia knockdown
In order to test the potential effect of the *Wolbachia* endosymbiont of *D. paulistorum* [47] on
mitotype persistence and levels, we generated lines with lowered *Wolbachia* infection titer.
Wolbachia was knocked down by keeping flies on food containing 0.05% tetracycline [w/v] for
four generations. Flies were repeatedly tested using PCR in order to confirm the absence of
the symbiont.　After four generations, knock-down flies were moved to tubes previously

occupied by males of the same line, so that they could feed on feces of flies with wild-type gut microbiota. This aimed to restore any gut flora affected by the antibiotic and keep the knockdown and wild-type fly lines as similar as possible with the exception of *Wolbachia* presence [49].

Single fly Southern blots

Single fly Southern blots were performed following the protocol by [83]. In brief, DNA was extracted from single individuals, digested with the restriction enzyme *Hin*dIII and processed for electrophoresis on a vertical 0.8% agarose gel. After vacuum-blotting onto a positively charged nylon membrane, fragments were hybridized with P_{32}-labelled mitochondrial *CO1* probe derived from a plasmid. Detection was performed by exposing the membranes to X-ray films at -80 degrees.

Restriction Fragment Length Polymorphism-PCR (RFLP-PCR)

Total DNA was extracted from single or multiple flies using the Gentra Puregene Tissue kit (Qiagen, Germany). Flies were homogenized using a TissueLyser LT (Qiagen) at 50 Hz for 50 sec and DNA was consequently isolated following the manufacturer's protocol. DNA concentration was determined with a NanoDrop OneC Spectrophotometer (Thermo Scientific). RFLP-PCR for the mitochondrial Cytochrome c oxidase genes *CO1* and *CO2* was set up in 20 µl reactions containing 1x Promega reaction buffer, 2.5 mM $MgCl_2$, 0.5 µM of each primer (**Table S11**), 200 µM of each dNTP, and 0.025 U of GoTaq G2 DNA Polymerase. *CO1/CO2* reactions were performed using the following thermal profile: initial denaturation for 2 min at 95°C followed by 30 cycles consisting of 45 sec denaturation at 95°C, 45 sec annealing at the appropriate primer conditions Tm°C (**Table S11**) and 30 sec extension at 72°C. Final extension was carried out at 72°C for 10 min. PCR products were consequently used for RFLP (digest plus undigested control) using the restriction enzymes *Hin*dIII and *Eco*RV-HF for *CO1*, and *Msp*I for *CO2*. Reactions were performed in 20 µl volume containing 10 or 20 U of enzyme, 1x reaction buffer and 10 µl PCR product. After overnight incubation at 37°C, enzymes were inactivated by either heat (80°C for 20 min) or by adding EDTA-containing gel loading dye. 10 µl of undigested and digested samples were then run on a 1.5% agarose gel and post-stained with 10 mg/ml ethidium bromide. Imaging was performed using a Molecular Imager ChemiDocTM XRS Imaging System and mitochondrial heteroplasmy levels were quantified using Image Lab 5.2.1 software.

Quantifying levels of heteroplasmy

Undigested samples representing the sum of both mitotypes were set as reference band with relative quantity 1. The relative levels of mitochondria were then quantified by summing up the two digested fragments plus the undigested fragment. Due to technical differences in band

1048   intensity, however, the sum of all fragments did not result in a relative quantity of 1. Hence the

1049   actual sum value was set to 100% to determine the percentage of mitotype levels. Smaller

1050   fragments with length up to approximately 100 bp were not detectable after ethidium bromide

1051   staining and we hence used the correction factor $Cf = \frac{undigested[bp]}{digested[bp]}$ to estimate their relative

1052   quantities.

1053

DNA extraction, genome sequencing and assembly

For each line 20 ovaries were dissected from 3-day old adult females and DNA extraction was

performed using the Gentra Puregene Blood and Tissue kit (Qiagen). DNA samples used for

PacBio sequencing were produced using the protocol described in [84]. Briefly, *Drosophila*

flies were transferred to apple juice agar plates and allowed to oviposit for 2 hours, after which

the eggs were collected, washed and dechorionated in 50% bleach and manually

homogenized using a plastic pestle. Following centrifugation and filtration of the homogenate,

the resulting cell pellet was subjected to whole genome amplification using the Repli-g® midi

kit (Qiagen) after which the DNA was purified using QIAamp® DNA mini kit (Qiagen) according

to the manufacturer's recommendations.

1064

Sequencing

Drosophila paulistorum lines from the classic six semispecies plus lines from more recent

collections were used for genome sequencing (**Table 1 and S1**). DNA extracted from ovaries

was used to construct 350 bp fragment TruSeq libraries, multiplexed and sequenced at the

Uppsala SNP and Seq platform in three different runs on an Illumina 2500 HiSeq machine

generating 2x125 bp reads. The Illumina reads were quality filtered and trimmed using

Trimmomatic-0.30 and error corrected using SPAdes-3.5.0. Amplified DNA extracted from

early embryos from three lines (O11, POA-1 and TP37) was used to create 5 kb fragment

SMRTbell libraries. Each library was run using P6-C4 chemistry in one SMRT cell on a RSII

PacBio instrument. PacBio libraries were produced and sequencing was performed at the

Uppsala Genome Center, Uppsala, Sweden.

1076

Mitochondrial genome assembly

The two different mitochondrial genomes α and β of the O11 line were generated by mapping

all Illumina reads to the published mitochondrial genome of *D. willistoni* (NCBI accession

number BK006338.1) using BWA-MEM [85] and thereafter extracting all aligned reads and

their mates using SAMtools [86]. The extracted Illumina reads were then subsampled to

achieve an approximate coverage of 100 and assembled together with PacBio filtered

subreads using SPAdes-3.10.1 [87] with multiple different k-mers settings. The β

mitochondrion consistently came out as a single full length contig, so the best assembly was

chosen based on the contiguity of the putative α mitochondrion. From this assembly the contig

representing the β mitochondrion and the contigs with a homogenous coverage likely

representing the α mitochondrion were taken out and used as trusted contigs together with

the Illumina data to generate a better assembly of the α mitochondrion. Finally, the Illumina

and PacBio data were aligned against the resulting contigs using BWA, MEM and SW

respectively. The resulting bam-file was converted to an ace-file that was used for manual

curation of the sequences in Consed [88]. The mitochondrial genomes of all other *Drosophila*

lines were generated by mapping Illumina reads to the two O11 mitochondria with BWA-MEM,

extracting all aligned reads and their mates using SAMtools and assembling the extracted

reads with SPAdes-3.10.1 with multiple different k-mers settings. Since PacBio data were only

generated from the O11, TP37 and POA-1 lines, the assemblies of all other lines were run

using only Illumina reads. For all lines, this assembly procedure generated one contig that

represented a complete (O11) or nearly complete circular mitochondrial genome.

To test for heteroplasmy and to correct possible errors in contigs, the Illumina reads

for each line were mapped back to the contig containing a complete or nearly complete

mitochondrion for that line using BWA-MEM and indels were realigned with the IndelRealigner

from GATK [89]. The resulting bam-file was then used to call variants using Freebayes with a

range of parameters and converted to ace-format for manual curation in Consed. A line was

determined to be heteroplasmic if 1) at least 80 sites (ca 0,5%) had a variant and 2) the

variants had a consistent alternative allele frequency over the whole or a significant part of the

mitochondrial genome. These settings were chosen to avoid calling variant genomes based

on sequence errors and to make sure that only complete mitochondrial genomes (not NUMTs)

were used to decide heteroplasmy status. If a line was seen to be heteroplasmic using these

criteria, two different alternatives were used to resolve the assembly of the second

mitochondrion. 1) The Illumina reads mapping to the O11 α and O11 β mitochondria were split

into two separate data sets and assembled separately, 2) in the case of MS that has two α

mitochondria, variant calling was performed with lofreq [90], variants with a frequency between

8-30% were selected and used to make a new consensus with bcftools and the reads were

mapped to the two variants of the mitochondrial genome. The same procedure was done in

multiple rounds until no more variant sites within the correct frequency were found. The final

bam-file was converted to ace-format and manually inspected in Consed.

1116

Alignments and phylogenetic analyses

Datasets of either whole mitochondrial genomes, nuclear markers or mitochondrial genes

were aligned with MAFFT v6.850b (mafft-linsi algorithm) [91] and trimmed with TrimAl

v1.2rev59 [92] using the automated1 for removal of poorly aligned regions. Concatenated sets

of either mitochondrial or nuclear genes were created by combining trimmed alignments of

single genes. Datasets of mitochondrial protein coding genes containing either only the 3rd or

a combination of 1st and 2nd codon positions were created by parsing the alignments with an

in-house script.

Maximum likelihood trees were built with RAxML version 7.2.6 [93] using -f a -m GTRGAMMA

-x 12345 and -# 1000. Bayesian trees were made with Mrbayes version 3.2.6 [94, 95] using

"nst= mixed" for estimating evolutionary models. Analyses were run for 10.000.000

generations with "nruns= 2", "nchains= 4" and "samplefreq= 1000". A burn-in of 25% was

applied. Trees were visualized in FigTree v. 1.4.3 [96] and rooted on the species of the *saltans*

group.

1131

Identification of synonymous and non-synonymous substitutions

Synonymous and non-synonymous substitutions in pairwise comparisons between lines of the

α x β and α1 x α2 clades were identified in single gene alignments using codeml (PAML

package v. 4.8) [97]. Values obtained from pairwise comparisons between lines were then

used to estimate an average for the clades. The same method was used for calculating the

number of nonsynonymous substitutions per non-synonymous site (dN), synonymous

substitutions per synonymous site (dS) and the dN/dS ratios. Synonymous and non-

synonymous mutations unique to α1 and α2 were counted through visual inspection of single

gene alignments using Aliview v. 2019 [98].

1141

Assessing rarity and functional impact of α2-unique mutations

We used the MutationAssessor server (www.mutationsassessor.org) to estimate the potential

impact of non-synonymous substitutions exclusive to α2 to protein function. Since the server

only analyses human sequences, we used the Uniprot website (www.uniprot.org) to align *D.*

melanogaster and human *CO3*, *ND3* and *ND4* amino acid sequences and identify the

positions in the human genes which correspond to the non-synonymous substitution sites

unique to α2. With that information we generated the following input strings which were then

used in the server: COX3_HUMAN H6N, NU3M_HUMAN T21M and NU4M_HUMAN I391V.

We used Uniprot to assess whether the α2-unique non-synonymous mutation in *CO3* was

found in other species. This was done by searching for "COX3" in the UniProtKB database,

aligning all entries of the first results page (250 entries) and visually checking for the presence

of an N residue in the position of interest. The process was repeated for the subsequent result

pages until 2000 entries had been aligned and checked.

1155

Quantitative Real-Time PCR (qPCR)

DNA was extracted by homogenizing whole flies in 200µl Chelex solution at 5% concentration

in distilled water. A total of 10µl of proteinase K (10mg/ml) was added per sample and vials

were incubated for 6 hours at 56°C. After digestion, vials were centrifuged at 16,200 x g for 3

minutes, and the supernatant was transferred to a new vial. DNA was stored at 4°C and

freeze-thaw cycles were avoided when possible.

Samples were run in duplicate using SYBR Green reagent on a StepOnePlus™ Real Time

PCR machine (Applied Biosystems). Technical repeats with >1 Ct difference were rerun or

discarded. Each reaction contained 10µl Maxima SYBR Green/ROX qPCR Master Mix (2x),

1µl of forward primer, 1µl of reverse primer, 6µl nuclease-free water and 2µl of the DNA

sample. Cycling conditions were 95°C for 20 seconds, followed by 40 cycles of: 95°C for 3

seconds and 60°C for 30 seconds for rps, and *Wolbachia* primers (**Table S11**). Discriminant

mitochondrial primers cycling conditions were 95°C for 20 seconds, followed by 40 cycles of:

95°C for 3 seconds and 55°C for 30 seconds (**Table S11**). Discriminant primers used linked

base technology to increase specificity (Exiqon). Melt curves and the fly control gene *rps17*

were run on each plate to control for successful single-peak amplification and to control for

DNA quality. All primers had >90% efficiency. For absolute quantification, each 96-well qPCR

plate was analyzed using StepOne Software v2.2.2 (Applied Biosystems) and Ct values were

obtained by comparing each primer sample to a single standard curve of known concentration

and using identical threshold and baseline levels for each primer target across plates.

Standard curves were created by amplifying positive control samples using PCR, calculating

DNA concentrations using QBIT, and then serially diluting the sample 1:10 with distilled water

to create a 5-sample curve comprising known concentrations decreasing from 10pmol/ ml.

Samples with Ct values over 30 were classed as negative, confirmed by our negative controls.

As a result, the detection limit was 50 copies per sample for *rps17*, 115 for *CO3* (SNP 1), 735

for *CO3* (SNP 2) and 1627 for *CO3* (SNP 3). This varied for the FG295 qPCR, where samples

with Ct values over 20 were classed as negative for the *CO3* (SNP 2) primer due to very high

amplification of some DNA segments. This was confirmed by negative flies, but led to a high

threshold of detection of 6150 for *CO3* (SNP 2). DNA concentration was calculated for each

sample by comparing sample Ct values back to the equation of the standard curve. DNA copy

numbers were then calculated by using DNA product length to determine DNA concentration.

To control for fly size and extraction efficiency, copy numbers for each sample were

normalized using the fly housekeeping gene as a control, giving a final value in terms of

mitochondrial genome copies to fly nuclear genome copy ratio.

Since we used three independent experimental approaches to determine levels of

heteroplasmy in *D. paulistorum*, i.e. NGS, RFLP-PCR, and qPCR, we compared the

robustness of these. We looked at the levels of $\alpha2$ in O11 females (**Fig S5**), and found that all

three techniques determined similar levels, hence confirming the robustness of our

experimental approach.

1195

Cross-mating and embryo collection

For mass-mating crosses (pools), virgin males and females from different *D. paulistorum* lines
were obtained by collecting newly emerged flies within 2-3 h intervals to avoid any possible
sibmating. Collected flies were kept separated for 5-7 days to ensure their virgin status.
Crosses were set up in pools using 10-20 females and 20-30 males per each vial and in 2-3
replicates. Flies were mated for one week and then transferred to new vials or used for
consequent embryo collecting. F_1 flies emerging from the first vials were collected for DNA
extraction. Mass-mating A28 x O11 hybrids were generated by mating 10 five-day-old virgin
A28 females to 10 O11 males. F_1 individuals were used for assays. For generating
interspecies hybrids *D. paulistorum* O11 males were mated either with *D. equinoxialis* or *D.*
willistoni virgin females in pools of 20 females and 60 males per cross. F_1 flies emerging from
the bottles were collected for DNA extraction. Single fly crosses were generated by collecting
virgin flies every 6-8 hours to avoid sibling mating. Collected flies were kept separated for 5-7
days to ensure their virgin status before use in crossings. Single pairs of male and female flies
were mated for two weeks in fresh vials. Emerging F_1 flies were collected, aged to 8-10 days
and frozen for subsequent DNA extraction. For FG295 x O11 and O11 x FG295 crosses,
single-pair sibling mating were continued for four consequent generations.
1213

Backcrossing
For backcrossing experiments, heteroplasmic F1 flies obtained from reciprocal crosses
between O11 and FG295 parents (G0) were crossed in pools to the homoplasmic FG295 line.
Virgin F_1 males and females were obtained from the G0 crosses by collecting newly emerged
flies within 2-3 h intervals to avoid sib mating. Collected flies were kept separated for 5-7 days
to ensure their virgin status. Backcross 1 (BC1) was set up in pools using 5-10 females and
5-10 males per each vial. The first 10 F_1 flies of each sex emerging from the vials were
collected for further backcrossing with FG295 line and the remaining F_1 flies were frozen in
liquid nitrogen for future DNA extraction. Backcross 2 (BC2) was set up similarly to BC1.
Between 10 and 20 flies were analyzed for each backcross.
1224

Collection of staged embryos
D. paulistorum embryos at early (0-30 min AEL), mid (3-4 hrs AEL) and late (18-20 hrs AEL)
stages of development were collected on homemade medium (agar, molasses, cornmeal and
yeast medium) in plastic Petri dishes (3 x 3 cm) using collection chambers at 25°C. Prior to
collecting embryos, 1-2 weeks old flies were kept in these chambers for 2-3 days to adapt to
the new environment. Embryos were then manually picked with a dissection needle and
transferred immediately to ice-cold lysis buffer (Qiagen). We collected 20 embryos for each
sample with at least five biological replicates. Embryos were immediately processed for DNA
extraction or stored at -20°C for no more than one week.
1234

Recombination analysis

We checked for intergenic recombination between lines by analyzing single gene trees built as described in the "Alignments and phylogenetic analyses" methods section. Tests for intragenic recombination on single-gene and whole-genome alignments were done using Geneconv v.1.81 and PhiPack with default parameters. Recombination over mutation rates were calculated for single-gene and whole-genome alignments using ClonalFrame v.1.2 with default settings.

Nuclear markers

Illumina reads from each fly line were separately assembled with both Megahit v.1.1.2 [99] and SPAdes v. 3.13.0 using default parameters. We then blasted the *D. willistoni* CDSs of the selected nuclear markers (**Table S12**) to each assembly and retrieved sequences with an alignment length of over 200 bp and above 70% identity. The obtained contigs were aligned to the corresponding *D. willistoni* gene with MAFFT v6.850b. The alignments were visualized with Aliview v. 2019 and inspected for assembly issues and, if necessary, multiple exons were joined into a single sequence using Aliviews' "merge" function. For each fly line and marker gene, results obtained from both assemblies were compared and only the one with highest quality was used in further analyses.

Identification of nuclear mitochondrial DNA (NUMTs)

Illumina reads of each line were separately mapped with BWA-MEM v.0.7.10 to the genome assemblies generated with Megahit v. 1.1.2. The average per-base coverage of each assembled contig was estimated with SAMtools mpileup and contigs with average coverage lower than 2 were excluded from the analysis.

Each genomic assembly was blasted to all previously assembled *D. paulistorum* mitochondrial genomes. Sequences corresponding to high scoring pairs (HSPs) with at least 200 bp length and 90% similarity to mitochondrial DNA were retrieved. Recovered sequences with more than 97% identity to either α or β genomes were deemed "conserved" and tentatively assigned to the clade they were most similar to, while those with 90-97% identity were considered divergent. Sequences retrieved from each assembly were mapped back to an α and a β mitochondrial genome (C2, O11 beta) with BWA-MEM and the number of bases covered by them were counted with Samtools mpileup. This was done in order to determine how much of the mitochondrial genome was covered by contigs belonging to each category (α, β, diverged). Genes present in the retrieved sequences were identified by blasting each sequence to a database containing all genes from the A28 mitochondrial genome (CDS, tRNA and rRNA). All contigs containing mitochondrial sequences were blasted against NCBI's nt database for identification of potential matches to non-mitochondrial references.

Sequences retrieved from each genomic assembly were considered NUMTs if the
contigs in which they were found a) also contained matches to non-mitochondrial references;
or b) had average coverage similar (± 30%) to the reference nuclear coverage of its line. The
reference nuclear coverage for each line was calculated as the average coverage of the
contigs containing the genes *CAD/r*, *Beta Spectrin* and *FASN3*. These were identified by
blasting *D. willistoni* CDSs of these genes to each assembly. The reference mitochondrial
coverage of each line was defined as the average per base coverage of the longest contig
which had 100% identity to the line's assembled mitochondrial genome(s).
1280
Statistics
Statistical analyses varied depending on the data source and distribution. For the qPCR tissue
distribution of α and β mitotypes between different ages and sexes we used ANOVA on log-
transformed data to increase normality. In the qPCR analysis of A28 x O11 hybrids, we used
ANOVA with post-hoc Tukey tests of significance. Due to low numbers of repeats and power
in the intraspecies FG295 x O11 hybrids, Wilcoxon Rank Sum tests were used for significance
testing. Assumptions and variance were checked before use. These analyses were
completed using R version 3.3.2 [100].
In the RFLP-PCR experiments, the α2-levels in embryos from different *D. paulistorum*
semispecies and crosses at different developmental stages were compared using One-way
ANOVA with post-hoc Tukey HSD tests. Quantifications were made with the help of an online
calculator (www.astatsa.com). The raw data showing levels of α2 in percent were obtained
with Image Lab 5.2.1 software and transformed from percentage to arcsine for normal
distribution before applying the statistics. RFLP-data were statistically analyzed using ANOVA
and Student's t tests.
1296

ACKNOWLEDGEMENTS
We thank Matthias Schaefer, Lee Ehrman, DeEtte Walker and Silvia Bulgheresi for critical
and constructive discussions during the whole period of the project and Wolfgang Däuble for
valuable contributions in the early stage of the project, Jonathan Haars for valuable
contributions to preliminary data for embryos and both Julia Mras and Christian Binnyei for
excellent technical support. Sequencing was performed at the SNP&SEQ Technology
Platform and Uppsala Genome Center in Uppsala, Sweden, which is part of the Swedish
National Genomics Infrastructure. Finally, we want to thank Aurélie Hua-Van for fly sampling
and the Nouragues research field station (managed by CNRS), which benefits from
"Investissement d'Avenir" grants managed by Agence Nationale de la Recherche (AnaEE
France ANR-11-INBS-0001; Labex CEBA ANR-10-LABX-25-01). The project was funded by

the Swedish research council VR grant 2014-4353 to LK, and by the Austrian Science Fund

FWF grant P28255-B22 to WJM.

1310

1311   **DATA AVAILABILITY**

1312   Raw sequence data used in this project as well as assembled whole mitochondrial genomes

1313   and nuclear markers are available in NCBI under the Bioproject PRJNA643793. Accession

1314   numbers of individual mitochondrial genomes are listed in Table S2.

1315

REFERENCES:

1.   Giles RE, Blanc H, Cann HM, Wallace DC. Maternal inheritance of human
mitochondrial DNA. Proceedings of the National Academy of Sciences of the United States of
America. 1980;77(11):6715-9. doi: 10.1073/pnas.77.11.6715. PubMed PMID: 6256757.
2.   Birky CW. Relaxed cellular controls and organelle heredity. Science.
1983;222(4623):468. doi: 10.1126/science.6353578.
3.   Stefano G, B., Bjenning C, Wang F, Wang N, Kream R, M. . Mitochondrial
Heteroplasmy. In: Santulli G, editor. Mitochondrial Dynamics in Cardiovascular Medicine
Advances in Experimental Medicine and Biology. 982. Cham: Springer; 2017.
4.   Burr SP, Pezet M, Chinnery PF. Mitochondrial DNA Heteroplasmy and Purifying
Selection in the Mammalian Female Germ Line. Development, Growth & Differentiation.
2018;60(1):21-32. doi: 10.1111/dgd.12420.
5.   White DJ, Wolff JN, Pierson M, Gemmell NJ. Revealing the hidden complexities of
mtDNA inheritance. Molecular Ecology. 2008;17(23):4925-42. doi: 10.1111/j.1365-
294X.2008.03982.x.
6.   Sharpley MS, Marciniak C, Eckel-Mahan K, McManus M, Crimi M, Waymire K, et al.
Heteroplasmy of mouse mtDNA is genetically unstable and results in altered behavior and
cognition. Cell. 2012;151(2):333-43. doi: 10.1016/j.cell.2012.09.004. PubMed PMID:
23063123.
7.   Stewart JB, Chinnery PF. The dynamics of mitochondrial DNA heteroplasmy:
implications for human health and disease. Nature Reviews Genetics. 2015;16(9):530-42. doi:
10.1038/nrg3966.
8.   Sloan DB, Warren JM, Williams AM, Wu Z, Abdel-Ghany SE, Chicco AJ, et al.
Cytonuclear integration and co-evolution. Nature Reviews Genetics. 2018;19(10):635-48. doi:
10.1038/s41576-018-0035-9.
9.   Christie JR, Schaerf TM, Beekman M. Selection against Heteroplasmy Explains the
Evolution of Uniparental Inheritance of Mitochondria. PLOS Genetics. 2015;11(4):e1005112.
doi: 10.1371/journal.pgen.1005112.
10.   Birky CW. Uniparental inheritance of mitochondrial and chloroplast genes:
mechanisms and evolution. Proceedings of the National Academy of Sciences.
1995;92(25):11331. doi: 10.1073/pnas.92.25.11331.
11.   Pitnick S, Karr TL. Paternal products and by-products in *Drosophila* development.
Proceedings of the Royal Society B: Biological Sciences. 1998;265(1398):821-6. doi:
10.1098/rspb.1998.0366. PubMed PMID: 9628039.
12.   Sutovsky P, Moreno RD, Ramalho-Santos J, Dominko T, Simerly C, Schatten G.
Ubiquitin tag for sperm mitochondria. Nature. 1999;402(6760):371-2. doi: 10.1038/46466.
13.   DeLuca Steven Z, O'Farrell Patrick H. Barriers to Male Transmission of Mitochondrial
DNA in Sperm Development. Developmental Cell. 2012;22(3):660-8. doi:
https://doi.org/10.1016/j.devcel.2011.12.021.
14.   Yu Z, O'Farrell PH, Yakubovich N, DeLuca SZ. The Mitochondrial DNA Polymerase
Promotes Elimination of Paternal Mitochondrial Genomes. Current Biology. 2017;27(7):1033-
9. doi: 10.1016/j.cub.2017.02.014.

15.   Sato K, Sato M. Multiple ways to prevent transmission of paternal mitochondrial DNA
for maternal inheritance in animals. The Journal of Biochemistry. 2017;162(4):247-53. doi:
10.1093/jb/mvx052.
16.   Wolff JN, Gemmell NJ. Lost in the zygote: the dilution of paternal mtDNA upon
fertilization. Heredity. 2008;101(5):429-34. doi: 10.1038/hdy.2008.74.
17.   Passamonti M, Ghiselli F. Doubly Uniparental Inheritance: Two Mitochondrial
Genomes, One Precious Model for Organelle DNA Inheritance and Evolution. DNA and Cell
Biology. 2009;28(2):79-89. doi: 10.1089/dna.2008.0807.
18.   Zouros E, Ball AO, Saavedra C, Freeman KR. Mitochondrial DNA inheritance. Nature.
1994;368(6474):818-. doi: 10.1038/368818a0.
19.   Doublet V, Souty-Grosset C, Bouchon D, Cordaux R, Marcadé I. A thirty million year-
old   inherited   heteroplasmy.   PLoS   One.   2008;3(8):e2938-e.   doi:
10.1371/journal.pone.0002938. PubMed PMID: 18698356.
20.   Doublet V, Raimond R, Grandjean F, Lafitte A, Souty-Grosset C, Marcadé I.
Widespread atypical mitochondrial DNA structure in isopods (Crustacea, Peracarida) related
to a constitutive heteroplasmy in terrestrial species. Genome. 2012;55(3):234-44. doi:
10.1139/g2012-008.
21.   Peccoud J, Chebbi MA, Cormier A, Moumen B, Gilbert C, Marcadé I, et al. Untangling
Heteroplasmy, Structure, and Evolution of an Atypical Mitochondrial Genome by PacBio
Sequencing. Genetics. 2017;207(1):269. doi: 10.1534/genetics.117.203380.
22.   Robison GA, Balvin O, Schal C, Vargo EL, Booth W. Extensive Mitochondrial
Heteroplasmy in Natural Populations of a Resurging Human Pest, the Bed Bug (Hemiptera:
Cimicidae). Journal of medical entomology. 2015;52(4):734-8. Epub 2015/05/23. doi:
10.1093/jme/tjv055. PubMed PMID: 26335484.
23.   Meza-Lázaro RN, Poteaux C, Bayona-Vásquez NJ, Branstetter MG, Zaldívar-Riverón
A. Extensive mitochondrial heteroplasmy in the neotropical ants of the *Ectatomma ruidum*
complex (Formicidae: Ectatomminae). Mitochondrial DNA Part A. 2018;29(8):1203-14. doi:
10.1080/24701394.2018.1431228.
24.   Mastrantonio V, Latrofa MS, Porretta D, Lia RP, Parisi A, Iatta R, et al. Paternal
leakage and mtDNA heteroplasmy in *Rhipicephalus spp* ticks. Scientific reports.
2019;9(1):1460. doi: 10.1038/s41598-018-38001-8.
25.   Kastally C, Mardulyn P. Widespread co-occurrence of two distantly related
mitochondrial genomes in individuals of the leaf beetle *Gonioctena intermedia*. Biology Letters.
2017;13(11):20170570. doi: 10.1098/rsbl.2017.0570.
26.   Solignac M, Monnerot M, Mounolou JC. Mitochondrial DNA heteroplasmy in
Drosophila mauritiana. Proceedings of the National Academy of Sciences of the United States
of America. 1983;80(22):6942-6. doi: 10.1073/pnas.80.22.6942. PubMed PMID: 6316335.
27.   Matsuura ET, Fukuda H, Chigusa SI. Mitochondrial DNA heteroplasmy maintained in
natural populations of *Drosophila simulans* in Réunion. Genetical Research. 1991;57(2):123-
6. Epub 2009/04/14. doi: 10.1017/S0016672300029189.
28.   Dean MD, Ballard KJ, Glass A, Ballard JW. Influence of two *Wolbachia* strains on
population structure of East African *Drosophila* simulans. Genetics. 2003;165(4):1959-69.
Epub 2004/01/06. PubMed PMID: 14704179; PubMed Central PMCID: PMC1462921.
29.   Wolff JN, Nafisinia M, Sutovsky P, Ballard JWO. Paternal transmission of
mitochondrial DNA as an integral part of mitochondrial inheritance in metapopulations of
Drosophila   simulans.   Heredity.   2013;110(1):57-62.   Epub   2012/09/26.   doi:
10.1038/hdy.2012.60. PubMed PMID: 23010820.
30.   Nunes MDS, Dolezal M, Schlötterer C. Extensive paternal mtDNA leakage in natural
populations of *Drosophila melanogaster*. Molecular ecology. 2013;22(8):2106-17. Epub
2013/03/04. doi: 10.1111/mec.12256. PubMed PMID: 23452233.
31.   Volz-Lingenhöhl A, Solignac M, Sperlich D. Stable heteroplasmy for a large-scale
deletion in the coding region of Drosophila subobscura mitochondrial DNA. Proceedings of
the National Academy of Sciences. 1992;89(23):11528. doi: 10.1073/pnas.89.23.11528.
32.   Lecher P, Béziat F, Alziari S. Tissular distribution of heteroplasmy and ultrastructural
studies of mitochondria from a *Drosophila subobscura* mitochondrial deletion mutant. Biology
of the Cell. 1994;80(1):25-33. doi: 10.1016/0248-4900(94)90013-2.

33. Kondo R, Matsuura ET, Chigusa SI. Further observation of paternal transmission of
Drosophila mitochondrial DNA by PCR selective amplification method. Genetical Research.
1992;59(2):81-4. Epub 2009/04/14. doi: 10.1017/S0016672300030287.
34. Sherengul W, Kondo R, Matsuura ET. Analysis of paternal transmission of
mitochondrial DNA in *Drosophila*. Genes & Genetic Systems. 2006;81(6):399-404. doi:
10.1266/ggs.81.399.
35. Dokianakis E, Ladoukakis ED. Different degree of paternal mtDNA leakage between
male and female progeny in interspecific *Drosophila* crosses. Ecology and evolution.
2014;4(13):2633-41. Epub 2014/06/19. doi: 10.1002/ece3.1069. PubMed PMID: 25077015.
36. Hahn A, Zuryn S. The Cellular Mitochondrial Genome Landscape in Disease. Trends
in Cell Biology. 2019;29(3):227-40. doi: 10.1016/j.tcb.2018.11.004.
37. Havird JC, Forsythe ES, Williams AM, Werren JH, Dowling DK, Sloan DB. Selfish
mitonuclear conflict. Current Biology. 2019;29(11):R496-R511. doi:
https://doi.org/10.1016/j.cub.2019.03.020.
38. Oliver KM, Degnan PH, Burke GR, Moran NA. Facultative Symbionts in Aphids and
the Horizontal Transfer of Ecologically Important Traits. Annual Review of Entomology.
2009;55(1):247-66. doi: 10.1146/annurev-ento-112408-085305.
39. Sloan DB, Moran NA. Genome reduction and co-evolution between the primary and
secondary bacterial symbionts of psyllids. Molecular biology and evolution. 2012;29(12):3781-
92. Epub 2012/07/19. doi: 10.1093/molbev/mss180. PubMed PMID: 22821013.
40. Ferrari J, Vavre F. Bacterial symbionts in insects or the story of communities affecting
communities. Philosophical Transactions of the Royal Society B: Biological Sciences.
2011;366(1569):1389-400. doi: 10.1098/rstb.2010.0226.
41. Lamelas A, Gosalbes MJ, Manzano-Marín A, Peretó J, Moya A, Latorre A. *Serratia*
symbiotica from the Aphid *Cinara cedri*: A Missing Link from Facultative to Obligate Insect
Endosymbiont. PLOS Genetics. 2011;7(11):e1002357. doi: 10.1371/journal.pgen.1002357.
42. McCutcheon JP, McDonald BR, Moran NA. Convergent evolution of metabolic roles in
bacterial co-symbionts of insects. Proceedings of the National Academy of Sciences. 2009.
Epub 2009/08/27. doi: 0906424106 [pii]
10.1073/pnas.0906424106. PubMed PMID: 19706397.
43. Burla H, Da Cunha AB, Cordeiro AR, Dobzhansky T, Malogolowkin C, Pavan C. The
willistoni group of sibling species of *Drosophila*. Evolution. 1949;3:300-14. doi: 10.1111/j.1558-
5646.1949.tb00031.x.
44. Ehrman L, Powell JR. The *Drosophila willistoni* species group. In: Ashburner M,
Carson HL, Thompson Jr. JN, editors. The Genetics and Biology of *Drosophila*. 3b. New York:
Academic Press Inc; 1982. p. 193-220.
45. Dobzhansky T, Ehrman L, Pavlovsky O, Spassky B. The superspecies *Drosophila*
paulistorum. Proceedings of the National Academy of Sciences. 1964;51(1):3. doi:
10.1073/pnas.51.1.3.
46. Dobzhansky T, Spassky B. *Drosophila paulistorum*, a cluster of species in *statu*
nascendi. Proceedings of the national Academy of Sciences. 1959;45(3):419-28.
47. Miller WJ, Ehrman L, Schneider D. Infectious speciation revisited: impact of symbiont-
depletion on female fitness and mating behavior of *Drosophila paulistorum*. PLoS Pathogens.
2010;6(12):e1001214. Epub 2010/12/15. doi: 10.1371/journal.ppat.1001214. PubMed PMID:
21151959; PubMed Central PMCID: PMCPMC2996333.
48. Miller W, J, , Schneider D. Endosymbiotic microbes as adaptive manipulators of
arthropod behaviour and natural driving sources of host speciation. In: Hughes D, Brodeur J,
Thomas F, editors. Host manipulation by parasites. Oxford: University Press; 2012. p. 119-39.
49. Schneider DI, Ehrman L, Engl T, Kaltenpoth M, Hua-Van A, Le Rouzic A, et al.
Symbiont-driven male mating success in the Neotropical *Drosophila paulistorum* superspecies.
Behavior Genetics. 2019;49(1):83-98. doi: 10.1007/s10519-018-9937-8.
50. Baião GC, Schneider DI, Miller WJ, Klasson L. The effect of *Wolbachia* on gene
expression in *Drosophila paulistorum* and its implications for symbiont-induced host speciation.
BMC Genomics. 2019;20(1):465. doi: 10.1186/s12864-019-5816-9.

51.   Strunov A, Schneider DI, Albertson R, Miller WJ. Restricted distribution and
lateralization of mutualistic *Wolbachia* in the *Drosophila* brain. Cellular Microbiology.
2017;19(1). doi: 10.1111/cmi.12639. PubMed PMID: 27353950.
52.   Gleason JM, Griffith EC, Powell JR. A molecular phylogeny of the *Drosophila willistoni*
group: Conflicts between species concepts? Evolution. 1998;52(4):1093-103. doi:
10.2307/2411239.
53.   Robe LJ, Cordeiro J, Loreto ELS, Valente VLS. Taxonomic boundaries, phylogenetic
relationships and biogeography of the *Drosophila willistoni* subgroup (Diptera: Drosophilidae).
Genetica. 2010;138(6):601-17. doi: 10.1007/s10709-009-9432-5.
54.   Zanini R, Müller MJ, Vieira GC, Valiati VH, Deprá M, Valente VLdS. Combining
morphology and molecular data to improve *Drosophila paulistorum* (Diptera, Drosophilidae)
taxonomic status. Fly. 2018:1-14. doi: 10.1080/19336934.2018.1429859.
55.   Ross CL, Harrison RG. A fine scale spatial analysis of the mosaic hybrid zone between
Gryllus firmus and *Gryllus pennsylvanicus* Evolution. 2002;56(11):2296-312. doi:
10.1111/j.0014-3820.2002.tb00153.x.
56.   Toews DPL, Brelsford A. The biogeography of mitochondrial and nuclear discordance
in animals. Molecular Ecology. 2012;21(16):3907-30. doi: 10.1111/j.1365-294X.2012.05664.x.
57.   Zhang D, Tang L, Cheng Y, Hao Y, Xiong Y, Song G, et al. "Ghost Introgression" As a
Cause of Deep Mitochondrial Divergence in a Bird Species Complex. Molecular Biology and
Evolution. 2019;36(11):2375-86. doi: 10.1093/molbev/msz170.
58.   Winge H. Interspecific hybridisation between the six cryptic species of the *Drosophila*
willistoni group. Heredity. 1965;20:9-19.
59.   Wolff JN, Sutovsky P, Ballard JWO. Mitochondrial DNA content of mature
spermatozoa and oocytes in the genetic model *Drosophila*. Cell and Tissue Research.
2013;353(1):195-200. doi: 10.1007/s00441-013-1628-4.
60.   Rubenstein JLR, Brutlag D, Clayton DA. The mitochondrial DNA of *Drosophila*
melanogaster exists in two distinct and stable superhelical forms. Cell. 1977;12(2):471-82. doi:
10.1016/0092-8674(77)90123-4.
61.   Yamane T, Miller Dl, Hopfield JJ. Interaction of elongation factor Tu with the aminoacyl
transfer ribonucleic acid dimer Phe-tRNA-Glu-tRNA. Biochemistry. 1981;20(20):449-52.
62.   Moazed D, Robertson JM, Noller HF. Interaction of elongation factors EF-G and EF-
Tu with a conserved loop in 23S RNA. Nature. 1988;334(6180):362-4. doi: 10.1038/334362a0.
63.   Marygold SJ, Attrill H, Lasko P. The translation factors of *Drosophila melanogaster*.
Fly. 2017;11(1):65-74. Epub 2016/08/05. doi: 10.1080/19336934.2016.1220464. PubMed
PMID: 27494710.
64.   Sato A, Suematsu T, Aihara K-k, Kita K, Suzuki T, Watanabe K, et al. Duplication of
Drosophila melanogaster mitochondrial EF-Tu: pre-adaptation to T-arm truncation and
exclusion of bulky aminoacyl residues. Biochemical Journal. 2017;474(6):957-69. doi:
10.1042/BCJ20160929.
65.   Bruen TC, Philippe H, Bryant D. A Simple and Robust Statistical Test for Detecting the
Presence of Recombination. Genetics. 2006;172(4):2665. doi: 10.1534/genetics.105.048975.
66.   Werner M, Simmons LW. Insect Sperm Motility. Biological Reviews. 2008;83(2):191-
208. doi: 10.1111/j.1469-185X.2008.00039.x.
67.   du Plessis SS, Agarwal A, Mohanty G, van der Linde M. Oxidative phosphorylation
versus glycolysis: what fuel do spermatozoa use? Asian J Androl. 2015;17(2):230-5. doi:
10.4103/1008-682X.135123. PubMed PMID: 25475660.
68.   Chiang ACY, McCartney E, O'Farrell PH, Ma H. A Genome-wide Screen Reveals that
Reducing Mitochondrial DNA Polymerase Can Promote Elimination of Deleterious
Mitochondrial Mutations. Current Biology. 2019;29(24):4330-6.e3. doi:
https://doi.org/10.1016/j.cub.2019.10.060.
69.   Politi Y, Gal L, Kalifa Y, Ravid L, Elazar Z, Arama E. Paternal Mitochondrial Destruction
after Fertilization Is Mediated by a Common Endocytic and Autophagic Pathway in *Drosophila*.
Developmental Cell. 2014;29(3):305-20. doi: https://doi.org/10.1016/j.devcel.2014.04.005.
70.   Birky CW. The Inheritance of Genes in Mitochondria and Chloroplasts: Laws,
Mechanisms, and Models. Annual Review of Genetics. 2001;35(1):125-48. doi:
10.1146/annurev.genet.35.102401.090231.

71.   St. John J. The control of mtDNA replication during differentiation and development. Biochimica et Biophysica Acta. 2014;1840(4):1345-54. doi: https://doi.org/10.1016/j.bbagen.2013.10.036.

72.   Gemmell NJ, Metcalf VJ, Allendorf FW. Mother's curse: the effect of mtDNA on individual fitness and population viability. Trends in Ecology & Evolution. 2004;19(5):238-44. doi: 10.1016/j.tree.2004.02.002.

73.   Petit N, Touraille S, Debise R, Morel F, Renoux M, Lécher P, et al. Developmental changes in heteroplasmy level and mitochondrial gene expression in a *Drosophila subobscura* mitochondrial deletion mutant. Current genetics. 1998;33(5):330-9.

74.   Huang DI, Hefer CA, Kolosova N, Douglas CJ, Cronk QCB. Whole plastome sequencing reveals deep plastid divergence and cytonuclear discordance between closely related balsam poplars, *Populus balsamifera* and *P. trichocarpa* (Salicaceae). New Phytologist. 2014;204(3):693-703. doi: 10.1111/nph.12956.

75.   Ehrman L, Kernaghan RP. Infectious Heredity in *Drosophila paulistorum*. Ciba Foundation Symposium 6 - Pathogenic Mycoplasmas. 1972:227-50. doi: doi:10.1002/9780470719893.ch11 10.1002/9780470719893.ch11.

76.   Dobzhansky T, Pavlovsky O. Experiments on the Incipient Species of the *Drosophila paulistorum* complex. Genetics. 1967;55(1):141-56. PubMed PMID: PMC1211364.

77.   Ehrman L. The transitional races of *Drosophila paulistorum*: a study of hybrid sterility Proceedings of the National Academy of Sciences. 1962;48(2):157. doi: 10.1073/pnas.48.2.157.

78.   Malogolowkin C, Simmons AS, Levene H. On the nature of the transitional strains in *Drosophila paulistorum*. Proceedings of the National Academy of Sciences. 1964;51(2):260. doi: 10.1073/pnas.51.2.260.

79.   Turelli M, Hoffmann AA, McKechnie SW. Dynamics of cytoplasmic incompatibility and mtDNA variation in natural *Drosophila simulans* populations. Genetics. 1992;132(3):713.

80.   Hurst GD, Jiggins FM. Problems with mitochondrial DNA as a marker in population, phylogeographic and phylogenetic studies: the effects of inherited symbionts. Proceedings of the Royal Society B: Biological Sciences. 2005;272(1572):1525-34. Epub 2005/07/29. doi: 10.1098/rspb.2005.3056. PubMed PMID: 16048766; PubMed Central PMCID: PMCPMC1559843.

81.   Wei W, Pagnamenta AT, Gleadall N, Sanchis-Juan A, Stephens J, Broxholme J, et al. Nuclear-mitochondrial DNA segments resemble paternally inherited mitochondrial DNA in humans. Nature communications. 2020;11(1):1740. doi: 10.1038/s41467-020-15336-3.

82.   Luo S, Valencia CA, Zhang J, Lee N-C, Slone J, Gui B, et al. Biparental Inheritance of Mitochondrial DNA in Humans. Proceedings of the National Academy of Sciences. 2018;115(51):13039. doi: 10.1073/pnas.1810946115.

83.   Junakovic N. Southern Blot Analysis of Individual *Drosophila* Flies. In: Miller WJ, Capy P, editors. Mobile Genetic Elements Methods in Molecular Biology. 260: Humana Press; 2004.

84.   Ellegaard KM, Klasson L, Naslund K, Bourtzis K, Andersson SG. Comparative genomics of *Wolbachia* and the bacterial species concept. PLoS Genetics. 2013;9(4):e1003381. Epub 2013/04/18. doi: 10.1371/journal.pgen.1003381. PubMed PMID: 23593012; PubMed Central PMCID: PMCPMC3616963.

85.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv:13033997v2. 2013.

86.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9. Epub 2009/06/08. doi: 10.1093/bioinformatics/btp352. PubMed PMID: 19505943.

87.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. Journal of Computational Biology. 2012;19(5):455-77. doi: 10.1089/cmb.2012.0021. PubMed PMID: 22506599; PubMed Central PMCID: PMC3342519.

88.   Gordon D, Abajian C, Green P. Consed: a graphical tool for sequence finishing. Genome Research. 1998;8(3):195-202. PubMed PMID: 9521923.

89.   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et al. The
Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA
sequencing data. Genome Research. 2010;20(9):1297-303. doi: 10.1101/gr.107524.110.
90.   Wilm A, Aw PPK, Bertrand D, Yeo GHT, Ong SH, Wong CH, et al. LoFreq: a sequence-
quality aware, ultra-sensitive variant caller for uncovering cell-population heterogeneity from
high-throughput sequencing datasets. Nucleic acids research. 2012;40(22):11189-201. Epub
2012/10/12. doi: 10.1093/nar/gks918. PubMed PMID: 23066108.
91.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Molecular Biology and Evolution. 2013;30(4):772-
80. doi: 10.1093/molbev/mst010.
92.   Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25(15):1972-3.
doi: 10.1093/bioinformatics/btp348. PubMed PMID: 19505945; PubMed Central PMCID:
PMC2712344.
93.   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large   phylogenies.   Bioinformatics.   2014;30(9):1312-3.   Epub   2014/01/21.   doi:
10.1093/bioinformatics/btu033. PubMed PMID: 24451623.
94.   Huelsenbeck JP, Ronquist F. MRBAYES: Bayesian inference of phylogenetic trees.
Bioinformatics. 2001;17(8):754-5. doi: 10.1093/bioinformatics/17.8.754.
95.   Ronquist F, Huelsenbeck JP. MrBayes 3: Bayesian phylogenetic inference under
mixed models. Bioinformatics. 2003;19(12):1572-4. doi: 10.1093/bioinformatics/btg180.
96.   Rambaut   A.   FigTree   v1.4:   Tree   Figure   Drawing   Tool.   Available   at:
http://tree.bio.ed.ac.uk/software/figtree/2009.
97.   Yang Z. PAML 4: phylogenetic analysis by maximum likelihood. Molecular Biology and
Evolution. 2007;24(8):1586-91. doi: 10.1093/molbev/msm088.
98.   Larsson A. AliView: a fast and lightweight alignment viewer and editor for large
datasets.   Bioinformatics.   2014;30(22):3276-8.   Epub   2014/08/05.   doi:
10.1093/bioinformatics/btu531. PubMed PMID: 25095880.
99.   Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node
solution for large and complex metagenomics assembly via succinct de Bruijn graph.
Bioinformatics. 2015;31(10):1674-6. doi: 10.1093/bioinformatics/btv033.
100.   R-Core-Team. R: A Language and Environment for Statistical Computing. Vienna,
Austria: R Foundation for Statistical computing; 2015.
1612