DNA Methylation Reprogramming during Sex Determination and Transition in Zebrafish

Xinxin Wang[1,2#], Xin Ma[1,2#], Gaobo Wei[2,3#], Weirui Ma[2,4], Zhen Zhang[5], Xuepeng Chen[1], Lei Gao[1], Zhenbo Liu[1], Yue Yuan[2,4], Lizhi Yi[1,2], Jun Wang[6], Toshinobu Tokumoto[6], Junjiu Huang[5], Dahua Chen[2,3*], Jian Zhang[2,4,8*], Jiang Liu[1,2,7*]

1. CAS Key Laboratory of Genome Sciences and Information, Beijing Institute of Genomics, Chinese Academy of Sciences, Beijing 100101, China.

2. University of Chinese Academy of Sciences, Beijing 100049, China.

3. State Key Laboratory of Membrane Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China.

4. State Key Laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China.

5. Key Laboratory of Reproductive Medicine of Guangdong Province, the First Affiliated Hospital and School of Life Sciences, Sun Yat-sen University, Guangzhou 510275, China.

6. Integrated Bioscience Section, Graduate School of Science and Technology, National University Corporation Shizuoka University, Shizuoka 422-8529, Japan.

7. Center for Excellence in Animal Evolution and Genetics, Chinese Academy of Sciences, Kunming 650223, China.

8. Center for Life Sciences, School of Life Sciences, State Key Laboratory for Conservation and Utilization of Bio-Resources in Yunnan, Yunnan University, Kunming 650091, China.

#These authors contributed equally to this work

*Correspondence to: Jiang Liu (liuj@big.ac.cn); Jian Zhang (jianzhang@ynu.edu.cn); Dahua Chen (chendh@ioz.ac.cn);

Abstract

It is a mystery about sex determination and sexual plasticity in species without sex chromosomes. DNA methylation is a prevalent epigenetic modification in vertebrates, which has been shown to involve in the regulation of gene expression and embryo development. However, it remains unclear about how DNA methylation regulates sexual development. To elucidate it, we used zebrafish to investigate DNA methylation reprogramming during juvenile germ cell development and adult female-to-male sex transition. We revealed that primordial germ cells (PGCs) undergo significant DNA methylation reprogramming during germline development and set to an oocyte/ovary-like pattern at 9 days post fertilization (9 dpf). When blocking DNMTs activity in juveniles after 9 dpf, the zebrafish preferably develops into females. We also show that *Tet3* involves in PGC development. Notably, we find that DNA methylome reprogramming during adult zebrafish sex transition is similar to the reprogramming during the sex differentiation from 9 dpf PGCs to sperm. Furthermore, inhibiting DNMTs activity can prevent the female-to-male sex transition, suggesting that methylation reprogramming is required for zebrafish sex transition. In summary, DNA methylation plays important roles in zebrafish germline development and sexual plasticity.

Keywords

Germ cell; DNA methylation; Sex determination; Sex transition

Introduction

Sex determination in animals has been one of the most fantastic mysteries in biology. In some species sex is determined by sex chromosomes, while in some other species no sex chromosomes can be found in their genomes [1]. It is rarely understood about the mechanism of sex determination in these species. Even more interestingly, some adult animals can change their sex in nature conditions. For example, sexual fate in *sea goldie* can be regulated by the presence or absence of male fish within the population. When a dominant male dies in one group, one of the remaining females in this group will transform into a functional male [2, 3].

Many fishes, such as domesticated laboratory zebrafish strains, do not have sex chromosomes [4]. Prior to the sex determination stage during zebrafish germline development, juvenile zebrafish have an ovary-like gonad which contains early stage oocytes [5]. If these early stage oocytes continue to mature, zebrafish will develop into females. On the contrary, if these oocytes start to degenerate, then "juvenile ovary to testis" transformation can be observed, and zebrafish will develop into males [6-8]. It has been found that zebrafish sexual fate can be affected by environmental factors, such as temperature and endocrine hormones [9-11]. Additionally, some studies have also reported that the germ cell number can influence sex determination in zebrafish [12]. In their natural growing environment, zebrafish sex transition can be observed. In the laboratory, aromatase inhibitor can induce female zebrafish to transform into male after sexual maturity [13]. However, the molecular mechanism about sexual plasticity in adults has not been largely elusive.

DNA methylation plays important roles in gene expression and development [14-18]. Two waves of DNA methylation reprogramming occur during mammalian early embryogenesis and primordial germ cell (PGC) development [19-24]. In mouse, abrogating DNA demethylation in primordial germ cells (PGCs) can lead to infertility [25], indicating the importance of DNA methylation programming during germ cell development. In zebrafish, the DNA methylation

reprogramming during early embryogenesis has been investigated, showing

that sperm but not oocyte DNA methylome is inherited by the offspring [26, 27].

Nevertheless, the role of DNA methylation during zebrafish sex differentiation

and transition is largely unknown.

To address these issues, we systematically analyzed DNA methylome and

transcriptome during zebrafish germline development and female-to-male

transition. Our study provides the mechanism insights in understanding

zebrafish sexual development.

Results

Mapping the global DNA methylome during zebrafish germ cell

development

In order to investigate the dynamics of DNA methylome during zebrafish

germline development, we collected the samples of 9 representative stages

during PGC development and germ cell differentiation. We used a transgenic

kop-gfp-nos1-3'UTR strain to collect early PGCs [28], including 4 hours post

fertilization (4 hpf) PGCs at specification stage, 6 hpf PGCs at migration stage,

and 24/36 hpf PGCs arriving at genital ridge (Figure 1A). kop GFP expression

cannot be detected in later germline development stages, but vasa can mark

later germ cells [29-31]. Thus, we generated *vasa::egfp* transgenic strain to

trace later germ cells. We collected PGCs at 4 days post fertilization (4 dpf), 9

dpf，17 dpf from juvenile gonads, and germ cells (stage-1B oocytes [32] with

30-50 um diameter) from 35 dpf female zebrafish (denoted as 35 dF germ cells)

and germ cells (GFP positive cells with less than 10 um diameter) from 35 dpf

male zebrafish (denoted as 35 dM germ cells) (Figure 1A and S1A). Zebrafish

starts sex differentiation at 17 dpf [33]. In this study, germ cells from 4 hpf to 17

dpf stages are uniformly described as "PGCs", and 35 dF and 35 dM germ cells

are called as female germ cells (FGCs) and male germ cells (MGCs)

respectively. Micromanipulation and fluorescence-activated cell sorting (FACS)

methods were employed to isolate the germ cells (Figure S1B; see Methods).

Highly purified germ cells were used to prepare DNA methylation libraries by

low-input WGBS method [34] with at least two biological replicates (Table S1).

CpGs covered by at least 3 reads were used for subsequent analysis. Our data

show that CpG density is anti-correlated with DNA methylation level across all

stages (Figure S2A). Most of genomic elements are hypermethylated except

for 5'UTR and promoter CpG islands (CGIs) (Figure S2B). While, half of

lncRNAs [35] show middle or low methylation levels, which is different from

tRNA and rRNA (Figure S2C).

Next, we plotted average methylation levels of PGCs at all examined stages.

The global methylation levels do not substantially change for the early PGCs

from 4 hpf to 24 hpf which are similar to that of midblastula (MBT) embryos

(Figure 1B). This observation is similar to the results from the previous study

[36]. Intriguingly, our data further show that after PGCs arrive at germinal ridge,

the average methylation levels (MLs) of PGCs at 36 hpf start to decrease and

reach the lowest point at 9 dpf (ML=81%) which is comparable to that of oocytes

(ML=79%) and ovaries (ML=77%). The hierarchical clustering analysis of DNA

methylomes also shows that 9 dpf PGCs, oocytes and ovaries cluster together,

separated from the other stages (Figure 1C). This is different from previous

study reporting that PGCs migrate into gonads maintain the stable methylome

pattern [37]. Furthermore, the methylation level of 35 dM increases dramatically

to a similar level of sperm. On the contrary, the methylation level of 35 dF is

lower than that of 35 dM (Figure 1B). In summary, a unique reprogramming of

DNA methylome occurs during zebrafish germline development, which is

distinct from that in mammals [38].

Transcription landscape in zebrafish germ cells

Besides DNA methylome, we also performed RNA-seq in zebrafish germ cells

(Table S2). Firstly, we identified the stage-specific expressed genes during

germline development (Figure S3A and Table S3). Gene ontology (GO)

enrichment analysis shows that 6 hpf specifically expressed genes are enriched

in the categories of somitogenesis and germ cell migration (Figure S3A), such as *G-protein signaling 14a* (*rgs14a*) (Figure S3B). It is consistent with the fact that germ cells at 6 hpf undergo the process of migration. Our data also show that pluripotency genes such as *zic3* and *klf17* are especially expressed at the stages of 4 hpf and 6 hpf (Figure 1D and S3B), suggesting that PGCs at 4 hpf and 6 hpf show more stemness. A significant number of genes are expressed in both 35dF and 35dM, but many genes are only highly expressed in 35dF or 35dM respectively. Further GO analyses show that 35 dF specifically expressed genes are enriched in the processes of egg coat formation and binding of sperm to zona pellucida (Figure S3A), such as *zp2.2* (Figure 1D and S3B), reflecting the character of oocytes. Notably, *zygote arrest 1-like* (*zar1l*), *pou5f3* and *carbonic anhydrase 15b* (*ca15b)* are only expressed in 35 dF but not in 35 dM. These genes may function specifically in female germ cells (Figure S3B). The stage of 35dM specifically expressed genes are enriched in cilium assembly (e.g., *daw1*) (Figure 1D; Figure S3A and B), which is in line with the fact that the flagellum of sperm is a modified cilium. In addition, germ cell genes, such as *piwil* and *tdrd* family genes, show limited expression in early PGCs but highly express at the later stages Supplementary information，Figure S3B. However, genes like *vasa* and *dead end 1(dnd1)*, keep highly expressed throughout the whole examined development period (Figure 1D). Taken together, gene expression can reflect the general cellular processes of zebrafish germline development (Figure 1D).

The role of methylation reprogramming during germ cell development

Promoters represent a vital regulatory level for transcription initiation [39]. DNA methylation in promoters can regulate gene expression [40]. Therefore, we focused on the DNA methylation dynamics at promoter regions. We performed clustering analysis based on differentially methylated promoters (DMPs) (Figure 2A). Our data show that the stage-specific hypomethylated promoters at the 9 dpf stage compared to the other developing stage are also

hypomethylated in oocytes/ovaries, and the stage-specific hypermethylated

promoters at the 9 dpf stage are hypermethylated in oocytes/ovaries as well.

These results indicate that the DNA methylation pattern of 9 dpf PGCs at

promoters is similar to that of the oocytes/ovaries.

Further, GO enrichment analysis shows that genes with hypomethylated

promoters in oocytes and 9 dpf PGCs are enriched in developmental growth,

hormone mediated signal pathway and cell motion (Figure 2A and Table S4),

such as *myoc* (Figure S4A), and *dcn* (Figure 2B) which regulates cell adhesion

and migration by binding to ECM molecules [41]. The result is consistent with

the observation that the PGCs migrate and then start to proliferate in gonads

between 6 hpf and 9 dpf. Our data show that stage-specific hypermethylated

promoters in 9 dpf PGCs are enriched in the reproductive process, germ cell

development (Figure 2A and Table S4). These promoters become

hypomethylated states at 17 dpf, 35 dF and 35 dM. The representative

examples of germ cell development related genes *dazl*, *tdrd1* and *ta*

respectively are shown in figures (Figure 2C; Figure S4B and C). We speculate

that these 9 dpf hypermethylated-promoter genes could help PGCs avoid the

precocious differentiation.

To examine the potential regulation of promoter methylation on gene

expression during zebrafish PGC development, we also analyzed the

relationship of differential gene expression and promoter methylation between

6 hpf and 9 dpf PGCs. Our results illustrate that gene expression usually

increases when the promoter methylation level decreases, including cell

migration, growth and hormone-mediated genes (e.g., *prlra*) (Figure 2D). By

contrast, gene expression usually decreases when the promoter methylation

level increases, including pluripotency gene (e.g., *nanog*) (Figure 2D). These

results suggest that the dynamic of DNA methylation from 6 hpf to 9 dpf can

inhibit the pluripotency capability of PGCs and facilitate PGCs migration and

proliferation.

At 35 dpf, zebrafish has determined its sex. Gonads of female zebrafish

at 35 dpf contain immature oocytes (Figure S1A) and show expression of

oocyte-related genes (Figure S3B). However, there are a large amount of DMPs

between 35 dF FGCs and oocytes, indicating that DNA methylome of 35dF

FGCs are different from that of oocytes (Figure 2A). In contrast, the methylome

pattern of 35 dM MGCs is quite similar to that of sperm (Figure 2A). These

results suggest that the DNA methylome of sperm is nearly established as early

as 35 dpf, whereas dramatic DNA methylation reprogramming is necessary for

oocytes maturation after 35 dpf.

To further elucidate the role of DNA methylation on sex determination, we

used 5-Aza-2'-deoxycytidine (inhibitors of DNA methyltransferases (DNMTs))

to block DNA methylation during zebrafish germ cell development. We utilized

three dose-response 5-Aza-2'-deoxycytidine (5-Aza-dC: 0.005mg/g, 0.01mg/g,

0.015mg/g) to treat *vasa::egfp* transgenic strain at 9 dpf and raised the treated

juvenile fish till 35 dpf, 60 dpf and 90 dpf. Then we checked the male/female

ratio in each group (see Methods). The male sex ratio in treated group from 35

dpf to 90 dpf is lower than that in the control (Figure 2E and Table S5), indicating

that fish treated with DNA methylation inhibitor are prefer to develop into

females, which is consistent with recent study [42]. Therefore, these results

suggested that DNA methylation plays a role during sex determination.

Tet3 involves in PGC development

Our data show there are more than ten thousand of demethylated differentially

methylated regions (DMRs) between 36 hpf stage and 9 dpf stage (Figure S5A).

Tet proteins were proposed to be involved in the DNA demethylation in

mammalian embryos [43, 44] and zebrafish phylotypic-stage embryos [45]. Our

data shows that *tet1* and *tet3* but not *tet2* were highly expressed in PGCs at the

36 hpf and 9 dpf stages (Figure 3A). To better define the role of *tet* during germ

cell development, we used the morpholino (MO) knock down approach to target

tet1 and *tet3* in *vasa::egfp* transgenic embryos and validate by western blot

(Figure S5B). The majority embryos injected with double morpholino (*tet1* MO

+ *tet3* MO) showed developmental deformity and lethality (Figure S5C). *Tet1*

MO embryos did not show abnormal phenotypes. By contrast, *tet3* MO

individuals showed much weaker fluorescence signal in gonads at 5 dpf and 9

dpf compared to the control (Figure S5D). Previous studies have reported that

the germ cell number can affect sex determination in zebrafish [12]. Our results

show that the number of PGCs in *tet3* MO juveniles is significantly reduced

(Figure S5E).

Next, we introduced mutations in the zebrafish *tet3* gene by using

CRISPR/Cas9 method. The *tet3*[c1] allele is a mutation allele with an 8-bp

deletion in exon3. This deletion leads to a premature stop codon in exon3 and

the loss of catalytic domain in C terminal (Figure 3B). Then, we counted the

number of PGCs in *tet3*[c1] mutants at 5 dpf under stereoscopic microscope by

in situ hybridization with *vasa* probes (Figure 3C and D). We observed that

tet3[c1] fish contain less *vasa*-positive cells compared to the control, indicating

that interrupting *tet3* gene can reduce germ cell numbers in zebrafish. Hence,

the *Tet3* involves in the regulation of PGC development.

Female-to-male sex transition in adult zebrafish

Inhibiting aromatase activity by aromatase inhibitor can revert adult zebrafish

female-to-male [13]. This female to male sex transition process provides us a

chance to explore the underlying molecular mechanisms for adult sexual

plasticity in the vertebrate. Thus, we used aromatase inhibitor aromasin

(Exemestane) to induce female-to-male sex reversal in *vasa::egfp* zebrafish

(Figure 4A and B; See Methods). Our results show that the EGFP fluorescence

intensity within adult gonads is gradually decreased after aromasin treatment,

suggesting that the female germ cells start degeneration (Figure 4B).

Histological staining also shows that the number of female germ cells is

gradually reduced (Figure 4B). After aromasin treatment for three months, the

gonad exhibits an intermediate transition state containing both degenerated

oocytes and emerged male germ cells. We then stopped the aromasin

treatment and found that the treated female fish would develop a testis-like

gonad in the next one or two months (Figure 4B). We further mated the sex

reversed fish with wild type females. Our result shows that the sperm from sex

reversal fish is able to successfully fertilize and produce normal embryos

(Figure S6A).

We then performed RNA sequencing to investigate the dynamics of gene

expression during female-to-male sex transition (Table S2). By the time-course

gene expression clustering analysis (see Methods), we identified four dynamic

gene clusters (Figure 4C and Table S6). Our data show that the expression

level for female sexual development genes (cluster1, e.g., *bone morphogenetic*

protein 15 (*bmp15)* and *hsd3b2*) are down-regulated during the sex transition

process (Figure 4C). It is worth noting that 3_mo stage is intermediate transition

stage (Figure 4B). Genes (cluster 2) which display uniquely high expression

levels at this stage but low expression in the ovary and the sex reversed/ wild

type testis were enriched in the tissue regeneration process (e.g., *igf2b*),

immune response and localization of cell process (e.g., *cxcr4b and cxcl12a*)

(Figure 4C). We also found genes which became highly expressed and kept

high levels in the sex reversed/wild type testis (cluster3, e.g., *notch2*) were

enriched in organ morphogenesis and hormone response (Figure 4C). In the

end, male sexual development genes (cluster4, e.g., *doublesex and mab-3*

related transcription factor 1 (*dmrt1)* and *gonadal somatic cell derived factor*

(gsdf)) are up-regulated. In summary, during female-to-male transition, genes

related to female sexual development stop expression at first, then genes

related to tissue regeneration, cell migration, organ morphogenesis and

hormone response start expression at the inter transition stage, and genes

related to male and sperm generation start expression in the end.

DNA methylome reprogramming during adult zebrafish female-to-male

sex transition.

Our results have shown that the DNA methylation is tightly associated with germ

cell development in juvenile zebrafish (Figure 2). To explore the role of DNA

methylation in adult zebrafish sex transition, we sequenced DNA methylomes

of gonads during sex transition process, including wild type ovary/testis, gonads

treated by aromasin for 1 month (1_mo), 2 months (2_mo), and 3 months (3_mo)

and fully sex reversal testis (sr testis) with at least two independent biological

replicates (Figure 4A; Figure S6B and Table S1). Our data show that the global

DNA methylation level has a significant change during female-to-male sex

transitions (Figure 5A), and the methylation level of 3_mo gonads (ML= 83%)

is between that of wild type ovary (ML=77%) and testis (ML=86%) (Figure 5A

and B; Figure S6C). When the treated females sexually revert to males, the

methylation level of gonads from sex reversal fish reached to a similar level of

wild type testis (Figure 5A and B; Figure S6C). The results above indicate that

sperm from sex reversal fish have same functions as wild type sperm (Figure

S6A). Then we also sequenced DNA methylomes for sex reversal sperm and

wild type sperm. Our data show that the DNA methylome of sperm from sex

reversal fish is similar to that of sperm from wild type fish, but different from that

of oocytes (Figure 5C). For examples, *dnmt6* is low-methylated in oocyte, but

becomes high-methylated in sex reversal sperm, which is similar to the state in

wild type sperm (Figure S6D) and *sycp3l* is high-methylated in oocyte, but show

low-methylated in sex reversal sperm (Figure S6E).

Our data illustrate that several well-known sex determination genes show

differentially methylated at their promoters between ovaries and testis (Figure

S7A and B; Table S7). Then, we investigated the relationship between

transcription and methylation status for these known sex determination genes

during sex transition. For example, a well-defined sex determination gene *dmrt1*,

a key regulator of sex determination which is necessary for male sexual

development in vertebrates [46-49], has a male-specific hypomethylated

promoter and starts to be highly expressed from 3_mo onwards (Figure 5D). In

addition, *cytochrome P450 superfamily member-cyp19a1a*, a critical gene in

promoting ovary development in fishes [50], shows a low methylation level at

the promoter region in ovaries, but gradually reprograms to a high methylation level at the later stages (Figure S7C).

We next compared the DNA methylation level dynamics between adult sex transition process and juvenile germline development in zebrafish. The dynamics of methylation level during female-to-male sex transition is very similar to that from 9 dpf germ cells to testis during normal development (Figure S7D and E). Perhaps, the methylome of female gonads similar to the 9 dpf-like bipotential DNA methylation pattern is the reason why the female gonad is able to transform into the male gonad.

Furthermore, we wanted to know the role of DNA methylome during zebrafish sex reversal process. We added 5-Aza-dC (0.035mg/g) to examine its effect on aromasin-induced female-to-male sex transition (see Methods). After aromasin treatment for three months, 31% of total female fish with only–aromasin-treatment start transiting to male as these fish have both ovary and testis, while none of females with both 5-Aza-dC and aromasin treatment start the transition. Furthermore, we raised the treated fish in the recirculation system till 5 months after the initial treatment began (fish sex reversed stage) with no more drug treatment and found that 33% of total females transited into males for the only-aromasin-treatment group, whereas merely 5% of females transited into males for both 5-Aza-dC and aromasin group (Figure 5E and Table S5).

In summary, our results indicate that blocking DNA methylation can prevent sex reversal in zebrafish, suggesting that DNA methylation dynamics is essential for sex reversal in adult zebrafish.

Discussion

Genome-wide erasure of DNA methylation during the germline development has been documented in mammals [21-24, 38], but it was limitedly reported in zebrafish germ cells. A recent study generated low depths of DNA methylome of zebrafish PGCs after 24 hpf [37]. Low depths and coverage of the DNA methylomes block the understanding of germ cell development. They

report that zebrafish did not experience significant reprogramming during PGC development. In contrast, we generated a relatively higher depths of DNA methylomes for zebrafish including early PGCs and late germ cells using modified library generation methods at single-base resolution [34]. Our results show that zebrafish PGCs undergo significant DNA methylation reprogramming during the germline development. More importantly, we further reveal that the methylome reset to an oocyte/ovary-like pattern at 9 dpf stage, and illustrate that zebrafish PGCs at 9 dpf have the bipotential to differentiate into male or female germ cells. It is excited for us to find that a similar methylation reprogramming during normal sexual development from 9 dpf PGC to male testis can be observed during aromasin-induced female ovary to male testis transition. Thus, we infer that the 9 dpf-like bipotential DNA methylation pattern in female gonads is the reason why the female gonad is able to transform into the male gonad. The role of DNA methylation in sex reversal had not been revealed previously. In this study, we inhibit the DNA methylation reprogramming after aromasin treatment, and demonstrate that DNA methylation reprogramming is required for the sex reversal in zebrafish.

During artificially induced zebrafish female-to-male sex transition, our results indicated that there exists DNA methylation dynamics for vital sex determination genes including dmrt1 and cyp19a1a, which shows a similar trend in blue head wrasses and half-smooth tongue sole[51, 52]. These results suggest that DNA methylation may have a conserved regulatory role in fish sex determination.

Recent study report that H3K27me3 plays roles in turtle sex determination [53]. The interplay of DNA methylation and other epigenetic mechanisms for zebrafish sex determination is worthy of further study.

Taken together, our research provides molecular mechanism insights for understanding the mystery of sex reversal, and we revealed that the DNA methylome reprogramming play roles in zebrafish germline development and female-to-male sex transition.

385 **Competing interests:** The authors declare no competing interests.

Reference

[1] Graves JA. Sex chromosome specialization and degeneration in mammals. Cell 2006;124:901-14.

[2] Shapiro DY. Behavioral influences on gene structure and other new ideas concerning sex change in fishes. Environ. Biol. Fish 1988;23:283-97.

[3] Fishelson L. Protogynous Sex Reversal in the Fish Anthias squamipinnis (Teleostei, Anthiidae) regulated by the Presence or Absence of a Male Fish. nature 1970;227:90-1.

[4] Walther T, Heinz W. Meiotic chromosomes and stages of sex chromosome evolution in fish: zebrafish, platyfish and guppy. Chromosome Research 2001;9:659-72.

[5] Takahashi H. Juvenile Hermaphroditism in the Zebrafish, Brachydanio rerio. Bull Fac Fish Hokkaido Univ 1977;28:57-65.

[6] Maack G, Segner H. Morphological development of the gonads in zebrafish. Journal of Fish Biology 2003;62:895-906.

[7] Wang XG, Bartfai R, Sleptsova-Freidrich I, Orban L. The timing and extent of 'juvenile ovary' phase are highly variable during zebrafish testis differentiation. Journal of Fish Biology 2007;70:33-44.

[8] Orban L, Sreenivasan R, Olsson PE. Long and winding roads: testis differentiation in zebrafish. Mol Cell Endocrinol 2009;312:35-41.

[9] Baroiller J, D'Cotta H. Environment and sex determination in farmed fish. Comparative Biochemistry and Physiology.part C: Toxicology & Pharmacology 2001;130:399-409.

[10] Örn S, Holbech H, Madsen TH, Norrgren L, Petersen GI. Gonad development and vitellogenin production in zebrafish (Danio rerio) exposed to ethinylestradiol and methyltestosterone. Aquatic Toxicology 2003;65:397-411.

[11] Abozaid H, Wessels S, Hörstgen-Schwark G. Elevated Temperature Applied during Gonadal Transformation Leads to Male Bias in Zebrafish (Danio rerio). Sexual Development 2012;6:201-9.

[12] Tzung KW, Goto R, Saju JM, Sreenivasan R, Saito T, Arai K, et al. Early depletion of primordial germ cells in zebrafish promotes testis formation. Stem Cell Reports 2015;4:61-73.

[13] Takatsu K, Miyaoku K, Roy SR, Murono Y, Sago T, Itagaki H, et al. Induction of female-to-male sex change in adult zebrafish by aromatase inhibitor treatment. Scientific Reports 2013;3:3400.

[14] Reik W. Stability and flexibility of epigenetic gene regulation in mammalian development. Nature 2007;447:425-32.

[15] Bird A. DNA methylation patterns and epigenetic memory. Genes Dev 2002;16:6-21.

[16] M. Azim Surani RK, Nicholas D . Allen. Genome imprinting and development in the mouse. Development Supplement 1990;108:89-98.

[17] Calarco JP, Borges F, Donoghue MT, Van Ex F, Jullien PE, Lopes T, et al. Reprogramming of DNA methylation in pollen guides epigenetic inheritance via small RNA. Cell 2012;151:194-205.

[18] Xie W, Barr CL, Kim A, Yue F, Lee AY, Eubanks J, et al. Base-resolution analyses of sequence and parent-of-origin dependent DNA methylation in the mouse genome. Cell 2012;148:816-31.

[19] Guo H, Zhu P, Yan L, Li R, Hu B, Lian Y, et al. The DNA methylation landscape of human early embryos. Nature 2014;511:606-10.

[20] Wang L, Zhang J, Duan J, Gao X, Zhu W, Lu X, et al. Programming and inheritance of parental DNA methylomes in mammals. Cell 2014;157:979-91.

[21] Seisenberger S, Andrews S, Krueger F, Arand J, Walter J, Santos F, et al. The dynamics of genome-wide DNA methylation reprogramming in mouse primordial germ cells. Molecular Cell 2012;48:849-62.

[22] Guo F, Yan L, Guo H, Li L, Hu B, Zhao Y, et al. The Transcriptome and DNA Methylome Landscapes of Human Primordial Germ Cells. Cell 2015;161:1437-52.

[23] Gkountela S, Zhang KX, Shafiq TA, Liao WW, Hargan-Calvopina J, Chen PY, et al. DNA Demethylation Dynamics in the Human Prenatal Germline. Cell 2015;161:1425-36.

[24] Tang WW, Dietmann S, Irie N, Leitch HG, Floros VI, Bradshaw CR, et al. A Unique Gene Regulatory Network Resets the Human Germline Epigenome for Development. Cell 2015;161:1453-67.

[25] Hargan-Calvopina J, Taylor S, Cook H, Hu Z, Lee Serena A, Yen M-R, et al. Stage-Specific Demethylation in Primordial Germ Cells Safeguards against Precocious Differentiation. Developmental Cell 2016;39:75-86.

[26] Jiang L, Zhang J, Wang JJ, Wang L, Zhang L, Li G, et al. Sperm, but not oocyte, DNA methylome is inherited by zebrafish early embryos. Cell 2013;153:773-84.

[27] Potok ME, Nix DA, Parnell TJ, Cairns BR. Reprogramming the Maternal Zebrafish Genome after Fertilization to Match the Paternal Methylation Pattern. Cell 2013;153:759-72.

[28] Blaser H, Eisenbeiss S, Neumann M, Reichman-Fried M, Thisse B, Thisse C, et al. Transition from non-motile behaviour to directed migration during early PGC development in zebrafish. Journal of Cell Science 2005;118:4027-38.

[29] Yoon C, Kawakami K, Hopkins N. Zebrafish vasa homologue RNA is localized to the cleavage planes of 2- and 4-cell-stage embryos and is expressed in the primordial germ cells. Development 1997;124:3157-65.

[30] Olsen LC, Aasland R, Fjose A. A vasa-like gene in zebrafish identifies putative primordial germ cells. Mechanisms of development 1997;66:95-105.

[31] Krovel AV, Olsen LC. Sexual dimorphic expression pattern of a splice variant of zebrafish vasa during gonadal development. Dev Biol 2004;271:190-7.

[32] Selman. K, Wallace. RA, Sarka. A, Qi X. Stages of oocyte development in the zebrafish, Brachydanio rerio. journal of morphology 1993;218:203–24.

[33] Sun D, Zhang Y, Wang C, Hua X, Zhang XA, Yan J. Sox9-related signaling controls zebrafish juvenile ovary-testis transformation. Cell Death and Disease 2013;4:e930.

[34] Li G, Yu Y, Fan Y, Li C, Xu X, Duan J, et al. Genome wide abnormal DNA methylome of human blastocyst in assisted reproductive technology. Journal Genetics Genomics 2017;44:475-81.

[35] Ulitsky I, Shkumatava A, Jan CH, Sive H, Bartel DP. Conserved function of lincRNAs in vertebrate embryonic development despite rapid sequence evolution. Cell 2011;147:1537-50.

[36] Skvortsova K, Tarbashevich K, Stehling M, Lister R, Irimia M, Raz E, et al. Retention of paternal DNA methylome in the developing zebrafish germline. Nature Communications 2019;10.

[37] Ortega-Recalde O, Day RC, Gemmell NJ, Hore TA. Zebrafish preserve global germline DNA methylation while sex-linked rDNA is amplified and demethylated during feminisation. Nat Commun 2019;10:3053.

[38] von Meyenn F, Reik W. Forget the Parents: Epigenetic Reprogramming in Human Germ Cells. Cell 2015;161:1248-51.

[39] Nepal C, Hadzhiev Y, Previti C, Haberle V, Li N, Takahashi H, et al. Dynamic regulation of the transcription initiation landscape at single nucleotide resolution during vertebrate embryogenesis. Genome Res 2013;23:1938-50.

[40] Xie W, Schultz MD, Lister R, Hou Z, Rajagopal N, Ray P, et al. Epigenomic analysis of multilineage differentiation of human embryonic stem cells. Cell 2013;153:1134-48.

[41] Lala PK, Nandi P. Mechanisms of trophoblast migration, endometrial angiogenesis in preeclampsia: The role of decorin. Cell Adh Migr 2016;10:111-25.

[42] Ribas L, Vanezis K, Imues MA, Piferrer F. Treatment with a DNA methyltransferase inhibitor

feminizes zebrafish and induces long-term expression changes in the gonads. Epigenetics Chromatin

2017;10:59.

[43] Guo F, Li X, Liang D, Li T, Zhu P, Guo H, et al. Active and passive demethylation of male and

female pronuclear DNA in the mammalian zygote. Cell Stem Cell 2014;15:447-59.

[44] Gu TP, Guo F, Yang H, Wu HP, Xu GF, Liu W, et al. The role of Tet3 DNA dioxygenase in epigenetic

reprogramming by oocytes. Nature 2011;477:606-10.

[45] Bogdanović O, Smits AH, de la Calle Mustienes E, Tena JJ, Ford E, Williams R, et al. Active DNA

demethylation at enhancers during the vertebrate phylotypic period. Nature Genetics 2016;48:417-26.

[46] Smith CA, Roeszler KN, Ohnesorg T, Cummins DM, Farlie PG, Doran TJ, et al. The avian Z-linked

gene DMRT1 is required for male sex determination in the chicken. Nature 2009;461:267-71.

[47] Matson CK, Murphy MW, Sarver AL, Griswold MD, Bardwell VJ, Zarkower D. DMRT1 prevents

female reprogramming in the postnatal mammalian testis. Nature 2011;476:101-4.

[48] Nanda I, Kondo M, Hornung U, Asakawa S, Winkler C, Shimizu A, et al. A duplicated copy of

DMRT1 in the sex-determining region of the Y chromosome of the medaka, Oryzias latipes. Proceedings

of the National Academy of Sciences of the United States of America, 2002;99:11778-83.

[49] Webster KA, Schach U, Ordaz A, Steinfeld JS, Draper BW, Siegfried KR. Dmrt1 is necessary for

male sexual development in zebrafish. Developmental Biology 2017;422:33-46.

[50] Guiguen Y, Fostier A, Piferrer F, Chang C-F. Ovarian aromatase and estrogens: A pivotal role for

gonadal sex differentiation and sex change in fish. General and Comparative Endocrinology

2010;165:352-66.

[51] Todd E, Ortega-Recalde O, Liu H, Lamm M, Rutherford K, Cross H, et al. Stress, novel sex genes,

and epigenetic reprogramming orchestrate socially controlled sex change. Science Advances

2019;5:eaaw7006.

[52] Shao C, Li Q, Chen S, Zhang P, Lian J, Hu Q, et al. Epigenetic modification and inheritance in

sexual reversal of fish. Genome Res 2014;24:604-15.

[53] Ge C, Ye J, Weber C, Sun W, Zhang H, Zhou Y, et al. The histone demethylase KDM6B regulates

temperature-dependent sex determination in a turtle species. science 2018;360:645-8.

[54] Li G, Yu Y, Fan Y, Li C, Xu X, Duan J, et al. Genome-wide abnormal DNA methylome of human

blastocyst in assisted reproductive technology. Journal of Genetics and Genomics 2017;44:475-81.

[55] Hruscha A, Krawitz P, Rechenberg A, Heinrich V, Hecht J, Haass C, et al. Efficient CRISPR/Cas9

genome editing with low off-target effects in zebrafish. Development 2013;140:4982-7.

[56] Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.

Bioinformatics 2014;30:2114-20.

[57] Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq

applications. Bioinformatics 2011;27:1571-2.

[58] Akalin A, Kormaksson M, Li S, Garrett-Bakelman F, Figueroa M, Melnick A, et al. MethylKit: A

comprehensive R package for the analysis of genome-wide DNA methylation profiles. Genome biology

2012;13:R87.

[59] Sun D, Xi Y, Rodriguez B, Park HJ, Tong P, Meong M, et al. MOABS: Model based analysis of

bisulfite sequencing data. Genome biology 2014;15:R38.

[60] Hansen K, Langmead B, Irizarry R. BSmooth: From whole genome bisulfite sequencing reads to

differentially methylated regions. Genome biology 2012;13:R83.

[61] Dobin A, Davis C, Schlesinger F, Drenkow J, Zaleski C, Batut P, et al. STAR: ultrafast universal

RNA-seq aligner. Bioinformatics 2012;29:15-21.

[62] Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using

DAVID bioinformatics resources. Nature Protocols 2008;4:44-57.

[63] Tripathi S, Pohl Marie O, Zhou Y, Rodriguez A, Wang G, Stein David A, et al. Meta- and Orthogonal

Integration of Influenza "oMICs" Data Defines a Role for UBR4 in Virus Budding. Cell Host & Microbe

2015;18:723-35.

Figure 1. Dynamics of DNA methylation during zebrafish germline development. A. The schematic diagram for zebrafish germline developmental stages in this study. Two transgenic strains were used to track the development of germ cells in zebrafish. Dotted boxes indicate the major developmental processes of germ cells at different stages. **B.** Global DNA methylation levels during zebrafish germ cell development. DNA methylation data of zebrafish sperm, oocyte, MBT and testis stages are from public data (GSE44075). Green triangle means the PGC samples, red triangle means female samples, blue triangle means male samples, grey triangle means embryonic samples. **C.** A dendrogram showing the clustering of DNA methylomes during zebrafish germ cell development (500 bp bin for each unit). **D.** A brief summary of key gene

553 expressions and key biological processes during zebrafish germline

554 development.

556
557

Figure 2. Methylome reprogramming during zebrafish germ cell development. A. DNA methylation level heatmap for differentially methylated promoters (DMPs). These DMPs are classified into groups by Partitioning Around Medoid (PAM) method. Corresponding GO biological process terms are shown in the right panel and the color represents the enrichment stastical significance. Promoters (Transcription Start Site, TSS±300 bp; n=385). **B.**

Snapshots for DNA methylation at 9 dpf-specific hypomethylated promoter gene *dcn* (cytokine-mediated signal pathway related gene). Dynamic regions around promoters are highlighted in grey. DNA methylation level from 0 to 1. Each vertical line represents one CpG site. **C.** Snapshot for methylation at 9 dpf-specific hypermethylated promoter gene *dazl* (germline gene). **D.** Heatmaps of DNA methylation level (left) and gene expression fold change (right) for the genes with DMPs in 6 hpf and 9 dpf. Only genes with |log2 (fold change)|>1 are included. **E.** Sex ratio within zebrafish population after treatment with different 5-Aza-dC concentration, including 0.005mg/g (control n=124; 35 dpf n=47; 60 dpf n=121; 90 dpf n=112), 0.01mg/g (control n=86; 60 dpf n=90; 90 dpf n=82), 0.015mg/g (control n=60; 60 dpf n=64; 90 dpf n=56). The error bar is mean±s.e.m., Statistical significance were calculated by one-way ANOVA.

Figure 3. The function of *tet3^{c1}* mutant during zebrafish germ cell development. A. Histogram of the gene expression levels of *tet1/tet2/tet3* in PGCs. Gene expression level is averaged from biological replicates. **B.** Schematic diagram shows CRISPR/Cas9 editing in zebrafish *tet3* gene, which delete 8 bps in exon3. **C.** Expression of *vasa* mRNA in situ hybridization for wild type and *tet3^{c1}* mutant fish at the 5 dpf stage. **D.** The PGC number of 5 dpf was counted under stereoscopic microscope (wild type n=43; tet3 n=27). The error bar represents mean±s.d.. The p-value was calculated by t test. * means p < 0.05. ** means p < 0.01, *** means p < 0.001.

Figure 4. Female-to-male sex transition in the adult zebrafish. A. The

schematic diagram for zebrafish sex reversal samples in this study. *vasa::egfp*

transgenic strain was used to induce sex reversal by aromasin treatment. The

samples in this figure include wild type ovary, gonads from zebrafish with one

month (1_mo), two month (2_mo), and three month (3_mo) drug treatment, sex

reversal testis and wild type testis. **B.** Observation of gonad morphology change

in *Tg* (*vasa::egfp*) wild type female and post aromasin treatment fish per month.

The image of gonads under bright filed, GFP fluorescence and HE staining. The

scale bar indicates 100 μm. **C.** Time-course clustering analysis of gene

expression patterns across zebrafish sex reversal process. In the four distinct

expression groups, the upper left panel of each group shows the gene

expression pattern, the upper right panel shows the GO enrichment results, and

the representative genes expression dynamics is shown in the lower panel.

Figure 5. DNA methylation reprogramming during zebrafish female-to-male transition. A. The global DNA methylation reprogramming during zebrafish sex reversal. The methylation level is calculated as the average across all covered CpGs. The error bar represents mean±s.d.. The samples in this figure include wild type ovary, gonads from zebrafish with one month (1_mo), two month (2_mo), and three month (3_mo) drug treatment, sex reversal testis and wild type testis. **B.** The dynamic changes for DMRs between zebrafish ovary and testis during sex transition process. The △ methylation heatmap displays the difference of DNA methylation level between chosen two

stages. **C.** The correlation analysis of CpG methylation level in sex reversal sperm (sr sperm), wild type sperm (wt sperm) and oocyte. Pearson correlation coefficients r is also shown in the figure. **D.** Snapshots of DNA methylation and gene expression tracks for sex determination gene *dmrt1* in ovary, 1_mo, 2_mo, 3_mo, sr testis and wt testis. **E.** Female sex ratio within populations after drug treatment for the 3_mo stage (aromasin n=36; aromasin+5-Aza-dC n=47) and 5_mo stage (sex reversed stage, aromasin n=36; aromasin+5-Aza-dC n=38). 3 independent experiments for each drug treatment group. The error bar is mean±s.e.m., the statistical significance was calculated by unpaired two-sided t test.

Supplementary Figure Legends

Supplementary Figure S1. PGC and germ cell purification. A. Fluorescent

photography and HE staining for immature ovary and immature testis at 35 dpf

stage (approximately equal length of males and females around 1.7±0.1cm).

The left panel represents the fluorescence intensity of ovary and testis at 35

dpf; the right panel represents the HE staining of ovary and testis at 35 dpf. The

scale bar is 50 µm. **B.** Sorting GFP positive cells by FACS. Left panel shows

GFP negative cells for control fishes. Right panel shows GFP positive cells at

each germline time point for transgenic fishes.

Supplementary Figure S2. Global analysis of DNA methylation at different

germline developmental stages in zebrafish. A. Boxplots show the

correlation between CpG density and methylation level around TSS±2Kb

regions. Boxes and whiskers represent the 25th/50th/75th percentiles and 1.5X

the interquartile range, respectively. X axis means CpG counts. **B and C.** The

Proportion of CpGs with high methylation level (ML ≥0.75), middle methylation

level (0.25 < ML < 0.75) and low methylation level (ML ≤0.25) for indicated

genomic features: 5' untranslated regions (UTRs), exons, introns, 3'UTRs,

gene body, intergenic regions and CpG islands (CGIs) are shown in **B** and

repeat elements, noncoding RNAs are shown in **C**.

A

	GO terms	p value
1	somiteogenesis	3.4e-13
	mRNA splicing, via spliceosome	1.3e-12
	cell migration involved in gastrulation	5.0e-7
	notch signaling pathway	8.8e-7
	germ cell migration	1.1e-2
2	transcription, DNA-templated	5.5e-10
	multicelluar organism development	1.7e-6
	brain development	1.3e-5
3	oxidation-reduction process	8.8e-4
	epidermal growth factor receptor signaling pathway	1.4e-2
	protein folding	2.4e-2
4	protein transport	4.5e-14
	cell cycle	2.7e-8
	methylation	1.7e-7
	egg coat formation	1.1e-5
	binding of sperm to zona pellucida	1.1e-5
	fatty acid metabolic process	1.2e-4
5	translation	1.0e-27
	DNA replication	4.6e-9
	translational initiation	1.2e-7
	maturation of SSU-rRNA from tricistronic rRNA transcript	1.8e-6
	DNA repair	5.6e-7
	cell division	6.9e-4
6	regulation of transcription, DNA-templed	1.6e-11
	steroid hormone mediated signaling pathway	1.1e-6
	cilium assembly	3.4e-6
	phosphorylation	7.1e-5
	cell differentiation	1.2e-4
	axon guidance	3.6e-4

B

Supplementary Figure S3. Transcription landscapes of zebrafish germ cells. A. Heatmap for stage specific gene expression from 4 hpf PGCs to 35 dpf germ cells. 6 clusters were identified by k-means method. 1253, 952, 989, 3191, 2717, 1902 genes for each cluster. The right panels show the corresponding cluster-enriched GO biological process terms by using DAVID. P-values are also shown. **B.** Barplots for the gene expression levels of germline-related genes at different stages. Gene expression levels are averaged from all biological replicates.

Supplementary Figure S4. DNA methylation dynamics at promoter

regions across zebrafish germline development. A-C. Snapshots for the

DNA methylation of 9 dpf-specific hypomethylated promoter gene *myoc* **A** and

9 dpf-specific hypermethylated promoter gene *tdrd1* **B** and *ta* **C**. Dynamic

regions around promoters are highlighted in grey. DNA methylation level from

0 to 1. Each vertical line represents one CpG site.

Supplementary Figure S5. The function of *tet3* morpholino in zebrafish germ cell development. A. Boxplots of CpG methylation level in differentially methylated regions (DMRs) identified by stage adjacent pairwise comparisons. The hypermethylated stage is denoted as red box and the hypomethylated stage is denoted as blue box, grey boxes stand for other stages. Boxes and whiskers represent the 25th/50th/75th percentiles and 1.5X the interquartile range, respectively. n represents the number of DMRs. **B**. The western blot for control larvas, control MO larvas and *tet3* MO larvas at 3 dpf. Tubulin is the loading control for western blot. **C.** The images for control larvas, *tet1* MO larvas, *tet3* MO larvas and *tet1/3* MO larvas at 3 dpf under microscopy. **D.** The fluorescence intensity for wild type and *tet3* MO germ cells at 5 dpf and 9 dpf (red box). **E.** Dot plot showing changes in the PGC number at 5 dpf (control

767 n=22; tet3 MO n=27) and 9 dpf (control n= 22 tet3 MO n=23), which was
768 counted by the squash method. The error bar represents mean±s.d.. The p-
769 value was calculated by t test. * means p < 0.05. ** means p < 0.01, *** means
770 p < 0.001.
771
772

Supplementary Figure S6. DNA methylation dynamics for zebrafish sex reversal samples. A. The pictures for embryos generated by sex reversal fish mating with wild type female. **B.** Hierarchical clustering analysis according to the DNA methylation pattern for the biological replicates of ovary, 1_mo, 2_mo, 3_mo, sr testis and testis. sr means sex reversal. **C.** Bimodal distribution of CpG methylation level across genome for zebrafish at the indicated stages: ovary, 1_mo, 2_mo and 3_mo drug treated gonad, sr testis and testis. **D** and **E.** Snapshots of DNA methylation tracks for genes with DMPs among oocyte, wt sperm and sr sperm, including *dnmt6* **D** and *sycp3l* **E.**

Supplementary Figure S7. The relationship between DNA methylation and gene expression during zebrafish sex transition. A. DMPs between testis and ovary. GO terms for the testis-specific hypomethylated and ovary-specific hypomethylated DMPs are shown in the right panel. Promoters (TSS±1000 bp). **B.** Heatmap shows DMPs between ovary and testis for key sex determination genes. **C.** Snapshots of DNA methylation and gene expression tracks for sex determination gene *cyp19a1a* in ovary, 1_mo, 2_mo and 3_mo drug treated gonad, sr testis and testis. **D.** The comparison of DNA methylation reprogramming during sex transition process and germline development process in zebrafish. **E.** The heatmap shows the dynamic changes of DMPs between 9 dpf PGCs and testis during sex transition process (DMPs, n=574).

SUPPLEMENTARY TABLES (tables separately in Excel format)

Table 1. DNA methylation data summary for zebrafish germline and sex reversal

samples

Table 2. Transcriptome data summary for zebrafish germline and sex reversal samples

Table 3. Go terms for stage specific gene in zebrafish germ cells

Table 4. Go terms for differentially methylated promoters (DMPs) in zebrafish germline

Table 5. Sex ratio summary for drug treatment fish

Table 6. Go terms for time-course clustering genes during zebrafish sex transition

Table 7. Go terms for DMPs between testis and ovary