1 **Title:** Capillary forces drive pioneer transcription factor-mediated DNA
2 condensation
3

Thomas Quail[1,2,3,4], Stefan Golfier[1,2,3,4], Maria Elsner[1,2,3,4], Keisuke Ishihara[1,2,3,4],

Frank Jülicher[2,3,4], Jan Brugués*[1,2,3,4]

Affiliations:

[1]Max Planck Institute of Molecular Cell Biology and Genetics, 01307 Dresden, Germany
[2]Max Planck Institute for the Physics of Complex Systems, 01187 Dresden, Germany
[3]Center for Systems Biology Dresden, 01307 Dresden, Germany
[4]Cluster of Excellence Physics of Life, TU Dresden, 01062 Dresden, Germany
*correspondence: brugues@mpi-cbg.de

Summary: Capillary forces are driven by interactions between liquids and

surfaces[1]. These forces are crucial for many processes ranging from biology and

physics to engineering, such as the motion of aquatic insects on the surface of

water[2], modulation of the material properties of spider silk[3,4], and self-assembly

of small objects and microstructures[5]. Recent studies have shown that cells

assemble biomolecular condensates in a manner similar to phase separation[6]. In

the nucleus, these condensates are thought to drive transcription[7-10],

heterochromatin formation[11,12], nucleolus assembly[13,14], and DNA repair[15,16].

Here, we test if the interaction between liquid-like condensates and chromatin

generates capillary forces, which might play a role in bringing distant regulatory

elements of DNA together, a key step in transcriptional regulation. We combine

quantitative microscopy, *in vitro* reconstitution, and theory to show that the

pioneer transcription factor FoxA1 mediates the condensation of a DNA-protein

phase via a mesoscopic first-order phase transition. Surprisingly, after

nucleation, capillary forces drive growth of this phase by pulling non-condensed

DNA. Altering the tension on the DNA strand enlarges or dissolves the

condensates, revealing their mechanosensitive nature. These findings show that

DNA condensation mediated by transcription factors could bring distant regions

of DNA in close proximity using capillary forces. Our work suggests that the

physics of DNA and protein condensation driven by capillary forces provides a

general regulatory principle for chromatin organization.

Main text: To investigate how transcription factors physically organize DNA, we attached linearized λ-phage DNA to a functionalized coverslip via biotin-streptavidin linkers (Fig. 1a). We used TIRF microscopy to image the interactions between DNA and Forkhead Box Protein A1 (FoxA1), a pioneer factor that regulates tissue differentiation across a range of organisms[17,18] (Fig. 1b). Upon addition of 10 nM FoxA1-mCherry (FoxA1) to the flow chamber in the presence of DNA, FoxA1 formed protein condensates that decorated the strand (Fig. 1c). In the absence of DNA, FoxA1 did not nucleate condensates in solution at concentrations ranging from 10 to 500 nM (Extended Data Fig. 1a). The requirement for DNA in condensate formation suggests that DNA mediates protein condensation of a thin layer of FoxA1 on DNA.

In our assay, DNA molecules displayed a broad distribution of end-to-end distances (L), which are determined by the locations of the biotin-streptavidin attachment points of the λ-phage DNA molecules to the coverslip (Fig. 1c, d). This end-to-end distance governs the tension on the DNA molecule, as the contour length of each individual λ-phage DNA molecule is the same[19]. For DNA strands with end-to-end distances greater than approximately 10 μm, FoxA1 generated protein condensates on DNA (Fig. 1c). In these conditions, however, FoxA1 condensation did not influence the DNA molecule (Fig. 1c). Strikingly, for DNA molecules with end-to-end distances below 10 μm, FoxA1 pulled DNA into highly enriched condensates of FoxA1 and DNA (Fig. 1c). To quantify FoxA1-mediated DNA condensation, we measured the cross-correlation of FoxA1-DNA intensities as a function of end-to-end distance (Fig. 1d, e; Extended Data Fig. 1b). This measure ranges from roughly zero to one, where a value of zero corresponds to FoxA1-only condensation and a value of one corresponds to DNA-FoxA1 condensation. Consistent with the ability of FoxA1 to form DNA-FoxA1 condensates at low tensions (corresponding to short end-to-end distances), the cross-correlation decayed from one to zero with increasing end-to-end distances (Fig. 1e). Thus, FoxA1 can induce the formation of a DNA–protein-rich microphase in a tension-dependent manner.

The observation that FoxA1 drives DNA condensation suggests that it can
overcome the entropic tension of the DNA molecule set by the end-to-end
distance[19]. Similar to tightening an elastic band, incorporating DNA into the
condensates will increase the tension on the non-condensed DNA by reducing
the amount of available slack. This increased tension should reduce transverse
DNA fluctuations of the non-condensed DNA, consistent with a reduction of slack
in the molecule. To quantify these fluctuations, we first generated time-averaged
projections from movies of DNA molecules with and without FoxA1. These
projections are single images that display the area explored by the molecule due
to its motion in a given time period (Fig. 1d, Extended Data Fig. 1c). From these
projections, we extracted line profiles of the intensities of non-condensed DNA in
the direction perpendicular to the long axis of the strand. Finally, we used
Gaussian fits to these line profiles to define the DNA envelope width (Extended
Data Fig. 1c). In buffer, the DNA envelope width decreased as a function of end-
to-end distance, consistent with the corresponding increase of DNA strand
tension for increasing end-to-end distances[19] (Fig. 1f). However, in the presence
of FoxA1, the envelope width remained constant for all end-to-end distances as
FoxA1 pulled DNA into one or more condensates. The magnitude of the DNA
envelope width was significantly lower than in buffer conditions for all end-to-
end distances (Fig. 1f). This result suggests that FoxA1-DNA condensates
generate forces that can overcome the entropic tension of the non-condensed
DNA and buffer its tension.
90
The observation that FoxA1 can mediate DNA condensation suggests that it
could bridge distant DNA strands. To investigate this possibility, we examined
DNA molecules that were bound to the same streptavidin molecule at one end
(Fig. 1g, Extended Data Fig. 1d). In the absence of FoxA1, these DNA molecules
form a v-shaped morphology and fluctuate independently of one another. Upon
addition of FoxA1, however, we observed that the two independent strands
zipped together, generating a y-shaped morphology as the condensation of
FoxA1 increased over time (Fig. 1g, Extended Data Fig. 1d). Taken together, these
data demonstrate that FoxA1 can physically bridge DNA strands in both *cis* and
trans.

Two mechanisms can be postulated to explain FoxA1-mediated DNA condensation in our experiments: (i) direct cross-linking via the multiple DNA-binding activities of FoxA1[20] or (ii) weak protein-protein interactions driven by disordered regions of FoxA1. Structurally, FoxA1 consists of a winged helix-turn-helix DNA-binding domain and two mostly disordered domains corresponding to the N and C termini[20,21]. The DNA-binding domain contains a sequence-specific binding region composed of three alpha helices and a non-sequence-specific binding region composed of two wings. Point mutations in the sequence-specific DNA-binding region (NH-FoxA1[20]) had virtually no influence on DNA condensation activity (Fig. 2a). Although the presence of two point mutations in one of the non-sequence-specific DNA-binding regions (RR-FoxA1[20]) partially inhibited FoxA1 localization to the strand (Fig. 2b), this mutant still condensed DNA. In this case, condensation occurred on a time scale of minutes rather than seconds (as in WT-FoxA1), which can be explained by the delay in condensing sufficient RR-FoxA1 to the strand. These data suggest that non-sequence-specific binding drives the localization of FoxA1 to DNA but does not mediate DNA condensation through cross-linking. Furthermore, the sequence-specific binding domain of FoxA1 is dispensable for its localization to DNA *in vitro.* To probe whether FoxA1 protein-protein interactions through disordered domains mediate DNA condensation, we truncated both the N and C termini of FoxA1. Although ΔN-FoxA1 retained DNA condensation activity (Fig. 2c), truncating the disordered C terminus of FoxA1 inhibited DNA condensation activity (Fig. 2d). Thus, non-sequence-specific binding drives FoxA1 localization to DNA, and the disordered C terminus of FoxA1 drives DNA condensation.

Our results show that the disordered C terminus of FoxA1 is functionally required for FoxA1-mediated DNA condensation. Disordered regions of proteins influence condensate formation via weak protein-protein interactions[22]. Thus, our results support the hypothesis that FoxA1 condenses onto DNA to generate a DNA–protein-rich condensate via weak protein-protein interactions. This condensate phase nucleates only if the capillary forces it generates are larger than the entropic tension on the strand. To explore this possibility, we developed

a theoretical description based on the thermodynamics of a semi-flexible
polymer partially forming a liquid-like condensate. In the present case, the semi-
flexible polymer is DNA and the condensation is mediated by the transcription
factor. The free energy of this process contains volume and surface
contributions, and the capillary work to pull DNA into the condensate (Fig. 3a).
The volume contribution, $\mu \frac{4}{3}\pi R^3$, is proportional to the volume of the
condensate, where μ is the condensation free energy per volume and R is the
radius of the condensate. We assume that DNA is fully collapsed inside the
condensate and thus its volume is proportional to the adsorbed DNA contour
length, $V = \alpha\, L_d$. Here, α describes the packing efficiency given as DNA strand
length per cross-sectional area. The surface contribution to the free energy,
$\gamma 4\pi R^2$, is characterized by the surface tension γ of the condensate. The work
$W = \int_0^{L_d} f\, dl$ is the work that must be performed by capillary forces f to bring
an amount L_d of DNA into the condensate. The free energy for fixed end-to-end
distance L depends on the contour length of DNA in the condensate, L_d, as

149
$$F(L, L_d) = -\mu\alpha L_d + \gamma 4\pi \left(\frac{3\alpha}{4\pi}\right)^{\frac{2}{3}} L_d^{\frac{2}{3}}$$

150
$$+ \frac{\kappa L}{4}\left(\frac{l l_d}{(l-1)(l+l_d-1)} + \log\left(\frac{(l-1)^2}{(1-l_d)^4(l+l_d-1)^2}\right)\right)$$

where $l = \frac{L}{L_t}$, $l_d = \frac{L_d}{L_t}$, $\kappa = \frac{k_B T}{P}$, k_B is the Boltzmann constant, T is the
temperature, and P is the persistence length of DNA (see Supplemental
information). This free energy predicts a stochastic first-order phase transition
for the formation of DNA–protein microphases (Fig. 3b) with a distribution of
condensate sizes $P(L_d) \sim e^{-\beta F(L, L_d)}$ for fixed L (Fig. 3c). Furthermore, it predicts a
sharp transition for DNA condensation controlled by the end-to-end distance and
thus the tension on the DNA molecule. The first-order nature of this behavior
implies regimes of hysteresis and bistability. Our theory also predicts that the
capillary forces on the DNA are buffered by the condensate.

To test this theory, we first measured DNA condensate volumes and found that
they increase linearly with the amount of condensed DNA (L_d), with α=0.04 ±

0.005 μm² (Fig. 3d, Extended Data Fig. 2a, Methods). This confirms that DNA is in

a collapsed conformation inside the condensates. Next, we simultaneously fit the

predictions to the average amount of DNA contained in the condensates (L_d), and

the probability of nucleating a DNA condensate (P_{cond}) as a function of end-to-

end distance (see Methods). We calculated L_d (Fig. 3e, Extended Data Fig. 2b,

Extended Data Fig. 3) and P_{cond} (Fig. 3f, Extended Data Fig. 2c) using the

Boltzmann probability distributions (Fig. 3c) from the free energy. Our fits agree

quantitatively with the data and show that the amount of DNA in the

condensates decreases with L until a critical end-to-end distance beyond which

DNA condensates do not form. Below this critical length, we observed that the

capillary force exerted by the condensate is buffered at 0.22 pN (0.18 – 0.27 pN

CI), consistent with the theory (Fig. 3f). Finally, P_{cond} exhibits a sharp transition

at $L = 10.7$ μm (9.2 – 10.8 μm CI), in agreement with a stochastic first-order

phase transition (Fig. 3f). Close to the transition point FoxA1-mediated DNA

condensation displayed bistability. This bistability was observed in strands that

contained multiple FoxA1 condensates, but where only some of them condensed

DNA (Extended Data Fig. 4a). Our fits allowed us to extract the physical

parameters associated with condensate formation, namely the condensation free

energy per volume μ = 5.9 pN/μm² (4.9 – 8.9 pN/μm² CI) and the surface tension

γ=0.05 pN/μm (0.04 – 0.38 pN/μm CI), see Methods section. These parameters

are consistent with previous measurements for *in vitro* and *in vivo*

condensates[23,24].

Our theory and experiments show that two key parameters govern DNA–protein

condensation behavior, namely the condensation free energy per volume (μ) of

the DNA–protein phase and the surface tension of the condensate (γ). We

reasoned that different DNA-binding proteins may exhibit a range of behaviors

depending on the values of these parameters. First we investigated the

sequence-specific DNA-binding region mutant (NH-FoxA1), which also

condensed DNA but to a lesser extent (Fig. 2a). Quantitatively, we found that the

surface tension of condensates formed with this mutant was roughly unchanged

compared to WT-FoxA1, γ=0.03 pN/μm (0.02 – 0.09 pN/μm CI), but the free

energy per volume of condensation was reduced consistent with reduced DNA

binding, μ=1.7 pN/μm^2 (1.6 – 2.1 pN/μm^2 CI), Extended Data Fig. 5b, Fig. 4a. This

was also reflected in an increase in the value of the packing efficiency parameter,

with α = 0.09 ± 0.02 μm^2, meaning that these condensates packed less DNA

compared to WT-FoxA1 given the same volume of DNA-protein condensate

(Extended Data Fig. 5a). We also observed that NH-FoxA1-mediated condensates

generated a force of 0.14 pN (0.09 – 0.15 pN, CI), lower than that for WT-FoxA1.

This moderate decrease in force was also reflected in a shift of the transition

point of condensate nucleation towards a lower end-to-end distance $L_c = 8.9$ μm

(8.8 – 9.0 μm, CI), Fig. 4a. In addition, NH-FoxA1 displayed bistable DNA-protein

condensation activity in the neighborhood of the transition point (Extended Data

Fig. 4b). Next, we examined the interactions of a different transcription factor,

Tata-Box-binding protein (TBP) with DNA. We found that TBP also formed small

condensates on DNA, but did not condense DNA even at the lowest imposed DNA

tensions (Fig. 4b). Instead, TBP performed a diffusive motion along the DNA

strand (Extended Data Fig. 6). This suggests that the free energy per volume of

condensation   is   positive,   and   DNA-protein   condensation   is   not

thermodynamically favored. Lastly, we analyzed somatic linker histone H1, a

protein that is structurally similar to FoxA1. However, in contrast to FoxA1, one

of the known functions of H1 is to compact chromatin[25], so we expected H1 to

strongly condense DNA. Consistent with this, we found that H1 displayed a

dramatically   increased   DNA   condensation   activity   compared   to   FoxA1,

condensing DNA for all measured end-to-end distances (Fig. 4c). Thus, we

propose that the competition between condensation free energy per volume of

the DNA–protein phase and surface tension regulate a spectrum of DNA

condensation activities, which may be tuned by the structure of transcription

factors.

Here, we show that FoxA1 can condense DNA to form a DNA–protein-rich

microphase. The DNA–protein condensates form through a force-dependent

first-order transition for forces below a critical value. This critical force, which is

on the order of 0.2 pN, is set by capillary forces that the DNA–protein condensate

exerts on the non-condensed DNA strand. These forces are similar in magnitude

to those recently measured for DNA loop extrusion of the order of 0.2-1 pN[26-30].

We speculate that these weak forces are of key relevance to the mechanics of chromatin organization. Thus, our work suggests that capillary forces can act as a chromatin remodeler in addition to molecular motors that extrude loops and complexes that remove or displace nucleosomes (Fig. 4d).

Transcription-factor-mediated DNA–protein condensation represents a possible mechanism by which transcription factors coordinate enhancer-promoter contacts in transcriptional hubs[31]. In this context, DNA–protein condensates could act as scaffolds, recruiting transcriptional co-factors. Furthermore, capillary forces generated by the condensate could also pull co-factors into the droplet (Fig. 4d). Our theoretical description reveals that these DNA–protein condensates are formed via a first-order phase transition, suggesting that they can be assembled and disassembled rapidly by changing mechanical conditions. We find that, near the transition point, assembly and disassembly of DNA–protein condensates becomes highly stochastic. This is reminiscent of the rapid dynamics associated with the initiation and cessation of transcriptional bursts observed in vivo[32], suggesting stochastic condensate dynamics as a candidate for mechanisms that underlie transcriptional bursting.

DNA–protein condensation not only provides mechanisms to facilitate enhancer–promoter contacts, but could also play a more general role in DNA compaction such as in mitotic chromatid compaction[33], and the formation of chromatin compartments[11,12]. Our work highlights the role of capillary forces in the interaction between condensates and surfaces. Such forces may also be relevant beyond chromatin in other biological contexts, including membranes and the cytoskeleton. Owing to the tension-dependent nature of transcription-factor-mediated DNA–protein condensation, our work suggests that mechanics and capillary forces from liquid-like condensates could play a key and, as yet, underappreciated role in the spatial organization of the genome and its transcriptional initiation. It is appealing to imagine that transcriptional outputs not only respond to concentrations of transcription factors in the nucleus, but also to mechanical cues from chromatin.

References:

1. De Gennes, P.-G., Brochard-Wyart, F. & Quere, D. *Capillarity and Wetting Phenomena*. Springer, (2013).
2. He, D. L. & Bush, J. W. M. The hydrodynamics of water-walking arthropods. *J. Fluid Mech.* **644,** 5–33 (2010).
3. Elettro, H., Neukirch, S., Vollrath, F. & Antkowiak, A. In-drop capillary spooling of spider capture thread inspires hybrid fibers with mixed solid–liquid mechanical properties. *Proc. Natl. Acad. Sci. USA* **113,** 6143–6147 (2016).
4. Vollrath, F. & Edmonds, D. T. Modulation of the mechanical properties of spider silk. *Nature* **340,** 305–307 (1989).
5. Mastrangeli, M. *et al.* Self-assembly from milli- to nanoscales: methods and applications. *J. Micromech. Microeng.* **19,** 083001–38 (2009).
6. Hyman, A. A., Weber, C. A. & Jülicher, F. Liquid-Liquid Phase Separation in Biology. *Annu. Rev. Cell Dev. Biol.* **30,** 39–58 (2014).
7. Sabari, B. R. *et al.* Coactivator condensation at super-enhancers links phase separation and gene control. *Science* **361,** eaar3958–13 (2018).
8. Hnisz, D., Shrinivas, K., Young, R. A., Chakraborty, A. K. & Sharp, P. A. A Phase Separation Model for Transcriptional Control. *Cell* **169,** 13–23 (2017).
9. Chong, S. *et al.* Imaging dynamic and selective low-complexity domain interactions that control gene transcription. *Science* **361,** eaar2555–11 (2018).
10. Cho, W.-K. *et al.* Mediator and RNA polymerase II clusters associate in transcription-dependent condensates. *Science* **361,** 412–415 (2018).
11. Strom, A. R. *et al.* Phase separation drives heterochromatin domain formation. *Nature* **547,** 241–245 (2017).
12. Larson, A. G. *et al.* Liquid droplet formation by HP1α suggests a role for phase separation in heterochromatin. *Nature* **547,** 236–240 (2017).
13. Brangwynne, C. P., Mitchison, T. J. & Hyman, A. A. Active liquid-like behavior of nucleoli determines their size and shape in Xenopus laevis oocytes. *Proc. Natl. Acad. Sci. USA* **108,** 4334–4339 (2011).
14. Feric, M. *et al.* Coexisting Liquid Phases Underlie Nucleolar Subcompartments. *Cell* **165,** 1686–1697 (2016).
15. Kilic, S. *et al.* Phase separation of 53BP1 determines liquid-like behavior of DNArepair compartments. *EMBO J.* **38,** 7–17 (2019).
16. Patel, A. *et al.* A Liquid-to-Solid Phase Transition of the ALS Protein FUS Accelerated by Disease Mutation. *Cell* **162,** 1066–1077 (2015).
17. Zaret, K. S. & Mango, S. E. Pioneer transcription factors, chromatin dynamics, and cell fate control. *Curr. Opin. Genet. Dev.* **37,** 76–81 (2016).
18. Kaestner, K. H. The FoxA factors in organogenesis and differentiation. *Curr. Opin. Genet. Dev.* **20,** 527–532 (2010).
19. Bustamante, C., Marko, J. F., Siggia, E. D. & Smith, S. Entropic elasticity of lambda-phage DNA. *Science* **265,** 1599–1600 (1994).
20. Sekiya, T., Muthurajan, U. M., Luger, K., Tulin, A. V. & Zaret, K. S. Nucleosome-binding affinity as a primary determinant of the nuclear

mobility of the pioneer transcription factor FoxA. *Genes Dev.* **23,** 804–809
(2009).
21. Iwafuchi, M. *et al.* Gene network transitions in embryos depend upon
interactions between a pioneer transcription factor and core histones. *Nat.*
Genet. **52,** 418–427 (2020).
22. Zhou, H.-X., Nguemaha, V., Mazarakos, K. & Qin, S. Why Do Disordered and
Structured Proteins Behave Differently in Phase Separation? *Trends*
Biochem. Sci. **43,** 499–516 (2018).
23. Jawerth, L. M. *et al.* Salt-Dependent Rheology and Surface Tension of
Protein Condensates Using Optical Traps. *Phys. Rev. Lett.* **121,** 258101
(2018).
24. Brangwynne, C. P. *et al.* Germline P granules are liquid droplets that
localize by controlled dissolution/condensation. *Science* **324,** 1729–1732
(2009).
25. Botao, X. *et* al. Histone H1 compacts DNA under force and during
chromatin assembly. **23,** 4864–4871 (2012).
26. Golfier, S., Quail, T., Kimura, H. & Brugués, J. Cohesin and condensin
extrude DNA loops in a cell-cycle dependent manner. *Elife* **9,** (2020).
27. Ganji, M. *et al.* Real-time imaging of DNA loop extrusion by condensin.
Science **360,** 102–105 (2018).
28. Davidson, I. F. *et al.* DNA loop extrusion by human cohesin. *Science* **366,**
1338–1345 (2019).
29. Kim, Y., Shi, Z., Zhang, H., Finkelstein, I. J. & Yu, H. Human cohesin compacts
DNA by loop extrusion. *Science* **366,** 1345–1349 (2019).
30. Banigan, E. J. & Mirny, L. A. Loop extrusion: theory meets single-molecule
experiments. *Curr. Opin. Cell Biol.* **64,** 124–138 (2020).
31. Cramer, P. Organization and regulation of gene transcription. *Nature* **573**,
45–54 (2019).
32. Fukaya, T., Lim, B. & Levine, M. Enhancer Control of Transcriptional
Bursting. *Cell* **166,** 358–368 (2016).
33. Yoshimura, S. H. & Hirano, T. HEAT repeats – versatile arrays of
amphiphilic helices working in crowded environments? *J. Cell. Sci.*
jcs.185710–8 (2016).
344

Acknowledgements: We thank Pavel Tomancak, Anthony Hyman, Stephan Grill,

Iain Patten, Martin Loose, David Oriola, Benjamin Dalton, Patrick McCall,

Vasanthnarayan Murugesan, and Claudia Meyer for helpful feedback and

stimulating discussions. We would like to thank Aliona Bogdanova for

discussions and help with cloning as well as both the Protein Expression and

Purification Facility and the Light Microscopy Facility at the Max Planck Institute

of Molecular Cell Biology and Genetics. We would like to acknowledge and thank

Nadine Vastenhouw for discussions and the construct containing the Tata-box-

binding protein template, and Christoph Zechner for help with statistical

analyses. Lastly, we would like to thank Kenneth Zaret for sending constructs,

information on mutant FoxA1 proteins, and advice on FoxA1 protein purification.

This work was supported by an EMBO long-term fellowship (ALTF-1456-2015)

and a Volkswagen "Life" grant number 94820.

Author contributions: T.Q. and J.B. conceived the project. T.Q. and S.G.

performed imaging experiments. T.Q. purified proteins, made constructs, and

performed data analysis. T.Q., J.B., and F.J. performed theoretical calculations.

M.E. made the TBP construct and purified the protein. K.I. made the initial FoxA1

construct and provided key biochemical support. J.B. and F.J. supervised the

work. T.Q., J.B., and F.J. wrote the manuscript and all authors contributed ideas

and reviewed the manuscript.

371 **Table of contents:**

**Fig. 1: FoxA1 forms DNA-FoxA1 condensates in a tension-dependent
manner.** (A) Schematic of single λ-phage DNA molecule assay. (B) Structure of
FoxA1, consisting of a structured DNA-binding domain flanked by mostly
disordered N and C termini. The DNA-binding domain has a sequence-specific
binding region (blue) and two non-sequence-specific binding regions (green). (C)
The extent of FoxA1-mediated DNA condensation depends on the end-to-end
distance of the strand. Representative time-averaged projections of FoxA1 and
DNA. Note that the total amount of DNA is the same in each example. Scale bar=2
μm. (D) Schematic displaying three main quantities used to characterize DNA-
FoxA1 condensation: the end-to-end distance L; Cross-correlation of DNA and
FoxA1 intensities; and DNA envelope width, a measure of transverse DNA
fluctuations. (E) Cross-correlation of FoxA1 and DNA signals shows that FoxA1
condenses DNA below a critical end-to-end distance. The gray dots represent
individual strands, n=107. The data is binned every 2-μm (black, mean ± SD). (F)

DNA envelope width measurements reveal that FoxA1-DNA condensation
buffers DNA tension (blue and black dots correspond to control and DNA+FoxA1
conditions, n=46 and n=50 respectively). The data is binned every 2-μm (mean $\pm$
SD). (G) Representative images of FoxA1 zipping two independent DNA strands
over time. Scale bar=1 μm.

460 **Figure 2**

461
462
Fig. 2: Mutant analysis reveals that the C terminus of FoxA1 drives DNA
condensation. Representative images of FoxA1 mutants and corresponding DNA
envelope width measurements. The data is binned every 2-µm and the mean ±
SD are shown in black for each mutant and in blue for control (n=46). Scale
bar=2 μm. (A) Sequence-specific DNA binding mutant NH-FoxA1 condenses DNA
(n=30). (B) Non-sequence-specific DNA-binding mutant RR-FoxA1 condenses
DNA (n=28). (C) N-terminal truncation of FoxA1 ΔN-FoxA1 condenses DNA
(n=15). (D) C-terminal truncation of FoxA1 ΔC-FoxA1 inhibits DNA condensation
(n=34).

Figure 3

Fig. 3: Thermodynamic description of a liquid phase condensing onto a semi-flexible polymer explains FoxA1-mediated DNA condensation. (A) Schematic representing DNA-FoxA1 condensation (orange). DNA can be in a condensed state (black) or a non-condensed state (green). DNA condensation depends on the condensate surface tension (γ), condensation free energy per volume (μ), and DNA packing efficiency (α). (B) Free energy profiles as a function of condensed DNA (L_d) for different L reveal a first-order phase transition for DNA–protein condensation (orange and blue correspond to favorable and unfavorable condensation, respectively). (C) Boltzmann distributions corresponding to the free energy profiles in (B). (D) Condensate volume linearly increases with L_d. The orange curve represents a linear fit to individual strands (n=53). For (D), (E), and (F), individual strands are represented as gray dots and binned mean±SEM is in black. (E) Amount of condensed DNA as a function of L

(n=63 strands) reveals sharp transition. Orange curve represents optimal theoretical fit. The gray dashed-line corresponds to the limit of maximum condensation where L_d is equal to the contour length of DNA (16.5 μm) minus L. (F) Capillary forces that DNA-protein condensates exert on non-condensed DNA are buffered (n=63 strands). Orange curve is the theoretical prediction. The gray dashed line represents the force when L_d=0. (G) Probability to nucleate a DNA-FoxA1 condensate (P_{cond}) reveals a sharp transition at a critical end-to-end distance. P_{cond} is computed from binned local correlation data (n=181 condensates). The end-to-end distance error bars are the SEM, and the P_{cond} error bars are the 95% confidence intervals from a Beta distribution.

Figure 4

D Capillary forces as chromatin remodellers

Fig. 4: Universality of protein-mediated DNA-protein condensation.
Representative images of sequence-specific DNA-binding mutant NH-FoxA1 (A),
Tata-box-binding protein (B), and Somatic linker histone H1 (C), with
corresponding P_{cond} data below. DNA is stained with 10 nM Sytox Green for NH-
FoxA1 and 25 nM Sytox Orange for both Tata-box-binding protein and Somatic
linker histone H1. P_{cond} is computed from local correlation data with n=174
condensates for NH-FoxA1 (A), n=154 condensates for Tata-box-binding protein
(B), and n=61 for H1 (C). Scale bar=2 μm. We found that NH-FoxA1 condensed
DNA less efficiently than WT-FoxA1, TBP could not condense DNA for any end-
to-end distance, and H1 condensed DNA for all measured end-to-end distances.
(D) Biomolecular condensates generate capillary forces that could serve to
recruit transcriptional regulators, and potentially remodel chromatin at
physiologically relevant force scales in order to properly regulate transcription.

Methods

<u>Cloning and protein purification</u>

FoxA1 was first cloned into a middle-entry Destination vector with a fluorescent

tag on the C terminus and a glycine-rich flexible linker region between FoxA1

and mCherry. A LR reaction was performed with pDEST17, a destination vector

with a N-terminal His$_6$ tag using Gateway Cloning to introduce FoxA1-mCherry

into this bacterial expression vector: pDEST17-FoxA1-mCherry. This vector was

transformed into T7 express bacterial cells (enhanced BL21 derivative, NEB

C2566I), and grown to OD~0.4-0.8, whereupon we added 1 mM IPTG and

expressed His$_6$-FoxA1-mCherry for 3-4 hours at 37 °C. We snap-froze the pellets

in liquid nitrogen and stored them at -80 °C. The pellets were redissolved in

binding buffer (1xBB) that contained 20 mM Tris-HCl (pH=7.9), 500 mM NaCl, 20

mM Imidazole, 1 mM MgCl$_2$, supplemented with a single cOmplete Protease

Inhibitor Cocktail tablet and Benzonase. The redissolved pellets were lysed using

a pressure-based homogenizer, and clarified via centrifugation. The supernatant

was discarded and the pellets were resuspended in 1xBB + 6 M Urea. The lysate

was spun again, and the supernatant was poured over an IMAC column, washed

with 10 column volumes (5 ml per column volume) of 1xBB + 6 M Urea, and then

eluted with 1xBB+6 M Urea+250 mM Imidazole. Protein-rich fractions were

detected using a Bradford assay, pooled, and then dialyzed overnight into

storage buffer (1xSB), 20 mM HEPES (pH=6.5), 100 mM KCl, 1 mM MgCl$_2$, 3 mM

DTT, and 5 M Urea. Next, the proteins were dialyzed in 1xSB+4 M Urea for two

hours, then 1xSB+3 M Urea for two hours, and finally 1xSB+2 M Urea for two

hours. The protein fraction was spun-concentrated to 4-5 mg/ml (~50 μM), and

then snap-frozen in liquid nitrogen and stored at -80 °C. Dialyzing the protein

from 2 M urea to lower amounts of urea led to protein precipitation. NH-FoxA1-

mCherry and RR-FoxA1-mCherry were obtained by modifying the pDEST17-

FoxA1-mCherry vector using the Q5 Site-Directed Mutagenesis Kit (NEB), and

purified using the same protocol given above. Specifically, for NH-FoxA1, we

mutated residue number 216 (where residue number 1 is start codon of the

open-reading frame of FoxA1) from an asparagine to an alanine, and residue

number 222 from a histidine to an alanine[20]. For RR-FoxA1, we changed residue

numbers 262 and 265 from arginines to alanines[20]. For the FoxA1 N-terminal

truncation, we first digested pDEST17-FoxA1-mCherry with NdeI and BamHI—

these digestion sites are unique and located right before the N-terminal His$_6$ tag

and directly following the C-terminal mCherry. Then we amplified the target

using a forward primer that contained the NdeI cut site, a His$_6$ tag, and the start

of the DNA-binding domain (residue number 144) and a reverse primer that had

the BamHI cut site and the end of the mCherry sequence. For the C-terminal

truncation, we again digested pDEST17-FoxA1-mCherry with NdeI and BamHI,

and amplified our target using two sets of primers: first we amplified the

purification tag, the N terminus, and the DNA-binding domain using a reverse

primer that targeted the end of the DNA-binding domain and had a unique SpeI

cut site. Second, we simply amplified the mCherry fluorescent tag where the

forward primer contained a SpeI cut site. For both constructs, we then cut the

template, performed a ligation, transformed into bacteria, and selected for clones

using ampicillin. We used somatic linker histone purified from calf thymus,

labeled with Alexa Fluor 488 dye, kept at 2 mg/ml (~70 μM) in 1xPBS, stored

snap-frozen at -80 °C (H-13188, ThermoFisher). To purify Tata-box binding

protein, we cloned a template of a zebrafish Tata-box-binding protein (zTBP)—a

gift from Nadine Vastenhouw—into a bacterial expression vector where a His$_6$

tag, a Maltose-binding protein purification tag, and an eGFP fluorescent tag were

on the N terminus (His$_6$-MBP-eGFP-zTBP). This vector was transformed into T7

express bacterial cells (enhanced BL21 derivative, NEB C2566I) and grown to

OD~0.4-0.8, whereupon we added 0.2 mM IPTG and expressed His$_6$-MBP-eGFP-

zTBP overnight at 18 °C. We lysed the cells into lysis buffer A containing 50 mM

Tris-HCl (pH=8.0), 1 M NaCl, 10% glycerol, 1 mM DTT, 1 mM MgCl$_2$

supplemented with a single cOmplete Protease Inhibitor Cocktail tablet. After

lysis, we added NP40 to 0.1% and clarified via centrifugation. We performed a

polyethylenimine precipitation to precipitate the remaining DNA in the soluble

fraction and then an ammonium sulfate precipitation to recover the protein

fraction, resuspending the precipitated proteins in lysis buffer B containing 50

mM Tris-HCl (pH=8.0), 1 M NaCl, 10% glycerol, 1 mM DTT, 0.1% NP40, and 20

mM imidazole and clarified the soluble fraction via centrifugation. We poured the

lysate over an IMAC column and eluted using 2xPBS, 250 mM imidazole, 10%

glycerol, and 1 mM DTT. We performed a Bradford assay to detect protein-rich

fractions, pooled these fractions, and then dialyzed the proteins overnight into

20 mM HEPES pH=7.7, 150 mM KCl, 10% glycerol, and 1 mM DTT. We spun-

concentrated the proteins to 10 mg/ml ($\sim$100 μM), snap-froze using liquid

nitrogen, and stored at -80 °C.

<u>DNA functionalization</u>

To biotinylate DNA purified from λ-phage (λ-phage DNA), we combined 10 μg of

λ-phage DNA (NEB, N3011S) and 5 μl of a 10X polymerase buffer (50 mM Tris-

HCl, pH=7.2, 10 mM $MgSO_4$, 100 μM DTT) to a total reaction volume of 50 μl. We

then heated the mixture up to 65 °C for 7 minutes to break apart the λ-phage

DNA's sticky ends. After heat treatment, we added 100x molar excess of

biotinylated dATP, biotinylated dUTP, and dGTP, and dCTP. We then added 1 unit

($\sim$1 μl) of Klenow enzyme, mixed well, and incubated overnight at room

temperature. We purified the biotinylated λ-phage DNA using ethanol

precipitation and stored at -20 °C.

<u>PEGylation of cover slips and DNA micro-channel preparation</u>

We functionalized glass cover slips with mPEG and PEG-Biotin. Briefly, we

sonicated coverslips first in acetone for 15 minutes followed by 5 rinses with

MilliQ water, and then another sonication step in 5 M KOH for 40 minutes. After

rinsing the coverslips 3 times with water and then 3 times with methanol, we

dried the coverslips with N_2. We silanized the coverslips combining 250 ml

methanol, 12.5 ml acetic acid, and 2.5 ml (3-aminopropyl)-trimethoxysilane,

incubating the coverslips in this mixture for 10 minutes at room temperature,

sonicating for 1 minute, and then incubating the coverslips for an additional 10

minutes. Next we rinsed the coverslips once with methanol, once with water, and

once again methanol, and dried with N_2. Then we mixed 100 mg mPEG and $\sim$1.5

mg Biotin-PEG with 450 μl PEGylation buffer (0.1M Sodium Bicarbonate,

pH=8.5), and spun the reaction at 10000 RPM for 1 minute. We pipetted 25 μl of

the PEG mixture onto a dried, silanized coverslip and put another coverslip on

top, generating a coverslip sandwich. We incubated these sandwiches over night

in distilled water-filled pipette tip-boxes in the dark. After incubation, we

carefully disassembled the coverslips, rinsed with MilliQ water, and dried with

N$_2$. To generate a channel for imaging, we first drilled holes through a cleaned

cover slide—these holes acted as channel inlets and outlets. We placed custom-

designed, laser-cut double-sided tape onto the coverslip, defining the channel

geometry. We then placed a functionalized PEG-biotinylated coverslip on top of

the double-sided tape, sealing the channel on either end with Valap. We filled the

channel with ~10-15 μl of 0.1 mg/ml free streptavidin, incubating the channel

with streptavidin for 1 minute. To remove the free, unbound streptavidin, we

flushed ~100 μl channel washing buffer (40 mM Tris-HCl, pH=8.0, 20 mM NaCl,

0.4 mM EDTA) through the channel, using the drilled holes as channel inlets and

outlets. We added 20 μl of 1:1000 biotinylated λ-phage DNA (~5 pM), incubating

it for ~1 min and then washed the channel with 3x100 μl of channel washing

buffer.

<u>DNA imaging</u>

For live imaging, we fluorescently stained immobilized DNA strands with 10-20

nM Sytox Green in imaging buffer (20 mM HEPES, pH=7.8, 50 mM KCl, 2 mM

DTT, 5% glycerol, 100 μg/ml BSA). For experiments with H1-Alexa-488 and

MBP-TBP-eGFP, we imaged DNA with 25 nM Sytox Orange. All experiments were

performed with this buffer and the concentration we used for all proteins was 10

nM. In general, protein concentrations above 10 nM led to an increase in the

number of strands that stuck to the coverslip. Before performing an experiment,

we pre-diluted our stock concentration 1:1000 and then performed a hard-spin

at 13K at 4°C for 1 minute. We then further diluted the proteins to 10 nM and

performed another hard-spin at 13K at 4°C for 1 minute. We excited Sytox

Green- or Sytox Orange-labelled DNA using a 488 nm and 561 nm laser

respectively, imaging the strands using a Nikon Eclipse microscope stand with a

Nikon 100x/NA 1.49 oil SR Apo TIRF and an Andor iXon3 EMCCD camera using a

frame-rate of 100 – 300ms. A highly inclined and laminated optical sheet (HILO)

microscopy mode was established using a Nikon Ti-TIRF-E unit mounted onto

the microscope stand to improve signal-to-noise ratio by cutting off background

fluorescence signal from unbound DNA dye in the buffer.

<u>Strand length calculation</u>

To calculate the end-to-end distance, we generated time-averaged projections of
FoxA1 and DNA and integrated these projections along the orthogonal axis to the
strand. To find the profiles' left edge, we computed the gradient of the DNA
signal's one-dimensional integrated intensity and computed the position when
the gradient signal went through a threshold (defined as 0.2). We then took all
the points from the start of the signal to this position, background-subtracted
this cropped signal, and fit an exponential to these points. To ensure that we
were including all of the DNA signal, we defined the threshold for both the left
and the right edges as three-quarters of the value of the fitted exponential value
at the point when the gradient had gone through the threshold. This threshold
value gave rise to end-to-end distances consistent from what we could observe
from the fluorescent images and from qualitative measurements that we
performed in FIJI. Using this threshold, we background-subtracted our signal and
simply computed the position values at which the signal went through the
threshold on the left and the right sides, and computed the end-to-end distance
as the difference between these two positions.

Global cross-correlation analysis
We generated time-averaged projections of both the FoxA1 and DNA signal from
movies roughly 30-60 sec in duration, and then summed the intensities in the
orthogonal axis to the strand, generating line profiles for both the FoxA1 and
DNA signals. We then normalized the amplitudes of the signals by their
Euclidean distances, and computed the zero-lag cross-correlation coefficient of
the normalized signals, which we defined as "Correlation". In general, Correlation
values range from -1 to 1, but, in our experimental data, the values range from
roughly 0 to 1, where 1 represents the formation of DNA-FoxA1 condensates and
0 represents the formation of only FoxA1 condensates (no DNA condensation).

DNA envelope width calculation
To compute the DNA envelope width, we first generated time-averaged
projections from movies of FoxA1 and DNA. We then selected segments of the
strand that did not contain FoxA1—regions of non-condensed DNA. Using these
segments, we then extracted a line profile of the DNA signal orthogonal to the

strand that gave the maximum width. We then subtracted off the background of

the DNA profile, scaled the signal's amplitude using the Euclidean distance, and

fit a Gaussian. We defined the DNA envelope width as 2σ, which represents two

times the standard deviation of the fitted Gaussian.

<u>Condensate volume analysis</u>

To calculate the volume, we first generated time-averaged DNA-FoxA1

projections, and then localized the peaks of the DNA condensates. Using the

location of the peaks, we extracted background-subtracted one-dimensional

profiles of the DNA condensates in the orthogonal axis to the strand—these

profiles went through the peak location. We then fit Gaussians to these profiles

without normalizing the amplitude. To define the radii of the condensates, we

computed the gradient of the fitted Gaussians and defined the condensate

"edges" as when the absolute value of the gradient of the Gaussian function

gradient went through a threshold value (defined as one, and determined by

comparing with fluorescence). Assuming condensates are spherical, we

computed the condensate volume as $V = \frac{4}{3}\pi R^3$, where R is the condensate's

radius. To compute a condensate volume for strands with multiple condensates,

we simply added up the volumes for each condensate.

<u>Condensed DNA amount analysis</u>

To compute the contour length of DNA contained within a DNA-protein

condensate—defined hereafter as condensed DNA amount (L_d)—we used a

fluorescence-based approach. We first generated time-averaged projections of

DNA and FoxA1 signals, and integrated the DNA signal in the orthogonal

direction to the DNA strand. We then defined condensed vs non-condensed DNA

with $Threshold_{drop}$: the median value of the profile plus a tolerance. Intensity

values below $Threshold_{drop}$ were defined as pixels associated with non-

condensed DNA, and intensity values above $Threshold_{drop}$ were defined as pixels

associated with condensed DNA. This assumption was also consistent with the

measured FoxA1 signal, where FoxA1 signals clearly localized to regions of

condensed DNA, as defined by the $Threshold_{drop}$. The tolerance value was used to

suppress artefactual fluctuations of the non-condensed DNA signal in the

neighborhood of the median. To optimize the tolerance value, we assume that L_d

as a function of L is linear for lower values of L (<5 μm) with a y-intercept equal

to the contour length of the DNA molecule (16.5 μm), as this is consistent with

our theoretical description. We plotted the y-intercepts of the linear fits as a

function of tolerance and found that tolerance=500 gives a y-intercept equal to

16.5 and generates DNA-FoxA1 condensates up to 10 μm consistent with our

data and analysis (Fig. S3). In order to calculate the DNA length contained within

the droplet, we integrated the intensities from pixels above Threshold$_{drop}$,

divided this value by the sum of the total intensity of the profile, and then

multiplied this ratio by the contour length of λ-phage DNA, 16.5 μm. The non-

condensed DNA length was calculated as simply the contour length minus L_d.

<u>Force analysis</u>

In order to calculate the force that the condensate exerts against the non-

condensed DNA strand, we used the worm-like chain model. The worm-like

chain model gives a relationship between the extension of λ-phage DNA —$E = \frac{L}{L_t}$,

where L is the end-to-end distance and L_t is the total contour length of the

molecule (16.5 μm, for λ-phage DNA)—and the force. Quantitatively, upon

addition of FoxA1, the amount of non-condensed DNA reduces, and the extension

changes as follows, $E = \frac{L}{L_t - L_d}$, where L_d is the amount of condensed DNA. With

this expression, we directly compute the force using the worm-like -chain model,

$F = \kappa \left(\frac{1}{4}(1 - E)^{-2} - \frac{1}{4} + E \right)$

<u>Condensate nucleation probability analysis</u>

To calculate the probability that a FoxA1 condensate would generate a DNA

condensate as a function of end-to-end distance, we first localized the peaks of

the FoxA1 condensates from time-averaged projections taken from movies of

FoxA1 and DNA. We then extracted 0.9 μm x 0.5 μm windows centered around

the localized FoxA1 peaks of both the FoxA1 and DNA signals—with the

window's long axis going with the strand and the short axis as orthogonal to the

strand. This cut the image up into smaller boxes. We then computed the zero-lag

normalized cross-correlation coefficient as follows:

809
$$C_{loc} = \frac{\sum\sum(f(x,y)-\mu_f)(g(x,y)-\mu_g)}{\sqrt{\sum f(x,y)^2 - \mu_f{}^2}\sqrt{\sum g(x,y)^2 - \mu_g{}^2}}$$

where $f(x,y)$ is the DNA, $g(x,y)$ is FoxA1, μ_f is the mean of the DNA image, and

μ_g is the mean of the FoxA1 image. This generates values from -1 to 1. For FoxA1-

mediated DNA condensation, the values for particular condensates are clearly

close to 1. When FoxA1 fails to mediate DNA condensation, owing to the

morphology of the underlying DNA strand and the small number of pixels, we

obtain values that range from -1 to roughly 0.5. To obtain a value for P_{nuc} as a

function of end-to-end distance, we selected a threshold of 0.75—C_{loc} values

above the threshold are considered as "condensed" and values below would be

considered "non-condensed". Hence, we binned the C_{loc} data in 2-μm increments

as a function of end-to-end distance, and calculated P_{cond} by simply taking the

number of "condensed" condensates and dividing it by the total number of

condensates within the bin. The confidence intervals for P_{cond} in each respective

bin are computed by computing the 95% confidence interval of a beta-

distribution, which represents the probability distribution for a Bernoulli

process that takes into account the total number of successes with respect to the

total number of attempts.

<u>Parameter fitting and confidence intervals</u>

Our thermodynamic description contains three parameters: α, the DNA packing

efficiency given as DNA strand length as a function of cross-sectional area; γ, the

surface tension of the condensate; and μ, the condensation free energy per

volume. To fit α, we computed condensate volumes for individual strands and

plotted these volumes as a function of condensed DNA (L_d) and fit a linear

function to the individual points, giving 0.04 μm^2 for WT-FoxA1 and 0.09 μm^2 for

NH-FoxA1. The confidence intervals presented in the main text represent the 95

per cent confidence intervals that were generated from fitting the individual

points directly. To fit γ and μ, we generated theoretical expressions for both L_d

and P_{cond} as a function of L, and minimized the error of these theoretical

expressions with respect to the experimental data to optimize the parameter

values. First, to compute a theoretical expression of $L_d(L)$, we generated free

energy profiles as a function of condensed DNA (L_d) for values of L from 0 to L_t,

where L_t is λ-phage DNA's contour length, $16.5\ \mu m$. For each individual free

energy profile, we computed the normalized Boltzmann distribution $P(L_d) =$

$\frac{e^{-\beta F(L,L_d)}}{\int_0^{L_t-L} e^{-\beta F(l)} dl}$ and then calculated the mean of the Boltzmann distribution $\overline{L_d} =$

$\int_0^{L_t-L} l P(l) dl$, giving rise to $\overline{L_d}(L)$. To compute theoretical expressions of

$P_{cond}(L)$, we again generated free energy profiles for a range of L from $[0, L_t]$

plotted as a function of L_d. We then localized the position of the local maximum

in the free energy, L_d^{max}, and then computed the probability to "not" nucleate a

droplet from the Boltzmann distribution $\int_0^{L_d^{max}} P(l) dl$. Following from this, we

then computed $P_{cond} = 1 - \int_0^{L_d^{max}} P(l) dl$, giving rise to $P_{cond}(L)$. To optimally fit

γ and μ, we computed weighted errors from the theoretical expressions as

compared to the experimental data. For the following error minimization

analysis, we compared the theoretical expressions against the mean of data that

was binned in 2-μm-width bins. Briefly, for a given set of (γ, μ), we computed

theoretical expressions for both $\overline{L_d}(L)$ and $P_{cond}(L)$. Then, for each "binned"

mean for both condensed DNA and condensation probability, we computed the

squared residual of the mean value with the theoretical expression. For residuals

calculated from $\overline{L_d}(L)$, we normalized each residual by the squared standard

error of the mean, and then summed the normalized residuals to obtain the

error. For residuals calculated from $P_{nuc}(L)$, we normalized each residual by the

variance of the beta distribution,

$P_{nuc\,cond}\,\sigma^2 = \frac{(1+k)(1-k+N)}{(2N^2(3+N))}$ and then summed the normalized residuals to obtain the

error. For the global error, we simply added the error from both deviations in

$\overline{L_d}(L)$ and $P_{cond}(L)$. We then iterated through a range of values for (γ, μ) and

computed the total error associated with each set of parameter values,

exponentiated the negative values of the total error matrix, and computed the

largest combined value to select the parameter values. To calculate confidence

intervals for the fitted parameters from the error minimization, we first obtained

one-dimensional profiles of the integrated exponentiated total error for μ as a

function of γ and γ as a function of μ. The peaks of these profiles represented the

values that we selected for our best-fit parameters. We assumed that these

profiles represented probability distributions for parameter selection, and then

calculated the left and right bounds where the area under the curve between these bounds represented 95 per cent of the area. These left and right bounds represent the lower and upper values of our confidence intervals. To compute the 95 per cent confidence intervals for the force for each respective end-to-end distance value, we scanned through (γ, μ) parameter space and computed the value of L_d for each set of parameters. We then plotted these values against the probability that these parameter values were the "true" values—simply the probability from the exponentiated error matrix. Integrating the points under the Probability vs. L_d curve and dividing this by the total area under this curve, we generated a probability distribution function from which we could compute the 95% confidence intervals for L_d. Because the force was constant, to compute the confidence intervals for the force, we calculated the force using the worm-like chain model using corresponding L_d values for an end-to-end distance that retained FoxA1-mediated DNA condensation. To compute the confidence intervals for L_{crit}, we scanned through (γ, μ) parameter space and computed L_{crit} for each set of parameters. We then plotted L_{crit} values with the corresponding values from the probability that these parameter values were true (again, the exponentiated error matrix). Integrating the points under the Probability vs L_{crit} curve and dividing this by the total area under this curve, we generated a probability distribution function from which we could compute the 95% confidence intervals for L_{crit}.

Extended Data Figure 1:

Extended Data Fig. 1: Quantification of FoxA1-mediated DNA condensation.
(A) Representative fluorescent images of FoxA1-mCherry in buffer (20 mM HEPES, pH=7.8, 50 mM KCl, 2 mM DTT, 5% glycerol, 100 μg/ml BSA) at different concentrations, 10-500 nM, in the absence of DNA reveals that FoxA1 does not form condensates in bulk. Using spinning disk microscopy and a 60x objective, we acquired images 70 μm x 70 μm in size with an exposure time of 250 msec and a time stamp of 500 msec to generate movies 30 seconds in duration. For all measured concentrations we generated n=3 movies and did not observe any FoxA1 condensation. (B) Global cross-correlation between FoxA1 and DNA reveals FoxA1-mediatd DNA condensation. Left, representative fluorescent time-averaged projections of DNA and FoxA1 at two different end-to-end distances.

Integrating both the DNA and FoxA1 signals along the axis orthogonal to the long axis of the strand gave rise to line profiles, which we normalized, and then plotted as a function of distance (DNA in black and FoxA1 in orange). We then computed the zero-lag cross-correlation coefficient (C_{global}). (C) DNA envelope width measure measures the transverse fluctuation of non-condensed DNA. Top box: DNA alone condition. Bottom box: DNA+FoxA1 condition. For both conditions, we display representative fluorescent images of single frames and time-averaged projections of the DNA and FoxA1 signals. The white dashed line represents the maximum width of the DNA signal along the orthogonal axis of the non-condensed DNA. The black dots in the profile represent the background-subtracted points from the white dashed line, and the black line represents a Gaussian fit. The DNA envelope width was defined as 2 times the standard deviation of the Gaussian fit. (D) Three representative examples of FoxA1-mediated zipping. These images are time-averaged projections of both FoxA1 and DNA.

Extended Data Figure 2:

Extended Data Fig. 2: Quantification of condensate volume, force and condensation probability. (A) Condensate volume calculation. Top, a representative time-averaged projection of a single DNA condensate. The black dashed cross represents the peak location. The white dashed line is the line profile position that we extract from the image governed by the peak location. Below, the black dots are the background-subtracted values of the line profile and the solid black line is a fit to a Gaussian function. The gray line represents the threshold value computed from the gradient of the Gaussian function that defines the edges of the condensate (see Methods). These positions define a condensate diameter that, assuming the condensates are spherical, allows to compute the volume of the condensate. (B) Condensed DNA calculation. Top, representative time-averaged projections of DNA and FoxA1. Below, a line profile (black) was computed from integrating the DNA signal along the orthogonal axis to the strand. To define condensed versus non-condensed regions of DNA, we defined a threshold given by the median of the signal (gray) plus a tolerance (black dashed). Above this threshold, pixels count towards condensed DNA, and, below this threshold to non-condensed DNA. (C) Local correlation calculation. In order to compute P_{cond} as a function of end-to-end distance, we computed the correlations between individual condensates. We first localized the peaks of FoxA1 condensates and then cropped 1 μm x 0.5 μm boxes, centered around these peaks, of both the FoxA1 and corresponding DNA signals. The black dashed crosses give the location of the peaks and the images below are the boxes for both the FoxA1 and the DNA signal (FoxA1 to the left, DNA to the right). We then computed the normalized correlation of these images.

1006   **Extended Data Figure 3:**

1007
1008
1009
Extended Data Fig. 3: Tolerance value calculation. To calculate the amount of
condensed DNA (L_d), we define a threshold for integrated DNA signal intensities
that controls how we differentiate between condensed versus non-condensed
DNA. The threshold is the median value of the signal plus a tolerance—a
tolerance is required to suppress artefactual fluctuations of the signal close to
the median value. We plotted L_d as a function of end-to-end distance L for
different tolerance values. The black dots represent the amount of condensed
DNA for individual strands and the orange curve represent linear fits to these
points for end-to-end distance below 5 μm. (B) To optimize the tolerance value,
we assume that L_d as a function of L is linear for lower values of L (< 5 μm) with a
y-intercept equal to the contour length of the DNA molecule (16.5 μm). We
plotted the y-intercepts of the linear as a function of the tolerance and found two
values with y-intercepts equal to 16.5. We selected a tolerance=500, as this was
consistent with FoxA1-mediated DNA-FoxA1 condensate formation up to
approximately 10 μm, which we experimentally observed, as opposed to a
tolerance~2000, that gave DNA-FoxA1 condensate formation up to only 5 μm,
inconsistent with our measurements and analysis (see Methods).
1027
1028
1029

1030   **Extended Data Figure 4:**

1031
1032
1033
Extended Data Fig. 4: Bistability of FoxA1-mediated DNA condensation. (A)
Representative time-averaged projections of DNA and FoxA1 signals show that
FoxA1 condenses DNA in an all-or-nothing manner. On the right side of each pair
of images, we localized the FoxA1 condensates and showed whether FoxA1
condenses DNA (filled-in gray circle) or not (open circle). Interestingly, there is a
mixed population, revealing the bistable nature of the condensation process. (B)
Representative images of condensation bistability for the sequence-specific DNA-
binding mutant, NH-FoxA1.

Extended Data Figure 5

Extended Data Fig. 5: Quantification of NH-FoxA1-mediated DNA condensation. (A) Condensate volume as a function of condensed DNA amount (L_d). The black dots represent individual strands (n=43) and the data is binned every 2 μm (mean ± SEM). Fitting the individual points, we obtained a linear curve with slope 0.09 μm^2 given in orange. The green dashed line is a guide to demonstrate the theoretically determined fit for WT-FoxA1 (Fig. 3D) with a slope of 0.04 μm^2. (B) Condensed DNA amount as a function of end-to-end distance (L). The black dots represent individual strands (n=57) and the data is binned every 2 μm (mean ± SEM). The orange curve is the expression computed from the theoretical description parameter values determined through error minimization (see Methods). (C) The force that the condensate exerts on the non-condensed DNA as a function of end-to-end distance. The black dots represent individual strands (n=57) and the data is binned every 2 μm (mean ± SEM). The orange curve is the expression computed from the theoretical expression of L_d versus L from panel B for the force. NH-FoxA1 generates forces at roughly 0.14 pN. The dashed black line represents the force exerted on the non-condensed strand when L_d=0. (D) The probability for FoxA1 to form a DNA-FoxA1 condensate as a function of end-to-end distance. Local correlations of individual FoxA1 condensates with DNA (Fig. S2 C) are first calculated and binned into 2-μm-width bins. P_{cond} is then computed by taking all of the values of the local correlations that are above the threshold value 0.75 in each bin and dividing this number by the total number of condensates in the bin. There are a total number of n=174 condensates used for this analysis. The dashed lines represent the P_{cond} value as computed within the bin with ± SEM for the end-to-end distance of the strand. The interval for P_{cond} are computed by computing the 95% confidence interval of a beta-distribution, which represents the probability distribution for a Bernoulli process that takes into account the total number of successes with respect to the total number of attempts.

1090

1091

Extended Data Fig. 6: Dynamics of DNA-binding proteins. (A) Representative
images of FoxA1 condensates on DNA. The kymograph reveals FoxA1
condensates do not move on DNA. (B) NH-FoxA1 condensates remain stable on
DNA and do not move. (C) TBP condensates exhibit diffusive-like behavior on
DNA. (D) Similar to FoxA1 condensation, H1 condensates do not exhibit
diffusive-like behavior on DNA.

Supplementary Information

1   Thermodynamic description of DNA-protein condensation

We consider the change of free energy associated with nucleating a condensate of radius R that contains DNA and FoxA1. The free energy of this process contains volume and surface contributions as well as the capillary work to pull DNA into the condensate,

$$F(R) = -\mu \frac{4}{3}\pi R^3 + \gamma 4\pi R^2 + W \tag{1}$$

where μ is the condensation free energy per volume, γ is the surface tension of the condensate, and W is the capillary work. We assume that the DNA in the condensate is in a collapsed state, which corresponds to the maximal packing allowed by excluded volume. With this assumption, the droplet volume and the amount of condensed DNA are linearly related, $V = \alpha L_d$, or $R^3 = \frac{3\alpha}{4\pi} L_d$, where α describes the DNA packing efficiency given as DNA strand length per cross-sectional area, and L_d is the amount of condensed DNA. We can then obtain the free energy of nucleating a condensate as a function of L_d,

$$F(L_d) = -\mu \alpha L_d + \gamma 4\pi \left(\frac{3\alpha}{4\pi}\right)^{2/3} L_d^{2/3} + W. \tag{2}$$

The work $W = \int_0^{L_d} f dx$ is the work that must be performed by capillary forces f to pull an

amount L_d into the condensate. To define f, we use a modified version of the phenomenological force-extension curve of the worm-like-chain model for λ-phage DNA that takes into account the amount of available non-condensed DNA (19). To this end, we modified the extension that the non-condensed DNA feels following the nucleation of a FoxA1-DNA condensate. Extension is a quantity that defines the ratio of the end-to-end distance of the DNA strand, L, to its contour length, L_t (for λ-phage DNA $L_t = 16.5$ μm). Here, FoxA1 mediates the nucleation of a DNA condensate phase, reeling in available slack into the condensate, which modifies the extension of the non-condensed strand: $\frac{L}{L_t - L_d}$. Using this extension, we can obtain an expression for the external force that the strand is subjected to, f, and define a modified version for the free energy of nucleating a FoxA1-DNA condensate:

$$F(L, L_d) = -\mu\alpha L_d + \gamma 4\pi \left(\frac{3\alpha}{4\pi}\right)^{2/3} L_d^{2/3} + \frac{k_B T}{P} \int_0^{L_d} \left(\frac{1}{4}(1 - \frac{L}{L_t - x})^{-2} - \frac{1}{4} + \frac{L}{L_t - x}\right) dx$$

(3)

where k_B is the Boltzmann constant, T is the temperature, and P is the persistence length of DNA. For what follows we define $\kappa = \frac{k_B T}{P}$. Solving the work integral we obtain,

$$F(L, L_d) = -\mu\alpha L_d + \gamma 4\pi \left(\frac{3\alpha}{4\pi}\right)^{2/3} L_d^{2/3} + \frac{\kappa L}{4} \left(\frac{l l_d}{(l-1)(l+l_d-1)} + \log\left(\frac{(l-1)^2}{(1-l_d)^4(l+l_d-1)^2}\right)\right),$$

(4)

where $l = \frac{L}{L_t}$ and $l_d = \frac{L_d}{L_t}$. Using this expression for the free energy, we can vary the end-to-end distance L, and obtain different profiles for the free energy as a function of L_d (see Fig. 3b). For example, at low L values—where the strand is not under much tension—we observe that there is a minimum value for L_d close to L_t. This means that, at this value of L, FoxA1 has mediated the generation of a FoxA1-DNA condensate using almost all of the strand's DNA. As L increases, however, the local minimum shifts to lower values of L_d and ultimately crosses zero, giving rise to a branch of metastable solutions. For even higher L values, the

metastable solution disappears and the global minimum is at $L_d = 0$ dominates (Fig. 3b and 3d). This sharp transition corresponds to a first-order phase transition. Notably, with optimized parameter values (see supplementary methods), the energy gained at low L is on the order of 2-3 $k_B T$, meaning that stochasticity is relevant for the condensation process. To account for the inherent stochastic nature of the condensation, we compute the probability of nucleating a DNA-protein condensate of size L_d using Boltzmann distributions from the corresponding energy profiles,

$$P(L_d) = \frac{e^{-\beta F(L_d)}}{\int_0^{L_t - L} e^{-\beta F(x)} dx} \tag{5}$$

where $\beta = \frac{1}{k_B T}$. To determine the relationship between L_d and L, we compute the mean L_d value of these Boltzmann distributions: $\bar{L}_d(L) = \int_0^{L_t - L} x P(x) dx$ which then allows us also to calculate the magnitude of the capillary forces using the modified worm-like chain model given by f.