Bronchus-associated macrophages efficiently capture and present soluble inhaled antigens and are capable of local Th2 cell activation

Xin-Zi Tang[1,2,3,5], Lieselotte S. M. Kreuk[1,2,6], Cynthia Cho[1,2,7], Ross J. Metzger[4,8], and Christopher D. C. Allen[1,2,4,*]

[1]Cardiovascular Research Institute, University of California, San Francisco, San Francisco, CA, USA
[2]Sandler Asthma Basic Research Center, University of California, San Francisco, San Francisco, CA, USA
[3]Biomedical Sciences Graduate Program, University of California, San Francisco, San Francisco, CA, USA
[4]Department of Anatomy, University of California, San Francisco, San Francisco, CA, USA
[5]Present address: Molecular Engineering Laboratory, Institute of Molecular and Cell Biology, Agency for Science, Technology and Research, Singapore
[6]Present address: Department of Microbiology, University of Washington School of Medicine, Seattle, WA, USA
[7]Present address: Nkarta Therapeutics, South San Francisco, CA, USA
[8]Present address: Department of Pediatrics (Cardiology), Stanford University School of Medicine, Stanford, CA, USA

*Correspondence should be addressed to:
        Christopher D. C. Allen, Ph.D.
        Address: UCSF
                        CVRI, Box 3122
                        555 Mission Bay Blvd S
                        San Francisco, CA 94143, USA
        Email: Chris.Allen@ucsf.edu
        Phone: 415-476-5178

Abstract

In allergic asthma, allergen inhalation leads to local Th2 cell activation and peribronchial

inflammation. However, the mechanisms for local antigen capture and presentation remain

unclear. By two-photon microscopy, we established that soluble antigens in the bronchial airway

lumen were efficiently captured and presented by a population of CD11c$^+$ interstitial

macrophages with high CX3CR1-GFP and MHC class II expression. We refer to these cells as

Bronchus-Associated Macrophages (BAMs) based on their localization underneath the

bronchial epithelium. BAMs were enriched in collagen-rich regions near some airway

branchpoints, where inhaled antigens are likely to deposit. BAMs engaged in extended

interactions with effector Th2 cells and promoted Th2 cytokine production. BAMs were also

often in contact with dendritic cells (DCs). After exposure to inflammatory stimuli, DCs migrated

to draining lymph nodes, whereas BAMs remained lung resident. We propose that BAMs act as

local antigen presenting cells in the lung and also transfer antigen to DCs.

Introduction

Immune surveillance of barrier surfaces is essential for the maintenance of tissue homeostasis

and defense against pathogens. Within the lung, the conducting bronchial airways represent

one of the first sites of exposure to inhaled substances and infectious agents. In allergic asthma,

the inhalation of harmless environmental allergens leads to peribronchial inflammation, in which

the local activation of type 2 helper T cells (Th2 cells) results in effector cytokine production that

promotes pathogenesis (Holgate, 2012; Lambrecht et al., 2019). The Th2 cells are activated by

antigen presenting cells (APCs) that survey the epithelial barrier and capture and present

antigen. However, the types of APCs contributing to surveillance of the conducting airways are

not fully characterized, and the anatomical organization that couples antigen capture to Th2 cell

activation remains unclear.

The lung tissue regions around the conducting airways may be ideal sites for antigen

surveillance and the initiation of immune responses. Unlike alveolar spaces, the conducting

airways do not participate in gaseous exchange but serve to conduct and filter air. As inspired

air passes through airway branchpoints, changes in the direction of airflow promote the

deposition of particulates and the dissolution of soluble substances in the overlying mucus

layer (Carvalho et al., 2011; Clarke and Yeates, 1994). In allergic asthma, eosinophils and

lymphocytes are recruited to regions around the conducting airways, resulting in peribronchial

inflammation (Jeffery, 1992). In several mouse models of airway infections, T and B cells

accumulate in organized foci near the airways, forming induced Bronchus-Associated Lymphoid

Tissue (BALT) structures (Fleige and Forster, 2017; Randall, 2010). The accumulation of

immune cells in tissue regions near the conducting airways may allow streamlined access to

inhaled antigens that have become deposited.

The mechanism by which antigen capture and immune activation occurs at the airways is

unclear. Most studies have focused on the potential contributions of dendritic cells (DCs) to

allergic inflammation in the lung (Wikstrom and Stumbles, 2007), based initially on observations

that MHC-II$^+$ cells with dendritic morphology were abundant along both human and rodent

airways at baseline (Demedts et al., 2005; Huh et al., 2003; Lambrecht et al., 1998; von Garnier

et al., 2007). Flow cytometric analysis of lung-draining lymph nodes, dissociated lung tissue

enriched for the conducting airways, and the trachea also showed that DCs, gated as CD11c$^+$

MHC-II$^+$ cells, were able to capture, process, and present fluorescent ovalbumin (OVA)

administered via the airways (del Rio et al., 2007; Fear et al., 2011; Hammad et al., 2010; Huh

et al., 2003; von Garnier et al., 2007). The uptake of OVA by CD11c$^+$ MHC-II$^+$ cells throughout

the lung has also been observed by confocal microscopy (Hoffmann et al., 2018); however, two-

photon microscopy studies of CD11c-YFP reporter mice found that DCs near the conducting

airways did not detectably capture soluble antigens (OVA or dextran), or particulate antigens in

the form of beads (Thornton et al., 2012; Veres et al., 2011). CD11c-YFP$^+$ DCs were reported to

capture beads at the alveoli in a mouse model of asthma, and then were proposed to migrate to

tissue regions proximal to the bronchial airways, where T cells are activated (Thornton et al.,

2012). This mechanism might be relevant for the uptake and presentation of particulate antigens

that reach the alveoli, but whether soluble antigen can be directly captured and presented by

APCs at the conducting airways remains unclear. The presentation of soluble antigens is

pertinent because it was reported that individuals allergic to aeroallergens had an increased

frequency of Th2 cells that preferentially recognized soluble antigen components of

aeroallergens, whereas regulatory T cells preferentially recognized particulate antigen

components (Bacher et al., 2016). Developing a better understanding of the uptake and

presentation of soluble antigens may therefore be of significant relevance to Th2 cell responses

in asthma.

In the intestine, soluble antigen in the lumen is efficiently captured by CX3CR1$^+$ macrophages in

the lamina propria (Farache et al., 2013a; Farache et al., 2013b; Mazzini et al., 2014; Schulz et

al., 2009). These intestinal macrophages were found to be distinct from conventional DCs

(cDCs) based on functional and behavioral differences, and were reported to contribute to local

intestinal immunity (Hadis et al., 2011; Mazzini et al., 2014; Schreiber et al., 2013; Schulz et al.,

2009). Recent gene expression and microscopy studies of the lung have identified CX3CR1+

interstitial macrophages (IMs) that are distinct from alveolar macrophages (AMs). Notably,

subpopulation(s) of these IMs express many of the surface markers previously thought to be

exclusive to DCs, such as MHC-II and CD11c (Chakarov et al., 2019; Gautier et al., 2012;

Gibbings et al., 2017; Sabatel et al., 2017; Schlitzer et al., 2013; Schyns et al., 2019; Ural et al.,

2020). While most studies on resident tissue macrophages have focused on comparing various

macrophage populations to each other, how lung DCs and IMs compare in their behaviors and

functions is yet unclear.

Here we show that in the steady state in naïve mice, lung myeloid APCs are primarily positioned

underneath the bronchial airway epithelium and are strategically enriched near some airway

branchpoints where inhaled allergens are likely to deposit. We found that at these regions, the

majority of inhaled soluble antigens were captured by CX3CR1$^+$ CD11c$^+$ MHC-II$^+$ IMs rather

than DCs. We observed frequent contacts between CX3CR1$^+$ CD11c$^+$ MHC-II$^+$ IMs and DCs,

suggesting the possibility of antigen transfer to DCs. Yet CX3CR1$^+$ CD11c$^+$ MHC-II$^+$ IMs

themselves were capable of antigen processing and presentation, activated T cells *in vitro*, and

interacted with T cells recruited to the airways in an asthma model. Unlike DCs, CX3CR1$^+$

CD11c$^+$ MHC-II$^+$ IMs did not migrate to the draining lymph node and remained lung resident. We

therefore propose that CX3CR1$^+$ CD11c$^+$ MHC-II$^+$ IMs located near the bronchial airways are

tissue-resident sentinel APCs that are able to drive local allergic inflammation, whereas DCs

may be more critical for priming T cell responses in lymph nodes.

Results

Myeloid cells are enriched near bifurcating airways prior to the induction of inflammation

Peribronchial inflammation is a hallmark of asthma that is recapitulated in mouse models of

allergic airway disease. Foci of leukocytes can be observed near the bronchi that include

numerous eosinophils. In our own studies of mouse models of allergic airway disease, such as

in a classical model with OVA as an allergen (Daubeuf and Frossard, 2013), we observed in thin

cryosections that these foci were not evenly distributed along the airway, but rather were

typically located near airway branchpoints and associated with large blood vessels (Figure 1A).

These areas were further visualized in 3D in agarose-inflated, thick lung sections that were

precision cut with a vibratome and imaged by two-photon laser scanning microscopy (hereafter

referred to as two-photon microscopy). Bronchial airway epithelial cells were revealed by GFP

expression when we crossed Sonic hedgehog (Shh)-Cre mice (Harfe et al., 2004) to ROSA26-

mTmG reporter mice, in which Cre-mediated recombination leads to the expression of

membrane-bound GFP (Muzumdar et al., 2007). Alveolar epithelial cells also express GFP in

these mice, with alveolar type II cells appearing more prominently due to their cuboidal shape,

whereas all other cells express membrane-bound tdTomato. Collagen was detected by second

harmonic generation. Our imaging revealed regions near some airway branchpoints with dense

accumulations of collagen surrounding blood vessels (Figure 1B and Video 1). These regions

likely represent bronchovascular bundles which are known to contain bronchi and arteries in a

common adventitial sheath (Townsley, 2012). Bifurcating blood vessels, visualized by

membrane-bound tdTomato, were also visible at some collagen-rich sites in which the airways

appeared to undergo planar bifurcation (Metzger et al., 2008) (Figure 1B and Video 1).

Various myeloid cells are known to accumulate in peribronchial regions during inflammation. To

visualize myeloid cell infiltration, we imaged thick lung sections from mice expressing the

MacGreen (CSF1R-GFP) reporter (Sasmono et al., 2003) by two-photon microscopy. In

MacGreen mice, we had previously observed that most myeloid cells including macrophages,

monocytes, neutrophils, and CD11b[+] DCs brightly expressed the GFP reporter, although

eosinophils exhibited much lower GFP expression (data not shown). As some B cells were also

found to express GFP in these mice (unpublished observations), we had crossed the MacGreen

mice to B-cell deficient μMT mice so that GFP expression was restricted to myeloid cells. At the

collagen-rich regions near some bifurcating airways, we observed dense accumulations of

myeloid cells with the in both OVA and house dust mite (HDM) models of allergic airway

disease (Figures 1C, 1D). Interestingly, in naïve mice, a sparse population of MacGreen[+] cells

were also observed at these collagen-rich regions near bifurcating airways (Figure 1E),

suggesting that myeloid cells may localize to these areas prior to the formation of foci of

leukocytes during inflammation. Notably, these myeloid cells were best visualized in naïve mice

in thick lung sections by two-photon microscopy, because in thin cryosections the very low

density of cells made it difficult to discern their distribution.

To further evaluate the types of myeloid cells located at these sites in the steady state, we

imaged IMs and DCs using the CX3CR1-GFP and CD11c-YFP reporters, respectively, in thick

lung sections by two-photon microscopy (Gibbings et al., 2017; Jung et al., 2000; Lindquist et

al., 2004; Rodero et al., 2015; Thornton et al., 2012; Veres et al., 2011). In order to specifically

visualize IMs and DCs, our analysis focused on cells with bright expression of these fluorescent

reporters using limiting laser power, as other cell types have weaker expression of these

fluorescent reporters. For example, AMs, which express high amounts of surface CD11c

protein, are known to have low expression of the CD11c-YFP transgene in naïve mice (Rodero

et al., 2015; Thornton et al., 2012). We simultaneously visualized airway epithelial cells with the

E-cadherin-mCFP reporter (Snippert et al., 2010), or with a combination of Shh-Cre with a

ROSA26-loxP-STOP-loxP-tdTomato (Ai14) reporter (Madisen et al., 2010), such that airway

epithelial cells expressed mCFP or tdTomato, respectively. In naïve mice, we observed clusters

of bright CX3CR1-GFP[+] IMs and CD11c-YFP[+] DCs at some airway bifurcations in collagen-rich

regions (Figures 1F and 1H, Videos 2 and 3). Scattered bright CX3CR1-GFP[+] IMs were also

observed underneath the entire bronchial airway epithelium (Figure 1F and Video 2). Bright

CX3CR1-GFP[+] cells were more abundant at the airways than at the alveoli (Figure 1G), and

bright CD11c-YFP[+] DCs were only rarely observed at the alveoli in naïve mice (Figure 1I).

Taken together, these findings indicate that in the steady-state in the absence of inflammation,

both IMs and DCs are located near bronchial airways and may be particularly enriched in

collagen-rich regions near bifurcations. These regions correspond to the sites where eosinophils

and other leukocytes accumulate in allergic airway disease, suggesting a potential role for these

IMs and/or DCs in the initiation of the local inflammatory response.

CX3CR1-GFP+ macrophages capture antigen from the bronchial airways

Our above findings regarding the positioning of CX3CR1-GFP+ IMs and CD11c-YFP+ DCs at

some airway branchpoints may be significant, as antigen is likely to deposit on the walls of the

airways at branchpoints where airflow changes directions. In addition, the distribution of

CX3CR1-GFP+ IMs underneath the entire bronchial airway epithelium suggests that these cells

may generally be involved in surveillance of the airways. We next considered whether inhaled

soluble antigens could be directly captured by the CX3CR1-GFP+ IMs and CD11c-YFP+ DCs

located near the bronchial airways. In order to visualize the uptake of soluble protein antigens

from the airway lumen, we instilled fluorescent OVA intranasally and prepared precision-cut lung

slices 1-2 h later for imaging by two-photon microscopy. A short time point was chosen to

ensure that the antigen capturing cells identified had not migrated from a different anatomical

location, as had been proposed by Thornton et al. (2012). Airway epithelial cells were visualized

by the E-cadherin-mCFP reporter. Our analysis revealed numerous bright CX3CR1-GFP+ IMs

located near the bronchial airways that had taken up fluorescent OVA (Figures 2A, 2B). These

bright OVA+ CX3CR1-GFP+ IMs had a dendritic morphology, extending long processes under

the bronchial airway epithelium. As expected, soluble antigen was also taken up at the alveoli,

primarily by AMs which had a distinct spherical appearance (Figure 2—figure supplement 1).

Thus, CX3CR1-GFP+ IMs with a dendritic morphology readily captured soluble antigen from the

bronchial airway lumen and were distinct from AMs.

In contrast to the CX3CR1-GFP+ IMs near the bronchial airways, the CD11c-YFP+ DCs at these

sites did not visibly appear to capture OVA by microscopy (Figure 2C). To confirm preferential

uptake in CX3CR1-GFP+ IMs rather than CD11c-YFP+ DCs, we administered fluorescent OVA

intranasally to mice simultaneously expressing both CX3CR1-GFP and CD11c-YFP reporters.

This combination of fluorescent reporters has been helpful in the intestine to distinguish DC and

macrophage subsets (Farache et al., 2013a; McDole et al., 2012), and we optimized

methodology to minimize the spectral overlap of GFP and YFP by two-photon microscopy (see

Methods section for details). Our imaging of dual reporter mice confirmed that OVA was

preferentially captured by bright CX3CR1-GFP[+] IMs rather than bright CD11c-YFP[+] DCs (Figure

2D). Quantification of OVA uptake indicated that CD11c-YFP[+] DCs captured much smaller

amounts of OVA than CX3CR1-GFP[+] IMs (Figure 2E). We also observed preferential uptake of

fluorescently labelled extracts from HDM, a relevant aeroallergen, by bright CX3CR1-GFP[+] IMs

(Figure 2F). Our imaging also confirmed that bright CX3CR1-GFP[+] and bright CD11c-YFP[+] cells

were largely non-overlapping populations (Figure 2D, 2F). When we increased the image

contrast, we observed that some of the bright CX3CR1-GFP+ cells did have dim expression of

CD11c-YFP, and vice versa, as expected from previous studies (Jung et al., 2000; Sen et al.,

2016) (Figure 2—figure supplement 2A). Occasionally, we also observed rare cells that were

bright for both the CX3CR1-GFP and CD11c-YFP reporters, yet the majority of cells in the naïve

state appeared to be only bright for only one of these reporters suggesting they were distinct

cell types (Figure 2—figure supplement 2B). Indeed, CX3CR1-GFP[+] IMs had a dendritic

morphology as we had noted above, whereas CD11c-YFP[+] DCs were rounder and did not

appear to extend long processes. We quantified these morphologic features by measuring

sphericity as well as the length of the longest axis of each cell to characterize the dendritic

projections. CX3CR1-GFP[+] IMs had lower sphericity values (Figure 2G) and increased mean

lengths of their longest axes (Figure 2H) compared to CD11c-YFP[+] DCs. The disparity in

morphology and antigen uptake capacity suggests that CX3CR1-GFP[+] IMs and CD11c-YFP[+]

DCs are two distinct cell populations present in airway-adjacent areas in the steady state, and

that CX3CR1-GFP[+] IMs are the primary antigen capturing cell type at the airways.

We confirmed these microscopy observations by flow cytometric analysis of CX3CR1-GFP

versus CD11c-YFP reporter mice after instillation of fluorescent OVA. This analysis confirmed

that the average intensity of fluorescent OVA was over three-fold brighter in CX3CR1-GFP[bright]

IMs compared with CD11c-YFP^bright DCs (Figure 2—figure supplement 3A, 3B). Microscopy

detects a comparably narrower dynamic range of fluorescent signals than flow cytometry, and

thus we most likely only visualized CX3CR1-GFP^bright IMs that carried the most OVA. Indeed,

when we gated specifically on OVA^bright MHC-II^+ cells by flow cytometry, the majority of cells

were CX3CR1-GFP^bright but not CD11c-YFP^bright (Figure 2—figure supplement 3C, 3D).

Therefore, our results show that DCs also take up soluble antigen from the bronchial airway

lumen, but that the majority of antigen is taken up by a distinct population of CX3CR1-GFP^+

IMs.

CX3CR1-GFP^bright antigen capturing cells are Bronchus-Associated Macrophages (BAMs)

We next evaluated whether the CX3CR1-GFP^+ cells that captured soluble inhaled antigen

expressed MHC-II and confirmed their macrophage identity by flow cytometry. This was

important given that CX3CR1-GFP is also expressed in other cell types, such as monocytes, a

subset of DCs, and some lymphocytes (Jung et al., 2000). For this analysis, we excluded AMs

using the marker Siglec-F, as these cells often confound the gating of GFP^+ cells by flow

cytometry due to their autofluorescence, yet based on their round morphology and alveolar

localization were clearly distinct from the antigen-capturing CX3CR1-GFP^+ cells near the

airways (Figure 2—figure supplement 1). After administering fluorescent OVA intranasally and

then isolating and enzymatically digesting the lungs, our flow cytometric analysis showed that

the CX3CR1-GFP^bright OVA^bright cells were predominantly MHC-II^hi and CD11c^mid (Figure 3A).

The presence of these cells in naïve mice and high MHC-II expression suggested they were not

derived from inflammatory monocytes. The CX3CR1-GFP^bright OVA^bright cells also predominantly

expressed the macrophage markers CD64 and MerTK, and had relatively low to undetectable

staining for the cDC marker CD24 (Figures 3A, 3B), consistent with surface markers previously

established for IMs (Gautier et al., 2012; Guilliams et al., 2013; Schlitzer et al., 2013).

Conversely, the majority of the CD11c-YFP^{bright} cells were MHC-II$^+$, CD11c^{mid-hi} and CD24^{hi} but

did not express CD64 and MerTK, thus representing cDCs (Figures 3A, 3B). Notably, CD64$^+$

OVA^{bright} cells were largely CD11c-YFP$^-$ but did express surface CD11c by antibody staining

(Figure 3—figure supplement 1A), albeit with reduced abundance compared to cDCs, indicating

the expression of the CD11c-YFP transgene is more restricted to DCs than the CD11c protein,

consistent with a prior study (Rodero et al., 2015). The small fraction of CD64$^+$ OVA^{bright} cells

that were also CD11c-YFP^{bright} (Figure 3—figure supplement 1A) may correspond to the rare

cells we had observed by microscopy that were bright for both CX3CR1-GFP and CD11c-YFP

(Figure 2—figure supplement 2B). We found that monocytes, which resolve into two subsets

based on CX3CR1-GFP expression (Geissmann et al., 2003), and the CX3CR1-GFP$^+$ subset of

cDC2s, all had lower CX3CR1-GFP expression than the MHC-II$^+$ CD64$^+$ MerTK$^+$ cells (Figure

3—figure supplement 1B), consistent with our imaging data (Figure 2—figure supplement 2A).

Again, as microscopy has a more limited dynamic range than flow cytometry and we had

focused our imaging on bright cells using limiting laser power, this likely favored our specific

visualization of CX3CR1-GFP^{bright} cells. To indeed verify that the CX3CR1-GFP^{bright} cells we had

observed near the bronchial airway epithelium by microscopy were represented by the

CX3CR1-GFP^{bright} CD64$^+$ MHC-II$^+$ IMs identified by flow cytometry, we stained precision-cut

thick lung slices from CX3CR1-GFP or CD11c-YFP mice with antibodies against CD64 and

MHC-II. By two-photon microscopy, we confirmed that the CX3CR1-GFP^{bright} cells located near

the bronchial airways were CD64$^+$ (Figures 3C and 3D). In contrast, the CD11c-YFP^{bright} cells

were CD64-negative (Figure 3E). Most of the CX3CR1-GFP^{bright} cells located near the airways

were MHC-II$^+$ (Figures 3C solid arrowheads and Figure 3D), although occasional MHC-II$^{lo/—}$

cells were also visualized (Figure 3C, open arrowheads). Taken together, these findings

indicate that the majority of the CX3CR1-GFP^{bright} cells localized near the bronchial airway

epithelium that capture soluble inhaled antigens are CD11c$^+$ CD64$^+$ MHC-II$^+$ IMs. Based on their

spatial localization near the bronchial airways, we propose that this subset of IMs be referred to

as Bronchus Associated Macrophages (BAMs).

We further characterized BAMs in an immunophenotyping analysis by flow cytometry comparing

CD64$^+$ MHC-II$^+$ cells to cDCs and AMs. CD64$^+$ MHC-II$^+$ BAMs had high surface expression of

CD11b and were uniformly MerTK$^+$, but had only minimal expression of a Zbtb46-GFP reporter

that was nearly 10-fold lower than CD24$^+$ DCs (Figure 3—figure supplement 2A-C), further

confirming these cells are not DCs despite sharing similar markers to cDC2s. Whereas AMs had

high surface expression of CD169 and Siglec-F as expected, we found that CD64$^+$ MHC-II$^+$

BAMs had low CD169 expression that partially overlapped with background staining, and did

not express Siglec-F (Figure 3—figure supplement 2D, 2F), further supporting that these are

two distinct macrophage populations.

To compare gene expression profiles between BAMs and other CD11c$^+$ cells, particularly DCs

and AMs, we further analyzed these populations by single cell RNA sequencing (scRNAseq)

analysis, using a multi-step protocol we devised to enrich for CD11c$^+$ cells from the lungs of

unimmunized mice (see Methods for details). Given the large abundance of CD11c$^+$ AMs in the

lung, we first depleted some of these cells to help enrich for IM and DC subsets. This allowed us

to sequence hundreds of IMs and DCs from each sample. A substantial number of AMs still

remained, allowing us to do a comparison of IMs, DCs, and AMs. Most of the cells sequenced

expressed *Ptprc*, the gene encoding CD45, confirming most of the cells analyzed were

leukocytes, with minor populations of structural cells (Figure 4A). Unsupervised clustering of

cells visualized by UMAP revealed multiple clusters of AMs, which expressed *Siglecf*, with one

predominant grouping of clusters that we defined as AMs (Figure 4A and Figure 4-supplement

1). One major cluster of IMs was detected, marked by the highest density of expression of

Cx3cr1, consistent with our imaging analysis (Figure 4B and Figure 4-supplement 1). The IM

cluster also had high expression of *Mafb* (Figure 4B), which is thought to largely be restricted to

cells of the macrophage lineage (Wu et al., 2016), with the exception of AMs (Gautier et al.,

2012; Gibbings et al., 2017). Within the major IM cluster two distinct populations were visible.

BAMs make up the majority of the population of IMs, which expressed comparable amounts of

H2-Ab1 (MHC-II) to the DC subsets, consistent with our flow cytometric analysis, as well as

Itgax (CD11c), *Mmp12* and *Ccr2* (Figure 4B). A minor population of IMs was visible with high

expression of *Lyve1* and *Folr2*, with weaker expression of *H2-Ab1* and *Itgax* (Figure 4B),

consistent with previous reports (Chakarov et al., 2019; Schyns et al., 2019; Ural et al., 2020).

Mrc1 (CD206) was widely expressed among IMs, with the highest expression on the *Lyve1*+

population (Figure 4B). *Siglec1* (CD169) was only detected in a small fraction of cells in the IM

cluster (Figure 4B), suggesting reads were near the threshold for detection, consistent with our

CD169 staining by flow cytometry (Figure 3—figure supplement 2D). Several clusters of DCs

were identified by a high density of expression of *Flt3* and *Zbtb46* (Figure 4C). These clusters

showed differential expression of *Mgl2*, *Cd207*, and *Ccr7*, suggesting they represented cDC2s,

cDC1s, and migratory DCs, respectively (Izumi et al., 2021; Sichien et al., 2017). We grouped

these DC clusters together for comparison to AMs and IMs (Figure 4-supplement 1). Several

genes were identified that were most highly expressed in IMs compared to AMs and DCs: these

included genes encoding chemokines (*Cxcl13*, *Ccl7*, and *Ccl2*), complement pathway

components (*C1q* genes and *C3ar1*), *Apoe*, and *Mmp12* (Figure 4D). This transcriptional profile

analysis demonstrates that despite similarities in their surface marker profiles, BAMs, AMs, and

DCs are functionally distinct subsets of cells.

Previous studies have reported lung IMs can be divided into subsets based on CD206

expression (Chakarov et al., 2019; Gibbings et al., 2017; Schyns et al., 2019). One of these

studies reported that airway associated IMs were largely MHC-II⁻ and CD206⁺, whereas MHC-

II⁺ CD206⁻ IMs were predominantly located near the alveolar parenchyma (Schyns et al., 2019).

300 Our microscopy data above, however, had clearly demonstrated CX3CR1-GFP^{bright} MHC-II$^+$ IMs

301 were positioned near the airways. Our scRNAseq data above further indicated that MHC-II$^+$ IMs

302 express *Mrc1*, the gene encoding CD206, although *Mrc1* was most highly expressed in the

303 *Lyve1$^+$* subset of IMs. Our flow cytometric analysis of CD64$^+$ MHC-II$^+$ cells showed that BAMs

304 had intermediate CD206 surface expression in a Gaussian distribution (Figure 3—figure

305 supplement 2E). By two-photon microscopy of precision-cut thick lung slices, we observed that

306 the CX3CR1-GFP^{bright} BAMs that predominantly captured antigen (OVA) near the airways were

307 CD206$^+$ (Figure 5A), although the expression of CD206 was variable. By co-staining with MHC-

308 II, we observed that CX3CR1-GFP^{bright} MHC-II^{hi} cells near the airways (Figure 5B, solid

309 arrowheads) expressed variable intermediate amounts of CD206, whereas the highest CD206

310 expression was observed on some CX3CR1-GFP$^+$MHC-II^{lo} cells that were also located near the

311 airways (Figure 5B, open arrowheads). Taken together, these data establish that BAMs, which

312 capture antigen near the airways and are MHC-II$^+$, express variable amounts of CD206.

313 Two recent studies reported that some IMs in the lung were located near nerves (Chakarov et

314 al., 2019; Ural et al., 2020), although this was less apparent in another study (Schyns et al.,

315 2019). Notably, bronchovascular bundles contain bronchi, arteries, lymphatic vessels, and

316 nerves. By two-photon microscopy of precision-cut thick lung slices stained with an antibody to

317 L1CAM, a neuronal cell adhesion molecule (Rathjen and Schachner, 1984; Tacke et al., 1987),

318 we readily visualized nerves near the bronchial airways and at airway branchpoints where

319 BAMs were abundant (Figures 5C and 5D). A subset of BAMs were directly in contact with

320 nerves, while other BAMs were not in contact with nerves (Figures 5C and 5D). One recent

321 study defined a subset of IMs as nerve and airway associated macrophages (NAMs) (Ural et al.,

322 2020). However, this population of NAMs was notable for very high CD169 expression, yet our

323 flow cytometric and scRNAseq analyses suggested the antigen-capturing BAMs near the

324 airways were low to negative for CD169 expression (Figure 3—figure supplement 2D and

Figure 4B). In two-photon microscopy of precision-cut thick lung slices, we observed punctate

CD169 staining on BAMs that was only detectable above background in a subset of cells

(Figure 5D, solid arrowheads). In contrast, we occasionally observed cells with bright uniform

CD169 staining in contact with nerves; however, these cells were not CX3CR1-GFP[bright] and

thus were distinct from BAMs (Figure 5D, open arrowhead). Interestingly, the punctate CD169

staining on BAMs was polarized toward one side of the cell, and this often seemed to be at the

interface with a CD11c-YFP[+] DC (Figure 5E). A previous study demonstrated an important role

for CD169 in cell-cell contacts between macrophages and DCs in the spleen and lymph nodes,

which enhanced antigen transfer and cross-presentation (van Dinther et al., 2018). In our static

imaging data above, we noted that BAMs and DCs were frequently in contact (see Figure 2D,

2F, Figure 2—figure supplement 2, and Figure 3E). We observed extensive, dynamic

interactions between BAMs and DCs in time-lapse imaging of precision-cut lung slices by two-

photon microscopy (Video 4). This suggests the possibility that BAMs may transfer antigen, or

peptide-MHC II complexes, to DCs. Taken together, in addition to their role in antigen capture

from the airways, BAMs may interact with nerves and/or communicate with DCs.

BAMs primarily capture lower molecular weight soluble proteins

We next further characterized the ability of BAMs to take up antigens from the airway lumen. To

compare the antigen uptake capacities of BAMs to individual subsets of DCs and monocytes,

fluorescent OVA was administered to mice intranasally 2 h before lungs were digested and

dissociated into single cell suspensions for flow cytometric analysis. We gated on CD45[+] CD19[−]

Ly6G[−] Siglec-F[−] CD11c[+] MHC-II[+] CD11b[+] CD24[−/lo] CD64[+] cells as a reasonable representation

of BAMs, as we had established based on our marker analysis above. Indeed, a large majority

(~80%) of BAMs gated by these markers had taken up OVA, whereas a smaller proportion

(~50%) of cDC2s and less than 5% of monocytes were OVA[+] (Figure 6A). By microscopy, we

observed that in addition to soluble OVA and HDM extract, CX3CR1-GFP$^+$ BAMs took up other

soluble fluorescent molecules of a range of sizes, such as TRITC-dextran molecules that were

10 kDa or 155 kDa, as well as phycoerythrin (PE), a 240 kDa molecule (Figures 6B-6D). To

quantify whether antigen size affected the efficiency of antigen capture by BAMs, we

administered FITC-dextran (FITC-DX) of various sizes ranging from 10 kDa to 2 MDa, and then

assessed the frequency of FITC-DX$^+$ cells among BAMs (Figure 6E, left) and cDC2s (Figure 6E,

right) by flow cytometry. To exclude the possibility that FITC-DX was taken up during the lung

digestion process, lungs from congenic mice that were not exposed to FITC-DX (control lung)

were processed in the same digestion reaction as the lungs from the mice that had received

FITC-DX (host lung). Very few (or less than 5% of) BAMs from the control lungs of congenic

mice that were co-digested were FITC-DX$^+$, confirming that minimal FITC-DX uptake occurred

during the digestion procedure (Figure 6E). We found that a greater proportion of BAMs were

FITC-DX$^+$ when smaller molecules were instilled compared to larger molecules (Figure 6E).

Therefore, the magnitude of antigen uptake by BAMs is inversely related to the antigen size.

It has been reported that DCs in the alveolar interstitium send projections into the alveolar

space to capture fluorescent bead particles (Thornton et al., 2012). To assess whether BAMs

captured particulate antigens from the bronchial airway lumen, we intranasally instilled

fluorescent beads and analyzed the lungs by microscopy. However, we rarely observed bead

capture by CX3CR1-GFP$^+$ BAMs or CD11c-YFP$^+$ DCs near the airways (Figure 6F). In contrast,

in the alveolar space, beads were taken up primarily by highly spherical cells that were dimly

CD11c-YFP$^+$ (Figure 6F). By location and morphology, we concluded that these cells were AMs,

not DCs. In summary, BAMs preferentially captured soluble antigens, whereas both BAMs and

DCs residing at the airway interstitium showed minimal capture of beads.

BAMs activate local effector T cells

Based on our findings that BAMs readily captured soluble antigens from the bronchial airways

and had high surface expression of MHC-II comparable to DCs, we considered whether BAMs

are capable of activating T cells at these sites. In order to activate T cells, APCs need to

process antigen and present antigen peptides on MHC-II on their cell surface. We used the Y-

Ae antibody to test whether BAMs were able to process and present antigen. The Y-Ae antibody

specifically recognizes the Eα peptide, a peptide fragment of the I-E MHC-II molecule,

presented on the I-A^b MHC-II molecule (Murphy et al., 1992). C57BL/6 mice do not express I-E

and thus Eα represents a foreign peptide. The Eα peptide was fused to OVA, administered

intranasally together with fluorescent OVA to label antigen-capturing cells, and then OVA$^+$ lung

APCs were analyzed 17-18 h later to allow time for antigen processing and presentation. Both

BAMs and cDC2s stained with the Y-Ae antibody by flow cytometry (Figure 7A), demonstrating

that BAMs processed and presented antigen, like DCs, rather than simply degrading it. Minimal

Y-Ae staining was observed 15 min after OVA-Eα was administered, confirming that antigen

processing and presentation were needed for detection of MHC-II:Eα complexes on the cell

surface (data not shown). Brighter Y-Ae staining was observed on BAMs compared with cDC2s

(Figure 7A), consistent with the superior soluble antigen uptake we had observed (Figures 2

and 6). AMs, with low MHC-II expression and a high degradative capacity, showed negligible Y-

Ae staining (Figure 7A), confirming these are not the relevant APCs for CD4 T cells. We also

tested whether BAMs could provide the necessary co-stimulatory signals for T cell activation

and found that BAMs had higher surface expression of the co-stimulatory molecules CD80 and

CD86 compared to cDC2s (Figure 7B). To determine whether BAMs were able to activate CD4

T cells, BAMs were sorted by FACS and co-cultured with OT-II TCR transgenic T cells, which

respond to an OVA peptide presented on I-A^b by APCs. As a positive control for APC function in

cell culture, cDC2s were sorted and co-cultured with OT-II T cells. When cultured in the

presence of OVA peptide, BAMs induced the robust proliferation of naïve OT-II T cells, albeit

somewhat less efficiently than cDC2s, as expected, given the high efficiency of naïve T cell

activation by DCs (Figure 7C). Interestingly, BAMs and cDC2s equivalently induced IL-13

production when co-cultured with Th2 polarized, effector OT-II cells and OVA peptide (Figure

7D). These findings indicate that BAMs possess the necessary machinery for antigen

presentation and activation of T cells, and further suggest that BAMs may equivalently activate

effector T cells compared with DCs.

We hypothesized that the primary function of MHC-II expression on BAMs might be to activate

local effector/memory T cells that have been recruited to the lung tissue. To test this hypothesis,

we utilized time-lapse two-photon microscopy to determine whether BAMs encountered and

interacted with effector T cells in the lungs. Naïve OT-II T cells expressing CFP (β-actin CFP) or

tdTomato (CD4-Cre Ai14) were adoptively transferred into CX3CR1-GFP X CD11c-YFP mice

and then primed *in vivo*, which resulted in expansion and recruitment to the lungs. Mice were

subsequently rechallenged with intranasal OVA alone, lungs were excised 4 h later, and

precision-cut lung slices were prepared for imaging. After allergen challenge, we observed

sustained interactions between OT-II T cells and CX3CR1-GFP[+] BAMs in the lung (Figures 7E-

G, Videos 5 and 6). These interactions between BAMs and OT-II T cells were comparable to the

interactions reported during antigen-specific T cell activation by DCs in the lymph

nodes (Mempel et al., 2004). Some OT-II T cells were also observed to engage in sustained

interactions with CD11c-YFP[+] DCs near the airways (data not shown). These observations

suggest that both BAMs and DCs can serve as APCs for the local activation of effector T cells in

lung peribronchial regions.

BAMs are non-migratory while DCs are migratory

Although BAMs and DCs both exhibited the capacity to serve as APCs for T cells, time-lapse

imaging by two-photon microscopy revealed fundamental differences in cellular dynamics.

BAMs had dendrites that were continuously extended and probing, but their cell bodies showed

very minimal displacement over time. In contrast, DCs were surveying their surroundings by

moving their entire cell bodies and traversed greater distances than BAMs (Figures 8A-D and

Video 7). Therefore, DCs were highly migratory whereas BAMs were probing by extending

dendrites with little net movement.

Based on our observations of their distinct migratory behavior, we hypothesized that BAMs may

remain at the airways, whereas DCs would migrate to the draining lymph node to prime T cells,

after antigen capture that leads to an inflammatory response. However, in other studies, cells

with similar markers as BAMs, referred to as monocyte-derived DCs, had been observed in the

draining lymph node harboring antigen (Plantinga et al., 2013). We therefore tested whether

BAMs were able to migrate to the right posterior mediastinal lymph node that drains the lung.

We first considered the expression of the chemokine receptor CCR7, which is necessary for

migration to lymph nodes and is upregulated following exposure to inflammatory stimuli. We

analyzed the cell surface expression of CCR7 on DCs and BAMs after administering LPS or

HDM as inflammatory stimuli. After 4 h, we found that CCR7 was upregulated on a significant

proportion of DCs in a dose dependent manner, but not on BAMs (Figures 8E-F). We also

tested whether OVA, which does not have intrinsic inflammatory properties, would induce CCR7

upregulation in previously sensitized mice in which OVA-specific effector/memory T cells could

induce maturation of the APCs. Indeed, after OVA challenge, DCs upregulated CCR7

expression in previously sensitized mice but not in naïve mice exposed to OVA for the first time

(Figure 8G). Conversely, BAMs did not upregulate CCR7 in either sensitized or naïve mice

(Figure 8G). As such, the expression of CCR7 only on the DCs, but not BAMs, would likely

allow DCs to selectively migrate to the lymph node. To further assess whether DCs or BAMs

entered and migrated in lymphatic vessels within the lung by two-photon microscopy, we

visualized lymphatics by breeding Prox1-tdTomato transgenic mice (Truman et al., 2012) to

CD11c-YFP or CX3CR1-GFP mice. After intranasal instillation of HDM or LPS, we frequently

observed CD11c-YFP$^+$ cells, but not CX3CR1-GFP$^+$ cells, migrating within lung lymphatics

(Figure 8H, Videos 8 and 9). To determine which cells carried antigen into the lymph node, we

intranasally instilled PE. With a molecular weight of 240 kDa, PE is too large to enter the conduit

system of the lymph node (Roozendaal et al., 2009) and thus can only reach the lymph node

parenchyma when carried by APCs. As an inflammatory stimulus to promote APC trafficking to

the lymph node, we administered HDM extract together with the PE. This led to the recruitment

of a population of CD64$^+$ MHC-II$^+$ cells that were Ly6C$^+$ to the draining mediastinal lymph node

1 d later. However, these CD64$^+$ MHC-II$^+$ Ly6C$^+$ cells were PE$^-$, whereas we were readily able

to identify a fraction of DCs that were PE$^+$ (Figure 8I-J). Thus, the CD64$^+$ MHC-II$^+$ Ly6C$^+$ cells

were not transporting PE from the lung to the lymph node. The high surface expression of Ly6C

on the CD64$^+$ MHC-II$^+$ cells also suggests that they were not BAMs, which have low to

moderate surface expression of Ly6C (Figure 8—figure supplement 1). Instead, the high

expression of Ly6C suggests that the CD64+ MHC-II+ cells were derived from monocytes that

were recruited to the lymph node from the bloodstream and had upregulated CD64 and MHC-II

in response to the inflammatory conditions. Taken together, these findings indicate that BAMs

capture antigen at the airways but do not migrate to the draining lymph node. In contrast, DCs

are induced to migrate to the draining LN in response to inflammatory stimuli or in the context of

adaptive immune responses.

Discussion

This work focused on the cell types involved in antigen surveillance at the airways in the steady

state to understand the early events that initiate allergic airway inflammation. Our analysis of

APCs at the airways revealed an abundant population of MHC-II^{hi} IMs positioned under the

bronchial airway epithelium that sampled soluble luminal antigen, which we term Bronchus-

Associated Macrophages (BAMs). We found that BAMs were particularly enriched in collagen-

rich regions near some airway branchpoints and engaged in extensive interactions with DCs. In

addition, we demonstrated that BAMs were capable of antigen presentation to T cells and

interacted with recruited T cells near the airways in an allergic airway disease model. While DCs

migrated to the draining lymph node, BAMs remained resident in the lung after antigen capture,

even in the presence of inflammatory stimuli. These findings support a model in which BAMs

may locally activate effector/memory T cells recruited to the lung, whereas migratory DCs are

likely critical for T cell priming in lymph nodes.

In characterizing the localization of lung cDCs and BAMs, we found that in naïve mice, both cell

types were positioned around the conducting airways, and were especially enriched at some

airway branchpoints in collagen-rich regions. This positioning likely facilitates the uptake of

antigens that have deposited at airway branchpoints, where airflow changes directions, leading

to the activation of adaptive immune responses at these locations. We found that these sites

corresponded to the regions where inflammatory clusters form in allergic airway disease

models. BALT has also been observed near airway bifurcations (Bienenstock and McDermott,

2005; Randall, 2010). Our observation that BAMs and DCs were enriched near airway

branchpoints in the steady state, prior to immunization, suggests that airway branchpoints are

ideal sites for continuous immunosurveillance of the bronchial airway lumen, enabling the rapid

local activation of antigen-specific T cells and the generation of an inflammatory response.

While most of the lung is filled with alveolar space to support gaseous exchange, these

collagen-rich adventitial regions of connective tissue and interstitial space in the

bronchovascular bundles represent a site where immune cells may be recruited. Recent work

indicates that similar interstitial spaces may be found in other organs and that the adventitial

"cuff" regions may generally be important sites for tissue immunity (Benias et al., 2018;

Dahlgren and Molofsky, 2019).

While both BAMs and DCs are strategically positioned in the lung for immunosurveillance, they

differ substantially in the efficiency of antigen uptake in the steady state. Consistent with

previous imaging studies of CD11c-YFP mice (Thornton et al., 2012; Veres et al., 2011), we

observed minimal fluorescent antigen uptake by CD11c-YFP$^+$ DCs near the airways by

microscopy. Our quantitative analysis of microscope images and flow cytometry revealed that

CD11c-YFP$^+$ DCs were nevertheless taking up low amounts of soluble antigen. In contrast,

BAMs took up higher amounts of antigen that were readily visible by imaging. In a similar

manner, IMs are superior at capturing soluble antigen from the intestinal lumen compared to

DCs (Mazzini et al., 2014; Schulz et al., 2009). . Although one group reported equivalent uptake

of antigen by DCs and IMs in the lung (Hoffmann et al., 2018), this was likely due to

contamination of their DC population, which was defined as CD11c$^+$ MHC-II$^+$ cells, by BAMs.

We have shown here that BAMs express surface CD11c and MHC-II, yet BAMs exhibit distinct

antigen uptake, morphology, and motility characteristics compared with cDCs. Considering that

BAMs look more dendritic than lung cDCs and are MHC-II^{hi}, it is likely that the MHC-II$^+$

dendritiform cells identified in rodent airways in earlier studies (Fear et al., 2011; Huh et al.,

2003; von Garnier et al., 2007) were actually BAMs rather than DCs. The stellate shape of

BAMs is reminiscent of recent imaging studies of IMs in other tissues, such as in the

kidney (Stamatiades et al., 2016), and may facilitate antigen sampling.

We observed that BAMs engaged in frequent, extensive interactions with DCs. The punctate

localization of CD169 at the interacting surface between BAMs and DCs suggests this cell

contact has biological significance, given precedent for a role for CD169 in antigen transfer from

macrophages to DCs in the spleen and lymph nodes that contributed to cross-presentation (van

Dinther et al., 2018). Phagocytosed antigen could potentially be passed from BAMs to DCs,

similar to intestinal macrophages which have been shown to pass antigen to DCs through gap

junctions (Mazzini et al., 2014). Peptide-MHC complexes could also potentially exchange from

BAMs to DCs through trogocytosis. BAMs could thus play an important role in

immunosurveillance, both directly and through antigen transfer to DCs.

During inflammatory exposures, BAMs stay resident near the bronchial airways, whereas some

DCs migrate to the draining lymph node. Consequently, BAMs may play an important role in the

local activation of effector/memory T cells recruited to the lung. After initial sensitization, effector

T cells were recruited to the airway interstitium where BAMs were abundant as early as 4-6 h

after allergen challenge and formed long interactions with BAMs, reminiscent of activating

interactions previously observed between T cells and DCs in the lymph node (Mempel et al.,

2004). These interactions may lead to the production of chemotactic factors that recruit

additional cells to these sites forming inflammatory foci. Thus, the recruitment and activation of

T cells at the bronchial airway interstitium, where BAMs display captured antigen, would provide

a basis for the initiation of allergic inflammation during allergen re-exposure in sensitized

individuals. In the intestine and adipose tissue, resident macrophages were also reported to

activate effector T cells and shape local immunity (Cho et al., 2014; Kamada et al., 2009;

Schreiber et al., 2013); BAMs may be the functional equivalent of these macrophages at the

airways.

Here we reported the rapid and efficient uptake of soluble antigen across a range of sizes by

BAMs, with the magnitude of antigen uptake inversely proportional to antigen size. We did not

readily observe dendrite extension through the epithelial barrier by live microscopy, making this

unlikely to account for the widespread antigen uptake by BAMs, and therefore suggesting that

BAMs capture soluble antigens by alternative mechanism(s). Soluble antigens may instead be

captured after diffusing across the airway epithelial barrier. This model may account for the

reduced uptake of larger molecular weight antigens, which would be less effective at diffusing

across the epithelium. In the intestine, it was reported that low molecular weight soluble

antigens passed through goblet cells (McDole et al., 2012). Although few goblet cells are

present in the murine lung in the steady state, whether soluble antigens could also pass through

the more abundant club cells remains to be determined. The increase in goblet cells that occurs

in allergic airway disease might also facilitate the passage of soluble antigens and thus further

promote the inflammatory response. At sites of BALT formation, the follicle associated

epithelium containing microfold (M) cells has also been reported to facilitate the uptake of

soluble antigens (Bienenstock and Befus, 1984). Alternatively, soluble antigen may pass

through gaps in the airway epithelium, such as tri-junctional complexes located where three

epithelial cells meet (Patton, 1992). The 200 nm bead particles that we tested would be too

large to pass through the gaps in tri-junctional complexes that average 27 nm, consistent with

our observation of minimal bead uptake around the airways by either BAMs or DCs.

Interestingly, one study reported that very small 20 nm beads were captured by some MHC-II[+]

cells underneath the tracheal epithelium, although the majority of beads of a range of sizes were

captured by AMs in the lung parenchyma (Blank et al., 2013). Indeed, we also observed that

200 nm beads were primarily captured within the alveolar spaces by AMs. Another study

observed the uptake of 1000 nm beads by DCs located at the alveoli, but not by DCs located

near the conducting airways (Thornton et al., 2012). In that study, previously sensitized mice

were rechallenged with antigen while beads were administered, creating an inflammatory state

in which monocyte-derived cells and DCs (Jenkins and Hume, 2014; Rodero et al., 2015) likely

were recruited to the alveolar regions and captured beads. In the steady state, DCs are largely

restricted to the regions near bronchial airways, yet monocytes have also been reported to

patrol the alveoli and capture beads (Rodero et al., 2015). Overall, our work reveals that at the

bronchial interstitium, small soluble antigens are preferentially captured by BAMs.

The form and size of an antigen was recently reported to affect the type of T cell responses

induced, as type 2 allergic responses were primarily directed to small soluble antigens whereas

regulatory T cell responses were primarily directed to particulate antigens (Bacher et al., 2016),

though the mechanism was not explored. Our finding that soluble and particulate antigens are

preferentially captured in different regions of the lung leads us to propose that the anatomic

location of antigen uptake ultimately influences the final T cell response. The tissue signals

inherent in different anatomic environments may prime divergent immune responses. In the

alveoli, the abundant AMs have been proposed in several studies to dampen allergic

responses (Bang et al., 2011; Mathias et al., 2013; Tang et al., 2001; Zaslona et al., 2014).

Macrophages that were earlier found in the alveolar interstitium were also reported to produce

IL-10 and reduce inflammation in allergic airway disease models (Bedoret et al., 2009; Kawano

et al., 2016). The tissue environment at the alveoli thus appears to favor reduced inflammation

in the delicate alveolar tissue, in the absence of overt infection in this region. Instead, the

airways may be more permissive to the induction of inflammation as indicated by the

concentration of immune responses at airway adjacent regions. Future work on the functional

capacities and/or cytokine profiles of BAMs would further inform our understanding of the

inflammatory milieu at the airway microenvironment.

We propose that BAMs represent a subset of IMs specialized for sampling the bronchial airway

lumen. This work extends the findings of Gibbings et al. (2017) and Rodero et al. (2015) who

first observed CX3CR1-GFP[+] IMs localized near the bronchial airways. IM populations near the

alveoli have also been reported in microscopy studies (Bedoret et al., 2009; Schyns et al., 2019;

Tan and Krasnow, 2016). Multiple subsets of IMs have also been described based on different

combinations of surface markers by flow cytometric analysis, as well as distinct transcriptional

profiles (Chakarov et al., 2019; Gibbings et al., 2017; Sabatel et al., 2017; Schyns et al., 2019).

There is a lack of consensus on the identification and localization of the IM subsets, and it

seems likely that diverse pathogen exposures in different mouse facilities may affect baseline

inflammation and the relative abundance of different subsets. This may explain, for example,

why we observed few CD169[bright] macrophages near the airways and nerves, whereas another

group found these were abundant (Ural et al., 2020). Of note, most of our mice for this study

were bred in a 'super-barrier' mouse facility that excludes additional pathogens compared to

standard SPF facilities. Here we have focused specifically on the subpopulation of IMs localized

near the bronchial airway epithelium that we found readily captures soluble inhaled antigens in

naïve mice. We propose that the field consider referring to these macrophages as BAMs based

on their anatomic localization near the bronchial airways, much as it has been useful to classify

AMs based on their anatomic location. We found that BAMs expressed high amounts of MHC-II;

were efficient at antigen capture, processing and presentation; and interacted with effector T

cells *in vivo*; suggesting that BAMs serve as APCs near the bronchial airways. Our finding that

BAMs expressed high amounts of surface MHC-II differs from a recent study by Schyns et

al (2019), which reported MHC-II[+] CX3CR1-GFP[+] IMs were primarily localized near the alveoli.

This difference is likely due to technical differences in our imaging approaches. We imaged

native GFP fluorescence in unfixed tissue allowing us to preferentially visualize CX3CR1-

GFP[bright] BAMs, whereas Schyns et al. used anti-GFP antibody amplification of CX3CR1-GFP

signal in fixed tissue, which would have likely resulted in the visualization of a much larger

population of cells with varying expression of CX3CR1-GFP, including monocytes and DCs. We

also imaged thick tissue vibratome sections by two-photon microscopy whereas Schyns et al.

imaged thin cryosections by confocal microscopy. We found the analysis of thick sections to be

particularly useful to try to discern the cellular distribution of BAMs in unimmunized mice,

whereas in thin sections the cells were too sparse.

Emerging data from microarray and RNAseq analyses have highlighted the distinct gene

expression patterns of IMs, AMs and DCs in the lung (Chakarov et al., 2019; Gautier et al.,

2012; Gibbings et al., 2017; Schlitzer et al., 2013; Schyns et al., 2019), which we have further

elaborated here by our scRNAseq analysis of CD11c$^+$ cells. Functional studies in other tissues

such as the intestine have revealed distinct contributions of macrophages and DCs to

immunity (Chang et al., 2013; Kamada et al., 2009; Mazzini et al., 2014; Schreiber et al., 2013;

Schulz et al., 2009; Shaw et al., 2012). Here we have shown that BAMs, as a subset of IMs, are

distinct from lung cDCs in morphology, antigen uptake capacity and the ability to migrate to the

draining lymph node. It is important to note that under inflammatory conditions, including in

mouse models of allergic airway disease, monocytes rapidly enter lung tissue and differentiate

into monocyte-derived cells that express surface markers that bear some resemblance to

resident DCs and macrophages (Guilliams et al., 2013; Jakubzick et al., 2008; Plantinga et al.,

2013). Although the MAR-1 antibody to FcɛRI was thought to be a marker of these monocyte-

derived cells, we recently established that this antibody also binds to CD64 and FcγRIV (Tang

et al., 2019), and thus would also detect resident CD64$^+$ macrophages. Recent studies of

respiratory viral infection have also revealed the recruitment of a subpopulation of cDCs to the

lung that stain with MAR-1 and express CD64 (Bosteels et al., 2020), further confounding the

use of these markers to distinguish macrophages and cDCs in the context of inflammation. One

microscopy study of the lungs of mice administered HDM reported an increase in the

abundance of CD64$^+$ "DCs" identified by CD11c and MHC-II expression (Hoffmann et al., 2018),

which likely consisted of a heterogeneous mixture of cDCs, resident BAMs and monocyte-

derived cells. Another study used an extensive analysis of the ratio of expression of CX3CR1-

GFP and CD11c-mCherry reporters to determine the relative differentiation state of infiltrating

monocytes (Sen et al., 2016). This paper showed an increase in the abundance of cells with

high expression of both CX3CR1-GFP and CD11c-mCherry reporters in allergic airway

inflammation models. These cells were proposed to represent differentiated monocyte-derived

cells, and we speculate that the rare cells we observed that were bright for CX3CR1-GFP and

CD11c-YFP reporters in naïve mice may also represent cells that had recently differentiated

from monocytes. One challenge, however, in relating this study to our findings presented here is

that the high expression of CX3CR1-GFP in resident BAMs versus the lower expression in

monocytes was not considered in the analysis of the ratio of CX3CR1-GFP to CD11c-mCherry

expression, which formed the basis of the monocyte differentiation model (Sen et al., 2016).

Similar to BAMs and DCs, monocyte-derived cells have been reported to activate T cells and

drive allergic inflammation (Plantinga et al., 2013). Conversely, monocyte-derived cells and IMs

have also been reported to promote tolerance and dampen inflammation in the lung (Bedoret et

al., 2009; Kawano et al., 2016; Sabatel et al., 2017). Although all these APCs may exhibit some

overlap in their ability to activate T cells, it seems likely that these different cell types may skew

T cell responses at the airways in distinct ways. Further studies with selective tools for

manipulation will be needed to elucidate the specific contributions of BAMs, DCs, and

monocyte-derived cells to lung immunity.While we have focused on the potential role of BAMs

as APCs in immune surveillance of the bronchial airways, our findings do not exclude other

potential functions of these cells. The association of some BAMs with nerves at

bronchovascular bundles is likely of significance to tissue function and/or immunity. BAMs might

also be involved in the clearance of epithelial cells or in the production of molecules involved in

tissue homeostasis. It will be interesting for future studies to resolve the potential distinctions

and functions of the various subsets of IMs recently defined in the lung.

In summary, our study demonstrates that the bronchial airways are a major site of soluble

antigen capture where BAMs serve as resident sentinel immune cells. Compared to DCs, BAMs

exhibited distinct morphology and enhanced soluble antigen uptake from the airways. BAMs

also closely interacted with cDCs, suggesting the potential for antigen transfer between these

cell types. In addition, DCs migrated to the draining lymph nodes whereas BAMs remained

resident near the airways. This disparity in migratory behavior suggests a division of labor

where migratory cDCs may prime naïve T cells in the lymph nodes while BAMs mediate local

airway immunity. Our data demonstrate that BAMs are able to activate effector Th2 cells and

interact with CD4 T cells recruited to the airways in mouse models of allergic airway disease.

We propose that BAMs may play a similar role in the local activation of T cells in context of

infection. Overall, BAMs are strategically positioned as sentinel APCs underlying the bronchial

airway epithelium. BAMs were enriched at collagen-rich regions near some airway

branchpoints, which may ultimately develop inflammatory foci when an adaptive immune

response is initiated.

Methods

Mice and chimeras

C57BL/6J (B6/J; 000664), μMT (002288; B6.129S2-*Ighm*^{tm1Cgn}/J), CX3CR1-GFP (005582;

B6.129P-*Cx3cr1*^{tm1Litt}/J), CD4-Cre (022071; B6.Cg-Tg(Cd4-cre)$^{1Cwi/Bflu}$/J), Shh-Cre (005622;

B6.Cg-*Shh*$^{tm1(EGFP/cre)Cjt}$/J), Ai14 (007914; B6.Cg-Gt(*ROSA)26*Sor$^{tm14(CAG-tdTomato)Hze}$/J), E-cadherin-

mCFP (016933; B6.129P2(Cg)-*Cdh1*^{tm1Cle}/J), MacBlue (026051; STOCK Tg(Csf1r*-

GAL4/VP16,UAS-ECFP)^{1Hume}/J), ROSA26-mTmG (007676; B6.129(Cg)-*Gt(ROSA)26Sor*$^{tm4(ACTB-}$

$^{tdTomato,-EGFP)Luo}$/J), Prox1-tdTomato (022766; C57BL/6-Tg(Prox1-tdTomato)12^{Nrud}/J), Thy1.1

(000406; B6.PL-*Thy1*^{a}/CyJ) and Zbtb46-GFP (018534; 129S-*Zbtb46*$^{tm1.1Kmm}$/J) mice were

originally from The Jackson Laboratories. All mice were maintained on B6 background except

Zbtb46-GFP, which was obtained on a 129 background and crossed one generation to B6/J.

MacGreen (CSF1R-GFP, Tg(Csf1r-EGFP)1^{Hume}) (Sasmono et al., 2003), CD11c-YFP (Tg(Itgax-

Venus)1^{Mnz}) (Lindquist et al., 2004), β-actin-CFP (Tg(CAG-ECFP)CK6^{Nagy}) (Hadjantonakis et al.,

2002), and OT-II (Y-chromosome insertion; Tg(TcraTcrb)426-6^{Cbn}) (Barnden et al., 1998) mice

were maintained on a B6 background. B6-CD45.1 congenic mice, which were used as wild-type

mice in some experiments, were bred from the Boy/J line (002014; B6.SJL-*Ptprc*^{a}*Pepc*^{b}/BoyJ)

from The Jackson Laboratories or purchased from Charles River (National Cancer Institute

Model 01B96; B6-Ly5.2/Cr, later renamed to B6-Ly5.1/Cr). All mice were maintained in specific-

pathogen-free facilities and protocols were approved by the Institutional Animal Care and Use

Committee of the University of California San Francisco. MacGreen mice were crossed to μMT

mice to generate a faithful reporter of myeloid cells because some B cells were also GFP^{+}

(unpublished observations). Fluorescent OT-II T cells were obtained by crossing OT-II mice to

β-actin-CFP mice or CD4-Cre X Ai14 mice.

Antigen administration

Intranasal (i.n.) droplets were administered to mice that had been lightly anesthetized with

isoflurane using a controlled vaporizer. To induce allergic airway inflammation with a classical

OVA model (Daubeuf and Frossard, 2013) for Figure 1A, mice were first sensitized by

intraperitoneal (i.p.) injections with 100 μg Endograde OVA (Hyglos) mixed with 100 μl alum

(Alhydrogel, Grade A 1.3%, Accurate Chemical and Scientific) once a week for a total of three

times. One week after the last sensitization, mice were challenged i.n. with 50 μg Endograde

OVA in 50 μl PBS daily over three days. lungs (see details below) were assessed two days after

the last challenge. For Figure 1C, 10^4 OT-II cells were adoptively transferred i.v. and then mice

were primed i.p. with 100 μg OVA grade VII (Sigma-Aldrich, this lot was found to have minimal

endotoxin activity) in 200 μl alum on d0 and d9, and then challenged i.n. with 100 μg OVA in 50

μl PBS on d16, 17, and 18, and then imaged on d20. In repeats of this particular experiment,

the mice were primed once rather than twice with OVA/alum, as this led to greater recovery of

OT-II cells and a similar inflammatory response. For experiments represented in Figure 1D,

mice received 10-100 μg of HDM extract (Greer) reconstituted in PBS i.n. once a week for a

total of four times (Kohl et al., 2006). Lung slices were imaged 2 days after the last exposure.

For live imaging, an abbreviated i.n. model of OVA allergic airway inflammation (Eisenbarth et

al., 2002) was used to study early events after allergen challenge. Mice were sensitized i.n. with

100 μg Endograde OVA mixed with low-dose (100 ng) LPS (ultrapure LPS, O55:B5, List

Biologicals) in 50 μl of PBS and OT-II cells were adoptively transferred i.v. on the same day. 9-

10 days later, mice were then challenged i.n. with 50 μg OVA in 50 μl of PBS , followed by

excision of lungs 4-6h after challenge.

For microscopy, fluorescent OVA conjugated to Texas Red or Alexa 647 (Life Technologies)

was administered in 20-50 μl PBS i.n. Smaller volumes were found to somewhat favor labeling

at the larger airways rather than at the alveoli, but also led to a more heterogeneous distribution

of antigen throughout the lung. To produce HDM conjugated to Texas Red or OVA conjugated

to TAMRA, 100 μg of HDM or OVA were reconstituted in PBS with 0.1 M sodium bicarbonate at

1 mg/ml and mixed with 18 μg Texas Red-succinimidyl ester or 36 μg 5,(6)-TAMRA-succinimidyl

ester (Life Technologies), respectively. Excess free Texas Red or TAMRA were removed with

Bio-spin 6 kDa size exclusion columns (Bio-Rad Laboratories) equilibrated with PBS according

to manufacturer's instructions. We noted that Texas Red-labeled antigens bound nonspecifically

to dead epithelial cells at the cut site of the vibratome sections, but this was not observed with

TAMRA- or Alexa 647-labeled antigens. However, we observed similar capture of all types of

fluorophore-labeled antigens by BAMs, confirming this result was not fluorophore-dependent.

TRITC-Dextran and FITC-Dextran were from Life Technologies (10 kDa) or Sigma-Aldrich (all

other molecular weights). R-Phycoerythrin (PE) (Life Technologies) was dialyzed for 48 h at 4°C

into PBS using 10 kDa MWCO Slide-a-lyzer cassettes (Thermo Fisher) before i.n.

administration. 0.2 μm Sky Blue fluorescent beads (Spherotech) were administered i.n. at

0.025-0.05% w/v in 30 μl PBS. To induce CCR7 upregulation and track lymphatic migration

(Figure 8), mice received either 1 μg ultrapure LPS or 100 μg HDM in 30-50 μl of PBS i.n., or

were primed and challenged with the abbreviated low-dose LPS i.n. OVA model described

above, and lungs were analyzed 4-6 h later.

To generate OVA-Eα, an Eα peptide (Murphy et al., 1992) synthesized with an added terminal

cysteine residue (New England Peptide) was dissolved in ultrapure water with 5 mM EDTA and

mixed with maleimide-activated OVA (Thermo Fisher Scientific) at room temperature for 2 h.

Excess Eα peptide was removed with Bio-spin 30 kDa size exclusion columns (Bio-Rad

Laboratories) equilibrated with PBS according to the manufacturer's instructions. 20 μg OVA-Eα

was administered i.n. together with 2.5 μg of OVA-Alexa 647 (Life Technologies) in 30-50 μl

PBS and lungs were analyzed 17-18 h later. For imaging of T cell activation, 5 x 10^5 fluorescent

OT-II T cells (see Mice and Chimeras above) were adoptively transferred i.v. into the retro-

orbital plexus of anesthetized CX3CR1-GFP X CD11c-YFP recipient mice, which were then

sensitized and challenged with the abbreviated low-dose LPS i.n. OVA model described above,

and lungs were excised for imaging 4-6 h after challenge.

Cryosection and immunofluorescence microscopy

Mouse lungs were inflated with 1 mL of Tissue-Tek optimum cutting temperature compound

(OCT, Sakura), then placed in a cryomold with OCT and "snap-frozen" in dry ice and ethanol.

Frozen tissue blocks were stored at -80°C. Acetone-fixed cryosections were prepared as

described (Allen et al., 2004). Slides were stained as described (Yang et al., 2018) in PBS or

TBS containing 0.1% BSA (Roche), 1% normal mouse serum (Jackson Immunoresearch) and a

diluted mixture of purified Siglec-F PE (E50-2440, BD) and SMA-FITC (1A4, Abcam) for 2 h.

Slides were washed for 3 min in PBS or TBS and mounted in Fluoromount G with DAPI

(SouthernBiotech). Slides were imaged on the Axio Scan.Z1 slide scanner equipped with an

HXP120 fluorescence microscope illuminator as well as 38 HE and 20 HE filter sets for FITC

and PE visualization (Carl Zeiss Microscopy). A Chroma filter set (GT370/60x, T425lpxr,

ET460/50m) was used for DAPI visualization. Images were collected and analyzed with Zen

Blue (Carl Zeiss Microscopy).

2-photon microscopy

In preparation for 2-photon microscopy, mouse lungs were inflated with 1 mL of 2% low melt

agarose, and 250-600 μm thick sections were precision cut using a vibratome (Leica VT1200 or

Oxford Laboratories Model G) as described (Sullivan et al., 2011) and then placed in Leibovitz's

L-15 medium at room temperature until imaging. Airways could be identified by transmitted light

based on the presence of a columnar epithelial lining and active ciliary beating. Imaging was

performed within several hours with the exception of slices stained with antibodies. For antibody

staining, lung slices were incubated at room temperature in PBS with TruStain FcX (Biolegend)

diluted to 5 μg/mL, mouse gamma globulin (Jackson ImmunoResearch) diluted to 100-400

μg/mL, and in some cases rat gamma globulin (Jackson ImmunoResearch) diluted to 100-200

μg/mL for 2 h and then stained with the antibodies (see Supplementary file 1) overnight at 4°C.

Lung slices were then washed in excess PBS before imaging. For time-lapse microscopy, lung

slices were incubated as described (Sullivan et al., 2011). Briefly, lungs slices were mounted in

a heated chamber (JG-23W/HP, Warner Instruments) with warm RPMI (Gibco) supplemented

with 1X Penicillin, Streptomycin, and L-Glutamine (Life Technologies) flowing over it at a rate of

1.5 mL/min. Media temperature in the imaging chamber was maintained at 35-37°C. 5%

CO_2/95% O_2 was bubbled into RPMI to maintain physiologic pH.

Lung slices were imaged on an LSM 7 MP INDIMO two-photon laser scanning microscope (Carl

Zeiss Microscopy) equipped with two Chameleon Ultra II lasers (Coherent), a Compact Optical

Parametric Oscillator (OPO, Coherent) and a W Plan-Apochromat 20x/1.0 objective. Images

were collected on non-descanned detectors consisting of four gallium arsenide phosphide

(GaAsP) detectors and one far-red sensitive detector. Transmitted illumination was detected on

a T-PMT. Emission filters used to record fluorescence signals are listed in Supplementary file 2.

Wavelengths used to excite particular fluorophores are listed in Supplementary file 3. Sequential

laser excitation at different wavelengths was used to enhance the spectral separation of

fluorophores, similar to our previous approach (Sullivan et al., 2011) with CFP and GFP.

Samples were imaged with multiple tracks, with rapid line or frame switching of laser emissions

by acousto-optical modulators in order to separate the excitation and detection of different

fluorophores and for second harmonic generation. For example, in order to improve the

separation of GFP and YFP: in the first track, one laser was tuned to 850-870 nm to excite CFP

and GFP with minimal excitation of YFP; and then in the second track, the other laser was tuned

to 1020-1040 nm to strongly excite YFP with minimal excitation of CFP and GFP. In some

cases, a third track was added with the OPO tuned to 1100-1200 nm to strongly excite PE

and/or Alexa 647 with minimal excitation of GFP/YFP. Second harmonic generation results in

an emission at half the incident laser wavelength and the signal was collected in various filters

(Supplementary file 2); the separation of the second harmonic from fluorophore emissions was

assisted by tuning lasers to particular wavelengths and the use of multiple tracks. For

experiments in which enhanced spectral separation was not needed, or for fast imaging for

time-lapse recordings, fluorophores were excited simultaneously with one or more laser lines. Z-

stacks were collected every 30 s for 20-45 min. *xy* planes were collected at variable zoom and

pixel sizes, with a typical resolution of 512 x 512 pixels (some high resolution images were

collected at 1024 x 1024 pixels). Bidirectional scanning was typically used for faster image

acquisition. The spacing between planes in the *z* dimension was typically set to three to five

times the *xy* pixel size for a given zoom setting. Images were collected with Zen Black (Carl

Zeiss Microscopy) and images were analyzed with Zen Black, Volocity (PerkinElmer) and Imaris

software (Bitplane). Videos were prepared with Volocity and then annotated in Adobe After

Effects CC.

Flow cytometry

To collect bronchoalveolar lavages (BAL), mice were euthanized with an overdose of isoflurane

and lungs were deflated and exposed by thoracotomy. A 20G x 1" SurFlash catheter (Terumo)

was inserted into the trachea and secured with a suture. Lungs were flushed twice with a

syringe with 1 mL of DPBS (Gibco) and then the lavage was added to 200 µl of PBS with 50%

FBS and 0.1 M EDTA (Gibco) and kept on ice. BALs were centrifuged at 350g for 7 min at 4°C

and resuspended in PBS with 2% FBS, 1 mM EDTA (Gibco) and 0.1% sodium azide (Gibco)

(FACS buffer) for antibody staining. Lungs were excised into 1 mL DMEM media supplemented

with HEPES (Gibco), Penicillin, Streptomycin and L-Glutamine (Life Technologies). For lung

digestion, 25 µg of DNase I (Sigma) and 0.28 Wunitz Units of Liberase TM (Roche) were added

to each mL of digest. Lungs were snipped into small pieces with scissors in a 1.7 mL eppendorf

tube and incubated on a 37°C shaker for 30-45 min at 700 rpm on a ThermoMixer F1.5

(Eppendorf). The digestion reaction was quenched by adding FBS and EDTA to a final

concentration of 5% FBS and 10 mM EDTA. Digested lungs were then gently dissociated and

passed through a 70 μm strainer to yield single cell suspensions that were then centrifuged and

resuspended in DMEM supplemented with 5% FBS, HEPES, Penicillin, Streptomycin and L-

Glutamine. For FACS staining, 3 x10^6 lung cells were added to a 96-well plate and washed with

150 μl of ACK Lysing Buffer (Quality Biological) before antibodies were added for staining. Flow

cytometric staining of BAL and lung cells was done as described (Allen et al., 2004; Yang et al.,

2012), except CCR7 was stained for 15 min at 37°C. Antibodies used for staining are listed in

Supplementary file 1. Flow cytometric data was acquired on a BD LSRFortessa with 405nm,

488nm, and 640nm lasers, using BD FACS DiVa software and analyzed with FlowJo software

(TreeStar).

scRNAseq analysis

Lungs were excised from four naive mice (two males and two females) and digested and

dissociated as above. MacBlue mice were used to allow for the analysis of CFP expression in

future studies. Single cell suspensions of two lung samples from mice of the same sex were

pooled and washed with a 2X volume of ACK Lysing Buffer (Quality Biological) as above to lyse

red blood cells and then were resuspended in PBS with 0.5% BSA (Roche) and 2 mM EDTA

(Gibco) (MACS buffer). Cells were depleted of B cells and partially depleted of AMs and cells

that had lost membrane integrity by first incubating with anti-IgD Biotin (Southern Biotech) and

anti-Siglec-F Biotin, as well as 0.25 μM of NHS-Biotin (Life Technologies, freshly reconstituted

in PBS), for 30 min on ice. Cells were washed in MACS buffer and incubated with 50 μl of

MyOne Streptavidin T1 Dynabeads (Thermo Fisher) for 30 min, during which tubes were

inverted and rotated at 4°C. Tubes were placed in a Dynamag-2 magnet (Thermo Fisher) for 2

min for bead isolation and the negative fraction containing unlabeled cells was collected and

diluted with MACS buffer to 5 ml. To further enrich for macrophages and DCs and to remove

stromal cells, 3ml of 14% OptiprepTM (Axis-Shield) was layered below the cell suspension, and

cells were centrifuged at 600 g for 20 min with no brake at 8°C. Cells at the interface were

collected and washed in MACS buffer. Finally, CD11c+ cells were enriched by incubating with

CD11c Mojosort Nanobeads (Biolegend) for 15 min at 4°C. Bead-labeled cells were purified with

the Mojosort magnet (Biolegend) according to the manufacturer's instructions. Bead-labeled

cells were washed in PBS with 0.5% BSA (Roche), washed, and submitted to the UCSF

Institute for Human Genetics for single cell RNA sequencing using 10x Genomics Chromium

technology. Data was processed in Cell Ranger, v5.0.1 (10x Genomics) through count and

aggregate pipelines according to manufacturer's instructions and visualized using Loupe Cell

Browser 5.0 (10x Genomics).

APC sorting

Lungs were excised from CD45.1 congenic mice into DMEM media supplemented with HEPES,

Penicillin, Streptomycin and L-Glutamine and digested and dissociated as above. Single cell

suspensions of lung digests were combined and then a 2X volume of ACK Lysing Buffer

(Quality Biological) was added and cells were centrifuged at 350g for 7 min at 4°C and

resuspended in PBS with 0.5% BSA (Roche) and 1% FBS (Sorting Buffer). Cells were

incubated with CD11c BV650 antibody together with CD11c Mojosort Nanobeads (Biolegend),

after which bead-labeled cells were purified with the Mojosort magnet (Biolegend) according to

the manufacturer's instructions. Cells were then washed and stained with sorting antibodies

(Viability dye eFluor780, CD45 Alexa 700, Ly6G PE-Cy7, Siglec-F BV421, CD19 PE-Dazzle

594, CD11b BV785, MHC-II I-A^b FITC, CD64 PE, CD24 APC) at 4 x10^7 cells/ml. Cells were

washed and diluted with Sorting Buffer, centrifuged through a 1 mL layer of 100% FBS,

resuspended in Sorting Buffer and passed through a 40 μm cell strainer. Cells were sorted on

the FACSAria III (BD) into Advanced RPMI (Gibco) with 20% FBS.

T cell culture

Inguinal, brachial, axillary and cervical lymph nodes were excised from OT-II mice with a Thy1.1

congenic marker. The tissues were then gently dissociated and passed through 40 μm strainers

to yield single cell suspensions that were centrifuged at 350g for 7 min at 4°C and resuspended

in DMEM supplemented with 5% FBS, HEPES, Penicillin, Streptomycin and L-Glutamine. To

purify CD4 T cells by negative selection, cells were resuspended at 1-2 x10^8 cells/ml and

incubated with biotinylated antibodies against B220, CD11b, CD11c, CD8, NK1.1 and MHC-II I-

A^b for 30 min on ice. Cells were washed to remove excess antibodies and then incubated with

250 μl MyOne Streptavidin T1 Dynabeads (Thermo Fisher) per 10^8 cells for 30 min, during

which tubes were inverted and rotated at 4°C. Tubes were placed in a Dynamag-2 magnet

(Thermo Fisher) for 2 min for bead isolation and the negative fraction containing unlabeled cells

was collected and washed. For Cell Trace Violet labeling, cells were centrifuged and

resuspended in PBS with 2% FBS at 2-20 x10^6 cells/ml. A 10 mM stock of Cell Trace Violet

(Thermo Fisher) in DMSO was diluted to 10 μM in PBS and added at 1:1 volumetric ratio to cells

while vortexing. Cells were incubated with Cell Trace Violet for 20 min at 37°C, then 100 μl of

100% FBS was layered at the bottom of the tube and cells were then centrifuged and

resuspended in Advanced RPMI (Gibco) culture media. 50,000 labelled naïve T cells were co-

cultured with 10,000 sorted APCs for 2.5 days in the presence or absence of 20 μg/mL

OVA$_{323-339}$ peptide (Invivogen) before analysis.

For Th2 polarization, CD4 T cells were purified by negative selection with magnetic beads as

described above and cultured in Advanced RPMI (Gibco) supplemented with 10% FBS,

HEPES, Penicillin, Streptomycin, L-glutamine, 1mM sodium pyruvate and 50 mM 2-

mercaptoethanol (culture media). Flat-bottom 96-well plates were coated with 2.5 µg/mL anti-

mouse CD3e (eBioscience, functional grade purified, eBio500A2) in PBS for 3 h at 37°C and

washed twice with PBS before use. T cells were added to anti-CD3 coated wells at 10^6 cells/mL

in the presence of 1 µg/mL anti-CD28 (Biolegend, LEAF purified, 37.51), 100 U/ml IL-2 (gift from

K. Mark Ansel's lab), 50 ng/mL IL-4 (Peprotech) and 10 µg/ml anti-IFNγ (Biolegend, LEAF

purified) for 3 days. T cells were then transferred into uncoated wells and replenished with IL-2,

IL-4 and anti-IFNγ for another 3 days before co-culturing with sorted APCs. 50,000 T cells were

co-cultured with 10,000 sorted APCs in 96-well V-bottom plates (Cellstar) in the presence or

absence of 20 µg/mL $OVA_{323-339}$ peptide (Invivogen). To assess cytokine production, 3 µg/mL

Brefeldin A (eBioscience) was added to cultures 1-2 h after the start of culture, for 14-18 h

before flow cytometric analysis. T cells were analyzed by flow cytometry as described above.

For intracellular cytokine staining, cells were fixed and permeabilized with Cytofix/cytoperm

solution (BD) for 20 min on ice and then washed with Perm/Wash solution (BD) according to the

manufacturer's instructions. Cells were stained with anti-cytokine antibodies (Supplementary file

1) in Perm/Wash solution for 20 min on ice. Cells were washed twice with Perm/Wash solution

and resuspended in FACS buffer, acquired on LSR Fortessa (BD) using FACS DiVA software

(BD) and analyzed with FlowJo software (Treestar).

Statistical analysis

Statistical analyses and graphical representations were done using GraphPad Prism v7 or v8.

Data were expected to have a log-normal distribution and all data were log transformed for

parametric statistical tests. The appropriate tests were chosen based on experimental design

after consulting the GraphPad Statistics Guide. All tests were two-tailed. Multiple biological

replicates and imaging fields were analyzed to draw conclusions. The number of samples

analyzed was based on the expected magnitude of differences, which was estimated from past

similar experiments or pilot studies. Results were further validated by repeating experiments

and/or by performing related experiments. Detailed information on samples and experiments is

provided in each figure legend. Full statistical results with exact p values for all figures are

provided in Supplementary file 4.

Acknowledgments

We thank D. Sheppard for providing CD11c-YFP mice; Z. Werb for providing β-actin-CFP and

MacGreen mice; H. Chapman, J. Cyster, R. Locksley, J-S. Shin, D. Sheppard, and X. Huang for

helpful advice; J. Gordon and the UCSF Laboratory for Cell Analysis (supported by National

Cancer Institute Cancer Center Support Grant P30CA082103) for cell sorting; and M. Shenoy

for contributions to some early experiments. Research reported in this publication was

supported by the National Heart, Lung, and Blood Institute (DP2HL117752) and the National

Institute of Allergy and Infectious Diseases (R21AI130495) of the National Institutes of Health,

the UCSF Cardiovascular Research Institute, the UCSF Sandler Asthma Basic Research

Center, and the UCSF Program for Breakthrough Biomedical Research, which is partially

funded by the Sandler Foundation. X.T. was funded by the Agency for Science, Technology and

Research (A*STAR) National Science Scholarship (PhD). L.S.M.K. was supported by a National

Institute of Allergy and Infectious Diseases training grant (T32AI007334-31). The content is

solely the responsibility of the authors and does not necessarily represent the official views of

the funding agencies.

References

Allen, CD, Ansel, KM, Low, C, Lesley, R, Tamamura, H, Fujii, N, and Cyster, JG. 2004. Germinal center dark and light zone organization is mediated by CXCR4 and CXCR5. *Nat Immunol* **5**: 943-952. doi:10.1038/ni1100

Bacher, P, Heinrich, F, Stervbo, U, Nienen, M, Vahldieck, M, Iwert, C, Vogt, K, Kollet, J, Babel, N, Sawitzki, B, Schwarz, C, Bereswill, S, Heimesaat, MM, Heine, G, Gadermaier, G, Asam, C, Assenmacher, M, Kniemeyer, O, Brakhage, AA, Ferreira, F, et al. 2016. Regulatory T Cell Specificity Directs Tolerance versus Allergy against Aeroantigens in Humans. *Cell* **167**: 1067-1078 e1016. doi:10.1016/j.cell.2016.09.050

Bang, BR, Chun, E, Shim, EJ, Lee, HS, Lee, SY, Cho, SH, Min, KU, Kim, YY, and Park, HW. 2011. Alveolar macrophages modulate allergic inflammation in a murine model of asthma. *Exp Mol Med* **43**: 275-280. doi:10.3858/emm.2011.43.5.028

Barnden, MJ, Allison, J, Heath, WR, and Carbone, FR. 1998. Defective TCR expression in transgenic mice constructed using cDNA-based alpha- and beta-chain genes under the control of heterologous regulatory elements. *Immunol Cell Biol* **76**: 34-40. doi:10.1046/j.1440-1711.1998.00709.x

Bedoret, D, Wallemacq, H, Marichal, T, Desmet, C, Quesada Calvo, F, Henry, E, Closset, R, Dewals, B, Thielen, C, Gustin, P, de Leval, L, Van Rooijen, N, Le Moine, A, Vanderplasschen, A, Cataldo, D, Drion, PV, Moser, M, Lekeux, P, and Bureau, F. 2009. Lung interstitial macrophages alter dendritic cell functions to prevent airway allergy in mice. *J Clin Invest* **119**: 3723-3738. doi:10.1172/JCI39717

Benias, PC, Wells, RG, Sackey-Aboagye, B, Klavan, H, Reidy, J, Buonocore, D, Miranda, M, Kornacki, S, Wayne, M, Carr-Locke, DL, and Theise, ND. 2018. Structure and Distribution of an Unrecognized Interstitium in Human Tissues. *Sci Rep* **8**: 4947. doi:10.1038/s41598-018-23062-6

Bienenstock, J, and Befus, D. 1984. Gut- and bronchus-associated lymphoid tissue. *Am J Anat* **170**: 437-445. doi:10.1002/aja.1001700316

Bienenstock, J, and McDermott, MR. 2005. Bronchus- and nasal-associated lymphoid tissues. *Immunol Rev* **206**: 22-31. doi:10.1111/j.0105-2896.2005.00299.x

Blank, F, Stumbles, PA, Seydoux, E, Holt, PG, Fink, A, Rothen-Rutishauser, B, Strickland, DH, and von Garnier, C. 2013. Size-dependent uptake of particles by pulmonary antigen-presenting cell populations and trafficking to regional lymph nodes. *Am J Respir Cell Mol Biol* **49**: 67-77. doi:10.1165/rcmb.2012-0387OC

Bosteels, C, Neyt, K, Vanheerswynghels, M, van Helden, MJ, Sichien, D, Debeuf, N, De Prijck, S, Bosteels, V, Vandamme, N, Martens, L, Saeys, Y, Louagie, E, Lesage, M, Williams, DL, Tang, SC, Mayer, JU, Ronchese, F, Scott, CL, Hammad, H, Guilliams, M, et al. 2020. Inflammatory Type 2 cDCs Acquire Features of cDC1s and Macrophages to Orchestrate Immunity to Respiratory Virus Infection. *Immunity* **52**: 1039-1056 e1039. doi:10.1016/j.immuni.2020.04.005

Carvalho, TC, Peters, JI, and Williams, RO, 3rd. 2011. Influence of particle size on regional lung deposition--what evidence is there? *Int J Pharm* **406**: 1-10. doi:10.1016/j.ijpharm.2010.12.040

Chakarov, S, Lim, HY, Tan, L, Lim, SY, See, P, Lum, J, Zhang, XM, Foo, S, Nakamizo, S, Duan, K, Kong, WT, Gentek, R, Balachander, A, Carbajo, D, Bleriot, C, Malleret, B, Tam, JKC, Baig, S, Shabeer, M, Toh, SES, et al. 2019. Two distinct interstitial macrophage populations coexist across tissues in specific subtissular niches. *Science* **363**: eaau0964. doi:10.1126/science.aau0964

Chang, SY, Song, JH, Guleng, B, Cotoner, CA, Arihiro, S, Zhao, Y, Chiang, HS, O'Keeffe, M, Liao, G, Karp, CL, Kweon, MN, Sharpe, AH, Bhan, A, Terhorst, C, and Reinecker, HC. 2013.

Circulatory antigen processing by mucosal dendritic cells controls CD8(+) T cell activation.
Immunity **38**: 153-165. doi:10.1016/j.immuni.2012.09.018

Cho, KW, Morris, DL, DelProposto, JL, Geletka, L, Zamarron, B, Martinez-Santibanez, G,
Meyer, KA, Singer, K, O'Rourke, RW, and Lumeng, CN. 2014. An MHC II-dependent
activation loop between adipose tissue macrophages and CD4+ T cells controls obesity-
induced inflammation. *Cell Rep* **9**: 605-617. doi:10.1016/j.celrep.2014.09.004

Clarke, SW, and Yeates, D. 1994. Deposition and Clearance. In Murray, JF, and Nadel, JA,
editors. *Textbook of Respiratory Medicine.* W.B. Saunders Company, p345-369.

Dahlgren, MW, and Molofsky, AB. 2019. Adventitial Cuffs: Regional Hubs for Tissue Immunity.
Trends Immunol **40**: 877-887. doi:10.1016/j.it.2019.08.002

Daubeuf, F, and Frossard, N. 2013. Acute Asthma Models to Ovalbumin in the Mouse. *Curr*
Protoc Mouse Biol **3**: 31-37. doi:10.1002/9780470942390.mo120202

del Rio, ML, Rodriguez-Barbosa, JI, Kremmer, E, and Forster, R. 2007. CD103- and CD103+
bronchial lymph node dendritic cells are specialized in presenting and cross-presenting
innocuous antigen to CD4+ and CD8+ T cells. *J Immunol* **178**: 6861-6866.
doi:10.4049/jimmunol.178.11.6861

Demedts, IK, Brusselle, GG, Vermaelen, KY, and Pauwels, RA. 2005. Identification and
characterization of human pulmonary dendritic cells. *Am J Respir Cell Mol Biol* **32**: 177-184.
doi:10.1165/rcmb.2004-0279OC

Eisenbarth, SC, Piggott, DA, Huleatt, JW, Visintin, I, Herrick, CA, and Bottomly, K. 2002.
Lipopolysaccharide-enhanced, toll-like receptor 4-dependent T helper cell type 2 responses
to inhaled antigen. *J Exp Med* **196**: 1645-1651. doi:10.1084/jem.20021340

Farache, J, Koren, I, Milo, I, Gurevich, I, Kim, KW, Zigmond, E, Furtado, GC, Lira, SA, and
Shakhar, G. 2013a. Luminal bacteria recruit CD103+ dendritic cells into the intestinal
epithelium to sample bacterial antigens for presentation. *Immunity* **38**: 581-595.
doi:10.1016/j.immuni.2013.01.009

Farache, J, Zigmond, E, Shakhar, G, and Jung, S. 2013b. Contributions of dendritic cells and
macrophages to intestinal homeostasis and immune defense. *Immunol Cell Biol* **91**: 232-
239. doi:10.1038/icb.2012.79

Fear, VS, Burchell, JT, Lai, SP, Wikstrom, ME, Blank, F, von Garnier, C, Turner, DJ, Sly, PD,
Holt, PG, Strickland, DS, and Stumbles, PA. 2011. Restricted aeroallergen access to airway
mucosal dendritic cells in vivo limits allergen-specific CD4+ T cell proliferation during the
induction of inhalation tolerance. *J Immunol* **187**: 4561-4570. doi:10.4049/jimmunol.1004189

Fleige, H, and Forster, R. 2017. Induction and Analysis of Bronchus-Associated Lymphoid
Tissue. *Methods Mol Biol* **1559**: 185-198. doi:10.1007/978-1-4939-6786-5_13

Gautier, EL, Shay, T, Miller, J, Greter, M, Jakubzick, C, Ivanov, S, Helft, J, Chow, A, Elpek, KG,
Gordonov, S, Mazloom, AR, Ma'ayan, A, Chua, WJ, Hansen, TH, Turley, SJ, Merad, M,
Randolph, GJ, and Immunological Genome, C. 2012. Gene-expression profiles and
transcriptional regulatory pathways that underlie the identity and diversity of mouse tissue
macrophages. *Nat Immunol* **13**: 1118-1128. doi:10.1038/ni.2419

Geissmann, F, Jung, S, and Littman, DR. 2003. Blood monocytes consist of two principal
subsets with distinct migratory properties. *Immunity* **19**: 71-82. doi:10.1016/s1074-
7613(03)00174-2

Gibbings, SL, Thomas, SM, Atif, SM, McCubbrey, AL, Desch, AN, Danhorn, T, Leach, SM,
Bratton, DL, Henson, PM, Janssen, WJ, and Jakubzick, CV. 2017. Three Unique Interstitial
Macrophages in the Murine Lung at Steady State. *Am J Respir Cell Mol Biol* **57**: 66-76.
doi:10.1165/rcmb.2016-0361OC

Guilliams, M, Lambrecht, BN, and Hammad, H. 2013. Division of labor between lung dendritic
cells and macrophages in the defense against pulmonary infections. *Mucosal Immunol* **6**:
464-473. doi:10.1038/mi.2013.14

Hadis, U, Wahl, B, Schulz, O, Hardtke-Wolenski, M, Schippers, A, Wagner, N, Muller, W, Sparwasser, T, Forster, R, and Pabst, O. 2011. Intestinal tolerance requires gut homing and expansion of FoxP3+ regulatory T cells in the lamina propria. *Immunity* **34**: 237-246. doi:10.1016/j.immuni.2011.01.016

Hadjantonakis, AK, Macmaster, S, and Nagy, A. 2002. Embryonic stem cells and mice expressing different GFP variants for multiple non-invasive reporter usage within a single animal. *BMC Biotechnol* **2**: 11.

Hammad, H, Plantinga, M, Deswarte, K, Pouliot, P, Willart, MA, Kool, M, Muskens, F, and Lambrecht, BN. 2010. Inflammatory dendritic cells--not basophils--are necessary and sufficient for induction of Th2 immunity to inhaled house dust mite allergen. *J Exp Med* **207**: 2097-2111. doi:10.1084/jem.20101563

Harfe, BD, Scherz, PJ, Nissim, S, Tian, H, McMahon, AP, and Tabin, CJ. 2004. Evidence for an expansion-based temporal Shh gradient in specifying vertebrate digit identities. *Cell* **118**: 517-528. doi:10.1016/j.cell.2004.07.024

Hoffmann, FM, Berger, JL, Lingel, I, Laumonnier, Y, Lewkowich, IP, Schmudde, I, and König, P. 2018. Distribution and Interaction of Murine Pulmonary Phagocytes in the Naive and Allergic Lung. *Front Immunol* **9**: 1046. doi:10.3389/fimmu.2018.01046

Holgate, ST. 2012. Innate and adaptive immune responses in asthma. *Nat Med* **18**: 673-683. doi:10.1038/nm.2731

Huh, JC, Strickland, DH, Jahnsen, FL, Turner, DJ, Thomas, JA, Napoli, S, Tobagus, I, Stumbles, PA, Sly, PD, and Holt, PG. 2003. Bidirectional interactions between antigen-bearing respiratory tract dendritic cells (DCs) and T cells precede the late phase reaction in experimental asthma: DC activation occurs in the airway mucosa but not in the lung parenchyma. *J Exp Med* **198**: 19-30. doi:10.1084/jem.20021328

Izumi, G, Nakano, H, Nakano, K, Whitehead, GS, Grimm, SA, Fessler, MB, Karmaus, PW, and Cook, DN. 2021. CD11b(+) lung dendritic cells at different stages of maturation induce Th17 or Th2 differentiation. *Nat Commun* **12**: 5029. doi:10.1038/s41467-021-25307-x

Jakubzick, C, Tacke, F, Ginhoux, F, Wagers, AJ, van Rooijen, N, Mack, M, Merad, M, and Randolph, GJ. 2008. Blood monocyte subsets differentially give rise to CD103+ and CD103- pulmonary dendritic cell populations. *J Immunol* **180**: 3019-3027. doi:10.4049/jimmunol.180.5.3019

Jeffery, PK. 1992. Histological features of the airways in asthma and COPD. *Respiration* **59 Suppl 1**: 13-16. doi:10.1159/000196096

Jenkins, SJ, and Hume, DA. 2014. Homeostasis in the mononuclear phagocyte system. *Trends Immunol* **35**: 358-367. doi:10.1016/j.it.2014.06.006

Jung, S, Aliberti, J, Graemmel, P, Sunshine, MJ, Kreutzberg, GW, Sher, A, and Littman, DR. 2000. Analysis of fractalkine receptor CX(3)CR1 function by targeted deletion and green fluorescent protein reporter gene insertion. *Mol Cell Biol* **20**: 4106-4114. doi:10.1128/mcb.20.11.4106-4114.2000

Kamada, N, Hisamatsu, T, Honda, H, Kobayashi, T, Chinen, H, Kitazume, MT, Takayama, T, Okamoto, S, Koganei, K, Sugita, A, Kanai, T, and Hibi, T. 2009. Human CD14+ macrophages in intestinal lamina propria exhibit potent antigen-presenting ability. *J Immunol* **183**: 1724-1731. doi:10.4049/jimmunol.0804369

Kawano, H, Kayama, H, Nakama, T, Hashimoto, T, Umemoto, E, and Takeda, K. 2016. IL-10-producing lung interstitial macrophages prevent neutrophilic asthma. *Int Immunol* **28**: 489-501. doi:10.1093/intimm/dxw012

Kohl, J, Baelder, R, Lewkowich, IP, Pandey, MK, Hawlisch, H, Wang, L, Best, J, Herman, NS, Sproles, AA, Zwirner, J, Whitsett, JA, Gerard, C, Sfyroera, G, Lambris, JD, and Wills-Karp, M. 2006. A regulatory role for the C5a anaphylatoxin in type 2 immunity in asthma. *J Clin Invest* **116**: 783-796. doi:10.1172/JCI26582

Lambrecht, BN, Hammad, H, and Fahy, JV. 2019. The Cytokines of Asthma. *Immunity* **50**: 975-
991. doi:10.1016/j.immuni.2019.03.018
Lambrecht, BN, Salomon, B, Klatzmann, D, and Pauwels, RA. 1998. Dendritic cells are required
for the development of chronic eosinophilic airway inflammation in response to inhaled
antigen in sensitized mice. *J Immunol* **160**: 4090-4097.
Lindquist, RL, Shakhar, G, Dudziak, D, Wardemann, H, Eisenreich, T, Dustin, ML, and
Nussenzweig, MC. 2004. Visualizing dendritic cell networks in vivo. *Nat Immunol* **5**: 1243-
1250. doi:10.1038/ni1139
Madisen, L, Zwingman, TA, Sunkin, SM, Oh, SW, Zariwala, HA, Gu, H, Ng, LL, Palmiter, RD,
Hawrylycz, MJ, Jones, AR, Lein, ES, and Zeng, H. 2010. A robust and high-throughput Cre
reporting and characterization system for the whole mouse brain. *Nat Neurosci* **13**: 133-140.
doi:10.1038/nn.2467
Mathias, LJ, Khong, SM, Spyroglou, L, Payne, NL, Siatskas, C, Thorburn, AN, Boyd, RL, and
Heng, TS. 2013. Alveolar macrophages are critical for the inhibition of allergic asthma by
mesenchymal stromal cells. *J Immunol* **191**: 5914-5924. doi:10.4049/jimmunol.1300667
Mazzini, E, Massimiliano, L, Penna, G, and Rescigno, M. 2014. Oral tolerance can be
established via gap junction transfer of fed antigens from CX3CR1(+) macrophages to
CD103(+) dendritic cells. *Immunity* **40**: 248-261. doi:10.1016/j.immuni.2013.12.012
McDole, JR, Wheeler, LW, McDonald, KG, Wang, B, Konjufca, V, Knoop, KA, Newberry, RD,
and Miller, MJ. 2012. Goblet cells deliver luminal antigen to CD103+ dendritic cells in the
small intestine. *Nature* **483**: 345-349. doi:10.1038/nature10863
Mempel, TR, Henrickson, SE, and Von Andrian, UH. 2004. T-cell priming by dendritic cells in
lymph nodes occurs in three distinct phases. *Nature* **427**: 154-159. doi:10.1038/nature02238
Metzger, RJ, Klein, OD, Martin, GR, and Krasnow, MA. 2008. The branching programme of
mouse lung development. *Nature* **453**: 745-750. doi:10.1038/nature07005
Murphy, DB, Rath, S, Pizzo, E, Rudensky, AY, George, A, Larson, JK, and Janeway, CA, Jr.
1992. Monoclonal antibody detection of a major self peptide. MHC class II complex. *J
Immunol* **148**: 3483-3491.
Muzumdar, MD, Tasic, B, Miyamichi, K, Li, L, and Luo, L. 2007. A global double-fluorescent Cre
reporter mouse. *Genesis* **45**: 593-605. doi:10.1002/dvg.20335
Patton, JS. 1992. Mechanisms of macromolecule absorption by the lungs. *Adv Drug Deliv Rev*
19: 3-36. doi:10.1016/0169-409X(95)00113-L
Plantinga, M, Guilliams, M, Vanheerswynghels, M, Deswarte, K, Branco-Madeira, F, Toussaint,
W, Vanhoutte, L, Neyt, K, Killeen, N, Malissen, B, Hammad, H, and Lambrecht, BN. 2013.
Conventional and monocyte-derived CD11b(+) dendritic cells initiate and maintain T helper
2 cell-mediated immunity to house dust mite allergen. *Immunity* **38**: 322-335.
doi:10.1016/j.immuni.2012.10.016
Randall, TD. 2010. Bronchus-associated lymphoid tissue (BALT) structure and function. *Adv
Immunol* **107**: 187-241. doi:10.1016/B978-0-12-381300-8.00007-1
Rathjen, FG, and Schachner, M. 1984. Immunocytological and biochemical characterization of a
new neuronal cell surface component (L1 antigen) which is involved in cell adhesion. *EMBO
J* **3**: 1-10. doi:10.1002/j.1460-2075.1984.tb01753.x
Rodero, MP, Poupel, L, Loyher, PL, Hamon, P, Licata, F, Pessel, C, Hume, DA, Combadiere, C,
and Boissonnas, A. 2015. Immune surveillance of the lung by migrating tissue monocytes.
Elife **4**: e07847. doi:10.7554/eLife.07847
Roozendaal, R, Mempel, TR, Pitcher, LA, Gonzalez, SF, Verschoor, A, Mebius, RE, von
Andrian, UH, and Carroll, MC. 2009. Conduits mediate transport of low-molecular-weight
antigen to lymph node follicles. *Immunity* **30**: 264-276. doi:10.1016/j.immuni.2008.12.014
Sabatel, C, Radermecker, C, Fievez, L, Paulissen, G, Chakarov, S, Fernandes, C, Olivier, S,
Toussaint, M, Pirottin, D, Xiao, X, Quatresooz, P, Sirard, JC, Cataldo, D, Gillet, L, Bouabe,
H, Desmet, CJ, Ginhoux, F, Marichal, T, and Bureau, F. 2017. Exposure to Bacterial CpG

DNA Protects from Airway Allergic Inflammation by Expanding Regulatory Lung Interstitial Macrophages. *Immunity* **46**: 457-473. doi:10.1016/j.immuni.2017.02.016

Sasmono, RT, Oceandy, D, Pollard, JW, Tong, W, Pavli, P, Wainwright, BJ, Ostrowski, MC, Himes, SR, and Hume, DA. 2003. A macrophage colony-stimulating factor receptor-green fluorescent protein transgene is expressed throughout the mononuclear phagocyte system of the mouse. *Blood* **101**: 1155-1163. doi:10.1182/blood-2002-02-1569

Schlitzer, A, McGovern, N, Teo, P, Zelante, T, Atarashi, K, Low, D, Ho, AW, See, P, Shin, A, Wasan, PS, Hoeffel, G, Malleret, B, Heiseke, A, Chew, S, Jardine, L, Purvis, HA, Hilkens, CM, Tam, J, Poidinger, M, Stanley, ER, et al. 2013. IRF4 transcription factor-dependent CD11b+ dendritic cells in human and mouse control mucosal IL-17 cytokine responses. *Immunity* **38**: 970-983. doi:10.1016/j.immuni.2013.04.011

Schreiber, HA, Loschko, J, Karssemeijer, RA, Escolano, A, Meredith, MM, Mucida, D, Guermonprez, P, and Nussenzweig, MC. 2013. Intestinal monocytes and macrophages are required for T cell polarization in response to Citrobacter rodentium. *J Exp Med* **210**: 2025-2039. doi:10.1084/jem.20130903

Schulz, O, Jaensson, E, Persson, EK, Liu, X, Worbs, T, Agace, WW, and Pabst, O. 2009. Intestinal CD103+, but not CX3CR1+, antigen sampling cells migrate in lymph and serve classical dendritic cell functions. *J Exp Med* **206**: 3101-3114. doi:10.1084/jem.20091925

Schyns, J, Bai, Q, Ruscitti, C, Radermecker, C, De Schepper, S, Chakarov, S, Farnir, F, Pirottin, D, Ginhoux, F, Boeckxstaens, G, Bureau, F, and Marichal, T. 2019. Non-classical tissue monocytes and two functionally distinct populations of interstitial macrophages populate the mouse lung. *Nat Commun* **10**: 3964. doi:10.1038/s41467-019-11843-0

Sen, D, Jones, SM, Oswald, EM, Pinkard, H, Corbin, K, and Krummel, MF. 2016. Tracking the Spatial and Functional Gradient of Monocyte-To-Macrophage Differentiation in Inflamed Lung. *PLoS One* **11**: e0165064. doi:10.1371/journal.pone.0165064

Shaw, MH, Kamada, N, Kim, YG, and Nunez, G. 2012. Microbiota-induced IL-1beta, but not IL-6, is critical for the development of steady-state TH17 cells in the intestine. *J Exp Med* **209**: 251-258. doi:10.1084/jem.20111703

Sichien, D, Lambrecht, BN, Guilliams, M, and Scott, CL. 2017. Development of conventional dendritic cells: from common bone marrow progenitors to multiple subsets in peripheral tissues. *Mucosal Immunol* **10**: 831-844. doi:10.1038/mi.2017.8

Snippert, HJ, van der Flier, LG, Sato, T, van Es, JH, van den Born, M, Kroon-Veenboer, C, Barker, N, Klein, AM, van Rheenen, J, Simons, BD, and Clevers, H. 2010. Intestinal crypt homeostasis results from neutral competition between symmetrically dividing Lgr5 stem cells. *Cell* **143**: 134-144. doi:10.1016/j.cell.2010.09.016

Stamatiades, EG, Tremblay, ME, Bohm, M, Crozet, L, Bisht, K, Kao, D, Coelho, C, Fan, X, Yewdell, WT, Davidson, A, Heeger, PS, Diebold, S, Nimmerjahn, F, and Geissmann, F. 2016. Immune Monitoring of Trans-endothelial Transport by Kidney-Resident Macrophages. *Cell* **166**: 991-1003. doi:10.1016/j.cell.2016.06.058

Sullivan, BM, Liang, HE, Bando, JK, Wu, D, Cheng, LE, McKerrow, JK, Allen, CD, and Locksley, RM. 2011. Genetic analysis of basophil function in vivo. *Nat Immunol* **12**: 527-535. doi:10.1038/ni.2036

Tacke, R, Moos, M, Teplow, DB, Fruh, K, Scherer, H, Bach, A, and Schachner, M. 1987. Identification of cDNA clones of the mouse neural cell adhesion molecule L1. *Neurosci Lett* **82**: 89-94. doi:10.1016/0304-3940(87)90176-5

Tan, SY, and Krasnow, MA. 2016. Developmental origin of lung macrophage diversity. *Development* **143**: 1318-1327. doi:10.1242/dev.129122

Tang, C, Inman, MD, van Rooijen, N, Yang, P, Shen, H, Matsumoto, K, and O'Byrne, PM. 2001. Th type 1-stimulating activity of lung macrophages inhibits Th2-mediated allergic airway inflammation by an IFN-gamma-dependent mechanism. *J Immunol* **166**: 1471-1481. doi:10.4049/jimmunol.166.3.1471

Tang, XZ, Jung, JB, and Allen, CDC. 2019. A case of mistaken identity: The MAR-1 antibody to
mouse FcεRIα cross-reacts with FcγRI and FcγRIV. *J Allergy Clin Immunol* **143**: 1643-
1646.e1646. doi:10.1016/j.jaci.2018.11.045

Thornton, EE, Looney, MR, Bose, O, Sen, D, Sheppard, D, Locksley, R, Huang, X, and
Krummel, MF. 2012. Spatiotemporally separated antigen uptake by alveolar dendritic cells
and airway presentation to T cells in the lung. *J Exp Med* **209**: 1183-1199.
doi:10.1084/jem.20112667

Townsley, MI. 2012. Structure and composition of pulmonary arteries, capillaries, and veins.
Compr Physiol **2**: 675-709. doi:10.1002/cphy.c100081

Truman, LA, Bentley, KL, Smith, EC, Massaro, SA, Gonzalez, DG, Haberman, AM, Hill, M,
Jones, D, Min, W, Krause, DS, and Ruddle, NH. 2012. ProxTom lymphatic vessel reporter
mice reveal Prox1 expression in the adrenal medulla, megakaryocytes, and platelets. *Am J*
Pathol **180**: 1715-1725. doi:10.1016/j.ajpath.2011.12.026

Ural, BB, Yeung, ST, Damani-Yokota, P, Devlin, JC, de Vries, M, Vera-Licona, P, Samji, T,
Sawai, CM, Jang, G, Perez, OA, Pham, Q, Maher, L, Loke, P, Dittmann, M, Reizis, B, and
Khanna, KM. 2020. Identification of a nerve-associated, lung-resident interstitial
macrophage subset with distinct localization and immunoregulatory properties. *Sci Immunol*
5: eaax8756. doi:10.1126/sciimmunol.aax8756

van Dinther, D, Veninga, H, Iborra, S, Borg, EGF, Hoogterp, L, Olesek, K, Beijer, MR, Schetters,
STT, Kalay, H, Garcia-Vallejo, JJ, Franken, KL, Cham, LB, Lang, KS, van Kooyk, Y,
Sancho, D, Crocker, PR, and den Haan, JMM. 2018. Functional CD169 on Macrophages
Mediates Interaction with Dendritic Cells for CD8(+) T Cell Cross-Priming. *Cell Rep* **22**:
1484-1495. doi:10.1016/j.celrep.2018.01.021

Veres, TZ, Voedisch, S, Spies, E, Tschernig, T, and Braun, A. 2011. Spatiotemporal and
functional behavior of airway dendritic cells visualized by two-photon microscopy. *Am J*
Pathol **179**: 603-609. doi:10.1016/j.ajpath.2011.04.039

von Garnier, C, Wikstrom, ME, Zosky, G, Turner, DJ, Sly, PD, Smith, M, Thomas, JA, Judd, SR,
Strickland, DH, Holt, PG, and Stumbles, PA. 2007. Allergic airways disease develops after
an increase in allergen capture and processing in the airway mucosa. *J Immunol* **179**: 5748-
5759. doi:10.4049/jimmunol.179.9.5748

Wikstrom, ME, and Stumbles, PA. 2007. Mouse respiratory tract dendritic cell subsets and the
immunological fate of inhaled antigens. *Immunol Cell Biol* **85**: 182-188.
doi:10.1038/sj.icb.7100039

Wu, X, Briseno, CG, Durai, V, Albring, JC, Haldar, M, Bagadia, P, Kim, KW, Randolph, GJ,
Murphy, TL, and Murphy, KM. 2016. Mafb lineage tracing to distinguish macrophages from
other immune lineages reveals dual identity of Langerhans cells. *J Exp Med* **213**: 2553-
2565. doi:10.1084/jem.20160600

Yang, Z, Jung, JB, and Allen, CDC. 2018. Study of IgE-Producing B Cells Using the Verigem
Fluorescent Reporter Mouse. *Methods Mol Biol* **1799**: 247-264. doi:10.1007/978-1-4939-
7896-0_19

Yang, Z, Sullivan, BM, and Allen, CD. 2012. Fluorescent in vivo detection reveals that IgE(+) B
cells are restrained by an intrinsic cell fate predisposition. *Immunity* **36**: 857-872.
doi:10.1016/j.immuni.2012.02.009

Zaslona, Z, Przybranowski, S, Wilke, C, van Rooijen, N, Teitz-Tennenbaum, S, Osterholzer, JJ,
Wilkinson, JE, Moore, BB, and Peters-Golden, M. 2014. Resident alveolar macrophages
suppress, whereas recruited monocytes promote, allergic lung inflammation in murine
models of asthma. *J Immunol* **193**: 4245-4253. doi:10.4049/jimmunol.1400580

1245
1246
1247

 Figure Legends

Figure 1: Myeloid cells are strategically positioned for surveillance of the airways
(A) Representative immunofluorescence staining of thin cryostat sections of the lung from wild-
type mice sensitized and challenged with OVA in a classical OVA model of allergic airway
disease. Sections were stained with an antibody to smooth muscle actin (SMA, green) to
differentiate between airways and blood vessels, and with an antibody to Siglec F (red) to
visualize tissue eosinophils (Siglec-F^{bright}) and AMs (Siglec-F^{int}). Sections were stained with
DAPI (blue) to visualize all nuclei. Airways (*) and blood vessels (BV) are labeled. (B)
Representative 3D fluorescence opacity rendering of branching airways (*) and blood vessels
(BV). The image is from a 516 µm z-stack collected by two-photon microscopy of a precision-cut
thick vibratome slice of the lung from a naïve Shh-Cre ROSA26-mTmG mouse. Bronchial
airway and alveolar epithelial cells express membrane-bound GFP (green), whereas all other
cells express tdTomato (red). Collagen (blue) was visualized by second harmonic generation
(2HG). (C, D, E) Representative two-photon microscopy images of precision-cut thick vibratome
slices of the lungs of MacGreen (CSF1R-GFP) µMT mice that were treated with OVA (C) or
HDM (D) allergic airway disease models, or were untreated (E). Maximum Intensity Projections
(MIPs) representing z-stacks of 165 µm (C), 122.5 µm (D), and 127.5 µm (E) are shown. The
bronchial airways were visualized by transmitted illumination and the approximate boundaries
are denoted by dashed lines, with the airway lumen denoted by *. Collagen (blue) was
visualized by 2HG. (F) Representative two-photon microscopy images of thick lung slices from
mice expressing CX3CR1-GFP (green) together with E-cadherin-mCFP (left, blue) or Shh-Cre
Ai14 (right, blue) to label airway epithelial cells. Collagen (magenta) was visualized by 2HG.
MIPs representing z-stacks of 210 µm (left) and 142 µm (right) are shown. (G) Quantification of
the density of CX3CR1-GFP^{bright} cells located in tissue regions near the bronchial airways (AW)
or alveoli (alv). (H) Representative two-photon microscopy images of thick lung slices from mice
expressing CD11c-YFP (green) together with E-cadherin-mCFP (left, blue) or Shh-Cre Ai14
(right, blue) to label airway epithelial cells. Collagen (magenta) was visualized by 2HG. MIPs
representing z-stacks of 231 µm (left) and 285 µm (right) are shown. (I) Quantification of the
density of CD11c-YFP$^+$ cells in tissue regions near the bronchial airways (AW) or alveoli (alv).
Data are representative of 3 experiments (A), 4 experiments (B), 3 experiments (C), 2
experiments (D), >5 experiments (E), and ≥3 experiments (F, H). Images in (B-H) are stitched
from tiled images. Each data point in (G, I) represents the quantification of cell density in one
region of interest collected from 6 tiled images (AW) and 5 tiled images (alv) from 5 different
lung lobes from 4 mice across 3 experiments. ****p<0.0001 (t-test). Scale bars, 100 µm.
1282
1283
Figure 2: Antigen administered via the airways is captured by CX3CR1-GFP^{bright} IMs
(A-E) Fluorescent OVA (OVA-FL, red) was administered by intranasal droplet to naïve mice
45min-2 h prior to lung isolation. Precision-cut, thick vibratome sections were then imaged by
two-photon microscopy. Shown are representative images of regions near the bronchial airways
from mice expressing CX3CR1-GFP (green) and E-cadherin-mCFP (blue) (A, B), CD11c-YFP
(green) and E-cadherin-mCFP (blue) (C) and CX3CR1-GFP (blue) and CD11c-YFP (green) (D).
In (B) and (D), single color and merged color images of the regions are shown. Collagen
(magenta) was visualized in (A) by 2HG. MIPs representing z-stacks of 6 µm (A, left), 21 µm (A,
right), 9 µm (B), 24 µm (C, left), 22 µm (C, right), and 24 µm (D) are shown. (E) Quantification of
OVA-FL uptake by CX3CR1-GFP$^+$ versus CD11c-YFP$^+$ cells; shown are the sum of the
fluorescence intensity (FI) per cell (left) for a representative imaging field, and geometric mean
fluorescence intensity (gMFI) per imaging field (right). (F) Fluorescent HDM extract (HDM-FL,
red) was instilled intranasally to naïve mice expressing CX3CR1-GFP (blue), CD11c-YFP
(green), and E-cadherin-mCFP (gray). MIPs representing z-stacks of 12-13 µm are shown. (G,

H) Analysis of the shapes of CX3CR1-GFP$^+$ and CD11c-YFP$^+$ cells in naïve mice by quantifying
sphericity (G) and the length of the longest cell axis (H). Data are shown for individual cells as
well as the median (G) or mean (H) of cells in each imaging field analyzed. Data are
representative of 5 experiments (A-B, D); 2 experiments (C,F); or were analyzed from 5
experiments (E, right), or 4 experiments (G-H). ** p<0.01, *** p<0.001, **** p<0.0001 (t-test).
Scale bars, 20 µm.
1304
Figure 2—figure supplement 1: AMs in CX3CR1-GFP mice appear distinct from BAMs.
Representative alveolar region in the lung slice from a mouse expressing CX3CR1-GFP (green)
that was imaged by two-photon microscopy 2 h after the intranasal administration of fluorescent
antigen (OVA, red). Single color and merged MIP images depicting an 80 µm z-stack are
shown. AMs (white arrows) are visible with a round morphology and capture large amounts of
fluorescent OVA, yet exhibit only punctate autofluorescence in the GFP channel at our imaging
settings. Data are representative of >5 experiments. Scale bar, 50 µm.
1312
Figure 2—figure supplement 2: Characterization of the relative intensity of CX3CR1-GFP
versus CD11c-YFP fluorescence.
(A) Two different contrast adjustments are depicted of the same original image, shown as
merged versus single color images. In the 'normal' settings (top row), the cells look exclusively
CX3CR1-GFP$^+$ (green) or CD11c-YFP$^+$ (red), but when the contrast is increased ('bright' setting,
second row) then dim CX3CR1-GFP expression is observed in some bright CD11c-YFP$^+$ cells,
and vice versa (white arrowheads). The MIP represents a 53 µm z-stack collected by two-
photon microscopy of a precision-cut thick vibratome section from a naïve mouse. (B) Three
images are shown depicting rare cells that have bright expression of both CX3CR1-GFP (green)
and CD11c-YFP (red) reporters, and thus appear yellow (white arrowheads). The MIPs
represent 60 µm (left) and 34 µm (middle, right) z-stacks collected by two-photon microscopy of
precision-cut thick vibratome sections. Images are from independent experiments. Scale bars,
20 µm.
1326
Figure 2—figure supplement 3: OVA-capturing BAMs do not express the CD11c-YFP
reporter.
Fluorescent OVA was administered by intranasal droplet to naïve mice 2-3 h before lung
isolation and enzymatic digestion. (A, B) Representative flow cytometric analysis (A) and
quantification (B) of the intensity of OVA uptake in CX3CR1-GFP^{bright} versus CD11c-YFP^{bright}
cells. Each data point represents one mouse (naïve). (C) Representative flow cytometric
analyses of CX3CR1-GFP (upper histogram) versus CD11c-YFP (lower histogram) expression
on OVA^{bright} (OVA^{++}) APCs. Cells were pre-gated to exclude AMs, B cells, and neutrophils. (D)
Quantification of the frequency of CX3CR1-GFP^{bright} versus CD11c-YFP^{bright} cells among OVA^{++}
cells. Note that the GFP and YFP reporters were analyzed in separate mice. Only the cells
capturing the most OVA, and the cells with bright GFP or YFP expression were gated,
corresponding to the cells visualized by microscopy. Each data point represents one mouse,
and data are representative of 3 experiments. *** p<0.001, **** p<0.0001 (t-test).
1340
1341
Figure 3: Antigen-capturing CX3CR1-GFP^{bright} cells are Bronchus-Associated Macrophages
(BAMs)
(A) Representative flow cytometric analysis of CX3CR1-GFP^{bright} OVA^{bright} cells (top) versus
CD11c-YFP^{bright} cells (bottom). Cells were pre-gated to exclude B cells, neutrophils, eosinophils,
and AMs (CD19$^-$ Ly6G$^-$ Siglec-F$^-$). (B) Quantification of the frequency of CD24$^+$ CD64$^-$
("CD24$^+$") or CD64$^+$ CD24$^{-/lo}$ ("CD64$^+$") cells among CX3CR1-GFP^{bright} OVA^{bright} cells (top) or

CD11c-YFP^{bright} cells (bottom) that were gated as in (A). (C-D) Representative two-photon
microscopy of MHC-II (blue) and CD64 (red) antibody staining in precision-cut vibratome lung
sections from CX3CR1-GFP mice, showing a region containing an airway as a 3D fluorescence
opacity rendering of a 54 μm z-stack (C) as well as single z-plane images of cells to confirm
colocalization of molecules (D); note C and D are from different imaging fields. In (C and D),
single color and merged images are shown. In (C), the bronchial airway was visualized by
transmitted illumination and the approximate boundaries are denoted by dashed lines, with the
airway lumen denoted by * in the merged image. In (C), most CX3CR1-GFP^{bright} CD64⁺ cells
are MHC-II^{hi} (solid arrowheads), yet some cells are MHC-II^{lo} (open arrowheads). (E)
Representative two-photon microscopy of CD64 antibody staining (red) in vibratome lung
sections from CD11c-YFP mice, showing a MIP of a 15 μm z-stack. Mice in all panels were
unimmunized. Each data point represents one mouse (B). Data in all panels are representative
of ≥2 experiments. Scale bars, 20 μm.
1361
Figure 3—figure supplement 1: CD11c antibody labeling, CD11c-YFP and CX3CR1-GFP
expression on CD64+ IMs.
(A) Representative flow cytometric analysis of CD11c-YFP reporter expression (upper
histogram) versus CD11c antibody surface staining (bottom histogram) in the same sample,
comparing OVA^{bright} (OVA⁺⁺) BAMs (blue) to DCs (red) gated as shown (left). (B) Representative
flow cytometric analysis comparing the fluorescence intensity of the CX3CR1-GFP reporter in
cDC2s (red), MHC-II⁺ CD64⁺ MerTK⁺ cells (blue), and monocytes (green). Data are
representative of 2 experiments (A) and 4 experiments (B).
1370
Figure 3—figure supplement 2: Phenotypic comparison of cDCs, BAMs and AMs.
Representative flow cytometric analysis of marker expression on all CD24⁺ DCs (red), CD64⁺
MHC-II⁺ BAMs (blue) and AMs (green). cDCs and BAMs were gated as CD19[−] Ly6G[−] Siglec-F[−]
CD11c⁺ MHC-II⁺ then CD24⁺ (cDCs) versus CD64⁺ (BAMs) as in (Figure 3A). AMs were gated
as CD19[−] CD11c⁺ Siglec-F⁺. (A) GFP expression on cells from the lung of a *Zbtb46-GFP*
mouse. Background signal in a GFP-negative control mouse is shown in gray. (B-E) Antibody
surface staining for CD11b (B), MerTK (C), CD169 (D), and CD206 (E) on each cell population
in a wild-type mouse. Background signals in Fluorescence Minus One (FMO) controls are
shown in gray. (F) Antibody surface staining for Siglec-F on each indicated cell population in a
wild-type mouse. Cells were gated as in Figure 4. Data are representative of ≥2 experiments.
1381
Figure 4: Comparison of transcriptional profiles of AMs, IMs, and DCs by scRNA-seq.
(A) UMAP plot depicting automated graph-based clustering of CD11c-enriched cells aggregated
from lungs of 4 naïve wild-type mice. The abundance of *Siglecf* and *Ptprc* (CD45) transcripts are
shown in middle and right plots. (B) Transcriptional profile of IMs identified as the cluster of cells
most highly expressing *Mafb* and *Cx3cr1*. Genes of interest that were previously reported to be
expressed in IMs are shown in the bottom panels. (C) Transcriptional profile of DCs identified as
the clusters of cells most highly expressing *Flt3* and *Zbtb46.* Specific transcripts for each of the
sub-clusters are shown in the bottom panels. (D) Heat map showing the top 25 most
upregulated, differentially-expressed genes from each of the AMs, IMs and DCs populations
gated according to Figure 4 – figure supplement 1.
1392
1393 **Figure 4—figure supplement 1**: Gating for AMs, IMs and DCs for scRNA-seq.
1394 Based on the expression levels of cell type specific genes, *Siglecf*, *Mafb*, *Cx3cr1*, *Flt3* and
1395 *Zbtb46*, clusters containing AMs (green), IMs (blue) and DCs (orange) were gated using
1396 polygonal selection in Loupe Browser. These gated populations were analyzed for differentially
1397 expressed genes to define the upregulated genes in Figure 4D.
1398

Figure 5: Characterization of BAM marker expression and positioning with respect to nerves.
Lungs were isolated from naïve mice expressing E-cadherin CFP, CX3CR1-GFP, and/or
CD11c-YFP (as indicated). Some mice received fluorescent OVA (OVA-FL) 2 h prior to imaging.
Precision-cut lung slices were labeled with the indicated antibodies (see Methods for details)
and then imaged by two-photon microscopy. Shown are 3D fluorescence opacity renderings of
image stacks of regions adjacent to airways. (A, B) Comparison of CD206 staining to OVA-FL
uptake (A) and MHC-II staining (B). Separate two-color images (each color with E-cadherin-
CFP) are shown for best visualization of the different fluorophores. In (B), CX3CR1-GFP^{bright}
MHC-II^{hi} cells are indicated by solid arrowheads and CX3CR1-GFP^{bright} MHC-II^{lo} cells are
indicated by open arrowheads. (C, D) Visualization of nerves with L1CAM staining together with
OVA-FL uptake (C) and CD169 staining (D). Merged color images are shown to enable clear
visualization of spatial localization of cells with respect to nerves. In (D), punctate CD169
staining on BAMs is indicated by solid arrowheads, whereas a CX3CR1-GFP^{neg} cell with
uniform, bright CD169 staining adjacent to the nerve is shown with an open arrowhead. (E)
Examples of punctate CD169 staining on BAMs, often at the interface between BAMs and DCs.
Images were derived from z-stacks of 182 μm (A), 102 μm (B), 127 μm (C), 219 μm (D), 165 μm
(E, left), 124 μm (E, middle), and 7.5 μm (E, right). Scale bars, 20 μm.
1416
Figure 6: Characteristics of molecules captured by BAMs versus cDCs
(A) Fluorescent OVA was administered by intranasal droplet to naïve mice 2 h before lung
isolation and enzymatic digestion. Representative flow cytometry (left) and quantification (right)
of the frequency of OVA$^+$ cells within each indicated cell subset. Cells were pre-gated to exclude
B cells, neutrophils, eosinophils, and AMs (CD19$^-$ Ly6G$^-$ Siglec-F$^-$). CD11c$^+$ MHC-II$^+$ cells were
resolved into CD24+ cDCs versus CD64+ BAMs as in Figure 4A, and then the remaining cells
that were not CD11c$^+$ MHC-II$^+$ cells were gated as CD11b+ to define monocytes (mono). cDC1s
were gated as CD11b$^-$ CD103$^+$ and cDC2s were gated as CD11b$^+$ CD103$^-$ subsets of cDCs.
Monocytes (mono) were gated as the remaining CD11b$^+$ cells after excluding CD11c$^+$ MHC-II$^+$
cells. Mice were either CX3CR1-GFP or CD11c-YFP mice. In CX3CR1-GFP mice, monocytes
appeared as two subsets with lo to intermediate GFP expression, consistent with classical and
non-classical monocytes, respectively (data not shown). (B, C, D) The indicated fluorescent
molecules were administered by intranasal droplet to naïve CX3CR1-GFP X CD11c-YFP mice
2-4 h before lung isolation and the preparation of precision-cut thick lung sections.
Representative images depict cells that captured the indicated fluorescent molecules in 3D
fluorescence opacity renderings of 73 μm (B), 16 μm (C), and 51 μm (D) z-stacks collected by
two-photon microscopy. DX, dextran; scale bars, 20 μm. (E) FITC-dextran of a range of sizes
was administered by intranasal droplet to naïve mice, and then 2-3 h later the lungs were
isolated and enzymatically digested. The frequencies of FITC$^+$ BAMs (left) and FITC$^+$ cDC2s
(right) were enumerated by flow cytometry as shown. To control for the possibility of FITC-
dextran uptake during lung digestion and dissociation, the lungs from congenic mice that did not
receive FITC-dextran were co-digested ("control lung") with the lungs of mice that had received
FITC-dextran ("host lung"). (F) Fluorescent beads were administered to naïve mice expressing
CX3CR1-GFP and CD11c-YFP. Representative two-photon microscopy images of an airway
branchpoint (left) and alveoli (right) are shown. Images are MIPs of 107 μm (left) and 45 μm
(right) z-stacks. Beads were primarily captured by AMs which appear dimly fluorescent in the
YFP channel (note that the image of the alveoli [right] is adjusted to show the dimmer signal in
the green channel). Collagen was visualized by 2HG. Each data point represents one mouse (A,
E). Data are representative of 2 experiments (A-D). Similar results to (F) were observed in other
experiments with smaller and larger beads. Scale bars, 20 μm.
1447
1448

Figure 7: BAMs act as APCs and interact with CD4 effector T cells in the lung
(A) Representative flow cytometric analysis of Y-Ae antibody staining to detect the presentation
of MHC-II:Eα peptide complexes. OVA conjugated to Eα peptide was administered intranasally
together with OVA-Alexa647 to fluorescently label antigen-capturing cells, and then after 17-18
h to allow time for antigen processing and presentation, the lungs were isolated, followed by
enzymatic digestion and analysis of Alexa647$^+$ BAMs (blue), cDC2s (red) and AMs (green).
Control mice received OVA without the Eα peptide (gray). (B) Representative flow cytometry
(top) and quantification (bottom) of the surface expression of the co-stimulatory molecules CD80
and CD86 on BAMs (blue) and cDC2s (red) in the lungs of naïve mice in the steady state. FMO
(gray), fluorescence minus one control. (C) Flow cytometric analysis of the proliferation of naïve
OT-II T cells that were co-cultured for 2.5 days with sorted cDC2s or BAMs in the presence
(red) or absence (black) of OVA peptide. Proliferation was assessed by dilution of CellTrace
Violet. (D) Flow cytometric analysis of IL-13 production in Th2-polarized OT-II T cells that were
cultured overnight with sorted cDC2s or BAMs, or no APCs as a control, in the presence (red) or
absence (black) of OVA peptide, together with Brefeldin A. (E-G) Time-lapse images (E, F) and
quantification (G) of CD4 T cell interactions with CX3CR1-GFP$^+$ BAMs. OT-II T cells (red)
expressing tdTomato (E) or CFP (F) were adoptively transferred to mice expressing CX3CR1-
GFP (green) and CD11c-YFP (yellow) that had been sensitized with OVA and low-dose LPS,
and then challenged with OVA alone intranasally 9-10 days later. Precision-cut thick vibratome
sections of the lung were prepared for imaging 4-6 h after OVA challenge, and two-photon
microscopy z-stacks of 15 μm (E) and 22 μm (F) are shown. (G) Quantification of interaction
times in which CX3CR1-GFP$^+$ BAMs were in direct contact with OT-II T cells. Each data point
represents a BAM-T cell interaction event. Due to the limitations of imaging in a specific volume
over a fixed period of time, only some cellular interactions could be tracked for their entire
duration from start to finish (complete, black dots), whereas other interactions could only be
tracked for a portion of the total interaction time (partial, gray dots) and therefore are
underestimates. Elapsed time is indicated as hh:mm:ss. Each data point represents one mouse
(B). Data are representative of 3 experiments (A, D), 2 experiments (B, C), and 5 experiments
(E-F). Data in (G) are pooled from 3 time-lapse imaging sessions from 3 separate experiments.
****p<0.0001 (Mann-Whitney test). Scale bars, 20 μm.
1479
Figure 8: BAMs remain lung resident whereas DCs migrate to the draining lymph node
(A-D) Motility measurements derived from time-lapse imaging of CX3CR1-GFP$^+$ BAMs and
CD11c-YFP$^+$ DCs by two-photon microscopy of precision-cut thick lung sections from naïve
mice. (E) Flow cytometric analysis of the frequency of CCR7$^+$ cells among BAMs and cDC2s in
the lung 4 h after the indicated concentrations of LPS were administered intranasally. (F)
Representative flow cytometry showing CCR7 expression on BAMs (blue) and cDC2s (red) in
the lung 6 h after intranasal HDM administration. (G) The frequency of CCR7$^+$ cells among
BAMs and cDC2s in the lung in mice that were naïve ("no prime") or had been primed 10 d
earlier with OVA and 100 ng LPS ("prime"), and were challenged once with intranasal OVA (i.n.
OVA) without adjuvant. Lungs were excised 6 h after OVA challenge, enzymatically digested,
and analyzed by flow cytometry. (H) Representative two-photon microscopy images of CD11c-
YFP$^+$ DCs or CX3CR1-GFP$^+$ BAMs migrating in lymphatic vessels (Prox1-tdTomato). Mice were
administered intranasal LPS 2-4 h before lung isolation, and then precision-cut thick lung slices
were imaged for several hours. Images are MIPs of 41 μm (left) and 32 μm (right) z-stacks. (I)
Representative flow cytometry gating in the lung-draining mediastinal lymph node in mice that
were untreated or had been given HDM and PE intranasally 1 d earlier. Left, gating of cDCs
(DC, red) and monocyte-derived cells (Ly6C+, blue). Cells were pregated as CD19$^-$ CD11c$^+$
MHC-II$^+$ CD11b$^+$. Right, gating of PE$^+$ cells. (J) Quantification of the frequency of PE$^+$ cells
among DCs and monocyte-derived cells (Ly6C$^+$) gated as in (I) 1 d after HDM and PE were

   administered intranasally. Each data point represents an individual cell, pooled from four time-
1500   lapse image sequences (A); the average of all cells from one time-lapse image sequence (B, C,
1501   and D); or one mouse (E, G, J). Data are from 2 experiments (A-D) or are representative of 2
1502   experiments (E, F, I, J). Similar results to the primed mice shown in (G) were observed 1 d after
1503   OVA challenge, and similar results for CD11c-YFP cells shown in (H) were observed with HDM
1504   treatment. *** p<0.001, **** p<0.0001 (t-test). Scale bars, 20 μm.
1505
Figure 8—figure supplement 1: Comparison of Ly-6C surface expression on BAMs, DCs, and
monocytes.
Representative flow cytometric analysis, comparing Ly-6C surface expression on DCs (red) and
monocytes (green) from the lung and right posterior mediastinal lymph node (LN) as well as
lung CD64$^+$ MHC-II$^+$ BAMs (blue). Cells were from naïve mice except LN monocytes, which
were analyzed 1 d after HDM was administered intranasally. Data are representative of 2
experiments.
1513

Video Legends

Video 1: Collagen-rich region at branchpoint. The video is a 3D rendering of the image depicted
in Figure 1B, showing branching airways and blood vessels (BV) in the lung of a naïve Shh-Cre
ROSA26-mTmG mouse. The image was stitched from tiled images collected by two-photon
microscopy of a precision-cut thick vibratome section. Bronchial airway and alveolar epithelial
cells express membrane-bound GFP (green), whereas all other cells express tdTomato (red).
Collagen (blue) was visualized by second harmonic generation.

Video 2: CX3CR1-GFP[bright] IMs are positioned under the bronchial airway epithelium and are
enriched at collagen-rich regions near branchpoints. The video is a 3D rendering of the image
depicted in Figure 1F (left panel), showing branching airways and blood vessels (BV) in the lung
of a naïve mouse expressing CX3CR1-GFP (green) to visualize IMs, together with E-cadherin-
mCFP (blue) to visualize epithelial cells. Collagen (magenta) was visualized by second
harmonic generation. The image was stitched from tiled images collected by two-photon
microscopy of a precision-cut thick vibratome section. The video begins by showing the lumen
of the airways with opacity rendering and then the image rotates to show CX3CR1-GFP[bright] IMs
positioned underneath the airway epithelium. Finally, a zoomed in view of CX3CR1-GFP[bright] IMs
in the collagen-rich region at the branchpoint is shown.

Video 3: CD11c-YFP[bright] DCs are positioned near the bronchial airway epithelium and are
enriched at collagen-rich regions near branchpoints. The video is a 3D rendering of the image
depicted in Figure 1H (right panel), showing branching airways in the lung of a naïve mouse
expressing CD11c-YFP (green) to visualize DCs, together with Shh-Cre and the Ai14
(tdTomato) reporter to show airway and alveolar epithelial cells (blue). Collagen (magenta) was
visualized by second harmonic generation. The image was stitched from tiled images collected
by two-photon microscopy of a precision-cut thick vibratome section.

Video 4: Antigen-capturing BAMs interact with DCs. The video is a time-lapse image sequence
showing the interactions of CX3CR1-GFP[bright] BAMs (green) with CD11c-YFP[bright] DCs (yellow).
The BAMs appear to have captured 10 kDa TRITC-dextran (red), which was administered to the
mouse by intranasal droplet approximately two hours prior to lung isolation. The image
sequence is rendered from a 32 µm z-stack collected by two-photon microscopy of a cultured
precision-cut vibratome section of the lung. Elapsed time is shown as hh:mm:ss.

Video 5: CD4 T cell interaction with a BAM (example 1). The video is a time-lapse image
sequence corresponding to Figure 6E, showing the interactions of an OVA-specific OT-II CD4 T
cell (red) with a CX3CR1-GFP[bright] BAM (green) in the proximity of CD11c-YFP[bright] DCs (yellow).
The OT-II T cell was visualized by the expression of CD4-Cre and the Ai14 reporter leading to
tdTomato expression. The image sequence is rendered from a 15 µm z-stack, collected by two-
photon microscopy of a cultured precision-cut vibratome section of the lung isolated 4 h after
intranasal OVA challenge. Elapsed time is shown as hh:mm:ss.

Video 6: CD4 T cell interaction with a BAM (example 2). The video is a time-lapse image
sequence corresponding to Figure 6F, showing the interactions of an OVA-specific OT-II CD4 T
cell (red) with a CX3CR1-GFP^{bright} BAM (green) in the proximity of CD11c-YFP^{bright} DCs (yellow).
The OT-II T cell was visualized by the expression of a β-actin-CFP transgene. The image
sequence is rendered from a 22 µm z-stack, collected by two-photon microscopy of a cultured
precision-cut vibratome section of the lung isolated 6 h after intranasal OVA challenge. Elapsed
time is shown as hh:mm:ss.

Video 7: Different motility characteristics of BAMs vs DCs. The video is a time-lapse image
sequence showing the dynamics of CX3CR1-GFP^{bright} BAMs versus CD11c-YFP^{bright} DCs in a
collagen-rich region proximal to an airway. Collagen (blue) was visualized by second harmonic
generation. The MIP represents a 27 µm z-stack collected by two-photon microscopy of a
precision-cut vibratome section of the lung of a naïve mouse. Elapsed time is shown as
hh:mm:ss.

Video 8: Some DCs enter lymphatics. The video is a time-lapse image sequence showing
CD11c-YFP^{bright} DCs (green) migrating within a Prox1-tdTomato$^+$ lymphatic vessel (red). One
DC can be seen that is entering the lymphatic vessel. Collagen (blue) was visualized by second
harmonic generation. The image sequence was rendered from a 58 µm z-stack collected by
two-photon microscopy. The mouse was administered intranasal LPS 2 h before lung isolation,
and then various precision-cut thick lung slices were imaged over a period of several hours prior
to this recording. Elapsed time is shown as hh:mm:ss.

Video 9 BAMs do not enter lymphatics. The video is a time-lapse image sequence
corresponding to Figure 7H, right panel. CX3CR1-GFP^{bright} BAMs can be observed proximal to a
Prox1-tdTomato$^+$ lymphatic vessel (red), but do not enter the lymphatic. Collagen (blue) was
visualized by second harmonic generation. The image sequence was rendered from a 32 µm z-
stack collected by two-photon microscopy. The mouse was administered intranasal LPS 3.5 h
before lung isolation and then various precision-cut lung slices were imaged over the next few
hours prior to this recording. Elapsed time is shown as hh:mm:ss.

Figure 1

Figure 2

Figure 2—figure supplement 1

Figure 2—figure supplement 2

Figure 2—figure supplement 3

A

B

C

D

Figure 3

A

CX3CR1-GFP / OVA
72% (CD11c / MHC-II)
65% (MerTK / CD64)
6.7% CD24 / 81% CD64

CD11c-YFP / MHC-II
91% (CD11c / MHC-II)
15% (MerTK / CD64)
62% CD24 / 20% CD64

B

Frequency among OVA⁺GFP⁺ cells (%) — CD24⁺, CD64⁺

Frequency among CD11c-YFP⁺ cells (%) — CD24⁺, CD64⁺

C CX3CR1-GFP MHC-II CD64 Merge

D CX3CR1-GFP MHC-II CD64 Merge

E CD11c-YFP CD64

Figure 3—figure supplement 1

A

B

Figure 3—figure supplement 2

Figure 4

A

B IM characterization

C DC characterization

D

Figure 4—figure supplement 1

AM   IM   DC

Figure 5

Figure 6

A BAMs cDC1s cDC2s mono OVA %OVA⁺ — $\%OVA^+$

BAMs cDC1s cDC2s mono

B TRITC-dextran 10 kDa **C** TRITC-dextran 155 kDa **D** PE 240 kDa

CX3CR1-GFP
CD11c-YFP
Antigen

E BAMs — cDC2s

Host lung   Control lung

Frequency of FITC⁺ cells (%)

Ctrl 10 40 150 500 2000
Dextran size (kDa)

F Airway branchpoint   Alveoli

CX3CR1-GFP
CD11c-YFP
Beads
Collagen

CX3CR1-GFP
CD11c-YFP
Beads

Figure 7

Figure 8

Figure 8—figure supplement 1