Cortical actin properties controlled by *Drosophila* Fos

aid macrophage infiltration against surrounding tissue resistance

Short title: Macrophage Cortical Actin counteracts Tissue Resistance

Vera Belyaeva[§1,2]**, Stephanie Wachner**[§1]**, Attila Gyoergy**[§1,3]**, Shamsi Emtenani**[1]**, Igor**

Gridchyn[†1,4]**, Maria Akhmanova**[†1]**, Markus Linder**[5,6]**, Marko Roblek**[1]**, Maria Sibilia**[5]**,**

Daria Siekhaus[1*]

§: These authors contributed equally

†: These authors contributed equally

*- Corresponding author, *daria.siekhaus@ist.ac.at*

1 – Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg,

Austria

2 – Current address: Molecular Devices, Urstein Süd 17, 5412, Austria

3 – Current address: Campus-Vienna-BioCenter 1, 1030 Vienna, Austria

4 – Current address: TetraScience, Talstrasse 11, 87637, Seeg, Germany

5 – Institute of Cancer Research, Department of Medicine 1, Comprehensive Cancer Center,

Medical University of Vienna, 1090 Vienna, Austria

6 – Current address: F. Hoffmann ‐ La Roche Ltd., Grenzacherstrasse 124, 4070 Basel,

Switzerland

Blurb: Like a tank using thick armor to move through rough terrain, macrophages require a
crosslinked cortical actin shell to efficiently penetrate into tissues during *Drosophila*
development
26

ABSTRACT

The infiltration of immune cells into tissues underlies the establishment of tissue resident

macrophages, and responses to infections and tumors. Yet the mechanisms immune cells

utilize to negotiate tissue barriers in living organisms are not well understood, and a role for

cortical actin has not been examined. Here we find that the tissue invasion of *Drosophila*

macrophages, also known as plasmatocytes or hemocytes, utilizes enhanced cortical F-actin

levels stimulated by the *Drosophila* member of the fos proto oncogene transcription factor

family (Dfos, Kayak). RNA sequencing analysis and live imaging show that Dfos enhances

F-actin levels around the entire macrophage surface by increasing mRNA levels of the

membrane spanning molecular scaffold tetraspanin TM4SF, and the actin cross-linking

filamin Cheerio which are themselves required for invasion. Both the filamin and the

tetraspanin enhance the cortical activity of Rho1 and the formin Diaphanous and thus the

assembly of cortical actin, which is a critical function since expressing a dominant active

form of Diaphanous can rescue the *Dfos* macrophage invasion defect. *In vivo* imaging shows

that Dfos enhances the efficiency of the initial phases of macrophage tissue entry. Genetic

evidence argues that this Dfos-induced program in macrophages counteracts the constraint

produced by the tension of surrounding tissues and buffers the properties of the macrophage

nucleus from affecting tissue entry. We thus identify strengthening the cortical actin

cytoskeleton through Dfos as a key process allowing efficient forward movement of an

immune cell into surrounding tissues.

Introduction

The classical model of cell migration on a surface postulated in the 1980's by

Abercrombie has been extended (Danuser et al., 2013) by studies showing that migrating

cells utilize diverse strategies depending on the architecture and physical properties of their

three dimensional (3D) surroundings (Paluch et al., 2016). Much of this work has been

conducted *in vitro*, where variations in the environment can be strictly controlled. However

most 3D migration occurs within the body, and much less research has elucidated the

mechanisms used to efficiently move in these diverse environments, particularly into and

through tissues. Such migration is crucial for the influence of the immune system on health

and disease. Vertebrate macrophages migrate into tissues during development where they

take up residence, regulating organ formation and homeostasis and organizing tissue repair

upon injury (Ginhoux and Guilliams, 2016; Theret et al 2019). A variety of types of immune

cells infiltrate into tumors, and can both promote or impede cancer progression (Greten and

Grivennikov 2019; Sharma and Allison, 2015). Responses to infection require immune cells

to traverse through the vascular wall, into the lymph node, and through tissues (Luster et al.,

2005). Yet the mechanisms utilized by immune cells to allow migration into such

challenging cellular environments *in vivo* are not well understood.

Migration in 2D and 3D environments requires actin polymerization to power

forward progress. The assembly of actin at the leading edge, when coupled to Integrin

adhesion to anchor points in the surrounding ECM, can allow the front of the cell to progress

(Mitchison and Cramer, 1996). This anchoring also allows the contraction of cortical actin

at the rear plasma membrane to bring the body of the cell forwards. But a role for

crosslinked actin at the cell surface in assisting forward progress by helping to counteract the

resistance of surrounding tissues and in buffering the nucleus has not been previously

identified.

Our lab examines *Drosophila* macrophage migration into the embryonic germband

(gb) to investigate mechanisms of immune cell tissue invasion. Macrophages, also called

plasmatocytes or hemocytes, are the primary phagocytic cell in *Drosophila* and share

striking similarities with vertebrate macrophages (Brückner et al., 2004; Evans & Wood,

2011; Lemaitre & Hoffmann, 2007; Ratheesh et al., 2015; Weavers et al., 2016). They are

specified in the head mesoderm at embryonic stages 4-6 and by stage 10 start spreading

along predetermined routes guided by platelet-derived growth factor- and vascular

endothelial growth factor-related factors (Pvf) 2 and 3 (Cho et al., 2002; Brückner et al.,

2004; Wood et al., 2006) to populate the whole embryo. One of these paths, the movement

into the gb, requires macrophages to invade confined between the ectoderm and mesoderm

(Ratheesh et al., 2018; Siekhaus et al., 2010). The level of tension and thus apparent

stiffness of the flanking ectoderm is a key parameter defining the efficiency of macrophage

passage into and within the gb (Ratheesh et al., 2018). Penetration of macrophages into the

gb utilizes Integrin, occurs normally without MMPs (Siekhaus et al., 2010) and is even

enhanced by ECM deposition (Valoskova et al., 2019; Sánchez-Sánchez et al., 2017) likely

because the basement membrane has not yet formed at this stage (Matsubayashi et al., 2017;

Ratheesh et al., 2018). Thus *Drosophila* macrophage gb invasion represents an ideal system

to explore the mechanisms by which immune cells and surrounding tissues interact with one

another to aid the invasion process.

Here we sought to identify a transcription factor that could control immune cell tissue

invasion and elucidate its downstream mechanisms. We identify a role for the *Drosophila*

ortholog of the proto-oncogene Fos in initial entry and migration within the tissue. We find

Dfos increases cortical macrophage F-actin levels through the formin Cheerio and a novel

target, the tetraspanin TM4SF, aiding macrophages to move forward against the resistance of

the surrounding tissues while buffering the nucleus.

Results

The transcription factor Dfos is required for macrophage germband invasion

To identify regulators of programs for invasion we searched the literature for

transcription factors expressed in macrophages prior to or during their invasion of germband

tissues (gb) (Fig 1A-B'). Of the 12 such factors (S1 Table, based on Hammonds et al., 2013)

we focused on Dfos, a member of the Fos proto-oncogene family, assigned by the Roundup

algorithm as being closest to vertebrate c-fos (Deluca et al., 2012; Thurmond et al., 2019)

(Fig 1C). Dfos contains the basic leucine zipper domain (bZIP) shown to mediate DNA

binding and hetero and homo dimerization (Glover and Harrison, 1995; Szalóki et al., 2015)

with the third leucine replaced by a methionine, a position also altered in the *C. elegans*

ortholog FOS-1A (Sherwood et al., 2005). Embryo *in situ* hybridizations reveal enriched

expression of the gene in macrophages at early stage 11 (Fig 1D, arrow) which is attenuated

by stage 13 matching what was seen in the BDGP *in situ* database

https://insitu.fruitfly.org/cgi-bin/ex/report.pl?ftype=1&ftext=FBgn0001297. Antibody

staining against Dfos protein appears in the nucleus in macrophages that are migrating

towards the gb at stage 10-12 (Fig 1E-F' yellow arrowheads, G-G''' white arrows) and is

still observed in stage 13 (S1A Fig). The *Dfos[1]* null mutant that removes exon 1 including

the translational start site (Riesgo-Escover and Hafen, 1997, Zeitlinger, et al., 1997)

eliminates the signal in macrophages, indicating antibody specificity (Fig 1H). To determine

if Dfos affects invasion, we examined the 70% of embryos that did not display

developmental defects at these early stages from *Dfos[1]* and the hypomorphic *Dfos[2]*

(Zeitlinger et al., 1997); we quantified macrophage numbers in the gb during a defined

developmental period in early stage 12 (Fig 1M). Both Dfos mutants displayed significantly

reduced numbers of macrophages in the gb compared to the control (Fig 1I-K, N) with

normal numbers in the pre-gb zone for *Dfos[2]* (S1B Fig) (S1 Data). Macrophage-specific

expression of *Dfos* rescues the *Dfos[2]* mutant (Fig 1L,N). Blocking Dfos function in

macrophages with a dominant negative (DN) Dfos (Fig 1O-Q) that lacks the activation

domain but retains the capacity to dimerize and bind DNA (Eresh et al., 1997) or two

different RNAis against *Dfos* (Fig 1R) recapitulates the decrease in gb macrophages seen in

the null while not affecting macrophage numbers in the whole embryo (S1C Fig), and along

the ventral nerve cord (vnc) (S1D-E Fig). However, macrophages expressing DfosDN or

the *Dfos* RNAis accumulate in the pre-germband area (S1F-G Fig), as if they are

accumulating there when unable to progress further. These results argue that Dfos is

required in macrophages for their migration into the gb. The tool we chose to examine this

capability was DfosDN for the following reasons. Dfos and DfosDN do not appear to inhibit

other bZIP proteins at higher levels of expression: overexpressing DfosDN in the midgut

does not inhibit another bZIP protein that acts there (Eresh et al., 1997) and overexpressing

Dfos in macrophages does not change gb numbers (S1H Fig). DfosDN should exert a

quicker effect than the RNAis. And finally, the *Dfos RNAi*s no longer exert an effect when a

second UAS construct is simultaneously expressed (S1I Fig). Thus our further experiments

examining Dfos' role in enhancing macrophage germband invasion utilized mostly the DN

form.

Fig 1. The bZIP transcription factor Dfos acts in macrophages to facilitate their
migration into the germband

Schematics of lateral **(A)** stage (St) 11 and **(A')** early St 12 embryos. The boxed region

magnified below indicates where macrophages (green) invade the germband (gb) after

moving there from the head **(B-B')**. Macrophages sit on the yolk sac (yellow) next to the

amnioserosa (black line) and then invade between the ectoderm (blue) and mesoderm

(purple).

(C) Dfos protein aligned with its human orthologs c-Fos and FosB; orange outlines the

bZIP region that has 48% identity to both proteins: identical amino acids shown in orange,

conserved ones in green. Stars indicate Leucines in the zipper; ^ the third leucine which in

Dfos is a methionine, a tolerated substitution (Garcia-Echeverria, 1997). The lower solid

line indicates the basic domain and the dotted line the leucine zipper (ZIP).

(D) *In situ* hybridization of St 11 and 13 embryos with a riboprobe for Dfos-RB

(Fbcl0282531) which also detects all Dfos isoforms. *Dfos* RNA expression is enriched in

macrophages (arrow) and the amnioserosa (arrowhead) before gb invasion, but is gone

thereafter.

(E-H') Confocal images of the boxed region in A from fixed embryos expressing *GFP* in

macrophages (green) stained with a Dfos Ab (red). **(E-F', H-H')** A white dashed line

indicates the gb edge. **(E-F)** The Dfos Ab (yellow arrowheads) stains **(E)** macrophages

moving towards the gb at St 11, and **(F)** early St 12, as well as the amnioserosa (white

arrowheads). **(G)** Higher magnification shows Dfos colocalizing with the nuclear marker

DAPI (white). **(H)** No staining is detected in macrophages or the amnioserosa in the null

Dfos^1 mutant.

(I-L) Lateral views of mid St 12 embryos from **(I)** the control, **(J)** the null allele *Dfos^1*, **(K)**

the hypomorphic allele *Dfos^2*, and **(L)** *Dfos^2* with *Dfos* re-expressed in macrophages.

(M) Schematic of St 12 embryo, gb region indicated by a black oval outline. **(N)**

Quantitation reveals that both *Dfos* alleles display fewer macrophages in the gb. Re-

expression of *Dfos* in macrophages in the *Dfos^2* hypomorph significantly rescues the defect.

Control vs. *Dfos^1* p=0.02 (30% reduction), Control vs. *Dfos^2* p=0.017 (25% reduction),

Control vs. *Dfos^2; mac>Dfos* p=0.334.

(O-P) Lateral views of mid St 12 embryos from **(O)** the control, or **(P)** a line expressing a

dominant negative (DN) form of Dfos in macrophages. **(Q)** Quantification of macrophage

numbers in the gb (see schematic) in the two genotypes visualized in **O-P**. p=0.0002 (***)

(40% reduction). Standard Deviation (SD): 25, 25.

(R) Quantification of macrophage numbers in the gb of the control and two different lines

expressing RNAi constructs against Dfos in macrophages. Quantification of macrophage

numbers in the gb for lines expressing one of two different *UAS-Dfos RNAi* constructs in

macrophages. Control vs. *mac>Dfos RNAi^1* (TRiP HMS00254) or vs. *mac>Dfos RNAi^2*

(TRiP JF02804), p<0.0001 (54 or 52% reduction). SD: 32, 19, 29. The data in **Q** and **R** argue

that Dfos is required within macrophages to promote gb tissue invasion.

Embryos are positioned with anterior to left and dorsal up in all images and histograms show

mean + standard error of the mean (SEM) throughout. Macrophages are labeled using *srp-*

Gal4 ("mac>") driving *UAS-GFP* in **E-H**, *UAS-GFP::nls* in **I-L** and *srpHemo-*

H2A::3xmCherry in **O-R**. ***p<0.005, **p<0.01, *p<0.05. One-way ANOVA with Tukey

post hoc was used for **N** and **R**, and unpaired t-test for **Q**. The embryo number analysed is

indicated within the relevant column in the graphs. Scale bar: 50 µm in **D**, 5 µm in **E-H** and

10 µm in **I-L, O-P**.

Dfos promotes macrophage motility and persistence during tissue entry

To examine the dynamic effects of Dfos on tissue invasion, we performed live imaging and

tracking of macrophages. We visualized macrophages with *srpHemo-H2A::3xmCherry*

(Gyoergy et al., 2018) in either a wild type or *mac>DfosDN* background, capturing the

initial stage of invasion (S1 Movie). The speed of macrophages moving in the area

neighboring the germband prior to invasion was not significantly changed (pre-gb, Fig

2B,C). However, the first *mac>DfosDN* macrophage to enter is delayed by 20 min in

crossing into the gb (Fig 2D). *mac>DfosDN* macrophages also displayed reduced speed and

directional persistence during entering as well as while moving along the first 20µm of the

ectoderm-mesoderm interface (gb entry, Fig 2E, S2A Fig). Macrophages in the *Dfos²* mutant

largely mirrored this phenotype, but displayed slower movement in the pre-gb zone

neighboring the amnioserosa in which Dfos is also expressed (Fig 1D-F), likely causing a

non-autonomous effect (S2B-C Fig, S2 Movie) (Fig 1D, black arrowhead, E-F, white

arrowheads). Macrophages expressing DfosDN moved with unaltered average speed as they

spread out along the non-invasive route of the vnc (Fig 2F, Fig 2G, S3 Movie), albeit with

reduced directional persistence (S2A Fig). We thus conclude from live imaging that Dfos in

macrophages aids their initial invasive migration into the gb, increases their speed within the

gb and does not underlie their progress along the vnc.

Fig 2. Dfos facilitates the initial invasion of macrophages into the gb tissue

(**A**) Movie stills of control embryos and those expressing DfosDN in macrophages (green,

labelled using *srpHemo-H2A::3xmCherry*). Area imaged corresponds to the black dashed

square in the schematic above. The germband (gb) border is outlined with a white dashed

line. The first entering macrophage is indicated with a white arrowhead, and time in minutes

in the upper right corner.

(**B**) Detailed schematic showing the different zones for which the parameters of macrophage

gb invasion were quantified. The pre-gb area is shown in yellow, the gb entry zone is outlined

in a solid line.

(**C**) Macrophage speed in the pre-gb area was not significantly changed in macrophages

expressing DfosDN (3.00 µm/min) compared to the control (3.61 µm/min), p= 0.58.

(**D**) Quantification shows a 68% increase in the total gb crossing time of DfosDN expressing

macrophages compared to the control. Total gb crossing time runs from when macrophages

have migrated onto the outer edge of the gb ectoderm, aligning in a half arch, until the first

macrophage has translocated its nucleus into the gb ecto-meso interface. p=0.008. SD: 4, 14.

(**E**) DfosDN expressing macrophages displayed a significantly reduced speed (1.53 µm/min)

at the gb entry zone compared to the control (1.98 µm/min), p= $1.11e^{-06}$. SD: 2, 2.

(**F**) Macrophages expressing DfosDN in a Stage 13 embryo move with unaltered speed along

the vnc in the region outlined by the dashed black box in the schematic above (4.93

µm/min), compared to the control (4.55 µm/min), p= 0.64. Corresponding stills shown in (**G**)

Macrophages are labeled by *srpHemo-Gal4* driving *UAS-GFP::nls*.

***p<0.005, **p<0.01, *p<0.05. Unpaired t-test used for **C-F**, a Kolmogorov-Smirnov test

for **D.** For each genotype, the number of tracks analysed in **C** and **F,** and the number of

macrophages in **D-E** are indicated within the graph columns. Tracks were obtained from

movies of 7 control and 7 *mac>DfosDN* expressing embryos in panel **D**, 3 each in **C, F**, and

4 each in **E**. Scale bar: 10 µm.

Dfos modulates Filamin and Tetraspanin to aid gb tissue invasion

To identify Dfos targets that promote macrophage invasion, we FACS isolated

macrophages from wild type and *mac>DfosDN* embryos during the time when invasion has

just begun, and conducted RNA-sequencing of the corresponding transcriptomes (Fig 3A, S1

Data). We first assessed reads that map to Dfos, which can correspond to both endogenous

and DfosDN mRNA; we found a 1.6 fold increase in the presence of the one copy of DfosDN

in this line, arguing that this transgene is expressed at levels similar to each endogenous copy

of Dfos and is unlikely to produce extraneous effects (S2 Data). We then examined genes that

displayed a $\log_2$ fold change of at least 1.5 with an adjusted P value less than 0.05 in the

presence of DfosDN. Ten genes were down-regulated (Fig 3B, S3A-B Fig) and 9 up-

regulated by DfosDN (S2 Table). Upregulated genes in DfosDN encoded mostly stress

response proteins, consistent with the role previously demonstrated for fos in *C. elegans* in

suppressing stress responses (Hattori et al., 2013). We concentrated on the downregulated

class. Of these, we focused on the actin crosslinking filamin Cheerio (Cher) and the

tetraspanin TM4SF from a group that can form membrane microdomains that affect

signalling and migration (Razinia et al., 2012; Yeung et al., 2018). No known role for TM4SF

had been previously identified in *Drosophila*. To determine if these Dfos targets were

themselves required for invasion, we knocked down Cher and TM4SF through RNAi

individually or simultaneously and observed significantly reduced macrophage numbers in

the gb, particularly upon the knockdown of both targets simultaneously (Fig 3C-G) while not

affecting macrophage numbers in the pre-gb zone (S3D Fig) or on the vnc (S3E Fig). Over-

expression of Cher or TM4SF along with *DfosDN* in macrophages increased the mean

macrophage numbers in the gb, and over-expression of TM4SF rescued the *DfosDN*

macrophage invasion defect (Fig 3H-L). Expression of a GFP control did not restore

macrophage invasion indicating that the rescue we observed through Cher or TM4SF

expression was not due to promoter competition leading to reductions in DfosDN expression.

We conclude that Dfos aids macrophage gb invasion by increasing the mRNA levels of the

filamin actin crosslinker Cher and the tetraspanin TM4SF.

Fig 3. Dfos regulates macrophage germband invasion through cytoskeletal regulators:
the Filamin Cheerio and the tetraspanin TM4SF

(A) Schematic representing the pipeline for analyzing mRNA levels in FACS sorted
macrophages.

(B) Table of genes down-regulated in macrophages expressing DfosDN. Genes are ordered
according to the normalized p-value from the RNA-Sequencing. The closest mouse protein
orthologs were found using UniProt BLAST; the hit with the top score is shown in the table.

(C-F) Lateral views of representative St 12 embryos in which the two targets with links to
actin organization, (**D**) the Tetraspanin TM4SF and (**E**) the Filamin Cheerio, have been
knocked down individually or (**F**) together, along with the control (**C**). Scale bar: 50 μm.

(G) Quantification shows that the number of macrophages in the germband is reduced in
embryos expressing RNAi against either *cher (KK 107451)* or *TM4SF (KK 102206)* in
macrophages, and even more strongly affected in the double RNAi of both. Control vs. *cher*
RNAi p=0.0005 (46% reduction). Control vs. *TM4SF RNAi* p=0.009 (37% reduction), Control
vs. *cher/TM4SF RNAi* p>0.0001 (61% reduction). *cher RNAi* vs. *TM4SF RNAi* p=0.15. SD:
29, 23, 17, 12.

(H-K) Lateral views of a representative St 12 embryo from (**H**) the control, as well as
embryos expressing DfosDN in macrophages along with either (**I**) GFP, (**J**) Cher, or (**K**)
TM4SF.

(**L**) Quantification shows that over-expression of TM4SF in DfosDN expressing
macrophages restores their normal numbers in the gb. Over-expression of Cher in this
background shows a strong trend towards rescue, but did not reach statistical significance.
Control vs. *DfosDN* p=0.015 (28% reduction); Control vs. *cher* p=0.74; Control vs. *TM4SF*
p>0.99; *DfosDN* vs. *DfosDN cher* p=0.14; DfosDN vs. *DfosDN, TM4SF* p<0.0001; Control
vs. *cher* p=0.97; Control vs. *TM4SF* p=0.35. SD: 22, 16, 16, 21, 22, 13.

(M-O) q-PCR analysis of mRNA extracted from the bones of mice that are wild type,
transgenic (tg) for *Fos* controlled by a Major Histocompatibility promoter and viral 3'UTR
elements, and those in which such c-Fos transgenesis has led to an osteosarcoma (OS).
Analysis of mRNA expression shows that higher levels of (**M**) *Fos* correlate with higher
levels of (**N-N''**) FlnA-C, and (**O**) Tspan6 in osteosarcomas. p values = 0.86, 0.001, 0.003,
SD: 0.7, 0.6, 0.3 in **M**, 0.98, 0.009, 0.007 and 0.4, 0.2, 1.5 in **N**, 0.39, < 0.0001, <0.0001 and
0.2, 0,3, 1.1 in **N'**, 0.76, 0.005, 0.002 and 0.8, 2.3, 2.4 in **N''**, 0.99, 0.004, 0.003 and 0.1, 0.2,
0.2 in **O**. Scale bar: 50 μm.

Macrophages are labeled using either **(C-F)** *srp::H2A::3xmCherry* or **(H-K)** *srpHemo-Gal4*
("*mac>*") driving *UAS-mCherry::nls*. ***p<0.005, **p<0.01, *p<0.05. Unpaired t-test or
one-way ANOVA with Tukey post hoc were used for statistics. Each column contains the
number of analyzed embryos.

In murine osteosarcoma c-fos mRNA level increases correlate with those of Filamins

and Tetraspanin-6

To determine if these Dfos targets in *Drosophila* could also be Fos targets in vertebrate

cells, we utilized a well-established murine transgenic model that over expresses c-fos. In

these mice transgenic c-fos expression from viral 3' UTR elements in osteoblasts (the bone

forming cells) leads to osteosarcoma development accompanied by a 5 fold increase in c-fos

mRNA expression (Fig 3M) (Linder et al., 2018). We examined by qPCR the mRNA levels

of our identified Dfos targets' orthologs, comparing their levels in osteosarcomas (Fos tg

OS) to neighboring, osteoblast-containing healthy bones from Fos tg mice (Fos tg bone) and

control bones from wild-type mice (wt bone). We saw 2.5 to 8 fold higher mRNA levels of

the three murine Filamin orthologs (Fig 3N-N") and a 15 fold increase in Tetraspanin-6 (Fig

3O) in osteosarcoma cells. mRNA levels of several of the orthologs of other Dfos targets we

had identified showed less strong inductions or even decreases; the Glutathione S transferase

Gstt3 and the Slit receptor Eva1c increased 4 and 2.8 fold respectively, while the

mitochondrial translocator Tspo was 25% lower (S3F-I Fig). These results suggest that

Dfos's ability to increase mRNA levels of two key functional targets for migration, a

Filamin and a Tetraspanin, is maintained by at least one vertebrate fos family member.

Dfos increases assembly of cortical actin through Cheerio and TM4SF to aid

macrophage invasion

We wished to determine what cellular properties Dfos could affect through such targets to

facilitate *Drosophila* macrophage invasion. Given Cheerio's known role as an actin

crosslinker, we examined actin in invading *mac>DfosDN* macrophages within live embryos.

To visualize actin in macrophages, we utilized a *srpHemo-moe::3xmCherry* reporter which

marks cortical F-actin (Edwards et al., 1997; Franck et al., 1993) and observed a reduction of

53% (Fig 4A-D) in its signal in invading m*ac>DfosDN* macrophages. We saw no change by

Western analysis in the levels of the Moe::3xmCherry protein itself upon DfosDN expression

(Fig S4A-A'). We hypothesized that the changes in cortical actin we observed in the

mac>DfosDN all could be due to the lower levels of Cheerio and/or TM4SF mRNA. Indeed,

we observed reductions in Moe::3xmCherry all around the edge of invading macrophages in

live embryos expressing RNAi against Cher or TM4SF in macrophages (Fig 4E-H). To test if

a decrease in actin assembly could underlie the reduced tissue invasion of *mac>DfosDN*

macrophages, we forced cortical actin polymerization by expressing a constitutively active

version of the formin Diaphanous (DiaCA) which localizes to the cortex (Gonzalez-Gaitan

and Peifer, 2009). Indeed, expressing DiaCA in macrophages completely rescued the *Dfos^1*,

Dfos^2 (Fig S4B), and *mac>DfosDN* invasion defect (Fig 4I-J). Given that Dia, like Dfos, does

not affect general macrophage migratory capacities along the ventral nerve cord (Davis et al.,

2015), we examined if Dia might normally play a role in invasion. We utilized two RNAis

against Dia and observed decreased macrophage numbers in the gb in each (Fig 4K-L) with

no effect on numbers in the pre-gb (S4C Fig) or on the vnc (S4D Fig). These results argue

that Dfos aids invasion by increasing levels of TM4SF and Cheerio to enhance assembly of

actin around the surface of the macrophage, potentially by increasing the activity of Dia.

Dfos stimulates the cortical activity of Rho1 and Diaphanous through its targets TM4SF

and Cheerio

We sought to examine if Dfos or its targets could stimulate Dia activity or that of its

known regulator, Rho, at the cortex. Dia's autoinhibition negatively regulates its cortical

localization and activity (Goode, Eck Ann Rev Biochemistry, 2007). For mDia, binding to

activated Rho GTPases as well as to other unknown membrane associated proteins can

release this autoinhibition (Seth, Otsomo, Rosen JCB 2006). We expressed a reporter for

active Rho1, the Rho1 binding domain of Dia (DiaRBD::GFP) (Abreu-Blanco et al., 2014),

in macrophages along with myristoylated Tomato to delineate the plasma membrane and

quantified intensity profiles of linescans across the membrane in various genetic backgrounds

(Fig 4M). To validate the assay we expressed Rho1DN and found, as expected, a significant

reduction, by 34%, in cortical DiaRBD compared to the control (mem/cyto=1.15 in control,

0.76 for Rho1DN) (Fig 4N). Interestingly expressing either DfosDN, or RNAis against Cher

or TM4SF also resulted in a reduction of cortical DiaRBD, by 21, 28, and 20%, respectively

(mem/cyto=0.91, 0.83, 0.92 respectively). We then tested if this increase in Rho1 activity

prompted by Dfos and its targets also resulted in a change in the activity and thus the

localization of Dia itself, utilizing Dia::GFP (Homem & Peifer, 2008) expressed in

macrophages along with Myr::Tomato to mark the membrane and quantification as above

(Fig 4O). As predicted by the prior results with mDia, upon the expression of Rho1DN we

observed a significant reduction, by 29%, in cortical Dia compared to the control

(mem/cyto=2.46 in control, 1.76 for Rho1DN) (Fig 4P). Consistent with our findings for their

effect on Rho1 activity, we observed that expressing either DfosDN, or RNAis against Cher

or TM4SF resulted in a reduction of cortical Dia, by 23, 23, and 20%, respectively

(mem/cyto=1.9, 1.88, 1.97 respectively). Our data argue that the Dfos targets TM4SF and

Cheerio increase Rho1's activity and Dia's localization at the cortex, stimulating cortical

actin assembly.

Fig 4. Dfos regulates the actin cytoskeleton through Cher, TM4SF, and the formin
Diaphanous

(A) Quantification of phalloidin intensity to detect F actin at the macrophage-macrophage contacts in
Stage 11/12 *Dfos[1]* embryos. F-actin is strongly reduced at these homotypic contacts.

(B-C) Representative confocal images of live embryos expressing in invading macrophages the F-
actin binding and homodimerizing portion of Moesin (*srpHemo-moe::3xmCherry*) to label F-actin,
presented as a maximum z-projection. Relative Moe-3xmCherry intensity is indicated with a pseudo-
color heat map as indicated on the left, with yellow as the highest levels and dark blue as the lowest as

372 indicated in the calibration bar to the left. Insets in the bottom left corner of each panel show a grey-

373 scale single z-plane corresponding to the white box in the main image. Embryo genotype indicated

374 below. Strong reductions in cortical actin are observed in macrophages expressing DfosDN compared

375 to the control.

376 **(D-E)** Quantification of the macrophage Moe:3xmCherry intensity as a measure of cortical F-actin,

377 normalized to the average fluorescence intensity of the control per batch.

378 **(D)** Quantification shows that macrophages expressing DfosDN display a 53% reduction in

379 Moe::3xmCherry intensity compared to the control when the two outliers shown as single dots are

380 excluded, 37% if they are included. Outliers identified by 10% ROUT. n of ROIs analysed = 650 for

381 control, 687 for *DfosDN*. p=0.0007 for analysis including outliers (Kolmogorov-Smirnov) and

382 p<0.0001 for analysis excluding outliers (Welch's t-test). SD: 0.2, 0.4.

383 **(E)** Quantification reveals that macrophage expression of an RNAi against either *cher* or *TM4SF*, the

384 two genes whose expression is reduced in *DfosDN*, also results in a decrease of Moe::3xmCherry

385 intensity (by 40% each). n of ROIs analysed = 549 for control, 423 for *cher RNAi,* 306 for *TM4SF*

386 *RNAi.* Control vs. *cher RNAi* p=0.006. Control vs. *TM4SF* p=0.003. SD: 0.2, 0.3, 0.2.

387 **(F-H)** Images and representation as in B-C. Strong reductions in cortical actin are observed in

388 macrophages expressing *cher RNAi* or *TM4SF RNAi* compared to the control.

389 **(I,I')** Representative confocal images of St 12 embryos from the control and a line in which

390 macrophages express DfosDN and a constitutively active (CA) form of the formin Dia to restore

391 cortical actin polymerization.

392 **(J)** Quantification shows that while macrophage expression of DiaCA does not significantly affect the

393 number of macrophages in the gb, expressing it in a DfosDN background rescues macrophage gb

394 invasion. Control vs. *DfosDN* p=0.017 (28% reduction), Control vs. *diaCA* p=0.18, Control vs.

395 *DfosDN, diaCA* p=0.010, *DfosDN* vs. *DfosDN, diaCA* p<0.0001. SD: 22, 16, 16, 24.

396 **(K,K')** Representative confocal images of St 12 embryos from the control and from a line expressing

397 an RNAi against *dia* in macrophages.

398 **(L)** Quantification of two RNAi lines against *dia* expressed in macrophages shows a 37% and 21%

399 reduction in macrophage numbers in the gb compared to control. Control vs. *dia RNAi[1]* (TRiP

400 HMS05027) p<0. 0001; control vs. *dia RNAi[2]* (TRiP HMS00308) p=0.0008. SD: 13, 20, 22.

401 **(M, O)** Examples of line profiles used for the determination of the membrane-to-cytoplasmic ratio of

402 DiaRBD in panel N and Dia in panel P. Line intensity profiles from fixed Stage 11 embryos of (M)

403 DiaRBD::GFP or (O) Dia::GFP (green) and membrane myr::Tomato (magenta) across the outward

404 facing edge of groups of macrophages sitting within ~ 40 µm of the germband that expressed either

lacZ (Control), Rho1DN, DfosDN, *cher RNAi*, or *TM4SF RNAi* as shown in the schematic in M. Line

length ~ 8 µm. Blue lines indicate mean GFP intensity on the membrane and in cytoplasm.

(**N, P**) Quantification of membrane-to-cytoplasmic intensity ratio of (N) DiaRBD::GFP or (P)

Dia::GFP expressed in macrophages under UAS control along with either lacZ (control, n=158 line

scans from 11 embryos or n=233 from 15), Rho1DN (n=123 from 7 or n=212 from 14), DfosDN

(n=135 from 8 or n=237 from 12), *cher RNAi* (n=128 from 8 or n=252 from 13), *TM4SF RNAi*

(n=205 from 11 or n=279 from 17). Control vs. Rho1DN ****$p<0.0001$ (34% or 29% reduction),

Control vs. DfosDN p=**0.0037 (21 or 23% reduction), Control vs. *cher RNAi* ***p=0.0007 (28 or

24% reduction), Control vs. *TM4SF RNAi* *p=0.026 or 0.024 (20% reduction). SD: 0.7, 0.5, 0.5, 0.5,

0.4 in N; 1.9, 0.9, 1.0, 0.9, 1.0 in P.

Macrophages are labeled using either *srpHemo-Gal4* driving *UAS-mCherry::nls* (**I-I'**), *srpHemo-*

H2A::3xmCherry (**K-K'**). *srpHemo-moe::3xmCherry*, *srpHemo-Gal4* (*mac>*) crossed to (**B**) *UAS-*

GFP as a Control, (**C**) *UAS-DfosDN*, (**F**) *w⁻* Control, (**G**) *UAS-cher RNAi* (KK 107451), (**H**) *UAS-*

TM4SF RNAi (KK 102206). *srpHemo-GAL4 UAS-Myr::tdTomato UAS-diaRBD::GFP* (M, O) *or*

UAS-dia::GFP (N, P) crossed to *UAS-lacZ* as a Ctrl, *UAS-Rho1DN* or the lines indicated above.

***$p<0.005$, **$p<0.01$, *$p<0.05$. Unpaired t-test used for **A.** Welch's t test of normalized average

mean intensity per embryo for **D** with the two indicated outliers excluded, for statistical assessment.

One way ANOVA with Tukey post hoc for **E**, **J, L.** Kruskal-Wallis for **N, P**. The number of analyzed

(**A**) macrophage-macrophage junctions, or (**D-E, J, L, N, P**) embryos is shown in each column. Scale

bar 10 µm in (**B-C, F-H**), 50 µm in (**I, K**).

We examined what consequence these lower cortical F-actin levels had on the cellular

behavior of macrophages during entry. Quantitation showed that the actin protrusion that

macrophages initially insert between the ectoderm and mesoderm during invasion was

actually longer in the *mac>DfosDN >LifeAct::GFP* macrophages than in the control (Fig 5A,

S5A Fig, S4 Movie). We then performed live imaging of macrophages labeled with

CLIP::GFP to visualize microtubules and thus cell outlines in both genotypes; we determined

the aspect ratio (maximal length over width) that the first entering cell displays as it enters

into the gb. The first DfosDN-expressing macrophage was extended even before it had fully

moved its rear into the gb (S5B Fig). We carried out measurements, taking only cells that had

entered the gb to be able to clearly distinguish the rear of the first macrophage from the tips

of following cells (Fig 5B). We also avoided including in this measurement the forward

protrusion and determined that the first macrophage inside the gb displays an average

increase of 23% in the maximal length (L) of the cell body and a 12% reduction in the

maximal width (W) (S5 Fig). Interestingly, in the pre-gb zone the aspect ratio (max L/W) of

mac>DfosDN macrophages was not different from control macrophages (Fig 5C-D) although

the *mac>DfosDN* cells were 9% smaller in both their length and width (S5D Fig). This

suggested that the gb could impose resistance on the entering macrophage, an effect which

mac>DfosDN macrophages have trouble overcoming due to their compromised cortical actin

cytoskeleton.

Fig 5. Dfos aids macrophage gb invasion against the resistance of surrounding tissues and
buffers the nucleus

(A) Quantification from live embryos shows that the length of the F-actin protrusion of the first
entering macrophage is longer in macrophages expressing DfosDN. p= 0.011. The F-actin protrusion
labelled with *srpHemo-Gal4* driving *UAS-LifeAct::GFP* was measured in the direction of forward
migration (see schematic). SD: 2.4, 3.7

(B-C) Stills from 2-photon movies of St 11 embryos showing **(B)** the first macrophages entering the
gb and **(C)** macrophages in the pre-gb zone in the control and in a line expressing DfosDN in
macrophages. Microtubules are labelled with *srpHemo-Gal4* driving *UAS-CLIP::GFP*. A blue arrow
indicates the front and a yellow arrow indicates the rear of the macrophage. Schematics above
indicate where images were acquired

(D) Schematic at left shows macrophage measurements: vertical line for the maximum length and
horizontal line for the maximum width. Histograms show the probability density distributions of the
aspect ratios (maximum length over maximum width) of the first macrophage entering the gb (left)
and macrophages in the pre-gb (right). Macrophages expressing *DfosDN* are more elongated the
mac>DfosDN line. Control vs. *DfosDN* aspect ratios at gb entry p=0.0011, in pre-gb p=0.53. SD: in
gb 1.0, 1.6; in pre-gb 0.5, 0.5.

Confocal images of St 12 embryos expressing RNAi against Lamin or LaminC in macrophages in **(E-**
E''') the control, or **(F-F''')** in embryos also expressing DfosDN in macrophages. *srpHemo-GAL4*
used as drover. *Lam RNAi[1]*: GD45636; *Lam RNAi[2]*KK107419. *Lam C RNAi:* TRiP JF01406

(G) Macrophage RNAi knockdown of Lamins which can increase nuclear deformability did not
affect macrophages numbers in the gb in the control. In embryos in which macrophages expressed
DfosDN, Lamin knockdown rescues their reduced numbers in the gb. Control vs. *DfosDN* p<0.0001.

Control vs. *Lam RNAi[1]* p>0.99, vs. *Lam RNAi[2]* p=0.83, vs. *LamC RNAi* p>0.99. Control vs. *DfosDN,*

Lam RNAi[1] p=0.024, vs. *DfosDN, Lam RNAi[2]* p>0.99, vs. *DfosDN, LamC RNAi* p>0.99. *DfosDN* vs.

DfosDN, Lam RNAi[1] p<0.0001, vs. *DfosDN, Lam RNAi[2]* p=0.0049, vs. *DfosDN, LamC RNAi*

p<0.0001. SD: 22, 10, 19, 11, 21, 23, 16, 20.

(**H**) Expressing *DfosDN* in macrophages reduces their number in the gb. Concomitantly reducing

tissue tension in the ectoderm (light blue in schematic) through Rho1DN substantially rescues

invasion. *srpHemo-QF QUAS* control (*mac<>)* governed macrophage expression and *e22C-GAL4*

ectodermal (*ecto>*). Control vs. *mac<>DfosDN* p<0.0001 (56% reduction), vs. *mac<>DfosDN;*

ecto>Rho1DN p>0.99, vs. *ecto>Rho1DN* p=0.11. *mac<>DfosDN* vs. *mac<>DfosDN; ecto>Rho1DN*

p<0.0001, vs. *ecto>Rho1DN* p=0.0044. *mac<>DfosDN; ecto>Rh1oDN* vs. *ecto>Rho1DN* p>0.99.

SD: 23, 16, 21, 18.

Macrophages are labeled in **B-C** by *srp-Gal4* driving *UAS-CLIP::GFP,* and in **E-F'''** by *srpHemo-*

Gal4 UAS-mCherry-nls. ***p<0.005, **p<0.01, *p<0.05. Unpaired t-test was used for **A,** one way

ANOVA with Tukey post hoc for **G-H**. The number shown within the column corresponds to

measurements in **A,** and analysed embryos in **G-H.** Scale bar 5μm in **B-C**, and 50μm in **E-F**'''.

Dfos promotes advancement of macrophages against the resistance of the surrounding

tissues and buffers the nucleus

We therefore examined how the properties of the gb tissues and macrophages interact

during invasion. We first investigated if the macrophage nucleus impedes normal invasion

by varying levels of the two *Drosophila* Lamin genes, Lam and LamC, both equally related

to the vertebrate lamins A and B1 (Muñoz-Alarcón et al., 2007) and both shown to affect

nuclear stiffness and deformability (Wintner et al., 2020; Zwerger et al., 2013). Over-

expressing Lam (S5E Fig) or knocking down either of these Lamins in macrophages

through RNAi (Perkins et al., 2015) did not change macrophage numbers in the gb of wild

type embryos (Fig 5E-E''', G), suggesting that the properties of the macrophage nucleus are

not a rate limiting parameter during normal tissue invasion into the narrow path between the

ectoderm and mesoderm. This result also argues that Lamins' capacity to alter gene

expression is not normally important for invasion (Andrés & González, 2009). However in

mac>DfosDN macrophages, knockdown of these Lamins was able to rescue the gb invasion

defect (Fig 5E-G), supporting the conclusion that the properties of the nucleus affect

invasion in the absence of the higher levels of cortical actin Dfos normally induces. To

directly test if reducing the tension of surrounding tissues can counteract the absence of

Dfos, we expressed Rho1DN in the ectoderm with the *e22C-GAL4* driver while expressing

QUAS-DfosDN in macrophages with the GAL4-independent Q-system driver we had

constructed, *srpHemo-QF2* (Gyoergy et al., 2018). Rho1 through ROCK is a key regulator

of Myosin activity, epithelial tension and tissue stiffness (Warner & Longmore, 2009; Zhou

et al., 2009); Myosin II is essential for actin contractility (Heer & Martin, 2017) and tension

in the *Drosophila* gb ectoderm (Ratheesh et al., 2018). Indeed, we found that this reduction

of ectodermal tension substantially rescued DfosDN expressing macrophage numbers in the

gb (Fig 5H). Taken together our results argue that Dfos aids *Drosophila* macrophages in

withstanding the resisting force of surrounding cells against the nucleus during invasion into

tissues.

Discussion:

We identify the ability to tune the state of the cortical actin cytoskeleton as a key

capacity for immune cells migrating into and within tissue barriers *in vivo*. We find that

macrophages upregulate a program governed by the transcription factor Dfos to enable this.

Dfos in *Drosophila* is known to regulate the movement during dorsal or wound closure of

epithelial sheets (Brock et al., 2012; Lesch et al., 2010; Riesgo-Escovar & Hafen, 1997;

Zeitlinger et al., 1997) as well as the development of epithelial tumors and their

dissemination (Külshammer et al., 2015; Uhlirova & Bohmann, 2006; Külshammer &

Uhlirova, 2013; Benhra et al., 2018). Here we define a different role, namely that Dfos

enables a stream of individual immune cells to efficiently push their way into tissues, a

process which is aided rather than hampered by the presence of the ECM (Sánchez-Sánchez

et al., 2017; Valoskova et al., 2019). This function appears to be specifically required for

invasion as we observe no defects in *DfosDN* macrophages' migratory speed in open

environments. *DfosDN* macrophages display decreased actin at the cell circumference and

an elongated shape within in the confinement of the germband, suggesting a defect in the

stiffness of the cortex. Strikingly, only in the presence of *DfosDN* does the state of the

nucleus become relevant, with reductions in Lamins shown to underlie nuclear stiffness

(Wintner et al., 2020) enhancing the ability of macrophages to invade. These findings along

with the ability of a softened ectoderm to substantially rescue the *DfosDN* macrophages'

germband invasion defect lead us to propose the model (Fig 6) that Dfos permits efficient

initial translocation of the macrophage body under ectodermal reactive load by forming a

stiff cortical actin shell that counteracts surrounding tissue resistance and protects the

nucleus from undergoing high levels of mechanical stress during tissue entry.

Fig. 6. Model: Dfos increases actin assembly and crosslinking through the tetraspanin TM4SF

and the Filamin Cheerio to counter surrounding tissue resistance

We propose a model for how Dfos tunes the cortical actin properties of *Drosophila* embryonic

macrophages to aid their infiltration against the resistance of the surrounding germband tissue. Dfos

leads to an increase of the tetraspanin TM4SF and the Filamin Cheerio (Cher). The binding of

TM4SF and Filamin to different partners (see Figure S6) forms a network at the cell surface of

Integrin, actin and upstream signaling molecules; this results in the recruitment of Rho GEFs and

activation of Rho1 GTPase and the formin Diaphanous, which can stimulate further actin

polymerization at the cortex. Thereby, F-actin is assembled into a more crosslinked and dense

network aiding the macrophage in moving its cell body into the ecto-meso interface. The presence of

Lamin around the nuclear membrane does not affect this process in the wildtype since the dense

cross-linked cortical actin network helps macrophages withstand the load of the surrounding tissues.

However, in the DfosDN-expressing macrophages, the loss of Cher and TM4SF lead to reduced

cross-linked actin levels at the cell cortex making the stiffness of the nucleus the rate limiting step

for macrophage infiltration of the gb tissue.

A molecular program for tissue invasion that strengthens cortical actin

Crucial mediators of this process are two actin regulators, the filamin Cher, known to

be a Dfos target in epithelia, and the previously uncharacterized membrane scaffold

tetraspanin TM4SF. We show that both require Dfos for higher mRNA levels in

macrophages and present correlative evidence that these classes of genes are also

upregulated by vertebrate c-fos. Each of these Dfos targets is required for macrophage

invasion; over-expression of TM4SF in macrophages can rescue the *DfosDN* tissue invasion

phenotype. We propose that these targets act together to strengthen the actin cytoskeleton for

tissue invasion. Higher Filamin levels cross-link actin filaments into resilient and stiffer

networks maintaining cell integrity during mechanical stress (Goldmann et al., 1997; Tseng

et al., 2004; Fujita et al., 2012). This aids the distribution of forces from focal adhesions

across the entire migrating cell body, since Filamins can bind directly to Integrin, and even

more strongly under strain (Ehrlicher et al., 2011; Glogauer et al., 1998; Kumar et al., 2019;

Razinia et al., 2012). Tetraspanins, self-associating multipass transmembrane proteins, also

can bind Integrin, forming microdomains of adhesion molecules, receptors and their

intracellular signaling complexes, including Rho GTPases (Zimmerman, et al., 2016;

Termini & Gillette, 2017; Yáñez-Mó et al., 2009; Berditchevski & Odintsova, 1999).

Filamins similarly bind receptors, regulators of actin assembly, Rho GTPases and the Rho

GEF Trio (Popowicz et al., 2006; Stossel et al., 2001; Vadlamudi et al., 2002; Ohta et al.,

1999; Bellanger et al., 2000). We observe reduced cortical levels of F-actin, active Rho1

GTPase, and the actin polymerizing formin Diaphanous in the absence of either Dfos, the

Filamin Cheerio or the Tetraspanin TM4SF. Thus our data supports the hypothesis that these

Dfos targets enhance the cortical recruitment and activation of the formin Dia to stimulate

actin polymerization at least in part through activating the Rho1 GTPase (Fig 6, S6 Fig)

(Rousso et al., 2013; Seth et al., 2006; Großhans et al., 2005; Williams et al., 2007;

Delaguillaumie et al., 2002). Dfos' upregulation of both targets could thus lead to a supra-

network in which ECM-anchored FAs connect to a strong cross-linked cortical actin lattice,

allowing Myosin contraction to be converted into cellular advancement despite resistance

from the flanking ectoderm.

We demonstrate that the actin nucleating formin Dia is important for *Drosophila*

macrophage invasion and capable of rescuing the defects in the *DfosDN* mutant. Unlike the

formin Ena which mediates chemotaxis (Davidson et al., 2019), Dia is not required for

general *Drosophila* macrophage migration, and instead allows macrophages to recoil away

from one another (Davis et al., 2015). Dia could be required for macrophages specifically

when they face resistance from their surroundings and need to increase their cortical tension.

Modeling indicates that Dia1's regulation of cortical tension requires an optimal

combination of actin cross linking and intermediate actin filament length (Chugh et al.,

2017). *Drosophila* Dia is a more processive nucleator than Ena (Bilancia et al., 2014) and

thus could create the intermediate length actin filaments that enable higher levels of

macrophage cortical tension and strain stiffening (Kasza et al., 2010) on all sides of the cell

during their invasion.

Our findings thus demonstrate that there are commonalities in the molecular

mechanisms by which *Drosophila* cells invade into either confluent tissues or the ECM.

Dfos's upregulation of the Filamin Cheerio is also required in tumor cells and aneuploid

epithelial cells to enhance ECM breaching (Külshammer & Uhlirova, 2013; Benhra et al.,

2018). Both cell types displayed enhanced levels of cortical filamentous actin, which in the

tumors is concomitant with Dia upregulation (Külshammer & Uhlirova, 2013). In the

oocyte, Filamin is required for follicle cell intercalation and border cells display higher

levels of Filamin and F-actin to maintain cellular integrity during migration between nurse

cells (Sokol & Cooley, 2003; Somogyi & Rørth, 2004). The mediator of these increased F-

actin levels, MAL-D, can be activated by Dia (Somogyi & Rørth, 2004). Thus while MMPs

may be specific to ECM crossing, a denser and more cross linked actin cortex due to

increased levels of the filamin Cheerio and activity of the formin Dia could be a common

feature of *Drosophila* cells moving through the resistance of either ECM or surrounding

tissues. Determining if such shifts in cell surface actin properties underlie some vertebrate

cancer cells' capacity to metastasize even in the presence of MMP inhibitors is an interesting

area of inquiry (Butcher et al 2009; Kessenbrock et al 2010).

Implications for vertebrate immune cell migration

Our work also suggests a new perspective on the migration of some vertebrate

immune cells. We find that altering lamin levels does not normally affect *Drosophila*

macrophage tissue invasion. This contrasts with results showing that nuclear deformability

from lower lamin levels underlies the migration of some immune cell types through narrow

constrictions engineered from rigid materials (Davidson et al., 2014; Thiam et al, 2016).

However, negotiation of such extremely challenging *in vitro* environments can lead to DNA

damage (Raab et al., 2016) and higher nuclear flexibility caused by lower lamin levels is

associated with increased cell death (Harada et al., 2014). A robust cell surface actin layer

would allow long-lived cells or those not easily replenished to protect their genome as they

move through resistant yet deformable environments. Embryonic *Drosophila* and vertebrate

tissue resident macrophages migrate into tissues during development, survive into the adult,

and serve as founders of proliferative hematopoetic niches (Holz et al., 2003; Makhijani et

al., 2011; Bosch et al., 2019; Ginhoux and Guilliams, 2016; Theret et al 2019; Guilliams et

al, 2020). Tissue resident memory T cells migrate in response to infection in mature animals,

are long-lived and not easily renewed from the blood (Szabo et al., 2019). Thus the

importance of nuclear mechanics for migration in challenging *in vivo* environments should

be explored for a broader range of immune cells as well as the utilization of cortical actin as

a strategy for genomic protection.

Materials and Methods

Fly strains and genetics

Flies were raised on standard food bought from IMBA (Vienna, Austria) containing agar,

cornmeal, and molasses with the addition of 1.5% Nipagin. Adults were placed in cages in a

fly room or a Percival DR36VL incubator maintained at 25ºC and 65% humidity or a Sanyo

MIR-153 incubator at 29ºC within the humidity controlled 25ºC fly room; embryos were

collected on standard plates prepared in house from apple juice, sugar, agar and Nipagin

supplemented with yeast from Lesaffre (Marcq, France) on the plate surface. Fly crosses and

embryo collections for RNA interference experiments (7 hour collection) as well as live

imaging (6 hour collection) were conducted at 29ºC to optimize expression under GAL4

driver control (Duffy, 2002). All fly lines utilized are listed below.

Fly stocks

srpHemo-GAL4 (mac>) was provided by K. Brückner (UCSF, USA)(Brückner et al., 2004).

Oregon R (control), P{CaryP}attP2 (control), P{CaryP}attP40 (control), kay^2 (Dfos^2),

(UAS-Fra)2 (Dfos), UAS-Rho1.N19 Rho1DN), UAS-fbz (DfosDN), UAS-kayak RNAi *(Dfos*

RNAi) TRiP HMS00254 and TRiP JF02804, *UAS-dia RNAi* TRiP HM05027, *UAS-LamC*

RNAi TRiP JF01406 and TRiP HMS00308, *e22c-GAL4 (ecto>), Resille::GFP, UAS-*

GFP.nls, UAS-mCherry.nls, UAS-CD8::GFP lines were obtained from the Bloomington

Stock Center (Indiana, USA). *kay^1 (Dfos^1)* line was provided by O. Schuldiner (WIS, Israel).

UAS-dia.deltaDad.EGFP (diaCA) and *srpHemo-GAL4 UAS-CLIP::GFP*

(mac>CLIP::GFP) lines were provided by B. Stramer (KCL, UK). *UAS-cher.FLAG (cher)*

line was provided by M. Uhlirova (CECAD, Germany). *w[1118] (control), UAS-cher* RNAi

KK107451, *UAS-TM4SF* RNAi KK102206, *UAS-Lam RNAi[1] GD45636, UAS-Lam RNAi[2]*

KK107419 lines were obtained from the Vienna *Drosophila* Resource Center (Austria).

Extended genotypes:

Here we list the lines used in each Fig; we state first the name from FlyBase; in parentheses

the name used in the Fig panels is provided.

Fig 1 and S1 Fig:

Fig 1D: *Oregon R.* Fig 1E-G: *srpHemo-GAL4, UAS-GFP (control).* S1A, F Fig: *srpHemo-*

Gal4, srpHemo-H2A::3xmCherry/P{CaryP}attP2 (control). Fig 1H: *srpHemo-GAL4, UAS-*

GFP; kay[1] (Dfos[1]). Fig 1I-L and S1B, G Fig: *srpHemo-GAL4, UAS-GFP::nls/+ (control 1).*

Fig 1H, 1J, 1N: *srpHemo-GAL4, UAS-GFP/+; kay[1] (Dfos[1]).* Fig 1K, 1N and S1B: *srpHemo-*

GAL4, UAS-GFP::nls/+; kay[2] (Dfos[2]). Fig 1L, 1N: *srpHemo-GAL4, UAS-GFP::nls/(UAS-*

Fra)2; kay[2] (Dfos[2];mac>Dfos). Fig 1O, 1Q: *10XUAS-IVS-myr::GFP/+; srpHemo-Gal4,*

srpHemo-H2A::3xmCherry/+ (control 2 and control). Fig 1P-Q: *UAS-DfosDN/+; srpHemo-*

Gal4, srpHemo-H2A::3xmCherry/+ (mac>DfosDN). S1C, F Fig: *srpHemo-Gal4, srpHemo-*

H2A::3xmCherry/ UAS GFP::nls (ctrl). *srpHemo-Gal4, srpHemo-H2A::3xmCherry/UAS-fbz*

(mac>DfosDN). S1D Fig: *srpHemo-Gal4, srpHemo-H2A::3xmCherry /+* (ctrl). *srpHemo-*

Gal4, srpHemo-H2A::3xmCherry/UAS-DfosDN (mac>DfosDN). Fig 1R and S1E, G, I Fig:

UAS-GFP; srpHemo-Gal4, srpHemo-H2A::3xmCherry (ctrl). UAS-Dfos RNAi

HMS00254/srpHemo-Gal4, srpHemo-H2A::3xmCherry (mac>DfosRNAi[1]). UAS-Dfos RNAi

JF02804/srpHemo-Gal4, srpHemo-H2A::3xmCherry (mac>DfosRNAi[2]). S1H Fig: *srpHemo-*

GAL4, UAS-GFP::nls/+ or /(UAS-Fra)2 (mac>Dfos). S1I Fig: *UAS-GFP; UAS-Dfos RNAi*

HMS00254/srp-GAL4, srpHemo-Gal4, srpHemo-H2A::3xmCherry (mac>DfosRNAi[1] + GFP).

UAS-GFP; UAS-Dfos RNAi JF02804/srpHemo-Gal4, srpHemo-H2A::3xmCherry

(mac>DfosRNAi[2] + GFP).

Fig 2 and S2 Fig:

Fig 2A, 2C-I and S2A-B, E Fig: *srpHemo-Gal4, srpHemo-H2A::3xmCherry/+ (control).* Fig

2D: *srpHemo-Gal4, srpHemo-H2A::3xmCherry/+* (3 movies) and *Resille::GFP/+; srpHemo-*

Gal4, srpHemo-H2A::3xmCherry/+ (4 movies, control) and *Resille::GFP/+; srpHemo-Gal4,*

srpHemo-H2A::3xmCherry/+ (3 movies) and *Resille::GFP/+; srpHemo-Gal4, srpHemo-*

H2A::3xmCherry/UAS-DfosDN (4 movies, *DfosDN*) Fig 2A, 2C-I and S2A-B, E Fig: *srpHemo-*

Gal4, srpHemo-H2A::3xmCherry/UAS-fbz (mac>DfosDN). S2C-D Fig: *srpHemo-GAL4, UAS-*

GFP.nls/+ (control). S2C-D Fig: *srpHemo-GAL4, UAS-GFP.nls/+; kay^2 (Dfos^2).*

Fig 3 and S3 Fig:

Fig 3C, G and S3D Fig: *UAS-Dicer2;; srpHemo-Gal4, srpHemo-H2A::3xmCherry/w^{1118}*

(control). Fig 3D, 3G and S3D Fig: *UAS-Dicer2; UAS-TM4SF RNAi* KK10220/+; *srpHemo-*

Gal4, srpHemo-H2A::3xmCherry/+ (mac>TM4SF RNAi). Fig 3E, G and S3D Fig: *UAS-*

Dicer2; UAS-cher RNAi KK107451/+; *srpHemo-Gal4, srpHemo-H2A::3xmCherry/+*

(mac>cher RNAi). Fig 3F-G: *UAS-Dicer2; UAS-cher RNAi* KK107451/*UAS-TM4SF RNAi*

KK102206; *srpHemo-Gal4, srpHemo-H2A::3xmCherry/+ (mac>TM4SF RNAi, cher RNAi).*

Fig 3H, L: *srpHemo-GAL4, UAS-mCherry::nls/UAS-mCD8::GFP (control).* Fig 3I, L:

srpHemo-GAL4, UAS-mCherry::nls/UAS-mCD8::GFP; UAS-fbz/+ (mac>DfosDN). Fig 3J,

L: *srpHemo-GAL4,UAS-mCherry::nls/UAS-cheerio::FLAG; UAS-fbz/+ (mac>DfosDN,*

cher). Fig 3K-L: *srpHemo-GAL4,UAS-mCherry.nls/UAS-TM4SF; UAS-fbz/+ (mac>DfosDN,*

TM4SF). Fig 3L: *srpHemo-GAL4, UAS-mCherry::nls/ UAS-TM4SF (mac>TM4SF).* Fig 3L:

srpHemo-GAL4, UAS-mCherry::nls/UAS-cher (mac>cher). S3A-C Fig: *srpHemo-Gal4,*

srpHemo-3xmCherry/+ (control). S3A-C Fig: *srpHemo-Gal4, srpHemo-3xmCherry/UAS-fbz*

(mac>DfosDN).

Fig 4 and S4 Fig:

Fig 4A: *srpHemo-3xmCherry; kay^1 (Dfos^1) and srpHemo-3xmCherry; +*. Fig 4B, D and S4A

Fig: *srpHemo-Gal4, srpHemo-moe::3xmCherry/+;UAS-mCD8::GFP/+*

(Control). Fig 4C-D and S4A Fig: *srpHemo-Gal4, srpHemo-moe::3xmCherry/UAS-fbz*

(mac>DfosDN). S4A Fig: *w^{118}*. Fig 4E-F: *srpHemo-Gal4, srpHemo-moe::3xmCherry/w^{118}*

(Control). Fig 4E, G: *srpHemo-Gal4, srpHemo-moe::3xmCherry/UAS-cher*

RNAi KK107451 *(mac>cher RNAi)*. Fig 4E, H: *srpHemo-Gal4, srpHemo-*

moe::3xmCherry/UAS-TM4SF RNAi KK102206 *(mac>TM4SF RNAi)*. Fig 4I-J: *srpHemo-*

GAL4, UAS-mCherry.nls/UAS-mCD8::GFP (control). Fig 4I', J: *srpHemo-GAL4, UAS-*

mCherry.nls/UAS-DiaΔDad::EGFP; UAS-fbz/+ (mac>DfosDN, diaCA). Fig 4J: *srpHemo-*

GAL4, UAS-mCherry.nls/UAS-mCD8::GFP; UAS-fbz/+ (mac>DfosDN). Fig 4J: *srpHemo-*

GAL4, UAS-mCherry.nls/ UAS-DiaΔDad::EGFP (mac>diaCA). S4B Fig: #1: *UAS-GFPnls;*

srpHemo-Gal4, srpHemo-H2A::3xmCherry. #2: *UAS-GFPnls/srpHemo-Gal4, srpHemo-*

H2A::3xmCherry; Dfos^1. #3: *UAS-GFPnls/ srpHemo-Gal4, srpHemo-H2A::3xmCherry;*

Dfos^2. #4: *UAS-DiaΔDad::EGFP/srpHemo-Gal4, srpHemo-H2A::3xmCherry; Dfos^1*. #5:

UAS-DiaΔDad::EGFP/srpHemo-Gal4, srpHemo-H2A::3xmCherry; Dfos^2.

Fig 4K-L and S4C-D Fig: *UAS-Dicer2;; srpHemo-Gal4, srpHemo-*

H2A::3xmCherry/P{CaryP}attP40 (control). Fig 4K', L and S4C-D Fig: *UAS-Dicer2;+;*

srpHemo-Gal4, srpHemo-H2A::3xmCherry/ UAS-dia RNAi HM05027 *(mac>dia RNAi^1)*. Fig

4L and S4C-D Fig: *UAS-Dicer2;+; srpHemo-Gal4, srpHemo-H2A::3xmCherry/UAS-dia*

RNAi HMS00308 *(mac>dia RNAi^2)*. Fig 4M-N and S4E Fig: *(control) UAS-*

diaRBD::GFP/+; UAS-nlacz/ srpHemo-Gal4, 10XUAS-IVS-myr::tdTomato. (mac>Rho1DN)

UAS-diaRBD::GFP/+; UAS-Rho1N.19)/srpHemo-Gal4, 10XUAS-IVS-myr::tdTomato.

(mac>DfosDN) UAS-diaRBD::GFP/+; UAS-fbz/srpHemo-Gal4, 10XUAS-IVS-

myr::tdTomato. (mac>cher RNAi) UAS-diaRBD::GFP/+; UAS-cher RNAi

KK107451/srpHemo-Gal4, 10XUAS-IVS-myr::tdTomato. (mac>TM4SF RNAi) UAS-

diaRBD::GFP/+; UAS-TM4SF RNAi KK102206/srpHemo-Gal4, 10XUAS-IVS-

myr::tdTomato. Fig 4O-P and S4F Fig: *(control) UAS-dia::GFP/+; srpHemo-Gal4, 10XUAS-*

IVS-myr::tdTomato/UAS-nlacZ. (mac>Rho1DN) UAS-dia::GFP/+; srpHemo-Gal4,

10XUAS-IVS-myr::tdTomato/UAS-Rho1N.19. (mac>DfosDN) UAS-dia::GFP/UAS-fbz;

srpHemo-Gal4, 10XUAS-IVS-myr::tdTomato/+. (mac>cher RNAi) UAS-dia::GFP/UAS-cher

RNAi KK107451; srpHemo-Gal4, 10XUAS-IVS-myr::tdTomato/+. (mac>TM4SF RNAi)

UAS-dia-GFP/UAS-TM4SF RNAi KK102206; srpHemo-Gal4, 10XUAS-IVS-

myr::tdTomato/+.

Fig 5 and S5 Fig:

Fig 5A and S5A Fig: *srpHemo-Gal4 UAS-LifeActGFP UAS-RedStinger/ srpHemo-Gal4*

UAS-LifeActGFP, UAS-RedStinger control*; srpHemo-Gal4 UAS-LifeActGFP UAS-*

RedStinger/ srpHemo-Gal4 UAS-LifeActGFP UAS-RedStinger; UAS-DfosDN/UAS-DfosDN.

Fig 5B-D and S5B-D Fig: *srpHemo-Gal4, UAS-CLIP::GFP, UAS-RedStinger (control).* Fig

5B-D and S5B-D Fig: *srpHemo-Gal4, UAS-CLIP::GFP, UAS-RedStinger; UAS-fbz*

(mac>DfosDN). Fig 5E, G: *srpHemo-GAL4, UAS-mCherry.nls/UAS-mCD8::GFP (control).*

Fig 5E'-E'', 5G: *srpHemo-GAL4, UAS-mCherry.nls/UAS-Lamin RNAi GD45636,*

KK107419 *(mac>Lam RNAi[1]* and *mac>Lam RNAi[2], respectively).* Fig 5E''', G: *srpHemo-*

GAL4, UAS-mCherry.nls/UAS-LaminC RNAi TRIP JF01406 (mac>LamC RNAi). Fig 5F-G:

srpHemo-GAL4, UAS-mCherry.nls/UAS-mCD8::GFP; UAS-fbz/+ (mac>DfosDN). Fig

5F',F'', G: *srpHemo-GAL4, UAS-mCherry.nls/UAS-Lam RNAi (Lam RNAi[1]*=GD45636, *Lam*

RNAi[2]=KK107419)*; UAS-fbz/+ (mac>DfosDN, Lam RNAi[1]* and *mac>DfosDN, Lam RNAi[2]).*

Fig 5F''', G: *srpHemo-GAL4, UAS-mCherry.nls/UAS-LaminC RNAi TRIP JF01406; UAS-*

fbz/+ (mac>DfosDN, LamC RNAi). Fig 5H: *e22CGal4,srpHemo-H2A::3xmCherry/+*

(control). Fig 5H: *srpQF/ srpHemo-H2A::3xmCherry; QUAS-fbz/UAS-Rho1.N12*

(mac<>DfosDN). Fig 5H: *e22CGal4, srpHemo-H2A::3xmCherry/srpQF; +/ UAS-Rho1.N12*

(ecto>Rho1DN). Fig 5H: srpQF/ e22C-Gal4, srpHemo-H2A::3xmCherry; UAS-

Rho1N12/QUAS-fbz (mac<>DfosDN, ecto>rho1DN). S5E Fig: *+;UAS-GFP::nls, srpHemo-*

GAL4 (control). +;UAS-GFP::Lamin, srpHemo-GAL4.

Cloning and generation of QUAS-DfosDN line

The fragment was amplified from genomic DNA of the published *UAS-fbz (UAS-Dfos DN)*

line (Eresh, Riese, Jackson, Bohmann, & Bienz, 1997) using primers encompassing a 5'

consensus translation initiation sequence followed by the bZIP fragment and containing

BglII and XhoI restriction sites: 5'-GAAGATCTATTGGGAATTCAACATGACCCCG-3'

and 5'-CCCTCGAGTCAGGTGACCACGCTCAGCAT-3'. The resulting fragment was

cloned into the pQUASt vector, a gift from Christopher Potter (Addgene plasmid # 104880).

The final construct was sequenced and injected into the attP2 landing site by BestGene

(Chino Hills, CA, USA).

Cloning and generation of UAS-TM4SF line

The TM4SF open reading frame was amplified from the DGRC GH07902 cDNA clone

(#3260, Fbcl0121651), using primers acagcgGAATTCATGGCATTGCCGAAGAAAT

and acagcgTCTAGATTAAAAGCTAATCGTCTGTCATT. The PCR product and the

pUASt-aTTB vector (DGRC plasmid #1419) were digested with EcoRI and XbaI, and

ligated. After sequencing, the construct was injected into the landing site line, (*y¹ M{vas-*

int.Dm}ZH-2A w*; *M{3xP3-RFP.attP}ZH-51D*, BL 24483), to produce second chromosome

inserts. All male survivors were crossed to *w*; *Sp/CyO*; *PrDr/TM3Ser* virgins. Transformants

were recognized by eye color and crossed again to *w*; *Sp/CyO*; *PrDr/TM3Ser* virgins to get

rid of the X chromosomal integrase.

Embryo staging:

Laterally oriented embryos with complete germband (gb) extension and the presence of

stomadeal invagination were staged based on gb retraction from the anterior as a percentage

of total embryo length. Embryos with no gb retraction were classified as Stage 11, 30%

retraction early Stage 12, 60% retraction Stage 12, and 70% Stage 13. Imaged embryos are

shown throughout paper in a lateral orientation with anterior to the left and dorsal up.

In situ **hybridization and immunofluorescence**

Embryos were dechorionated by 5 min treatment with 50% Chlorox bleach. After extensive

washing with water, embryos were fixed with 3.7% formaldehyde/heptane for 20 min

followed by methanol devitellinization for *in situ* hybridization and visualization of

3xmCherry or tdTomato. The *Dfos* cDNA clone SD04477 was obtained from the DGRC. T7

or T3 polymerase-synthesized digoxigenin-labelled anti-sense probe preparation and *in situ*

hybridization was performed using standard methods (Lehmann & Tautz, 1994). Images

were taken with a Nikon-Eclipse Wide field microscope with a 20X 0.5 NA DIC water

Immersion Objective. Embryos were mounted after immunolabeling in Vectashield

Mounting Medium (Vector Labs, Burlingame, USA) and imaged with a Zeiss Inverted

LSM700 and LSM800 Confocal Microscope using a Plain-Apochromat 20X/0.8 Air

Objective or a Plain-Apochromat 63X/1.4 Oil Objective as required.

Antibody	Source animal	Dilution	Provided by
Anti-Dfos	Rabbit	1:50	J. Zeitlinger (Stowers Institute, USA)
Anti-GFP	Chicken	1:500	Abcam (ab13970)
Anti-mCherry	Goat	1:200	Invitrogen (M11217)

Dfos antibody

The Dfos rabbit polyclonal antibody was produced for the lab of Julia Zeitlinger. It was

raised by Genescript (Piscataway, NJ, USA) against the C-terminal end of *Drosophila* Kayak

found in all isoforms and was purified against an N terminally His tagged antigen

corresponding to aa 73 to 595 of Kay isoform A. The internal Genescript order number is

163185-30, and in the Zeitlinger lab is referred to as anti-kay/fos Ab.

Western

Cages were prefed on fresh yeast plates for two days. Late stage 11/ early Stage 12 embryos

were hand picked using a Leica M205 fluorescent microscope on ice cold apple juice plates.

They were transferred to RIPA buffer (50mM Tris, 150mM NaCl, 1% NP-40, 1mM EDTA,

0.5% Na-Deoxycholate, 0.1% SDS) with a Halt Protease/Phosphatase inhibitor cocktail

(ThermoFisher, #78440) and lysed. After a 30 min incubation on ice, they were centrifuged

15 min at 4°C at 15,000 g. 10 µg of the cleared lysate were separated by SDS-PAGE using

4%-15% Mini-PROTEAN TGX Precast Protein gels (Bio-Rad, #4561085) and blotted onto a

Amersham Protran Premium Western blotting nitrocellulose membrane (Sigma,

#GE10600003). The nitrocellulose membrane was blocked with Pierce Clear Milk blocking

buffer (ThermoFisher, #37587) and incubated in blocking buffer with anti-mCherry (Novus

Biologicals, #NBP1-96752) at 1:1000, and anti-Profilin (DSHB, #chi 1J) at 1:50 antibodies

over night at 4°C. The membrane was washed three times for 10 minutes with 1x PBS and

incubated with Goat Anti-Mouse IgG (H+L)-HRP Conjugate (BioRad, #172-1011).

Chemiluminescence was induced by incubation with SuperSignal West Femto Maximum

Sensitivity Substrate (ThermoFisher, #34096) and recorded with a ChemieDoc MP (BioRad)

molecular imager. Densitometric quantification of bands was done with ImageJ.

Time-Lapse Imaging

Embryos were dechorionated in 50% bleach for 5 min, washed with water, and mounted in

halocarbon oil 27 (Sigma) on a 24x50mm high precision coverslip (Marienfeld Laboratory

Glassware, No. 1.5H) between two bridges (~0.5 cm high) of coverslips glued on top of

each other, or mounted in halocarbon oil 27 (Sigma) between a 18x18mm coverslip

(Marienfeld Laboratory Glassware, No. 1.5H) and an oxygen permeable membrane (YSI).

The embryo was imaged on an upright multiphoton microscope (TrimScope, LaVision)

equipped with a W Plan-Apochromat 40X/1.4 oil immersion objective (Olympus). GFP and

mCherry were imaged at 860 nm and 1100 nm excitation wavelengths, respectively, using a

Ti-Sapphire femtosecond laser system (Coherent Chameleon Ultra) combined with optical

parametric oscillator technology (Coherent Chameleon Compact OPO). Excitation intensity

profiles were adjusted to tissue penetration depth and Z-sectioning for imaging was set at

1µm for tracking. For long-term imaging, movies were acquired for 60 - 150 minutes with a

frame rate of 25-45 seconds. A temperature control unit set to 29°C was utilized for all

genotypes except kay^2 for which the setting was 25°C.

Image Analysis

Macrophage cell counts: Autofluorescence of the embryo revealed the position of the

germband (gb) for staging of fixed samples. Embryos with 40% (±5%) gb retraction (Stage

12) were analysed for macrophage numbers in the pre-gb, within the germband, along the

ventral nerve cord (vnc) and in the whole embryo. For the *kay RNAi*.embryos with 70% gb

retraction (Stage 13) were used for vnc counts. The pre-gb zone was defined based on

embryo and yolk autofluorescence as an area on the yolk sac underneath the amnioserosa

with borders defined posteriorly by the gb ectoderm and anteriorly by the head.

Macrophages were visualized using confocal microscopy with a Z-stack step size of 2 µm

and macrophage numbers within the gb or the segments of the vnc were calculated in

individual slices (and then aggregated) using the Cell Counter plugin in FIJI. Total

macrophage numbers were obtained using Imaris (Bitplane) by detecting all the macrophage

nuclei as spots.

Macrophage Tracking, Speed, Persistence. Mode of Migration and Macrophage gb

crossing Analysis

Embryos with macrophage nuclei labelled with *srpHemo-H2A::3XmCherry* and the

surrounding tissues with *Resille::GFP,* or with only macrophages labelled by *srpHemo-*

H2A::3XmCherry, or srpHemo>GFP.nls were imaged and 250x250x40µm^3 3D-stacks were

typically acquired with ~0.2x0.2x1µm^3 voxel size every 39-41 seconds for ~2 hours. For

imaging macrophages on vnc frames were acquired at every 40-43 seconds for 30 min after

macrophages started spreading into abdominal segment 2 (see Fig 2G). Multiphoton

microscopy images were initially processed with ImSpector software (LaVision Bio Tec) to

compile channels, and exported files were further processed using Imaris software (Bitplane)

for 3D visualization.

Each movie was rotated and aligned along the embryonic AP axis for tracking analysis. For

analysis of migration in the pre-gb and gb in the control and *kay^2* mutant, embryos were

synchronized using the onset of germ and retraction. For vnc migration analysis,

macrophages were tracked for 30 minutes from when macrophages started moving into the

second abdominal segment. Only macrophages migrating along the inner edge of the vnc

were analyzed.

Gb crossing time was calculated from when the macrophages align in front of the gb

ectoderm in a characteristic arc, until the first macrophage had transitioned its nucleus inside

the ecto-meso-interphase. To see the gb edge and yolk in movies of *srpHemo-*

3xH2A::mCherry, either *Resille::GFP* labelling the outlines of all cells, or the auto-

fluorescence of the yolk was used.

For analysis of gb migration in the *DfosDN* vs control macrophages, macrophages were

tracked from when the first macrophage appeared between the ectoderm and the yolk sac

until gb retraction started, typically 60 minutes. In the head and pre-gb, macrophage nuclei

were extracted using the spot detection function, and tracks generated in 3D over time. The

pre-gb and gb were defined as for macrophage counts described above. The mean position

of the tracks in X- and Y restrict analysis to each migratory zones.

Cell speed and persistence were calculated from nuclei positions using custom Python

scripts as described elsewhere (Smutny et al., 2017). Briefly, instantaneous velocities from

single cell trajectories were averaged to obtain a mean instantaneous velocity value over the

course of measurement. The directional persistence of a trajectory was calculated as the

mean cosine of an angle between subsequent instantaneous velocities:

$$I(v_1, ..., v_l) = \frac{1}{l-1} \sum_{k=1}^{l-1} \cos(v_k, v_{k+1}),$$

where l is duration of the trajectory and $v_1, ..., v_l$ are its instantaneous velocities. Only

trajectories with a minimal duration of 15 timeframes were used. Calculated persistence

values were averaged over all trajectories to obtain a persistence index (I) for the duration

of measurement (with -1 being the lowest and 1 the maximum). 3-6 embryos were

recorded and analyzed for each genotype, numbers of control and perturbed embryos are

equal in each pairwise comparison.

Measurement of junctional Phalloidin

The junctional intensity of F-actin (Phalloidin) was calculated using linescan analysis as

previously described (Smutny et al., 2010) with the following changes. The line was ~5 μm

and was always drawn in the middle slice of the Z stack (1 μm resolution) of the

macrophage-macrophage junction. For every line, a Gaussian fit was applied and maximum

intensities across the cell junction were then normalized against average intensities of F-

actin (Phalloidin) staining in the stereotypical gb area of ~50x50µm^2 in each embryo.

Analyses were carried out using standard Fiji software. 4-5 embryos were analysed per

genotype. Macrophages in the pre-gb or gb entry zones were analyzed.

Measurement of F-actin reporters

To quantify cortical F-actin intensity in living embryos, a *srpHemo-moe::3xmCherry*

reporter line (Gyoergy et al., 2018) was crossed into a background of macrophages

expressing *DfosDN*, *cher RNAi*, or *TM4SF* RNAi. Embryos were collected for 5h 30min at

29°C, de-chorionated in 50% bleach for 5 min, rinsed thoroughly with water, and aligned

laterally side by side under a stereomicroscope using a fluorescence lamp to check for the

presence of mCherry. Aligned embryos were then mounted as described in the live imaging

section above. To image Moe::3xmCherry, a Zeiss LSM800 inverted microscope was used

with the following settings: Plan-APOCHROMAT 40x/1.4 Oil, DIC, WD=0.13 objective,

1.5x zoom, 1025x1025 pixel, speed 8, heating chamber set to 29°C, z-interval 1µm. Laser

settings were kept constant in all experiments. Images were acquired during macrophage

invasion into the gb (St 12). Pseudo-coloring was conducted for the mCherry red channel.

Each pixel in the image has a color ascribed to it via the fire "Look Up Table" translating the

level of intensity of the mCherry channel into a defined amount of each color. The highest

intensity of the image is represented as very bright yellow and all other grey values are

depicted as colors on the scale accordingly.

For quantification of Moe::3xmCherry intensity, an ROI was drawn in Fiji software

around macrophages at the germband entry site in 20 z-stacks for each embryo. The area

mean intensity was measured in all ROIs and the average/embryo was calculated. To

normalize fluorescence intensities per batch, the average intensity/embryo of all ROIs in

each sample was divided by the arithmetic mean of the average intensity/embryo of all ROIs

in the control per batch. The normalized average intensities/embryo were then compared to

each other using a t-test with Welch's correction for *DfosDN* and one way-ANOVA for *cher*

RNAi and *TM4SF RNAi*.

Quantification of membrane localization of DiaRBD::GFP and Dia::GFP

Methanol fixed St 11 embryos were mounted either after staining with GFP antibody

(Dia::GFP) or without staining(DiaRBD::GFP) and imaged with a Zeiss Inverted LSM800,

Plain-Apochromat 63X/1.4 Oil Objective at an XY-resolution of 0.1 μm and a Z-resolution

of 1 μm (~15 μm total stack). All macrophages within 40μm of the germband were

analysed. For the quantification of the levels of DiaRBD or the complete Dia protein at the

plasma membrane versus the cytoplasm, confocal images were processed using Fiji and

MATLAB-R2017b (MathWorks). Individual focal planes were used to segment a profile

corresponding to an 8 pixel wide line drawn across the single outer membrane of individual

macrophages chosen such that the extracellular portion of the line extended into surrounding

tissue or space and not another macrophage. The corresponding intensity profiles of the

mem::Tomato and Dia::GFP or DiaRBD::GFP channels were extracted in Fiji using a

custom macro and analyzed further using a custom MATLAB script. The membrane region

was defined by finding the maximal value in the Tomato intensity profile and centering a

0.8 μm interval around it. The background was calculated for each GFP profile as the mean

intensity in the 2 μm outside the cell, flanking the membrane region, and substracted from

the entire profile. The integrated Dia::GFP or DiaRBD::GFP intensity at the membrane was

calculated within the 0.8 μm interval defined above. The integrated cytoplasmic Dia::GFP or

DiaRBD::GFP level was calculated as the mean intensity of 2 μm of the GFP profile inside

the cell flanking the membrane region. Image analysis scripts are publicly available at

https://github.com/Axmasha/Image_analysis_scripts.

Cell aspect ratio analysis and imaging actin dynamics

Laterally oriented embryos were used to measure the maximal length and width of

macrophages expressing *UAS-CLIP::GFP* under the control of *srpHemoGal4*. Briefly, 3D-

stacks with 1 µm Z resolution were acquired every 35-45 seconds for approximately 1 hour.

As the strength of the GAL4 expression increased over time, laser power was adjusted

during acquisition to reach the best possible quality of visualization. Images acquired from

mutiphoton microscopy were initially processed with ImSpector software (LaVision Bio

Tec) to compile channels from the imaging data.

We started measuring from the time the cell body of the first macrophage fully appeared at

the interface between the ectoderm and mesoderm and yolk sac until it had moved 30 µm

along the ectoderm mesoderm interface. At each timeframe, a line was drawn in Fiji along

the longest dimension of the macrophage in the direction of its front-rear polarization axis,

denoted the maximal cell length, and along the orthologonal longest dimension, which was

considered maximal cell width. We did not observe long CLIP::GFP protrusions, but when a

small protrusion was present, it was not included in the length measurement; within this gb

region the front of the first macrophage was clearly outlined with CLIP::GFP. The border

between the first and second entering macrophages was drawn based on the uninterrupted

intense line of CLIP::GFP at the base of the first macrophage; only cells with a clearly

visible border were measured. The length to width ratio was quantified for each timeframe

and a probability density function was plotted: 5 embryos were recorded for each genotype.

Imaging the actin protrusion

Laterally oriented embryos expressing *srpHemo-Gal4 UAS-LifeAct::GFP* were used to image

macrophage actin live with a 3D-stack resolution of 1µm. See above description of

CLIP::GFP labeled macrophage imaging for laser power and image compilation. Laser power

was also increased further in the DfosDN samples to enhance actin visualization. We

measured the length of the filopodia-like protrusion of the first entering macrophage with

Imaris software (Bitplane) from the time when the protrusion was inserted into the ectoderm,

mesoderm and yolk sac interface until the macrophage started to translocate its cell body into

that location.

FACS sorting of macrophages

Adult flies of either *w;+;srpHemoGal4,srpHemo::3xmCherry/+* or *w;+;*

srpHemoGal4,srpHemo::3xmCherry /UASDfosDN genotypes were placed into plastic cages

closed with apple juice plates with applied yeast to enhance egg laying. Collections were

performed at 29°C for 1 hour, then kept at 29°C for additional 5 hours 15 minutes to reach

stage 11-early stage 12. Embryos were harvested for 2 days with 6-7 collections per day and

stored meanwhile at +4°C to slow down development. Collected embryos were dissociated

and the macrophages sorted as previously described (Gyoergy et al., 2018). About $1-1.5 \times 10^5$

macrophages were sorted within 30 minutes.

Sequencing of the macrophage transcriptome

Total RNA was isolated from FACS-sorted macrophages using Qiagen RNeasy Mini kit (Cat

No. 74104). The quality and concentration of RNA was determined using Agilent 6000 Pico

kit (Cat No. 5067-1513) on an Agilent 2100 Bioanalyzer: on average about 100 ng of total

RNA was extracted from 1.5×10^5 macrophages. RNA sequencing was performed by the CSF

facility of Vienna Biocenter according to standard procedures

(https://www.vbcf.ac.at/facilities/next-generation-sequencing/) on three replicates. Briefly,

the cDNA library was synthesized using QuantSeq 3' mRNA-seq Library Prep kit and

sequenced on the Illumina HiSeq 2500 platform. The reads were mapped to the *Drosophila*

melanogaster Ensembl BDGP6 reference genome with STAR (version 2.5.1b). The read

counts for each gene were detected using HTSeq (version 0.5.4p3). Flybase annotation

(r6.19) was used in both mapping and read counting. Counts were normalised to arbitrary

units using the TMM normalization from edgeR package in R. Prior to statistical testing the

data was voom transformed and then the differential expression between the sample groups

was calculated with limma package in R. The functional analyses were done using the topGO

and gage packages in R (Anders, Pyl, & Huber, 2015; Dobin et al., 2013). RNA sequencing

data has been deposited at GEO as GSE182470.

qRT-PCR analysis of mRNA levels in murine bones and osteosarcomas

RNA isolation and qPCR was performed from bones of wild-type C57BL/6 mice and from

bones and osteosarcomas (OS) of H2-c-fosLTR as previously described with the below

primers (Rüther et al., 1989).

Primer	Sequence
Fos fw	ATGGTGAAGACCGTGTCAGG
Fos_rv	GTTGATCTGTCTCCGCTTGGA
Flna_fw	GTCACAGTGTCAATCGGAGGT
Flna_rv	TTGCCTGCTGCTTTTGTGTC
Flnb_fw	TTCTACACTGCTGCCAAGCC
Flnb_rv	CTGTAACCCAGGGCCTGAATC
Flnc_fw	CATCACCCGGAGTCCTTTCC
Flnc_rv	CTCTGTGCCCTTTGGACCTT
Tspan6_fw	TCGAACTAGTTGCCGCCATT
Tspan6_rv	CCGCAACAATGCAACGTACT
Gstt3_fw	GGAGCTCTACCTGGACCTGA
Gstt3_rv	AAGATGGCCACACTCTCTGC
Eva1c_fw	GTTGCCTACGCATGTGTTCC
Eva1c_rv	CCGATGCAGACACTGGACAT
Tspo_fw	GTATTCAGCCATGGGGTATGG
Tspo_rv	AAGCAGAAGATCGGCCAAGG
Tbp_fw	GGGGAGCTGTGATGTGAAGT
Tbp_rv	CCAGGAAATAATTCTGGCTCAT

Statistics and Repeatability

Mouse experiments:

Data are shown as mean±SEM. One-way ANOVA followed by Tukey's multiple

comparisons post-test was applied to compare experimental groups. Statistical analysis was

performed using GraphPad Prism 6.0 software. A p-value <0.05 was considered statistically

significant (*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001).

Drosophila experiments:

Statistical tests as well as the number of embryos/cells/tracks/contacts assessed are listed in

the Figure legends. All statistical analyses were performed using GraphPad PRISM or R

Studio and significance was determined using a 95% confidence interval. No statistical

method was used to predetermine sample size.

Representative images of Dfos antibody staining were analyzed per replicate per genotype

and *in situ* hybridization are from experiments that were repeated 2 times with many

embryos with reproducible results. *Dfos* mutants analysis in Fig 1 and S1 Fig are from

experiments that were repeated 2-3 times. In live imaging experiments in Fig 2 and S2 Fig,

3-7 embryos for each genotype were analyzed, each embryo was recorded on a separate day.

FACS sorting of macrophages from embryos was conducted in three replicates, from which

RNA samples were prepared for RNA sequencing. Experiments in Fig 4 and S4 Fig were

repeated at least 3 times, with representative images and plots of phalloidin immunostaining

from experiments that were repeated 4 times. In the LifeAct::GFP protrusion live imaging

experiment in Fig 5 and S5 Fig, 3-5 embryos were analyzed for each genotype. In

CLIP::GFP live imaging experiments in Fig 5 and S5 Fig, 5-6 embryos were analyzed for

each genotype for the cell aspect ratio in the germband zone, and 2 embryos in the pre-

germband zone and for tracking of the front vs rear speed. Each embryo was recorded on a

separate day. The Lamin over expression in S5 Fig and the Lamin knockdown rescue

experiments in Fig 5G were repeated at least 3 times. The gb rescue experiment in Fig 5H

was repeated at least 4 times.

Acknowledgements

We thank the following for their contributions: The *Drosophila* Genomics Resource Center

supported by NIH grant 2P40OD010949-10A1 for plasmids, K. Brueckner. B. Stramer, M.

Uhlirova, O. Schuldiner, the Bloomington *Drosophila* Stock Center supported by NIH grant

P40OD018537 and the Vienna *Drosophila* Resource Center for fly stocks, FlyBase

(Thurmond et al., 2019) for essential genomic information, and the BDGP *in situ* database

for data (Tomancak et al., 2002, 2007). For antibodies, we thank the Developmental Studies

Hybridoma Bank, which was created by the Eunice Kennedy Shriver National Institute of

Child Health and Human Development of the NIH, and is maintained at the University of

Iowa, as well as J. Zeitlinger for her generous gift of Dfos antibody. We thank the Vienna

BioCenter Core Facilities for RNA sequencing and analysis and the Life Scientific Service

Units at IST Austria for technical support and assistance with microscopy and FACS

analysis. We thank C. P. Heisenberg, P. Martin, M. Sixt and Siekhaus group members for

discussions and T. Hurd, A. Ratheesh and P. Rangan for comments on the manuscript.

References

Anders, S., Pyl, P. T., & Huber, W. (2015). HTSeq-A Python framework to work with high-throughput
sequencing data. *Bioinformatics*, *31*(2), 166–169. https://doi.org/10.1093/bioinformatics/btu638

Abreu-Blanco, M.T., Verboon, J.M., & Parkhurst S.M. (2014) Coordination of Rho Family GTPase activities to
orchestrate cytoskeleton responses during wound repair. *Current Biology*, *24*(2), 144-155.
https://doi.org/10.1016/j.cub.2013.11.048

Andrés, V., & González, J. M. (2009). Role of A-type lamins in signaling, transcription, and chromatin
organization. *Journal of Cell Biology*, *187*(7), 945–957. https://doi.org/10.1083/jcb.200904124

Bellanger, J. M., Astier, C., Sardet, C., Ohta, Y., Stossel, T. P., & Debant, A. (2000). The Rac1- and RhoG-

specific GEF domain of trio targets filamin to remodel cytoskeletal actin. *Nature Cell Biology*, *2*(12),

888–892. https://doi.org/10.1038/35046533

Benhra, N., Barrio, L., Muzzopappa, M., & Milán, M. (2018). Chromosomal Instability Induces Cellular

Invasion in Epithelial Tissues. *Developmental Cell*, *47*(2), 161-174.e4.

https://doi.org/10.1016/j.devcel.2018.08.021

Berditchevski, F., & Odintsova, E. (1999). Characterization of integrin-tetraspanin adhesion complexes: Role of

tetraspanins in integrin signaling. *Journal of Cell Biology*, *146*(2), 477–492.

https://doi.org/10.1083/jcb.146.2.477

Bershadsky, A. D., Balaban, N. Q., & Geiger, B. (2003). Adhesion-Dependent Cell Mechanosensitivity. *Annual*

Review of Cell and Developmental Biology, *19*(1), 677–695.

https://doi.org/10.1146/annurev.cellbio.19.111301.153011

Bilancia, C. G., Winkelman, J. D., Tsygankov, D., Nowotarski, S. H., Sees, J. A., Comber, K., … Peifer, M.

(2014). Enabled negatively regulates diaphanous-driven actin dynamics in vitro and in vivo.

Developmental Cell, *28*(4), 394–408. https://doi.org/10.1016/j.devcel.2014.01.015

Bosch, P.S., Makhijani, K., Herboso, L., Gold, K.S., Baginsky, R., Woodcock, K.J., Alexander, B., Kukar, K.,

Corcoran, S., Jacobs, T., Ouyang, D., Wong, C., Ramond, E.J.V., Rhiner, C., Moreno, E., Lemaitre, B.,

Geissmann, F., & Brueckner, K. (2019). Adult *Drosophila* lack hematopoiesis but rely on a blood cell

reservoir at the respiratory epithelia to relay infection signals to surrounding tissues. *Developmental Cell*,

51(6)787-803.

https://doi.org/10.1016/j.devcel.2019.10.017

Brock, A. R., Wang, Y., Berger, S., Renkawitz-Pohl, R., Han, V. C., Wu, Y., & Galko, M. J. (2012).

Transcriptional regulation of profilin during wound closure in Drosophila larvae. *Journal of Cell Science*,

125(23), 5667–5676. https://doi.org/10.1242/jcs.107490

Brückner, K., Kockel, L., Duchek, P., Luque, C. M., Rørth, P., & Perrimon, N. (2004). The PDGF/VEGF

receptor controls blood cell survival in Drosophila. *Developmental Cell*, *7*(1), 73–84.

https://doi.org/10.1016/j.devcel.2004.06.007

Brzozowski, J. S., Bond, D. R., Jankowski, H., Goldie, B. J., Burchell, R., Naudin, C., … Weidenhofer, J.

(2018). Extracellular vesicles with altered tetraspanin CD9 and CD151 levels confer increased prostate

cell motility and invasion. *Scientific Reports*, *8*(1). https://doi.org/10.1038/s41598-018-27180-z

Butcher, D.T., Alliston, T. & Weaver, V.M.(2009). A tense situation: forcing tumour progression. *Nature*

Reviews Cancer, 9(2), 108-22. . https://doi.org/10.11038/nrc2544

Cho, N. K., Keyes, L., Johnson, E., Heller, J., Ryner, L., Karim, F., & Krasnow, M. A. (2002). Developmental

control of blood cell migration by the Drosophila VEGF pathway. *Cell*, *108*(6).

https://doi.org/10.1016/S0092-8674(02)00676-1

Chugh, P., Clark, A. G., Smith, M. B., Cassani, D. A. D., Dierkes, K., Ragab, A., … Paluch, E. K. (2017). Actin

cortex architecture regulates cell surface tension. *Nature Cell Biology*, *19*(6), 689–697.

https://doi.org/10.1038/ncb3525

Danuser, G., Allard, J., & Mogilner, A. (2013). Mathematical Modeling of Eukaryotic Cell Migration: Insights

Beyond Experiments. *Annual Review of Cell and Developmental Biology*, *29*(1).

https://doi.org/10.1146/annurev-cellbio-101512-122308

Davidson, A. J., Millard, T. H., Evans, I. R., & Wood, W. (2019). Ena orchestrates remodelling within the actin

cytoskeleton to drive robust Drosophila macrophage chemotaxis. *Journal of Cell Science*, *132*(5).

https://doi.org/10.1242/jcs.224618

Davidson, P.M., Denais, C., Bakshi, M., & Lammerding, J. (2014).. Nuclear deformability constitutes a rate-

limiting step during cell migration in 3-D environments. *Cellular and Molecular Bioengineering,*

7(3)293-306. https://doi.org/10.1007/s12195-014-0342-y

Davis, J. R., Luchici, A., Miodownik, M., Stramer, B. M., Davis, J. R., Luchici, A., … Stramer, B. M. (2015).

Inter-Cellular Forces Orchestrate Contact Inhibition of Locomotion Article Inter-Cellular Forces

Orchestrate Contact Inhibition of Locomotion. *Cell*, *161*(2), 361–373.

https://doi.org/10.1016/j.cell.2015.02.015

Delaguillaumie, A., Lagaudrière-Gesbert, C., Popoff, M. R., & Conjeaud H., H. (2002). Rho GTPase link

cytoskeletal rearrangements and activation processes induced via the tetraspanin CD82 in T lymphocytes.

Journal of Cell Science, *115*(2).

Deluca, T. F., Cui, J., Jung, J. Y., St. Gabriel, K. C., & Wall, D. P. (2012). Roundup 2.0: Enabling comparative

genomics for over 1800 genomes. *Bioinformatics*, *28*(5). https://doi.org/10.1093/bioinformatics/bts006

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., … Gingeras, T. R. (2013). STAR:

Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21.

https://doi.org/10.1093/bioinformatics/bts635

Duffy, J. B. (2002). GAL4 system in Drosophila: A fly geneticist's Swiss army knife. *Genesis*, *34*(1–2), 1–15.

https://doi.org/10.1002/gene.10150

Edwards, K. A., Demsky, M., Montague, R. A., Weymouth, N., & Kiehart, D. P. (1997). GFP-moesin

illuminates actin cytoskeleton dynamics in living tissue and demonstrates cell shape changes during

morphogenesis in Drosophila. *Developmental Biology*, *191*(1). https://doi.org/10.1006/dbio.1997.8707

Ehrlicher, A. J., Nakamura, F., Hartwig, J. H., Weitz, D. A., & Stossel, T. P. (2011). Mechanical strain in actin

networks regulates FilGAP and integrin binding to filamin A. *Nature*, *478*(7368), 260–263.

https://doi.org/10.1038/nature10430

Eresh, S., Riese, J., Jackson, D. B., Bohmann, D., & Bienz, M. (1997). A CREB-binding site as a target for

decapentaplegic signalling during Drosophila endoderm induction. *EMBO Journal*, *16*(8), 2014–2022.

https://doi.org/10.1093/emboj/16.8.2014

Evans, I. R., & Wood, W. (2011). Drosophila embryonic hemocytes. *Current Biology*, *21*(5), R173–R174.

https://doi.org/10.1016/j.cub.2011.01.061

Franck, Z., Gary, R., & Bretscher, A. (1993). Moesin, like ezrin, colocalizes with actin in the cortical

cytoskeleton in cultured cells, but its expression is more variable. *Journal of Cell Science*, *105*(1).

Fujita, M., Mitsuhashi, H., Isogai, S., Nakata, T., Kawakami, A., Nonaka, I., … Kudo, A. (2012). Filamin C

plays an essential role in the maintenance of the structural integrity of cardiac and skeletal muscles,

revealed by the medaka mutant zacro. *Developmental Biology*, *361*(1), 79–89.

https://doi.org/10.1016/j.ydbio.2011.10.008

García-Echeverría, C. Methionine-containing zipper peptides. *Lett Pept Sci* **4,** 135–140 (1997).

https://doi.org/10.1007/BF02443525

Ginhoux, F.& Guilliams, M. (2016). Tissue-resident macrophage ontogeny and homeostasis. *Immunity 44*(3),

439-449. https://doi.org/10.1016/j.immuni.2016.02.024

Glogauer, M., Arora, P., Chou, D., Janmey, P. A., Downey, G. P., & McCulloch, C. A. G. (1998). The role of

actin-binding protein 280 in integrin-dependent mechanoprotection. *Journal of Biological Chemistry*,

273(3), 1689–1698. https://doi.org/10.1074/jbc.273.3.1689

Glover, J.N.M. and Harrison, S.C. (1005). Crystal structure of the heterodimeric bZIP transcritpion factor c-Fos-

c-Jun bound to DNA. *Nature, 373*(6511):257-261. https://doi.org/10.1038/373257a0

Goldmann, W. H., Tempel, M., Sprenger, I., Isenberg, G., & Ezzell, R. M. (1997). Viscoelasticity of actin-

gelsolin networks in the presence of filamin. *European Journal of Biochemistry*, *246*(2), 373–379.

https://doi.org/10.1111/j.1432-1033.1997.00373.x

Gonzalez-Gaitan, M. & Peifer, M. (2009) Exploring the roles of Diaphanous and Enabled activity in shaping the

balance between filopodia and lamellipodia. *Molecular Biology of the Cell 20*(24).

https://doi.org/10.1091/mbc.e09-02-0144

Goode, B.L. & Eck. M. (2007) Mechanism and Function of Formins in the Control of Actin Assembly. *Annual*

Review of Biochemistry 78, 593-627. https://doi.org/10.1146/annurev.biochem.75.103004.142647

Greten, F.R. & Grovennikov, S.I. (2019). Inflammation and Cancer: Triggers, Mechanisms and Consequences.

Immunity 51(1), 27-41. https://doi.org/10.1016/j.immuni.2019.06.025

Großhans, J., Wenzl, C., Herz, H. M., Bartoszewski, S., Schnorrer, F., Vogt, N., … Müller, H. A. (2005).

RhoGEF2 and the formin Dia control the formation of the furrow canal by directed actin assembly during

Drosophila cellularisation. *Development*, *132*(5), 1009–1020. https://doi.org/10.1242/dev.01669

Guilliams, M., Thierry, G.R., Bonnardel, J., & Bajenoff, M. (2020). Establishment and Maintenance of the

Macrophage Niche. *Immunity 52*(3), 434-451. https://doi.org/10.1016/j.immuni.2020.02.015

Gyoergy, A., Roblek, M., Ratheesh, A., Valoskova, K., Belyaeva, V., Wachner, S., … Siekhaus, D. E. (2018).

Tools allowing independent visualization and genetic manipulation of Drosophila melanogaster

macrophages and surrounding tissues. *G3: Genes, Genomes, Genetics*, *8*(3).

https://doi.org/10.1534/g3.117.300452

Hammonds, A. A. S., Bristow, C. C. a, Fisher, W. W., Weiszmann, R., Wu, S., Hartenstein, V., … Celniker, S.

E. (2013). Spatial expression of transcription factors in Drosophila embryonic organ development.

Genome Biology, *14*(12), R140. https://doi.org/10.1186/gb-2013-14-12-r140

Harada, T., Swift, J., Irianto, J., Shin, J, Spinler, K.R., Athirasala, A., Diegmiller, R., Dingal, P.C.D.P.,

Ivanovska, I.L., & Discher, D.E. (2014). Nuclear lamin stiffness is a barrier to 3D migration, but softness

can limit survival. *J. Cell Biology,* (5)669-82. https://doi.org/10.1083/jcb.201308029.

Hattori, A., Mizuno, T., Akamatsu, M., Hisamoto, N., Matsumoto, K. (2013). The Caenorhabditis elegans JNK

signaling pathway activates expression of stress response genes by depressing the Fos/HDAC repressor

complex. *Plos Genetics 9*(2):e1003315. https://doi.org/10.1371/journal.pgen.1003315

Heer, N. C., & Martin, A. C. (2017). Tension, contraction and tissue morphogenesis. *Development (Cambridge)*,

144(23)4249-4260. https://doi.org/10.1242/dev.151282

Holz, A., Bossinger, B., Strasser, T., Janning, W., & Klapper, R. (2003). The two origins of hemocytes in

Drosophila. Development, 130(20), 4955-62. https://doi.org/10.1242/dev.007202

Homem, C.C., Peifer, M. (2008) Diaphanous regulates myosin and adherens junctions to control cell

contractility and protrusive behavior during morphogenesis. *Development*, *135*(6)1005-1018.

https://doi.org/10.1242/dev.151282

Hong, I. K., Jeoung, D. Il, Ha, K. S., Kim, Y. M., & Lee, H. (2012). Tetraspanin CD151 stimulates adhesion-

dependent activation of Ras, Rac, and Cdc42 by facilitating molecular association between β1 integrins

and small GTPases. *Journal of Biological Chemistry*, *287*(38), 32027–32039.

https://doi.org/10.1074/jbc.M111.314443

Hu, J., Lu, J., Lian, G., Ferland, R. J., Dettenhofer, M., & Sheen, V. L. (2014). Formin 1 and filamin B

physically interact to coordinate chondrocyte proliferation and differentiation in the growth plate. *Human*

Molecular Genetics, *23*(17), 4663–4673. https://doi.org/10.1093/hmg/ddu186

Kasza, K. E., Broedersz, C. P., Koenderink, G. H., Lin, Y. C., Messner, W., Millman, E. A., … Weitz, D. A.

(2010). Actin filament length tunes elasticity of flexibly cross-linked actin networks. *Biophysical Journal*,

99(4), 1091–1100. https://doi.org/10.1016/j.bpj.2010.06.025

Kessenbrock, K., Plaks, V., & Werb, Z. (2010). Matrix metalloproteinases: regulators of the tumor

microenvironment. *Cell 141*(1), 52-67. https://doi.org/10.1016/j.cell.2010.03.015

Kühn, S., & Geyer, M. (2014). Formins as effector proteins of rho GTPases. *Small GTPases*, *5*(JUNE).

https://doi.org/10.4161/sgtp.29513

Külshammer, E., Mundorf, J., Kilinc, M., Frommolt, P., Wagle, P., & Uhlirova, M. (2015). Interplay among

Drosophila transcription factors Ets21c, Fos and Ftz-F1 drives JNK-mediated tumor malignancy. *DMM*

Disease Models and Mechanisms, *8*(10), 1279–1293. https://doi.org/10.1242/dmm.020719

Külshammer, E., & Uhlirova, M. (2013). The actin cross-linker Filamin/Cheerio mediates tumor malignancy

downstream of JNK signaling. *Journal of Cell Science*, *126*(4), 927–938.

https://doi.org/10.1242/jcs.114462

Kumar, A., Shutova, M. S., Tanaka, K., Iwamoto, D. V., Calderwood, D. A., Svitkina, T. M., & Schwartz, M.

A. (2019). Filamin A mediates isotropic distribution of applied force across the actin network. *Journal of*

Cell Biology, *218*(8), 2481–2491. https://doi.org/10.1083/jcb.201901086

Lehmann, R., & Tautz, D. (1994). In Situ Hybridization to RNA. *Methods in Cell Biology*, *44*(C), 575–596.

https://doi.org/10.1016/S0091-679X(08)60933-4

Lemaitre, B., & Hoffmann, J. (2007). The Host Defense of *Drosophila melanogaster*. *Annual Review of*

Immunology, *25*(1), 697–743. https://doi.org/10.1146/annurev.immunol.25.022106.141615

Lesch, C., Jo, J., Wu, Y., Fish, G. S., & Galko, M. J. (2010). A Targeted *UAS-RNAi* Screen in Drosophila

Larvae Identifies Wound Closure Genes Regulating Distinct Cellular Processes. *Genetics*, *186*(3), 943–

957. https://doi.org/10.1534/genetics.110.121822

Lian, G., Dettenhofer, M., Lu, J., Downing, M., Chenn, A., Wong, T., & Sheen, V. (2016). Filamin A- and

formin 2-dependent endocytosis regulates proliferation via the canonical wnt pathway. *Development*

(Cambridge), *143*(23). https://doi.org/10.1242/dev.139295

Linder, M., Glitzner, E., Srivatsa, S., Bakiri, L., Matsuoka, K., Shahrouzi, P., … Sibilia, M. (2018). EGFR is

required for FOS☐dependent bone tumor development via RSK2/CREB signaling. *EMBO Molecular*

Medicine, *10*(11). https://doi.org/10.15252/emmm.201809408

Luster, A. D., Alon, R., & von Andrian, U. H. (2005). Immune cell migration in inflammation: Present and

future therapeutic targets. *Nature Immunology*, *6*(12), 1182–1190. https://doi.org/10.1038/ni1275

Makhijani, K., Alexander, B., Tanaka, T., Rulfson, E., & Brückner, K. (2011). The peripheral nervous system

supports blood cell homing and survival in the Drosophila larva. *Development, 138*(24), 5379-91.

https://doi.org/10.1242/dev.067322

Matsubayashi, Y., Louani, A., Dragu, A., Sánchez-Sánchez, B. J., Serna-Morales, E., Yolland, L., … Stramer,

B. M. (2017). A Moving Source of Matrix Components Is Essential for De Novo Basement Membrane

Formation. *Current Biology*, *27*(22), 3526-3534.e4. https://doi.org/10.1016/j.cub.2017.10.001

Min, P. I., Spurney, R. F., Qisheng, T. U., Hinson, T., & Darryl Quarles, L. (2002). Calcium-sensing receptor

activation of Rho involves filamin and rho-guanine nucleotide exchange factor. *Endocrinology*, *143*(10),

3830–3838. https://doi.org/10.1210/en.2002-220240

Mitchison, T.J., and Cramer, L.P. (1996) Actin-based cell motility and cell locomotion. *Cell 84*(3)371-9).

https://doi.org/10.1016/s0092-8864(00)81281-7.

Muñoz-Alarcón, A., Pavlovic, M., Wismar, J., Schmitt, B., Eriksson, M., Kylsten, P., & Dushay, M. S. (2007).

Characterization of lamin mutation phenotypes in Drosophila and comparison to human laminopathies.

PloS One, *2*(6). https://doi.org/10.1371/journal.pone.0000532

Ohta, Y., Suzuki, N., Nakamura, S., Hartwig, J. H., & Stossel, T. P. (1999). The small GTPase RaIA targets

filamin to induce filopodia. *Proceedings of the National Academy of Sciences of the United States of*

America, *96*(5), 2122–2128. https://doi.org/10.1073/pnas.96.5.2122

Paluch, E. K., Aspalter, I. M., & Sixt, M. (2016). Focal Adhesion–Independent Cell Migration. *Annual Review*

of Cell and Developmental Biology, *32*(1), 469–490. https://doi.org/10.1146/annurev-cellbio-111315-

125341

Perez-Hernandez, D., Gutiérrez-Vázquez, C., Jorge, I., López-Martín, S., Ursa, A., Sánchez-Madrid, F., …

Yañez-Mó, M. (2013). The intracellular interactome of tetraspanin-enriched microdomains reveals their

function as sorting machineries toward exosomes. *Journal of Biological Chemistry*, *288*(17), 11649–

11661. https://doi.org/10.1074/jbc.M112.445304

Perkins, L. A., Holderbaum, L., Tao, R., Hu, Y., Sopko, R., McCall, K., … Perrimon, N. (2015). The transgenic

RNAi project at Harvard medical school: Resources and validation. *Genetics*, *201*(3), 843–852.

https://doi.org/10.1534/genetics.115.180208

Popowicz, G. M., Schleicher, M., Noegel, A. A., & Holak, T. A. (2006). Filamins: promiscuous organizers of

the cytoskeleton. *Trends in Biochemical Sciences*, *31*(7), 411–419.

https://doi.org/10.1016/j.tibs.2006.05.006

Raab, M., Gentili, M., de Belly, H., Thiam, H.R., Vargas, P., Jimenez, A.J., Lautenschlaeger, F., Voituriez, R.,

Lennon-Duménil, A.M., Manel, N, Piel, M. (2016). ESCRT III repairs nuclear envelope ruptures during

cell migration to limit DNA damage and cell death. Science, 352(6283), 359-62.

https://doi.org/10.1126/science.aad7611

Ratheesh, A., Belyaeva, V., & Siekhaus, D. E. (2015). Drosophila immune cell migration and adhesion during

embryonic development and larval immune responses. *Current Opinion in Cell Biology*, *36*, 71–79.

https://doi.org/10.1016/j.ceb.2015.07.003

Ratheesh, A., Biebl, J., Vesela, J., Smutny, M., Papusheva, E., Krens, S. F. G., … Siekhaus, D. E. (2018).

Drosophila TNF Modulates Tissue Tension in the Embryo to Facilitate Macrophage Invasive Migration.

Developmental Cell, *45*(3), 331-346.e7. https://doi.org/10.1016/j.devcel.2018.04.002

Razinia, Z., Mäkelä, T., Ylänne, J., & Calderwood, D. A. (2012). Filamins in Mechanosensing and Signaling.

Annual Review of Biophysics, *41*(1), 227–246. https://doi.org/10.1146/annurev-biophys-050511-102252

Riesgo-Escovar, J. R., & Hafen, E. (1997). Common and distinct roles of DFos and DJun during Drosophila

development. *Science (New York, N.Y.)*, *278*(5338), 669–672.

https://doi.org/10.1126/science.278.5338.669

Riveline, D., Zamir, E., Balaban, N. Q., Schwarz, U. S., Ishizaki, T., Narumiya, S., … Bershadsky, A. D.

(2001). Focal contacts as mechanosensors: Externally applied local mechanical force induces growth of

focal contacts by an mDia1-dependent and ROCK-independent mechanism. *Journal of Cell Biology*,

153(6), 1175–1185. https://doi.org/10.1083/jcb.153.6.1175

Rose, R., Weyand, M., Lammers, M., Ishizaki, T., Ahmadian, M. R., & Wittinghofer, A. (2005). Structural and

mechanistic insights into the interaction between Rho and mammalian Dia. *Nature*, *435*(7041), 513–518.

https://doi.org/10.1038/nature03604

Rousso, T., Shewan, A. M., Mostov, K. E., Schejter, E. D., & Shilo, B. Z. (2013). Apical targeting of the formin

diaphanous in Drosophila tubular epithelia. *ELife*, *2013*(2). https://doi.org/10.7554/eLife.00666

Sánchez-Sánchez, B. J., Urbano, J. M., Comber, K., Dragu, A., Wood, W., Stramer, B., & Martín-Bermudo, M.

D. (2017). Drosophila Embryonic Hemocytes Produce Laminins to Strengthen Migratory Response. *Cell*

Reports, *21*(6), 1461–1470. https://doi.org/10.1016/j.celrep.2017.10.047

Seth, A., Otomo, C., & Rosen, M. K. (2006). Autoinhibition regulates cellular localization and actin assembly

activity of the diaphanous-related formins FRLα and mDia1. *Journal of Cell Biology*, *174*(5), 701–713.

https://doi.org/10.1083/jcb.200605006

Sharma, P. & Allison, J.P. (2015) The future of immune checkpoint therapy. *Science 348*(6230), 56-61.

https://doi.org/10.1126/science.aaa8172.

Shigeta, M., Sanzen, N., Ozawa, M., Gu, J., Hasegawa, H., & Sekiguchi, K. (2003). CD151 regulates epithelial

cell-cell adhesion through PKC- and Cdc42-dependent actin cytoskeletal reorganization. *Journal of Cell*

Biology, *163*(1), 165–176. https://doi.org/10.1083/jcb.200301075

Siekhaus, D., Haesemeyer, M., Moffitt, O., & Lehmann, R. (2010). RhoL controls invasion and Rap1

localization during immune cell transmigration in Drosophila. *Nature Cell Biology*, *12*(6), 605–610.

https://doi.org/10.1038/ncb2063

Sit, S. T., & Manser, E. (2011). Rho GTPases and their role in organizing the actin cytoskeleton. *Journal of Cell*

Science, *124*(5), 679–683. https://doi.org/10.1242/jcs.064964

Smutny, M., Ákos, Z., Grigolon, S., Shamipour, S., Ruprecht, V., Čapek, D., … Heisenberg, C. P. (2017).

Friction forces position the neural anlage. *Nature Cell Biology*, *19*(4), 306–317.

https://doi.org/10.1038/ncb3492

Sokol, N. S., & Cooley, L. (2003). Drosophila filamin is required for follicle cell motility during oogenesis.

Developmental Biology, *260*(1), 260–272. https://doi.org/10.1016/S0012-1606(03)00248-3

Somogyi, K., & Rørth, P. (2004). Evidence for tension-based regulation of Drosophila MAL and SRF during

invasive cell migration. *Developmental Cell*, *7*(1), 85–93. https://doi.org/10.1016/j.devcel.2004.05.020

Stossel, T. P., Condeelis, J., Cooley, L., Hartwig, J. H., Noegel, A., Schleicher, M., & Shapiro, S. S. (2001).

Filamins as integrators of cell mechanics and signalling. *Nature Reviews Molecular Cell Biology*, *2*(2),

138–145. https://doi.org/10.1038/35052082

Szabo, P.A., Miron, M., & Farber, D.L. (2019). Location, location, location: Tissue residence memory T cells in

mice and humans. *Science Immunology 4*(34), https://doi.org/10.1126/scieimunol.aas9673

Szalóki, N., Krieger, J. W., Komáromi, I., Tóth, K., & Vámosi, G. (2015). Evidence for Homodimerization of

the c-Fos Transcription Factor in Live Cells Revealed by Fluorescence Microscopy and Computer

Modeling. *Molecular and Cellular Biology*, *35*(21). https://doi.org/10.1128/mcb.00346-15

Tejera, E., Rocha-Perugini, V., López-Martín, S., □rez-Hernández, D., Bachir, A. I., Horwitz, A. R., … Yáñez-

Mo, M. (2013). CD81 regulates cell migration through its association with Rac GTPase. *Molecular*

Biology of the Cell, *24*(3), 261–273. https://doi.org/10.1091/mbc.E12-09-0642

Termini, C. M., & Gillette, J. M. (2017). Tetraspanins Function as Regulators of Cellular Signaling. *Frontiers in*

Cell and Developmental Biology, *5*, 34. https://doi.org/10.3389/fcell.2017.00034

Theret, M., Mounier, R., & Rossi, F. (2019). The origins and non-canonical functions of macrophages in

development and regeneration. *Development, 146*(9). https://doi.org/10.1242/dev.156000

Thiam, H., Vargas, P., Carpi, N., Crespo, C.L., Raab, M., Terriac, E., King, M.C., Jacobelli, J., Alberts, A.S.,

Stradal, T., Lennon-Dumenil, A., Piel, M. (2016). Perinuclear Arp2/3-driven actin polymerization enables

nuclear deformation to facilitate cell migration through complex environments. Stremmel:10997,

https://doi.org/10.1038/ncomms10997

Thurmond, J., Goodman, J. L., Strelets, V. B., Attrill, H., Gramates, L. S., Marygold, S. J., … Baker, P. (2019).

FlyBase 2.0: The next generation. *Nucleic Acids Research*, *47*(D1). https://doi.org/10.1093/nar/gky1003

Tomancak, P., Beaton, A., Weiszmann, R., Kwan, E., Shu, S. Q., Lewis, S. E., … Rubin, G. M. (2002).

Systematic determination of patterns of gene expression during Drosophila embryogenesis. *Genome*

Biology, *3*(12). https://doi.org/10.1186/gb-2002-3-12-research0088

Tomancak, P., Berman, B. P., Beaton, A., Weiszmann, R., Kwan, E., Hartenstein, V., … Rubin, G. M. (2007).

Global analysis of patterns of gene expression during Drosophila embryogenesis. *Genome Biology*, *8*(7).

https://doi.org/10.1186/gb-2007-8-7-r145

Tseng, Y., An, K. M., Esue, O., & Wirtz, D. (2004). The Bimodal Role of Filamin in Controlling the

Architecture and Mechanics of F-actin Networks. *Journal of Biological Chemistry*, *279*(3), 1819–1826.

https://doi.org/10.1074/jbc.M306090200

Uhlirova, M., & Bohmann, D. (2006). JNK- and Fos-regulated Mmp1 expression cooperates with Ras to induce

invasive tumors in Drosophila. *EMBO Journal*, *25*(22), 5294–5304.

https://doi.org/10.1038/sj.emboj.7601401

Vadlamudi, R. K., Li, F., Adam, L., Nguyen, D., Ohta, Y., Stossel, T. P., & Kumar, R. (2002). Filamin is

essential in actin cytoskeletal assembly mediated by p21-activated kinase 1. *Nature Cell Biology*, *4*(9),

681–690. https://doi.org/10.1038/ncb838

Valoskova, K., Biebl, J., Roblek, M., Emtenani, S., Gyoergy, A., Misova, M., … Siekhaus, D. E. (2019). A

conserved major facilitator superfamily member orchestrates a subset of O-glycosylation to aid

macrophage tissue invasion. *ELife*, *8*. https://doi.org/10.7554/eLife.41801

Vetter, I.R., & Wittinghofer, A. (2001). The guanine nucleotide-binding switch in three dimensions. *Science*

*294(*5545),1299-1304. https://doi.org/10.1126/science.1062023

Warner, S. J., & Longmore, G. D. (2009). Cdc42 antagonizes Rho1 activity at adherens junctions to limit

epithelial cell apical tension. *Journal of Cell Biology*, *187*(1), 119–133.

https://doi.org/10.1083/jcb.200906047

Weavers, H., Evans, I. R., Martin, P., & Wood, W. (2016). Corpse Engulfment Generates a Molecular Memory

that Primes the Macrophage Inflammatory Response. *Cell*. https://doi.org/10.1016/j.cell.2016.04.049

Williams, M. J., Habayeb, M. S., & Hultmark, D. (2007). Reciprocal regulation of Rac1 and Rho1 in Drosophila

circulating immune surveillance cells. *Journal of Cell Science*, *120*(3), 502–511.

https://doi.org/10.1242/jcs.03341

Wintner, O., Hirsch-Attas, N., Schlossberg, M., Brofman, F., Friedman, R., Kupervaser, M., … Buxboim, A.

(2020). A Unified Linear Viscoelastic Model of the Cell Nucleus Defines the Mechanical Contributions of

Lamins and Chromatin. *Advanced Science*, *7*(8). https://doi.org/10.1002/advs.201901222

Wood, W., Faria, C., & Jacinto, A. (2006). Distinct mechanisms regulate hemocyte chemotaxis during

development and wound healing in Drosophila melanogaster. *Journal of Cell Biology*, *173*(3).

https://doi.org/10.1083/jcb.200508161

Yáñez-Mó, M., Barreiro, O., Gordon-Alonso, M., Sala-Valdés, M., & Sánchez-Madrid, F. (2009). Tetraspanin-

enriched microdomains: a functional unit in cell plasma membranes. *Trends in Cell Biology*, *19*(9), 434–

446. https://doi.org/10.1016/j.tcb.2009.06.004

Yeung, L., Hickey, M.J., & Wright, M.D. (2018) The many and varied roles of tetraspanins in immune cell

recrutment and migration. Frontiers in Immunology.9:1644 .https:// doi: 10.3389/fimmu.2018.01644.

Zeitlinger, J., Kockel, L., Peverali, F. A., Jackson, D. B., Mlodzik, M., & Bohmann, D. (1997). Defective dorsal

closure and loss of epidermal decapentaplegic expression in Drosophila fos mutants. *EMBO Journal*,

16(24), 7393–7401. https://doi.org/10.1093/emboj/16.24.7393

Zhang, X. A., Bontrager, A. L., & Hemler, M. E. (2001). Transmembrane-4 Superfamily Proteins Associate

with Activated Protein Kinase C (PKC) and Link PKC to Specific β1 Integrins. *Journal of Biological*

Chemistry, *276*(27), 25005–25013. https://doi.org/10.1074/jbc.M102156200

Zhou, J., Kim, H. Y., & Davidson, L. A. (2009). Actomyosin stiffens the vertebrate embryo during crucial

stages of elongation and neural tube closure. *Development*, *136*(4), 677–688.

https://doi.org/10.1242/dev.026211

Zimmerman, B., Kelly, B., McMillan, B. J., Seegar, T. C. M., Dror, R. O., Kruse, A. C., & Blacklow, S. C.

(2016). Crystal Structure of a Full-Length Human Tetraspanin Reveals a Cholesterol-Binding Pocket.

Cell, *167*(4). https://doi.org/10.1016/j.cell.2016.09.056

Zimmerman, B., Kelly, B., McMillan, B., Seegar, T., Kruse, A., & Blacklow, S. (2016). Crystal Structure of

Human Tetraspanin CD81 Reveals a Conserved Intramembrane Binding Cavity. *Cell*, *167*(4), 1041–1051.

https://doi.org/10.1016/j.cell.2016.09.056

Zwerger, M., Jaalouk, D. E., Lombardi, M. L., Isermann, P., Mauermann, M., Dialynas, G., … Lammerding, J.

(2013). Myopathic lamin mutations impair nuclear stability in cells and tissue and disrupt nucleo-

cytoskeletal coupling. *Human Molecular Genetics*, *22*(12), 2335–2349. https://doi.org/10.109

1378

1379

1380

1381   **Supplementary Information Titles/ Legends**

1382

1383   **S1 Data. Data used to plot all graphs and to perform statistical analyses**

1384

1385   **S2 Data. RNA sequencing data**

1386

S1 Fig. Dfos does not affect the total number of macrophages, or their number in the

pre-gb zone and along the vnc

(A) Dfos protein (red) is detected with an antibody in macrophages (green) in embryos from

the stages as indicated.

(**B-I**) Quantification in mid St 12 embryos.

(**B**) The number of macrophages (green) in the pre-gb zone (outlined by a black dotted line in

the schematic on the left) showed no significant change in $Dfos^2$ mutant embryos compared

to the control (p=0.37) SD: 6,7.

(**C**) The total number of macrophages (see schematic at left) was not altered from that in the

control embryos expressing DfosDN in macrophages (p=0.12). SD: 60, 120.

(**D-E**) The number of macrophages (green) along the vnc (outlined by black dotted line in the

schematic on the left) shows no significant difference between the control and (**D**)

macrophages that express DfosDN or (**E**) either of two RNAi lines against *Dfos*. (**D**) *DfosDN*

p=0.88, 0.99, >0.99. *Dfos RNAi^1* (TRiP HMS00254) p=0.21, 0.06, 0.11, 0.072, 0.033, 0.30,

0.56. *Dfos RNAi^2* (TRiP JF02804) p=0.34, 0.15, 0.83, 0.27, 0.47, 1.0, 0.45. (D) SD: Ctrl 3, 3,

3, 0.8; *DfosDN* 6, 3, 0.7. (E) SD: Ctrl 6, 3, 3, 3, 2, 0.3; *Dfos RNAi^1* 6, 3, 3, 3, 2, 2, 0.3; *Dfos*

RNAi^2 6, 2, 3, 2, 3, 1, 0.4.

(**F, G**) Macrophage numbers in the pre-gb (see schematic at left) are increased compared to

the control for lines expressing (**F**) DfosDN or (**G**) one of two different *UAS-Dfos RNAi*

constructs in macrophages under *srpHemo-GAL4* control. (F) p=.04, SD: 19, 29. (G) *Dfos*

RNAi[1] p<0.0009, *Dfos RNAi[2]* p<0.0001. SD: 12, 9, 14.

(**H**) Macrophage numbers in the germband are not significantly altered compared to the

control upon overexpression of Dfos in macrophages (p=0.14). SD: 22, 14.

(**I**) Macrophage numbers in the gb for lines expressing one of two different *UAS-Dfos RNAi*

constructs in macrophages under *srpHemo-GAL4* control and lines which additionally

express *UAS-GFP*. Control vs. *mac>Dfos RNAi[1]* (TRiP HMS00254) or Control vs.

mac>Dfos RNAi[2] (TRiP JF02804), p<0.0001. *mac>Dfos RNAi[1]* vs. *mac>Dfos RNAi[1] + GFP*

or *mac>Dfos RNAi[2]* vs. *mac>Dfos RNAi[2] + GFP*, p>0.99. SD: 33, 47, 34. The effect of each

Dfos RNAi was eliminated upon simultaneous expression of another UAS construct.

Macrophages are labeled using either *srpHemo-Gal4* driving *UAS-GFP* or *srpHemo-*

H2A::3xmCherry. "*mac>*" indicates *srpHemo-GAL4* driver expressing UAS constructs

specifically in macrophages. Histograms show mean + SEM ***p<0.005, **p<0.01,

*p<0.05. Unpaired t-test was used for statistics, except for G, I, which used One-Way

ANOVA. The number of embryos analysed for that genotype is shown within each column in

the graphs. In D n=6 embryos for the control and n=9 for Dfos DN. In E n=9 embryos for

control, 15 and 11 for *Dfos RNAis*. Scale bar in A: 10 µm.

S2 Fig. Dfos facilitates macrophage motility during initial invasion into the tissue

(**A**) Quantification reveals that the directional persistence of macrophages expressing

DfosDN (0.58) is unchanged (0.56) in the pre-gb area (p=0.66) but decreased during gb entry

(0.65) (0.72), p=0.038 and along the vnc (0.54) compared to the control (0.61), p=0.00026.

Left schematic shows pre-gb area in yellow, gb entry outlined in solid line. Boxed area in

right schematic shows analyzed area of vnc.

(B) Movie stills showing wild type and *Dfos²* macrophages entering the gb (outlined by the

dashed line). Time in min shown in the top right corner of each image.

(C) Quantification of macrophage speed shows a significant reduction in the speed of *Dfos²*

macrophages in the pre-gb zone and at gb entry, but none in the head. Regions analysed

indicated in left schematic. Speed in head: control=2.59 µm/min, *Dfos²*=2.68 µm/min,

p=0.40; speed in pre-gb=3.38 µm/min, *Dfos²*=2.47 µm/min, p=2.38e^{-06}; speed in gb entry:

control=2.35 µm/min, *Dfos²*=1.62 µm/min, p= 0.0003.

Macrophages are labeled using *srpHemo-H2A::3xmCherry*. Histograms show mean + SEM

***p<0.005, **p<0.01, *p<0.05. Unpaired t-test was used for statistics. The number of

analyzed macrophages for each genotype shown within each graph column. Tracks were

obtained from movies of 3 embryos each for control and *mac>DfosDN* for pre gb entry in **A**,

4 each for gb entry in **A**, 3 each for the vnc in **A**, 4 each of control and 4 *Dfos²* embryos for

head and pre-gb in **C**, and 3 embryos each for gb entry in **C**. Scale bars: 10 µm.

S3 Fig. Dfos regulates macrophage germband invasion through actin cytoskeleton

associated proteins

(A-C) Comparative mRNA expression levels as determined from RNA sequencing analysis

of FACS sorted wild type macrophages and those expressing DfosDN, n=3 biological

replicates. **(A-B)** Genes down-regulated in macrophages expressing DfosDN are shown,

separated into those with **(A)** strong and **(B)** moderate expression in wild type macrophages.

(C) Expression levels of *Drosophila* formin family genes are unchanged. Fold enrichment is

normalized. p-values: *Dhc36C* 0.02, *CG14204* 0.03, *CG42402* 0.04, *CR43767* 0.046, *TM4SF*

0.03, *CG42260* 0.0011, *cher* 0.046, *GstT4* 0.018, *Xrp1* 0.0011, *Tspo* 0.046, CG31337 0.046.

Frl, *DAAM*, *dia*, *capu* all >0.99.

(D-E) Quantification of the macrophage numbers in **(D)** the pre-gb zone and **(E)** along the

vnc from embryos expressing RNAi against *cher* (*KK 107451*), or *TM4SF* in macrophages

(*KK 102206*) driven by *srpHemo-Gal4* shows no significant alteration. The number in the

column in **(D)** corresponds to the number of embryos analyzed. Control vs. *cher RNAi*

p=0.33. Control vs. *TM4SF RNAi* p=0.05. Control vs. *cher/TM4SF RNAi* p=0.67. (D) SD: 20,

20, 19, 13.

For **(E)** n=13 embryos for control and n=15 for each *cher RNAi* and *TM4SF RNAi*. Control

vs. *cher RNAi* p=0.97 for T1, p=0.33 for T2, p=0.88 for T3. Control vs. *TM4SF RNAi* p=0.52

for T1, p=0.76 for T2, p=0.35 for T3. SD: ctrl 6.5, 5.4, 0.6; *cher RNAi* 5.0, 3.3, 0.8; *TM4SF*

RNAi 4.4, 4.9, 1.9.

(F-I) q-PCR analysis of mRNA extracted from the bones of mice that are wild type,

transgenic (tg) for MHC c-fos, viral 3'UTR, and those in which c-fos transgenesis has led to

an osteosarcoma (OS). Analysis of mRNA expression shows that **(F)** higher *Fos* levels in

osteosarcomas correlate with hgher levels of (**G**) the glutathione S transferase *Gstt3*, and **(H)**

the slit receptor *Eva1c*. but not **(I)** *Tspo*. Bone and osteosarcoma RNA isolated from the

same transgenic mouse, n=4 mice per group, age 5 to 6 months. p-values= 0.86, 0.0028,

0.0013 in **(F)**, 0.79, 0.0001, 0.0003 in **(G)**, 1.0, 0.054, 0.049 in **(H)**, 0.37, 0.33, 0.040 in **(I)**.

SD: 0.7, 0.6, 2.6 in **(F)**; 0.2, 0.3, 1.1 in **(G)**; 0.4, 0.2, 1.5 in **(H)**; 0.1, 0.2, 0.2 in **(I)**.

Histograms show mean + SEM ***p<0.005, **p<0.01, *p<0.05.

Unpaired t-test or one-way ANOVA with Tukey post hoc were used for statistics of

quantifications. Significance is based on adjusted p values.

S4 Fig. Dia does not affect macrophage numbers in the pre-gb zone and along the vnc

(**A**) Three Western blots probed with an mCherry antibody of St 11 embryo extracts from

srpHemo-moe::3xmCherry expressing either *CD8::GFP* (ctrl) or *DfosDN* in macrophages.

Left Western blot also contains *w-* lane. **A'**) Quantitation of the Western blots. We observed

no significant change in the expression of the Moe protein reporter when Dfos function is

inhibited.

(**B**) Expressing Dia-CA in macrophages in $Dfos^1$ or $Dfos^2$ embryos completely rescued the

macrophage germband invasion defect. p-values: Control vs. $Dfos^1$ or vs. $Dfos^2$ p=0.0004 or

p=0.0055 respectively; Control vs. $Dfos^1$ *mac>DiaCA* or vs. $Dfos^2$ *mac>DiaCA* p >0.999;

$Dfos^1$ vs. $Dfos^1$ *mac>DiaCA* p=0.0005; $Dfos^2$ vs. $Dfos^2$ *mac>DiaCA* p=0.035. SD: 20, 23,

18, 19, 7.8.

(**C-D**) There was no significant change in the number of macrophages in (**C**) the pre-gb zone

or (**D**) along the vnc in embryos expressing either of two different RNAi lines against *dia*

expressed in macrophages. Pre-gb: Control vs. *dia RNAi^1* p=0.54, Control vs. *dia RNAi^2*

p=0.77. vnc: Control vs. *dia RNAi^1* p=0.99, Control vs. *dia RNAi^2* p=0.95. *RNAi^1* =TRiP

HMS05027, *RNAi^2* = TRiP HMS00308. (C) SD: 9, 12, 13. (D) SD: Ctrl 5.2, 6.4, 2.5, 0.4; *dia*

RNAi^1 5.6, 6.8, 1.7, 0.2; *dia RNAi^2* 5.1, 4.9, 2.1, 0.6.

(**E-F**) Two further examples of line profiles used for the determination of the membrane-to-

cytoplasmic ratios in Fig 4N, P. Line intensity profiles of (E) DiaRBD::GFP or (F) Dia::GFP

(green) and membrane myr::Tomato (magenta) across the edge of macrophages expressing

either lacZ (Control), RhoDN, DfosDN, *cher RNAi*, or *TM4SF RNAi* as shown in the

schematic in E. Line length ~ 8 µm. Blue lines indicate mean GFP intensity on the membrane

and in cytoplasm.

Histograms show mean + SEM ***p<0.005, **p<0.01, *p<0.05. One way ANOVA with

Tukey post hoc was used for statistics of quantification. The number in each column

corresponds to the number of analysed embryos. "mac>" indicates *srpHemo-GAL4* driver

expressing *UAS* constructs in macrophages. Macrophages are labeled using *srpHemo-*

H2A::3xmCherry.

S5 Fig. Dfos controls cell shape in macrophages

(A) Representative image showing actin protrusions of the first macrophage entering the

germband in the control and in lines expressing DfosDN in macrophages. Actin was

visualized by *srpHemo-Gal4* ("mac>") driving *UAS-LifeActGFP*. White stars indicate the tip

of each actin protrusion. Scale bar 5μm.

(B) Microtubules are labeled with *srpHemo-Gal4* driving *UAS-CLIP::GFP*. Spatially

matched stills of the first macrophage expressing DfosDN and control extending protrusions

into the gb slightly before entering with the body of the cell. As DfosDN macrophages have a

delay in entry, the stills from the DfosDN movie are from a later developmental time point

than the control.

(C) Quantification of macrophage maximum length and maximum width shows that DfosDN

expressing macrophages are 23% longer and 12% thinner than wild type macrophages inside

the gb (indicated in schematic above by dashed box). Control vs. *DfosDN* maximum length

p=0.0005, SD: 3.4, 5.7; control vs. *DfosDN* maximum width p=0.0025, SD: 1.3, 1.0.

(D) Quantification of the maximum length and maximum width of macrophages in the pre-gb

zone (indicated in schematic by dashed box) shows that macrophages expressing DfosDN are

9% shorter and 9% thinner than wild type macrophages. Control vs. *DfosDN* maximum

length p=0.0095, SD: 2.2, 2.0; control vs. *DfosDN* maximum width p=0.005, SD: 2.3, 1.9.

(**E**) Overexpression of *UAS-Lam* in macrophages through *srpHemo-Gal4* (*mac>*) causes no

change in their number in the gb compared to the control. p=0.65, SD: 15, 18.

Histograms show mean + SEM ***p<0.005, **p<0.01, *p<0.05. Unpaired t-test was used for

statistics of quantification. The number of measurements per genotype is shown in each

columns

S6 Fig: Model of protein interactions at the macrophage cortex

Proposed interactions of proteins at the cell cortex in wildtype macrophages during gb

infiltration as shown in Fig 6. Direct binding between two proteins is indicated by a line,

signalling between the interaction partners is represented as an arrow. These interactions and

the resulting model in Figure 6 are based on the papers at the end of this legend next to the

corresponding number shown for each linkage. The Tetraspanin TM4SF can cluster adhesion

receptors such as Integrins at the membrane and lead to the recruitment and activation of Rho

GTPases. Rho GTPases can bind and activate the formin Dia leading to F-actin

polymerization. In addition, Integrin can bind filamins (Cher), which can bind to and thereby

recruit RhoGEF to the membrane. Rho GEFs can in turn bind to and activate Rho GTPases.

References for listed interactions: **1, Tetraspanins-Integrin**) Berditchevski & Odintsova,

1999; Termini & Gillette, 2017. **2, Tetraspanins-Rho GTPases**) Shigeta et al., 2003; Sit &

Manser, 2011; Hong et al., 2012; Tejera et al., 2013. **3, Tetraspanins-Filamins**) Brzozowski

et al., 2018; Perez-Hernandez et al., 2013. **4, Integrin-Filamins**) Razinia et al., 2012;

Ehrlicher et al., 2011. **5, Rho GTPases-Formins**) Ku□hn & Geyer, 2014; Rose et al., 2005.

6 & 8, Rho GEF-Rho GTPases) Vetter & Wittinghofer, 2001 **7, Formins-Filamins**) Hu et

al., 2014; Lian et al., 2016. **9, Filamins-RhoGEFs**) Bellanger et al., 2000; Min et al., 2002.

1541

S1 Movie. Dfos facilitates macrophage motility during initial invasion into the

germband tissue

Movies corresponding to stills shown in **Fig 2A**. Macrophages (green) labelled using

srpHemo-H2A::3xmCherry are imaged while entering the gb in control embryos (left) and

embryos in which macrophages express a dominant negative (DN) version of Dfos (DfosDN)

(right). Time in minutes is indicated in the upper right corner. Scale bar: 10 µm.

S2 Movie. Dfos does not affect macrophage migration along the vnc

Movies corresponding to stills shown in **Fig 2G**. Macrophages (green) labelled by *srpHemo-*

Gal4 driving *UAS-GFP::nls* are imaged during their migration along the segments of the vnc

in control embryos (left) and embryos in which DfosDN is expressed in macrophages (right).

Time in minutes is indicated in the upper right corner. Scale bar: 10 µm.

S3 Movie. Macrophages in *Dfos²* mutants invade germband more slowly

Movies corresponding to stills shown in **S2B Fig**. Macrophages (green) labelled by *srpHemo-*

Gal4 driving *UAS-GFP::nls* are imaged while entering the gb in control embryos (left) and

Dfos² mutant embryos (right). Time is indicated in minutes. Scale bar: 10 µm.

S4 Movie. DfosDN expressing macrophages make long actin protrusions during

germband entry

Movies corresponding to stills shown in **S5A Fig**. F-actin in macrophages (green) labelled

with *srpHemo-Gal4* driving *UAS-LifeAct::GFP* is imaged during gb entry in control embryos

(left) and embryos with macrophages expressing DfosDN (right). Note the extended

protrusion of the DfosDN expressing macrophages. Time is indicated in minutes. Scale bar:

10 µm.

S5 Movie. Dfos controls cell shape in macrophages

Movies corresponding to stills shown in **Fig 5B-C** and **S5B Fig**. Microtubules of

macrophages are labelled with *srpHemo-Gal4* driving *UAS-CLIP::GFP*. They are imaged

1567 during gb entry in control embryos (left) and embryos with macrophages expressing DfosDN

1568 (right). Note the extended shape of the DfosDN expressing macrophages. Time is indicated in

1569 minutes. Scale bar: 10 µm.

Stage 11 A

head germband

early Stage 12 A'

head germband

B

B'

macrophages   yolk sac   ectoderm   mesoderm

C

FosB
c-Fos
Dfos

48% Id

FosB KRRVRRERNKLAAAKCRNRRRELTDRLQAETDQLEEEKAAELESEIAELQKEKERLEFVLVAH
c-Fos KRRIRRERNKMAAAKCRNRRRELTDTLQAETDQLEDEKSALQTEIANLLKEKEKLEFILATH
Dfos KRAVRRERNKQAAARCKRRVDQTNELTEEVEQLEKRGESMRKEIEVLTNSKNQLEYLLAAH

bZIP domain

D **Dfos in situ**

St 11   St 13

control, St 11 E E'

control, early St 12 F F'

Dfos¹, early St 12 H H'

macrophages   Dfos Ab   DAPI-nucleus

ctrl, early St 12 G G' G'' G'''

I **control 1**

J **Dfos¹**

O **control 2**

K **Dfos²**

L **Dfos²; mac>Dfos**

P **mac>DfosDN**

macrophages

M

St 12

N

macrophage #
in germband

ctrl 1   Dfos¹   Dfos²   Dfos²;
              mac>
              Dfos

34   12   23   22

Q

macrophage #
in germband

ctrl 2   mac>
         DfosDN

19   18

R

macrophage #
in germband

ctrl   mac>   mac>
       Dfos   Dfos
       RNAi¹ RNAi²

29   27   28

A Macrophage nuclei (H2A::3xmCherry)

control

0'   30'   90'

mac>DfosDN

0'   30'   90'

B Macrophage invasion into germband (gb)

pre-gb -> gb entry

C speed, µm/min   pre-gb   ns
130   118

D time, min   gb crossing   *
7   7

E speed, µm/min   gb entry   ***
28   24

F speed, µm/min   vnc   ns
105   117

control
mac>DfosDN

G

ctrl   0'   ctrl   30'

mac>DfosDN   0'   *mac>DfosDN*   30'

Macrophage nuclei (H2A::3xmCherry)

A
Early stage 12

control   mac>DfosDN

↓   ↓
FACS
macrophages
↓   ↓
RNA-Seq
gene expression

B

Gene	Possible Cellular Function	Mouse Ortholog Protein	Identity
cher	actin crosslinking	Filamin A	53%
CG42402	homophilic adhesion	Protein eva-1 homolog	35%
TM4SF	cell membrane organisation	Tetraspanin-6	24%
Xrp1	adhesion down-regulation	Epiglycan	22%
Dhc36C	cargo transport	Dynein axonemal, heavy chain 7C	57%
CG42260	ion transport	Cyclic nucleotide-gated olfact. channel	50%
Tspo	mitochondrial transport	Translocator protein	46%
GstT4	detoxification (glutathione transfer)	Glutathione S-transferase theta-3	39%
CG14204	acetyl-CoA metabolism	O-acetyltransferase like protein	27%
CG31337	proteolysis	no ortholog	NA

C *ctrl*

D *mac>TM4SF RNAi*

E *mac>cher RNAi*

F *mac>TM4SF RNAi,cher RNAi*

G
macrophage # in germband

ctrl: 28
mac>cher RNAi: 23
mac>TM4SF RNAi: 29
mac>TM4SF RNAi, cher RNAi: 27

*** ** **** ns *

H *ctrl*

I *mac>DfosDN*

J *mac>DfosDN,cher*

K *mac>DfosDN,TM4SF*

L
macrophage # in germband

ctrl: 38
mac>DfosDN: 19
mac>cher: 28
mac>TM4SF: 27
mac>DfosDN, cher: 34
mac>DfosDN, TM4SF: 29

* ns ns ns ns ns ****

M *Fos*
mRNA fold change
wt bone / Fos tg bone / Fos tg OS
** ** ns

N *FlnA*
wt bone / Fos tg bone / Fos tg OS
** ** ns

N' *FlnB*
wt bone / Fos tg bone / Fos tg OS
**** **** ns

N" *FlnC*
wt bone / Fos tg bone / Fos tg OS
** ** ns

O *Tspan6*
wt bone / Fos tg bone / Fos tg OS
** ** ns

A

Actin protrusion length, μm (in *mac>Lifeact::GFP*)

ctrl   *mac>DfosDN*

B   germband entry

mac>CLIP::GFP

control   *mac>DfosDN*

C   pre-germband

mac>CLIP::GFP

control   *mac>DfosDN*

D   control   *mac>DfosDN*

Probability density

Max L/MaxW germband entry

Probability density

Max L/MaxW pre-germband

E   ctrl   >GFP   E'   >Lam RNAi¹   E''   >Lam RNAi²   E'''   >LamC RNAi

F   *mac>DfosDN*   >GFP   F'   >Lam RNAi¹   F''   >Lam RNAi²   F'''   >LamC RNAi

G   ctrl *(mac> Lam RNAi)*   *mac> DfosDN*   *mac> DfosDN Lam RNAi*

macrophage # in germband

ctrl   *mac> DfosDN*   *mac> Lam RNAi¹*   *mac> Lam RNAi²*   *mac> LamC RNAi*   *mac> DfosDN Lam RNAi¹*   *mac> DfosDN Lam RNAi²*   *mac> DfosDN LamC RNAi*

H   ectoderm   macrophages

macrophage # in germband

ctrl   *mac<> DfosDN*   *mac<> DfosDN ecto> Rho1DN*   *ecto> Rho1DN*

wildtype	*mac>DfosDN*

ecto — ectoderm
mac — macrophage
meso — mesoderm

mac macrophage
ecto ectoderm
meso mesoderm

— cortical actin
— Lamin

TM4SF
Integrin
Cher/ Filamin
Rho GEF
Rho1
Dia/ Formin
F-actin

Dfos → ↑*Cher*
↑*TM4SF* → *Dia* → cortical
actin
assembly

DfosDN ⟶ ↓*Cher*
↓*TM4SF*