Adaptive fitness in *APC*-mutant cells requires chromosome rearrangements

Authors

Yoshihiro Kawasaki[1,2],Tomoko Hamaji[1], Koji Owada[1], Akiko Hayashi[1], Yuping Wu[1],
Taisaku Nogi[1], Miwa Okada[1,†], Shoko Sakai[1], Naoko Tokushige[1], Shota Sasagawa[3],
Hidewaki Nakagawa[3], Yuta Kouyama[4, ††], Atsushi Niida[5], Koshi Mimori[4], Toshihiko
Kuroda[6], Takao Senda[7], Miho Ohsugi[8], Katsumi Fumoto[9], Akira Kikuchi[10], Per O.
Widlund[11], Kazuyuki Kiyosue[12], Norio Yamashita[13], Masahiko Morita[13], Hideo Yokota[13],
Satya N. V. Arjunan[14], Wei-Xiang Chew[14,15], Koichi Takahashi[14], Wesley R. Legant[16,†††],
Bi-Chang Chen[16,††††], Eric Betzig[16,†††††], Ron Smits[17], Riccardo Fodde[18], Hiroko Oshima[19],
Masanobu Oshima[19], M. Mark Taketo[20], Tetsu Akiyama[2], Yuko Mimori-Kiyosue[1,21,*]

Affiliations

[1]Laboratory for Molecular and Cellular Dynamics, RIKEN Center for Biosystems
Dynamics Research, Kobe 650-0047, Japan

[2]Laboratory of Molecular and Genetic Information, Institute of Molecular and Cellular
Biosciences, The University of Tokyo, Tokyo 113-0032, Japan

[3]Laboratory for Cancer Genomics, RIKEN Center for Integrative Medical Sciences,
Yokohama 230-0045, Japan

[4]Department of Surgery, Kyushu University Beppu Hospital, Beppu 874-0838, Japan

[5]Division of Health Medical Computational Science, Health Intelligence Center, Institute of
Medical Science, The University of Tokyo, Tokyo 108-8639, Japan

[6]Nico-tama Coloproctology Clinic, Tokyo 158-0094, Japan

[7]Department of Anatomy, Graduate School of Medicine, Gifu University, Gifu 501-1194,
Japan

[8]Department of Life Sciences, Graduate School of Arts and Sciences, The University of

Tokyo, Tokyo 153-8902, Japan

[9]Department of Molecular Biology and Biochemistry, Graduate School of Medicine, Osaka

University, Osaka 565-0871, Japan

[10]Center for Infectious Disease Education and Research, Osaka University, Osaka 565-

0871, Japan

[11]Department of Microbiology and Immunology, University of Gothenburg, Gothenburg

405 30, Sweden

[12]Biomedical Research Institute, National Institute of Advanced Industrial Science and

Technology, Osaka 563-0026, Japan

[13]Image Processing Research Team, Center for Advanced Photonics, RIKEN, Saitama 351-

0198, Japan

[14]Laboratory for Biologically Inspired Computing, RIKEN Center for Biosystems

Dynamics Research, Osaka 565-0874, Japan

[15]Department of Physics, Faculty of Science, University of Malaya, 50603 Kuala Lumpur,

Malaysia

[16]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA 20147, USA

[17]Department of Gastroenterology and Hepatology, Erasmus MC Cancer Institute,

University Medical Center Rotterdam, 3015 CN Rotterdam, The Netherlands

[18]Department of Pathology, Erasmus MC Cancer Institute, Erasmus University Medical

Center, 3000 CA Rotterdam, The Netherlands

[19]Division of Genetics, Cancer Research Institute, Kanazawa University, Ishikawa 920-

1192, Japan

[20]Colon Cancer Project, Kyoto University Hospital-iACT, Graduate School of Medicine,

Kyoto University, Kyoto 606-8501, Japan

[21]Department of Molecular Genetics, Institute of Biomedical Science, Kansai Medical

University, Osaka 573-1010, Japan

†Present address: Department of Immunology, Laboratory of Immunology, Institute of

Biomedical Research and Innovation (IBRI), 6-3-7, Minatojima-minamimachi, Chuo-ku,

Kobe 650-0047 Japan

††Present address: Showa University Northern Yokohama Hospital, Digestive Disease

Center, Kanagawa 224-8503, Japan

†††Present address: Joint Department of Biomedical Engineering, UNC, Chapel Hill, NC

27599, USA, and NCSU Raleigh, NC 27695, USA; Department of Pharmacology,

University of North Carolina, Chapel Hill, NC, USA

††††Present address: Research Center for Applied Sciences, Academia Sinica, Taipei 115-

29, Taiwan

†††††Present address: Departments of Physics and Molecular and Cell Biology, University

of California, Berkeley, CA 94720, USA; Howard Hughes Medical Institute, Berkeley, CA

94720, USA

*Correspondence to: Yuko Mimori-Kiyosue, Ph.D., Department of Molecular Genetics

Institute of Biomedical Science,

Kansai Medical University

2-5-1 Shinmachi, Hirakata, Osaka 573-1010, Japan

Tel.: +81-72-804-2420

Fax: +81-72-804-2429

E-mail: kiyosuey@hirakata.kmu.ac.jp

Abstract

Cancer cells tolerate copy number alterations (CNAs) of genomic regions that are lethal to non-cancer cells. Certain CNAs are preferentially associated with specific cancer types and lineages, but the mechanisms underlying the emergence and selection of specific CNAs remain unclear. *Adenomatous polyposis coli* (*APC*) mutations induce mitotic errors, but their impact on tumor evolution remains elusive. We investigated APC function in cultured cells and tumors and found that its loss led to β-catenin accumulation at centrosomes, suppressing its maturation through inhibition of key centrosome regulators, including Aurora kinase A (AURKA) that promotes tumor growth. These defects collectively reduced cellular fitness, leading to impaired mitotic fidelity and delayed cell cycle progression. However, in *APC*-mutant tumors, AURKA activity was maintained, at least in part, through the amplification of chromosomes harboring *AURKA* and its activator genes, yet this alone was insufficient to fully restore proliferation: aberrant chromosomal reorganization also emerged and contributed to the adaptive fitness of *APC*-mutant cells. Such a process of adaptive CNA selection provides a framework for understanding how specific CNAs are selected to counteract disadvantages imposed by genetic alterations during tumor progression, providing one key insight into how specific CNAs are selected in this context.

Introduction

Chromosome instability (CIN) induces aneuploidy, a hallmark of cancer (1). However, the extent to which aneuploidy itself contributes to tumor initiation and progression remains unclear. Advances in genome characterization technologies have revealed the genomic landscape of cancer, identifying recurrent copy number alterations (CNAs), which include whole chromosome-, arm-level, and focal CNAs (2,3). Certain CNAs are preferentially associated with specific cancer types and lineages (2,4-6), raising critical questions regarding the mechanisms underlying the selection of these tumor-specific CNAs and their functional consequences. While chromosome gains and losses typically reduce cellular fitness, cancer cells can tolerate such alterations, although the mechanisms enabling this adaptation remain largely unknown (7-9).

The tumor suppressor adenomatous polyposis coli (APC) is well recognized for its role in the WNT/β-catenin pathway as a key component of the β-catenin destruction complex. Its primary function is to regulate β-catenin levels by promoting its degradation (10-12). Loss of APC function leads to aberrant nuclear accumulation of β-catenin, hyperactivation of Wnt signaling, and uncontrolled proliferation, thereby driving tumorigenesis (10-12).

Beyond its role in Wnt signaling, APC is also a critical regulator of mitosis, and its dysfunction induces CIN (13-16). APC inactivation in various cell lines and mouse embryonic stem (ES) cells often results in chromosome mis-segregation due to premature mitotic exit (13-15,17). The CIN phenotype of *APC*-mutant cells is characterized by abnormal chromosome numbers and inhibition of apoptosis in both mouse and human

tumors (18,19). However, the molecular mechanisms underlying APC-driven CIN remain

controversial due to the complexity of its interactions with multiple mitotic regulators (20).

Moreover, it is largely unknown whether and how CIN resulting from APC mutations

influences tumor progression, potentially through the selection of specific CNAs.

Here, through a comparative analysis of APC function in cultured cells and tumors,

we elucidated a functional relationship between APC deficiency and tumor-specific CNAs.

First, we examined the global impact of APC deficiency on mitotic regulation in cultured

cells. Second, we investigated the correlation between *APC* mutations and chromosomal

abnormalities in mouse tumors and human cancers. Finally, we analyzed the process by

which *APC* mutations lead to chromosomal reorganization and subsequent changes in cell

fitness, allowing cells to mitigate the proliferative stress associated with aneuploidy. These

findings provide new perspectives on how CNAs are selected in *APC*-mutant cancers.

Results

APC deficiency impairs centrosome maturation

APC cooperates with various mitotic regulators, including spindle assembly checkpoint

proteins (Bub1/R1) and microtubule end-binding (EB) family proteins, to ensure accurate

chromosome segregation (13-15) (Fig. S1A). Notably, detailed observations of HeLa cells

have shown that APC depletion does not significantly disrupt mitotic spindle formation but

leads to chromosome misalignment due to premature mitotic exit (15). The precise

mechanism underlying this process remains unclear.

To detect subtle phenotypic differences in spindle architecture caused by APC

depletion in HeLa cells, we employed lattice light-sheet microscopy (21) combined with

GFP-fused EB1 (EB1-GFP), a marker of newly assembled microtubule ends (22). This

approach enabled three-dimensional (3D) imaging of entire spindles with high

spatiotemporal resolution (23) (Fig. 1A, B; Fig. S2A, B; and Movie 1–3). Following APC

depletion, microtubule growth probability increased predominantly in the outer equatorial

plane of the spindle, while it significantly decreased around centrosomes (Fig. 1C). These

findings indicate that the primary effect of APC depletion is centrosome immaturity.

Consistent with this finding, γ-tubulin accumulation at centrosomes and the

activation of the central mitotic kinases AURKA (autophosphorylated at T288) and polo-

like kinase 1 (PLK1; phosphorylated at T210), which cooperatively regulate centrosome

maturation (24,25), were significantly reduced in APC-depleted cells (Fig. 2A).

APC together with AXIN1 and β-catenin regulate AURKA activity

To elucidate the molecular mechanism by which APC regulates centrosomes, we screened

centrosomal proteins involved in centrosome maturation for their association with APC.

APC was found to bind AURKA, PCNT (pericentrin), and ch-TOG through its N- and/or

C-terminal regions (Fig. S3). Since AURKA functions upstream of these interactors in

centrosome maturation (26,27), we further examined APC's role in AURKA regulation.

We did not detect direct promotion of AURKA activation by APC in vitro.

However, given that β-catenin and AXIN1, a core scaffold protein in the β-catenin

destruction complex (10), bind centrosomal proteins such as AURKA and PLK1 and

regulate mitosis (28-30), and that Drosophila APC2 and Axin cooperate to promote mitotic

fidelity (31), we investigated whether AXIN1 and β-catenin influence AURKA activation.

In vitro and HEK293T cell-based assays revealed that the middle portion of AXIN1

promoted AURKA activation, similar to TPX2, a well-characterized AURKA activator (32)

(Fig. 2B, C; Fig. S4A, B). Elevated β-catenin levels inhibited AXIN1-mediated AURKA

activation (Fig. 2D), consistent with the competitive interaction of AURKA and β-catenin

with AXIN1 (33). Furthermore, the N-terminal region of APC enhanced AXIN1-mediated

AURKA activation, suggesting that APC facilitates scaffold formation between AXIN1 and

AURKA (Fig. 2E; Fig. S3A, B).

Technical limitations prevented us from expressing a full-length APC fragment

lacking only the C-terminal region. Instead, we utilized mouse embryonic fibroblasts

derived from Apc^{1638T} mice, in which the C-terminal half of APC is deleted (Fig. S1B) (34).

The truncated APC protein in these cells lacks two of the three AXIN-binding domains and

the C-terminal region that interacts with AURKA. These fibroblasts exhibited reduced total

AURKA levels and a lower p/T-AURKA ratio, indicating that the APC C-terminal region

also contributes to AURKA activation (Fig. S1B; Fig. S4C, D).

Consistently, in HeLa cells, AXIN1 depletion and expression of a dominant stable

form of β-catenin (MMβ-catenin), in which multiple GSK phosphorylation sites required

for degradation were mutated (10), reduced the active AURKA ratio (Fig. 2F; Fig. S2C).

Stabilized β-catenin accumulated abnormally around centrosomes along with AXIN1,

suggesting disrupted turnover of these molecules (Fig. 2G; Fig. S2C). These findings

indicate that under normal conditions, APC and AXIN1 function as positive regulators of

AURKA (Fig. 2H). Conversely, APC inactivation impairs AURKA activation, at least

partially through β-catenin stabilization. However, since β-catenin and AXIN1 interact with

multiple centrosomal factors, β-catenin stabilization likely has broader effects on the

dynamics of these interacting factors.

Upregulation of AURKA in *APC*-mutant tumors

The above results seem to contradict the traditional view that AURKA, which promotes

cell cycle progression at centrosomes, is hyperactive in various tumor types and functions

as an oncogene driving tumor growth (27). To examine AURKA activation in *APC*-mutant

tumors, we analyzed $Apc^{\Delta716}$ mice, which carry a heterozygous *Apc* truncation mutation

leading to severe polyposis after loss of the wild-type allele, mimicking early tumor

formation in humans (35) (Fig. S1B). Comparison of multiple polyps and crypts revealed

that, unexpectedly, the active AURKA-to-total AURKA ratio was higher in many polyps

than in crypts (Fig. 3A–C), despite the increase in β-catenin levels (Fig. 3C, E). Detailed

analysis of individual polyps showed that active and total AURKA levels varied between

polyps, exhibiting some degree of intra-tumor heterogeneity (Fig. 3C). Additionally,

regions with exceptionally high AURKA expression were often observed within portions of

the polyps (Fig. 3D). Consistently, in early-stage human *APC*-mutant tumors, increased

AURKA activity was accompanied by enhanced proliferative activity and heterogeneity

(Fig. S5). APC deficiency induces chromosomal instability (CIN) due to a combination of

defects in mitosis and apoptosis (18). Given this, we considered whether AURKA-activated

cells might arise as a consequence of copy number alterations (CNAs) involving AURKA-

activating genes.

Recurrent amplification of *AURKA* loci in human and mouse

To investigate the possibility of *AURKA* amplification, CNAs in human and mouse *APC*-

mutant tumors were examined and compared. To investigate CNAs in *APC*-mutant human

colorectal cancers, we interrogated The Cancer Genome Atlas (TCGA) dataset (3) for *APC*

mutations and identified recurrent gain/loss of certain chromosomes in *APC*-mutant

colorectal cancers (Fig. 4A). The differences in 20q, 13q and 7q gains and 18p, 18q, 14q,

4q, and 17p losses were significant, and there were also differences in 7p and 19q gains and

9q, 20p, and 1p losses in *APC*-mutant colorectal cancers. Although gains/losses of these

chromosomes were also observed in cancers without detectable *APC* mutations, it is

possible that APC expression is suppressed by methylation (36).

We also performed high resolution array-based comparative genomic hybridization

(aCGH) analysis to detect CNAs in individual polyps from $Apc^{\Delta 716}$ mice (Fig. S6). The

structures of human and mouse chromosomes are different; therefore, we examined the

gain or loss regions of chromosomes that overlap between the two, and found that human

chromosome 20 and part of mouse chromosome 2, and human chromosome 13 and part of

mouse chromosome 14 are commonly gained (Fig. 4B and Fig. S6). It is noteworthy that

human chromosome 20q contains *AURKA* and *TPX2* at 20q13.2 and 20q11, respectively,

and chromosome 13q contains *BORA* at 13q21.33, which encodes another AURKA-

activating factor (37). Furthermore, qPCR analysis of the relative DNA content of *Aurka*,

Tpx2, and *Bora* in multiple polyps from *Apc^{Δ716}* mice showed that these genes were

frequently increased in copy number (Fig. 4C). The syntenic comparison of human and

mouse chromosomes, which shows cross-species overlap in the amplified regions, indicates

that these AURKA-related loci were selected by functional relatedness rather than

structural factors.

CNAs induced by CIN in *APC*-mutant tumors

In the TCGA dataset analysis, global CNA scores, which provide an overall assessment of

the CNA burden, were significantly higher in colorectal cancers with *APC* mutations than

in colorectal cancers without *APC* mutations (Fig. 4D). In particular, in *APC*-mutant

colorectal cancers, the frequency of broad CNAs, which are defined as whole-chromosome-

level alterations, was markedly increased compared with focal CNAs, which are alterations

of limited size that range from a few kilobases to part of a chromosome arm (38).

Furthermore, the most frequently altered chromosomes showed a high frequency of the

microsatellite stable phenotype, which is characterized by CIN (Fig. S7A).

It is important to note that the CNA scores for cases with *CTNNB1* (*β-catenin*)

mutations were significantly lower than for cases without *CTNNB1* mutations (Fig. S7B)

and no distinctive recurrent CNA pattern similar to those seen for *APC* mutations was

observed (Fig. S7C), indicating that the effects of *APC* and *CTNNB1* mutations on CNA

formation are distinct.

APC-deficient MCF10A undergo tetraploidisation and maintain chromosome 20 trisomy

To investigate the effect of human chromosome 20 amplification, we used MCF10A, a nearly diploid, noncancerous p53-proficient human mammary epithelial cell line that spontaneously develops chromosome 20 trisomy (39,40) (Fig. S8A and Table S3). We isolated MCF10A subclones with chromosome 20 disomy [MCF10A(20di)] or trisomy [MCF10A(20tri)] (Table S3). Contrary to the expectation that AURKA amplification promotes cell proliferation, MCF10A(20tri) cells exhibited a longer doubling time than MCF10A(20di) cells (Fig. 5A, B and Movie S4). This is consistent with the observation that the gain of an extra chromosome often leads to poor proliferation due to aneuploidy-induced fitness deficits (41). Although AURKA levels at centrosomes were higher in MCF10A(20tri) cells than in MCF10A(20di) cells as expected, the proportion of activated AURKA was lower in MCF10A(20tri) cells, indicating that chromosome-level amplification of AURKA alone is insufficient for enhanced proliferation (Fig. 5C, D). This suggests that additional regulatory mechanisms are required to balance AURKA activity for optimal cell fitness.

Next, we analyzed APC883 cells, a MCF10A-derived cell line carrying a homozygous *APC* mutation leading to a short, truncated APC protein lacking all domains involved in β-catenin regulation (Fig. S1A, B). This cell line is near-diploid but consistently exhibits chromosome 20 trisomy (Fig. S2D, S8B and Table S3). APC883 cells displayed an even longer doubling time than MCF10A(20tri) cells (Fig. 5B, E and Movie S4). Notably, β-catenin accumulated around the centrosome with AURKA (Fig. 5F, G), and

despite high AURKA levels at centrosomes, the proportion of activated AURKA was low

(Fig. 5D). The requirement for AURKA activity in these cells was evident, as moderate

knock-down of AURKA, at a level that did not inhibit MCF10A cell proliferation (Fig.

S2F, Movie S5), led to immediate growth arrest. Similarly, knock-down of truncated APC,

which retains AURKA-activating ability, also resulted in growth arrest (Fig. 5B, Fig. S2E,

F and Movie S4). In contrast, the expression of active AURKA (T288D) or β-catenin

knock-down, both of which increase AURKA activation, partially rescued proliferation and

mitotic fidelity (Fig. 5B, D, E and Fig. S2D, Movie S4). Despite these interventions,

chromosome 20 trisomy was maintained (Fig. S8C, D). These findings indicate that in

APC883 cells, chromosome 20 trisomy alone is insufficient to support proliferation due to

inadequate AURKA activation caused by high β-catenin levels. This results in heightened

sensitivity to AURKA perturbations, necessitating the maintenance of chromosome 20

trisomy. Importantly, the effect of chromosome 20 amplification on proliferation differs

between MCF10A and APC883 cells; chromosome 20 trisomy provides a fitness advantage

only in the context of APC mutation.

Additionally, the expression of stable MMβ-catenin in MCF10A cells did not slow

proliferation (Fig. S2D, F, Movie S5), indicating that functional APC counteracts the

negative effects of β-catenin. This may explain why β-catenin-mutant human colorectal

cancers (CRCs) do not exhibit a CIN phenotype or generate a CNA pattern similar to that

seen with APC mutations (Fig. 4A, Fig. S7C).

Chromosomal alterations in *APC*-mutant cells with extended passage

As APC883 cells were passaged over time, a subpopulation with an increased growth rate

gradually emerged and became dominant in the culture ("transformed APC883," Fig. 5B,

D, E, and Movie S4). These cells exhibited newly acquired karyotypic abnormalities,

including a near-tetraploid chromosome content with recurrent numerical imbalances (Fig.

5H; Fig. S8E and Table S3). Notably, chromosome 20 was present at the highest copy

number (six copies per cell) in transformed APC883 cells (Fig. 5H; Fig. S8E). This

coincided with an increase in total AURKA levels, but the proportion of activated AURKA

remained low compared to MCF10A(20di) cells, indicating that AURKA activation alone

does not drive increased proliferation (Fig. 5D). These findings suggest that an increase in

AURKA levels or activity alone is insufficient to fully restore proliferation. Instead,

maintaining a balanced chromosomal composition is necessary to support the growth of

APC-mutant cells, highlighting the importance of chromosomal alterations in adapting to

fitness impairments. However, despite these chromosomal changes, the proliferation rate of

transformed APC883 cells remained lower than that of MCF10A(20di) cells (Fig. 5B).

Instead of directly enhancing proliferation, these chromosomal changes may serve as an

adaptive mechanism that enables *APC*-mutant cells to survive and continue growing under

selective pressure.

Increased sensitivity of *APC*-mutant tumor cells to AURKA inhibition

Finally, we confirmed the necessity of AURKA activity for tumor growth in *APC*-mutant

tumors by inhibiting AURKA. In *Apc*^{A716} mice treated with the AURKA inhibitor alisertib

(MLN8237), the proportion of Ki67-positive tumor cells, a marker of proliferation, was

reduced, whereas Ki67 levels in crypts remained unchanged (Fig. 6A). These results

demonstrate that *APC*-mutant tumor cells are more dependent on AURKA activity,

consistent with observations in cultured cells. Polyp counting further showed that AURKA

inhibition reduced the total number of polyps (Fig. 6B), indicating that AURKA inhibition

slows tumor growth rather than inducing complete regression. Together with our findings

in MCF10A cells, these results suggest that AURKA activity is a necessary but not

sufficient factor for sustaining proliferation in *APC*-mutant cells, and that additional

chromosomal alterations may play a role in tumor adaptation.

Tetraploid or near-tetraploid cells have been observed in early-stage cancers,

including *APC*-mutant tumors (18,19,42,43), and are thought to serve as a transitional state

during the acquisition of CNAs (43,44). Consistent with this, our data show that near-

tetraploid *APC*-mutant MCF10A cells exhibited improved proliferation compared to their

parental counterparts, whereas amplification of only chromosome 20 in parental MCF10A

cells was associated with reduced fitness. These findings support a model in which

tetraploidization facilitates the emergence of complex CNAs, including those that influence

AURKA activity (Fig. 6C).

Discussion

Cells must maintain a proper chromosomal composition to sustain proliferative fitness, yet

APC-mutant cells take advantage of chromosomal imbalances to acquire adaptive fitness

for tumor formation. APC normally cooperates with AXIN1 to enhance AURKA

activation; however, when APC is dysfunctional, stabilized β-catenin plays a major role in

inhibiting AURKA function. In *APC*-mutant cells, as well as in cultured cells expressing

stable mutant MMβ-catenin, β-catenin accumulated near the centrosome. Phosphorylated β-

catenin is rapidly degraded (10-12); therefore, stable β-catenin is no longer removed from

the centrosome and may persistently block AURKA. Our findings suggest that this

sustained β-catenin accumulation disrupts mitotic fidelity and cell cycle progression, in

contrast to its well-established role in promoting proliferation and differentiation during

normal development. However, in the presence of functional APC, the effects of MMβ-

catenin expression were minimal, indicating that APC also regulates β-catenin through

mechanisms beyond phosphorylation-mediated degradation. This additional level of

regulation may underlie the distinct CNA patterns observed in *APC*-mutant versus *β-*

catenin/CTNNB1-mutant cancers. Further research is required to elucidate APC's role in

pathways beyond WNT/β-catenin signaling.

The mechanisms driving distinct CNA patterns across different cell and cancer

types remain largely unknown. By reanalyzing the TCGA dataset, we identified a strong

correlation between *APC* mutations and specific chromosomal gains and losses,

demonstrating how a single molecular alteration can shape CNA formation. Notably,

syntenic comparisons between mice and humans, despite their differing chromosomal

architectures, revealed shared amplification loci in *APC*-mutant contexts. This suggests that

APC mutations drive CNA selection through functional cellular changes rather than

through structural constraints imposed by chromosome organization. Indeed, our

experimental data indicate that CNA selection in *APC*-mutant cells is linked to the

restoration of AURKA activity. These findings highlight a mechanistic basis for the

correlations observed in TCGA analyses, suggesting that specific CNAs provide a selective

advantage by compensating for the proliferation deficits caused by *APC* loss.

APC mutations negatively impact proliferation, necessitating compensatory

mechanisms for tumor cell survival. Our findings indicate that *APC*-mutant cells experience

strong selection pressure for CNAs that restore AURKA activity, as insufficient AURKA

function is detrimental to their viability. Since AURKA plays a central role in mitotic

fidelity and cell cycle progression, its loss reduces cellular fitness, creating a disadvantage

for *APC*-mutant cells. However, our results also show that increasing AURKA activity

alone is insufficient to fully restore proliferation. Instead, maintaining a balanced

chromosomal composition is required for optimal cell fitness in *APC*-mutant cells.

Although the selection for CNAs that enhance AURKA function was observed in

both our in vitro culture model and in vivo *APC*-mutant tumors, the specific CNA patterns

differed. This discrepancy may be attributed to the fact that MCF10A cells were originally

established with a stable but imbalanced karyotype (39,45), making it unsurprising that

different chromosomal patterns emerged. Nonetheless, even in MCF10A cells,

chromosome 20 was never lost following *APC* mutation, suggesting a strong selective

pressure to retain this chromosome. Importantly, amplification of the *AURKA* locus was

consistently observed across MCF10A cells, mouse tumors, and human cancers, indicating

that in the context of *APC* mutations, *AURKA* amplification represents a common adaptive

strategy to improve cellular fitness, which may subsequently contribute to tumor

progression. MCF10A cells, unlike colorectal cancer cell lines, allow us to isolate the

effects of *APC* mutations on CNA selection without the confounding influences of pre-

existing genomic instability. This enables us to extend beyond intestinal tumorigenesis,

potentially contributing to a broader understanding of how APC loss influences

chromosomal alterations in cancer.

Moreover, although reducing β-catenin can enhance proliferation in vitro, *β-*

catenin/CTNNB1 mutations are rare in *APC*-mutant tumors, likely because β-catenin

remains essential for tumorigenesis (46,47). We observed that AURKA activation not only

improves proliferative capacity but also enables β-catenin to exert its tumorigenic effects,

demonstrating that both factors must act in concert to drive tumor progression. Thus, CNAs

that enhance AURKA function not only mitigate the disadvantages caused by *APC*

mutations but also facilitate β-catenin-dependent tumor evolution.

Our TCGA analysis also revealed significant correlations between *APC* mutations

and chromosomal losses, particularly on chromosomes 18, 14, 4, and 17—alterations

frequently detected in early colorectal cancer (4). These chromosomal deletions may

contribute to tumorigenesis through yet unidentified pathways or by altering gene dosage

and regulatory networks, potentially activating key proliferative factors such as AURKA.

Mechanisms underlying AURKA activation could involve transcriptional upregulation,

RNA stabilization, or other post-transcriptional modifications. Alternatively, these

deletions may be selected in response to the additional selective pressures required for

tumor growth in vivo, such as immune evasion (48,49) and hypoxic adaptation (50). Our

findings demonstrate the necessity of CNAs that enhance AURKA activation in *APC*-

mutant cells, but further investigation is required to elucidate how other selective pressures

contribute to the acquisition of tumor type- or context-dependent CNA patterns in tumor

evolution.

Although CNAs are common in cancers, they typically impose a fitness burden in

normal cells (51). While it has long been unclear why tumor cells tolerate such fitness-

impairing CNAs, our findings reveal that *APC* mutations shape CNA selection by

influencing cellular fitness. We found that APC-mutant cells display distinct aneuploidy

preferences compared with APC-active cells. In APC-active MCF10A cells, chromosome

20 amplification slowed proliferation, whereas APC-mutant MCF10A cells overcame their

proliferative disadvantage by maintaining chromosome 20 while transitioning toward near-

tetraploidy. While the specific CNA patterns observed in MCF10A cells differ from those

in human tumors, MCF10A remains a valuable model for experimentally dissecting the

selective pressures that drive chromosome 20 amplification and polyploidization. Despite

differences in overall CNA patterns, the transition toward near-tetraploidy observed in

APC-mutant MCF10A cells suggests a conserved mechanism by which polyploidization

mitigates the fitness costs of chromosomal instability. Polyploidy, frequently observed in

early-stage cancers (42,52,53), has been proposed to buffer against fitness defects arising

from chromosomal imbalances (8,54). However, polyploid cells are inherently unstable and

often serve as an intermediate state leading to aneuploidy and ultimately tumor progression

(43,44). Our observations in *APC*-mutant cells support the idea that polyploidization

mitigates the fitness costs of chromosomal instability, thereby promoting tumor evolution.

In summary, our findings suggest a functional relationship between *APC* mutations,

tumor-promoting CNAs, and adaptive fitness. *APC* mutations initially reduce AURKA

activity and proliferative potential, but *APC*-mutant cells adapt by exploiting compensatory

CNAs that arise from mitotic errors, recovering AURKA activity and facilitating fitness

adaptation during tumor evolution. This underscores how genetic background influences

CNA formation and enables tumor cells to acquire fitness traits that support their

progression. While AURKA remains a promising therapeutic target, *APC*-mutant cells'

ability to acquire CNAs suggests they may rapidly develop drug resistance. Thus, rather

than targeting AURKA, effective colorectal cancer treatments should focus on preventing

chromosomal rearrangements to constrain tumor adaptation and maintain therapeutic

efficacy.

Materials and Methods

Cell culture and treatment

All cell lines tested negative for mycoplasma contamination. Parental HeLa cell line was

authenticated by Bio-Synthesis Inc (Cell Culture STR Profiling and Comparison Analysis

(Cat. No.:CL1003). Other cell lines used were obtained from reliable distributors. HeLa

cells and HeLa cells expressing EB1-GFP and H2B-TagRFP (clone A1) were maintained as

previously described (55,56). A parental MCF10A cell line and an MCF10A cell line with

the *APC* gene knocked out (MCF10A cells APC$^{-/-}$) were obtained from Sigma-Aldrich-

Aldrich (CLLS1069, St. Louis, MO, USA), and cultured in as described previously (57).

The APC$^{-/-}$ cells, which were generated by zinc finger nuclease (ZFN) technology, are

homozygous for a 17 bp deletion (GCAGCCCAGATTGCCAA) in the endogenous *APC*

gene at 2650-2666 bp of transcript variant 3 (NM_000038). This mutation results in a

truncated 883-amino acid APC protein followed by an unrelated 21 amino acid sequence at

its C-terminus (SHGRSVSHSYLSGRQKFWVYH). In this paper we refer to this cell line

as APC883. The HEK293-based Lenti-X 293T (HEK293T) cell line was obtained from

Takara Bio (Shiga, Japan) and was cultured in DMEM (10566024, Thermo Fisher

Scientific/Gibco, Waltham, MA, USA) supplemented with 10% FCS at 37°C in a 5% CO_2

atmosphere. Colorectal cancer cell lines SW480 (CCL-228), HT-29 (HTB-38), and

HCT116 (CCL-247) were obtained from ATCC (Manassas, VA, USA). SW480 was

cultured in Leibovitz's L-15 Medium (11415064, Thermo Fisher Scientific/Gibco)

supplemented with 10% FCS at 37°C without CO_2. HT-29 and HCT116 were cultured in

McCoy's 5a Medium (16600082, Thermo Fisher Scientific/Gibco) supplemented with 10%

FCS at 37°C in a 5% CO_2 atmosphere. DLD-1 (JCRB9094) was obtained from JCRB Cell

Bank (Osaka, Japan) and cultured in RPMI1640 medium (11875093, Thermo Fisher

Scientific/Gibco) with 10% FCS at 37°C in a 5% CO_2 atmosphere. Mouse embryonic

fibroblasts (MEFs) of wild-type C57BL/6 mice and homozygous Apc^{1638T} mice were

isolated from embryonic day (E) 13.5 embryos, and were cultured in DMEM supplemented

with 10% FCS in a 5% CO_2 atmosphere. The number of cells was counted using an

automated cell counter (Countess; Invitrogen, Carlsbad, CA, USA).

Lattice light sheet microscopy and data analysis

Three-dimensional live imaging by lattice light sheet microscopy and quantitative data

analysis were performed as described previously (21,23). Briefly, timelapse images of

HeLa cells that stably expressed EB1-GFP and H2B-TagRFP (clone A1) (56) were

collected at 0.755-s intervals for EB1-GFP only single channel imaging, 1.550-s intervals

for EB1-GFP and H2B-TagRFP dual channel imaging in fast scanning mode (3D voxel

pitch: $0.100 \times 0.100 \times 0.217$ μm), or 10.000-s intervals for EB1-GFP and H2B-TagRFP

dual channel imaging in high resolution mode (3D voxel pitch: $0.100 \times 0.100 \times 0.195$ μm).

The former data were used for quantitative analyses and the latter data were used to

visualize mitosis progression. The data were displayed in 3D and processed using Imaris

software (Bitplane, Zurich, Switzerland). Supplementary video editing of projected images

exported from Imaris and image processing were performed using ImageJ software,

QuickTime software (Apple. Cupertino, CA, USA), and Adobe Photoshop software

(Adobe, San Jose, CA, USA).

To interpret multidimensional data, images were processed for visualization and

mathematical computation. The XYZ coordinate positions of EB1-GFP comets were

determined automatically. The results are presented as spots or density, or analysed

mathematically using a program developed with MATLAB (MathWorks, Natick, MA,

USA) as described previously (23) and Python using libraries NumPy, Math, and

Matplotlib. An EB1-GFP comet density map (Fig. 1C) was created from the 2D rotational

projected comet numbers and the volume associated with the 2D grid. To create a 2D

rotational projection of all comets, the coordinates of comets and centrosomes were first

transformed so that the spindle axis (vector that connected the two centrosomes) aligned

with the x-axis of the Cartesian coordinate. Then, all comet points were rotated around the

x-axis and projected onto an arbitrary 2D plane with its normal vector orthogonal to the

spindle axis. The result was a half spindle that consisted of comets and centrosomes in 2D

coordinates. Before assembling comet points from all cells, the x-coordinates of 2D comets

were rescaled so that the inter-centrosomal distance was one. Then, a regular 2D grid that

covered all comet spots was created and the number of comets in each grid element was

counted. Average comet density of each element was calculated by normalizing the number

of comets with i) the timelapse duration and ii) calculation of the 3D volume that

corresponded to each grid element. The latter is a function of distance from the spindle axis

and is expressed as $v=\pi h^2 w(2n-1)$ where $n=1,2,3,...,Ny$ with Ny that represents the total

number of grid elements perpendicular to the spindle axis. w and h are the width and height

of the elements, respectively. For cell type comparison, comet density was normalized so

that the summed density of all elements was one. Normalized density was then rescaled by

the global maximum to produce a value between zero and one. Rescaled normalized comet

density of each element and interpolated density values of in-between grid elements were

coded by color and plotted. Because the centrosomes had much higher densities relative to

other regions, we imposed an upper cutoff color value to better illustrate the density

variation of non-centrosomal regions. The density difference graph was constructed by the

difference between the normalized density of each grid element of APC KD and control

cells. A significance test of the density difference was performed by the two-sided t-test to

obtain the p-value. The inputs for the test were the mean and variance value for each grid of

the density map. The output is the p-value for each grid. The code was deposited in Github

(https://github.com/wxchew/cometdensity/blob/master/2DProjectedCometDensity.ipynb).

Antibodies and immunolabeling

No untested antibody was used. Rat anti-E-cadherin mAb (ECCD-2 (58)) was a kind gift

from Dr. M. Takeichi (RIKEN CDB). The following commercial primary antibodies were

used: mouse anti-APC (Ab-1) mAb for western blotting, clone FE9 (OP44; Lot#

D00138185; EMD Millipore/Calbiochem, Billerica, MA, USA); mouse anti-Aurora A

mAb, clone 35C1 (ab13824; Lot# GR121501-2; Abcam, Cambridge, UK); rabbit anti-

Aurora A (phospho T288) polyclonal antibody (pAb), (ab195748, Lot# GR212301-1;

Abcam); rabbit phospho-Aurora A (Thr288)/Aurora B (Thr232)/Aurora C (Thr198)

(D13A11) mAb (#2914; Lot# 3; CST, Danvers, MA, USA); mouse anri-PLK1 mAb, clone

36-298 (ab17057; Lot# GR136555-1; Abcam); rabbit anti-PLK1 (phospho T210) mAb,

clone EPNCIR167 (ab155095; Lot# YJ092901CS; Abcam); rabbit anti-AXIN1 mAb, clone

C76H11 (#2087; Lot# 4; CST); purified human anti-Centromere Protein IgG (CREST) (15-

235; Lot# 441.13BK.20; Antibodies Incorporated, Davis, CA, USA); rabbit anti-β-catenin

pAb (C2206; Lot# 059K4754; Sigma-Aldrich); mouse anti-β-catenin mAb, clone 14

(610154; Lot# 5113978; BD Transduction Lab.); rabbit anti-Ki67 pAb (ab15580; Lot#

GR264768-1; abcam); mouse anti-α-tubulin mAb, clone DM1A (T9026; Lot# 078K4763;

Sigma-Aldrich); rat anti-tubulin mAb, clone YL1/2 (ab6160; Lot# GR57687-1; Abcam);

rabbit anti-γ-tubulin pAb (ab11320; Lot# 628882; Abcam); mouse anti-γ-tubulin mAb,

clone GTU-88 (T6557; Lot# 079K4861; Sigma-Aldrich); mouse anti-GAPDH antibody-

loading control (HRP), clone mAbcam 9484 (ab9482; Lot# GR149747-1; Abcam); rabbit

anti-GFP, HRP-conjugated (598-7; Lot# 002; MBL, Aichi, Japan); mouse anti-GST, clone

B-14 (sc-138; Lot# K1808; Santa Cruz Biotechnology); mouse anti-penta-His antibody

(34660; Lot# 136244018; Qiagen); mouse anti-Myc tag mAb, clone 4A6 (05-724; Lot#

DAM1724025; Millipore); mouse anti-Myc tag mAb, clone 4A6, peroxidase-conjugated

(16-213; Lot# DAM1646292; Millipore); Hoechst 33342 solution (346-07951; Lot#

FN027; Dojin, Japan) was used as a nuclear marker. Immunofluorescent staining of fixed

cultured cells and western blotting were performed as described previously (55-57).

Intestinal tissues from 16-weeks old $Apc^{\Delta716}$ mice and human adenomas were fixed in

formalin and processed for paraffin-embedding and sectioning as described previously (35).

Antigen retrieval for β-catenin (610154, BD), E-cadherin (ECCD-2), AURKA, phosphor-

AURKA(T288) , Ki67 and anti-α-tubulin (YL1/2) immunostaining was performed by

boiling sections for 30 min in Tris-EDTA pH 9.0.

cDNAs, plasmids and RNA interference

The APC cDNA used was described previously (59). Wild-type mouse β-catenin and

dominant-stable β-catenin (MMβ-catenin), in which the NH_2-terminal four Ser/Thr residues

recognized by GSK-3β were substituted for Ala (60) were gifts from Dr. A. Nagafuchi

(Nara Medical University). A human AXIN1 cDNA were PCR cloned from MCF10A cells.

cDNAs for rat AXIN1 were described previously (12,61). The following plasmids were

obtained: pWZL Neo Myr Flag AURKA (addgene plasmid # 20427), a gift from William

Hahn & Jean Zhao; mEmerald-TPX2-N-10 (addgene plasmid #54285) and mEmerald-

Beta-Catenin-20 (mouse) (addgene plasmid #54017), gifts from Michael Davidson.

AURKA(T288D) mutation was introduced by PCR engineering. EGFP-fused APC, β-

catenin, AXIN1 and AURKA were expressed by inserting these cDNAs into pEGFP-C

(Takara/Clontech). Transient transfection of cells with plasmids was performed using

Effectene transfection reagent (Qiagen, Germany).

Lentiviral vectors expressing genes of interest under EF1α promoters (pLVSIN-

EF1a-IRES-pur, pLVSIN-EF1a-IRES-bsr) were described previously (56). Myc-tagged

AURKA(T288D) and MMβ-catenin were generated by PCR engineering and inserted into

pLVSIN-EF1a-IRES series vectors. Lentiviral vectors expressing shRNAs against human

APC under U6 promoters (pLVSIN-U6-pur, bsr) were described previously (62). The target

sequences of shRNAs for human *AURKA was* CAGGACCTGTTAAGGCTACAG. The

lentiviral plasmids for shRNA against β-catenin pLKO.1 puro shRNA beta-catenin was

obtained from addgene (plasmid #18803, a gift from Bob Weinberg).. The target sequences

of siRNA for human AXIN1 were AAGCCAGCCACCAAGAGCTTCATAA (Stealth

RNAi, invitrogen). Transfection of cells with siRNA was performed using HiPerFect

Transfection Reagent (Qiagen).

Lentiviral gene transfer

Lentiviruses that carried genes of interest were generated in HEK293T(Lenti-X) cells

(Takara) using Lentiviral High Titer Packaging Mix (Takara) as described previously (56).

Cells were infected for 24 h with the virus in the presence of 10 mg/ml polybrene (Santa

Cruz Biotechnology), washed, and allowed to recover for 48 h before selection with

appropriate antibiotics. For selection of gene-transduced cells, puromycin (InvivoGen, San

Diego, CA, USA) and blasticidin S (InvivoGen) or bleocin (Wako) were used for selection

because of the *bsr* resistance gene. HeLa cells transfected with cDNA and shRNA were

maintained in medium that contained appropriate concentrations of selection drugs: APC

shRNA/pLVSIN-U6-pur, 10–20 µg/ml puromycin; APC shRNA/pLVSIN-U6-bsr, 100

µg/ml blasticidin S; myc-MMβ-catenin/pLVSIN-EF1a-IRES bsr, 1 µg/ml blasticidin S.

MCF10A cells transfected with cDNA and shRNA were maintained in medium that

contained appropriate concentrations of selection drugs: APC883 + myc-

AURKA(T288D)/pLVSIN-EF1a-IRES bsr, 0.05 μg/ml blasticidin S; APC883 + β-catenin

shRNA, , 1 μg/ml puromycin;. To screen for genes able to stimulate proliferation of

APC883 cells because of rescue of mitotic defects and delayed doubling time, APC883

cells were screened after transfection by continuous culture for more than 1 month without

drug selection, which allowed unconstrained proliferation. For prolonged culture, the

rescued cells were maintained in medium that contained appropriate selection drugs as

described above.

Proteins

For bacterial expression, DNA fragments encoding APC proteins were inserted into pGEX-

5X-1 (GE Healthcare Life Sciences, Pittsburgh, PA, USA). For *in vitro* transcription and

translation reaction, AURKA cDNA was inserted into pcDNA3.1 (Invitrogen) and pT7-

FLAG-1 expression vectors (Sigma-Aldrich) to generate HA-AURKA and FLAG-

AURKA, respectively, and proteins were expressed with a TNT Quick Coupled

Transcription/Translation System (Promega, Madison, WI, USA) according to the

manufacturer's protocol. Purified human AXIN1-MYC/DDK (TP308349) and TPX2-

MYC/DDK (TP305821) proteins were obtained from OriGene Technologies (Rockville,

MD, USA). GST-tagged AURORA A, Active (A28-18G) and Inactive (A28-14G) proteins

were obtained from SignalChem (Richmond, Canada). Purified XMP215-His protein has

been described previously (63).

In-cell fluorescence colocalization assay, immunoprecipitation and in vitro pull-down

assays and cell proliferation assays

For preliminary detection of protein-protein association, target molecules were tagged with

GFP (EGFP or Emerald GFP) or RFP (TagRFP-T or mApple) and overexpressed in

HEK293T cells, and their colocalization analyzed under fluorescence microscopy (in-cell

colocalization assay). For *in vitro* pull-down assays, proteins fused to GST were

synthesized in *E. coli* and isolated by absorption to Glutathione Sepharose (GE Health

Care). GST and GST-fusion proteins (2 μg) immobilized to beads were incubated for 1 h at

4°C with *in vitro* transcription/translation products (HA-AURKA and FLAG-AURKA) or

purified XMAP215-His protein in 500 μl of binding buffer (0.1% NP-40, 50 mM HEPES

(pH 7.0), 150 mM NaCl, 50 mM NaF, 5 mM EDTA, 1 mM DTT, 1 μg/ml aprotinin, 1

μg/ml leupeptin, 50 μg/ml PMSF) and then washed extensively three times with binding

buffer. Proteins adhering to the beads were resolved by sodium dodecyl sulfate

polyacrylamide gel electrophoresis (SDS-PAGE), transferred to a polyvinylidene difluoride

membrane filters (Millipore), and analyzed by immunoblotting. For the pull-down

experiments, HEK293 cells were transfected with FLAG-tagged AXIN1-DIX(600-862) and

extracted in lysis buffer (50 mM Tris HCl pH 7.5, 150 mM NaCl, 5 mM EDTA, 1% Triton

X-100, 2 mM Na3VO4, 10 mM NaF, 1 mM DTT). Cell lysates (500 μl) were incubated

with 2 μg GST or GST-APC-221-880 bound to Glutathione Sepharose beads for 1 h at 4°C.

The beads were then washed three times with wash buffer (20 mM Tris HCl pH 8.0, 150

mM NaCl, 0.5% NP-40) and boiled in SDS sample loading buffer. Protein complexes were

separated by SDS-PAGE and detected by immunoblotting.

In vitro kinase assay

AURKA autophosphorylation was evaluated using the ADP-Glo Kinase Assay (Promega)

according to the manufacturer's instructions. The ADP-Glo Kinase Assay is a luminescent

kinase assay that measures ADP formed via a kinase reaction. Briefly, inactive AURKA

(0.3 pmol; SignalChem) was mixed with various concentrations (0.5-2 pmol) of purified

GST, Axin1 (OriGene Technologies) or TPX2 (SignalChem) for 10 min at RT. The

mixture was further incubated with 50 µM ATP in Reaction Buffer A (40 mM Tris, pH 7.5,

20 mM $MgCl_2$, 0.1 mg/ml BSA) supplemented with 50 µM dithiothreitol (DTT) in a 25 µl

volume for 30 min at RT. The reaction was terminated by adding 25 µl of ADP-Glo

Reagent. After incubation for 40 min at RT, 50 µL of the Kinase Detection Reagent was

added and incubated for 30 min at RT. The luminescence was measured using a

luminometer (Berthold Technologies, Bad Wildbad, Germany). Data from three

independent experiments were averaged after normalization against the control value

(AURKA only) which was set to 1.0.

Western blotting

Total cell lysates were prepared in Laemmli SDS-sample buffer supplemented with

Protease Inhibitor Cocktail (Nacalai Tesque, Kyoto, Japan) and protein concentrations were

quantified by the DC protein assay (Bio-Rad, CA, USA). Protein concentrations were

normalized if necessary and boiled for 5 min with 2-mercaptoethanol, loaded the same

amount of protein in each lane, separated by SDS-PAGE, and transferred to 0.45 µm

Immobilon-P polyvinylidene fluoride membranes (Millipore). Proteins were visualized

using Amersham ECL Select Western Blotting Detection Reagents (GE Healthcare) or

Chemi-Lumi One Super reagent (Nacalai Tesque) and detected with a chemiluminescence

imaging system Davinch-Chemi (CoreBio, Taoyuan, Taiwan) or Amersham Imager 680

(GE Healthcare). For detection of phosphorylated proteins, cell lysates were immediately

boiled for 5 min with 2-mercaptoethanol without concentration adjustment.

Fluorescence microscopy of immunostained specimens and data analysis

The immunostained specimens were observed using an LSM780 confocal microscope with

a Plan-APOCHROMAT 63×/1.4 NA oil immersion objective, a Plan-APOCHROMAT

40×/1.3NA oil immersion objective, a Plan-APOCHROMAT 10×/0.45 NA objective, four

laser lines (405 nm; Multi-Ar, 458, 488, 514 nm; DPSS, 561 nm; He-Ne, 633 nm), and a

highly-sensitive gallium arsenide phosphide (GaAsP) detector (Carl Zeiss) and an LSM880

confocal microscope with a Plan-APOCHROMAT 63×/1.4 NA oil immersion for SR

objective, a Plan-APOCHROMAT 40×/1.3 NA oil immersion for SR objective, a Plan-

APOCHROMAT 10×/0.45 NA objective, four laser lines (405 nm; Multi-Ar, 458, 488, 514

nm; DPSS, 561 nm; He-Ne, 633 nm), and a GaAsP detector (Carl Zeiss). Detection and

quantitative analysis of phosphorylated proteins were performed using cultured cells and

paraffin-embedded tissues fixed in 4% paraformaldehyde or formalin, respectively. To

compare immunostaining intensities in each cell type cultured on different cover slips, the

cover slips were processed under the same conditions and intensity values were obtained as

relative values against that of control cells, which was set to 1. To compare immunostaining

intensities in different tissue sections, the sections were bound to a single slide glass to be

processed under the same staining conditions. For tumor cells analysis in the intestinal

tissue specimens, tumor cells in each polyp were specified by their epithelial features

shown with E-cadherin or β-catenin immunostaining.

To measure fluorescence on centrosomes, maximum intensity projections of z-stack

images were produced, and centrosomal regions were manually selected to obtain total

fluorescence values in the region of interest using the ImageJ software. The average

fluorescence intensities and phospho-to-total protein ratios were normalized to the control

values, which were set to 1.0. In the analysis of different MCF10A derivatives, data from

different experiments were standardized and normalized using parental MCF10A cells as a

control to compare AURKA total intensity, phospho intensity, and the phospho-to-total

ratio. Finally, the values for MCF10A(20di) cells were normalized to 1.0. For mouse

intestines, average values from crypt cells were used as standards. For β-catenin-labeling in

mouse intestine, values from crypt cells in the neighbourhood of the polyps along the

intestinal tract were used as standards. For human tumors, the average values in polyp#2

were normalized to those of polyp#1 without *APC* mutation. Sample sizes indicate the

numbers of cells (β-catenin)/centrosomes (AURKA) analysed. To define the percentage of

Ki67-positive cells in AURKA inhibitor–treated mouse tumors, paraffin-embedded tissues

fixed in formalin were stained for β-catenin, Ki67, and nuclear DNA (with Hoechst 33342),

imaged on a confocal microscope, and analysed using the MetaMorph software. The Ki67-

and Hoechst 33342–positive areas were determined by intensity thresholding in the crypts

and tumor tissues, and the percentage of Ki67-positive areas were calculated in each image

obtained from multiple crypt/tumor areas. Sample sizes indicate the numbers of images

analysed. To count mitotic cells in mouse intestines, the tissue sections were stained with

Hoechst33342 and imaged using an Imager Z1 fluorescence microscope (Carl Zeiss), and

total and mitotic cells were manually counted based on the shapes of chromosomes.

Live imaging and data analysis

Time-lapse phase-contrast live imaging of cells was performed using an IX81-ZDC

inverted microscope with a 20× objective lens LUCPLFN 20×PH (Olympus) controlled by

MetaMorph software (Molecular Devices, Sunnyvale, CA, USA). The microscope was

installed in a temperature-controlled dark box and was equipped with a motorized XYZ

stage, a ZDC (Z drift compensator) autofocus system, a cooled CCD camera ORCA-AG

(Hamamatsu Photonics, Shizuoka, Japan), and a heat controller (Tokai Hit, Shizuoka,

Japan) and gas mixer (Olympus/Tokken, Chiba, Japan) for cell culture. During imaging

cells were maintained at 37°C in a 5% CO_2 atmosphere. Duration of mitosis and doubling

time were manually measured. Time-lapse images were edited using ImageJ and

QuickTime Pro software (Apple).

Animals and human colorectal adenoma

Animals were used in accordance with governmental and institutional regulations and

protocols were approved by the Animal Care and Use Committee of the RIKEN Kobe

Institute, Kanazawa University and the University of Tokyo. All animals were cared for in

accordance with the ethical standards of the institutions. $Apc^{\Delta716}$ mice and paraffin-

embedded tissue preparation were described previously (64). Apc^{1638T} mice (backcrossed to

C57BL/6) were described previously (34,65). Wild-type C57BL/6J mice were purchased

from Charles River Laboratories Japan Inc. (Kanagawa, Japan). For AURKA inhibitor

treatment, wild-type and $Apc^{\Delta716}$ mice were dosed orally twice daily for 14 consecutive

days for Ki67 staining (two mice in each group) and 28 consecutive days for polyp number

counting (ten mice in each group), with vehicle, 20 mg/kg alisertib (CS-0106, Chemscene

LLC, Monmouth Junction, NJ, USA) as a solution in 10% 2-HP-β-CD (66). For polyp

counts, incised intestinal tracts were briefly stained in 0.1% methylene blue, and counted

under a dissecting microscope and were categorized according to size (<1 , 1–2 and >2

mm).

Human colorectal polyps were excised by polypectomy at the Nico-tama Coloproctology

Clinic with informed consent and the protocol was approved by the Ethical Committee of

the Riken Kobe Institute. Samples were de-identified and analyzed anonymously. A small

portion of the polyp was used for genomic DNA isolation using a DNeasy Blood & Tissue

Kit (Qiagen, Hilden, Germany), and the remaining portion was immediately fixed in

formalin and embedded in paraffin. For genomic sequencing, the last exon of *APC*,

encoding from the armadillo repeat region to the stop codon, was cloned by PCR as a

fragment of 1000~1700 bp using specific primers, inserted into pCR-Blunt II-TOPO

(Thermo Fisher Scientific, Waltham, MA, USA) and analyzed by Sanger sequencing. In

this study, a hyperplasia polyp without APC mutation (polyp#1, Asian female, early 60s)

and a low-grade tubular adenoma with two truncating mutations (c.2698G>T and c4558G

del) (polyp#2, Asian female, late 50s) were used. Images of hematoxylin-eosin (HE)

stained sections of a formalin-fixed, paraffin-embedded Swiss roll of the intestine of an

$Apc^{\Delta716}$ mouse and of human polyps were collected using a SZX12 stereomicroscope with a

DF PLAPO 1.2× PF2 objective (Olympus, Tokyo, Japan) or a Primo Vert Monitor with a

Plan-ACHROMAT 4×/0.10 NA objective (Carl Zeiss, Jena, Germany), respectively.

Mutation and copy number analyses using TCGA data

Exome and SNP array data from The Cancer Genome Atlas (TCGA)-colon

adenocarcinomas (COAD) and rectum adenocarcinomas (READ) paired tumor/normal

samples were prepared as described previously (67). Briefly, mutation calling from the

exome data was performed using the *EBCall* algorithm (68) and absolute copy number

profiles were obtained from the SNP array data using the ASCAT algorithm (69). CNA

scores (focal CNA score, FCS; broad CNA scores, BCS; global CNA scores, GCS) and

arm-level copy number profiles were obtained from the SNP array data using the default

settings (including re-segmentation) of CNApp tools (38). In total, 319 COAD and 127

READ TCGA patient samples were analysed. The 446 patient samples were divided into

four groups as follows. Class 1: Lesions without a truncating mutation and LOH in the *APC*

locus; Class 2: Lesions with one truncating mutation or one LOH in the *APC* locus, which

are expected to have a single *APC* mutation and retain a single wildtype copy of *APC*;

Class 3: Lesions with two or more (two to three as a result) truncating mutations in the *APC*

locus, or one or more (one to three as a result) truncating mutations, and LOH in the *APC*

locus, which are expected to have no wildtype *APC*. *CTNNB1*: patients with mutations in β-

catenin gene (*CTNNB1*). The microsatellite instable status and clinicopathological

parameters of all COAD and READ patient samples were also obtained from TCGA

database. The samples were divided into two subgroups, microsatellite stable (MSS), which

included microsatellite instable (MSI) low and MSI high in accordance with the MSI status,

using TCGAbiolinks (70) and then the presence or absence of an *APC* mutation was

compared between each subgroup. Survival curves and CNA frequencies were analysed as

described in the available code. These data were analysed by Fisher's exact test. More

details are in the available code.

Comparative genomic hybridization (CGH) microarray analysis

CGH microarray analysis was performed by Takara Bio using a SurePrint G3 Mouse CGH

Microarray Kit, 1×1M (Agilent Technologies, CA, USA). Genomic DNA from $APC^{\Delta716}$

mouse normal intestines (control) and polyps were fragmented, amplified, and then labelled

with Cy3 or Cy5 dyes. Dye-swap replication experiments were performed for each polyp

sample vs. control DNA. The hybridised DNAs were applied to the microarray and scanned

on an Agilent SureScan Microarray Scanner. The results were analysed using Agilent

Genomic Workbench 7.0 software (Agilent Technologies. Experimental parameters were

set by considering the mosaicism of mutations in tumor cells and the presence of a

considerable amount of normal cells in each sample. For preprocessing: Feature Filter, ON;

Design Filter, v2; GC correction, ON; Centralization, ON; Intra array, ON. For Aberration

Filter: $Log_2Ratio \geq 0.01$; minimum number of probes in region, 5. For analysis, the AMD-2

algorithm with threshold 6 was used. Array CGH profiles derived from two dye-swap

experiments were plotted together on the same plot. Synteny between human and mouse

chromosomes were analysed on the Ensembl Comparative Genomics/Synteny website.

Ideograms of human chromosomes 13 and 20, and mouse chromosomes 2 and 14 shown in

Fig. 4B were generated by tracing illustrations in Ideogram View on the NCBI Genome

Data Viewer website. Syntenic comparison of human and mouse chromosomes was

performed using the Comparative Genomics/Synteny tool on the Ensembl website.

Genomic DNA isolation and quantitative PCR (qPCR)

Genomic DNA was extracted from mouse tissues using ISOGENOME (Nippon Gene,

Tokyo, Japan) in accordance with the manufacturer's instructions. Five nanograms of

genomic DNA was applied to qPCR using PowerUp SYBR Green Master Mix (Applied

Biosystems, Foster City, CA, USA) and the QuantStudio 5 Real-time PCR System

(Applied Biosystems). The reaction was first incubated at 50°C for 2 min and then at 95°C

for 2 min, followed by 40 cycles at 95°C for 15 sec and 60°C for 1 min. *Zfp438* located on

mouse chromosome 18A1 was used as the internal reference gene for qPCR because our

CGH microarray analysis did not show the presence of genetic alterations in the A1–A2

region of chromosome 18 in intestinal polyps of $Apc^{\Delta716}$ mice. The results were normalized

to the values obtained for *Zfp438* and analysed by the $2^{-\Delta\Delta CT}$ method. The primer sequences

were as follows: m*Aurka* 5′-TGGCTCAAACGGAGAGTTGA-3′ and 5′-

CCTGGAATCCCATACCTGGA-3′; m*Bora*, 5′-TAGGTTGGACGGTCCCTCAC-3′ and

5′-GCAGCACAGACATGGAGAGC-3′; m*Tpx2*, 5′-GCATGCGTTTTGTTCCTCTG-3′

and 5′-GCTGCATGATCCCAATGCTA-3′; and m*Zfp438*, 5′-

GGCTGTTGCTGTGCTGTACC-3′ and 5′-CTCCCTGGGCCATCAGTAAG-3′.

Karyotyping of chromosomes in MCF10A and APC883 cells

The APC833 cell line was generated from the MCF10A cell line, a genetically stable

human mammary gland epithelial cell line (71), by introducing a truncation mutation into

the APC locus by genome editing. The MCF10A cell line was established in 1990 from

human fibrocystic mammary tissue (39). Karyotyping prior to immortalization showed

normal diploidy, but later passages showed near-diploidy with some rearrangements

interpreted as 48,XX,3p-,6p+,8+,9p+,16+, including a balanced reciprocal translocation

t(3;9)(p13;p22) and an unbalanced translocation t(5q;der[9]) (39) In later studies, gain at

chromosome 1q was reported (45,72), as observed in this study. Trisomy 20 is recurrent in

subsequent studies (40,45,72,73),

In this study, Q-banding (20 cells for each cell type) were performed by

Chromosome Science Labo (Hokkaido, Japan). Complete description of karyotyping results

is shown in Table S3. In our early analysis, detailed rearrangement of one chromosome 8

was not determined and was described as "dup8(p?)" In later analysis, the structure of the

rearranged chromosome 8 was analyzed by FISH using specific probes (8p11, 8p23, 8q13,

8q21, 8q24) and the description was updated to

"der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3)". For the transformed, near-tetraploid

APC883 cells, common number imbalances are shown by letter color for karyotype

description: cyan, three chromosome 6s; brown, five chromosome 8s; red, six chromosome

20s (Table S3), and by box color for chromosome images: cyan, three chromosome 6s;

brown, five chromosome 8s; red, six chromosome 20s. To analyze newly generated

chromosome aberrations, types of abnormalities were classified as tetraploidy, gain, loss,

deletion and fusion. The fusion includes all types of chromosomes generated by fusions

such as translocation, robertsonian translocation and telomere association. To separate

subclones MCF10A cells carrying chromosome 20 disomy [MCF10A(20di)] and trisomy

[MCF10A(20tri)], clones were isolated by colony picking followed by Q-banding to

determine karyotype.

Statistics and reproducibility

Statistical evaluation was carried out by the Student's t-test (two sided), Tukey–Kramer

method (two sided), Fisher's exact test, Wilcoxon rank-sum test, or Mann–Whitney U-test

(two sided) as indicated. P-values and sample size are shown in each figure. Biochemical

experiments were repeated with at least three independent preparations with similar results.

For quantification of image data to ensure adequate power to detect significance, sufficient

numbers of regions/cells were analysed as indicated in each figure.

Data and materials availability: Data are available on request from the authors. Materials

used in this study are available for distribution after a materials transfer agreements (MTA),

except for materials from human research participants. The aCGH data have been deposited

in the NCBI Gene Expression Omnibus (GEO; http://www.ncbi.nlm.nih.gov/geo/) and are

accessible through GEO series accession number GSE3642. accession GSE63306

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc= GSE63306). The "Fig1c.zip" file

containes multiple csv files used to generate the plots in Fig. 1C.

Code availability: Computer codes are available on request from the authors. The code to

create the 2D rotational projection of EB1 comet data has been deposited in Github

(https://github.com/wxchew/cometdensity/blob/master/2DProjectedCometDensity.ipynb).

Codes for TCGA data analysis are available in Code S1. This folder also contains the

TCGA data used for the analysis with the codes attached.

References

1.   Boveri T. Concerning the origin of malignant tumours by Theodor Boveri. Translated

and annotated by Henry Harris. J Cell Sci **2008**;121 Suppl 1:1-84

2.   Beroukhim R, Mermel CH, Porter D, Wei G, Raychaudhuri S, Donovan J*, et al.* The

landscape of somatic copy-number alteration across human cancers. Nature

2010;463:899-905

3.   Cancer Genome Atlas N. Comprehensive molecular characterization of human colon

and rectal cancer. Nature **2012**;487:330-7

4.   Gerstung M, Jolly C, Leshchiner I, Dentro SC, Gonzalez S, Rosebrock D*, et al.* The

evolutionary history of 2,658 cancers. Nature **2020**;578:122-8

5.   Zack TI, Schumacher SE, Carter SL, Cherniack AD, Saksena G, Tabak B*, et al.* Pan-

cancer patterns of somatic copy number alteration. Nature genetics **2013**;45:1134-40

6.   Sack LM, Davoli T, Li MZ, Li Y, Xu Q, Naxerova K*, et al.* Profound Tissue

Specificity in Proliferation Control Underlies Cancer Drivers and Aneuploidy

Patterns. Cell **2018**;173:499-514 e23

7.   Sheltzer JM, Amon A. The aneuploidy paradox: costs and benefits of an incorrect

karyotype. Trends in genetics : TIG **2011**;27:446-53

8.   Ben-David U, Amon A. Context is everything: aneuploidy in cancer. Nature reviews

Genetics **2020**;21:44-62

9.   Gordon DJ, Resio B, Pellman D. Causes and consequences of aneuploidy in cancer.

Nature reviews Genetics **2012**;13:189-203

10. Stamos JL, Weis WI. The beta-catenin destruction complex. Cold Spring Harbor

perspectives in biology **2013**;5:a007898

11. Clevers H, Nusse R. Wnt/beta-catenin signaling and disease. Cell **2012**;149:1192-205

12. Ikeda S, Kishida S, Yamamoto H, Murai H, Koyama S, Kikuchi A. Axin, a negative

regulator of the Wnt signaling pathway, forms a complex with GSK-3beta and beta-

catenin and promotes GSK-3beta-dependent phosphorylation of beta-catenin. The

EMBO journal **1998**;17:1371-84

13.   Fodde R, Kuipers J, Rosenberg C, Smits R, Kielman M, Gaspar C*, et al.* Mutations in

the APC tumour suppressor gene cause chromosomal instability. Nature cell biology

2001;3:433-8

14.   Kaplan KB, Burds AA, Swedlow JR, Bekir SS, Sorger PK, Näthke IS. A role for the

Adenomatous Polyposis Coli protein in chromosome segregation. Nature cell biology

2001;3:429-32

15.   Draviam VM, Shapiro I, Aldridge B, Sorger PK. Misorientation and reduced

stretching of aligned sister kinetochores promote chromosome missegregation in

EB1- or APC-depleted cells. The EMBO journal **2006**;25:2814-27

16.   Banks JD, Heald R. Adenomatous polyposis coli associates with the microtubule-

destabilizing protein XMCAK. Current biology : CB **2004**;14:2033-8

17.   Tighe A, Johnson VL, Taylor SS. Truncating APC mutations have dominant effects

on proliferation, spindle checkpoint control, survival and chromosome stability. J

Cell Sci **2004**;117:6339-53

18.   Dikovskaya D, Schiffmann D, Newton IP, Oakley A, Kroboth K, Sansom O*, et al.*

Loss of APC induces polyploidy as a result of a combination of defects in mitosis and

apoptosis. Journal of Cell Biology **2007**;176:183-95

19.   Bezrookove V, Smits R, Moeslein G, Fodde R, Tanke HJ, Raap AK*, et al.* Premature

chromosome condensation revisited: a novel chemical approach permits efficient

cytogenetic analysis of cancers. Genes, chromosomes & cancer **2003**;38:177-86

20. Rusan NM, Peifer M. Original CIN: reviewing roles for APC in chromosome

instability. The Journal of cell biology **2008**;181:719-26

21. Chen BC, Legant WR, Wang K, Shao L, Milkie DE, Davidson MW*, et al.* Lattice

light-sheet microscopy: imaging molecules to embryos at high spatiotemporal

resolution. Science **2014**;346:1257998

22. Mimori-Kiyosue Y, Shiina N, Tsukita S. The dynamic behavior of the APC-binding

protein EB1 on the distal ends of microtubules. Current biology : CB **2000**;10:865-8

23. Yamashita N, Morita M, Legant WR, Chen BC, Betzig E, Yokota H*, et al.* Three-

dimensional tracking of plus-tips by lattice light-sheet microscopy permits the

quantification of microtubule growth trajectories within the mitotic apparatus. Journal

of biomedical optics **2015**;20:101206

24. Marumoto T, Zhang D, Saya H. Aurora-A - a guardian of poles. Nature Reviews

Cancer **2005**;5:42-50

25. Lens SMA, Voest EE, Medema RH. Shared and separate functions of polo-like

kinases and aurora kinases in cancer. Nature Reviews Cancer **2010**;10:825-41

26. Joukov V, De Nicolo A. Aurora-PLK1 cascades as key signaling modules in the

regulation of mitosis. Science signaling **2018**;11

27. Nikonova AS, Astsaturov I, Serebriiskii IG, Dunbrack RL, Jr., Golemis EA. Aurora

A kinase (AURKA) in normal and pathological cell division. Cellular and molecular

life sciences : CMLS **2013**;70:661-87

28. Kim SM, Choi EJ, Song KJ, Kim S, Seo E, Jho EH*, et al.* Axin localizes to mitotic

spindles and centrosomes in mitotic cells. Experimental cell research **2009**;315:943-

54

29. Ruan K, Ye F, Li CY, Liou YC, Lin SC, Lin SY. PLK1 Interacts and Phosphorylates

Axin That Is Essential for Proper Centrosome Formation. PloS one **2012**;7

30. Mbom BC, Nelson WJ, Barth A. beta-catenin at the centrosome: discrete pools of

beta-catenin communicate during mitosis and may co-ordinate centrosome functions

and cell cycle progression. BioEssays : news and reviews in molecular, cellular and

developmental biology **2013**;35:804-9

31. Poulton JS, Mu FW, Roberts DM, Peifer M. APC2 and Axin promote mitotic fidelity

by facilitating centrosome separation and cytoskeletal regulation. Development

2013;140:4226-36

32. Bayliss R, Sardon T, Vernos I, Conti E. Structural basis of Aurora-A activation by

TPX2 at the mitotic spindle. Mol Cell **2003**;12:851-62

33. Xia Z, Wei P, Zhang H, Ding Z, Yang L, Huang Z*, et al.* AURKA governs self-

renewal capacity in glioma-initiating cells via stabilization/activation of beta-

catenin/Wnt signaling. Molecular cancer research : MCR **2013**;11:1101-11

34. Smits R, Kielman MF, Breukel C, Zurcher C, Neufeld K, Jagmohan-Changur S*, et al.*

Apc1638T: a mouse model delineating critical domains of the adenomatous polyposis

coli protein involved in tumorigenesis and development. Genes & development

1999;13:1309-21

35. Oshima M, Oshima H, Kitagawa K, Kobayashi M, Itakura C, Taketo M. Loss of Apc

heterozygosity and abnormal tissue building in nascent intestinal polyps in mice

carrying a truncated Apc gene. Proceedings of the National Academy of Sciences of

the United States of America **1995**;92:4482-6

36. Esteller M, Sparks A, Toyota M, Sanchez-Cespedes M, Capella G, Peinado MA, *et*

al. Analysis of adenomatous polyposis coli promoter hypermethylation in human

cancer. Cancer research **2000**;60:4366-71

37. Hutterer A, Berdnik D, Wirtz-Peitz F, Zigman M, Schleiffer A, Knoblich JA. Mitotic

activation of the kinase Aurora-A requires its binding partner Bora. Developmental

cell **2006**;11:147-57

38. Franch-Expósito S, Bassaganyas L, Vila-Casadesus M, Hernandez-Illan E, Esteban-

Fabro R, Diaz-Gay M*, et al.* CNApp, a tool for the quantification of copy number

alterations and integrative analysis revealing clinical implications. eLife **2020**;9

39. Soule HD, Maloney TM, Wolman SR, Peterson WD, Jr., Brenz R, McGrath CM*, et*

al. Isolation and characterization of a spontaneously immortalized human breast

epithelial cell line, MCF-10. Cancer research **1990**;50:6075-86

40. Zientek-Targosz H, Kunnev D, Hawthorn L, Venkov M, Matsui S, Cheney RT*, et al.*

Transformation of MCF-10A cells by random mutagenesis with frameshift mutagen

ICR191: a model for identifying candidate breast-tumor suppressors. Molecular

cancer **2008**;7:51

41. Sheltzer JM, Ko JH, Replogle JM, Habibe Burgos NC, Chung ES, Meehl CM*, et al.*

Single-chromosome Gains Commonly Function as Tumor Suppressors. Cancer cell

2017;31:240-55

42. Caldwell CM, Green RA, Kaplan KB. APC mutations lead to cytokinetic failures in

vitro and tetraploid genotypes in Min mice. The Journal of cell biology

2007;178:1109-20

43.   Ganem NJ, Storchova Z, Pellman D. Tetraploidy, aneuploidy and cancer. Current

opinion in genetics & development **2007**;17:157-62

44.   Holland AJ, Cleveland DW. Boveri revisited: chromosomal instability, aneuploidy

and tumorigenesis. Nature reviews Molecular cell biology **2009**;10:478-87

45.   Cowell JK, LaDuca J, Rossi MR, Burkhardt T, Nowak NJ, Matsui S. Molecular

characterization of the t(3;9) associated with immortalization in the MCF10A cell

line. Cancer genetics and cytogenetics **2005**;163:23-9

46.   Sparks AB, Morin PJ, Vogelstein B, Kinzler KW. Mutational analysis of the

APC/beta-catenin/Tcf pathway in colorectal cancer. Cancer research **1998**;58:1130-4

47.   Scholer-Dahirel A, Schlabach MR, Loo A, Bagdasarian L, Meyer R, Guo R, *et al.*

Maintenance of adenomatous polyposis coli (APC)-mutant colorectal cancer is

dependent on Wnt/beta-catenin signaling. Proceedings of the National Academy of

Sciences of the United States of America **2011**;108:17135-40

48.   Vinay DS, Ryan EP, Pawelec G, Talib WH, Stagg J, Elkord E*, et al.* Immune evasion

in cancer: Mechanistic basis and therapeutic strategies. Semin Cancer Biol **2015**;35

Suppl:S185-S98

49.   Davoli T, Uno H, Wooten EC, Elledge SJ. Tumor aneuploidy correlates with markers

of immune evasion and with reduced response to immunotherapy. Science **2017**;355

50.   Hockel M, Vaupel P. Tumor hypoxia: definitions and current clinical, biologic, and

molecular aspects. Journal of the National Cancer Institute **2001**;93:266-76

51.   Torres EM, Williams BR, Amon A. Aneuploidy: cells losing their balance. Genetics

968   **2008**;179:737-46

969   52.   Shackney SE, Smith CA, Miller BW, Burholt DR, Murtha K, Giles HR, *et al.* Model

970   for the genetic evolution of human solid tumors. Cancer research **1989**;49:3344-54

971   53.   Olaharski AJ, Sotelo R, Solorza-Luna G, Gonsebatt ME, Guzman P, Mohar A, *et al.*

972   Tetraploidy and chromosomal instability are early events during cervical

973   carcinogenesis. Carcinogenesis **2006**;27:337-43

974   54.   Davoli T, de Lange T. The causes and consequences of polyploidy in normal

975   development and cancer. Annual review of cell and developmental biology

976   **2011**;27:585-610

977   55.   Mimori-Kiyosue Y, Grigoriev I, Lansbergen G, Sasaki H, Matsui C, Severin F, *et al.*

978   CLASP1 and CLASP2 bind to EB1 and regulate microtubule plus-end dynamics at

979   the cell cortex. The Journal of cell biology **2005**;168:141-53

980   56.   Nakamura S, Grigoriev I, Nogi T, Hamaji T, Cassimeris L, Mimori-Kiyosue Y.

981   Dissecting the nanoscale distributions and functions of microtubule-end-binding

982   proteins EB1 and ch-TOG in interphase HeLa cells. PloS one **2012**;7:e51442

983   57.   Hotta A, Kawakatsu T, Nakatani T, Sato T, Matsui C, Sukezane T, *et al.* Laminin-

984   based cell adhesion anchors microtubule plus ends to the epithelial cell basal cortex

985   through LL5alpha/beta. The Journal of cell biology **2010**;189:901-17

986   58.   Shirayoshi Y, Nose A, Iwasaki K, Takeichi M. N-linked oligosaccharides are not

987   involved in the function of a cell-cell binding glycoprotein E-cadherin. Cell Struct

988   Funct **1986**;11:245-52

989   59.   Kawasaki Y, Senda T, Ishidate T, Koyama R, Morishita T, Iwayama Y, *et al.* Asef, a

990   link between the tumor suppressor APC and G-protein signaling. Science

991   **2000**;289:1194-7

992   60.   Takahashi N, Ishihara S, Takada S, Tsukita S, Nagafuchi A. Posttranscriptional

993   regulation of alpha-catenin expression is required for Wnt signaling in L cells.

994   Biochem Bioph Res Co **2000**;277:691-8

995   61.   Yamamoto H, Kishida S, Uochi T, Ikeda S, Koyama S, Asashima M, *et al.* Axil, a

996   member of the Axin family, interacts with both glycogen synthase kinase 3 beta and

997   beta-catenin and inhibits axis formation of Xenopus embryos. Mol Cell Biol

998   **1998**;18:2867-75

999   62.   Pronobis MI, Deuitch N, Posham V, Mimori-Kiyosue Y, Peifer M. Reconstituting

1000   regulation of the canonical Wnt pathway by engineering a minimal beta-catenin

1001   destruction machine. Molecular biology of the cell **2017**;28:41-53

1002   63.   Widlund PO, Stear JH, Pozniakovsky A, Zanic M, Reber S, Brouhard GJ, *et al.*

1003   XMAP215 polymerase activity is built by combining multiple tubulin-binding TOG

1004   domains and a basic lattice-binding region. Proceedings of the National Academy of

1005   Sciences of the United States of America **2011**;108:2741-6

1006   64.   Oshima H, Oshima M, Kobayashi M, Tsutsumi M, Taketo MM. Morphological and

1007   molecular processes of polyp formation in Apc(Delta 716) knockout mice. Cancer

1008   research **1997**;57:1644-9

1009   65.   Yokoyama A, Nomura R, Kurosumi M, Shimomura A, Onouchi T, Iizuka-Kogo A, *et*

1010   *al.* The C-terminal domain of the adenomatous polyposis coli (Apc) protein is

involved in thyroid morphogenesis and function. Medical molecular morphology

2011;44:207-12

66. Sells TB, Chau R, Ecsedy JA, Gershman RE, Hoar K, Huck J*, et al.* MLN8054 and

Alisertib (MLN8237): Discovery of Selective Oral Aurora A Inhibitors. ACS

medicinal chemistry letters **2015**;6:630-4

67. Uchi R, Takahashi Y, Niida A, Shimamura T, Hirata H, Sugimachi K*, et al.*

Integrated Multiregional Analysis Proposing a New Model of Colorectal Cancer

Evolution. PLoS genetics **2016**;12:e1005778

68. Shiraishi Y, Sato Y, Chiba K, Okuno Y, Nagata Y, Yoshida K*, et al.* An empirical

Bayesian framework for somatic mutation detection from cancer genome sequencing

data. Nucleic acids research **2013**;41:e89

69. Van Loo P, Nordgard SH, Lingjaerde OC, Russnes HG, Rye IH, Sun W*, et al.* Allele-

specific copy number analysis of tumors. Proceedings of the National Academy of

Sciences of the United States of America **2010**;107:16910-5

70. Colaprico A, Silva TC, Olsen C, Garofano L, Cava C, Garolini D*, et al.*

TCGAbiolinks: an R/Bioconductor package for integrative analysis of TCGA data.

Nucleic acids research **2016**;44:e71

71. Debnath J, Muthuswamy SK, Brugge JS. Morphogenesis and oncogenesis of MCF-

10A mammary epithelial acini grown in three-dimensional basement membrane

cultures. Methods **2003**;30:256-68

72. Jonsson G, Staaf J, Olsson E, Heidenblad M, Vallon-Christersson J, Osoegawa K*, et*

al. High-resolution genomic profiles of breast cancer cell lines assessed by tiling

BAC array comparative genomic hybridization. Genes, chromosomes & cancer

2007;46:543-58

73.   Marella NV, Malyavantham KS, Wang J, Matsui S, Liang P, Berezney R.

Cytogenetic and cDNA microarray expression analysis of MCF10 human breast

cancer progression cell lines. Cancer research **2009**;69:5946-53

1038

Acknowledgements

We are grateful to Dr. A. Ochiai (NCC-EPOC) for his guidance on tumor analysis. Drs. A.

Nagafuchi (Nara Medical Univ), T. Toda (Hiroshima Univ), M. Toya (Waseda Univ), M.

Takeichi (Riken BDR), C. Furusawa (Univ. of Tokyo), and Y. Takai (Kobe Univ) for

critical reading of the manuscript and valuable suggestions. Our special thanks are also due

to Ms. M. Yamaguchi and Mr. Y. Sato (Carl Zeiss Japan), Mr. M. Yamaguchi (Molecular

Devices Japan), and Mrs. Y. Imai and M. Emi (Olympus) for excellent technical assistance

in operating the microscope systems and image analysis software.

1047

Funding: Y.M.-K. was supported by the Japan Society for the Promotion of Science–

NEXT program (LS128), the Takeda Science Foundation, the Uehara Memorial

Foundation, a Grant-in-Aid for Challenging Research (Pioneering) (JSPS KAKENHI Grant

Number 20K20379), and Japan Science and Technology Agency Core Research for

Evolutional Science and Technology (no. JPMJCR1863), Extramural Collaborative

Research Grant of Cancer Research Institute, Kanazawa University, and an intramural grant

from RIKEN. The results published here are in part based upon data generated by the

TCGA Research Network: http://cancergenome.nih.gov/. Imaging experiments were in part

supported by the RIKEN Kobe Light Microscopy Facility. Animal experiments were in part

supported by the Laboratory for Animal Resources and Genetic Engineering at the RIKEN

BDR.

1059

Author contributions: Y.K., T.H., K.O., A.H. and T.N. performed experiments and edited

the manuscript. Y.W., M.O., S.S., and N.T. performed experiments. Y.K., A.N., and K.M.

analyzed TCGA data using informatics techniques. T.K., T.S., M.O., K.F., A.K., P.O W.

and H.O. prepared biological samples. K.K. performed image data analysis and statistical

data analysis. N.Y., M.M., H.Y., C.W.X., S.A., K.T., W.R.L., B-C.C. and E.B. performed

imaging experiments, image processing and data analysis. R.S. and R.F. prepared

biological samples and edited the manuscript. M.O., M.M.T and T.A gave conceptual and

technical advice. Y.M.-K. conceived and managed the project, performed experiments, and

wrote the manuscript.

1069

1070   **Competing interests:** Portions of the technology described herein are covered by U.S.

1071   Patent 7,894,136 issued to Eric Betzig (EB) and assigned to Lattice Light, LLC of Ashburn,

1072   VA, U.S. Patents 8,711,211 and 9,477,074 issued to EB and assigned to HHMI, U.S. Patent

1073   application 13/844,405 filed by EB and Kai Wang (KW) and assigned to HHMI, and U.S.

1074   Patent 9,500,846 issued to EB and KW and assigned to HHMI.

1075

1076

1077

Fig. 1. Identification of the site affected by APC depletion. (A) Three-dimensional

representation of control (Ctrl) and APC-depleted (APC KD) mitotic HeLa cells expressing

EB1-GFP and H2B-TagRFP as visualized by lattice light sheet microscopy. Original

images (left) and examples of automated detection of EB1-GFP positions and chromosome

surfaces (right) are shown. See also Fig. S2A and Movies S1–S3. (B) Procedure of

quantitative analysis of the timelapse 3D data. The rotational projection method is shown

diagrammatically in the bottom. (C) Rotationally projected density map of control and

APC-depleted cells (left). The centrosome positions are indicated by green dots at positions

0.0 and 1.0 on the x-axis. The number of cells and time points (tps) in time lapse videos

1087   used are indicated in the panel. Density difference map (middle), and p-value map (right)

1088   are shown.

1089

Figure 2 Kawasaki et al.

Fig. 2. Regulation of AURKA by APC, AXIN1 and β-catenin.   (**A**) Immunostaining of

γ-tubulin, α-tubulin, AURKA and autophosphorylated AURKA (p-AURKA) with Hoechst

33342 to stain DNA (left). Arrowheads indicate spindle poles. Thereafter, see Table S1 for

all representative images. Fluorescence intensities of γ-tubulin at centrosomes and ratios of

fluorescence intensities (phosphorylated vs. total protein) of AURKA (T288) and PLK1

(T210) are shown (right). (**B**) In vitro AURKA autophosphorylation at T288 detected by

western blotting. (**C**) In vitro kinase assay of AURKA autophosphorylation. (**D, E**)

HEK293T cells were transfected with GFP-fused AURKA, AXIN1, β-catenin, and APC

fragments (see Fig. S3A) as indicated and then subjected to western blotting using the

indicated antibodies. APC bands are marked with asterisks. Note that the phosphorylated

AURKA band always appeared as multiple bands. AXIN1 and AURKA levels were

increased when coexpressed with binding APC fragments and AXIN1, respectively,

probably due to stabilizing effect. Thereafter, see Table S2 for uncropped versions of blots.

(**F**) AURKA status in AXIN1 depleted- and MMβ-catenin-expressing HeLa cells. (**G**)

Distribution of β-catenin and AXIN1 in APC depleted- and MM-catenin-expressing HeLa

cells. Arrowheads indicate spindle poles. (**H**) APC positively regulate AURKA together

with AXIN1 and β-catenin (up). When APC is dysfunctional, its ability to activate AURKA

reduced and stabilized β-catenin inhibits normal AURKA behavior and its activation

(bottom). Scale bars: 5 μm. * $P < 0.05$;** $P < 0.01$; *** $P < 0.001$ (Student's t-test in **A**,

Tukey–Kramer -test in **B, C, F**).

1111

A $Apc^{\Delta716}$ mouse intestine

B

C

D

E

1112

1113

Fig. 3. Upregulation of AURKA in mouse tumors. (**A**) HE-stained section of the

intestines of a 16-weeks old $Apc^{\Delta716}$ mouse. Polyps analyzed in this study are indicated by

numbers. (**B**) Immunostaining of a formalin-fixed, paraffin-embedded *Apc*^{Δ716} mouse

intestine tissue section for microtubules (MTs), AURKA and p-AURKA with nuclear DNA

stained by Hoechst 33342 to visualize crypt cells (top) and tumor cells (bottom) is shown.

Low magnification images are shown on the left, and magnified images of mitotic cells

indicated with arrowheads are shown in the three panels on the right. (**C**) Quantitative

analysis of immunofluorescence signals for AURKA/p-AURKA and β-catenin in crypts

and tumor tissues of nine polyps. (**D**, **E**) Immunostaining of a formalin-fixed, paraffin-

embedded *Apc*^{Δ716} mouse intestine section for AURKA, E-cadherin and β-catenin with

nuclear DNA stained by Hoechst 33342 to visualize patchy region in tumors

overexpressing AURKA (D) and crypt cells (top) and tumor cells (bottom) (E). In (E), Low

magnification images are shown on the left and magnified images of the boxed areas are

shown in the four panels on the right. Scale bars: 2 mm (A); 100 μm (D); 50 μm (E, low

magnification, left), and 10 μm (E, high magnification, right). *$P < 0.05$; ** $P < 0.01$; ***

$P < 0.001$ (Tukey–Kramer -test) .

1130

1131

Figure 4 Kawasaki et al.

A TCGA COAD / READ dataset (Total case number: 406)

Class 1: *APC* mutation = 0, Class 2: *APC* mutation = 1, Class 3: *APC* mutation ≥ 2

B Chromosome amplification

C

D

1132

Fig. 4. Recurrent CNAs in *APC*-mutant human colorectal cancers and mouse tumors.

(**A**) The Cancer Genome Atlas (TCGA)-colon adenocarcinomas (COAD) and rectum adenocarcinomas (READ) samples are arranged along the x-axis and ordered in accordance with their *APC* mutations groups (Class 1: *APC* mutation = 0; Class 2: *APC* mutation = 1; Class 3: *APC* mutation ≥2). The y-axis numbered with 1–22 denotes chromosome numbers. The intensity of red and blue matches the frequency of copy number gain and loss, respectively. The ratio of gain to loss for each class is displayed as a stacked graph with the *p*-value of the Fisher's exact test for each class on the far right. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$. (**B**) Synteny alignment of human chromosomes 20 and 13, and mouse chromosomes 2 and 14, and aCGH analysis of $Apc^{\Delta716}$ mouse polyps shown as frequency plots. Red and green indicate regions of gain and loss, respectively. Positions of mouse *Tpx2* (arrowhead, top), *Aurka* (arrow, top), and *Bora* (arrow, bottom) are shown. The three examined polyps all exhibited amplification of *Tpx2*. Amplification of *Aurka* and *Bora* loci was found in two polyps. See also Fig. S6. (**C**) Relative DNA content of *AURKA*, *TPX2*, and *BORA* genes in normal tissues (N) of WT and $Apc^{\Delta716}$ mice and polyps (P) of $Apc^{\Delta716}$ mice. Seventeen polyps from four $Apc^{\Delta716}$ mice were analysed. (**D**) Global CNA score (GCS), broad CNA score (BCS), and focal CNA score (FCS) distribution by each class obtained from TCGA dataset analysis shown in (A). Wilcoxon rank-sum test significance is shown as * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$ (Tukey–Kramer-test in **C**, Wilcoxon rank-sum test in **D**).

Figure 5 Kawasaki et al.

1155

Fig. 5. Proliferation and AURKA activities of MCF10A derived cell lines. (**A**) Selected

images from phase contrast timelapse videos of MCF10A(20di) and MCF10A(20tri) cells.

See also Fig. S8A and Movie S4, 5. (**B**) Doubling time of MCF10A- and APC883-derived

cell lines. Note that no data are available for AURKA or APC depleted APC883 cells, as

their proliferation is arrested. (**C**) Immunostaining of MCF10A(20di) and MCF10A(20tri)

cells for AURKA and p-AURKA. (**D**) AURKA status at centrosomes obtained from

immunofluorescence images. Total AURKA protein levels (t-AURKA),

autophosphorylated AURKA levels (p-AURKA), the auto-phosphorylation ratio (p-

AURKA/t-AURKA ratio), and mitotic error rate in MCF10A- and APC883-derived cell

lines are shown as indicated. The values were normalized to 1 using MCF10A(20di) cells

as a reference. An example of a chromosome separation error (arrow) is shown in the image

on the right. (**E**) Selected images from phase contrast timelapse videos of the indicated cell

lines. See also Movie S4, 5. (**F**) Immunostaining of MCF10A and APC883 cells as

indicated. Position of spindle poles are indicated with arrow heads. β-catenin was up-

regulated in the whole cytoplasm and at the spindle pole in APC883 cells, but only faintly

detected on the spindles in MCF10A cells. In APC883 cells, AXIN1 and AURKA were

also increased, whereas the proportion of p-AURKA is low. (**G**) Immunostaining of

MCF10A expressing WTβ-catenin or MMβ-catenin for AURKA, showing accumulation of

MMβ-catenin and AURKA around the centrosomes. (**H**) Representative karyotype of

spontaneously transformed APC883 cells that contained a near tetraploid karyotype with

six copies of chromosome 20 (red box) and other common numerical imbalances (cyan and

brown boxes). Scale bars: 5 μm (C, F, G); 20 μm (A, E). *** $P < 0.001$; [##] $P < 0.01$; [###] $P <$

0.001 (Tukey–Kramer test).

1179

1180

Figure 6 Kawasaki et al.

1181

Fig. 6. Requirement of AURKA activity for tumor cell growth. (**A**) Effect of AURKA

inhibitor on the proliferative activity of mouse intestinal tissues. $Apc^{\Delta 716}$ mice were treated

with the vehicle or AURKA inhibitor alisertib (20 mg/kg) for 14 consecutive days.

Representative images of Ki67-labelled tissue sections from $Apc^{\Delta 716}$ mice (left). Analysis of

the Ki67-positive cell population (right). (**B**) Effect of AURKA inhibitor on polyp

development in $Apc^{\Delta716}$ mice treated with the vehicle or AURKA inhibitor for 28

consecutive days. Examples of HE-stained sections of intestines from vehicle- or alisertib-

treated $Apc^{\Delta716}$ mice (left). Analysis of the polyp number and size distribution in the

intestines of vehicle- or alisertib-treated $Apc^{\Delta716}$ mice (right). Scale bars: 50 μm (A); 5 mm

(B). ***$P < 0.001$ (Student's t-test). (**C**) Model showing the steps of malignant

transformation during early tumorigenesis. *APC* mutations induce CIN partly due to

AURKA inactivation and growth retardation. Some cells improve fitness via AURKA-

activating CNAs, adapting to promote tumor growth, which may contribute to the

formation of cancer-specific CNAs in *APC*-mutant tumors.

1196

Supplementary Materials for

Adaptive fitness in *APC*-mutant cells requires chromosome

rearrangements

Yoshihiro Kawasaki *et al.*

*Corresponding author. Email: kiyosuey@hirakata.kmu.ac.jp

This PDF file includes:

Figs. S1 to S8

Legend for Tables S1 to S3, Movies S1 to S5, and Code S1

References 1 to 25

Other Supplementary Materials for this manuscript include the following:

Tables S1 to S3

Movies S1 to S5

Fig1c.zip

Code S1

Fig. S1.

Structure and function of APC. (A) Domain structures of full length APC and major

binding partners (1-7). The *APC* gene is responsible for many sporadic cases of

gastrointestinal cancers and familial adenomatous polyposis, a dominantly inherited

colorectal tumor predisposition that results from germline mutations in *APC* (8-14).

Tumors start developing upon mutation of both alleles. In general, *APC* mutations that

result in stable proteins truncated in the region between the armadillo repeat and the first

SAMP motif are associated with severe polyposis phenotypes and, in the human colon, are

associated with a very malignant colorectal cancer type (reviewed in (4,15)). (B) Truncated

APC mutants. Phenotypes of cells and mice are shown on the right. Heterozygous *Apc*

mutations, such as *Apc^{Min}* and *Apc^{Δ716}*, lead to severe polyposis after loss of the wildtype

allele (>~30 polyps), whereas their homozygosity results in embryonic lethality (16-18).

Homozygous Apc^{1638T} mice with a truncating mutation just after the first SAMP motif,

which retains the β-catenin destruction ability, are viable and do not develop any tumors,

thereby demonstrating that this N-terminal fragment that retains the first SAMP motif is

sufficient for the tumor-suppressing function (19). Nevertheless, Apc^{1638T} homozygous ES

cells have a CIN phenotype, although it is weaker than that in Apc^{Min} homozygous ES cells

(20). MCF10A-derived cell line APC883 with a homozygous *APC* truncation was

generated by genome editing.

Characterization of RNAi tools and
expression construct in HeLa cells

MCF10A and APC883 derivatives

Characterization of RNAi tools
in MCF10A and APC883 cells

F   Culture duration

Fig. S2.

Characterization of RNAi tools, expression constructs, and cell lines used in this

study. (**A, B**) Western blot (WB) analysis of HeLa cells and HeLa cells that expressed

EB1-GFP and H2B-TagRFP (clone A1) (21) before and after constitutive transfection with

shRNA against APC (APC KD). APC knockdown efficiency was ~70%. (**C**)

Characterization of RNAi tools and expression construct in HeLa cells as indicated using

the indicated antibodies. (**D**) Characterization of MCF10A cells, APC883 cells, and

APC883 cell-derived transfectants as indicated using the indicated antibodies. Full-length

APC and truncated APC883 protein are indicated. For Myc-AURKA (T288D), endogenous

and exogenous Myc-tagged proteins are shown. (**E**) Characterization of RNAi tools in

MCF10A and APC883 cells as indicated. (**F**) Doubling time of parental MCF10A cells,

AURKA knockdown MCF10A cells and MMβ-catenin expressing MCF10A cells. n.s.

(Student's t-test showed not significant, *P*>0.05).

Fig. S3.

(**A, B**) Domain structures of APC and AXIN1, and the analyzed fragments. Amino acid

numbering is based on human APC transcript variant 3 (NM_000038) and human AXIN1

transcript variant 1 (NM_003502). Associations of APC with AURKA and AXIN1 were

assessed by in-cell colocalisation analysis using a series of deletion mutants. The results are

shown on the right. (**C**) Direct binding between APC fragments and AURKA/AXIN1/ch-

TOG predicted in (A) was confirmed by in vitro pull-down assays using purified proteins.

The APC arm and APC-C6 bound to AURKA. The APC arm also bound to the C-terminal

DIX domain of AXIN1. XMAP215 (22), the *Xenopus* homologue of ch-TOG, bound to

APC-C3. We used Xenopus XMAP215 because we could not purify full length human ch-

TOG from E. coli. The C-terminal one-third bound to the APC arm. See also Table S2. (**D**)

Yeast two-hybrid screening using the APC arm region as bait identified pericentrin. The

two identified pericentrin clones are shown.

Fig. S4.

Functions of AXIN1 and *Apc^1637T* gene expression products. (**A**) AXIN1 structure and

AURKA-activating region that also binds to β-catenin (23). (**B**) Effects of AXIN1 on

AURKA phosphorylation were analyzed using the HEK293T cell overexpression assay

system. HEK293T cells were transfected with GFP-fused AURKA and full length AXIN1

or fragments shown in Fig. S3B, lysed, and then subjected to western blot analysis using

anti-GFP and anti-p-T288 AURKA antibodies. The left panel is the same as Fig. 2E. See

also Table S2. (**C, D**) Analysis of primary MEFs from wildtype (WT) and Apc^{1638T} mice.

Immunostaining of microtubules (MTs), γ-tubulin, AURKA, and p-T288 AURKA (C).

Immunostaining intensity of γ-tubulin, the autophosphorylation ratio (p-AURKA/t-

AURKA ratio), total AURKA protein levels (t-AURKA), and autophosphorylated AURKA

levels (p-AURKA) were normalized to those of the wildtype, which were set to 1.0 (**D**).

Scale bars: 5 μm. ***$P < 0.001$ (Student's t-test).

Supplementary Figure S5   Kawasaki et al.

A

Human colon polyp #1
(Hyperplasia polyp)

Human colon polyp #2
(Low-grade tubular adenoma)

Truncations

APC

B

Hoechst Ki67
E-cadherin

Low   High Ki67
area

Hoechst   Ki67   E-cadherin

C

Hoechst
MTs   AURKA   p-AURKA

Polyp #1

Polyp #2
Low Ki67

High Ki67

p-AURKA
/ t-AURKA
ratio

t-AURKA
intensity

p-AURKA
intensity

(n, sample size)

D

Hoechst
E-cadherin
AURKA

Hoechst   E-cadherin   AURKA

Fig. S5.

Immunofluorescence analysis of paraffin-embedded sections of *APC*-mutant human

tumor tissues. (**A**) HE-stained sections of a human hyperplasia polyp without an APC

mutation (polyp #1) (left) and a low grade tubular adenoma with two APC truncation

mutations (polyp #2) (right). Two mutant APC sequences detected in polyp #2 are shown

(arrows). For details, see Materials and Methods. (B) Ki67 proliferation index in a low

grade tubular adenoma with two APC truncation mutations (polyp #2). On the basis of the

Ki67 intensity, the area was roughly divided into a highly proliferative region (outer, high

Ki67 area) and moderately proliferative region (inner, low Ki67 area) as indicated by the

yellow line. (C) Representative images of mitotic cells in polyp #1 and low and high Ki67

areas of polyp #2 (left). The centrosomes are indicated by yellow arrowheads. Quantitative

analysis of immunofluorescence signals of AURKA in polyp #1 and polyp #2 (right) at the

centrosomes. (D) Immunostaining of polyp #2 for E-cadherin and AURKA with nuclear

DNA stained by Hoechst 33342, showing appearance of a patchy region with extremely

high AURKA expression levels (yellow arow heads). Scale bars: 200 μm (A), 100 μm (C,

E), 5 μm (B). **$P < 0.01$; ***$P < 0.001$ (Tukey–Kramer test).

Fig. S6.

aCGH analysis of mouse tumors and syntenic comparison with human TCGA

analysis. aCGH analysis of chromosomes in three independent $Apc^{\Delta716}$ mouse polyps. A

penetration profile of copy number changes from duplicate analyzes is shown.

Chromosomal regions showing cross-species overlaps showing similar gain/loss pattern in

two or more samples are indicated by cyan lines. Arrow and arrowhead on chromosome 2

indicate the loci of *Aurka* and *Tpx2*, respectively. Arrow on chromosome 14 indicates the

locus of Bora.

Fig. S7.

TCGA data analysis for MSI status. (**A**) Frequency of microsatellite stable (MSS) and

microsatellite instable (MSI) in copy number changes in chromosomes with the most

frequent alterations. Fisher's extract test, **P < 0.01; ***P < 0.001. (**B**) Global CNA score

124   (GCS) distribution in *CTNNB1* (*β-catenin*) wildtype group (Classes 1–3) and *CTNNB1*

125   mutant group. Class 1: *APC* mutation = 0; Class 2: *APC* mutation = 1; Class 3: *APC*

126   mutation ≥2. Mann-Whitney test significance is shown as **$P < 0.01$. (**C**) Class1 and

127   CTNNB1 mutant were without significant difference. *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$

128   ****$P < 0.0001$ (Fisher's extract test).

Supplementary Figure S8   Kawasaki et al.

Fig. S8.

Karyotypes of MCF10A and APC883 cells. Representative karyotype images of each cell line (20 cells were analyzed in each cell line). For complete karyotype descriptions of all analyzed cells, see Table S3. The karyotypes were determined by Q-banding with Hoechst/quinacrine staining. Note that commonly observed abnormalities in chromosome 1 (deletion and partial duplication), chromosome 3 (deletion and translocation), chromosome 8 (duplication), and chromosome 9 (translocation) have been observed previously (24,25).

(**A–D**) Karyotype of parental MCF10A cells, APC883 cells, and APC883 derivatives.

Major karyotypes in each cell type and representative examples with chromosome

aberrations are shown. Chromosome 20 is indicated by red arrows. The parental MCF10A

cell line has two major populations with chromosome 20 disomy or chromosome 20

trisomy (types 1 and 2, respectively). All APC883 cells and APC883-derive cells observed

have chromosome 20 trisomy, the same as MCF10A type No. 2. Newly derived abnormal

chromosomes are indicated by red arrowheads. (**E**) Representative karyotypes of

transformed APC883 cells appeared after >5 months of passaging are shown. In this

transformed subpopulation, all analyzed cells had a near tetraploid karyotype with some

numerical imbalances. Common numerical imbalances are shown by the letter color for the

karyotype description: cyan, three chromosome 6s; brown, five chromosome 8s; red, six

chromosome 20s. Examples of randomly occurring imbalances are indicated by arrows.

Legend for Tables S1 to S3, Movies S1 to S5, and Code S1 (separate files)

Table S1.

Representative images used in the analysis, which could not be presented in the main body.

Table S2.

Uncropped versions of all gels and blots.

Table S3.

Full descriptions of the karyotypes of cell lines MCF10A and APC883, their subclones, and

transfectants. Newly generated chromosomes, which included truncation and chromosome

rearrangements, such as translocation, duplication, and telomere association, are shown in

bold. For the spontaneously transformed, near tetraploid APC883 cells, the common

number imbalance of chromosomes 6, 8, and 20 is indicated by color codes identical to the

box color in Fig. S8E: cyan, three chromosome 6s; brown, five chromosome 8s; red, six

chromosome 20s.

Movie S1.

Three-dimensional timelapse images of a mitotic HeLa cell that expressed EB1-GFP and

H2B-TagRFP collected by lattice light sheet microscopy at 10-s intervals are represented in

a 3D space. Time scale: h.m.s.

Movie S2.

Three-dimensional timelapse images of an APC-depleted mitotic HeLa cell that expressed

EB1-GFP and H2B-TagRFP collected by lattice light sheet microscopy at 10-s intervals are

represented in a 3D space. Time scale: h.m.s.

Movie S3.

Three-dimensional timelapse images of control and APC-depleted mitotic HeLa cells

shown in Movies S1 and S2 were projected onto a 2D plane and combined for comparison.

Time scale: h.m.s.

Movie S4.

Representative phase contrast timelapse videos of MCF10A(20di), MCF10A(20tri),

APC883, and APC883+β-catenin KD, APC883+AURKA(T288D), APC883+APC KD,

APC883+AURKA KD and transformed APC883 cells that spontaneously appeared after >5

months of passaging, shown in Fig. 5A, E. Timelapse images at 10-min intervals for 50 h

are displayed from videos that recorded at 5-min intervals. Time scale: h:m:s.ms.

Movie S5.

Representative phase contrast timelapse videos of MCF10A, MCF10A+AURKA KD and

MCF10A+MMβ-catenin cells, shown in Fig. S2F. Timelapse images at 10-min intervals for

50 h are displayed from videos that recorded at 5-min intervals. Time scale: h:m:s.ms.

Code S1.

Codes for TCGA data analysis. This folder also contains the data used to create the diagram

with the attached codes.

References

1. Aoki K, Taketo MM. Adenomatous polyposis coli (APC): a multi-functional tumor suppressor gene. J Cell Sci **2007**;120:3327-35

2. Fearnhead NS, Britton MP, Bodmer WF. The ABC of APC. Human molecular genetics **2001**;10:721-33

3. Polakis P. The adenomatous polyposis coli (APC) tumor suppressor. Biochimica et biophysica acta **1997**;1332:F127-47

4. Fodde R, Smits R. Disease model: familial adenomatous polyposis. Trends in molecular medicine **2001**;7:369-73

5. Pronobis MI, Rusan NM, Peifer M. A novel GSK3-regulated APC:Axin interaction regulates Wnt signaling by driving a catalytic cycle of efficient betacatenin destruction. eLife **2015**;4:e08022

6. Näthke IS. The adenomatous polyposis coli protein: the Achilles heel of the gut epithelium. Annual review of cell and developmental biology **2004**;20:337-66

7. Banks JD, Heald R. Adenomatous polyposis coli associates with the microtubule-destabilizing protein XMCAK. Current biology : CB **2004**;14:2033-8

8. Powell SM, Zilz N, Beazer-Barclay Y, Bryan TM, Hamilton SR, Thibodeau SN*, et al.* APC mutations occur early during colorectal tumorigenesis. Nature **1992**;359:235-7

9. Horii A, Nakatsuru S, Miyoshi Y, Ichii S, Nagase H, Kato Y*, et al.* The APC gene, responsible for familial adenomatous polyposis, is mutated in human gastric cancer. Cancer research **1992**;52:3231-3

10.   Boynton RF, Blount PL, Yin J, Brown VL, Huang Y, Tong Y*, et al.* Loss of

heterozygosity involving the APC and MCC genetic loci occurs in the majority of

human esophageal cancers. Proceedings of the National Academy of Sciences of the

United States of America **1992**;89:3385-8

11.   Nishisho I, Nakamura Y, Miyoshi Y, Miki Y, Ando H, Horii A*, et al.* Mutations of

chromosome 5q21 genes in FAP and colorectal cancer patients. Science

1991;253:665-9

12.   Kinzler KW, Nilbert MC, Vogelstein B, Bryan TM, Levy DB, Smith KJ*, et al.*

Identification of a gene located at chromosome 5q21 that is mutated in colorectal

cancers. Science **1991**;251:1366-70

13.   Joslyn G, Carlson M, Thliveris A, Albertsen H, Gelbert L, Samowitz W*, et al.*

Identification of deletion mutations and three new genes at the familial polyposis

locus. Cell **1991**;66:601-13

14.   Groden J, Thliveris A, Samowitz W, Carlson M, Gelbert L, Albertsen H*, et al.*

Identification and characterization of the familial adenomatous polyposis coli gene.

Cell **1991**;66:589-600

15.   Miyoshi Y, Nagase H, Ando H, Horii A, Ichii S, Nakatsuru S*, et al.* Somatic

mutations of the APC gene in colorectal tumors: mutation cluster region in the APC

gene. Human molecular genetics **1992**;1:229-33

16.   Oshima M, Oshima H, Kitagawa K, Kobayashi M, Itakura C, Taketo M. Loss of

Apc heterozygosity and abnormal tissue building in nascent intestinal polyps in

mice carrying a truncated Apc gene. Proceedings of the National Academy of

241   Sciences of the United States of America **1995**;92:4482-6

242 17. Su LK, Kinzler KW, Vogelstein B, Preisinger AC, Moser AR, Luongo C*, et al.*

243   Multiple intestinal neoplasia caused by a mutation in the murine homolog of the

244   APC gene. Science **1992**;256:668-70

245 18. Moser AR, Pitot HC, Dove WF. A dominant mutation that predisposes to multiple

246   intestinal neoplasia in the mouse. Science **1990**;247:322-4

247 19. Smits R, Kielman MF, Breukel C, Zurcher C, Neufeld K, Jagmohan-Changur S*, et*

248   *al.* Apc1638T: a mouse model delineating critical domains of the adenomatous

249   polyposis coli protein involved in tumorigenesis and development. Genes &

250   development **1999**;13:1309-21

251 20. Fodde R, Kuipers J, Rosenberg C, Smits R, Kielman M, Gaspar C*, et al.* Mutations

252   in the APC tumour suppressor gene cause chromosomal instability. Nature cell

253   biology **2001**;3:433-8

254 21. Chen BC, Legant WR, Wang K, Shao L, Milkie DE, Davidson MW*, et al.* Lattice

255   light-sheet microscopy: imaging molecules to embryos at high spatiotemporal

256   resolution. Science **2014**;346:1257998

257 22. Widlund PO, Stear JH, Pozniakovsky A, Zanic M, Reber S, Brouhard GJ*, et al.*

258   XMAP215 polymerase activity is built by combining multiple tubulin-binding TOG

259   domains and a basic lattice-binding region. Proceedings of the National Academy of

260   Sciences of the United States of America **2011**;108:2741-6

261 23. Xia Z, Wei P, Zhang H, Ding Z, Yang L, Huang Z*, et al.* AURKA governs self-

262   renewal capacity in glioma-initiating cells via stabilization/activation of beta-

catenin/Wnt signaling. Molecular cancer research : MCR **2013**;11:1101-11

24.   Soule HD, Maloney TM, Wolman SR, Peterson WD, Jr., Brenz R, McGrath CM*, et*

al. Isolation and characterization of a spontaneously immortalized human breast

epithelial cell line, MCF-10. Cancer research **1990**;50:6075-86

25.   Cowell JK, LaDuca J, Rossi MR, Burkhardt T, Nowak NJ, Matsui S. Molecular

characterization of the t(3;9) associated with immortalization in the MCF10A cell

line. Cancer genetics and cytogenetics **2005**;163:23-9

Figure: **Figure 2A**

Specimen: Control and APC KD HeLa cells
Labeling: 1   Hoechst33342
                    2   Anti-PLK1 antibody [36-298] (ab17057)
                    3   Anti-PLK1 (phospho T210) antibody (ab155095)
                         Note the nonspecific staining of chromosomes.
Scale bars: 5 μm

Figure: **Figure 5D**

Specimen: MCF10A control for the correction of MCF10A subclones
Labeling: 1   Hoechst33342
                    2   Anti-Aurora A antibody [35C1] (ab13824)
                    3   Anti-Aurora A (phospho T288) antibody (ab195748)
                          MCF10A data were used as references
                          to normalize data from independet experiments.

Scale bars: 5 μm

Specimen: MCF10A and APC883+β-catenin KD cells
Labeling: 1   Hoechst33342
                    2   Anti-Aurora A antibody [35C1] (ab13824)
                    3   Anti-Aurora A (phospho T288) antibody (ab195748)
                          MCF10A data were used as references
                          to normalize data from independet experiments.

Scale bars: 5 μm

Specimen: MCF10A and transformed APC883 cells
Labeling: 1   Hoechst33342
                    2   Anti-Aurora A antibody [35C1] (ab13824)
                    3   Anti-Aurora A (phospho T288) antibody (ab195748)
                    MCF10A data were used as references
                    to normalize data from independet experiments.

Scale bars: 5 μm

Figure: **Figure 2B**

Specimen: *in vitro* phosphorylation assay
Lane: 1   AURKA (A28-14G)
                 2   AURKA (A28-14G) + GST (for GST protein, see p.16)
                 3   AURKA (A28-14G) + AXIN1(TP308349)
                 4   AURKA (A28-14G) + TPX2 (T40-30H)

Blot 1:   **Blot 2:**
Antibody: anti-AURKA (ab13824)   **Antibody:**   anti-p-AURKA (ab195748)

Figure: **Figure 2D**

Specimen: HEK293T cell lysate transfected with the plasmids shown below
Lane: 1   GFP-AURKA (0.6 ug)
                2   GFP-AURKA (0.6 ug) + GFP-AXIN1 (0.5 ug)
                3   GFP-AURKA (0.6 ug) + GFP-AXIN1 (0.5 ug)
                4   GFP-AURKA (0.6 ug) + GFP-AXIN1 (0.5 ug) + GFP-β-catenin(0.25 ug)
                5   GFP-AURKA (0.6 ug) + GFP-AXIN1 (0.5 ug) + GFP-β-catenin(0.5 ug)
                6   GFP-AURKA (0.6 ug) + GFP-AXIN1 (0.5 ug) + GFP-β-catenin(0.75 ug)
                7   GFP-AURKA (0.6 ug) + GFP-AXIN1 (0.5 ug) + GFP-β-catenin(1 ug)

Blot 1:   Top left panel   **Blot 2:**   Top left panel
Antibody: anti-GFP-HRP (598-7)   **Antibody:**   anti-p-AURKA (ab195748)

Blot 3:   Top right panel   **Blot 4:**   Bottom right panel
Antibody: anti-GFP-HRP (598-7)   **Antibody:**   anti-p-AURKA (ab195748)

Figure: **Figure 2E**

Specimen: HEK293T cell lysate transfected with the plasmids shown below
Lane: 1   GFP-AURKA (0.6 ug)
                       2   GFP-AURKA (0.6 ug) + GFP-AXIN1 (0.2 ug)
                       3   GFP-AURKA (0.6 ug) + GFP-AXIN1 (0.2 ug) + GFP-APC-N (0.3 ug)
                       4   GFP-AURKA (0.6 ug) + GFP-AXIN1 (0.2 ug) + GFP-APC-M (0.8 ug)
                       5   GFP-AURKA (0.6 ug) + GFP-AXIN1 (0.2 ug) + GFP-APC-C (0.6 ug)
                       6   GFP-AURKA (0.6 ug) + GFP-AXIN1 (0.2 ug) + GFP-APC-SAMP (0.6 ug)

Blot 1:   Top panel
Antibody: anti-GFP-HRP (598-7)

Blot 2:   Bottom panel
Antibody: anti-p-AURKA (ab195748)

Figure: **Figure S2A**

Specimen: Lysate of HeLa / EB1-GFP + H2B-TagRFP cell line (clone A1)
Lane: 1 HeLa, control
2 HeLa + APC KD
To detect APC protein, 12 times the protein concentration of the control was loaded.

Blot 1: Top panel
Antibody: Anti-APC [Ab-1] (OP44)

Blot 2: Bottom panel
Antibody: anti-GAPDH-HRP (ab9482)

Figure: **Figure S2B**

Specimen: Lysate of HeLa cells
Lane: 1   HeLa, control
    2   HeLa + APC KD
   To detect APC protein, 12 times the protein concentration of the control was loaded.

Blot 1:   Top panel
Antibody: Anti-APC [Ab-1] (OP44)

← Full-length APC

Blot 2:   Bottom panel
Antibody: anti-GAPDH-HRP (ab9482)

Figure: **Figure S2C, left**

Specimen: Lysate of HeLa cells
Lane: 1   HeLa, control
                          2   HeLa + APC KD
                          3   HeLa + AXIN1 KD

Blot 1:   Top panel   **Blot 2:**   2nd panel from top
Antibody: Anti-AXIN1 [C76H11] (#2087)   **Antibody:**   Anti-β-catenin (610154)

Blot 3:   3rd panel from top
Antibody: anti-GAPDH-HRP (ab9482)

Figure: **Figure S2C, right**

Specimen: Lysate of HeLa cells
Lane: 1   HeLa, control
                        Unrelated sample
                        Unrelated sample
                   2   HeLa + Myc-MMβ-catenin
                Lanes of unrelated samples were cut out.

Blot 1:   Top panel   **Blot 2:**   2nd panel from top
Antibody: Anti-β-catenin (610154)   **Antibody:** Anti-Myc Tag [4A6] (05-724)

Blot 3:   3rd panel from top   **Blot 3:**   4th panel from top
Antibody: Anti-AXIN1 [C76H11] (#2087)   **Antibody:** anti-GAPDH-HRP (ab9482)

Figure:　　**Figure S2D, left**

Specimen:　Lysate of MCF10A/APC883 cells
Lane:　　　1　MCF10A
　　　　　　　2　APC883
　　　　　　　3　APC883 + Myc-AURKA(T288D)
　　　　　　　4　APC883 + β-catenin KD
　　　　　　　The same amount of protein was loaded in each lane.

Blot 1:　　Top panel　　　　　　　　　**Blot 2:**　　2nd panel from top
Antibody:　Anti-APC [Ab-1] (OP44)　　　**Antibody:**　Anti-β-Catenin (610154)

Blot 3:　　3rd panel from top
Antibody:　Anti-AXIN1 [C76H11] (#2087)

Figure: **Figure S2D, left, continue**

Specimen: Lysate of MCF10A/APC883 cells
Lane: 1   MCF10A
                        2   APC883
                        3   APC883 + Myc-AURKA(T288D)
                        4   APC883 + β-catenin KD
                The same amount of protein was loaded in each lane.

Blot 4:   4th panel from top   **Blot 5:**   5th panel from top
Antibody: Anti-Aurora A [35C1] (ab13824)   **Antibody:**   Anti-Myc Tag [4A6] (05-724)

Blot 6:   6th panel from top
Antibody: anti-α-tubulin [DM1A] (T9026)

Figure: **Figure 2D, right**

Specimen: Lysate of MCF10A cells
Lane: 1   MCF10A
                 2   MCF10A + Myc-MMβ-catenin
        The same amount of protein was loaded in each lane.

Blot 1:   Top panel
Antibody: Anti-β-Catenin (610154)

Blot 2:   2nd panel from top
Antibody: Anti-Myc Tag [4A6] (05-724)

Blot 3:   3rd panel from top
Antibody: Anti-AXIN1 [C76H11] (#2087)

Blot 4:   4th panel from top
Antibody: anti-GAPDH-HRP (ab9482)

Figure: **Figure S2E**

Specimen: Lysate of APC883 cells
Lane: 1   APC883, control
                2   APC883 + APC KD

Blot 1:   Top panel   **Blot 2:**   2nd panel from top
Antibody: Anti-APC [Ab-1] (OP44)   **Antibody:** anti-GAPDH-HRP (ab9482)

Specimen: Laysates of MCF10A cells
Lane: 1   MCF10A, control
                2   MCF10A + AURKA KD

Blot 1:   Top panel   **Blot 2:**   2nd panel from top
Antibody: Anti-Aurora A [35C1] (ab13824)   **Antibody:** anti-GAPDH-HRP (ab9482)

Figure: **Figure S3C**

Specimen: *in vitro* pull-down assay
Lane: 1   Input (HA-AURKA)
                 2   + GST
                 3   + GST-APC-arm
                     (for GST proteins, see p.16)

Specimen: *in vitro* pull-down assay
Lane: 1   Input (FLAG-AURKA)
                 2   + GST
                 3   + GST-APC-C4
                 4   + GST-APC-C5
                 5   + GST-APC-C6
                     (for GST proteins, see p.16)

Blot 1:   Left panel
Antibody: Anti-HA [3F10] (11867423001)

Blot 2:   2nd panel from left
Antibody: Anti-FLAG [M2] (F1804)

Specimen: *in vitro* pull-down assay
Lane: 1   Input (FLAG-AXIN1-DIX)
                 2   + GST
                 3   + GST-APC-arm
                     (for GST proteins, see p.16)

Specimen: *in vitro* pull-down assay
Lane: 1   Input (XMAP215/ch-TOG-His)
                     (PNAS, 108 (7), 2741-2746, 2011)
                 2   + GST
                 3   + GST-APC-C3
                     (for GST proteins, see p.16)

Blot 3:   3rd panel from left
Antibody: Anti-FLAG [M2] (F1804)

Blot 4:   4th panel from left
Antibody: Anti-penta-His (34660)

Figure: **Figure S4B**

Specimen: HEK293T cell lysate transfected with the plasmids shown below
Lane: 1   GFP-AURKA (0.6 ug) + GFP-AXIN1-N (0.5 ug)
                           2   GFP-AURKA (0.6 ug) + GFP-AXIN1-M (0.5 ug)
                           3   GFP-AURKA (0.6 ug) + GFP-AXIN1-C (0.5 ug)
                    The left panel of Fig. S4B is the same as Fig. 2E.

Blot 1:   Top right panel   **Blot 2:**   Bottom right panel
Antibody: anti-GFP-HRP (598-7)   **Antibody:**   anti-p-AURKA (ab195748)

Figure: **Figure S9**

Specimen: HEK293T cell lysate transfected with the plasmids shown below
Lane: 1   GFP-fused wild type (WT) AURKA
                2-6   GFP-fused AURKA mutant#1 and subsequent mutnats
                           cloned from CRC cell lines

Blot 1:   HT-29, top panel   **Blot 2:**   HT-29, bottom panel
Antibody: anti-GFP-HRP (598-7)   **Antibody:**   anti-p-AURKA (ab195748)

Blot 4:   SW480, top panel   **Blot 5:**   SW480, bottom panel
Antibody: anti-GFP-HRP (598-7)   **Antibody:**   anti-p-AURKA (ab195748)

Figure: **Figure S9, continue**

Specimen: HEK293T cell lysate transfected with the plasmids shown below
Lane: 1 GFP-fused wild type (WT) AURKA
  2-7 or 8 GFP-fused AURKA mutant#1 and subsequent mutnats
    cloned from CRC cell lines

Blot 5: DLD-1, top panel
Antibody: anti-GFP-HRP (598-7)

Blot 6: DLD-1, bottm panel
Antibody: anti-p-AURKA (ab195748)

Blot 7: HCT116, top panel
Antibody: anti-GFP-HRP (598-7)

Blot 8: HCT116, bottm panel
Antibody: anti-p-AURKA (ab195748)

Figure: **Materials used in Figure 2B, C and Figure S3C**

Specimen: Coomassie Brilliant Blue (CBB) staining of the recombinant protein
Lane: 1   GST
               2   GST-APC-arm
               3   GST-APC-C4
               4   GST-APC-C5
               5   GST-APC-C6

MCF10A   The amplified chromosome 20 is underlined. Same as below, except transformed APC883.

Type No.	Karyotype	Number of cells
1	47,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24)	5
2	48,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),<u>+20</u>	10
3	46,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),**der(5)ins(5)(q10?)inv(5)(q11.2q35)**,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),-21	1
4	47,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),<u>+20</u>,-21	1
5	47,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),-19,<u>+20</u>	1
6	48,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3)	1
7	49,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p	1

MCF10A(20di)   Subclone of MCF10A carrying chromosome 20 disomy

Type No.	Karyotype	Number of cells
1	47,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24)	20

MCF10A(20tri)   Subclone of MCF10A carrying chromosome 20 trisomy

Type No.	Karyotype	Number of cells
1	48,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),+20	18
2	47,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),-22	1
3	49,XX,del(1)(q11q32),+del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),+20	1

APC883

Type No.	Karyotype	Number of cells
1	48,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),<u>+20</u>	15
2	47,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),**<u>+dic(20;22)(q13.3;p13)</u>**	1
3	47,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),**der(7)dic(7;9)(p22;q34)add(9)(p24)**,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),<u>+20</u>	1
4	47,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),-15,<u>+20</u>	1
5	48,XX,**der(1)del(1)(q11q32)add(1)(q44)**,+i(1)(q10),del(3)(p21),**dic(6;14)(q27;q32)**,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),<u>+20</u>	1
6	48,XX,del(1)(q11q32),**+der(1)i(1)(q10)t(1;13)(q44;p13)**,del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),<u>+20</u>,**+mar**	1

APC883 + β-catenin KD

Type No.	Karyotype	Number of cells
1	48,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),<u>+20</u>	15
2	47,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),<u>+20</u>,-21	3
3	47,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),<u>+20</u>,-22	1
4	48,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),-18,<u>+20</u>,**+mar**	1

APC883 + AURKA(T288D)

Type No.	Karyotype	Number of cells
1	48,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),+20	18
2	47,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24)	1
3	47,XX,del(1)(q11q32),+i(1)(q10),del(3)(p21),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),-14,+20	1

Transformed (Near-tetraploid) APC883

Type No.	Karyotype	Number of cells
1	95,XXXX,del(1)(q11q32),del(1)(q11q32),+i(1)(q10),del(3)(p21),del(3)(p21),-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),add(9)(p24),+20,+20	3
2	96,XXXX,del(1)(q11q32),del(1)(q11q32),+i(1)(q10),+i(1)(q10),del(3)(p21),del(3)(p21),-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),add(9)(p24),+20,+20	3
3	94,XXXX,del(1)(q11q32),del(1)(q11q32),+i(1)(q10),del(3)(p21),del(3)(p21),-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),add(9)(p24),-13,+20,+20	2
4	93,XXXX,del(1)(q11q32),del(1)(q11q32),+i(1)(q10),del(3)(p21),del(3)(p21),-6, der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),add(9)(p24),-15,-18,+20,+20	1
5	93,XXXX,del(1)(q11q32),del(1)(q11q32),i(1)(q10),-2,del(3)(p21),del(3)(p21),-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),add(9)(p24),+20,+20	1
6	94,XXXX,del(1)(q11q32),del(1)(p10),+i(1)(q10),del(3)(p21),-3,-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),add(9)(p24),+20,+20	1
7	94,XXXX,del(1)(q11q32),del(1)(q11q32),i(1)(q10),del(3)(p21),del(3)(p21),-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),add(9)(p24),+20,+20	1
8	94,XXXX,del(1)(q11q32),del(1)q11q32),+i(1)(q10),del(3)(p21),del(3)(p21),-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),add(9)(p24),**add(11)(p15)**,+20,+20,-21	1
9	95,XXXX,del(1)(q11q32),del(1)(q11q32),+i(1)(q10),+i(1)(q10),del(3)(p21),del(3)(p21),-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),add(9)(p24),-13,+20,+20	1
10	95,XXXX,del(1)(q11q32),del(1)(q11q32),i(1)(q10),+i(1)(q10),del(3)(p21),del(3)(p21),-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),add(9)(p24),+20,+20	1
11	96,XXXX,del(1)(q11q32),del(1)(q11q32),+i(1)(q10),+i(1)(q10),del(3)(p21),del(3)(p21),-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),**+i(8)(q10)**,add(9)(p24),add(9)(p24),-19,+20,+20	1
12	96,XXXX,del(1)(q11q32),del(1)(q11q32),+i(1)(q10),+i(1)(q10),del(3)(p21),del(3)(p21),-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),add(9)(p24),**add(13)(p13)**,+20,+20	1

13	97,XXXX,del(1)(q11q32),del(1)(q11q32),+del(1)(p10),+i(1)(q10),+i(1)(q10),del(3)(p21),del(3)(p21),-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3) der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),add(9)(p24),**del(19)(q10)**,+20,+20	1
14	97,XXXX,del(1)(q11q32),del(1)(q11q32),+del(1)(p10),dup(2)(q11q37),del(3)(p21),+del(3)(p21),-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3), der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),+add(9)(p24),+20,+20	1
15	97,XXXX,del(1)(q11q32),del(1)(q11q32),del(1)(q11q32),+i(1)(q10),+2,del(3)(p21),-3,-6,der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3), der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),+der(8)dup(8)(q22q24.3)ins(8)(p24.3q22q24.3),add(9)(p24),add(9)(p24),+20,+20	1