Reporting quality of trend analyses published in leading medicine and oncology journals during 2008-2018

Xiaoling Yuan,[1] Fei Deng,[2] Yating Wang,[3*] Lanjing Zhang,[2, 4, 5*]

[1]Department of Infectious Disease, Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China; [2]Department of Chemical Biology, Ernest Mario School of Pharmacy, Rutgers University, Piscataway, New Jersey; [3]Department of Oncology, Ascension All Saint Cancer Center, Racine, Wisconsin, USA; [4]Department of Pathology, Princeton Medical Center, Plainsboro, NJ; [5]Rutgers Cancer Institute of New Jersey, New Brunswick, New Jersey;

Correspondence: Lanjing Zhang, MD, Department of Pathology, Princeton Medical Center, 1 Plainsboro Rd., Plainsboro, NJ 08563. Email: lanjing.zhang@rutgers.edu

*Equal contribution.

Running head: Reporting quality of trend analyses

Word count: 1547

Key words: Trend, reporting quality, medical journal, oncology journal

Abstract

Reporting quality of clinical research is critical for evidence-based medicine and reproducibility of clinical research. Most of the works focused on reporting quality of clinical trials and observational longitudinal studies. However, few focused on that of trend analyses. The reporting of recommended statistic metrics in trend analyses was also largely unclear. Therefore, we examined reporting quality of the trend analyses based on reporting of recommended statistic metrics. We systemically searched the PubMed for the trend-analysis articles published in 10 leading medicine and oncology journals during the 11 years from 2008 to 2018. The studies published after 2019 were not included due to the sudden, significant increase of publication number during and immediately after the COVID-19 pandemic. Only original articles, research letters and meta-analyses/systematic reviews were included. We scored the reporting quality of these articles based on whether they reported p-values/effect-sizes, and beta/co-efficient/slope/ annual-percentage-change (APC). There were 297 qualified articles, among which 193 (66.0%) and 216 (72.7%) articles reported P-value and effect-size, respectively. Only 13 (5.8%) analyses reported neither p-value/effect size nor beta/coefficient/slope/APC. In multivariable regression models, author affiliation of epidemiology department was associated with less reporting effect-size, but that of statistics department with more reporting. Interestingly, U.S. senior-authors (versus non-U.S.) more likely reported p-values. No factors were independently linked to reporting APC. The reporting quality of trend analyses in leading medicine and oncology journals appear moderate and should be further improved. We thus call for more research and awareness of reporting-quality in trend analyses in oncology research and beyond.

Introduction

Reporting quality of clinical research is critical for evidence-based medicine and reproducibility of clinical research. Most of the works focused on reporting quality of clinical trials, including those in neuro-oncology, urology, cardiology, nephrology, pharmaceutics and infectious disease.[1-7] Although the research quality of clinical trials was improving,[8 9] that of observational longitudinal studies remains low.[10-12] However, few studies concerned the reporting quality of trend analyses.

Trend analyses are critical for assessing changes and predicting future of epidemiological parameters.[13 14] Recent trend-analysis guidelines recommended reporting slopes or beta/coefficient if possible.[15] American Statistical Association and others recommended reporting effect-size.[16 17] However, reporting of these recommended statistic metrics in trend analyses was largely unclear. Therefore, we examined reporting quality of the trend analyses in leading medicine and oncology journals, with reported p-value, effect-size or beta/coefficient/Annual percent change (APC) as the quality metrics. We also identified factors associated with reporting quality in these trend analyses.

Methods

We systemically searched the PubMed for the articles published during the 11 years from January 1, 2008 to December 31, 2018, whose titles included "trend" or "trends" among the following medicine and oncology journals: *Ann Intern Med, Ann Oncol, BMJ, J Clin Oncol, J Natl Cancer Inst, JAMA Oncol, JAMA, Lancet, Lancet Oncol* and *N Engl J Med*. It was considered to include articles published after 2019. However, there were surges in publication number during the pandemic of COVID-19 (also known as SARS-CoV-2 coronavirus) and after.[18 19] We felt the

reporting quality might be influenced by the pandemic and surges in publication number. Therefore, we remain focused on the 11 years of 2008-2018.

We only included original articles, research letters and meta-analyses/systematic reviews of trend analysis. We limited our search to title words to ensure articles focused primarily on trend analysis. We acknowledge this approach might miss some relevant articles and discuss this limitation. Three authors independently reviewed full-texts using a standardized data extraction form: publication year, journal specialty (medicine/oncology), model type, reporting of *P-value*, effect-size (defined as quartiles/confidence/credible/uncertainty intervals) and co-efficient/slope/Annual Percent Change (APC), senior author(s)' location, and any author in School of Public Health, statistics department, or epidemiology department. When there was discrepancy, they would discuss and reconcile the differences. In the rare events that a consensus could not be reached, Dr Zhang would make the final decision.

According to guidelines,[15] co-efficient/slope/APC should be reported in (piece-wise) linear models. We assessed reporting whether co-efficient/slope/APC was reported in the articles using linear model. We also scored the reporting quality of these articles by assigning 1 point for reporting a p-value or effect-size, and another point for reporting a beta/co-efficient/slope/APC. For articles reporting the same analysis, each article would be assessed independently. The sum of each article's scores was considered as its reporting-quality score, which was up to 2 points. Points/scores were unweighted since the metrics cover different aspects of the statistics. The clinical significance of various metrics was to our knowledge not differentiated or compared by known recommendations,[15-17] although they are all clinically important and useful.

We used Chi-square test, Fisher exact test, and (ordinal) logistic regression to examine potential associations (Stata, version 15). Only the factors with P<0.10 shown in univariate analysis were included in multivariable logistic regression analyses. Two-sided P-values were reported. Statistical significance was considered when P<0.05.

Results and Discussion

Among the 398 identified reports of trend analysis published during 2008-2018, there were 297 qualified reports (**Supplementary Figure**), including 38 (12.8%) analyses using non-parametric model, 226 (76.1%) analyses using (piece-wise) linear model, 32 (10.8%) analyses using non-linear parametric model and 1 (0.3%) analysis using semi-parametric model (Cox regression). Among these analyses, 193 (66.0%) and 216 (72.7%) analyses reported P-value and effect-size, respectively. Subgroup analyses showed that U.S. senior authors more likely reported P-value or effect-size than non-U.S. senior authors (**Figure 1**), while reporting of these parameters was not linked to any of the other factors.

Among the 226 trend analyses using (piecewise) linear model (**Table 1**), 169 (74.8%) reported p-value, 183 (81.0%) reported effect-size, 94 (41.6%) reported APC and 34 (15.0%) reported Beta/coefficient/slope. No multiple articles reported the same or similar analysis. Only 13 (5.8%) analyses reported neither p-value/effect size nor beta/coefficient/slope/APC. Ordinal logistic regression showed only author affiliation with school of public health was linked to higher reporting-quality scores (Odds ratio=7.44, 95% confidence interval, 3.22 - 31.17). In multivariable regression models (**Table 2**), author affiliation with epidemiology or statistics

department was associated with reporting effect-size, and U.S. senior-authors (versus non-U.S.) more likely reported p-value. No factors were independently linked to reporting APC (**Table 2**).

Reporting quality of the included trend analyses was overall good. This is consistent with reported increasing quality in clinical trials.[9] However, several reporting-quality issues are concerning. Reporting p-value or effect-size did not change with publication years, despite the recommendation on using effect-size.[16][17] The underlying reasons are worth additional investigation.

Moreover, Non-U.S. senior authors less frequently reported p-value or effect-size than U.S. ones, which warrants more research and training. Furthermore, more than half of the trend analyses using linear model did not report p-value/effect-size, slope/beta/APC or either, which is not consistent with recommendations.[15] Without these metrics, quantification and comparison of linear models would be difficult or impossible. Therefore, the scientific rigor in these papers is significantly reduced.

In addition, we show that involvement of statistics department appears to link to more reporting effect sizes of oncological trend analyses, while that of epidemiology department linked to fewer reporting effect sizes. The rigorous training in statistics and its departments may contribute to this finding. Indeed, involvement of biostatisticians or epidemiologists was associated with higher methodological quality,[7] higher acceptance rates,[20] and shorter time to publication.[21] Thus, it has also been advocated to include statisticians in more clinical research and trials.[10][22][23]

The paradoxical finding that epidemiology department involvement was associated with less effect-size reporting (OR=0.38, 95% CI: 0.18-0.80, p=0.011) warrants careful interpretation. One possible explanation is that the emphasis on effect size reporting is relatively recent (2016 and 2019), [16][17] yet epidemiologists may historically prefer other metrics. Given our sole focus on trend analyses in this study, this finding may not be applicable to, but should be examined in, other types of epidemiological and clinical studies. Strikingly, few studies on reporting quality of clinical trials or longitudinal studies separated biostatisticians and epidemiologists. [7][20][21] Thus, future works are warranted to fill this knowledge gap.

There are few studies focused on reporting quality of trend analyses, while there are many on that of clinical trials or epidemiological studies. [7][9][11][12][24] Some experiences and data in clinical trials and trend analyses may be applicable to each other. First, development and implementation of reporting guidelines appear to improve reporting quality of randomized clinical trials. [24] However, there are no official reporting guidelines on trend analyses which may be needed. We thus recommend developing and implementing reporting guidelines on trend analyses, beyond recommendations. [15] Second, librarians and information specialists did not impact reporting quality of systematic reviews but have helped with the journal review process. [25] It will be interesting to understand whether librarians and information specialists will help improve reporting quality of trend analyses. Third, our findings seem to call for journal editors and peer reviewers to more rigorously enforce reporting standards and improve reporting quality. Indeed, one of the recommendations was published by a leading medical journal. [17] Publishers and professional societies may also be more actively involved in enforcing reporting standards of trend analyses. Finally, our data seem to call for more involvement of statisticians in trend analyses in oncology research, probably also medical research at large. It is

concerning that involvement of epidemiological department was associated with less reporting of effect size. More works are needed to confirm our findings and understand the reason. It is noteworthy that involvement of either department was not linked to better reporting scores.

This study has several limitations. First, our search strategy using only 'trend/trends' in titles may have missed relevant analyses using alternative terminology (e.g., 'temporal changes,' 'secular patterns'). Second, we focused on high-impact journals, limiting generalizability to the broader literature. However, it is possible that journals of intermediate or low impact may publish trend analyses with worse quality than those of high impact. Future works are warranted to test this hypothesis. Third, we did not include any trend analyses published after 2018. These recent works are interesting due to the surges in publication numbers during and after the COVID-19 pandemic, [18][19] but may require a longer timeframe and more data than currently available data to reliably examine the biases and impact associated with the COVID-19 pandemic. Otherwise, the conclusions may be inaccurate, if not misleading. Third, our title-based search may have missed some relevant articles, that mentioned trend analyses only in the abstract or text. Although these may be a small portion of truly relevant articles, this limitation might affect the generalizability of the findings. Finally, the used reporting-quality metrics may not be applicable to all reports. Indeed, some trends are difficult to accurately model with a single algorithm, and thus may not report some of the quality metrics.

In summary, the reporting quality of trend analyses in leading medicine and oncology journals could be further improved. We thus call for more research and awareness of reporting-quality in trend analyses in oncology research and beyond.

Author Contributions: Drs Zhang and Wang had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. They equally supervised the works, and share the senior authorship.

Concept and design: Yuan, Wang, Zhang.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Yuan.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Yuan, Deng, Zhang.

Supervision: Zhang, Wang.

Funding:

This work was supported by the National Cancer Institute, National Institutes of Health (grant number R37CA277812 to LZ). The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Conflict of interest

Dr Zhang is a co–editor in chief of *Exploratory Research and Medical Hypothesis*.

Figure Legend

Proportions of the trend analyses published during 2008-2018 reporting p-value and effect-size and their associated factors. Only senior author's affiliation (U.S.A. vs non-U.S.A.) was linked to reporting either p-value (Left) or effect size (Right) among the 397 trend analyses published in leading medicine and oncology journals during 2008-2018.

Bibliography

1. Yang Z, He H, Wang R, Liu D, Li G, Sun F. Application and Quality of Model-Based Meta-Analysis in Pharmaceutical Research: A Systematic Cross-Sectional Analysis and Practical Considerations. Clin Pharmacol Ther 2024;116(2):397-407. doi: 10.1002/cpt.3290.

2. Suppree JS, Hart S, Keshwara SM, Trichinopoly Krishna S, Gillespie CS, Richardson GE, et al. Assessing the reporting quality of pediatric neuro-oncology protocols, abstracts, and trials: Adherence to the SPIRIT and CONSORT statements. Neurooncol Pract 2024;11(5):617-32. doi: 10.1093/nop/npae042.

3. Norling B, Edgerton Z, Bakker C, Dahm P. The Quality of Literature Search Reporting in Systematic Reviews Published in the Urological Literature (1998-2021). J Urol 2023;209(5):837-43. doi: 10.1097/ju.0000000000003190.

4. Deng J, He J, Wang J, Cheng CW, Jiao Y, Wang N, et al. Reporting quality of randomized controlled trials of angina pectoris with integrated traditional Chinese and western medicine interventions: a cross-sectional study. BMC Med Res Methodol 2023;23(1):124. doi: 10.1186/s12874-023-01953-1.

5. Chatzimanouil MKT, Wilkens L, Anders HJ. Quantity and Reporting Quality of Kidney Research. J Am Soc Nephrol 2019;30(1):13-22. doi: 10.1681/asn.2018050515.

6. Nagai K, Saito AM, Saito TI, Kaneko N. Reporting quality of randomized controlled trials in patients with HIV on antiretroviral therapy: a systematic review. Trials 2017;18(1):625. doi: 10.1186/s13063-017-2360-2.

7. Delgado-Rodriguez M, Ruiz-Canela M, De Irala-Estevez J, Llorca J, Martinez-Gonzalez A. Participation of epidemiologists and/or biostatisticians and methodological quality of published controlled clinical trials. J Epidemiol Community Health 2001;55(8):569-72. doi: 10.1136/jech.55.8.569.

8. Dechartres A, Charles P, Hopewell S, Ravaud P, Altman DG. Reviews assessing the quality or the reporting of randomized controlled trials are increasing over time but raised questions about how quality is assessed. Journal of clinical epidemiology 2011;64(2):136-44. doi: 10.1016/j.jclinepi.2010.04.015.

9. Dechartres A, Trinquart L, Atal I, Moher D, Dickersin K, Boutron I, et al. Evolution of poor reporting and inadequate methods over time in 20 920 randomised controlled trials included in Cochrane reviews: research on research study. BMJ (Clinical research ed) 2017;357:j2490. doi: 10.1136/bmj.j2490.

10. Van Calster B, Wynants L, Riley RD, van Smeden M, Collins GS. Methodology over metrics: current scientific standards are a disservice to patients and society. Journal of clinical epidemiology 2021;138:219-26. doi: 10.1016/j.jclinepi.2021.05.018.

11. Trentini B, Steindel M, Marlow MA. Low quality evidence of epidemiological observational studies on leishmaniasis in Brazil. PLoS One 2014;9(9):e106635. doi: 10.1371/journal.pone.0106635.

12. Tooth L, Ware R, Bain C, Purdie DM, Dobson A. Quality of reporting of observational longitudinal research. American journal of epidemiology 2005;161(3):280-8. doi: 10.1093/aje/kwi042.

13. Zhang J, Lin Y, Zhang L. Trends in Alcoholic Fatty Liver Disease. Jama 2019;322(10):979-80. doi: 10.1001/jama.2019.10347.

14. Wong T, Dang K, Ladhani S, Singal AK, Wong RJ. Prevalence of Alcoholic Fatty Liver Disease Among Adults in the United States, 2001-2016. Jama 2019;321(17):1723-25. doi: 10.1001/jama.2019.2276.

15. Ingram DD, Malec DJ, Makuc DM, Kruszon-Moran D, Gindi RM, Albert M, et al. National Center for Health Statistics Guidelines for Analysis of Trends. Vital Health Stat 2 2018(179):1-71.

16. Wasserstein RL, Lazar NA. The ASA statement on p-values: context, process, and purpose. The American Statistician 2016;70(2):129-33. doi: 10.1080/00031305.2016.1154108.

17. Harrington D, D'Agostino RB, Sr., Gatsonis C, Hogan JW, Hunter DJ, Normand ST, et al. New Guidelines for Statistical Reporting in the Journal. The New England journal of medicine 2019;381(3):285-86. doi: 10.1056/NEJMe1906559.

18. Aviv-Reuven S, Rosenfeld A. Publication patterns' changes due to the COVID-19 pandemic: a longitudinal and short-term scientometric analysis. Scientometrics 2021;126(8):6761-84. doi: 10.1007/s11192-021-04059-x.

19. Rousseau R, Garcia-Zorita C, Sanz-Casado E. Publications during COVID-19 times: An unexpected overall increase. Journal of Informetrics 2023;17(4):101461. doi: https://doi.org/10.1016/j.joi.2023.101461.

20. Altman DG, Goodman SN, Schroter S. How statistical expertise is used in medical research. Jama 2002;287(21):2817-20. doi: 10.1001/jama.287.21.2817.

21. Sebo P, Fournier JP, Maisonneuve H. Is statistician involvement as co-author associated with reduced time to publication of quantitative research in general medical journals? A bibliometric study. Fam Pract 2019;36(4):431-36. doi: 10.1093/fampra/cmy115.

22. Stocken DD, Mossop H, Armstrong E, Lewis S, Dutton SJ, Peckitt C, et al. Good Statistical Practice-development of tailored Good Clinical Practice training for statisticians. Trials 2024;25(1):113. doi: 10.1186/s13063-024-07940-1.

23. Li Z, Wang X, Barnhart HX, Wang Y. Working With Statisticians in Clinical Research. Stroke 2018;49(11):e311-e13. doi: 10.1161/STROKEAHA.118.022266.

24. Tan ZW, Tan AC, Li T, Harris I, Naylor JM, Siebelt M, et al. Has the reporting quality of published randomised controlled trial protocols improved since the SPIRIT statement? A methodological study. BMJ open 2020;10(8):e038283. doi: 10.1136/bmjopen-2020-038283.

25. Rethlefsen ML, Schroter S, Bouter LM, Kirkham JJ, Moher D, Ayala AP, et al. Improving peer review of systematic reviews and related review types by involving librarians and information specialists as methodological peer reviewers: a randomised controlled trial. BMJ Evid Based Med 2025;30(4):241-49. doi: 10.1136/bmjebm-2024-113527.

Subgroup	Proportion of articles reporting p-value (%)		Odds of reporting p value (95% CI)	P value	Proportion of articles reporting effect-size (%)		Odds of reporting effect-size (95% CI)	P value
Publication year								
2008-2011	52 / 75 (69.3)		1.00		51 / 75 (68.0)		1.00	
2012-2015	82 / 119 (68.9)		0.98 (0.52, 1.83)	0.950	89 / 119 (74.8)		1.40 (0.74, 2.64)	0.305
2016-2018	59 / 103 (57.3)		0.59 (0.32, 1.11)	0.103	76 / 103 (73.8)		1.32 (0.69, 2.55)	0.400
Journal specialty								
Medicine	128 / 203 (63.1)		1.00		147 / 203 (72.4)		1.00	
Oncology	65 / 94 (69.1)		1.31 (0.78, 2.21)	0.306	69 / 94 (73.4)		1.05 (0.61, 1.82)	0.859
Senior author affiliation								
Non-USA	65 / 117 (55.6)		1.00		74 / 117 (63.2)		1.00	
USA	128 / 180 (71.1)		1.97 (1.21, 3.20)	0.006	142 / 180 (78.9)		2.17 (1.29, 3.65)	0.003
Any author in school of public health								
No	125 / 191 (65.4)		1.00		136 / 191 (71.2)		1.00	
Yes	68 / 106 (64.2)		0.94 (0.58, 1.55)	0.823	80 / 106 (75.5)		1.24 (0.72, 2.14)	0.429
Any author in epidemiology								
No	134 / 204 (65.7)		1.00		152 / 204 (74.5)		1.00	
Yes	59 / 93 (63.4)		0.91 (0.54, 1.51)	0.707	64 / 93 (68.8)		0.75 (0.44, 1.30)	0.308
Any author in statistics department								
No	143 / 230 (62.2)		1.00		162 / 230 (70.4)		1.00	
Yes	50 / 67 (74.6)		1.79 (0.97, 3.30)	0.062	54 / 67 (80.6)		1.74 (0.89, 3.40)	0.103

Table 1. Reporting quality of the trend analyses with linear model published in leading medicine and oncology journals, 2008-2018

	Proportion of reporting quality metrics, n /total by strata (%)											
Subgroup	Reporting P value	*P* value	Reporting effect size	*P* value	Reporting APC	*P* value	Reporting Beta/coefficient	*P* value	Score 0	Score 1	Score 2	*P* value*
Overall	169 / 226 (74.8)		183 / 226 (81.0)		94 / 226 (41.6)		34 / 226 (15.0)		13 / 226 (5.8)	119 / 226 (52.7)	94 / 226 (41.6)	
Publication year		0.352		0.203		0.082		0.823				0.801
2008-2011	41 / 56 (73.2)		41 / 56 (73.2)		30 / 56 (53.6)		8 / 56 (14.3)		2 / 56 (3.6)	28 / 56 (50.0)	26 / 56 (46.4)	
2012-2015	71 / 89 (79.8)		73 / 89 (82.0)		31 / 89 (34.8)		15 / 89 (16.9)		5 / 89 (5.6)	50 / 89 (56.2)	34 / 89 (38.2)	
2016-2018	57 / 81 (70.4)		69 / 81 (85.2)		33 / 81 (40.7)		11 / 81 (13.6)		6 / 81 (7.4)	41 / 81 (50.6)	34 / 81 (42.0)	
Journal specialty		0.724		0.326		0.025		0.911				0.108
Medicine	114 / 151 (75.5)		125 / 151 (82.8)		55 / 151 (36.4)		23 / 151 (15.2)		12 / 151 (7.9)	80 / 151 (53.0)	59 / 151 (39.1)	
Oncology	55 / 75 (73.3)		58 / 75 (77.3)		39 / 75 (52.0)		11 / 75 (14.7)		1 / 75 (1.3)	39 / 75 (52.0)	35 / 75 (46.7)	
Senior-author location		0.012		0.016		0.058		0.428				0.827
Non-USA	52 / 80 (65.0)		58 / 80 (72.5)		40 / 80 (50.0)		10 / 80 (12.5)		5 / 80 (6.3)	40 / 80 (50.0)	35 / 80 (43.8)	
USA	117 / 146 (80.1)		125 / 146 (85.6)		54 / 146 (37.0)		24 / 146 (16.4)		8 / 146 (5.5)	79 / 146 (54.1)	59 / 146 (40.4)	
Any author in school of public health		0.89		0.682		0.064		0.217				0.004
No	105 / 141 (74.5)		113 / 141 (80.1)		52 / 141 (36.9)		18 / 141 (12.8)		7 / 141 (5.0)	86 / 141 (61.0)	48 / 141 (34.0)	
Yes	64 / 85 (75.3)		70 / 85 (82.4)		42 / 85 (49.4)		16 / 85 (18.8)		6 / 85 (7.1)	33 / 85 (38.8)	46 / 85 (54.1)	
Any author in epidemiology department		0.163		0.028		0.015		0.497				0.684
No	115 / 148		126 / 148		53 / 148		24 / 148 (16.2)		8 / 148	81 / 148	59 / 148 (39.9)	

	(77.7)	(85.1)	(35.8)		(5.4)	(54.7)		
Yes	54 / 78 (69.2)	57 / 78 (73.1)	41 / 78 (52.6)	10 / 78 (12.8)	5 / 78 (6.4)	38 / 78 (48.7)	35 / 78 (44.9)	
Any author in statistics department	0.044	0.089	0.151	0.413				0.226
No	123 / 172 (71.5)	135 / 172 (78.5)	67 / 172 (39.0)	24 / 172 (14.0)	11 / 172 (6.4)	95 / 172 (55.2)	66 / 172 (38.4)	
Yes	46 / 54 (85.2)	48 / 54 (88.9)	27 / 54 (50.0)	10 / 54 (18.5)	2 / 54 (3.7)	24 / 54 (44.4)	28 / 54 (51.9)	

*Fisher exact test; APC, annual percent change.

Table 2. Multivariable models showing factors associated with reporting quality metrics of the trend analyses with linear model published in leading medicine and oncology journals, 2008-2018

Subgroup	Reporting P value OR (95% CI)	P value	Reporting effect size OR (95% CI)	P value	Reporting APC OR (95% CI)	P value
Publication year						
2008-2011					reference	
2012-2015					0.51 (0.25 - 1.05)	0.066
2016-2018					0.72 (0.35 - 1.51)	0.390
Journal specialty						
Medicine					reference	
Oncology					1.77 (0.97 - 3.22)	0.063
Senior author affiliation						
Non-USA	reference		reference		reference	
USA	2.17 (1.17 - 4.03)	0.014	1.93 (0.96 - 3.88)	0.065	0.80 (0.44 - 1.47)	0.475
Any author in school of public health						
No					reference	
Yes					1.67 (0.92 - 3.02)	0.090
Any author in epidemiology department						
No			reference		reference	
Yes			0.38 (0.18 - 0.80)	0.011	1.62 (0.90 - 2.91)	0.108
Any author in statistics department						
No	reference		reference			
Yes	2.29 (1.00 - 5.24)	0.051	3.28 (1.21 - 8.90)	0.020		

OR, Odds ratio; CI, confidence intervals.