Seed morphology and emergence variability in a conservation collection of *Pinus torreyana*

Lionel N Di Santo[1,4], Monica Polgar[1], Storm Nies[1], Paul Hodgkiss[2], Courtney A Canning,[3]

Jessica W Wright[2], and Jill A Hamilton[1]

[1] North Dakota State University, Department of Biological Sciences, Fargo, ND, USA

[2] USDA Forest Service, Pacific Southwest Research Station, Davis, CA, USA

[3] USDA Forest Service, Pacific Southwest Research Station, Placerville CA, USA

[4] Correspondence: lionel.disanto@ndsu.edu

ABSTRACT

PREMISE: Optimizing the amount of genetic diversity captured in seed collections is a long-

standing objective of ex situ conservation. Particularly for rare species where limited genetic

information is available, it poses a significant challenge. However, understanding the within and

among population distribution of trait variation within seed collections may provide a means to

approximate standing genetic variation and inform conservation efforts.

METHODS: We quantified seed morphological variation and seedling emergence both within

and among populations for existing seed collections of Torrey pine (*Pinus torreyana*), a critically

endangered pine endemic to California. Using a simulation-based approach, we used estimates of

within-population variance to estimate the number of maternal families required to captured 95%

of trait variation within each existing seed collection.

KEY RESULTS: On average, 21% and 22% of seed morphological variation in Torrey pine was

explained by population origin and maternal family within populations respectively. Despite

significant morphological differences, seedling emergence was similar across populations.

Simulations revealed that 80% and 68% of all maternal families within island and mainland seed

collection respectively needed to be resampled to ensure 95% of seed trait variation within

existing collections was captured.

CONCLUSIONS: Overall, these results suggest that substantial structure exists for seed

morphological traits both within and between populations. From a conservation perspective, this

indicates that to optimize genetic diversity captured in Torrey pine *ex situ* conservation

collections, maximizing the number of maternal families sampled within each population will be

necessary.

INTRODUCTION

Ex situ seed collections preserve species genetic diversity outside of their native range, providing the raw material for species reintroductions and germplasm to augment restoration (Guerrant Jr et al., 2013; Potter et al., 2017). An invaluable conservation resource, particularly for rare species, ex situ collections protect against biodiversity loss in the wild while preserving species evolutionary potential. Ensuring ex situ collections represent genetic variation found in natural populations is critical to both contemporary conservation and potential future restoration efforts (Schaal and Leverich, 2004; Basey et al., 2015). However, the cost and logistical constraints associated with seed collection pose a significant challenge. Given this challenge, identifying means to optimize ex situ sampling efforts are needed (Hoban and Schlarbaum, 2014; Di Santo and Hamilton, 2020). One approach may be to use the distribution of trait variation existing within contemporary ex situ collections as a proxy for quantifying standing genetic variation within and among populations. These data may then be used to optimize and target supplemental ex situ efforts. In this way, quantifying within and between population variation for traits largely considered heritable within existing ex situ population collections may be invaluable to optimizing future collections.

The distribution of heritable genetic variation estimated via common garden experiments (Weber and Kolb, 2014; Hamilton et al., 2017; Yoko et al., 2020) or genetic data (Zhang and Zhou, 2013; Hausman et al., 2014; Tamaki et al., 2018) can be used to quantify the distribution of

standing genetic variation. However, when common garden experiments or genetic data are

unavailable, quantifying trait variation within and among ex situ seed population collections may

provide a reasonable proxy for the distribution of genetic variation. Seed collections established

to preserve genetic variation, as well as to provide restoration or research material are numerous

(Guerrant Jr et al., 2013; Westengen et al., 2013; Etterson et al., 2016). Millions of seed

accessions of known provenance have been stored in gene banks internationally (FAO, 2010),

representing a large conservation and research resource. Although common garden experiments

are preferred when available, heritability of seed morphological traits and ease of access to seeds

through existing ex situ collections suggests assessment of morphological trait variation may

provide a reasonable potential proxy to quantifying the distribution of standing genetic variation.

Seed traits, such as seed size (H^2 = 0.47-0.99, h^2 = 0.19-0.56), relative embryo size (H^2 = 0.89-

0.99), seed shape (h^2 = 0.59-0.79), seed coat thickness (H^2 = 0.59), and embryo length (H^2 =

0.92-0.96) are often highly heritable (Pandey et al., 1994; Cober et al., 1997; Mera et al., 2004;

Hakim and Suyamto, 2017). Given these observations, variation in seed morphological traits

likely has a genetic basis, and may reflect standing genetic variation responding to natural

selection (Barrett and Schluter, 2008; Carlson et al., 2014). Particularly for rare species where

existing genetic data is often limited, the distribution of trait variation within existing seed

collections may provide a valuable resource, even if estimates of variation may not necessarily

be conservative. For these species, where the goal is to limit the loss of biodiversity while

preserving evolutionary potential, quantifying the distribution of morphological variation within

and among seed populations provides a timely approach to evaluating variation preserved in

collections. Leveraging existing seed morphological data from conservation collections may aid

in understanding the distribution of variability currently maintained ex situ, and may be used to

optimize supplemental conservation collections where needed.

Pinus torreyana Parry ex Carr. (Torrey pine), is one of the rarest pines in the world (Critchfield

and Little, 1966; Dusek, 1985), endemic to two discrete natural populations in California. Torrey

pine occupies one mainland population of approximately 6000 trees at the Torrey Pine State

Reserve in La Jolla, CA, and an island population of approximately 3000 reproductively mature

trees on Santa Rosa Island, CA one of the Channel Islands (Ledig and Conkle, 1983; Hamilton et

al., 2017) (Fig. 1). Listed as critically endangered by the IUCN (2020), Torrey pine is of critical

conservation concern due to multiple factors; including low population size (Franklin and

Santos, 2011; Hall and Brinkman, 2015), low genetic diversity (Ledig and Conkle, 1983;

Whittall et al., 2010), climate change, and environmentally and human-mediated disturbances

(Franklin and Santos, 2011; Hamilton et al., 2017). While in situ conservation has preserved the

whole of the species' range, with fewer than 10,000 reproductively mature individuals in native

populations, there are undeniable risks to population-level extirpation events. To mitigate

potential losses in the wild, conservation efforts have focused on preservation of seed ex situ.

While ex situ seed collections provide an invaluable conservation resource, they may be also

used to understand within and among population variation to ultimately inform future

conservation efforts.

In this study, we evaluate morphological trait variation in a large ex situ conservation collection

of Torrey pine seed sourced from the two native extant populations. Specifically, we quantify the

distribution of variation for 14 seed morphology traits and assess differences in emergence

between island and mainland seedlings. In addition, we use existing ex situ collection data to

provide supplemental population sampling guidance for future Torrey pine collections. For this

latter objective, we use simulations to estimate the number of maternal families required to

capture 95% of seed morphological variation existing in contemporary ex situ collections, for

both island and mainland population independently. This study provides baseline understanding

of population- and family-level variability for an extremely rare endemic pine, and uses these

data to inform population sampling and aid in optimizing conservation efforts and preservation

of evolutionary potential in this species.

MATERIAL AND METHODS

Cone collection and seed processing – Reproductively mature Torrey pine (*Pinus torreyana*

Parry ex. Carr) cones were collected as part of a large ex situ conservation collection between

June and July of 2017. Cones were collected from 157 trees on Santa Rosa Island (Channel

Islands National park), CA (island population) and 201 trees at Torrey Pine State Reserve in La

Jolla, CA (mainland population), representing the species entire native distribution (Fig. 1).

Where possible, we collected between five to ten cones per maternal tree at each location. Seeds,

organized by individual maternal tree, were then extracted from cones using a combination of

mallet and pliers and processed for inclusion in a long-term ex situ conservation collection (see

below).

Seed viability tests – A float test was used first to quickly assess seed viability within the ex situ

collection. Floating seeds were presumed to lack an endosperm or embryo, while seeds that sunk

were presumed filled. Seeds were dropped into water for a period of approximately 15 seconds to

differentiate presumed non-viable, floating seed from presumed viable, sinking seeds (Gribko

and Jones, 1995; Morina et al., 2017). Those seeds classed as likely viable were organized by

maternal tree using paper bags, and then placed in a Blue M drying oven (Thermal Product

Solutions, White Deer, Pennsylvania, USA) maintained at 37°C for 24 hours to remove potential

surface moisture. Following this, x-ray photographs were taken to visually assess viability and

morphological variation of each seed and its contents at the Placerville Nursery, CA. Acrylic

seed trays [20.3 cm x 25.4 cm x 0.48 cm], with a 9 x 11 array of wells, were used to separate and

position each Torrey pine seed over the x-ray film. Kodak x-OMAT HBT film (20.3 cm x 25.4

cm) was placed in a lightproof x-ray film cassette which was positioned in the x-ray machine

with the seed tray centered on top of the film, with a shelf height of 55.9 cm. The x-ray was

taken using a 17 kVP exposure for a total of two minutes, based on standardized conditions

established previously for *Pinus coulteri* (Sara Wilson, USDA Forest Service, personal

communication). X-ray images were digitized using a Nikon D40 digital camera mounted on a

tripod over a light box. Because size standards were internal to each image (see below), no

additional calibrations or corrections were required.

Morphological measurement of seed traits – Using ImageJ (Abràmoff et al., 2004), eight seed

morphological traits were measured across 80 mainland maternal families and 30 island maternal

families, representing a haphazard subset of the complete collection (Fig. 2; Table 1). Each x-ray

picture was scaled using the diameter of a seed tray well (1.87 cm) to express pixels as trait

values in centimeters. Directly measured seed traits included seed length (SL, cm), seed width

(SW, cm), embryo length (EL, cm), embryo width (EW, cm), seed coat width (SCW, cm), seed

area (SA, cm^2), endosperm area (ESA, cm^2), and embryo area (EA, cm^2). Using these traits, six

additional traits were derived (Table 1); including seed length/width ratio (SLW), embryo

length/width ratio (ELW), relative embryo size (RES), relative endosperm size (REndS), seed

coat area (SCA, cm^2), and relative seed coat size (RSCS). We measured five randomly selected

seeds per maternal tree, including three technical replications per seed for each trait, with the

exception of seven mainland (TPSR 442: 30 seeds, TPSR 457: 20 seeds, TPSR 477: 15 seeds,

TPSR 1176: 23 seeds, TPSR 3005: 20 seeds, TPSR 3202: 10 seeds, and TPSR 3204: 10 seeds)

and seven island maternal trees (SRI 143: 10 seeds, SRI 243: 10 seeds, SRI 1088: 10 seeds, SRI

1136: 10 seeds, SRI 2028: 9 seeds, SRI 2048: 10 seeds, and SRI 2124: 10 seeds). Measurements

were averaged across technical replicates to summarize the mean trait value per seed. In total,

677 seeds were measured from across 110 maternal trees spanning the two Torrey pine

populations.

Seed emergence test – In January 2018, a seed emergence trial was conducted using a random

subset of seeds from the ex situ collection, including seeds sourced from Torrey Pine State

Reserve and Santa Rosa Island, CA. Following x-ray, seeds were stored at 4°C in sealed mylar

bags (USA emergency supply, Beaverton, Oregon, USA) placed in plastic boxes, each box

contained desiccant crystals to decrease ambient moisture and reduce likelihood of mold. Seeds

from eight maternal families per population were selected for the emergence trial. Between eight

to ten seeds per maternal tree were weighed and then stratified under cold, moist conditions for

30 days (placed in plastic boxes on a moist paper towel at 4°C). Seeds were sown directly into a

164 ml Ray Leach "Cone-tainer" ™ (Stuewe & Sons, Tangent, Oregon, USA) filled with

Sunshine® Mix #4 (Sungro horticulture, Agawam, Massachusetts, USA), pressed halfway into

the soil, and then covered with a thin layer of gravel. For approximately one month following

planting, seeds were misted for one minute at hourly intervals over a daily eight-hour period

(9am – 4pm). Following emergence, seedlings were hand watered to saturation weekly to

biweekly as needed. Emergence was quantified across three separate timepoints (02/06/2018,

02/16/2018 and 03/07/2018) per maternal family as the proportion of seeds that had emerged

from the total initially planted.

Evaluating the distribution of seed trait variation – We conducted a principal component

analysis (PCA) using all 14 measured and derived seed traits averaged by maternal family to

evaluate population-specific differentiation in seed morphology. Prior to performing the PCA, to

account for differences in measurement units, all seed traits were standardized using the *scale()*

function in R implementing the z-score standardization: $z_{ij} = \frac{x_{ij} - \mu_j}{\sigma_j}$ (1), where x_{ij} is the non-

transformed trait value, μ_j is the mean of a given seed trait across populations, and σ_j the

standard deviation of the same seed trait across populations. Subsequently, to test for seed trait

differences between population means, we used either Student's two-sample test or its non-

parametric equivalent when normality was not met, Wilcoxon's two-sample test, within the R

package "exactRankTests" (Hothorn and Hornik, 2019). Normality was assessed using Shapiro-

Wilk's test of normality within each population. In total, five of the fourteen traits were

distributed normally in both Torrey pine populations, including seed width (mainland: W = 0.97,

P = 0.11; island: W = 0.97, P = 0.57), embryo length (mainland: W = 0.98, P = 0.28; island: W =

0.96, P = 0.37), embryo area (mainland: W = 0.98, P = 0.46; island: W = 0.97, P = 0.59), seed

length/seed width ratio (mainland: W = 0.99, P = 0.5; island: W = 0.98, P = 0.75), and embryo

length/embryo width ratio (mainland: W = 0.97, P = 0.05; island: W = 0.97, P = 0.67).

To evaluate the distribution of morphological trait variation within and between Torrey pine

populations, we quantified the proportion of variation attributed to population and maternal tree

families using measured and derived morphological traits summarized by seeds. For each trait,

we fit a linear mixed model using the R package "lme4" (Bates et al., 2014) with population

considered a fixed effect and maternal families within populations considered a nested random

effect: $Y_{ij} = \mu + \pi_i + r_{i/j} + e_{ij}$, where Y_{ij} is the observed seed trait value, μ is the seed trait

overall mean, π_i is the effect of population origin on the seed trait mean, $r_{i/j}$ is the effect of

maternal family within populations on the observed seed trait value, and e_{ij} are the effects on the

seed trait value of any other variables unaccounted for in the model (residual error). For each

model, normality of residual errors was visually assessed and significance of fixed- and random-

effect terms was tested using the functions *anova()* and *ranova()* respectively, implemented in

the R package "stats" (R Core Team 2020) and "lmerTest" (Kuznetsova et al., 2016).

Proportions of seed morphological variance explained by populations (marginal R^2, R^2_m), both

populations and maternal families (conditional R^2, R^2_c), and maternal families alone ($R^2_c - R^2_m$)

were quantified for each model independently using the function *r.squaredGLMM()*

implemented in the R package "MuMIn" (Bartoń, 2020).

Assessing differences in seedling emergence across populations – To test for differences in the

probability and the timing of seed emergence in Torrey pine, we evaluated the proportion of

seedling emergence both within and between populations across timepoints. First, we used

Friedman's rank sum test (non-parametric repeated measures ANOVA) followed by Wilcoxon'

paired two-sample test, both implemented in the R package "rstatix" (Kassambara, 2020), to

assess differences in the proportion of emerged seeds between timepoints within populations. We

used a non-parametric approach for both Torrey pine populations because normality could not be

assumed at select timepoints due to high frequency of zero values. We accounted for multiple

testing using Benjamini & Hochberg (1995) False Discovery Rate (FDR) correction

implemented in the *wilcox_test()* function. Following this, we evaluated timepoint-specific

population differences in emergence. We used Shapiro-Wilk's test to assess populations'

deviation from normality at each timepoint and either Student's (for timepoints passing the

normality test) or Wilcoxon's two-sample test (for timepoints failing the normality test) to

evaluate differences in population emergence. Timepoints 02/16/18 (mainland: $W = 0.94$, $P =$

0.57; island: $W = 0.9$, $P = 0.28$) and 03/07/2018 (mainland: $W = 0.96$, $P = 0.85$; island: $W =$

0.88, $P = 0.21$) passed the normality test, while timepoint 02/06/2018 (mainland: $W = 0.52$, $P <$

0.001; island: $W = 0.73$, $P = 0.004$) failed the normality test. All statistical analyses were

performed using R version 4.0.2 (R Core Team 2020).

Simulating variation captured in* ex situ *collection using seed morphological traits – For each

of the 14 measured and derived seed traits, we conducted a separate simulation quantifying

morphological variation captured when increasing the number of maternal families sampled from

contemporary Torrey pine seed collections. Simulations were conducted in R using a customized

script (Appendix S3). Resampling of ex situ collections were performed for island and mainland

Torrey pine populations independently, using between one and the total number of maternal

families available within each ex situ population collection (mainland: 80 maternal families,

island: 30 maternal families) (N_{fam}). Maternal trees were sampled randomly without replacement

from the pool of available families. All seeds within each selected maternal family were sampled

as part of this simulation, except those with missing values for the trait simulated.

To evaluate the number of maternal families needed to capture 95% of seed trait variation in both

island and mainland populations, we estimated the number of unique seed trait values captured in

a sample of N_{fam} maternal families (N_c) relative to the total number of unique seed trait values

present in a seed population (N_t). Here, we define "unique seed trait values" as the number of

non-redundant standardized measurements for the seed trait simulated rounded to the first digit.

Seed morphological measurements were rounded to the first digits as we believe that seed trait

variation estimated using additional digits is more likely to fail to capture meaningful biological

variation. Standardization of the data was performed so that all seed traits share the same unit

(the number of standard deviations a value is from the overall trait mean across populations, see

equation (1) above) and become comparable. Sampling of maternal families and estimation of

the summary statistic, defined as the proportion of total seed trait variance captured (N_c/N_t), were

repeated 500 times for each seed morphological trait and Torrey pine population to account for

the variation introduced through randomly sampling maternal families.

Finally, for each number of maternal trees sampled (N_{fam}), we averaged the summary statistic

across all 500 replicates. This process was repeated for each of the 14 seed morphological traits

and performed for each Torrey pine population separately. Following this, the summary statistic

was averaged across all seed traits and separated by populations (see Results below). Proportions

of total seed trait variance captured (N_c/N_t) are provided based on proportions of maternal

families sampled (instead of the number of maternal families sampled) as sample sizes varied

across Torrey pine populations.

RESULTS

Island-mainland differentiation in seed morphology – A principal component analysis (PCA)

using all 14 measured and derived seed traits revealed substantial differences in seed

morphology between island and mainland populations of Torrey pine (Fig. 3). The first PC axis

explained 57.5% of variation in seed morphological traits, primarily separating the island from

the mainland population. Seed length, seed width, seed area, endosperm area, and seed coat area

exhibited the five highest loadings (absolute values) on PC1 (Appendix S1; see Supplemental

data with this article), indicating that seed size and seed coat thickness can largely discriminate

island from mainland individuals. On average, seeds collected on island trees were longer, wider,

larger and thicker than seeds collected on mainland trees (Table 1). The second PC axis

explained 16.5% of seed trait variation and summarizes within population variability in seed

morphology (Fig. 3). Relative seed coat size, relative endosperm size, and relative embryo size

had the three highest loadings (absolute values) on PC2 (Appendix S1). This suggests that once

corrected for seed size, seed coat thickness, endosperm size, and embryo size are traits

contributing to within-population variation.

Contribution of population origin and maternal family to seed trait variation – Consistent with

our principal component analysis, linear mixed models constructed for each of the 14 measured

and derived seed traits demonstrated that considerable variation in seed morphology in Torrey

pine is explained by population origin (Fig. 4). On average, population origin explained 21%

(range = 0.04–0.52) of variation across the species' distribution (Appendix S2). Traits associated

with seed size and seed coat thickness exhibited the highest proportion of variance explained by

population origin. These include seed coat area (0.52; $F = 202.5$, $df = 1$, $P < 0.001$), seed area

(0.45; F = 143.21, df = 1, P < 0.001), seed width (0.36; F = 130.71, df = 1, P < 0.001),

endosperm area (0.34; F = 95.47, df = 1, P < 0.001), seed coat width (0.29; F = 87.54, df = 1, P <

0.001), and seed length (0.28; F = 76.18, df = 1, P < 0.001). Overall, this suggests seed size and

seed coat thickness are major discriminants of island and mainland Torrey pine seeds.

While population origin explained substantial variation across populations, assessment of

maternal seed families within populations indicated substantial family structure to seed trait

variation (Fig. 4). On average, maternal seed family explained 22% (range = 0.05–0.34) of

variation within populations (Appendix S2). Endosperm area (0.34; χ^2 = 272.03, df = 1, P <

0.001), seed length (0.33; χ^2 = 217.24, df = 1, P < 0.001), seed area (0.3; χ^2 = 288.62, df = 1, P

< 0.001), seed coat width (0.29; χ^2 = 180.45, df = 1, P < 0.001), and embryo area (0.29; χ^2 =

142.9, df = 1, P < 0.001) exhibited the highest proportion of seed trait variation explained by

within-population maternal families. This suggests that there is substantial family-level structure

to seed size, endosperm size, embryo size and seed coat thickness within Torrey pine populations

Impact of population seed trait differentiation on seed emergence – The proportion of emerged

seeds increased over time for both island (Q = 15.5, df = 2, P < 0.001) and mainland (Q = 15.2,

df = 2, P < 0.001) populations (Fig. 5). However, we found no significant differences in the

proportion of individuals emerging between populations across observed time points. On

average, 7% and 9% of mainland and island seeds emerged a month after sowing (02/06/18; W =

28, P = 0.64), 63% and 53% of mainland and island seeds emerged a month and a half after

sowing (02/16/18; t = 0.81, df = 14, P = 0.43), and 78% of mainland and island seeds emerged

two months after sowing (03/07/18; t = −0.06, df = 14, P = 0.95). Overall, this indicates that

under controlled conditions, the timing and probability of emergence for Torrey pine is not

impacted by population differences in seed morphology. Alternatively, this may also suggest that

island and mainland seeds respond similarly to artificial germination protocol.

Morphological variation captured in simulated seed collections – Simulations revealed that to

capture 95% of seed trait variation present in our existing ex situ collections, on average 80% (24

out of 30) and 68% (54 out of 80) of all island and mainland families would need to be

resampled respectively (Fig. 6). This indicates that both island and mainland populations harbor

considerable within-population structure for seed morphological traits. Interestingly, capturing

equal morphological variation across seed collections always required a higher proportion of

island maternal families to be collected relative to the mainland population. This suggests that

the island population may exhibit higher variability in seed morphology compared to the

mainland population, and thus may be worth of additional sampling efforts.

DISCUSSION

Evaluating between- and within-population morphological variation in contemporary ex situ seed

collections may aid in understanding the distribution of variation needed to guide future

conservation efforts. Here, we quantified the distribution of trait variation within an ex situ

collection of Torrey pine seeds, with an aim to optimize future supplemental ex situ collections.

Morphologically, island and mainland seeds were significantly different from each other. Island

seeds were larger on average and had thicker seed coats relative to their mainland counterparts.

These morphological differences may be explained by a combination of stochastic and

deterministic factors associated with population origin. Interestingly, despite substantial

morphological differentiation, emergence did not vary among populations, suggesting that either

the probability and timing of emergence under controlled conditions is not impacted by

differences in seed morphology or that island and mainland seeds respond similarly to an

artificial germination protocol. In addition to population origin, a considerable proportion of seed

trait variation within each population was explained by maternal family. This suggests that there

is substantial within-population variation that will be important to conserve and maintain

populations' evolutionary potential. Finally, simulations demonstrated that 80% and 68% of all

maternal families within island and mainland seed collection respectively were necessary to

capture 95% of morphological variation within the existing collections. To ensure we maintain

representative levels of trait variation in future seed collections, our results thus suggest that the

number of maternal families sampled within natural populations should be maximized, with the

island population potentially requiring additional sampling efforts.

Species distributed between island and mainland origins often exhibit marked among-population

phenotypic differentiation, including differentiation in seed morphology (Burns et al., 2012; Lens

et al., 2013; Kavanagh and Burns, 2014; Burns, 2016). Our results revealed considerable seed

morphological differences between Torrey pine populations (Fig. 3), primarily in seed size and

seed coat thickness (Fig. 4; Appendix S1). On average, islands seeds were larger and had thicker

seed coats than seeds collected on the mainland (Table 1). These results are consistent with

previous studies of island-mainland systems that noted island populations exhibited larger seeds

relative to mainland populations (Kavanagh and Burns, 2014; Burns, 2016). A combination of

different factors could contribute to morphological variation among seed populations, including

both stochastic and deterministic forces.

On islands, seeds traits associated with long-distance dispersal may be selected against (Cody

and Overton, 1996; Schenk, 2013; Kavanagh and Burns, 2014), as they could increase the

probability an individual would disperse beyond the islands' limits. For potentially

anemochorous species like Torrey pine, increased seed size on islands may limit potential losses

via wind-dispersal and promote zoochory (Greene and Johnson, 1993; Reader, 1993; Benkman,

1995; Gómez, 2004). Seed predation may also have contributed to differences in seed size

between populations. On the island, *Peromyscus maniculatus* (island deer mouse) is the only

small mammal seed predator alongside fewer than a dozen bird species

(https://www.nps.gov/chis/learn/nature/birds.htm;

https://www.nps.gov/chis/learn/nature/terrestrial-animals.htm). This contrasts with the mainland,

where already 11 mammalian seed predators exist; including squirrels, rats, mice, and gophers

(https://torreypine.org/nature-center/mammals/). If large seeds are preferentially targeted by

seed-eating animals (Reader, 1993; Gómez, 2004), smaller seed size on the mainland may have

evolved in response to increased predation pressures from granivorous species.

While selection may contribute to population differences, differentiation in seed morphology

may result from stochastic evolutionary forces. Founder effects associated with the colonization

of Santa Rosa island by mainland individuals, as well as genetic drift combined with limited

gene flow may have led to seed morphological differentiation between Torrey pine populations

(Ledig and Conkle, 1983). While these factors may contribute to population differences in seed

morphology, further experimentation is needed to adequately test these hypotheses. As seeds

were collected from natural populations, measuring seeds produced by maternal plants grown in

a common environment is required to tease apart the contribution of environment and genetics to

seed trait differences observed among populations. Additionally, a reciprocal transplant

experiment would be the most effective test of the action of natural selection in shaping

morphological differences between island and mainland seeds.

Despite significant differences in seed morphology between Torrey pine populations, timing and

probability of emergence were similar across populations (Fig. 5). Emergence rates were high

throughout the trial, with 78% of island and mainland seeds germinating two months after

sowing. The absence of differences in emergence between populations was surprising, as seed

size often negatively correlates with time to emergence (Daws et al., 2005; Tanveer et al., 2013).

However, seed coat thickness can also influence rates of emergence. Seeds with thick seed coats

relative to their mass often emerge later than seeds with thinner seed coats (Daws et al., 2005;

Hamilton et al., 2013). Interestingly, island seeds were not only larger, but also had thicker seed

coats relative to mainland seeds (Table 1). Even after correcting for differences in seed size, seed

coat thickness (RSCS) remained moderately higher in island seeds. Together, these results

predict that island seed could emerge at similar timepoints as mainland seed, or later than seeds

collected on the mainland (Hamilton et al., 2017), which is consistent with our observations.

In addition, similar emergence rates may result from our experimental design. Abe & Matsunaga

(2011), in a mainland-island comparison study, observed that cold stratification attenuates

differences in seed emergence between populations of *Rhaphiolepis umbellata*. Additionally,

complete and rapid germination of pine seeds is generally observed when pretreated under cold

and moist conditions (Bonner and Karrfalt, 2008). Overall, this suggests that cold stratification

may mask population-specific differences in seedling emergence. Concurrent emergence of

seeds from both Torrey pine populations coupled with high emergence success suggests a cold

stratification protocol is valuable for Torrey pine, particularly where simultaneous emergence for

nursery-grown seedlings is desired.

Although population origin explained a substantial proportion of seed trait variation, linear

mixed models demonstrated that maternal seed families within populations explained nearly as

much variation (Fig. 4). Given generally high heritability for seed morphological traits and the

half-sib design of our collection (Pandey et al., 1994; Cober et al., 1997; Mera et al., 2004;

Hakim and Suyamto, 2017), family-level seed trait variation likely provides a useful proxy for

assessing within-population genetic diversity. With more than a fifth of variation explained on

average by maternal families (Appendix S2), this suggest there is substantial genetic structure

within Torrey pine populations. These results were surprising, as previous studies using

allozymes and chloroplast DNA both suggested that the species exhibits little to no within-

population genetic variability (Ledig and Conkle, 1983; Waters and Schaal, 1991; Whittall et al.,

2010). However, the common garden experiment conducted by Hamilton and colleagues (2017)

indicated substantial family-level variation in tree height within both island and mainland

populations. Overall, these results indicate that Torrey pine populations may possess within-

population genetic variation necessary for natural selection to act upon. From a conservation

perspective, these findings suggest that a strategy maximizing the number of maternal families

sampled would optimize genetic diversity preserved in future ex situ seed collections.

Generally, ex situ seed collections aim to capture 95% of genetic diversity present throughout a

species' distribution (Marshall and Brown, 1975; Brown and Marshall, 1995; Li et al., 2002;

Gapare et al., 2008). Simulations revealed that, in order to captured 95% of morphological

variation currently maintained ex situ, 24 (80% of island collection) and 54 (68% of mainland

collection) maternal families within each seed collection would need to be resampled (Fig. 6).

These data indicate that sampling more maternal families from the island population may be

necessary to achieve the same level of representation of morphological variation. If we presume

that these phenotypes result from the combination of different alleles, capturing 95% of genetic

variation within the island population will always require more maternal families relative to the

mainland population. For these simulations, we assumed that contemporary ex situ collections

captured all morphological variation both within and between populations, including seed

phenotype frequencies. However, if this is not the case, these recommendations may result in

suboptimal sampling of standing variation within targeted populations. This caveat is all the

more important because sample sizes differed between island (30 maternal families) and

mainland (80 maternal families) seed collections. An understanding of the breadth of variation

within native populations captured in contemporary ex situ seed collections would be necessary

to fully optimize recommendations regarding within-population sampling efforts using this

approach.

CONCLUSION

Practical and cost-effective, long-term storage of seeds ex situ is widely used to capture and

maintain rare species genetic diversity. These seed collections represent an invaluable resource

to estimate the distribution of seed trait variation within and between populations that may be

used to guide future ex situ sampling efforts. This research leverages ex situ seed collections of

Torrey pine to understand existing variation within and among populations and inform sampling

for future seed collections. Incorporating existing information from ex situ collections provides a

unique opportunity to optimize conservation objectives, particularly important for rare species.

ACKNOWLEDGEMENTS

The authors thank Annette Delfino Mix, Valerie Gallup, Andrew Bower, Drew Peterson, Emma

Ordemann, Conner Harrington, Forest Swaciak, Jill Wulf, and Stephen Johnson for their help

collecting the Torrey pine cones. The collections were funded by The USDA Forest Service,

State and Private Forestry, Gene Conservation program to J.W.W. and J.A.H. Sara Wilson

provided training and assistance for the use of the x-ray machine. We also thank the USDA

Forest Service, Pacific Southwest Research Station for their support of this project. This work

was also supported by a new faculty award from the office of the North Dakota Experimental

Program to Stimulate Competitive Research (ND-EPSCoR NSF-IIA-1355466) and funding from

the NDSU Environmental and Conservation Sciences Program to JAH. Any use of product

names is for information purposes only and does not imply endorsement by the US Government.

DATA AVAILABILITY

The data for this article, including seed morphological measurements and the integrality of R

scripts used are available from GitHub: https://github.com/LionelDiSanto/Di-Santo-et-al-2020

SUPPORTING INFORMATION

Supplementary information to this article may be found in the supporting information tab.

LITERATURE CITED

Abe, T., and M. Matsunaga. 2011. Geographic variation in germination traits in Melia azedarach

and Rhaphiolepis umbellata. *American Journal of Plant Sciences* 2: 52.

Abràmoff, M. D., P. J. Magalhães, and S. J. Ram. 2004. Image processing with ImageJ.

Biophotonics international 11: 36–42.

Barrett, R. D. H., and D. Schluter. 2008. Adaptation from standing genetic variation. *Trends in*

ecology & evolution 23: 38–44.

Bartoń, K. 2020. MuMIn: Multi-Model Inference. R package version 1.43.17.

Basey, A. C., J. B. Fant, and A. T. Kramer. 2015. Producing native plant materials for

restoration: 10 rules to collect and maintain genetic diversity. *Native Plants Journal* 16: 37–

53.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2014. Fitting linear mixed-effects models using

lme4. arXiv. org. *preprint]. doi* 10.

Benjamini, Y., and Y. Hochberg. 1995. Controlling the false discovery rate: a practical and

powerful approach to multiple testing. *Journal of the Royal statistical society: series B*

(Methodological) 57: 289–300.

Benkman, C. W. 1995. Wind dispersal capacity of pine seeds and the evolution of different seed

dispersal modes in pines. *Oikos*: 221–224.

Bonner, F. T., and R. P. Karrfalt. 2008. The woody plant seed manual. *Agric. Handbook No. 727.*

Washington, DC. US Department of Agriculture, Forest Service. 1223 p. 727.

Brown, A. H. D., and D. R. Marshall. 1995. A basic sampling strategy: theory and practice.

Collecting plant genetic diversity: technical guidelines. CAB International, Wallingford:

75–91.

Burns, K. C. 2016. Size changes in island plants: independent trait evolution in Alyxia ruscifolia

(Apocynaceae) on Lord Howe Island. *Biological Journal of the Linnean Society* 119: 847–

855.

Burns, K. C., N. Herold, and B. Wallace. 2012. Evolutionary size changes in plants of the

south☐west Pacific. *Global Ecology and Biogeography* 21: 819–828.

Carlson, S. M., C. J. Cunningham, and P. A. H. Westley. 2014. Evolutionary rescue in a

changing world. *Trends in Ecology and Evolution* 29: 521–530.

Cober, E. R., H. D. Voldeng, and J. A. Fregeau-Reid. 1997. Heritability of seed shape and seed

size in soybean. *Crop Science* 37: 1767–1769.

Cody, M. L., and J. M. Overton. 1996. Short-term evolution of reduced dispersal in island plant

populations. *Journal of Ecology*: 53–61.

Critchfield, W. B., and E. L. Little. 1966. Geographic distribution of the pines of the world. US

Department of Agriculture, Forest Service.

Daws, M. I., N. C. Garwood, and H. W. Pritchard. 2005. Traits of recalcitrant seeds in a

semi☐deciduous tropical forest in Panamá: some ecological implications. *Functional*

Ecology 19: 874–885.

Dusek, K. H. 1985. Update on our rarest pine. *American forests (USA)*.

Etterson, J. R., S. J. Franks, S. J. Mazer, R. G. Shaw, N. L. Soper Gorden, H. E. Schneider, J. J.

Weber, et al. 2016. Project baseline: An unprecedented resource to study plant evolution

across space and time. *American Journal of Botany* 103: 164–173.

FAO. 2010. The second report on the state of the world's plant genetic resources for food and

agriculture. Food & Agriculture Org, Rome.

Franklin, J., and E. V Santos. 2011. A spatially explicit census reveals population structure and

recruitment patterns for narrowly endemic pine, Pinus torreyana. *Plant Ecology* 212: 293–

306.

Gapare, W. J., A. D. Yanchuk, and S. N. Aitken. 2008. Optimal sampling strategies for capture

of genetic diversity differ between core and peripheral populations of Picea sitchensis

(Bong.) Carr. *Conservation Genetics* 9: 411–418.

Gómez, J. M. 2004. Bigger is not always better: conflicting selective pressures on seed size in

Quercus ilex. *Evolution* 58: 71–80.

Greene, D. F., and E. A. Johnson. 1993. Seed mass and dispersal capacity in wind-dispersed

diaspores. *Oikos*: 69–74.

Gribko, L. S., and W. E. Jones. 1995. Test of the float method of assessing northern red oak

acorn condition. *Tree Planters' Notes* 46: 143–147.

Guerrant Jr, E. O., K. Havens, and P. Vitt. 2013. Sampling for effective ex situ plant

conservation. *International Journal of Plant Sciences* 175: 11–20.

Hakim, L., and S. Suyamto. 2017. Gene action and heritability estimates of quantitative

characters among lines derived from varietal crosses of soybean. *Indonesian Journal of*

Agricultural Science 18: 25–32.

Hall, T., and A. Brinkman. 2015. Population Dynamics of the Island Torrey Pine (Pinus

torreyana ssp. insularis) on Santa Rosa Island, CA. California State University Chanel

Islands.

Hamilton, J. A., R. Royauté, J. W. Wright, P. Hodgskiss, and F. T. Ledig. 2017. Genetic

conservation and management of the California endemic, Torrey pine (Pinus torreyana

Parry): Implications of genetic rescue in a genetically depauperate species. *Ecology and*

Evolution.

Hamilton, K. N., C. A. Offord, P. Cuneo, and M. A. Deseo. 2013. A comparative study of seed

morphology in relation to desiccation tolerance and other physiological responses in 71

Eastern Australian rainforest species. *Plant Species Biology* 28: 51–62.

Hausman, C. E., M. M. Bertke, J. F. Jaeger, and O. J. Rocha. 2014. Genetic structure of green

ash (Fraxinus pennsylvanica): implications for the establishment of ex situ conservation

protocols in light of the invasion of the emerald ash borer. *Plant Genetic Resources* 12:

286–297.

Hoban, S., and S. Schlarbaum. 2014. Optimal sampling of seeds from plant populations for ex-

situ conservation of genetic biodiversity, considering realistic population structure.

Biological Conservation 177: 90–99.

Hothorn, T., and K. Hornik. 2019. exactRankTests: Exact Distributions for Rank and

Permutation Tests. R package version 0.8-31.

IUCN 2020. The IUCN Red List of Threatened Species. Version 2020-2.

https://www.iucnredlist.org. Downloaded on 09 July 2020.

Kassambara, A. 2020. rstatix: Pipe-Friendly Framework for Basic Statistical Tests. R package

version 0.4.0.

Kavanagh, P. H., and K. C. Burns. 2014. The repeated evolution of large seeds on islands.

Proceedings of the Royal Society B: Biological Sciences 281: 20140675.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2016. Tests in linear mixed effects

models version. *Cran. URL https://cran. r-project. org/web/packages/lmerTest/lmerTest.*

pdf.

Ledig, F. T., and M. T. Conkle. 1983. Gene diversity and genetic structure in a narrow endemic,

Torrey Pine (Pinus torreyana Parry ex Carr.). *Evolution* 37: 79–85.

560 Lens, F., N. Davin, E. Smets, and M. del Arco. 2013. Insular woodiness on the Canary Islands: a

561   remarkable case of convergent evolution. *International Journal of Plant Sciences* 174: 992–

562   1013.

563 Li, Q., Z. Xu, and T. He. 2002. Ex situ genetic conservation of endangered Vatica guangxiensis

564   (Dipterocarpaceae) in China. *Biological Conservation* 106: 151–156.

565 Marshall, D. R., and A. H. D. Brown. 1975. Optimum sampling strategies in genetic

566   conservation. *In* O. H. Frankel, and J. G. Hawkes [eds.], Crop Genetic Resources for Today

567   and Tomorrow, 53–80. Cambridge University Press, Cambridge.

568 Mera, M., R. Jerez, H. Miranda, and J. L. Rouanet. 2004. Seed coat specific weight in Lupinus

569   angustifolius: influence of genotype and environment and relationship with seed coat

570   proportion. *Australian journal of agricultural research* 55: 1189–1195.

571 Morina, D. L., M. A. Lashley, M. C. Chitwood, C. E. Moorman, and C. S. DePerno. 2017.

572   Should we use the float test to quantify acorn viability? *Wildlife Society Bulletin* 41: 776–

573   779.

574 Pandey, M. P., D. V Seshu, and M. Akbar. 1994. Genetics of embryo size and its relationship

575   with seed and seedling vigour in rice (Oryza sativa L.). *Indian Journal of Genetics and*

576   *Plant Breeding* 54: 68.

577 Potter, K. M., R. M. Jetton, A. Bower, D. F. Jacobs, G. Man, V. D. Hipkins, and M. Westwood.

578   2017. Banking on the future: progress, challenges and opportunities for the genetic

579   conservation of forest trees.

580 R Core Team (2020). R: A language and environment for statistical computing. R

581   Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

582 Reader, R. J. 1993. Control of seedling emergence by ground cover and seed predation in

relation to seed size for some old-field species. *Journal of Ecology*: 169–175.

Di Santo, L. N., and J. A. Hamilton. 2020. Using environmental and geographic data to optimize

ex situ collections and preserve evolutionary potential. *Conservation Biology*.

Schaal, B., and W. Leverich. 2004. Population genetic issues in ex situ plant conservation. Island

Press: Washington, DC, USA.

Schenk, J. J. 2013. Evolution of limited seed dispersal ability on gypsum islands. *American*

journal of botany 100: 1811–1822.

Tamaki, S., K. Isoda, M. Takahashi, H. Yamada, and Y. Yamashita. 2018. Genetic structure and

diversity in relation to the recently reduced population size of the rare conifer, Pseudotsuga

japonica, endemic to Japan. *Conservation Genetics*: 1–13.

Tanveer, A., M. Tasneem, A. Khaliq, M. M. Javaid, and M. N. Chaudhry. 2013. Influence of

seed size and ecological factors on the germination and emergence of field bindweed

(Convolvulus arvensis). *Planta daninha* 31: 39–51.

Waters, E. R., and B. A. Schaal. 1991. No variation is detected in the chloroplast genome of

Pinus torreyana. *Canadian Journal of Forest Research* 21: 1832–1835.

Weber, A., and A. Kolb. 2014. Differences in heritable trait variation among populations of

varying size in the perennial herb Phyteuma spicatum. *Conservation genetics* 15: 1329–

1337.

Westengen, O. T., S. Jeppson, and L. Guarino. 2013. Global ex-situ crop diversity conservation

and the Svalbard Global Seed Vault: Assessing the current status. *PloS one* 8: e64146.

Whittall, J. B., J. Syring, M. Parks, J. Buenrostro, C. Dick, A. Liston, and R. Cronn. 2010.

Finding a (pine) needle in a haystack: chloroplast genome sequence divergence in rare

widespread pines. *Molecular Ecology* 19: 100–114.

Yoko, Z. G., K. L. Volk, N. A. Dochtermann, and J. A. Hamilton. 2020. The importance of

quantitative trait differentiation in restoration: landscape heterogeneity and functional traits

inform seed transfer guidelines. *AoB Plants* 12: plaa009.

Zhang, D.-Q., and N. Zhou. 2013. Genetic diversity and population structure of the endangered

conifer Taxus wallichiana var. mairei (Taxaceae) revealed by Simple Sequence Repeat

(SSR) markers. *Biochemical systematics and ecology* 49: 107–114.

TABLES

Table 1. Morphological measurements of Torrey pine seeds sourced from Santa Rosa Island (Island, n=30) and Torrey Pine State

Reserve (Mainland, n=80), CA. Listed are measured (A) and derived (B) seed trait values averaged over maternal families given per

populations ($\pm$SE). Measurable traits: seed length (SL), seed width (SW), embryo length (EL), embryo width (EW), seed coat width

(SCW), seed area (SA), endosperm area (ESA), and embryo area (EA). Derived traits: seed length/width ratio (SLW), embryo

length/width ratio (ELW), relative embryo size (RES), relative endosperm size (REndS), seed coat area (SCA), and relative seed coat

size (RSCS). Differences in seed morphology between mainland and island population were significant (α=0.05) for all 14 seed traits.

A – Measured seed traits

Population	SL (cm)	SW (cm)	EL (cm)	EW (cm)	SCW (cm)	SA (cm^2)	ESA (cm^2)	EA (cm^2)
Mainland	1.42$\pm$0.016	0.77$\pm$0.009	1.09$\pm$0.013	0.15 $\pm$0.003	0.09$\pm$0.002	0.89$\pm$0.023	0.55$\pm$0.014	0.17$\pm$0.004
Island	1.67$\pm$0.019	0.97$\pm$0.014	1.18 $\pm$0.019	0.18 $\pm$0.003	0.12$\pm$0.002	1.39 $\pm$0.032	0.81$\pm$0.021	0.23$\pm$0.007

621 B – Derived seed traits

Population	SLW (SL/SW)	ELW (EL/EW)	RES (EA/ESA)	REndS (ESA/SA)	SCA (cm^2) (SA-ESA)	RSCS (SCA/SA)
Mainland	1.86$\pm$0.013	7.3 $\pm$0.092	0.31$\pm$0.005	0.62$\pm$0.003	0.34$\pm$0.009	0.38$\pm$0.003
Island	1.74 $\pm$0.021	6.59$\pm$0.132	0.28 $\pm$0.005	0.58$\pm$0.004	0.59$\pm$0.013	0.42$\pm$0.004

Fig. 1. *Pinus torreyana* distribution map. Top left dark grey area indicates the distribution of

Torrey pine on Santa Rosa Island, CA (*Pinus torreyana* sp. *insularis*). Bottom right dark grey

area indicates the distribution of Torrey pine at the Torrey Pine State Reserve, CA (*Pinus*

torreyana sp. *torreyana*).

Fig. 2. Visual of morphological measurements taken using ImageJ for seeds collected on Santa

Rosa Island and at the Torrey pine State Reserve. (A) Seed length [cm]. (B) Seed width [cm]. (C)

Embryo length [cm]. (D) Embryo width [cm]. (E) Seed coat width [cm]. (F) Embryo area [cm^2].

(G) Endosperm area [cm^2]. (H) Seed area [cm^2].

Fig. 3. Principal components analysis (PCA) using all 14 seed morphological traits measured

from maternal plants collected on Santa Rosa Island (black) and at the Torrey Pine State Reserve

(grey).

Fig. 4. Proportion of variance in seed morphology explained by populations (light grey),

maternal families within populations (grey), and other variables not accounted for in the model

(residuals; dark grey) for each of the 14 measured and derived seed traits. SW, seed width (cm);

SLW, seed length/width ratio; SL, seed length (cm); SCW, seed coat width (cm); SCA, seed coat

area (cm^2); SA, seed area (cm^2); RSCS, relative seed coat size; RES, relative embryo size;

REndS, relative endosperm size; EW, embryo width (cm); ESA, endosperm area (cm^2); ELW,

embryo length/width ratio; EL, embryo length (cm); EA, embryo area (cm^2). See Appendix S2

for numerical estimates.

Fig. 5. Seed emergence, measured as the proportion of emerged seeds (y axis), recorded at three

different timepoints (x axis) for seeds sampled on Santa Rosa Island (black) and at the Torrey

Pine State Reserve (gray). Significant differences in emergence time across timepoints within

populations are indicated with different letters. Comparisons between populations at each

timepoint is indicated with square brackets. ns, non-significant differences (α=0.05).

Fig. 6. Average proportion of phenotypic variation captured across seed traits estimated for

various proportions of maternal families sampled. P_{island} and $P_{mainland}$ represent the proportion of

maternal families required to capture 95% of morphological variation (horizontal dashed line)

present in island (black) and mainland (gray) ex situ seed populations respectively.