A Global Temporal Genetic Program for Neural Circuit Formation

Saumya Jain[1,4], Ying Lin[1,2,4], Yerbol Kurmangaliyev[3], Parmis Mirshahidi[1], Brianna Parrington[1], and S. Lawrence Zipursky[1,3,*]

[1]Department of Biological Chemistry, University of California, Los Angeles, USA. [2]Molecular Biology Institute, University of California, Los Angeles, USA. [3]Howard Hughes Medical Institute, David Geffen School of Medicine, University of California, Los Angeles, USA. [4]These authors contributed equally to this work. [*]e-mail: lzipursky@mednet.ucla.edu

Wiring a complex brain relies on cell-type and temporal-specific expression of genes

encoding cell recognition molecules regulating circuit formation[1–3]. Though genetic

programs driving cell-type specificity have been described[4–6], how precise temporal control

is achieved remains unknown. Here, we describe a global program for the temporal

regulation of cell-type-specific wiring genes in the *Drosophila* visual system. We show that

the Ecdysone Receptor induces a synchronous cascade of transcription factors in neurons

throughout the visual system in a highly stereotyped fashion. Single-cell RNA-seq analysis of

neurons lacking transcription factors in the cascade revealed that targets are cell-type

dependent and these are enriched for wiring genes. Knock-down of different transcription

factors in this cascade led to defects in sequential steps in wiring. Taken together, this work

identifies a synchronous, global program for temporal control of different sets of wiring

genes in different neurons. We speculate that this global program coordinates development

across cell types to choreograph specific steps in circuit assembly.

MAIN

Animal behavior is dependent upon the formation of neuronal circuits with high fidelity.

Many cell-surface proteins, particularly of the Immunoglobulin (Ig), Leucine-Rich Repeat and

Cadherin families, promote interactions between neurons during circuit assembly[1,7]. Several

developmental strategies have evolved to meet the demands of cellular recognition specificity

during circuit construction. These include the expression of combinations of different recognition

molecules in different neurons, and reusing the same molecules to determine multiple, spatially

and temporally separated neuron-neuron interactions[2,3,8]. Wiring specificity, thus, relies critically

on genetic programs which regulate the cell-type and temporal specificity of genes encoding

recognition molecules. Many programs that define cell-type specificity have been described[4–6].

Here we set out to assess how temporal specificity is determined.

Dynamic expression of wiring genes

To identify genetic programs regulating the temporal control of wiring genes, we profiled

the transcriptomes of post-mitotic lamina neurons (L1-L5) in the *Drosophila* optic lobe (analogous

to bipolar cells in the mammalian retina[8]), every 12h from 24hAPF (hours After Pupa Formation)

to adults (Fig. 1a, b). These time points encompass all aspects of circuit formation for lamina

neurons once their growth cones have reached their target neuropil[9,10]. For transcriptional

profiling, we modified a recently developed single-cell RNA-Seq-based approach that utilizes the

natural genetic variation (i.e detected by single nucleotide polymorphisms (SNPs)) amongst

different *Drosophila* wildtype lines (DGRP[11]) to determine gene expression in many different

samples and different time points in one experiment (Kurmangaliyev *et al*. revision under review,

Fig. 1b, Extended data Fig. 1, Supplementary Table 1). This approach eliminates issues due to

batch effects that limit transcriptome analyses combining data from independent experiments.

Using this approach, we acquired transcriptomic data for 8269 cells distributed between seven

timepoints (24hAPF – Adult) (Fig. 1b, Extended data Fig. 1b, Supplementary Table 2). Using the

SNPs within the transcripts encoded by the different DGRP strains, we assigned each cell to a

specific time point. We integrated all cells and performed unsupervised clustering independent of

global changes occurring over time yielding five clusters (see Methods). We used previously

identified lamina neuron-type specific genes to assign each cluster to a cell-type[12] (Extended data

Fig. 1c). For all subsequent analyses, we used normalized gene expression values prior to

integration. The expression patterns were similar to expression data from bulk RNA-seq (this

study, Supplementary Table 3) and from a large single cell sequencing study of visual system

neurons from our laboratory (Kurmangaliyev *et al.* revision under review) (Extended data Fig. 2).

Application of k-means clustering to group genes based on their expression dynamics

revealed several groups that showed dynamic patterns of expression (e.g. gene groups 5, 7, 9 and

10 in Fig. 1c; also see Extended data Fig. 3 and Supplementary Table 4). Dynamically expressed

genes (especially early-peaking gene groups) were enriched (over expected by chance) for genes

encoding Immunoglobulin Superfamily (IgSF) proteins, as well as GO terms related to wiring (Fig.

1c, Extended data Fig. 3, Supplementary Table 5). Genes encoding IgSF proteins expressed in a

dynamic temporal fashion (see Methods), were also expressed in a cell-type-dependent manner

(Extended data Fig. 4a). Thus, as previously described[13,14], wiring genes are expressed in a cell-

type and temporally dynamic fashion during circuit assembly.

Synchronous expression of the EcR Cascade

To identify transcription factors (TFs) that drive gene expression in a dynamic fashion, we looked at TFs that were differentially expressed between pairs of consecutive time points for each neuron-type across development (Extended data Fig. 4b, see Methods). Interestingly, for each neuron-type, dynamically expressed TFs were enriched for the *Nuclear Receptor Transcription Factor Pathway* (Reactome Pathway Enrichment Analysis[15], p-value < 10^{-5}, Extended data Fig. 4c). These TFs, including the **Ec**dysone **R**eceptor (EcR) and others regulated by Ecdysone[16,17], exhibit synchronous patterns of expression in lamina neurons over development (red lines in Fig. 2a, Extended data Fig. 5a, b). In addition, EcR also undergoes a temporal-specific isoform switch due to the use of an alternative transcription start site, from predominantly EcR-B1 to EcR-A in these neurons (Fig. 2a, Extended data Fig. 5), as determined by staining with isoform-specific antibodies and bulk RNA sequencing. This temporal switch in isoform expression has been reported previously for other neurons in the central nervous system[18].

Immunostaining for EcR-B1, EcR-A, Hr3 and Ftz-f1 (Extended data Fig. 5a), and our analysis of the whole optic lobe transcriptome (Kurmangaliyev *et al.* revision under review) revealed that this transcriptional cascade has near-synchronous dynamics in all optic lobe neurons throughout development (Fig. 2a). The timing and order of expression of these TFs correlates with a mid-pupal pulse of Ecdysone[16,17,19–21] (Fig. 2a). A role for this transcriptional cascade in various developmental contexts, including neuronal remodeling of specific neuron-types during metamorphosis has been studied in detail[22,23]. The coordinate and visual-system-wide expression of these transcription factors during pupal development raised the intriguing possibility that this cascade acts as a temporal regulator of circuit formation more broadly.

Temporal control of cell-type specific wiring genes

We next sought to address whether the EcR cascade drives the temporal expression of genes in lamina neurons. Analysis of putative cis-regulatory regions[24] for dynamically expressed genes in lamina neurons showed a significant enrichment for binding sites for TFs in the EcR cascade (Supplementary Table 6, see Methods), consistent with their direct role in driving these patterns of gene expression. Additionally, ATAC-Seq[25] at 40hAPF, 60hAPF and 72hAPF in L1 neurons (see Methods and Extended data Fig. 6, Supplementary Table 7) identified an enrichment for predicted binding sites for TFs in this cascade (e.g. EcR, Hr3, Hr4 and Ftz-f1) in regions with dynamic patterns of accessibility[24] (Extended data Fig. 6d, Supplementary Table 6). This is consistent with these TFs controlling chromatin structure, transcription of specific genes or both[26].

To identify genes regulated by the EcR cascade, we targeted expression of a dominant negative allele of EcR (EcR^{DN}, EcR-B1 W650A[27]), EcR RNAi and Hr3 RNAi to lamina neurons. Transcriptional profiling was performed using a single-cell RNA-Seq-based assay (Fig. 2b) to compare transcriptomes of wildtype and mutant lamina neurons at four timepoints (24hAPF, 48hAPF, 72hAPF and Adult) in a pooled fashion using the DGRP strategy (Supplementary Table 1). By comparing the transcriptomes of 3606 wildtype and 1503 EcR^{DN}-expressing lamina neurons, we identified 921 differentially expressed genes (fold change ≥ 2, p-value ≤ 0.05), out of a total of 3200 genes expressed in lamina neurons across development. These were distributed between the five lamina neuron-types and four time-points (Extended data Fig. 7, Supplementary Table. 8). Importantly, EcR^{DN}-expressing lamina neurons continued to express their cell-type specific TFs, suggesting no changes in cell-fate (Extended date Fig. 7b). In contrast to L1 and L3-L5, EcR^{DN} had modest effects on gene expression in L2. This is consistent with lower EcR^{DN} expression in L2 neurons at 48hAPF using this GAL4 driver line (Extended data Fig. 7e). Analysis

of the 921 differentially expressed transcripts led to four conclusions. First, EcR-dependent genes

are enriched for those showing dynamic expression in wildtype cells (Extended data Fig. 7f, 11).

Second, a large set of genes dependent upon this cascade is expressed in a cell-type-dependent

fashion in wildtype lamina neurons (Fig. 2c, Extended data Fig. 8). Third. the cell-type dependent

genomic targets of this cascade were highly enriched for IgSF genes (Hypergeometric test, p-value

$= 1.25 \times 10^{-18}$, see Methods) as well as GO terms associated with wiring (Fig. 2d, e, Supplementary

Table 9). And finally, it was not uncommon for EcR^{DN} to have different effects on the same gene

in different neuron types. This may reflect different levels of expression of EcR^{DN}. In some cases,

however, opposite effects in different lamina neurons were observed (Fig. 2d).

Three lines of evidence indicated that genes affected by the expression of EcR^{DN} were

largely bona fide targets of this transcriptional cascade. First, most genes affected by EcR^{DN} were

affected in a similar way by EcR RNAi, although the change was typically less pronounced

(Extended data Fig. 9, 12a; Supplementary Table 10). Second, changes in gene expression in Hr3

RNAi and EcR^{DN} were similar, especially at later time points, as expected from their genetic

relationship (Fig. 2a; Extended data Fig. 10, 12b; Supplementary Table 11). And third, expression

of several putative EcR-dependent genes in brains cultured *ex vivo*[28] depended on the inclusion of

Ecdysone in the medium (see Methods, Extended data Fig. 13, 16b). Thus, EcR, either directly or

indirectly, controls the temporal expression of wiring genes in a cell-type-specific manner.

The cell-type-specific, temporal control of wiring genes by the EcR cascade suggests that

it plays a role in circuit formation. Consistent with this notion, we observed a disorganization of

lamina neuron arbors in the medulla neuropil upon pan-lamina expression of EcR^{DN}, which was

largely rescued by overexpression of wildtype EcR-B1 (Extended data Fig. 14a). A disorganized

neuropil was also seen upon targeted expression of EcR RNAi. Disruption of the axon terminal

morphology of T4/T5 neurons in the lobula plate neuropil, a more central visual system structure,

was also seen upon targeted expression of EcR[DN] to these neurons (Extended data Fig. 14b).

Together these data support the idea that the EcR transcriptional cascade regulates genetic

programs governing the appropriate expression of cell surface and secreted proteins required for

proper wiring of the visual system.

EcR regulates the L3-specific expression of Netrin

We next sought to assess whether loss of EcR function gave rise to similar phenotypes seen

for loss-of-function mutations in cell surface or secreted proteins regulating wiring in L1-L5. Loss

of EcR activity resulted in phenotypes similar to *Netrin* mutants in L3[29,30] (Fig. 3a-d) and mutations

in *DIP-β,* encoding an IgSF protein, in L4[31] (Extended data Fig. 15). Here we focus on the

relationship between *EcR* and *Netrin*.

Netrin protein is required in L3 axon terminals in the M3 layer to promote adhesion of

photoreceptor R8 growth cones to processes in the same layer. In *Netrin* mutants, R8 terminals

extend into the developing M3 layer, but are unable to stably adhere to it and retract to the surface

of the medulla[30]. At least one other, yet to be identified, signal is necessary for R8 to recognize the

M3 layer[30]. From sequencing of mutants and anti-Netrin antibody staining, EcR activity is required

for Netrin expression in L3 (Fig. 3a, b, Extended data Fig. 16b). Targeting expression of EcR[DN] in

L3 leads to an R8 mis-targeting phenotype. Here most R8 terminals extend beyond the M3 layer

and some remain at the surface of the medulla. The EcR[DN] phenotype is partially rescued by

overexpression of Netrin in developing EcR-mutant L3 neurons (Fig. 3c). Differences between the

Netrin mutant and EcR[DN] phenotypes are consistent with the notion that EcR regulates one or more

signals in L3, in addition to Netrin, necessary for R8 targeting[30] (Fig. 3d).

As EcR is expressed in the same way in all lamina neurons, how is the cell-type specificity

of wiring genes determined? Interestingly, genes specifically expressed in L3, including *Netrin*,

are enriched for targets of the L3-specific TF Erm identified in a previous study[12] (Hypergeometric

test, p-value = 2.7 X 10^{-18}; Fig. 3e; Extended data Fig. 17). Erm is expressed selectively in L3 early

in lamina development and remains expressed into the adult[12,32] (Extended data Fig. 1c). Thus,

here the EcR pathway confers temporal specificity to a continuously expressed cell-type-specific

TF to ensure the precise spatial and temporal expression of wiring genes.

Stepwise assembly of L5 connectivity

We next sought to assess whether the EcR cascade regulates sequential steps in wiring. For

this we turned to L5. L5 axon morphology in the medulla forms in a stepwise manner[33] (Fig. 4b).

L5 targets to M5 and then arborizes in M1 and M2 (M1/2) (Fig. 1a, 4b). Arborization occurs in

two steps: 1) initial branching in M1 (between 48hAPF and 60hAPF) and, 2) growth from M1 to

M2 (between 75hAPF and 90hAPF) (Fig. 4b). EcR^{DN} or EcR RNAi expression blocks step 1, while

Hr3 and ftz-f1 RNAi each blocks step 2 (Fig. 4a, b). These findings are consistent with the notion

that EcR regulates the expression of genes that promote extension of dendrites into M1 and that

Hr3 and Ftz-f1 execute a subsequent developmental step, some 24 hours later (Fig. 4b). Hr3

directly regulates *ftz-f1* expression[17] (see Fig. 2a, Extended data Fig. 10d). Thus, Ftz-f1 may act

alone or in combination with Hr3, to regulate the expression of a set of genes necessary to control

targeting of dendrites to M2. As L5 makes synaptic connections with different partners in M1 and

M2[34], EcR may control cell surface proteins regulating synaptic specificity in M1, whereas

synaptic specificity in M2 may be controlled by cell recognition proteins regulated by Hr3, Ftz-f1,

or both.

The phenotypes resulting from Hr4 RNAi are qualitatively different from knocking down

levels of EcR, Hr3 and Ftz-f1. Loss of Hr4 results in promoting extension of L5 branches beyond

M2 to M5 (Fig. 4a). Hr4 may regulate a stop signal, preventing extension beyond M2 to deeper

layers. Alternatively, as Hr4 is expressed at a much earlier step in development (see Fig. 2a), it

may repress a growth pathway acting in the initial extension into M1. As knock down experiments

of EcR pathway components led to marked changes in the expression of genes encoding different

cell surface and secreted proteins implicated in wiring (Fig. 4c, Supplementary Table 12), these or

discrete combinations of them, may function at different steps or in specific spatial domains of L5

axons to establish specific patterns of synaptic connections.

Discussion

Wiring the brain requires precise cell-type and temporally restricted expression of genes

encoding cell surface proteins[2,3,12,13,35]. Our studies uncovered a strategy that plays a widespread,

yet specific role, in regulating this process. We describe a genetic program expressed in most, if

not all, visual system neurons that controls the precise temporal patterns of expression of wiring

genes (Fig. 4d). These have cell-type specific outputs which reflect the integration of a shared

temporal program with diverse pathways controlling the development of different cell types. We

show that this temporal program regulates discrete steps in wiring specificity.

The timing of expression of wiring genes results from the activation of a transcriptional

cascade by the steroid hormone Ecdysone, comprising multiple transcription factors (such as EcR,

Hr3 and Ftz-f1) including members of the nuclear receptor family. These produce sequential waves

of gene expression[16] (Fig. 2, 4d) which regulate different steps in wiring in different neurons. The

control of a cell-type-specific set of wiring genes is made possible, at least in part, by a

combinatorial mechanism in which many members of this transcriptional cascade work with

different cell-type-specific TFs to produce different outputs[36–38]. Mechanisms that establish

cellular transcriptional landscapes early in development have been identified[39–41], including cell-

type-specific TFs (e.g. Erm). These are expressed in postmitotic neurons and determine multiple

cell-type-dependent features, including transcription of genes required for circuit formation[4,42,43].

By extrapolation of the dual control of Netrin by Erm and EcR in L3, we speculate that it is the

coordinate activities of these two classes of TFs that regulates developmental programs specifying

connectivity.

Dynamic expression of wiring genes, including cell-recognition molecules, poses an

additional challenge for gene expression control. How are the transcriptomes of diverse neuron-

types coordinated to ensure proper wiring? A global temporal signal, such as Ecdysone, is a

possible mechanism to coordinate the development of multiple neuron types. For instance, the

Ecdysone-dependent timing of Netrin expression in L3 growth cones matches the arrival of R8

growth cones in M3 (Fig. 3d; Extended data Fig. 16a). Further experiments will be necessary to

determine whether Ecdysone not only regulates the timing of developmental steps, but also ensures

that these are coordinated between different neuron-types in the complex choreography leading to

circuit assembly.

Our studies rest on decades of genetic and biochemical experiments establishing the

Ecdysone-pathway as a temporal regulator of major developmental transitions, including larval

molts and metamorphosis, as well as controlling sequential divisions of postembryonic neuroblasts

and neuronal re-modelling[16,23,37,38,44]. Here we show that this program has been co-opted for the

temporal regulation of differentiation, and the coordinate development of different neuron types

leading to the complex circuitry of the fly visual system. The ability of the EcR cascade to function

as a flexible temporal regulatory module used in multiple developmental contexts may be similar

to modules regulating spatial patterning (e.g. Hox genes and BMP/Wnt/Hh pathways[45–47]) which

act in context-dependent ways to generate different morphological outcomes. These findings raise

the intriguing possibility that similar timing mechanisms may also regulate the sequence of events

regulating wiring in other systems[48] including the maturational sequence in the assembly of neural

circuits in the mammalian brain[49].

Fig. 1 | A developmental transcriptome reveals dynamic expression of wiring genes. a, Lamina neurons (L1-L5) in the *Drosophila* visual system project axons into the medulla. b, Scheme for single-cell RNA-Seq of lamina neurons (see Methods) using wildtype strains (DGRP) with unique SNPs as bar codes. Each time point includes lamina neurons from three animals, each heterozygous for a different SNP bar code. In total 21 different DGRP lines were used in this experiment. Thus, the transcripts of cells from each animal and their developmental stage (hours after pupal formation (hAPF)) were identified by unique SNPs. c, Gene groups generated via k-means clustering using expression dynamics in L1 neurons (see Methods; for other lamina neurons see Extended data Fig. 3). Relative expression over development for each gene in the group is shown as a line plot (purple lines), with the mean shown as a black line. n, number of genes in the

229 group. A gene from each group is shown (e.g. Xbp1). Heat map depicts enrichment (-$\log_{10}$ p-

230 values) of Immunoglobulin Superfamily genes over that expected by chance (see Methods).

Fig. 2 | Wiring genes in different cell-types are controlled by an Ecdysone-induced transcriptional cascade. a, Schematic of the genetic interactions between transcription factors of

the Ecdysone-induced transcriptional cascade. Only a subset of the known interactions is shown.

In the plots, red lines show normalized expression of these transcription factors in lamina neurons

and grey lines show expression dynamics in 83 other neuron-types (data from Kurmangaliyev *et*

al. revision under review) (bottom). Isoform change from EcR-B1 to EcR-A during development

is shown. Concentration of circulating Ecdysone over development is shown on the left (adapted

from Pak and Gilbert[21]) **b,** Scheme for scRNA-Seq-based analysis of genes affected by

perturbation of EcR or Hr3 activity in lamina neurons. Mutant and control brains for each

perturbation were analyzed together in three separate experiments. UAS-tdTom and UAS-wRNAi

were use as controls for EcR^{DN} and EcR (and Hr3) RNAi, respectively. **c,** Heat map showing

relative expression in wildtype cells of genes affected by EcR^{DN} at 48hAPF. Targets of EcR are

divided into two categories: Cell-type dependent, in any pairwise comparison fold difference ≥ 2,

p-value ≤ 0.05, 303 genes; Common targets, all remaining EcR targets, 208 genes. **d,** Expression

of some wiring genes affected by EcR^{DN}. Solid line, expression in wildtype cells; Dashed line,

expression in EcR^{DN} cells. **e,** Expression of some wiring genes altered by Hr3 RNAi. Solid line,

expression in wildtype cells; Dashed line, expression in Hr3 RNAi cells; *, difference in

expression between wildtype and perturbation ≥ 2-fold, p-value ≤ 0.05.

Fig. 3 | EcR-dependent program in L3 neurons controls R8 targeting. a, Normalized expression of NetrinA (NetA) and NetrinB (NetB) over development in WT (solid lines) and

EcR^{DN} (dashed lines) expressing L3 neurons. *, difference in expression between wildtype and

$EcR^{DN} \geq$ 2-fold, p-value < 0.05. **b,** Anti-NetB staining (magenta) at 48hAPF $\pm$ EcR^{DN} expression

in L3 neurons (green). Netrin expression in M3 is lost in response to EcR^{DN}. Note, as an internal

control, that NetB expression in the lobula (lo) is not affected. la, lamina; me, medulla; yellow

arrowhead, M3 layer; white arrowheads, L3 growth cones (some other cell types with axons in the

lobula are also labeled (white arrows)). Scale bar, 50μm. **c,** Morphology of R8 axons (magenta)

with wildtype L3 neurons (top). Other panels from top to bottom show L3s expressing EcR^{DN},

NetB and both EcR^{DN} and NetB. Left panels: R8 axons extending beyond M3 (white arrows); R8

axons terminals above the medulla neuropil (yellow arrowheads). Scale bar, 10μm. Right column,

distribution of R8 axon depths represented as ratio of depth of R8 to M6 (medulla layer 6 indicated

in green in the left panels). Red line, median of the distribution in wildtype. Distribution with WT

and EcR^{DN} expressing L3 is significantly different (Kolmogorov-Smirnov test, p-value < 10^{-15}, see

Extended data Fig. **16c** for other comparisons). n, number of neurons observed (6-10 animals for

all conditions). **d,** Model for control of R8 targeting by L3. EcR controls expression of Netrin and

one or more other molecules (X) in L3. Netrin is required for adhesion of R8 growth cones to

processes in M3 and X is required for recognition of M3 by R8 growth cones to terminate

extension[30]. **e,** Heat map showing relative expression of L3-specific targets of EcR (see Methods).

Aqua, targets of transcription factor Erm identified in Peng *et al.*[32]; Magenta, Immunoglobulin

Superfamily genes; Dashed lines, NetA and NetB.

Fig. 4 | The EcR cascade controls the stepwise arborization of L5 axons. a, L5 axonal arborization defects (left) and their distributions in wildtype L5's, or under pan-lamina expression of EcR[DN], EcR RNAi, Hr3 RNAi, Hr4 RNAi or ftz-f1 RNAi (right). *, Fisher's exact test p-value

< 0.01. n, number of neurons observed (4 -13 animals for all conditions). In wildtype only the axon

extends to M5; L5 neurons in Hr4 RNAi animals extend an additional process to M5. Scale bars,

5μm. **b,** Schematic of the stepwise morphogenesis of L5 in the medulla[33] showing the expected

roles for EcR, Hr3 and Ftz-f1 based on data in panel a) and previous biochemical and gene

expression studies (see Fig. 2a). The role of Hr4 in this process is unclear (see text). **c,** Volcano

plots showing candidate genes involved in step1 (Left, 131 genes, significantly affected in EcR^{DN}

at 48hAPF and not affected in Hr3 RNAi at any timepoint) and step2 (Right, 124 genes,

significantly affected in Hr3 RNAi at 72hAPF, regardless of effect of EcR^{DN}) of L5 arborization.

Immunoglobulin Superfamily genes are highlighted. **d,** Model: Control of wiring genes through a

global temporal program acting in combination with cell-type-specific transcription factors (see

Discussion). Other temporal regulators may also contribute to wiring.

References

1. Südhof, T. C. Towards an Understanding of Synapse Formation. *Neuron* **100**, 276--293

(2018).

2. Hassan, B. A. & Hiesinger, P. R. Beyond Molecular Codes: Simple Rules to Wire Complex

Brains. *Cell* **163**, 285--291 (2015).

3. Yogev, S. & Shen, K. Cellular and Molecular Mechanisms of Synaptic Specificity. *Annu Rev*

Cell Dev Bi **30**, 417--437 (2014).

4. Hobert, O. Chapter Twenty-Five - Terminal Selectors of Neuronal Identity. in *Current Topics*

in Developmental Biology (ed. Wassarman, P. M.) vol. 116 455--475 (Academic Press, 2016).

5. Hong, W. & Luo, L. Genetic Control of Wiring Specificity in the Fly Olfactory System.

Genetics **196**, 17--29 (2014).

6. Dasen, J. S. & Jessell, T. M. Chapter Six Hox Networks and the Origins of Motor Neuron

Diversity. in *Current Topics in Developmental Biology* vol. 88 169--200 (Academic Press,

2009).

7. Zipursky, S. L. & Sanes, J. R. Chemoaffinity Revisited: Dscams, Protocadherins, and Neural

Circuit Assembly. *Cell* **143**, 343--353 (2010).

8. Sanes, J. R. & Zipursky, S. L. Design Principles of Insect and Vertebrate Visual Systems.

Neuron **66**, 15--36 (2010).

9. Pecot, M. Y. *et al.* Sequential Axon-Derived Signals Couple Target Survival and Layer

Specificity in the Drosophila Visual System. *Neuron* **82**, 320--333 (2014).

10. Chen, Y. *et al.* Cell-type-Specific Labeling of Synapses In Vivo through Synaptic Tagging

with Recombination. *Neuron* **81**, 280--293 (2014).

11. Mackay, T. F. C. *et al.* The extitDrosophila melanogaster Genetic Reference Panel. *Nature*

482, 173--178 (2012).

12. Tan, L. *et al.* Ig Superfamily Ligand and Receptor Pairs Expressed in Synaptic Partners in

Drosophila. *Cell* **163**, 1756--1769 (2015).

13. Li, H. *et al.* Classifying Drosophila Olfactory Projection Neuron Subtypes by Single-Cell

RNA Sequencing. *Cell* **171**, 1206--1220.e22 (2017).

14. Favuzzi, E. *et al.* Distinct molecular programs regulate synapse specificity in cortical

inhibitory circuits. *Sci New York N Y* **363**, 413--417 (2019).

15. Jassal, B. *et al.* The reactome pathway knowledgebase. *Nucleic Acids Res* **48**, D498--D503

(2020).

16. Riddiford, L. M., Cherbas, P. & Truman, J. W. Ecdysone receptors and their biological

actions. *Vitamins Hormones* **60**, 1--73 (2000).

17. White, K. P., Hurban, P., Watanabe, T. & Hogness, D. S. Coordination of Drosophila

metamorphosis by two ecdysone-induced nuclear receptors. *Sci New York N Y* **276**, 114--117

(1997).

18. Truman, J. W., Talbot, W. S., Fahrbach, S. E. & Hogness, D. S. Ecdysone receptor

expression in the CNS correlates with stage-specific responses to ecdysteroids during Drosophila

and Manduca development. *Dev Camb Engl* **120**, 219--234 (1994).

19. Agawa, Y. *et al.* Drosophila Blimp-1 Is a Transient Transcriptional Repressor That Controls

Timing of the Ecdysone-Induced Developmental Pathway. *Mol Cell Biol* **27**, 8739--8747 (2007).

20. Rabinovich, D., Yaniv, S. P., Alyagor, I. & Schuldiner, O. Nitric Oxide as a Switching

Mechanism between Axon Degeneration and Regrowth during Developmental Remodeling. *Cell*

164, 170--182 (2016).

21. Pak, M. D. & Gilbert, L. I. A Developmental Analysis of Ecdysteroids During the

Metamorphosis of extitDrosophila Melanogaster. *J Liq Chromatogr* **10**, 2591--2611 (1987).

22. Boulanger, A. & Dura, J.-M. Nuclear receptors and Drosophila neuronal remodeling.

Biochimica Et Biophysica Acta Bba - Gene Regul Mech **1849**, 187--195 (2015).

23. Alyagor, I. *et al.* Combining Developmental and Perturbation-Seq Uncovers Transcriptional

Modules Orchestrating Neuronal Remodeling. *Dev Cell* **47**, 38--52.e6 (2018).

24. Imrichová, H., Hulselmans, G., Atak, Z. K., Potier, D. & Aerts, S. i-cisTarget 2015 update:

generalized cis-regulatory enrichment analysis in human, mouse and fly. *Nucleic Acids Res* **43**,

W57--W64 (2015).

25. Buenrostro, J. D., Wu, B., Chang, H. Y. & Greenleaf, W. J. ATAC-seq: A Method for

Assaying Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biology Ed Frederick M*

Ausubel Et Al **109**, 21.29.1--9 (2015).

26. Olivares, A. M., Moreno-Ramos, O. A. & Haider, N. B. Role of Nuclear Receptors in Central

Nervous System Development and Associated Diseases. *J Exp Neurosci* **9**, 93--121 (2016).

27. Cherbas, L., Hu, X., Zhimulev, I., Belyaeva, E. & Cherbas, P. EcR isoforms in Drosophila:

testing tissue-specific requirements by targeted blockade and rescue. *Development* **130**, 271--284

(2003).

28. Özel, M. N., Langen, M., Hassan, B. A. & Hiesinger, P. R. Filopodial dynamics and growth

cone stabilization in Drosophila visual circuit development. *Elife* **4**, e10721 (2015).

29. Timofeev, K., Joly, W., Hadjieconomou, D. & Salecker, I. Localized Netrins Act as

Positional Cues to Control Layer-Specific Targeting of Photoreceptor Axons in Drosophila.

Neuron **75**, 80--93 (2012).

30. Akin, O. & Zipursky, S. L. Frazzled promotes growth cone attachment at the source of a

Netrin gradient in the Drosophila visual system. *Elife* **5**, e20762 (2016).

31. Xu, C. *et al.* Control of Synaptic Specificity by Establishing a Relative Preference for

Synaptic Partners. *Neuron* **103**, 865--877.e7 (2019).

32. Peng, J. *et al.* Drosophila Fezf coordinates laminar-specific connectivity through cell-

intrinsic and cell-extrinsic mechanisms. *Elife* **7**, (2018).

33. Nern, A., Zhu, Y. & Zipursky, S. L. Local N-Cadherin Interactions Mediate Distinct Steps in

the Targeting of Lamina Neurons. *Neuron* **58**, 34--41 (2008).

34. Takemura, S. *et al.* A visual motion detection circuit suggested by extitDrosophila

connectomics. *Nature* **500**, 175--181 (2013).

35. Kurmangaliyev, Y. Z., Yoo, J., LoCascio, S. A. & Zipursky, S. L. Modular transcriptional

programs separately define axon and dendrite connectivity. *Elife* **8**, e50822 (2019).

36. Barolo, S. & Posakony, J. W. Three habits of highly effective signaling pathways: principles

of transcriptional control by developmental cell signaling. *Gene Dev* **16**, 1167--1181 (2002).

37. Shlyueva, D. *et al.* Hormone-responsive enhancer-activity maps reveal predictive motifs,

indirect repression, and targeting of closed chromatin. *Mol Cell* **54**, 180--192 (2014).

38. Uyehara, C. M. & McKay, D. J. Direct and widespread role for the nuclear receptor EcR in

mediating the response to ecdysone in Drosophila. *Proc National Acad Sci* **116**, 9893--9902

(2019).

39. Erclik, T. *et al.* Integration of temporal and spatial patterning generates neural diversity.

Nature **541**, 365--370 (2017).

40. McDonald, J. A. *et al.* Dorsoventral patterning in the Drosophila central nervous system: the

vnd homeobox gene specifies ventral column identity. *Gene Dev* **12**, 3603--3612 (1998).

41. Cochella, L. & Hobert, O. Embryonic Priming of a miRNA Locus Predetermines Postmitotic

Neuronal Left/Right Asymmetry in C. elegans. *Cell* **151**, 1229--1242 (2012).

42. Kratsios, P. *et al.* Transcriptional Coordination of Synaptogenesis and Neurotransmitter

Signaling. *Curr Biol* **25**, 1282--1295 (2015).

43. Philbrook, A. *et al.* Neurexin directs partner-specific synaptic connectivity in C. elegans.

Elife **7**, e35692 (2018).

44. Syed, M. H., Mark, B. & Doe, C. Q. Steroid hormone induction of temporal gene expression

in Drosophila brain neuroblasts generates neuronal and glial diversity. *Elife* **6**, e26287 (2017).

45. Altmann, C. R. & Brivanlou, A. H. Neural patterning in the vertebrate embryo. *Int Rev Cytol*

203, 447–482 (2001).

46. Briscoe, J. & Small, S. Morphogen rules: design principles of gradient-mediated embryo

patterning. *Development* **142**, 3996–4009 (2015).

47. Gaunt, S. J. Hox cluster genes and collinearities throughout the tree of animal life. *Int J Dev*

Biol **62**, 673–683 (2018).

48. Lawson, H. *et al.* The Makorin lep-2 and the lncRNA lep-5 regulate lin-28 to schedule sexual

maturation of the C. elegans nervous system. *Elife* **8**, (2019).

49. McCarthy, M. M. Estradiol and the Developing Brain. *Physiol Rev* **88**, 91--124 (2008).

Methods

Fly husbandry and stocks

Maintenance and rearing of fly lines, as well as staging of pupae was done as described in

Tan et al.[12] Fly lines used in this work are listed in Supplementary Table 1.

Multiplexed single cell transcriptomic analysis

For transcriptomic analysis of developing wildtype lamina neurons, w; UAS-H2A-GFP;

9B08-Gal4/Tm6B, tb females were crossed with males from different DGRP backgrounds

(wildtype, see Supplementary Table 1 for list of DGRPs used). F1 prepupae were staged as in

Kurmangaliyev et al.[35] Males were only included in the analysis if no significant differences in

gene expression were found with female pupae of the same genotype and developmental time

point. Pupae corresponding to different developmental stages (see Fig. 1b) were all dissected,

dissociated and processed at the same time. Each developmental time point was represented by $\geq$

2 DGRP heterozygotes, and only one animal/ DGRP heterozygote was dissected. Tissue

dissociation, FACS and preparation of single-cell libraries using 10X Genomics Chromium (v3)

were carried out similar to Kurmangaliyev et al., except that H2A-GFP expression was used to

enrich for lamina neurons (see Extended data Fig. 1a). All libraries were sequenced on a

NextSeq500 platform (single-end 75bp).

For transcriptomic analysis involving UAS EcR^{DN} (BDSC #6872), males carrying a single

DGRP autonomous chromosome – w; (UAS EcR^{DN} or UAS tdTom); DGRP/ Tm6b, tb were

crossed with w; UAS-H2A-GFP; 9B08-Gal4/Tm6B females. For experiments involving UAS EcR

RNAi (BDSC #9326) or UAS Hr3 RNAi (BDSC #27253), w; DGRP; (UAS wRNAi or UAS EcR

RNAi or UAS Hr3 RNAi)/ Tm6B, tb males were crossed with UAS Dcr2; UAS H2A-GFP; 9B08

Gal4/ Tm6B, tb females (see Fig. 2b and Supplementary Table 1). UAS tdTom and UAS wRNAi

were used as controls.

scRNA-seq data pre-processing

Raw fastq read files were processed using Cell Ranger (3.1.0) with default parameters.

Seurat V3 was used for all preliminary analyses. FlyBase reference genome (release 6.29[50]) was

used for alignment and annotation. Profiled single cells were identified as coming from a particular

developmental time-point and genetic background based on the natural genomic variance within

the different parent DGRP lines. This was done using the pipeline described in Kurmangalyev *et*

al., with the following modifications: the count of minor allele that equals to 2 among the analyzed

DGRP strains were used as genomic variants to assign the profiled single cells to different DGRP

parent lines. Only single cells with: 1) number of genes between 200 and 3000, 2) number of

mitochondrial transcripts < 20% of all transcripts, and 3) assignment to a unique DGRP parent

line; were used for all downstream analyses.

To identify different lamina neuron subtypes, all cells from a particular experiment were

integrated as described in Seurat v3 workflow and then subjected to unsupervised clustering, thus

disregarding global temporal gene expression changes (similar to Kurmangaliyev *et al.*).

Previously identified lamina neuron subtype specific genes[12] were used to assign each cluster to a

cell-type. Cells not assigned to a lamina neuron-type cluster were removed from subsequent

analyses. Average expression for each gene for each cell-type at a particular timepoint and genetic

background was calculated using normalized expression values prior to integration.

Bulk RNA-seq

Bulk RNA-Seq analysis of L1 neurons at 40hAPF, 60hAPF and 72hAPF was performed

as in Tan *et al.*[12] At least 2 replicates were generated for each timepoint. A minimum of 2000 cells

were sorted for each experiment. cDNA libraries were created using the SMART-Seq2 protocol[51],

which were then sequenced together using the HiSeq4000 platform (paired-end 50bp).

Raw fastq reads files were mapped to FlyBase reference genome (release 6.29) using STaR

(2.6.0) and only uniquely mapped reads were collected. Genes with counts per million (CPM) ≥ 4

in more than 2 samples were used for normalization with R package edgeR (3.26.8). Gene

expression was quantified using RPKM units (Reads Per Kilobase of exon per Million reads

mapped), calculated based on reads in the sum of exons using customized scripts. The correlation

between biological replicates were calculated using Spearman correlation for top 500 genes with

the highest variation across all samples.

Bulk ATAC- seq

At least 8000 L1 neurons were FACS purified at 40hAPF, 60hAPF and 72hAPF as

described in Tan *et al.*[12] Biological duplicates were generated for each timepoint. ATAC-Seq

libraries were generated as per the following protocol, which was modified from the one described

in Buenrostro *et al.*[52]: 1) FACS purified cells were collected in 300ul 1X PBS and spun at 800xg

for 8' at 4°C. 2) Cell pellet was then directly resuspended in 50µl of a tagmentation enzyme mix

(25µl Nextera DNA library prep kit – Tagment DNA Buffer, 19µl Nuclease-free water, 5µl 1%

IGEPAL CA630, 1µl Nextera DNA library prep kit - Tagment DNA Enzyme) and incubated at

37°C for 30' with constant agitation at 400rpm. 3) DNA was purified from the tagmentation mixes

using the Qiagen minElute Reaction Cleanup Kit as per the manufacturer's protocol. 4) A 10µl

qPCR reaction was setup to determine the optimum number of amplification cycles (similar to

Buenrostro *et al.*). 5) A larger PCR reaction (40μl) was then setup and PCR products in the 200-

500bp range were gel purified. 6) Steps 4 and 5 were repeated, this time with barcoded primers

(see Buenrostro *et al.*) to allow multiplexing of all libraries onto a single lane of HiSeq4000

(paired-end 50bp).

Raw fastq reads files were mapped to FlyBase reference genome (release 6.29) using

Bowtie2 (2.2.9) and uniquely mapped genes were kept. All samples were pooled together prior to

peak calling. Read start positions were shifted +4 or -5bp and used for peak calling using MACS2

(2.1.1) with parameters "-q 0.01 --nomodel --shift -100 --extsize 200". There were 26,122 peaks

identified. Then bedtools multicov (2.27.1) was used to sum the total reads within each peak

separately for each sample. Peaks with CPM ≥ 4 were used for counts normalization with edgeR.

Correlations between biological replicates were calculated using Spearman correlation for the top

500 peak regions with the highest variation of peak levels across all samples. Distribution of the

top 5000 peaks across different genomic features at 40hAPF, 60hAPF and 72hAPF was

determined using the R package ChIPseeker (1.20.0). Differential peak analysis was performed

using edgeR. Fold-change ≥ 2 and adjusted p-value ≤ 0.05 were chosen as the cutoffs to define

peaks with differential accessibility between time points. Each ATAC peak was associated to the

nearest gene based on proximity to the transcription start site using ChIPseeker. To compare

change of ATAC-seq peak accessibility over time and change in expression of nearest gene, ln(fold

change ATAC-seq peak read coverage) was plotted against ln(fold change expression of nearest

gene) for each ATAC-Seq peak (Extended data Fig. 6). Bulk RNA-Seq data was used for this, as

this transcriptome dataset had been generated for the same time points as for ATAC-Seq. This was

done separately for peaks differentially accessible between 40hAPF and 60hAPF (Extended data

Fig. 6b (left)) and between 60hAPF and 72hAPF (Extended data Fig. 6b (right)).

Comparison of scRNA-Seq with other datasets

For comparison of lamina neuron transcriptomes generated here with lamina neuron

transcriptomes generated in Kurmangaliyev *et al.*, 500 genes with the highest variance across all

cell types and all timepoints were used to calculate Spearman correlation. Comparison of L1

transcriptome generated using scRNA-Seq and L1 transcriptome obtained from bulk RNA-Seq

was done similarly.

k-means cluster analysis

Prior to clustering, all average gene expression from scRNA-seq transcriptomic analysis

were normalized to maximum expression in each cell type across development. k-means clustering

was performed using kmeans function in R (centers=10 and nstart=20).

Analysis of gene group traits

IgSF enrichment

A list of IgSF members was obtained from the InterPro database (IPR036179). Enrichment

within each gene group over expected by chance was then calculated using hypergeometric

distribution analysis. The set of genes expressed in a cell-type across development was taken as

the universal set (N). Then number of genes in a group (n) and number of IgSFs expressed in a

cell-type (m) were used as the sample size and number of successes in population respectively.

Enrichment was calculated based on actual overlap between n and m, and the expected overlap

between the two, defined by (n x m)/N. For enrichment of IgSF members within cell-type

dependent targets of EcR, universal set was taken as all genes expressed in all lamina neurons over

development. Sample size and number of successes in the population were taken as: the total

number of IgSF genes expressed over development in all lamina neurons (n), and the total number

of cell-type dependent targets of EcR (m) respectively. Enrichment was calculated as described

above.

Gene Ontology enrichment analysis

Gene Ontology enrichment analysis for each gene group was performed using a R package

TopGO (2.36.0) using GO terms obtained from FlyBase (GO-gene association file version 2.1).

Weight algorithm and Fisher's exact test were used to calculate p-value. Categories that have p-

value ≤ 0.01 were considered as enriched.

Effect of perturbation of EcR and Hr3

For each perturbation, expression of each gene was normalized to maximum expression in

each cell type across development and genetic background. The normalized gene expression of all

genes was averaged separately for each genetic background to plot average behaviors for each

gene group (Extended data Fig. 11).

Differential gene expression analysis

Wilcoxon rank-sum test (fold change ≥ 2, adjusted p-value ≤ 0.05) was used to identify

genes differentially expressed between timepoints, cell-types or as a consequence of EcR/ Hr3

perturbations. Dynamic TFs and IgSF genes were defined as ones with differential expression

between consecutive timepoints. Cell-type dependent genes were defined as ones with differential

expression in at least one pairwise comparison amongst all lamina neurons. Cell-type specific

genes were defined as ones that are differentially expressed in all pairwise comparison including

one lamina neuron-type. Bonferroni adjustment was used for multiple testing correction.

Reactome pathway analysis

Dynamic transcription factors (separately for each cell-type) were entered into the

'Analyze Data' tool on reactome.org (https://reactome.org/PathwayBrowser/#TOOL=AT).

Reactome pathway analysis for *Drosophila melanogaster* was then run with default parameters.

TF binding site analysis

i-cisTarget (https://gbiomed.kuleuven.be/apps/lcb/i-cisTarget/)[24] was used (using default

parameters) to identify TF binding sites enriched within putative cis-regulatory regions and

ATAC-Seq peaks. dm6 version of the *Drosophila* genome, and version 5 of the i-cisTarget

database were used for all analyses. All ATAC-Seq peaks overlapping with transcription start sites

were excluded from TF binding site analyses to reduce the occurrence of promoter-enriched

sequence motifs in i-cisTarget results.

All EcR-pathway TF binding motifs enriched within different sets of ATAC-Seq peaks

were additionally verified using the Homer (v4.7) mergePeaks utility[53]. Briefly: first, the top 5000

occurrences of the motif across the genome were called using the homer scanMotifGenomeWide

utility. Second, Homer suite mergePeaks was then used to evaluate if the extent of overlap between

the top 5000 occurrences of the motif and the set of ATAC-Seq peaks is higher than expected by

chance. p-value ≤ 0.01 was considered as significantly more overlap than expected by chance.

Immunohistochemistry and microscopy

Immunohistochemistry and microscopy were performed as described in Xu *et al.*[54] with the following modifications: 1) For experiments involving staining of lamina neuron presynaptic sites, brains were fixed using glyoxal (3.12% glyoxal, 0.75% acetic acid and 20% ethanol, pH adjusted to 5.0) for 30' at RT and then washed 3X with PBST. 2) All images were acquired using an LSM880 confocal microscope.

Primary antibodies used in this study were: mAb24B10 (1:20, DSHB), rabbit anti-NetB (1:500, gift from Akin Lab), rabbit anti-dsRed (1:400, Clontech 632496), chicken anti-GFP (1:1000, Abcam ab13970), rat anti-Flag (1:200, Novus Biologicals, NBP1-06712), mouse anti-EcR-B1 (1:20, DSHB AD4.4), mouse anti-EcR-A (1:10, DSHB 15G1a), rabbit anti-Hr3 (1:50, gift from Thummel Lab), mouse anti-svp (1:20, DSHB 5B11), rabbit anti-erm (1:100, gift from Wang Lab), rat anti-bab2 (1:500, gift from Laski Lab), mouse anti-V5 (1:200, BioRad MCA1360), guinea pig anti-Bsh (1:400, see Tan *et al.* 2015).

Secondary antibodies used in this study were: goat anti-mouse 488 (1:500, ThermoFisher A-32723), goat anti-mouse 568 (1:500, ThermoFisher A-11031), goat anti-mouse 647 (1:500, ThermoFisher A-21235), goat anti-chicken 488 (1:1000, ThermoFisher A-11039), goat anti-rabbit 568 (1:500, ThermoFisher A-11011), goat anti-guinea pig 647 (1:500, ThermoFisher A-21450), goat anti-rabbit 647 (1:200, ThermoFisher A-32733), goat anti-rat 568 (1:500, ThermoFisher A-11077).

Image analysis

All images for figures were created using ImageJ (v2.1.0). Details for quantification of phenotypes are given below.

R8 targeting

Imaris (v9.1.2) was used to measure the distance of R8 terminals from the top of the

medulla (M0), as well as the distance of 6^{th} medulla layer (M6) from the top of the medulla. The

latter was estimated using mAb24B10 staining, which labels R7 neurons that terminate in M6.

Values reported on the X-axis of Fig. 3c are (R8-M0)/ (M6-M0).

L5 morphological defects

Imaris was used to visualize individual neurons. Each neuron was then manually assigned

to one of the stated categories.

Quantification of BRP puncta in the lamina

The distance between the proximal end of the lamina and the distal end of the lamina was

measured using Imaris individually for all lamina cartridges. A line was drawn at the measured

distance divided by 2. All BRP puncta more distal of this line were counted manually.

***Ex vivo* culture of dissected brains**

We used a protocol similar to the one described in Özel *et al.*[28] Briefly, brains were

dissected from 22hAPF pupae in pre-warmed medium (ThermoFisher #2172004) with either

1:1000 dilution of 1mg/ml 20-HydroxyEcdysone (dissolved in 100% Ethanol, Sigma H5142) or

equivalent volume of 100% ethanol. Brains were then incubated in ~ 200ul of medium ± 20-

HydroxyEcdysone in 96 well plates for 26h at 25ºC in a humidified chamber. Thereafter, brains

were fixed, stained and imaged using the aforementioned protocols for immunohistochemistry and

microscopy.

In the presence of 20-HydroxyEcdysone in the media, strong expression of EcR-B1 and

Hr3 is observed throughout the optic lobe, while no expression of Ftz-f1 is seen (Extended data

Fig. 13a). When 20-HydroxyEcdysone is omitted from the culture medium, no expression of Hr3

is seen (as expected), however weak induction of Ftz-f1 is observed. *In vivo,* by 22hAPF (when

brains are removed from pupae), Ecdysone levels have begun to rise and may be sufficient to

activate EcR-B1 in neurons[21]. As Hr3 activation requires a subsequent drop in Ecdysone titers[17],

we expect the weak expression of Ftz-f1 may be due to the transfer of 22hAPF brains to an

Ecdysone-free medium.

Other statistics

To compare the dynamics of genes affected by EcR^{DN} (fold change ≥ 2, p-value ≤ 0.05

between WT and EcR^{DN} at any time point) and genes unaffected by EcR^{DN}, normalized wildtype

expression for each gene was divided by the maximum expression at any time point throughout

development. Variance was then calculated for each gene over time. Two-tailed Student's t-test

was then used to calculate the statistical difference between the distributions of variance for these

two sets of genes. This was done separately for each cell-type.

Hypergeometric test was used to evaluate enrichment of targets of Erm within L3-specific

targets of EcR. The set of genes expressed in a L3 neurons across development was taken as the

universal set (N). Then number of targets of Erm from Peng *et al.*[32] (m) and number of L3-specific

targets of EcR (n) were used as the number of successes in population and sample size respectively.

For all box-plots, solid line depicts median, while the upper and lower bounds of the box

depict the third and first quantile of the data spread respectively.

Kolmogorov-Smirnov (KS) test (Fig. 3c, 16c), two-tailed Student's t-test (Extended data

Fig. 7e, 7f, 11, 15a), two-tailed Fisher's exact test (Fig. 3e, 4a) were performed using the following

basic R functions (R 3.6.1) respectively: ks.test, t.test, fisher.test. Raw data for all analyses is

available upon request.

Data and code availability

All raw sequencing data and codes will be provided upon request. They will be made

publicly available prior to publication.

Extended data Fig. 1 | scRNA-Seq-based transcriptomic analysis of lamina neurons across

development. a, GFP vs forward scatter 2-D plot showing criteria used to enrich for lamina

610 neurons by FACS. 'Cells' highlighted in purple were excluded despite being GFP+ due to their

611 small size. **b,** tSNE plots showing L1-L5 clusters at 24hAPF, 36hAPF, 48hAPF, 60hAPF, 72hAPF,

612 84hAPF and 96hAPF (Adult). **c,** Normalized expression of previously identified lamina neuron-

613 type-specific genes in L1-L5 clusters identified at each time point over development (see Tan *et*

614 *al.*[12]).

Extended data Fig. 2 | Comparison of scRNA-Seq-derived transcriptome with other datasets.
a, Comparison of lamina neuron transcriptomes generated by scRNA-Seq in this study and by scRNA-Seq in Kurmangaliyev *et al*. **b,** Comparison of replicates of bulk-RNA-Seq of L1 neurons at 40hAPF, 60hAPF and 72hAPF (this study). **c,** Comparison of lamina neuron transcriptomes generated by scRNA-Seq in this study and by bulk RNA-Seq, also this study. See Methods for calculation of Spearman correlation.

623   expression over development for each gene in the group is shown as a line plot (colored lines),

624   with the mean shown as a black line. n, number of genes in the group. A gene from each group is

625   shown. Heat map depicts enrichment (-$\log_{10}$ p-values) of Immunoglobulin Superfamily genes over

626   that expected by chance (see Methods).

a Expression of dynamic Immunoglobulin Superfamily genes

b Expression of dynamic transcription factors

c Reactome Pathway Enrichment Analysis

Reactome Pathway in L1	p-value
Nuclear Receptor transcription pathway	7.5X10⁻⁶
SUMOylation of intracellular receptors	1.1X10⁻⁴

Reactome Pathway in L2	p-value
Nuclear Receptor transcription pathway	7.4X10⁻⁸
SUMOylation of intracellular receptors	1.7X10⁻⁷

Reactome Pathway in L3	p-value
Nuclear Receptor transcription pathway	1.3X10⁻⁷
SUMOylation of intracellular receptors	7.4X10⁻⁷
SUMO E3 ligases SUMOylate target proteins	1.2X10⁻³
SUMOylation	1.3X10⁻³
Generic Transcription Pathway	3.6X10⁻³

Reactome Pathway in L4	p-value
Nuclear Receptor transcription pathway	1.3X10⁻⁹
SUMOylation of intracellular receptors	6.0X10⁻⁹
SUMO E3 ligases SUMOylate target proteins	6.4X10⁻⁵
SUMOylation	7.7X10⁻⁵
Generic Transcription Pathway	4.4X10⁻⁴
RNA polymerase II transcription	2.1X10⁻³
Gene expression (transcription)	5.3X10⁻³

Reactome Pathway in L5	p-value
Nuclear Receptor transcription pathway	9.4X10⁻⁷
SUMOylation of intracellular receptors	1.0X10⁻⁵
Transcription of E2F targets under negative control by DREAM complex	1.0X10⁻⁴
G0 and Early G1	3.5X10⁻⁴
Binding of TCF/LEF:CTNNB1 to target gene promoters	6.0X10⁻⁴
RUNX3 regulates WNT signaling	6.0X10⁻⁴
Repression of WNT target genes	1.5X10⁻³
Transcriptional regulation of granulopoeisis	2.1X10⁻³
Transcription of E2F targets under negative control by p107 (RBL1) and p130 (RBL2) in complex with HDAC1	2.2X10⁻³
TP53 Regulates Transcription of Genes Involved in G1 Cell Cycle Arres	2.4X10⁻³
SUMO E3 ligases SUMOylate target proteins	5.4X10⁻³
SUMOylation	6.2X10⁻³
Generic Transcription Pathway	7.8X10⁻³
SMAD2/SMAD3:SMAD4 heterotrimer regulates transcription	8.6X10⁻³

Extended data Fig. 4 | IgSF coding genes and some TFs show dynamic expression over

development. a, Heat map showing relative expression of genes encoding Immunoglobulin

Superfamily proteins with dynamic expression in at least one lamina neuron-type over

development (64 genes, ≥ 2-fold change in expression between any consecutive time points, p-

value <0.05). Each row represents a different gene. Each column is a different time-point every

12h from 24 to 96 hAPF (Adult). Note the cell-type dependent patterns of expression of IgSF

coding genes. **b,** Heat map showing relative expression of dynamic transcription factors (see

Methods). Many members of the Ecdysone signaling pathway are highlighted in red. **c,** Reactome

pathway analysis of dynamic TFs showing all enriched categories (p-value < 0.01). Enrichment of

the Nuclear Receptor Transcription Pathway is highlighted in L1-L5 neurons.

 Extended data Fig. 5 | Expression of TFs in the Ecdysone cascade. a, Images showing immuno-

 staining for EcR-B1, EcR-A, Hr3 and Ftz-f1 at 12hAPF, 24hAPF, 48hAPF, 72hAPF and in adults.

Note that an anti-GFP antibody was used to stain for Ftz-f1 in fly lines carrying a BAC with GFP-

tagged Ftz-f1 and endogenous cis-regulatory sites (see Supplementary Table 1). Scale bar, 50μm.

b, Line plots of expression of TFs. **c,** Read pile-up in the EcR locus showing change of isoform

from predominantly EcR-B1 to EcR-A between 40hAPF and 60hAPF.

Extended data Fig. 6 | ATAC-Seq analysis of developing L1 neurons. a, Comparison of

replicates of bulk ATAC-Seq data of L1 neurons at 40hAPF, 60hAPF and 72hAPF. Values shown

are Spearman correlation values (see Methods). **b**, Comparison of change of ATAC-seq peak

coverage and change in expression of nearest gene. ln(fold change RPKM of nearest gene) vs

ln(fold change ATAC-seq peak coverage) between 40hAPF and 60hAPF, and 60hAPF and

72hAPF. r, Pearson's correlation coefficient. **c,** Distribution of the top 5000 peaks at each time

point between various genomic landmarks. **d,** Normalized ATAC-Seq read pile-ups are shown for

L1 neurons at 40hAPF, 60hAPF and 72hAPF for three gene loci – Fas1, CG42450, CG13698.

ATAC-Seq peaks with predicted binding sites for EcR, Hr3 and Ftz-f1 are highlighted. Shown

along with is normalized expression of these genes in control (solid line) vs EcR^{DN} or Hr3 RNAi

expressing L1 neurons (dashed lines).

72hAPF, and 96hAPF (Adult). **b,** Expression of lamina neuron-type-specific TFs ± EcR^{DN}

expression. **c,** Volcano plots showing significant gene expression changes in L1-L5 neurons

throughout development. Red dots: fold change ≥ 2 and p-value ≤ 0.05. **d,** Normalized expression

of TFs in the Ecdysone-signaling pathway in WT (solid lines) and EcR^{DN}-expressing (dashed

lines) L1-L5 neurons. *, p-value ≤ 0.05, fold change ≥ 2. **e.** Expression of EcR in EcR^{DN}-expressing

lamina neurons at 48hAPF normalized to mean expression of EcR in wildtype cells at 48hAPF

(done separately for each lamina neuron-type). Red dots, mean of data spread. Increase in EcR

expression in EcR^{DN}-expressing cells over wildtype is expected to be due to the expression of the

EcR^{DN} transgene. Note the poor induction of EcR^{DN} in L2 neurons. p-value from student's t-test

are stated in the figure for comparison between L2 and other lamina neuron-types. The difference

between EcR^{DN} expression in L2 and L3 neurons is not significant likely due to the low cell

numbers of EcR^{DN}-expressing L3 neurons. **f,** Comparison of variance of wildtype expression of

genes affected or unaffected by EcR^{DN} expression (see methods). ****, p-value < 0.0001, student's

t-test.

Expression of EcR^{DN} affected genes in wildtype lamina neurons

Extended data Fig. 8 | Cell-type dependent targets of EcR. Heat map showing relative

expression in wildtype cells of genes affected by EcR^{DN} at 24hAPF, 72hAPF and in adults. Targets

of EcR are divided into two categories: Cell-type dependent (in any pairwise comparison fold

change ≥ 2, p-value ≤ 0.05) and Common targets (no significant expression difference in any

pairwise comparison) (see Fig. 2c and Methods).

Extended data Fig. 9 | scRNA-Seq-based analysis of wRNAi (control) and EcR RNAi

expressing lamina neurons. a, tSNE plots showing wRNAi and EcR RNAi-expressing L1-L5

clusters at 24hAPF, 48hAPF, 72hAPF, and 96hAPF (Adult). **b,** Expression of lamina neuron-type-

specific TFs ± EcR RNAi expression. **c,** Volcano plots showing significant gene expression

changes in L1-L5 neurons throughout development. Red dots: fold change ≥ 2 and p-value ≤ 0.05.

d, Average expression of TFs in the Ecdysone-signaling pathway in wRNAi (solid lines) and EcR

RNAi-expressing (dashed lines) L1-L5 neurons. *, p-value ≤ 0.05, fold change ≥ 2. **e,** Image

showing optic lobe (top) stained using an antibody targeting all EcR isoforms (grey) at 24hAPF.

Box with green dotted outline marks the region containing lamina neuron cell-bodies. This region

is magnified in bottom two panels. Lamina neurons are labeled in magenta. Scale bar, 50μm.

Extended data Fig. 10 | scRNA-Seq-based analysis of wRNAi (control) and Hr3 RNAi

expressing lamina neurons. a, tSNE plots showing wRNAi and Hr3 RNAi-expressing L1-L5

clusters at 24hAPF, 48hAPF, 72hAPF, and 96hAPF (Adult). **b,** Expression of lamina neuron-type-

specific TFs ± Hr3 RNAi expression. **c,** Volcano plots showing significant gene expression

changes in L1-L5 neurons throughout development. Red dots: fold change ≥ 2 and p-value ≤ 0.05.

d, Average expression of TFs in the Ecdysone-signaling pathway in wRNAi (solid lines) and Hr3

RNAi-expressing (dashed lines) L1-L5 neurons. *, p-value ≤ 0.05, fold change ≥ 2. **e,** Image

showing optic lobe (top) stained using an antibody targeting Hr3 (grey) at 48hAPF. Box with green

dotted outline marks the region containing lamina neuron cell-bodies. This region is magnified in

bottom two panels. Lamina neurons are labeled in magenta. Scale bar, 50μm. Also see panel d for

efficiency of Hr3 knockdown.

Extended data Fig. 11 | Effect of EcR^DN, EcR RNAi and Hr3 RNAi on k-means cluster

generated gene groups. Relative expression of genes in control (aqua) or perturbation of EcR or

Hr3 activity (orange) for each gene group in L1-L5 neurons generated via k-means clustering (see

Fig. 1c, Extended data Fig. 3). Average behavior of genes in a group for control and perturbation

of EcR or Hr3 activity are shown in bold lines, while standard deviation is shown in shaded region

of the same color. Heat map shows Log_2 (Fold change) between average of WT and average of

perturbation for all comparisons where student's t-test p-value < 0.01.

Extended data Fig. 12 | Comparison of transcriptomic changes in EcR^{DN}, EcR RNAi and Hr3 RNAi. a, ln(normalized expression in wRNAi/EcR RNAi) vs ln(normalized expression in WT/EcR^{DN}) for L1-L5 neurons throughout development. **b,** ln(normalized expression in wRNAi/Hr3 RNAi) vs ln(normalized expression in WT/EcR^{DN}) for L1-L5 neurons throughout development. **a, b,** r, Pearson's correlation coefficient.

EcR-B1, Hr3 and Ftz-f1 in brain explants ± inclusion of 20E in the culture medium. Scale bar,

50μm. See Methods. **c,** Staining for protein-traps or GFP-Mimic reporter lines for dpr15, CARPB,

dpr17 and dpr6 (genes with cell-type-dependent patterns of expression and predicted to require

Ecdysone for expression from our sequencing data) ± inclusion of 20E in the medium. Scale bar,

10μm. Different lamina neurons are labeled using antibodies against lamina neuron-type-specific

TFs (see Extended data Fig. 1c). **d,** Normalized expression of dpr15, CARPB, dpr17 and dpr6 in

wildtype (solid lines) and EcR^{DN}-expressing (dashed lines) L1-L5 neurons throughout

development. *, p-value ≤ 0.05, fold change ≥ 2. Note that dpr6 shows Ecdysone-dependent

expression in L2 neurons in *ex vivo* cultured brains, but seems to be unaffected by EcR^{DN}

expression based on our sequencing data. This is consistent with low induction of EcR^{DN} in L2

neurons using the pan-lamina Gal4 driver (see Extended data Fig. 7e).

Extended data Fig. 14 | Effect of EcR^DN on neuronal morphology. a, Morphology of lamina

neurons (L1-L5) in wildtype (WT) brains and upon pan-lamina expression of EcR^{DN}, EcR^{DN} +

EcR-B1 cDNA and EcR RNAi. mAb24B10 labels M3 and M6. M3 is highlighted with yellow

arrowhead. Scale bar, 50μm. La, lamina neuropil. Me, medulla neuropil. **b,** Effect of EcR^{DN}

expression on morphology of T4/T5 neurons. Four layers in the lobula plate, a, b, c and d, are

marked with red, yellow, blue and pink arrowheads, respectively. Cartoons of one T4 (purple

neuron, top) and one T5 neuron (purple neuron, bottom) are shown to highlight wildtype

morphology. Scale bar, 20μm.

Extended data Fig. 15 | Distribution of presynaptic sites in the lamina ± EcR^{DN} expression.

a, Distribution of Brp puncta (presynaptic sites) in the lamina neuropil for wildtype (WT) and

EcR^{DN}-expressing lamina neurons. Magenta, all lamina neurons, Green, Brp puncta. Dotted line

differentiates proximal and distal halves of the lamina neuropil (see Methods). Scale bars, 50μm.

Distribution of distal BRP puncta/ cartridge for both conditions is also shown. ***, student's t test

p-value < 0.001. n, number of cartridges quantified (7 animals for wildtype, 6 animals for EcR^{DN}).

b, Normalized expression of DIP-β in wildtype (solid lines) and EcR^{DN}-expressing L1-L5 neurons

(dashed lines). *, p-value ≤ 0.05, fold change ≥ 2.

c P-value of KS test for R8 phenotype distribution

	WT	L3> EcR^{DN}	L3> NetB	L3> EcR^{DN}+ NetB
WT	1	<10^{-15}	0.65	2.2X10^{-15}
L3> EcR^{DN}	<10^{-15}	1	<10^{-15}	<10^{-15}
L3> NetB	0.65	<10^{-15}	1	<10^{-15}
L3> EcR^{DN}+ NetB	2.2X10^{-15}	<10^{-15}	<10^{-15}	1

Extended data Fig. 16 | NetB expression dynamics and dependence on Ecdysone for

expression. a, Staining for NetB (magenta) in the optic lobe at 24hAPF, 48hAPF, 72hAPF and in

Adult. M3 is highlighted with a yellow arrowhead. mAb24B10 is used as a layer marker in the

medulla. Note that maximum expression in M3 is seen at 48hAPF. **b,** Anti-NetB antibody staining

(magenta) in *ex vivo* cultured brains ± inclusion of 20E in the medium (see Methods and Extended

data Fig. 13a). In **a** and **b,** L3 neurons are labeled using myr-GFP. Scale bar, 50μm. **c,** Table

showing p-values from KS test for comparison of R8 axon depth distributions between all genetic

conditions in Fig. 3c. Note that p-value = $2.2X10^{-15}$ for comparison between L3>EcR^{DN} + NetB

and WT is indicative of an incomplete rescue of EcR^{DN} phenotype by NetB overexpression.

Extended data Fig. 17 | L1, L2, L4 and L5 – specific targets of EcR. Heat map showing relative expression of L1, L2, L4 and L5-specific targets of EcR (see Methods). Aqua, targets of transcription factor Erm identified in Peng *et al.*[32]. Magenta, Immunoglobulin Superfamily genes.

The relatively small number of L2-specific targets of EcR is likely to reflect low expression of

EcR^{DN} in these neurons (Extended data Fig. 7e).

Supplementary Tables

Supplementary Table 1. Fly strains used in this study

Supplementary Table 2. Developmental transcriptome of lamina neurons. Normalized

expression values are given for all expressed genes.

Supplementary Table 3. Transcriptome of L1 neurons at 40hAPF, 60hAPF and 72hAPF by bulk

RNA-Seq. Numbers in the table are RPKM values.

Supplementary Table 4. Gene groups generated by k-means clustering. Relative expression

values are given for all expressed genes.

Supplementary Table 5. Gene Ontology (GO) analysis of gene groups generated by k-means

clustering.

Supplementary Table 6. TF binding site analysis using i-cisTarget (see Methods).

Supplementary Table 7. ATAC-Seq analysis of L1 neurons at 40hAPF, 60hAPF and 72hAPF.

Measure of reads mapped to each peak are given as RPKM values. Peaks with differential

accessibility between consecutive time points are also given.

Supplementary Table 8. Normalized expression in WT and EcR^{DN}-expressing lamina neurons.

Genes differentially expressed between WT and EcR^{DN} are also given.

Supplementary Table 9. Gene Ontology (GO) analysis of common and cell-type dependent

targets of EcR.

Supplementary Table 10. Normalized expression in wRNAi and EcR RNAi-expressing lamina

neurons. Genes differentially expressed between wRNAi and EcR RNAi are also given.

Supplementary Table 11. Normalized expression in wRNAi and Hr3 RNAi-expressing lamina

neurons. Genes differentially expressed between wRNAi and Hr3 RNAi are also given.

Supplementary Table 12. Lists of genes likely to be involved in step 1 or step 2 of L5 axonal

morphogenesis (see Fig. 4c).

Acknowledgements

We thank Dr. Pecot (Harvard), Dr. Yamanaka (UC Riverside), Dr. Black (UCLA), Dr. De Robertis

(UCLA), Dr. Riddiford (Univ. of Washington) and Dr. Truman (Janelia Research Campus) for

helpful discussions. We thank Dr. Schuldiner (Weizmann Institute of Science) and members of the

Zipursky lab for feedback on the manuscript, and Dr. Diaz de la Loza for help with figure

illustrations. We would like to specifically acknowledge Juyoun Yoo's (Zipursky Lab) and Rachel

Hodge's (Jones Lab, UCLA) help with ATAC-Seq library prep and immunostaining. We also

thank the BSCRC Sequencing Core (UCLA) and the TCGB core (UCLA) for help with library

preparation and sequencing; the BSCRC FACS core (UCLA) and the Witte lab (UCLA) for

assistance with FACS purification of lamina neurons; and the IDRE Statistics Consulting (UCLA)

and Dr. Balliu (UCLA) for assistance with the statistical analysis of data. Reagents provided by

Dr. Akin (UCLA), Dr. Laski (UCLA), Dr. Wang (Duke-NUS), Dr. Pecot (Harvard), Dr. Thummel

(Univ. of Utah) and Dr. Bashaw (Univ. of Penn.) were critical for this work. This work was

supported by NIH T32-NS048004 Neurobehavioral Genetics Training Grant (S.J.), Helen Hay

Whitney Foundation (S.J.) and Whitcome Fellowship (Y.L.). S.L.Z is an investigator of the

Howard Hughes Medical Institute.

Author contributions

S.J., Y.L., Y.K. and S.L.Z. designed experiments. S.J., Y.L., P.M. and B.P. acquired data. S.J.,

Y.L. and Y.K. analyzed the data. S.J., Y.L. and S.L.Z. wrote the manuscript with input from all

co-authors.

Competing interest declaration

The authors declare no competing interests.

Additional information

Supplementary Information is available for this paper.

Correspondence and requests for materials should be addressed to: Dr. S. Lawrence Zipursky -

lzipursky@mednet.ucla.edu.

Additional References

50. Hoskins, R. A. *et al.* The Release 6 reference sequence of the Drosophila melanogaster genome. *Genome Res* **25**, 445–458 (2015).

51. Picelli, S. *et al.* Full-length RNA-seq from single cells using Smart-seq2. *Nat Protoc* **9**, 171–181 (2014).

52. Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nat Methods* **10**, 1213--1218 (2013).

53. Heinz, S. *et al.* Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. *Mol Cell* **38**, 576–589 (2010).

54. Xu, S. *et al.* Interactions between the Ig-Superfamily Proteins DIP-α and Dpr6/10 Regulate Assembly of Neural Circuits. *Neuron* **100**, 1369--1384.e6 (2018).