Inhibitory synaptic vesicles have unique dynamics and exocytosis properties

Chungwon Park[a], Xingxiang Chen[a], Chong-Li Tian[b, c], Gyu Nam Park[d, e], Nicolas Chenouard[f], Hunki Lee[a, g], Xin Yi Yeo[h, i], Sangyong Jung[h, j], Guoqiang Bi[b, c], Richard W. Tsien[k], and Hyokeun Park[a, l, m, 1]

[a]Division of Life Science

The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong

[b]School of Life Sciences, University of Science and Technology of China, Hefei, Anhui, China

[c]Chinese Academy of Sciences, Key Laboratory of Brain Function and Disease, University of Science and Technology of China, Hefei, Anhui, China

[d]Department of Chemistry Chung-Ang University, Seoul 06974, Korea

[e]Creative Research Initiative Center for Chemical Dynamics in Living Cells, Chung-Ang University, Seoul 06974, Korea

[f]Univ. Bordeaux, CNRS, Interdisciplinary Institute for Neuroscience, IINS, UMR 5297, F-33000 Bordeaux, France

[g]Institute of Medical Physics and Biophysics, University of Münster, Münster, 48149, Germany.

^hSingapore Bioimaging Consortium, Agency for Science, Technology and Research,

11 Biopolis Way, Singapore

ⁱDepartment of Psychological Medicine, Yong Loo Lin School of Medicine,

National University of Singapore, Singapore

^jDepartment of Physiology, Yong Loo Lin School of Medicine, National

University of Singapore, Singapore

^kNYU Neuroscience Institute and Department of Physiology and Neuroscience,

New York University, New York, NY, USA

^lDepartment of Physics,

The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon,

Hong Kong

^mState Key Laboratory of Molecular Neuroscience

The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon,

Hong Kong

¹To whom correspondence should be addressed. Email: hkpark@ust.hk

Classification

Biological Sciences, Neuroscience

real-time imaging

Author contributions

C.P. and H.P. designed the experiments. C.P., X.C., and C.-L.T., performed the

experiments. C.P, C.-L.T., G.N.P., N.C., and H.L. analyzed the data. C.P., C.-L.T.,

X.Y.Y., S.J., G.B., R.W.T., and H.P. discussed and interpreted the results. C.P., X.C.,

C.L.T., S.J., R.W.T., and H.P. wrote the manuscript. S.J., G.B., and H.P. supervised

students.

This PDF file includes:

Main Text

Figures 1 to 5

Abstract

Maintaining the balance between neuronal excitation and inhibition is essential for proper function of the central nervous system, with inhibitory synaptic transmission playing an important role. Although inhibitory transmission has higher kinetic demands compared to excitatory transmission, its properties are poorly understood. In particular, the dynamics and exocytosis of single inhibitory vesicles have not been investigated, due largely to both technical and practical limitations. Using a combination of quantum dots (QDs) conjugated to antibodies against the luminal domain of the vesicular GABA transporter (VGAT) to selectively label GABAergic (i.e., inhibitory) vesicles together with dual-focus imaging optics, we tracked the real-time three-dimensional position of single inhibitory vesicles up to the moment of exocytosis (i.e., fusion). Using three-dimensional trajectories, we found that inhibitory synaptic vesicles traveled a short distance prior to fusion and had a shorter time to fusion compared to synaptotagmin-1 (Syt1)-labeled vesicles, which were mostly from excitatory neurons. Moreover, our analysis revealed a close correlation between the release probability of inhibitory vesicles and both the proximity to their fusion site and the total travel length. Finally, we found that inhibitory vesicles have a higher prevalence of kiss-and-run fusion compared than Syt1-labeled vesicles. These results indicate that inhibitory synaptic vesicles have a unique set of dynamics and fusion properties to support rapid synaptic inhibition, thereby maintaining a tightly regulated balance between excitation and inhibition in the central nervous system.

Significance

Despite playing an important role in maintaining brain function, the dynamics of inhibitory synaptic vesicles are poorly understood. Here, we tracked the three-dimensional position of single inhibitory vesicles up to the moment of exocytosis in real time by loading single inhibitory vesicle with QDs-conjugated to antibodies against the luminal domain of the vesicular GABA transporter (VGAT). We found that inhibitory synaptic vesicles have a smaller total travel length before fusion, a shorter fusion time, and a higher prevalence of kiss-and-run than synaptotagmin-1-labeled vesicles. Our findings provide the first evidence that inhibitory vesicles have a unique set of dynamics and exocytosis properties to support rapid inhibitory synaptic transmission.

Introduction

Neurons communicate with other neurons and other cell types by releasing neurotransmitters from their presynaptic terminal via the exocytosis (i.e., fusion) of synaptic vesicles at the presynaptic membrane, subsequently activating postsynaptic receptors to mediate downstream effects (1-4). Synapses in the central nervous system can be broadly classified as either excitatory or inhibitory, depending on the type of neurotransmitters that they release and the effects of those neurotransmitters. While excitatory synapses cause the generation, propagation, and potentiation of neuronal responses for processing information (5), inhibitory synapses play an essential role in feedback and feedforward inhibition in order to control neural excitability (6). In the central nervous system, inhibitory synaptic transmission is mediated primarily by release of the neurotransmitter GABA and serves to coordinate the pattern of excitation and the synchronization of the neuronal network, thereby regulating neuronal excitability (7, 8). Thus, maintaining a tightly regulated balance between excitatory and inhibitory neurotransmission is essential for proper brain function.

An extensive analysis of the components and molecular events involved in vesicle fusion and neurotransmitter release has yielded general models describing the organization and functional properties of both presynaptic and postsynaptic components (9-11). On the presynaptic side, a transient increase in local Ca^{2+} concentration due to activation of voltage-gated Ca^{2+} channels triggers the localized buckling of the plasma membrane via a direct interaction between the C2B domain in the protein synaptotagmin-1 (Syt1) and lipids in the membrane (12-14). This leads to

the synchronous fusion between the synaptic and plasma membranes and release of the

vesicle's contents into the synaptic cleft (15), where excitatory and inhibitory

neurotransmitters act upon postsynaptic glutamate and GABA receptors, respectively.

Importantly, our general understanding of vesicle release stems from studying

excitatory neurotransmission and is currently unable to adequately explain the distinct

features associated with inhibitory transmission. For example, the size of the readily

releasable pool (RRP) of synaptic vesicles in striatal inhibitory GABAergic neurons—

probed by a hypertonic sucrose solution—is three times larger than the RRP in

excitatory hippocampal glutamatergic neurons (16). Furthermore, inhibitory neurons

have both higher average vesicular release probability (P_r) and more release sites

compared to excitatory neurons (16-18). These results suggest that quantitative

differences exist between inhibitory and excitatory synapses with respect to vesicle

dynamics, release, and recycling. In addition, disproportionate effects between

inhibitory and excitatory synapses following knockout of some or all synapsin genes

provide a molecular perspective on the putative differences between these two types of

synapses (19-22).

Despite the crucial role that inhibitory neurotransmission plays in processing

information in the neuronal network, the dynamics of inhibitory synaptic vesicles are

poorly understood. In particular, studying the dynamics of single inhibitory synaptic

vesicles in hippocampal neurons in real time has been hampered by both technical and

practical limitations, including a lack of markers to label individual inhibitory vesicles,

the complex three-dimensional structure of inhibitory synapses, and the extremely

small size of synaptic vesicles, which have an average diameter of approximately 40–

50 nm, well below the resolution of conventional light microscopy (2).

Here, we report the real-time three-dimensional tracking of single inhibitory

synaptic vesicles in cultured hippocampal neurons using dual-focus optics and QD-

conjugated antibodies against the luminal domain of the vesicular GABA transporter

(VGAT) to selectively label single inhibitory vesicles. We found significant differences

between inhibitory synaptic vesicles and Syt1-labeled vesicles with respect to the

fusion time, the total travel length prior to fusion, and the prevalence of kiss-and-run

fusion. Moreover, we found that the release probability (P_r) of inhibitory synaptic

vesicles was closely correlated with the vesicles' proximity to their fusion site and with

their total length traveled prior to fusion. These findings indicate that inhibitory

synaptic vesicles have unique dynamics and fusion properties that support their ability

to facilitate rapid neurotransmission.

Results

Real-time three-dimensional tracking of single inhibitory synaptic vesicles in

cultured hippocampal neurons.

To study the dynamics of inhibitory synaptic vesicles in mature synapses, we

tracked single inhibitory synaptic vesicles in three dimensions in real time up to the

moment of exocytosis in dissociated hippocampal neurons cultured for 14-21 days

using our previously reported strategy of loading each synaptic vesicle with a single

streptavidin-conjugated quantum dot (QD)-conjugated to biotinylated antibodies

against the luminal domain of endogenous synaptotagmin-1 (Syt1) (or Syt1-labeled synaptic vesicles); this approach provides with a spatial accuracy on the order of 20-30 nm, which is less than the diameter of a synaptic vesicle (23). To specifically load the QDs in inhibitory vesicles, we conjugated streptavidin-coated QDs to commercially available biotinylated antibodies against the luminal domain of the vesicular GABA transporter (VGAT), as shown in Fig. 1A. Upon exocytosis induced during electrical stimulation, the QD-conjugated antibodies bind to the luminal domain of VGAT, causing the QD to be taken up into the vesicle upon endocytosis. To confirm selective labeling of inhibitory vesicles, we used a CypHer5E-labeled antibodies against the luminal domain of VGAT (VGAT-CypHer5E) (24) to label spontaneously released inhibitory vesicles, revealing high colocalization between the QD and VGAT-CypHer5E fluorescence signals (Fig. 1B). Only vesicles that were colocalized with both VGAT-CypHer5E were used for analyzing inhibitory vesicles.

In our experiments, exocytosis of a QD-loaded vesicle is reflected by a sudden, irreversible loss of QD fluorescence caused by quenching with extracellular trypan blue (23). To measure the position of single QD-loaded inhibitory synaptic vesicles in the *x-y* plane, we used fluorescence imaging with one-nanometer accuracy (FIONA) (23, 25, 26); the vesicle's *z*-position was localized using dual-focus imaging (23, 27). Figure 1C shows the real-time three-dimensional position of a single QD-loaded inhibitory synaptic vesicle in a cultured neuron stimulated at 10 Hz up until the time of exocytosis; the *x-y* plane projection is overlaid on the fluorescence image of an inhibitory presynaptic terminal labeled with VGAT-CypHer5E, showing that the inhibitory

vesicle remained near the edge of the presynaptic terminal before undergoing

exocytosis. Fluorescence images of the QD-loaded inhibitory vesicle just before and

after exocytosis show near-complete, irreversible quenching of the QD 32.0 s after the

start of imaging (Fig. 1D). In addition to measuring the three-dimensional position, we

also analyzed the average fluorescence intensity (F) within the region of interest (ROI)

and the radial distance (R) between the momentary position and the fusion site using

the equation $R = \sqrt{\Delta X^2 + \Delta Y^2 + \Delta Z^2}$ (Fig. 1E). A close examination of the fluorescence

trace in Fig. 1E reveals several quantal "photoblinking" events (e.g., at approximately

8, 13, and 15 s), consistent with the presence of a single QD inside the vesicle lumen,

followed by a sharp, irreversible loss of fluorescence at 32 s due to exposure of the QD

to the quencher (trypan blue) in the external solution, indicating exocytosis. Thus, the

example shown in Fig. 1 demonstrates that the three-dimensional position of a single

inhibitory vesicle labeled with a single QD can be tracked in real time up to the moment

of exocytosis during electrical stimulation.

To determine whether QD-conjugated antibodies against the luminal domain of

Syt1 label inhibitory synaptic vesicles, we measured the colocalization between

VGAT-CypHer5E-labeled inhibitory presynaptic terminals and synaptic vesicles

loaded with QD-conjugated antibodies against Syt1. As shown in Fig. S1, we found a

relatively low percentage of colocalization between VGAT-CypHer5E terminals and

Syt1-QD–loaded vesicles (12 ± 1.4 % (N = 13 images)), suggesting that our previous

measurements of Syt1-labeled vesicles in the hippocampal neurons likely reflect the

dynamics of excitatory vesicles (23). In contrast, we found a significantly higher degree

of colocalization (61 ± 3.4% (N = 13)) between VGAT-CypHer5E–labeled terminals

and VGAT-QD–loaded vesicles (*SI Appendix,* Fig. S1B-C), confirming that VGAT

antibody–conjugated QDs label inhibitory synaptic vesicles preferentially.

Next, we examined the relative proportion of inhibitory neurons in our dissociated

hippocampal cultures by co-immunostaining the neurons for the neuronal marker

MAP2 and glutamic acid decarboxylase 67 (GAD67), a marker of inhibitory neurons

(Fig. 2A). Our analysis revealed 6 ± 0.8% (N = 23) colocalization between MAP2-

postive and GAD67-positve puncta (Fig. 2B), suggesting that our cultured hippocampal

cultures contain predominantly excitatory neurons. Moreover, we found high

colocalization (82 ± 7.7% (N = 19)) between GAD67 and parvalbumin (PV)

immunostaining (Fig. 2C), indicating that the majority of inhibitory neurons in our

cultures are PV-expressing fast-spiking GABAergic interneurons. By extension, we

conclude that our measurements of VGAT-labeled synaptic vesicles are likely to reflect

the dynamics of inhibitory synaptic vesicles in fast-spiking PV-expressing GABAergic

interneurons.

Inhibitory synaptic vesicles have unique dynamics.

To understand pre-fusion dynamics of inhibitory synaptic vesicles, we measured

the net displacement between the initial position and the fusion site of individual

vesicles while applying electrical stimuli at 10 Hz. The three-dimensional net

displacement of each vesicle was calculated as the Pythagorean displacement using our

real-time three-dimensional traces of single QD-loaded inhibitory synaptic vesicles.

Figure 3A shows the net displacements of inhibitory synaptic vesicles compared with

our previously reported net displacements of Syt1-labeled synaptic vesicles (23). Net

displacements between the initial and fusion sites of inhibitory synaptic vesicles were

not significantly different from that of Syt1-labeled synaptic vesicles ($p>0.7$,

Kolmogorov–Smirnov (K-S) test) (Fig. 3A).

Next, we measured fusion time, which we calculated as the interval between the

start of stimulation and the moment of fusion. We found that the average fusion time

was significantly shorter for inhibitory vesicles (26.2 ± 2.24 s) compared to Syt1-

labeled vesicles (44.4 ± 4.89 s) ($p<0.01$, K-S test) (Fig. 3B) indicating that inhibitory

vesicles undergo exocytosis more readily (i.e., have higher release probability) than

excitatory vesicles. Consistent with this finding, 82.5% of inhibitory vesicles (66 out of

80 vesicles measured) underwent fusion within 40 s of the start of stimulation,

compared to 53.1% (26 out of 49) excitatory vesicles, which is consistent with

inhibitory vesicles having a higher release probability compared to excitatory vesicles

(16, 17).

To understand why inhibitory vesicles fuse more readily than excitatory vesicles

yet have similar net displacement prior to fusion, we calculated the total three-

dimensional travel distance prior to fusion. We found that inhibitory vesicles traveled a

significantly shorter travel length before undergoing exocytosis compared to Syt1-

labeled vesicles (19.2 ± 1.13 μm vs. 27.9 ± 2.47 μm, respectively; $p<0.05$, K-S test;

(Fig. 3C). Taken together, these results indicate that inhibitory vesicles moved their

fusion site more straightly compared to excitatory vesicles, thereby increasing their

release probability.

248   The release probability of inhibitory vesicles is correlated with their position and

249   travel length.

Whether a synaptic vesicle's proximity to its fusion site determines its release

probability is a fundamental question in neuroscience. Using real-time tracking of Syt1-

labeled vesicles, we previously reported that the vesicle's proximity to its fusion site is

a key factor in determining the release probability (P_r) (23). However, whether the

proximity of inhibitory vesicles to their fusion sites is correlated with P_r is currently

unknown. We therefore addressed this question and found that net displacement was

significantly correlated with fusion time for inhibitory vesicles (Pearson's r = 0.77) (Fig.

4A). Moreover, the slope of the linear fit was shallower for inhibitory vesicles

compared to excitatory vesicles (0.08 ± 0.007 (Fig. 4A) vs. 0.11 ± 0.016 (*SI Appendix,*

Fig. S2A), respectively), reflecting earlier exocytosis of inhibitory vesicles. Using net

displacement, we then calculated the net velocity of releasing vesicles. We found no

significant difference in net velocity between inhibitory and excitatory vesicles (5.4 ±

0.54 nm/s vs. 4.7 ± 0.74 nm/s, respectively) (*p>0.2, K-S test*) (Fig. 4B). On the other

hand, the coefficient of variation (CV) for net velocity was smaller for inhibitory

vesicles (89%) compared to that of excitatory vesicles (110%), indicating that the net

velocity of inhibitory vesicles was less variable.

Next, we examined the correlation between fusion time and total travel length prior

to fusion and found a significant correlation for inhibitory vesicles (Pearson's r = 0.94)

(Fig. 4C). Consistent with our analysis of net displacement and fusion time, we found

that the slope of the linear fit was shallower for inhibitory vesicles compared to

excitatory vesicles (1.43 ± 0.06 (Fig. 4C) vs. 1.58 ± 0.09 (*SI Appendix,* Fig. S2B),

respectively), indicating earlier exocytosis of inhibitory synaptic vesicles. Using the

total travel length prior to fusion, we then calculated the speed (= total travel length/time

prior to fusion) of releasing inhibitory synaptic vesicles and found that the speed of

inhibitory vesicles were significantly slower compared to excitatory vesicles (405 ±

13.9 nm/s vs. 448 ± 28.1 nm/s, respectively) (*p<0.01, K-S test*) (Fig. 4D). As with net

velocity, we found that the CV for the speed was smaller for inhibitory vesicles

compared to excitatory vesicles (31% vs. 44%, respectively), indicating that the

movement of inhibitory vesicles prior to fusion is regulated more tightly compared to

excitatory vesicles. Taken together, these results suggest that inhibitory vesicles move

toward their fusion site in a more directed fashion than excitatory vesicles.

Kiss-and-run fusion is more prevalent in inhibitory synaptic vesicles.

Finally, we investigated the exocytotic fusion mode of single inhibitory

synaptic vesicles. In kiss-and-run fusion, the fusion pore opens briefly, causing a partial

drop in fluorescence due to quenching by extracellular trypan blue; in contrast, full-

collapse fusion causes the near complete loss of fluorescence (23). Figure 5A shows

the fluorescence images of a QD loaded in an inhibitory vesicle undergoing fusion,

taken before a drop in fluorescence (70.2 s), after partial drops in fluorescence (71.2

and 105 s), and after the subsequent loss of the remaining fluorescence (105.9 s). The

time course of the QD's fluorescence—obtained by analyzing an ROI containing the

QD—shows a sudden drop in fluorescence at 70.5 s (red arrow), followed by a further,

irreversible drop in fluorescence at 105.2 s (blue arrow), representing full-collapse

fusion (Fig. 5B). The fluorescence of the QD for the period after 70.5 s was reduced to

35% of the initial level, larger than the expected level for QDs steadily exposed to 2

μM TB (12%), which indicates that the inhibitory synaptic vesicle underwent kiss-and-

run fusion at 70.5 s.

We then classified all of the fluorescence traces based on their quenching

pattern and aligned the traces relative to the drop in fluorescence, revealing three

patterns: 1) near-complete loss of fluorescence (presumably representing full-collapse

fusion), 2) partial quenching followed by near-complete quenching (presumably

representing a kiss-and-run fusion event followed by full-collapse fusion or reuse after

the first kiss-and-run event), and 3) partial quenching that was not followed by near-

complete quenching (representing a kiss-and-run fusion event that was not followed by

full-collapse fusion); these three patterns are shown in Figure 5C1, C2, and C3,

respectively. The average remaining normalized fluorescence following full-collapse

fusion was smaller 0.12, suggesting near-complete quenching; in contrast, the

remaining fluorescence after kiss-and-run fusion (in both Fig. 5C2 and 5C3) was

considerably higher than 0.12. This clear difference in the remaining normalized

fluorescence after the first fusion depending on fusion mode confirms that the partial

quenching is a good indicator of kiss-and-run fusion in our experiments.

Using the degree of quenching as a measure of full-collapse fusion versus kiss-

and-run fusion, we then measured the prevalence of kiss-and-run fusion in releasing

inhibitory and excitatory vesicles. We found that approximately two thirds of inhibitory

vesicles that underwent exocytosis exhibited kiss-and-run fusion (Fig. 5D1) compared

to approximately one quarter of excitatory vesicles (Fig. 5D2), indicating that releasing

inhibitory vesicles undergo kiss-and-run fusion more frequently than excitatory vesicles.

This relatively high prevalence of kiss-and-run fusion in inhibitory vesicles may

contribute to the ability of inhibitory neurons to repeatedly release GABA during

prolonged stimulation.

Finally, we examined the dynamics of inhibitory synaptic vesicles based on

fusion modes and compared them with excitatory vesicles. We found that among the

vesicles that underwent full-collapse fusion, inhibitory vesicles traveled a shorter

distance compared to excitatory vesicles (16.9 ± 1.66 μm (n = 28) vs. 30.4 ± 3.09 μm

(n = 36), respectively) ($p < 0.05$, K-S test) (Fig. 5E). In contrast, when we examined the

vesicles that underwent kiss-and-run fusion, we found no significant difference between

inhibitory and excitatory vesicles (20.4 ± 1.48 μm (n = 52) vs. 21.0 ± 3.09 μm (n = 13),

respectively) ($p > 0.7$, K-S test) (Fig. 5F). Similarly, we found that fusion time for

inhibitory vesicles undergoing full-collapse fusion was significantly shorter than

excitatory vesicles (20.3 ± 3.04 s vs. 44.7 ± 5.84 s, respectively) ($p < 0.01$, K-S test) (Fig.

5G), but not for vesicles undergoing kiss-and-run fusion ($p > 0.4$, K-S test) (Fig. 5H).

On the other hand, the net displacement of inhibitory synaptic vesicles was not

significantly different compared excitatory vesicles regardless of the fusion mode of a

vesicle (*SI Appendix,* Fig. S4). Taken together, these results indicate that the unique

dynamics of inhibitory synaptic vesicles can be attributed primarily to the subgroup of

inhibitory vesicles that undergo full-collapse fusion.

Discussion

GABAergic inhibitory synapses differ from glutamatergic excitatory synapses with

respect to both structure and function. In the central nervous system, GABAergic

neurons play a critical role in controlling the number and activity of glutamatergic

neurons via rapid feedforward and feedback mechanisms (17, 28, 29). To allow for

rapid synaptic transmission with high fidelity, inhibitory neurons have a distinct set of

properties, including morphology, multiple release sites, biogenesis, and the trafficking

of synaptic vesicles in the presynaptic terminal (19, 30-32). Importantly, voltage-gated

Ca^{2+} channels are tightly coupled to the release machinery at GABAergic synapses,

thereby maximizing the speed and efficacy of GABAergic transmission (33). Although

the relatively high release probability and fidelity of action potential–evoked Ca^{2+}

signaling clearly play an important role in maintaining temporally precise GABAergic

transmission, the actual exocytosis mode and dynamics of single inhibitory vesicles

ranging from their fusion and recycling have not been investigated due to technical

limitations.

Here, we selectively loaded inhibitory synaptic vesicles in hippocampal neurons

with QDs conjugated to antibodies against the luminal domain of VGAT and then

tracked their three-dimensional position in real time up to the moment of exocytosis.

We then compared these dynamics with glutamatergic excitatory vesicles labeled with

QDs conjugated to antibodies against the luminal domain of Syt1. Although

approximately 10% of Syt1-labeled vesicles colocalized with VGAT-CypHer5E-

labeled presynaptic terminals, and given that Syt1 serves as the putative Ca^{2+} sensor in

both glutamatergic and GABAergic presynaptic terminals (34), our previous results

obtained with Syt1-labeled vesicles in cultured hippocampal neurons can be attributed

primarily to excitatory vesicles (23). Interestingly, in hippocampal neurons prepared

from Syt1 knockout mice, fast synchronous vesicle release was abolished in

glutamatergic presynaptic terminals but was spared in GABAergic presynaptic

terminals (35). Moreover, a subset of PV-expressing hippocampal interneurons use Syt2

as the primary Ca^{2+} sensor (36), and GABAergic inhibitory neurons can also utilize the

high-affinity Ca^{2+} sensor DOC2β (double C2-domain protein beta) to regulate

spontaneous release (34, 36). Even though the hippocampus contains more than 21

subtypes of interneurons (7), our immunostaining results indicate that less than 10% of

our cultured hippocampal neurons are inhibitory, and approximately 80% of these

inhibitory neurons express PV. Thus, our previous data obtained using Syt1-labeled

synaptic vesicles in cultured hippocampal neurons likely reflect excitatory vesicles (23).

Overall, we found that the total travel length by inhibitory vesicles was smaller

compared to excitatory (i.e., Syt1-labeled) vesicles. We also found that inhibitory

vesicles underwent exocytosis earlier than excitatory vesicles, with a larger percentage

of inhibitory vesicles undergoing release within 40 s of the onset of stimulation.

Moreover, we found that the release probability of inhibitory vesicles was closely

related to the proximity to their fusion site and their total travel length prior to fusion.

These results indicate that inhibitory synaptic vesicles reach their fusion sites by

following a more direct trajectory compared to excitatory vesicles. Interestingly, we

also found that kiss-and-run fusion is more prevalent among inhibitory vesicles

compared to excitatory vesicles. Taken together, these differences between inhibitory

and excitatory vesicles indicate that inhibitory vesicles have a distinct set of dynamics

and exocytosis properties, consistent with previous reports that inhibitory synapses

have unique electrophysiological and ultrastructural features, including a larger RRP,

rapid-onset short-term depression, and higher vesicle release probability (16, 17).

Furthermore, these properties support the efficient and rapid release of neurotransmitter

from PV-expressing inhibitory neurons, enabling increased microcircuit functions in

the brain, including feedforward and feedback inhibition as well as high-frequency

network oscillations on a rapid time scale (37). The differences between inhibitory and

excitatory vesicles with respect to their release dynamics suggest differences in how

these vesicles affect synaptic function, despite having (at least partially) overlapping

molecular release machinery. Consistent with the importance of vesicle localization and

movement, changes in the relationship between release probability and vesicle position

may alter the release of neurotransmitters in neurodegenerative diseases such as

Huntington's disease (38, 39)

The unique dynamics of single inhibitory synaptic vesicles likely contribute to

rapid inhibitory neurotransmission. Specifically, their shorter travel length prior to

exocytosis and high release probability enable inhibitory vesicles to undergo exocytosis

earlier and with higher fidelity compared to excitatory vesicles, features reflected by

their significantly shorter time to fusion. In addition, other mechanisms may also

contribute to rapid inhibitory neurotransmission. Although the precise mechanism

underlying the unique dynamics associated with inhibitory vesicles is currently

unknown, several possibilities come to mind, including distinct types and/or states of

voltage-gated Ca^{2+} channels, a larger RRP, and tight coupling between Ca^{2+} channels

and the Ca^{2+} sensor, features that may support rapid inhibitory neurotransmission and

minimize synaptic fatigue at inhibitory presynaptic terminals.

In summary, we provide the first direct evidence that inhibitory vesicles in cultured

hippocampal neurons travel less distance to reach their fusion site, undergo exocytosis

earlier, and have a higher prevalence of kiss-and-run fusion compared to excitatory

vesicles. These unique properties may explain the highly precise manner in which input

signals are converted to output signals in inhibitory neurons, providing new insights

into how inhibitory neurotransmission regulates the balance between excitation and

inhibition in the central nervous system.

Materials and Methods

Primary hippocampal neuron culture.

417 Rat (Sprague Dawley) pups at the postnatal 0-day (P0) were sacrificed to harvest CA1

418 and CA3 of the hippocampal regions of the brain tissue as previously described (23,

419 40). All procedures were performed according to the animal protocol approved by the

420 Department of Health, Government of Hong Kong. Around 2×10^4 neurons were plated

421 on each Poly D-Lysine (P7886, Sigma-Aldrich, USA)-coated cover glass with a

diameter of 12 mm (Glaswarenfabrik Karl Hecht GmbH & Co KG, Germany) in 24-well plates. Two days after plating the neurons, 10 µM uridine (U3003, Sigma-Aldrich) and 10 µM 5-fluorodeoxyuridine (F0503, Sigma-Aldrich) were added to inhibit the proliferation of glial cells in cultured neurons (41). The neurons were cultured in the Neurobasal^{TM}-A Medium (10888022, Thermo Fisher Scientific, USA) supplemented with 2.5 % FBS, 1 % Penicillin-Streptomycin (15140122, Thermo Fisher Scientific), 500 µM GlutaMAX^{TM}-I Supplement (A1286001, Thermo Fisher Scientific), and 2 % B-27^{TM} Supplement (17504001, Thermo Fisher Scientific). Cultured neurons were maintained in a 5% CO_2 incubator at 37°C before experiments were performed.

Microscopy setup.

A real-time three-dimensional tracking microscope with a custom-built dual-focus imaging optics was built as previously described (23). The dual-focus imaging optics was made up of an aperture, a beam splitter, lenses, mirrors and a right angle mirror. In the dual-focus imaging optics, the beam splitter divided fluorescent signals into two beam pathways, the focus plane of each beam pathway was determined by its lens, and two fluorescence signals from each beam pathway were imaged side-by-side in an electron multiplying (EM) CCD camera. An Olympus IX-73 inverted fluorescence microscope (Olympus, Japan) equipped with an EMCCD camera (iXon Ultra, Andor, UK) was used to acquire fluorescence signals as previously described (42). An oil immersion 100x objective with NA = 1.40 (UPlanSAPO, Olympus) was used for imaging Streptavidin-coated quantum dots (QD 625, A10196, Life Technologies)-

conjugated antibodies against VGAT and CypHer5E-conjugated antibodies against

VGAT. In order to locate two-dimensional centroids and to calculate the fluorescent

peak intensity at different focal planes (I_1 and I_2), we used custom-written IDL coded

programs (L3Harris Geospatial, USA) to fit a two-dimensional Gaussian function to

the fluorescence image. An objective scanner (P-725.2CD, Physik Instrumente (PI),

Germany) was used to generate a calibration curve based on the z-position dependence

of the normalized intensity difference $((I_1-I_2)/(I_1+I_2))$. Custom-made programs written

in LabVIEW (National Instruments, USA) were used to control an objective scanner

and acquire data on the z-position. The z-positions of quantum dots (QDs) were

obtained from the relative peak intensity differences using the empirically determined

calibration curve, as previously described (23, 27). A 405 nm laser (OBIS, Coherent

Inc., USA) and a 640 nm laser (OBIS, Coherent Inc.) were used to illuminate

streptavidin coated QDs and CypHer5E, respectively. A 15× beam expander (Edmund

Optics) and a focus lens (Newport Corporation, USA) were used to illuminate the

sample uniformly.

Real-time imaging of single inhibitory synaptic vesicles.

Imaging experiments were performed using 14 - 21 days in vitro (DIV) rat primary

hippocampal neurons at room temperature. Neurons were pre-incubated with 3 nM of

CypHer5E-tagged anti-VGAT antibodies (131 103CpH, Synaptic Systems, Germany)

in the culture medium and incubated in the 5% CO_2 incubator for 2 h at 37°C to

specifically label inhibitory synaptic vesicles in presynaptic terminals. Then, single

inhibitory synaptic vesicles were labeled by QDs conjugated with antibodies against the luminal domain of VGAT. To conjugate QDs to antibodies, Streptavidin-coated quantum dots 625 (A10196, Thermo Fisher Scientific) were incubated with biotinylated antibodies against the luminal domain of VGAT (131 103BT, Synaptic Systems) in a modified Tyrode's solution (4 mM KCl, 150 mM NaCl, 2 mM $CaCl_2$, 2 mM $MgCl_2$, 10 mM D-glucose, 10 mM HEPES, 310-315 mOsm/kg, and pH 7.3-7.4) with casein (C7078, Sigma-Aldrich) for 50 min at room temperature. After the incubation, the conjugated solution was added to a sample chamber containing a cover glass with neurons dipped in the modified Tyrode's solution. The labeling of single inhibitory synaptic vesicles with QDs was conducted by triggering evoked release of synaptic vesicles using 10 Hz electric field stimulation for 120 s, which was induced by parallel platinum electrodes connected with the SD9 Grass stimulator (Grass Instruments, USA). Following 2 min incubation after stimulation, the Tyrode's solution was superfused for 15 min at 1.0 ~ 2.0 ml/min. Subsequently, 2 µM of trypan blue (15250061, Thermo Fisher Scientific) were superfused before imaging. During the imaging, trypan blue (quencher) was kept in the chamber to quench the fluorescence of QDs exposed to the external solution during exocytosis. Fluorescence images of CypHer5E-labeled and QD-labeled synaptic vesicles were acquired using an Olympus IX-73 inverted microscope. A ZT640rdc-UF1 (Chroma Technology, USA) and an ET690/50M (Chroma Technology) were used to acquire fluorescence images of CypHer5E-labeled presynaptic boutons. A ZT405rdc-UF1 (Chroma Technology) and an ET605/70M (Chroma Technology) were used to acquire fluorescence images of QD-labeled

inhibitory synaptic vesicles. 2000 frame images were obtained using a frame-transfer

mode at 10 Hz for 200 s using an EMCCD camera. The EMCCD camera and the

stimulator were synchronized with Axon Digidata 1550 (Molecular Devices, USA) to

trigger an electric field stimulation at 10 Hz for 120 s while recording. Clampex

(program, Molecular Devices) was used to generate stimulation protocols and control

stimulation based on the protocol.

Analysis of the real-time single vesicle-tracking images.

Custom-written IDL coded programs were used to localize the centroids of raw

fluorescence images. Two-dimensional centroids and peak intensities of synaptic

vesicles loaded with QDs at two different focal planes were calculated by fitting a

Gaussian function to fluorescence images using custom-written programs using IDL.

The position along the z-axis was calculated based on a calibration curve while the

average intensity of the fluorescence of a QD within a region of interest (ROI) was

calculated using MetaMorph (Molecular Devices). Fusion positions and times of single

synaptic vesicles were determined as the first quenching positions and times of

fluorescence intensity by trypan blue during electrical stimulation similar to the

previous report (23). Two-sample Kolmogorov-Smirnov tests (K-S test) were used for

statistical analysis, using OriginPro 9.1 (OriginLab, USA). Data are presented as the

mean $\pm$ standard error of the mean (SEM). Differences with $p<0.05$ were considered

significant.

Immunofluorescence.

For immunocytochemistry, neurons at DIV14 were washed with PBS once, then fixed

with ice-cold 100% methanol for 10 minutes at -20°C. After three washes in PBS for

30 min at room temperature, cells were incubated with Chicken antibody against MAP2

(1:1000, ab5392, Abcam), Rabbit antibody against Parvalbumin (1:500, PA1-933,

Thermo Fischer Scientific) and Mouse antibody against GAD67 (1:500, MAB5406,

Sigma-Aldrich) in staining buffer (0.2% BSA, 0.8 M NaCl, 0.5% Triton X-100, 30 mM

phosphate buffer, pH 7.4) overnight at 4°C. Neurons were then washed three times in

PBS for 30 min at room temperature and incubated with Alexa Fluor conjugated

secondary antibodies (Goat anti-Chicken-Alexa 488, A11039, Life Technologies; Goat

anti-Rabbit-Alexa 568, A11011, and Donkey anti-Mouse-Alexa 647, A31571, Life

Technologies) with 1:1000 dilution in staining buffer for 2 hours at room temperature

and stained with DAPI (300 nM, D1306, Thermo Fisher Scientific) for 10 minutes at

room temperature. After washed three times in PBS for 30 min, Cover glasses were

mounted onto microscope slides with HydroMount medium (National Diagnostics).

Confocal images were acquired using a SP8 confocal microscope (Leica) with a 40x

oil objective.

Quantification of colocalization.

The biotinylated monoclonal mouse anti-Syt1 antibody (105 311BT, Synaptic

Systems) or the biotinylated anti-VGAT antibody (131 103CpH, Synaptic Systems)

was conjugated to streptavidin-conjugated quantum dots (cat. A10196, Thermo Fisher

Scientific), and vesicles were loaded as described previously (23). The rabbit polyclonal

anti-VGAT antibody conjugated to CypHer5E (cat. 131 103CpH, Synaptic Systems) were used to label inhibitory presynaptic terminals by incubation for 3 hours. QD-Syt1– loaded vesicles localized to VGAT-CypHer5E–labeled presynaptic boutons were manually counted based on colocalization between the QD and CypHer5E fluorescence signals.

Supporting information

This article contains supporting information available online.

Acknowledgments

We thank Ching Yeung Fan, Cheuk Long Frank, Lee, and Keun Yang Park for help with data analysis, and members of Park lab for helpful discussion and comments. We also thank Chenglu Tao for discussion. We thank Dr. Curtis Barrett for critically reading the manuscript and providing constructive comments. This work was supported by the Research Grants Council of Hong Kong (Grants 26101117, 16101518, A-HKUST603/17, and N_HKUST613/17) and the Innovation and Technology Commission (ITCPD/17-9).

Declaration of interest

The authors declare no conflict of interest.

REFERENCES

1. T. C. Sudhof, Calcium control of neurotransmitter release. *Cold Spring Harbor perspectives in biology* **4**, a011353 (2012).

2. C. Yu, M. Zhang, X. Qin, X. Yang, H. Park, Real-time imaging of single synaptic vesicles in live neurons. *Frontiers in Biology* **11**, 109-118 (2016).

3. V. N. Murthy, P. D. Camilli, CELL BIOLOGY OF THE PRESYNAPTIC TERMINAL. *Annu. Rev. Neurosci* **26**, 701–728 (2003).

4. T. C. Sudhof, Neurotransmitter release: the last millisecond in the life of a synaptic vesicle. *Neuron* **80**, 675-690 (2013).

5. A. Malgaroli, R. W. Tsien, Glutamate-induced long-term potentiation of the frequency of miniature synaptic currents in cultured hippocampal neurons. *Nature* **357**, 134-139 (1992).

6. S. Kim *et al.*, Loss of IQSEC3 Disrupts GABAergic Synapse Maintenance and Decreases Somatostatin Expression in the Hippocampus. *Cell reports* **30**, 1995-2005.e1995 (2020).

7. T. Klausberger, P. Somogyi, Neuronal diversity and temporal dynamics: the unity of hippocampal circuit operations. *Science (New York, N.Y.)* **321**, 53-57 (2008).

8. S. F. Owen *et al.*, Oxytocin enhances hippocampal spike transmission by modulating fast-spiking interneurons. *Nature* **500**, 458-462 (2013).

9. H. L. Atwood, S. Karunanithi, Diversification of synaptic strength: presynaptic elements. *Nature reviews. Neuroscience* **3**, 497-516 (2002).

10. P. S. Kaeser, W. G. Regehr, The readily releasable pool of synaptic vesicles. *Current opinion in neurobiology* **43**, 63-70 (2017).

11. M. Kneussel, T. J. Hausrat, Postsynaptic Neurotransmitter Receptor Reserve Pools for Synaptic Potentiation. *Trends in neurosciences* **39**, 170-182 (2016).

12. M. Yoshihara, J. T. Littleton, Synaptotagmin I functions as a calcium sensor to synchronize neurotransmitter release. *Neuron* **36**, 897-908 (2002).

13. O. D. Bello *et al.*, Synaptotagmin oligomerization is essential for calcium control of regulated exocytosis. *Proceedings of the National Academy of Sciences of the United States of America* **115**, E7624-e7631 (2018).

14. K. Grushin *et al.*, Structural basis for the clamping and Ca(2+) activation of SNARE-mediated fusion by synaptotagmin. *Nature communications* **10**, 2413 (2019).

15. G. Voglis, N. Tavernarakis, The role of synaptic ion channels in synaptic plasticity. *EMBO reports* **7**, 1104-1110 (2006).

16. R. Nair *et al.*, Neurobeachin regulates neurotransmitter receptor trafficking to synapses. *The Journal of cell biology* **200**, 61-80 (2013).

17. U. Kraushaar, P. Jonas, Efficacy and stability of quantal GABA release at a hippocampal interneuron-principal neuron synapse. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **20**, 5594-5607 (2000).

18. K. Jensen, J. D. Lambert, M. S. Jensen, Activity-dependent depression of GABAergic IPSCs in cultured hippocampal neurons. *Journal of neurophysiology* **82**, 42-49 (1999).

19. D. Gitler *et al.*, Different presynaptic roles of synapsins at excitatory and inhibitory synapses. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **24**, 11368-11380 (2004).

20. J. Feng *et al.*, Regulation of neurotransmitter release by synapsin III. *The Journal of*

neuroscience : the official journal of the Society for Neuroscience **22**, 4372-4380 (2002).

21. Y. Humeau *et al.*, Synapsin controls both reserve and releasable synaptic vesicle pools during
neuronal activity and short-term plasticity in Aplysia. *The Journal of neuroscience : the official*
journal of the Society for Neuroscience **21**, 4195-4206 (2001).

22. S. Terada, T. Tsujimoto, Y. Takei, T. Takahashi, N. Hirokawa, Impairment of inhibitory synaptic
transmission in mice lacking synapsin I. *The Journal of cell biology* **145**, 1039-1048 (1999).

23. H. Park, Y. Li, R. W. Tsien, Influence of synaptic vesicle position on release probability and
exocytotic fusion mode. *Science (New York, N.Y.)* **335**, 1362-1366 (2012).

24. Y. Hua *et al.*, A readily retrievable pool of synaptic vesicles. *Nat. Neurosci.* **14**, 833–839 (2011).

25. H. Park, E. Toprak, P. R. Selvin, Single-molecule fluorescence to study molecular motors.
Quarterly reviews of biophysics **40**, 87-111 (2007).

26. A. Yildiz, P. R. Selvin, Fluorescence imaging with one nanometer accuracy: application to
molecular motors. *Acc Chem Res* **38**, 574-582 (2005).

27. T. M. Watanabe, T. Sato, K. Gonda, H. Higuchi, Three-dimensional nanometry of vesicle
transport in living cells using dual-focus imaging optics. *Biochemical and biophysical research*
communications **359**, 1-7 (2007).

28. Y. Chagnac-Amitai, B. W. Connors, Horizontal spread of synchronized activity in neocortex and
its control by GABA-mediated inhibition. *Journal of neurophysiology* **61**, 747-758 (1989).

29. F. Manseau *et al.*, Desynchronization of neocortical networks by asynchronous release of GABA
at autaptic and synaptic contacts from fast-spiking interneurons. *PLoS Biology* **8** (2010).

30. B. High, A. A. Cole, X. Chen, T. S. Reese, Electron microscopic tomography reveals discrete
transcleft elements at excitatory and inhibitory synapses. *Frontiers in synaptic neuroscience* **7**,
9 (2015).

31. C. L. Tao *et al.*, Differentiation and Characterization of Excitatory and Inhibitory Synapses by
Cryo-electron Tomography and Correlative Microscopy. *The Journal of neuroscience : the*
official journal of the Society for Neuroscience **38**, 1493-1510 (2018).

32. X. Li *et al.*, Presynaptic Endosomal Cathepsin D Regulates the Biogenesis of GABAergic Synaptic
Vesicles. *Cell reports* **28**, 1015-1028.e1015 (2019).

33. C. Williams *et al.*, Coactivation of multiple tightly coupled calcium channels triggers
spontaneous release of GABA. *Nature neuroscience* **15**, 1195-1197 (2012).

34. N. A. Courtney, J. S. Briguglio, M. M. Bradberry, C. Greer, E. R. Chapman, Excitatory and
Inhibitory Neurons Utilize Different Ca(2+) Sensors and Sources to Regulate Spontaneous
Release. *Neuron* **98**, 977-991.e975 (2018).

35. A. M. Kerr, E. Reisinger, P. Jonas, Differential dependence of phasic transmitter release on
synaptotagmin 1 at GABAergic and glutamatergic hippocampal synapses. *Proceedings of the*
National Academy of Sciences of the United States of America **105**, 15581-15586 (2008).

36. B. Bouhours, E. Gjoni, O. Kochubey, R. Schneggenburger, Synaptotagmin2 (Syt2) Drives Fast
Release Redundantly with Syt1 at the Output Synapses of Parvalbumin-Expressing Inhibitory
Neurons. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **37**,
4604-4617 (2017).

37. H. Hu, F. C. Roth, D. Vandael, P. Jonas, Complementary Tuning of Na(+) and K(+) Channel Gating
Underlies Fast and Energy-Efficient Action Potentials in GABAergic Interneuron Axons. *Neuron*
98, 156-165.e156 (2018).

38. S. Chen *et al.*, Abnormal motion and vesicle pools of synaptic vesicles in neurons of

Huntington's Disease revealed by 3D tracking of single vesicles.   (Submitted).

39.   S. Chen *et al.*, Altered Synaptic Vesicle Release and Ca(2+) Influx at Single Presynaptic Terminals

of Cortical Neurons in a Knock-in Mouse Model of Huntington's Disease. *Frontiers in molecular*

neuroscience **11**, 478 (2018).

40.   G. Liu, R. W. Tsien, Synaptic transmission at single visualized hippocampal boutons.

Neuropharmacology **34**, 1407-1421 (1995).

41.   S. Leow-Dyke *et al.*, Neuronal toll-like receptor 4 signaling induces brain endothelial activation

and neutrophil transmigration in vitro. *J Neuroinflamm* **9**, 230 (2012).

42.   A. Alsina *et al.*, Real-time subpixel-accuracy tracking of single mitochondria in neurons reveals

heterogeneous mitochondrial motion. *Biochemical and biophysical research communications*

493, 776-782 (2017).

 FIGURE LEGENDS

Fig. 1. Real-time three-dimensional tracking of a single inhibitory synaptic vesicle

loaded with a single quantum dot (QD). (A) Schematic diagram depicting a

streptavidin-conjugated quantum dot (QD) conjugated to biotinylated antibodies

against the luminal domain of VGAT. **(B)** Colocalization of VGAT-QD–loaded

inhibitory vesicles (green) and CypHer5E-VGAT–labeled presynaptic boutons (red) in

cultured hippocampal neurons. Scale bar: 1 µm. **(C)** Three-dimensional trajectory of a

VGAT-QD–loaded inhibitory vesicle overlaid on the *x-y* plane of a CypHer5E-VGAT–

labeled presynaptic bouton. The color bar represents elapsed time; electrical stimulation

(10 Hz) started at 20 s, and the vesicle underwent exocytosis at 32.0 s. **(D)** Fluorescence

images of the VGAT-QD–loaded vesicle shown in panel C taken at the indicated times.

Scale bar: 0.5 µm. **(E)** Three-dimensional position, radial distance from the momentary

position to the fusion site (R), and fluorescence intensity (F) of the VGAT-QD–loaded

vesicle shown in panel C. Note the photoblinking events (e.g., at approximately 8 s, 13

s and 15 s), confirming the presence of one QD inside the vesicle. Electrical stimuli (10

Hz) were applied for 120 s starting at 20 s (green horizontal bar).

Fig. 2. Inhibitory neurons in cultured hippocampal cultures. (A) Representative

confocal images of cultured hippocampal neurons immunostained for microtubule-

associated protein 2 (MAP2, green), glutamic acid decarboxylase 67 (GAD67, red),

and parvalbumin (PV, yellow); the nuclei were counterstained with DAPI (blue). Scale

bar: 50 µm. **(B)** Percentage of GAD67-positive neurons among all MAP2-positive

neurons. **(C)** The percentage of PV-positive neurons among all GAD67-positive

neurons.

Fig. 3. Inhibitory synaptic vesicles have distinct spatiotemporal dynamics.

Cumulative distribution of the net displacement between the initial location and fusion

site **(A)**, fusion time **(B)**, and total length traveled **(C)** for VGAT-QD–labeled synaptic

vesicles (n = 80 vesicles) and Syt1-QD–loaded synaptic vesicles (n = 49 vesicles).

*$p<0.05$, **$p<0.01$, and NS, not significant (Kolmogorov-Smirnov test (K-S test)).

Fig. 4. The fusion time of inhibitory synaptic vesicles is correlated with net

displacement and total travel length. **(A)** Fusion time is plotted against the net

displacement; each symbol represents an individual VGAT-QD–loaded inhibitory

vesicle, and the solid red line represents the linear regression (Pearson's r = 0.77). **(B)**

Cumulative distribution of net velocity measured for VGAT-QD–labeled synaptic

vesicles and Syt1-QD–loaded synaptic vesicles. **(C)** Fusion time is plotted against the

total travel length; each symbol represents an individual VGAT-QD–loaded inhibitory

vesicle, and the solid red line represents the linear regression (Pearson's r = 0.94). **(D)**

Cumulative distribution of vesicle speed measured for VGAT-QD–labeled synaptic

vesicles and Syt1-QD–loaded synaptic vesicles. **$p<0.01$ and NS, not significant (K-

S test).

Fig. 5. Exocytotic fusion mode of inhibitory synaptic vesicles. (A) Fluorescence images taken at the indicated times for an inhibitory vesicle loaded with a VGAT-conjugated QD; electrical stimuli (10 Hz) were applied at 20 s. The intersection of the perpendicular lines marked the position before the first fusion. Scale bar: 0.5 μm. **(B)** Three-dimensional position, radial distance (R), and fluorescence (F) of a VGAT-QD–loaded vesicle that underwent kiss-and-run (K&R) fusion (red arrow) followed by full-collapse fusion (FCF, blue arrow). Electrical stimuli (10 Hz) were applied for 120 s starting at 20 s (green horizontal bar). **(C)** Average normalized fluorescence intensity traces (with SEM) time-aligned to the first fusion event for vesicles that underwent full-collapse fusion (FCF) (C1), vesicles that underwent kiss-and-run (K&R) fusion followed by FCF (C2), and vesicles that underwent K&R fusion but never underwent FCF (C3). The dotted horizontal line represents a normalized fluorescence intensity value of 0.12, (the expected normalized fluorescence right after full-collapse fusion). **(D)** Relative distribution of the fusion modes measured in VGAT-QD–labeled vesicles (D1) and Syt1-QD–loaded vesicles (D2) that underwent exocytosis, showing that 65% and 27% of vesicles, respectively, underwent K&R fusion. **(E-F)** Cumulative distribution of total travel length for inhibitory and excitatory vesicles that underwent either FCF (E) or K&R fusion (F). **(G-H)** Cumulative distribution of fusion time for inhibitory and excitatory vesicles that underwent either FCF (G) or K&R fusion (H). *$p<0.05$, **$p<0.01$, and NS, not significant (K-S test).

A Streptavidin-coated quantum dot (QD) | Biotin-labeled IgG against VGAT

B VGAT-CypHer5E | VGAT-QD | Merge

D 20.0 s | 31.9 s | 32.3 s

C

Z (nm)

X (nm)

Y (nm)

Time (s)

E Stim. at 10 Hz

Pos. (nm)

X
Y
Z

R (nm)

F (a.u.)

Time (s)