Dynamic tracking and identification of tissue-specific secretory proteins in the circulation of live

mice

Authors

Kwang-eun Kim[1,6], Isaac Park[2,6], Jeesoo Kim[3,4], Myeong-Gyun Kang[5], Won Gun Choi[1], Hyemi Shin[1],

Jong-Seo Kim[3,4*], Hyun-Woo Rhee[2,4*] and Jae Myoung Suh[1*]

Affiliation

[1] Graduate School of Medical Science and Engineering, KAIST, Daejeon, Republic of Korea

[2] Department of Chemistry, Seoul National University, Seoul, Republic of Korea

[3] Center for RNA Research, Institute for Basic Science, Seoul, Republic of Korea

[4] School of Biological Sciences, Seoul National University, Seoul, Republic of Korea

[5] Department of Chemistry, Ulsan National Institute of Science and Technology (UNIST), Ulsan,

Republic of Korea

[6] These authors contributed equally

[*] Co-corresponding authors (e-mail: jongseokim@snu.ac.kr, rheehw@snu.ac.kr, jmsuh@kaist.ac.kr)

Abstract

Here we describe *i*SLET (*in situ* Secretory protein Labeling via ER-anchored TurboID) which labels

secretory pathway proteins as they transit through the ER-lumen to enable dynamic tracking of tissue-

specific secreted proteomes *in vivo*. We expressed *i*SLET in the mouse liver and demonstrated efficient

in situ labeling of the liver-specific secreted proteome which could be tracked and identified within

circulating blood plasma. *i*SLET is a versatile and powerful tool for studying spatiotemporal dynamics

of secretory proteins, a valuable class of biomarkers and therapeutic targets.

Main text

Secretory proteins released into the blood circulation play essential roles in physiological

systems and are core mediators of interorgan communication[1]. To investigate this critical class of

proteins, previous studies analyze conditioned media from *in vitro* or *ex vivo* culture models to identify

cell type-specific secretory proteins, but these models often fail to fully recapitulate the intricacies of

multi-organ systems and thus do not sufficiently reflect *in vivo* realities[2]. In other approaches,

bioinformatic tools such as QENIE (Quantitative Endocrine Network Interaction Estimation) have been

developed[3], however, *in silico* predictions of endocrine protein factors still require many additional

layers of experimental validation. These limitations provide compelling motivation to develop *in vivo*

techniques that can identify and resolve characteristics of tissue-specific secretory proteins along time

and space dimensions.

To address this gap, we sought to utilize recently developed proximity-labeling enzymes such

as engineered biotin ligase (BioID)[4] or ascorbate peroxidase (APEX)[5]. When provided appropriate

substrates, these enzymes generate reactive biotin species, leading to *in situ* biotinylation of proximal

proteins on lysine or tyrosine residues, respectively. Thereafter, the biotinylated proteins are readily

enriched through streptavidin affinity purification and can be identified through mass spectrometry.

Recently, TurboID, a newly engineered biotin ligase, was developed to overcome the low biotinylation

efficiency of BioID in subcellular compartments[6]. TurboID exhibits a 100-fold improvement in

efficiency compared to that of BioID in endoplasmic reticulum (ER) of cultured human cells[6]. Here, we

introduce a new tool to profile tissue-specific secretory proteins by *in situ* proximity labeling of ER

lumen proteins through the catalytic actions of an ER-anchored TurboID.

To engineer a TurboID based tool for labeling secretory proteins located in the ER lumen, we

first tested the functionality of two ER lumen-targeted TurboIDs, an ER lumen-localized TurboID

(TurboID-KDEL) and an ER membrane-anchored TurboID (Sec61b-TurboID), in cultured cells. We

transfected either TurboID-KDEL or Sec61b-TurboID expression constructs, both of which also express

a V5 epitope tag, to cultured mammalian cells and analyzed biotinylated proteins in cell lysates and

culture supernatant (**Fig. 1a**). Immunofluorescence analysis of transfected cells with anti-V5 antibody

and fluorescence-conjugated streptavidin confirmed expected patterns of ER localization for both

TurboID-KDEL and Sec61b-TurboID along with their biotinylated targets (**Fig. 1b**), consistent with

results from previous ER localization studies for APEX2-KDEL and Sec61b-APEX2[7]. Analysis of

biotinylated proteins in control cell lysates revealed the presence of several endogenous biotinylated

carboxylases[8] which were not detected in culture supernatant, indicating that these carboxylases are not

secreted (**Fig. 1c**). In contrast to control cells, a broad array of biotinylated proteins was detected in

both the cell lysate and culture supernatant of TurboID-expressing cells in a biotin treatment-dependent

manner (**Fig. 1c**).

Somewhat unexpectedly, we found that TurboID-KDEL localization was not exclusive to the

ER compartment and TurboID-KDEL itself was secreted and readily detectable in the culture

supernatant of biotin treated cells (**Fig. 1c**). In contrast, the ER-anchored Sec61b-TurboID was

undetectable in the culture supernatant (**Fig. 1c** and **Supplementary Fig. 1**). These data indicate

effective retention of Sec61b-TurboID, but not TurboID-KDEL, in the ER compartment through ER

membrane-tethering action of the single transmembrane domain of Sec61b. We also confirmed that

Sec61b-TurboID robustly labeled secretory proteins without self-secretion in a HepG2 human liver cell

line, whereas TurboID-KDEL was again found to be secreted into the culture supernatant

(**Supplementary Fig. 2**).

Notably, the pattern of biotinylated proteins generated by Sec61b-TurboID in the culture

supernatant was clearly different from that of whole cell lysate, which is expected as ER-resident

proteins and secretory proteins differ in composition **(Fig. 1d)**. To further confirm the secretory pathway

origin of Sec61b-TurboID biotinylated proteins, we treated HepG2 cells expressing Sec61b-TurboID

with Brefeldin A (BFA), an inhibitor of ER to Golgi protein transport, and observed a uniform reduction

in the amount of biotinylated proteins detected in the culture supernatant **(Fig. 1e)**. Taken together, these

data indicate that Sec61b-TurboID is faithfully retained in the ER-lumen, a necessary property for *in*

vivo applications that require efficient and accurate labeling of tissue-specific secretory proteins.

Labeling kinetics determined by biotin treatment time course studies indicate that Sec61b-

TurboID efficiently labels secretory proteins in HepG2 cells by 10 min with increased labeling up to 4

hr (**Fig. 1f**). Conversely, biotin washout time course studies indicate that Sec61b-TurboID labeled

secretory proteins are largely sustained for 8 hr (**Fig. 1g and 1h**). Therefore, Sec61b-TurboID can

efficiently label classical secretory proteins in a biotin-dependent manner indicating compatibility with

kinetic studies such as classical pulse-chase labeling analyses.

Next, we applied our method, named *i*SLET, *in situ* Secretory protein Labeling via ER-

anchored TurboID, in live mice to demonstrate its *in vivo* functionality. Sec61b-TurboID adenovirus

was delivered to mice via tail vein injection to establish a liver *i*SLET mouse model and biotin was

administered to these mice to induce labeling of liver secretory proteins (**Fig. 2a**). Because TurboID has

faster labeling kinetics than BioID, we administered 24 mg/kg biotin to mice for 3 days, as compared

to 5 to 7 days of biotin administration used for previous BioID labeling studies in the postnatal brain[9,10].

Four days after Sec61b-TurboID adenovirus delivery, we observed that Sec61b-TurboID expression

was restricted to the liver (**Supplementary Fig. 3**). Liver tissues examined by histological analysis did

not reveal any obvious adverse effects due to adenoviral overexpression of TurboID and biotin

administration (**Supplementary Fig. 3**).

As expected, consistent with results obtained from the culture supernatant of Sec61b-TurboID-

expressing cell lines, endogenous biotinylated proteins were not detected in plasma samples from liver

*i*SLET mice (**Fig. 2b**). Thus, we could unambiguously detect TurboID-dependent biotinylated liver

secretory proteins in the plasma without any background (**Fig. 2b**). Interestingly, the pattern of

biotinylated proteins secreted from the liver *in vivo* was unique and clearly distinct from that of the

secretory protein profile of hepatocyte cell lines, human HepG2 and mouse AML12 (**Fig. 2c**). These

data confirm the *in vivo* functionality of Sec61b-TurboID in liver tissues as demonstrated by the

detection of biotinylated secretory protein species in the plasma of liver *i*SLET mice.

We next performed proteomic analysis of biotinylated proteins enriched from liver *i*SLET mice

plasma via liquid chromatography and tandem mass spectrometry (LC-MS/MS). Here, we followed a

previously optimized mass spectrometric identification workflow[11,12] which provides direct evidence

for biotinylated peptides identified by the mass shift of the biotinylated lysine residue. From our LC-

MS/MS data, 27 biotinylated proteins were identified in Sec61b-TurboID mouse plasma (**Fig. 2d and**

Supplementary Table 1). Representative MS/MS spectra of the biotinylated peptides are shown in

Supplementary Fig. 4. Signal peptide analysis for the biotinylated proteins with SignalP 5.0[13] revealed

that all of the detected proteins contain signal peptides required for cotranslational transport to the ER-

lumen. (**Fig. 2e**).

As expected, serum albumin (ALB) was the most abundant biotinylated protein detected from

liver *i*SLET mice plasma samples (**Fig. 2d**). Interestingly, the second most abundant protein was

pregnancy zone protein (PZP, Q61838) (**Fig. 2d**), which is also annotated under the alias alpha-2-

macroglobulin (A2M, Q6GQT1) in the UniProt database. However, *Pzp* and *A2m* are independent genes

in the mouse genome[14], and the identified peptides in our analysis were a precise match to the sequence

of PZP but not A2M (**Supplementary Fig. 4**).

We found that 93% of the proteins identified in this study are annotated as liver-enriched and

predicted as secreted plasma proteins in the Human Protein Atlas database (**Fig. 2e**). We next compared

the secretory protein profiles from liver *i*SLET mice plasma with *ex vivo* secretome studies using

primary hepatocytes[15]. While a considerable fraction (81%) of proteins were common in both (**Fig. 2e**),

fibrinogen gamma chain (FGA), complement component C8 alpha chain (C8A), histidine-rich

glycoprotein (HRG), inter alpha-trypsin inhibitor, heavy chain 4 (ITIH4) and serine protease inhibitor

A3M (SERPINA3M) were only detected in mouse plasma of liver *i*SLET mice (**Supplementary Table.**

1). Taken together, our results indicate that the liver-specific secretory protein profiles obtained from

liver *i*SLET mice are conserved in human and more accurately reflect *in vivo* physiology compared to

conventional *ex vivo* secretome analyses.

We next applied *i*SLET to characterize secreted proteomes associated with *in vivo*

pathophysiology in which endocrine signaling play an important role such as insulin resistance. S961

is an insulin receptor antagonist that induces systemic insulin resistance within 2 weeks of treatment [16].

We administered S961 and biotin for 8 consecutive days to liver *i*SLET mice generated by adenoviral

delivery of the Sec61-TurboID transgene (**Fig. 2f**). As expected, blood glucose was dramatically

increased following S691 administration (**Fig. 2g**). Proteomic analysis of biotinylated proteins from the

vehicle (PBS) or S961 group plasma identified 30 and 47 protein species, respectively (**Fig. 2h**).

Notably, 17 of the identified proteins were exclusively found in the S961 administered insulin resistant

group. Among these proteins, many have been reported to play a role in the development of insulin

resistance (**Fig. 2i**).

Alpha-2-HS-glycoprotein (AHSG), also known as Fetuin-A is elevated in serum of obese

diabetic human subjects and induces insulin resistance[17]. Fetuin-B (FETUB) is increased in type 2

diabetes patients and causes glucose intolerance[15]. Inter-alpha-trypsin inhibitor heavy chain H1 (ITIH1)

is increased in human subjects with impaired glucose tolerance or diabetes and antibody neutralization

of ITIH1 overcomes systemic insulin resistance in mice[18]. Afamin (AFM) is strongly associated with

insulin resistance, prevalence and incidence of type 2 diabetes in pooled analysis in >20,000

individuals[19]. Beta-2-glycoprotein 1 (APOH), also known as apolipoprotein H is increased in the plasma

of type 2 diabetic patients[20]. These results demonstrate that *i*SLET technology can be successfully

applied to animal disease models for the discovery of tissue-specific secreted proteins with potential

value as therapeutic targets or biomarkers.

*i*SLET is the first application of proximity labeling to dynamically track tissue-specific

secretory proteins in the circulation of live mice. Liver *i*SLET mice may be utilized to deepen our

understanding of liver endocrine signaling by investigating secretory protein profiles under various

physiological or disease conditions. Another valuable feature of *i*SLET technology is that it can be

applied to longitudinal secretome profiling studies by drawing blood samples, which contain labeled

secretome, at multiple time points from the same individual. Pre-immunodepletion of abundant plasma

proteins such as ALB and PZP can further enhance coverage of secretory protein profiles identified

from *i*SLET studies.

Furthermore, *i*SLET is a versatile and adaptable approach to discover tissue-specific secretory

proteins as tissue-specific *i*SLET expression can be achieved using a variety of existing conditional

gene expression strategies[21]. *i*SLET will be a valuable experimental tool for the identification of tissue-

specific endocrine proteins and the deconvolution of complex interorgan communication networks.

Methods

Animals. All animal experiments complied with relevant ethical regulations. Experimental protocols

for this study were approved by the institutional animal care and use committee at the Korea Advanced

Institute of Science and Technology. 10 weeks old C57BL6/J (JAX, 000664) male mice were used for

all animal experiments. Mice were housed in climate-controlled facilities under a 12 h light-dark cycle.

Standard chow diet (Envigo, 2018S) and water were provided *ad libitum*. Tissues were dissected and

fixed for histological analysis or snap-frozen in liquid nitrogen until further analysis.

Cell culture and transfection. All cell lines were purchased from the American Type Culture

Collection (ATCC; www.atcc.org) and cultured according to standard mammalian tissue culture

protocols at 37℃, 5% CO_2 in a humidified incubator. NIH-3T3 cells were cultured in DMEM (Hyclone,

SH30243.01) supplemented with 10% bovine serum (Invitrogen, 16170-078) and antibiotics (100

units/mL penicillin, 100 μg/mL streptomycin). HepG2 cells were cultured in DMEM (Hyclone,

SH30243.01) supplemented with 10% fetal bovine serum (Gibco, 16000-044), 1% GlutaMax (Gibco,

35050061) and antibiotics (100 units/mL penicillin, 100 μg/mL streptomycin). AML12 cells were

cultured in DMEM/F12 (Gibco, 11320-033) supplemented with 10% FBS, 1% Insulin-Transferrin-

Selenium (Gibco, 41400-045) and antibiotics. 293AD cells and HeLa cells were cultured in DMEM

supplemented with 10% FBS and antibiotics. For transient plasmid transfection, cells were plated at

$2.5x10^5$ cells/well in a 6-well culture plate. 24 h after plating, cells were transfected using 6 μL jetPEI

(Polyplus) and 2.5 μg GFP, TurboID-KDEL, or Sec61b-TurboID plasmids according to manufacturer

protocols.

***In vitro* biotin labeling and cell lysate preparation.** 5 mM Biotin (Sigma, B4639) stock was prepared

in DPBS with NaOH titration. 24 h after transfection or adenoviral infection, cells were washed with

PBS and culture medium was replaced with 50 μM biotin and incubated for 16 hr. For the biotin washout

experiment, the culture medium was replaced after biotin labeling. Cells were lysed by RIPA (Pierce,

89901) with Xpert Protease Inhibitor Cocktail (GenDEPOT, P3100-010) and incubated 30 min at 4℃.

Lysates were cleared by centrifugation at 16,000 g for 20 min at 4℃. The clear supernatant was used

for western blots. Protein concentrations were determined by BCA assay (Pierce, 23225).

Culture supernatant protein preparation. Cells were washed with PBS twice and the culture medium

was changed to phenol red free DMEM (Hyclone, SH30284.01) supplemented 1mM pyruvate (Sigma,

S8636) with or without 50 μM biotin. For secretory pathway inhibition, 1X GolgiPlug™ (BD, 555029),

which contains Brefeldin A, was treated with biotin. 16 h after biotin incubation, culture supernatant

was centrifuged 400 g for 5 min and the supernatant was filtered by 0.22 μm PES syringe filter

(Millipore, SLGP033RB). The filtered supernatant was concentrated by Amicon Ultra 2 mL 10K

(Millipore, UFC201024) with buffer exchange to 50 mM Tris-HCl pH 6.8. Concentrated supernatant

was used for western blots. Protein concentrations were determined by BCA assay.

Western blots. Denatured proteins were separated on 12% SDS-PAGE gels. Separated proteins were

transferred to PVDF membrane (Immobilon-P, IPVH00010). Membranes were stained with Ponceau S

for 15 min, washed with TBS-T (25 mM Tris, 150 mM NaCl, 0.1% Tween 20, pH 7.5) twice for 5 min,

and photographed. Membranes were blocked in 3% BSA in TBS-T for 1 h, washed with TBS-T five

times for 5 min each and incubated with primary antibodies, Anti-V5 (Invitrogen, R960-25, 1:10000),

Anti-GAPDH (CST, 14C10, 1:5000), in 3% BSA in TBS-T for 16 h at 4°C. Then, membranes were

washed five times with TBS-T for 5 min each and incubated with secondary anti-mouse antibodies

(Vector, PI-2000, 1:10000) or anti-rabbit antibodies (Vector, PI-1000, 1:10000) for 1 h at room

temperature. For detecting biotinylated proteins, blocked membranes were incubated with streptavidin-

HRP (Thermo, 21126, 1:15000) in 3% BSA in TBS-T for 1 h at room temperature. The membranes

were washed five times in TBS-T before detection with chemiluminescent HRP substrate (Immobilon,

P90720) and imaged on a ChemiDoc™ XRS+ system (Bio-Rad, 1708265).

Immunofluorescence staining. HeLa cells were plated on coverslips (thickness no. 1 and radius: 18

mm) and transfected with plasmids. Cells were treated with 50 μM biotin for 30 min. Cells were fixed

with 4% paraformaldehyde and permeabilized with ice-cold methanol for 5 min at -20°C. Next, cells

were washed with DPBS and blocked for 1 h with 2% dialyzed BSA in DPBS at room temperature.

Cells were incubated 1 h at room temperature with the primary antibody, Anti-V5 (Invitrogen, R960-

25, 1:5000), in the blocking solution. After washing four times with TBS-T each 5 min, cells were

simultaneously incubated with secondary Alexa Fluor 488 goat anti-mouse immunoglobulin G (IgG)

(Invitrogen, A-11001, 1:1000) and Streptavidin-Alexa Fluor 647 IgG (Invitrogen, S11226, 1:1000) for

30 min at room temperature. Cells were then washed four times with TBS-T each 5 min.

Immunofluorescence images were obtained and analyzed using a Confocal Laser Scanning Microscope

(Leica, SP8X) with White Light Laser (WLL): 470 – 670 nm (1 nm tunable laser) and HyD detector.

Adenovirus production and infection. Recombinant adenoviruses were generated as previously

described[22]. Briefly, Sec61b-TurboID was cloned to the pAdTrack-CMV shuttle vector by KpnI and

NotI digestion. The cloned shuttle vector was linearized with PmeI and transformed to BJ5183-AD-1

cells. The recombinant adenoviral plasmid was linearized with PacI and transfected to 293AD cells.

Stepwise amplification of adenovirus was performed, and adenovirus was concentrated by ViraBind™

adenovirus purification kit (Cell Biolabs, VPK-100). Adenovirus titer was measured by counting GFP-

positive cells 24 h after infection with serial dilution. For adenoviral infection, cells were plated at 2.5

x 10^5 cells/well in a 6-well culture plate. 24 h after plating, cells were infected with 1.25 x 10^6 adenoviral

GFP or Sec61b-TurboID particles.

In vivo **biotin labeling and protein sample preparation.** Approximately 10^8 adenoviral GFP or

Sec61b-TurboID particles were injected to mice via the tail vein. 24 mg/ml biotin stock was prepared

in DMSO. Vehicle (10% DMSO in PBS) or Biotin solution (2.4 mg/mL) was filtered through a 0.22

μm PES syringe filter and injected 10 μL/g (24 mg/kg) by daily intraperitoneal injection for 3

consecutive days. For acute insulin resistance model, S961 (Novo Nordisk) is injected 10 μL/g (100

nmol/kg) by daily intraperitoneal injection for 8 consecutive days, 2 hours prior to daily biotin injection.

Biotin was not administered on the last day to minimize residual biotin in blood. Blood samples were

obtained by cardiac puncture and plasma was separated in BD Microtainer® blood collection tubes (BD,

365985). Tissues were lysed and homogenized in RIPA buffer with Xpert Protease Inhibitor Cocktail

(GenDEPOT, P3100-010) by FastPrep-24™ bead homogenizer (MP Biomedicals). Lysates were

clarified by three rounds of centrifugation at 16,000 g for 20 min at 4℃ and supernatant collection. The

clear supernatant was used for western blots. Protein concentrations were determined by BCA assay.

Peptide sample preparation and enrichment of biotinylated peptides. Plasma samples were first

subjected to buffer exchange with PBS to completely remove residual free biotin via 10k MWCO

filtration for three times. The biotin depleted plasma samples were transferred and denatured with 500

μL of 8 M urea in 50 mM ammonium bicarbonate for 1 h at 37°C, and followed by reduction of disulfide

bonds with 10 mM dithiothreitol for 1 h at 37°C. The reduced thiol groups in the protein samples were

subsequently alkylated with 40 mM iodoacetamide for 1 h at 37°C in the dark. The resulting alkylated

samples were diluted eight times using 50 mM ABC and subjected to trypsinization at 2% (w/w) trypsin

concentration under 1 mM $CaCl_2$ concentration for overnight in Thermomixer (37°C and 500 rpm).

Samples were centrifuged at 10,000 g for 3 min to remove insoluble material. Then, 150 μL of

streptavidin beads (Pierce, 88816) per replicate was washed with 2 M urea in TBS four times and

combined with the individual digested sample. The combined samples were rotated for 1 h at room

temperature. The flow-through fraction was kept, and the beads were washed twice with 2 M urea in 50

mM ABC and finally with pure water in new tubes. The bound biotinylated peptides were eluted with

400 μL of 80% acetonitrile containing 0.2 % TFA and 0.1 % formic acid after mixing and heating the

bead slurry at 60°C. Each eluate was collected into a new tube. Repeat the elution process four more

times. Combined elution fractions were dried using Vacufuge® (Eppendorf) and reconstituted with 10

μL of 25 mM ABC for further LC-MS/MS experiment.

LC-MS/MS experiment of enriched biotinylated peptides. The enriched samples were analyzed with

an Orbitrap Fusion Lumos mass spectrometer (Thermo Scientific) coupled with a NanoAcquity UPLC

system (Waters, Milford) in sensitive acquisition settings. Precursor ions were acquired at a range of

m/z 400–1600 with 120 K resolving power and the isolation of precursor for MS/MS analysis was

performed with a 1.4 Th. Higher-energy collisional dissociation (HCD) with 30% collision energy was

used for sequencing with a target value of 1e5 ions determined by automatic gain control. Resolving

power for acquired MS2 spectra was set to 30k at m/z 200 with 150 ms maximum injection time. The

peptide samples were loaded onto the trap column (3 cm x 150 µm i.d) via the back-flushing technique

and separated with a 100 cm long analytical capillary column (75 µm i.d.) packed in-house with 3 µm

Jupiter C18 particles (Phenomenex, Torrance). The long analytical column was placed in a dedicated

95 cm long column heater (Analytical Sales and Services) regulated to a temperature of 45°C.

NanoAcquity UPLC system was operated at a flow rate of 300 nL/min over 2 h with a linear gradient

ranging from 95% solvent A (H_2O with 0.1% formic acid) to 40% of solvent B (acetonitrile with 0.1%

formic acid).

LC-MS/MS data processing and the identification of biotinylated peptides. All MS/MS datasets

were first subject to peak picking and mass recalibration processed with RawConverter[23]

(http://fields.scripps.edu/rawconv) and MZRefinery[24] (https://omics.pnl.gov/software/mzrefinery)

software, respectively, and then were searched by MS-GF+[25] algorithm (v.9979) at 10 ppm precursor

ion mass tolerance against the UniProt reference proteome database (55,152 entries, Mouse). The

following search parameters were applied: semi-tryptic digestion, fixed carbamidomethylation on

cysteine, dynamic oxidation of methionine, and dynamic biotinylation of a lysine residue (delta

monoisotopic mass: +226.07759 Da). The False discovery rate (FDR) was set at < 0.5% for non-

redundantly labeled peptide level and the resulting protein FDR was near or less than 1%. MS/MS

spectrum annotation for biotinylated peptides was carried out using LcMsSpectator software

(https://omics.pnl.gov/software/lcmsspectator).

Histological analysis. Mouse livers were fixed in 10 % neutral buffered formalin (Sigma, HT501128)

for 24 hr and embedded in paraffin by an automated tissue processor (Leica, TP1020). 4 μm-thick tissue

sections were obtained, deparaffinized, rehydrated, and stained with hematoxylin and eosin.

22. Luo, J. et al. *Nat. Protoc.* **2**, 1236-1247 (2007).

23. He, L. et al. *Anal. Chem.* **87**, 11361-11367 (2015).

24. Gibbons, B.C. et al. *Bioinformatics* **31**, 3838-3840 (2015).

25. Kim, S. & Pevzner, P.A. *Nat. Commun.* **5**, 5277 (2014).

Acknowledgments

We thank Dr. K.-J. Oh for assistance in adenovirus generation, Dr. K. Kim for assistance in mass

spectrometry analysis and H.-S. Jung, J.E. Kim, and H. Jung for technical support. We thank all lab

members for helpful discussions and technical assistance. This work was supported by the National

Research Foundation (NRF) of Korea grants (NRF-2018R1A2A3075389, NRF-2016M3A9B6902871)

and Global Research Laboratory (GRL) Program through the NRF funded by the Ministry of Science

and ICT (No. NRF-2017K1A1A2013124). J.-S.K. thanks to the support from Institute for Basic Science

from the Ministry of Science and ICT of Korea (IBS-R008-D1).

Author contributions

K.K., I.P., J.-S.K., H.-W.R. and J.M.S. designed research, K.K., I.P., J.K., M.K., W.G.C., and H.S.

conducted research. K.K., I.P., J.-S.K., H.-W.R. and J.M.S. wrote the paper.

Competing interests

The authors declare no competing financial interests.

References

1. Droujinine, I.A. & Perrimon, N. *Annu. Rev. Genet.* **50**, 539-570 (2016).

2. Brown, K.J. et al. *Expert Rev. Proteomic* **9**, 337-345 (2012).

3. Seldin, M.M. et al. *Cell Metab.* **27**, 1138-1155 (2018).

4. Roux, K.J. et al. *J. Cell Biol.* **196**, 801-810 (2012).

5. Rhee, H.-W. et al. *Science* **339**, 1328-1331 (2013).

6. Branon, T.C. et al. *Nat. Biotechnol.* **36**, 880-887 (2018).

7. Lee, S.Y. et al. *Cell Rep.* **15**, 1837-1847 (2016).

8. Chandler, C.S. & Ballard, F.J. *Biochem. J.* **232**, 385-393 (1985).

9. Uezu, A. et al. *Science* **353**, 1123-1129 (2016).

10. Spence, E.F. et al. *Nat. Commun.* **10**, 386 (2019).

11. Lee, S.Y. et al. *ACS Cent. Sci.* **2**, 506-516 (2016).

12. Lee, S.Y. et al. *J. Am. Chem. Soc.* **139**, 3651-3662 (2017).

13. Armenteros, J.J.A. et al. *Nat. Biotechnol.* **37**, 420-423 (2019).

14. Krause, K. et al. *Front. Microbiol.* **10**, 259 (2019).

15. Meex, R.C. et al. *Cell Metab.* **22**, 1078-1089 (2015).

16. Aguayo-Mazzucato, C. et al. *Cell Metab.* **30**, 129-142 (2019).

17. Pal, D. et al. *Nat Med.* **18**, 1279-1285 (2012).

18. Kim, T.H. et al. *Sci. Transl. Med.* **11**, eaan4735 (2019).

19. Kollerits, B. et al. *Diabetes Care* **40**, 1386-1393 (2017).

20. Castro, A. et al. *Atherosclerosis* **209**, 201-205 (2010).

21. Schnutgen, F. et al. *Nat. Biotechnol.* **21**, 562-565 (2003).

 Figure legends

Fig. 1 Proximity labeling of secretory pathway proteins using ER-anchored TurboID. a, Schematic

illustration for secretory protein labeling by ER-localized TurboID (TurboID-KDEL) or ER-anchored

TurboID (Sec61b-TurboID). **b**, Immunofluorescence localization of TurboID (Anti-V5) and

biotinylated proteins (Streptavidin-Alexa) in HeLa cells transfected with TurboID-KDEL or Sec61b-

TurboID expression plasmids. **c**, Western blots for biotinylated proteins (Streptavidin-HRP) and

TurboID (Anti-V5) in cell lysates or culture supernatant of NIH-3T3 cells transfected with GFP

(Control), TurboID-KDEL (KDEL), or Sec61b-TurboID (Sec61b) expression plasmids. Anti-GAPDH

is a loading control. Asterisk indicates self-biotinylated TurboID-KDEL or Sec61b-TurboID. **d**, Line-

scan analysis of biotinylated proteins in cell lysate (orange) or culture supernatant (black) from NIH-

3T3 cells transfected with Sec61b-TurboID expression plasmids and treated with biotin. PC, Pyruvate

carboxylase; MCC/PCC, Methylcrotonyl-CoA Carboxylase/ Propionyl-CoA carboxylase. **e**, Effect of

Brefeldin A (BFA) on biotinylated protein secretion in HepG2 cells transfected with Sec61b-TurboID

expression plasmids. **f**, Time course blot for biotinylated-labeled proteins (Streptavidin-HRP) in cell

lysates of HepG2 cells transfected with Sec61b-TurboID expression plasmids. **g**, Time course blot for

biotinylated protein (Streptavidin-HRP) turnover in cell lysates of HepG2 cells transfected with Sec61b-

TurboID expression plasmids following biotin washout. **h**, Quantitation and plotting of the time course

blot for biotinylated protein turnover shown in **g**. Asterisk indicates Sec61b-TurboID.

Fig. 2 Identification of liver-specific secretory proteins in mouse plasma from liver *i*SLET mice.

a, Experimental scheme for adenoviral expression of Sec61b-TurboID and biotin labeling in mouse

liver tissue. **b**, Streptavidin-HRP detection of biotinylated proteins and Ponceau S detection of proteins

from mouse plasma after adenoviral delivery of Sec61b-TurboID. **c**, Biotinylated secretory protein

profiles generated by Sec61b-TurboID in supernatants of hepatocyte cell lines, HepG2 and AML12,

and plasma of liver *i*SLET mice. **d**, Relative abundance of biotinylated secretory proteins detected in

plasma of liver *i*SLET mice. ALB, Serum albumin; PZP, Pregnancy zone protein; TF, Serotransferrin;

SERPINA3K, Serine protease inhibitor A3k; MUG1, Murinoglobulin-1; FGA, Fibrinogen alpha chain;

APOA1, Apolipoprotein A-I; FGG, Fibrinogen gamma chain; HPX, Hemopexin. **e,** Specificity analysis

for biotinylated proteins with SignalP 5.0, human protein atlas and literature. **f,** Experimental scheme

for adenoviral expression of Sec61b-TurboID and biotin labeling in the S961-induced insulin resistance

model. **g,** Blood glucose in Vehicle or S961 injected mouse. n=3 per group. **h,** Biotinylated secretory

proteins detected in plasma of Vehicle or S961 injected liver *i*SLET mice. AHSG, Alpha-2-HS-

glycoprotein; FETUB, Fetuin-B; ITIH1, Inter-alpha-trypsin inhibitor heavy chain H1; AFM, Afamin;

APOH, Beta-2-glycoprotein 1; ORM1, Alpha-1-acid glycoprotein 1; EGFR, Receptor protein-tyrosine

kinase; CFB, Complement factor B; CES1B, Carboxylic ester hydrolase; C4B, Complement C4-B; C5,

Complement C5; C6, Complement component 6; C8B, Complement component C8 beta chain; F13B,

Coagulation factor XIII B chain; ITIH4, Inter alpha-trypsin inhibitor, heavy chain 4; KLKB1, Plasma

kallikrein; PON1, Serum paraoxonase/arylesterase 1. **i**. Representative candidates related with insulin

resistance detected in this study.

Kim et al. Figure 1

a

ER Lumen   Cytosol

KDEL

Sec61b

Cell lysate

Culture supernatant

b

| TurboID-KDEL | Sec61b-TurboID |

Anti-V5

Streptavidin

c

	Cell lysate			Culture supernatant		
	Control	KDEL	Sec61b	Control	KDEL	Sec61b
Biotin	− +	− +	− +	− +	− +	− +

Streptavidin-HRP

Anti-GAPDH
37
kDa

Anti-V5
37

* Self biotin-labeled TurboID

d

Biotin-labeled proteins by Sec61b-TurboID

Cell lysate
Culture supernatant

PC   MCC/PCC

Normalized intensity (AU)

Molecular weight (kDa)

e

	Cell lysate		Culture supernatant
Sec61b-TurboID	+	+	+ +
	−	BFA	− BFA

Streptavidin-HRP

kDa

Anti-V5   n.s.
50
37

* Sec61b-TurboID
n.s. Non-specific band

f

Biotin treatment

| 0 | 10min | 0.5hr | 1hr | 2hr | 4hr |

kDa

Streptavidin-HRP   *

37
Anti-V5

* Sec61b-TurboID

g

Untransfected   **Biotin washout**

| 0hr | 1hr | 2hr | 4hr | 8hr | 24hr | 48hr |

kDa

Streptavidin-HRP   *

37
Anti-GAPDH

* Sec61b-TurboID

h

$y = -0.324\ln(x) + 1.2911$
$R^2 = 0.9706$

Streptavidin/GAPDH (AU)

Washout time (hour)

Kim et al. Figure 2

a Day 0 — Adeno-TurboID delivery; Day 1~3 — Biotin administration; Day 4 — Plasma harvest

b Plasma — AdV: -, GFP, TurboID; Biotin: -, +, -, +; Streptavidin-HRP; Ponceau S

c Biotin-labeled secretory proteins — HepG2, AML12, Liver; Streptavidin-HRP

d *In vivo* secretory protein composition of liver — ALB 33%, PZP 11%, TF 11%, SERPINA3K 9%, MUG1 6%, FGA 5%, APOA1 4%, FGG 4%, HPX 3%

e Signal peptide (SignalP 5.0): 27/0; Liver elevated: 25/2; Predicted secreted: 25/2; Plasma: 25/2 (Human Protein Atlas); Primary hepatocyte: 22/5 (Meex et al, 2015)

f Day 0 — Adeno-TurboID delivery; Day 1~8 — Vehicle or S961 injection 2 hours after, Biotin administration; Day 9 — Plasma harvest

g Blood glucose (mg/dl); Vehicle, S961; Day 1 (Basal), Day 8

h S961 (47); Vehicle (30); 30, 17
S961 only detected proteins (17)
AHSG, FETUB, ITIH1, AFM, APOH, ORM1, EGFR, CFB, CES1B, C4B, C5, C6, C8B, F13B, ITIH4, KLKB1, PON1

i

Gene	Reported description
AHSG	Increased in serum of diabetic human subjects Causes insulin resistance
FETUB	Increased in plasma of diabetic human subjects Causes impaired glucose metabolism
ITIH1	Increased in serum of diabetic human subjects Causes impaired glucose metabolism
AFM	Associated with insulin resistance, prevalence and incident of type 2 diabetes
APOH	Associated with metabolic syndrome in type 2 diabetic patients