Robust differentiation of human enteroendocrine cells from intestinal stem cells

Daniel Zeve[1,2], Eric Stas[1,†], Joshua de Sousa Casal[3,4,5,†], Prabhath Mannam[1], Wanshu Qi[1],

Xiaolei Yin[6,7,#], Sarah Dubois[1,8], Manasvi S. Shah[1,2], Erin P. Syverson[2,3], Sophie Hafner[1], Jeffrey

M. Karp[5,7,9], Diana L. Carlone[1,2,9], Jose Ordovas-Montanes[3,4,5,9], David T. Breault[1,2,9,*]

[1]Division of Endocrinology, Boston Children's Hospital, Boston, MA 02115, USA

[2]Department of Pediatrics, Harvard Medical School, Boston, MA 02115, USA

[3]Division of Gastroenterology, Hepatology, and Nutrition, Boston Children's Hospital, Boston,

MA 02115, USA

[4]Program in Immunology, Harvard Medical School, Boston, MA 02115, USA

[5]Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA

[6]David H. Koch Institute for Integrative Cancer Research, Massachusetts Institute of

Technology, Cambridge, MA 02139, USA

[7]Center for Nanomedicine and Division of Engineering in Medicine, Department of Medicine,

Brigham and Women's Hospital, Harvard Medical School, Harvard-MIT Division of Health

Sciences and Technology, Boston, MA 02115, USA

[8]School of Arts and Sciences, MCPHS University, Boston, MA 02115, USA

[9]Harvard Stem Cell Institute, 7 Divinity Avenue, Cambridge, MA 02138, USA

[†]Authors contributed equally to this work

[*]Corresponding author, email: david.breault@childrens.harvard.edu

[#]Current address: Institute for Regenerative Medicine, Shanghai East Hospital, Frontier Science

Center for Stem Cell Research, School of Life Sciences and Technology, Tongji University,

Shanghai, China

ABSTRACT

Enteroendocrine (EE) cells are the most abundant hormone-producing cells in humans and are critical regulators of energy homeostasis and gastrointestinal function. Challenges in converting human intestinal stem cells (ISCs) into functional EE cells, *ex vivo*, have limited progress in elucidating their role in disease pathogenesis and in harnessing their therapeutic potential. To address this, we employed small molecule targeting of the endocannabinoid receptor signaling pathway, JNK and FOXO1, known to mediate endodermal development and/or hormone production, together with directed differentiation of human ISCs from the duodenum and rectum. We observed marked induction of EE cell differentiation and gut-derived expression and secretion of SST, 5HT, GIP, CCK, GLP-1 and PYY upon treatment with various combinations of three small molecules: rimonabant, SP600125 and AS1842856. Robust differentiation strategies capable of driving human EE cell differentiation is a critical step towards understanding these essential cells and the development of cell-based therapeutics.

INTRODUCTION

Enteroendocrine (EE) cells are found throughout the gastrointestinal (GI) tract and represent the most abundant hormone-producing cell type within mammals. EE cells, as a whole, secrete a large variety of hormones, including glucose-dependent insulinotropic polypeptide (GIP, from K cells), cholecystokinin (CCK, from I cells), serotonin (5HT, from enterochromaffin cells), somatostatin (SST, from D cells), peptide YY (PYY, from L cells), and glucagon-like peptide-1 (GLP-1, from L cells), among others[1,2]. In response to physiological and nutritional cues, EE cells, through the production of these various hormones, are responsible for regulating multiple aspects of GI activity and nutritional homeostasis[1,2]. Because of this, EE cells have been implicated in the pathogenesis of GI diseases such as irritable bowel syndrome and inflammatory bowel disease, as well as metabolic diseases such as type 2 diabetes mellitus[3-5].

Similar to other mature intestinal epithelial cells, EE cells are derived from ISCs, which reside

within the crypts of Lieberkuhn[6]. In recent years, much progress has been made understanding

the mechanisms underlying ISC self-renewal and differentiation using 3D-organoid culture[7-10].

Using combinations of growth factors and small molecules targeting specific transcriptional

regulators and signaling pathways, intestinal organoids can either be maintained predominantly

as ISCs or differentiated into mature intestinal cells of either the absorptive or secretory

lineages[11,12]. For example, maintenance of ISC self-renewal requires activation of canonical Wnt

signaling using WNT3a and R-spondin, suppression of bone morphogenic protein (BMP)

signaling using Noggin, and inhibition of p38 MAPK signaling using the small molecule

SB202190[9-11]. By altering these pathways along with transcriptional regulators, strategies have

begun to emerge to direct ISC differentiation into mature intestinal epithelial cell types[8,9,13].

EE cells are defined by expression of the specific hormone each produces and their location

along the GI tract[2,14-16]. These cells also express multiple neuroendocrine secretory proteins,

including Chromogranin A (CHGA), some of which are more specific to certain EE cell types

than others[14,15]. Further, multiple transcription factors are critical for EE cell differentiation and

function, including neurogenin 3 (NEUROG3), neuronal differentiation 1 (NEUROD1), and

paired box 4 (PAX4)[1,2,10]. Pancreatic and duodenal homeobox 1 (PDX1) has also been shown to

regulate gene expression in a subpopulation of EE cells located within the duodenum[17]. The

directed differentiation of human EE cells requires the removal of Wnt ligands and inhibition of

Notch signaling but, in contrast to mice, human EE cells do not differentiate in the presence of

p38 MAPK inhibitors[9,10,16]. Through the use of an inducible NEUROG3 system, multiple studies

have shown induction of the human EE cell lineage, with expression and secretion of multiple

hormones in organoid cultures of both the small and large intestine[14,18-21]. However, while strong

mRNA expression of human mature EE markers has been observed multiple times using small

molecule differentiation protocols[8,9,16,22-24], confirmation using protein expression levels is less

frequent[16,22,24], being most thoroughly explored in studies evaluating the roles of short chain

fatty acids[22] and isoxazole-9[24] in human intestinal organoid differentiation and hormone

production.

Additional factors and signaling pathways have also been implicated in EE cell formation and

function. For example, inhibition of Forkhead box protein O1 (FOXO1), a transcription factor

critical for stem cell function and energy homeostasis[25,26], has been associated with

upregulation of multiple endocrine-associated transcription factors and hormones, including

NEUROG3 and GIP[27-29]. In human intestinal organoids (HIOs) derived from induced pluripotent

stem cells, transduction of a dominant negative FOXO1 yielded increased numbers of CHGA+

cells, less 5HT-positive cells, and no change in GLP-1 or SST-positive cells when compared to

control organoids at 230 days[30]. The endocannabinoid signaling pathway, acting through the

cannabinoid receptor type 1 (CB1), has also been shown to regulate EE cell function, with

studies showing activation of CB1 inhibiting CCK secretion[31,32]. Additionally, rimonabant (Rim),

a highly selective CB1 inverse agonist, has been shown to increase human serum GIP levels[33].

C-Jun N-terminal Kinase (JNK) signaling, which has been implicated in regulating ISCs[34,35], also

regulates beta cell function through its actions on PDX1[36,37], and has been suggested to

upregulate EE cell differentiation, with deletion of JNK2 reducing the number of CHGA-positive

(CHGA+) cells in mouse organoids[38]. The role of JNK in the regulation of EE cell differentiation

and PDX1 expression has not been studied in directed differentiation of human ISCs. Further,

BMP4 has been shown to induce the expression of various EE hormones in human intestinal

organoids[16]. However, aside from BMP4, how the above factors and pathways impact directed

differentiation of EE cells has been largely unstudied.

Here, we establish robust EE cell differentiation protocols for human duodenal organoids

(enteroids) and rectal organoids (rectoids) using various combinations of small molecules

(rimonabant, SP600125 and AS1842856) suggested to alter the activity of CB1, JNK, and

FOXO1, respectively. In enteroids, treatment with Rim and SP600125 (SP) leads to marked

induction of *CHGA* expression and a corresponding increase in the number of CHGA+ cells, as

well as in the expression of multiple hormones, including SST, 5HT, CCK, and GIP. Separately,

treatment with AS1842856 (AS) also induced SST, 5HT, and GIP, with an even stronger

induction of CHGA+ cells. Comparing levels of hormone secretion, AS leads to a greater

induction of 5HT secretion, but less GIP, whereas the combination of Rim and SP (RSP) leads

to a greater induction of GIP secretion, but less 5HT. The combination of AS treatment followed

by RSP leads to even higher expression of *SST*, and increased secretion of GIP and 5HT, when

compared to RSP treatment alone. In rectoids, treatment with base differentiation media (DM)

alone induces an increase in the number of CHGA+ cells, as well as increased expression,

production and secretion of GLP-1 and PYY, EE hormones predominantly produced in the distal

GI tract[2]. Together, these robust EE cell differentiation protocols, employing only small

molecules, provide important insights into human EE cell differentiation, with potentially

important implications for understanding disease pathogenesis and development of cell-based

therapeutics.

RESULTS

Differentiation Media Induces *CHGA* Expression in Human Duodenal Enteroids

To optimize human EE cell differentiation, we developed a DM (Supplementary Table 1) by

combining specific components from published protocols used to induce EE cells[9,12,16,23],

including Wnt3a and two additional small molecules associated with endocrine cell

differentiation: (1) betacellulin, a ligand of both EGFR and the ErbB4 receptors[39,40] and (2)

PF06260933 (PF), a small molecule inhibitor of MAP4K4[41]. Human enteroids were first cultured

in growth media (GM, Supplementary Table 1) for two days, allowing for ISC expansion,

followed by DM for 12 days (G2D12). Enteroids maintained for 14 days in GM (G14) were used

as controls to assess for changes in gene expression. In contrast to previous strategies aimed

at generating EE cells over a five-day period[9,16], exposure to DM for 12 days was compatible

with maintenance of enteroid structural integrity (Fig 1a), possibly due to the continued

presence of WNT3a.

To quantify the effectiveness of DM to induce EE cell differentiation, we profiled the gene

expression of *CHGA* and other lineage markers, including atonal BHLH transcription factor 1

(*ATOH1*, secretory progenitor cells), mucin 2 (*MUC2*, goblet cells), lysozyme (*LYZ*, Paneth

cells), intestinal alkaline phosphatase (*ALPI*, enterocytes) and leucine-rich repeat-containing G-

protein coupled receptor 5 (*LGR5*, ISCs). In addition, we assessed the expression of

transcription factors required for duodenal EE differentiation and function (*PDX1, PAX4,*

NEUROG3, and *NEUROD1*), as well as hormones secreted from the duodenum (*CCK*, *SST*,

and *GIP*). To allow for relative comparison to native tissue expression levels, total RNA from

whole duodenal mucosal biopsies was included in each qPCR analysis. Exposure of enteroids

to G2D12 induced consistent expression of *CHGA*, *PAX4*, *SST*, and *GIP*, with significantly

higher levels of *ALPI* and significantly lower expression of *LGR5* when compared to G14 (Fig 1b

and Supplementary Fig 1a). Despite G2D12's ability to induce EE and enterocyte markers when

compared to G14, their overall expression levels remained considerably lower than whole

duodenal mucosa. Levels of *PDX1*, *NEUROD1*, *NEUROG3*, *CCK*, *ATOH1*, and *LYZ* were

unchanged compared to G14, with *MUC2* showing a trend towards higher expression in G2D12

enteroids (Fig 1b and Supplementary Fig 1a). Finally, despite induction in *CHGA* mRNA levels

in response to G2D12, analysis of CHGA protein using immunofluorescence staining and flow

cytometric analysis revealed fewer than 0.1% of all cells to be CHGA+, showing a trend towards

higher expression compared to G14 (Fig 1b-1e).

Removal of WNT3a has been shown to aid EE differentiation[9,10,16]. Analysis of enteroids

cultured in DM without WNT3a (G2D12-Wnt) revealed undetectable expression levels of *CHGA*

and *SST*, reduced *MUC2* expression, and increased *ALPI* expression when compared with

enteroids differentiated with WNT3a (G2D12+Wnt) (Fig 1f). To determine whether *CHGA*

expression was increased at an earlier time-point during the 14-day differentiation protocol, we

performed time-course studies and found that *CHGA* expression was undetectable in enteroids

cultured in G2D12-Wnt, while those exposed to G2D12+Wnt showed expression after six days

of starting DM (Supplementary Fig 1b). Furthermore, two of the three enteroid lines exposed to

G2D12-Wnt showed low total RNA levels around the eighth day of differentiation, consistent

with failure to maintain these lines (Supplementary Fig 1c). Therefore, we concluded that the

presence of WNT3a in DM is beneficial to sustain enteroids in long-term culture (14 days) and is

not detrimental to EE differentiation. We also evaluated the impact of including betacellulin and

PF in our DM and found that both factors led to increased expression of EE cell markers

compared with GM and DM without betacellulin or PF, as well as maintaining *ALPI* levels when

compared to DM (Supplementary Fig 1d and 1e). Together, these data indicate that our

differentiation protocol markedly induced expression of some EE cell marker genes (e.g., *CHGA*

and *PAX4*), but is not sufficient to induce a significant increase in the number of CHGA+ EE

cells compared to undifferentiated controls.

Treatment with Rimonabant and SP600125 Induces Duodenal EE Lineage

Differentiation

To identify additional strategies that might further induce EE cell differentiation, we focused on

small molecules shown to target endocannabinoid signaling and PDX1 activity. First, we utilized

Rim, a highly selective CB1 inverse agonist. In parallel, we used the small molecule SP to inhibit

JNK signaling, which has been shown to suppress PDX1 activity[36,37].

Separately, both Rim and SP induced expression of multiple EE lineage markers (*CHGA,*

NEUROG3, SST, and *GIP*) when added to DM, with Rim having a much larger effect

(Supplementary Fig 2a). Together, RSP yielded even further increases in *SST* and *GIP*

expression compared to Rim or SP alone. PDX1 expression was unchanged under all

experimental conditions (Supplementary Fig 2a). Based on these results, Rim and SP were

used in combination for all subsequent experiments. The addition of RSP to DM maintained

overall enteroid structural integrity during the 14-day differentiation protocol (Fig 2a and

Supplementary Fig 2b). Moreover, compared to enteroids grown in G14 and G2D12, treatment

with RSP led to the upregulation of multiple EE markers (*CHGA*, *PAX4*, *NEUROD1*, *NEUROG3*,

CCK, *SST*, and *GIP*) to levels approximating whole duodenal mucosa (Fig 2b). Other lineage

markers were also increased with RSP exposure, including *ATOH1*, *MUC2*, *LYZ*, *ALPI*, and

LGR5 when compared to G14 and *ATOH1*, *MUC2*, and *LGR5* when compared to G2D12

(Supplementary Fig 2c). Immunofluorescence staining for CHGA showed multiple positive cells

within individual enteroids (Fig 2c), with a large majority of enteroids (83%) containing CHGA+

cells (Fig 2d). By comparison, only 1.0% of enteroids grown in G2D12 were CHGA+ (Fig 2d).

Quantitative flow cytometric analysis revealed 1.4% of all cells treated with RSP were CHGA+,

almost seven times the number seen with G2D12 alone (Fig 2e).

Treatment with AS1842856 Induces Duodenal EE Lineage Differentiation

Next, we assessed the impact of adding AS, a well-described FOXO1 inhibitor[42], to our

differentiation protocol. Addition of AS led to the formation of small, spherical enteroids (Fig 3a

and Supplementary Fig 3a). Compared to enteroids grown in G14 and G2D12, AS treatment led

to the upregulation of multiple EE markers (*CHGA, PAX4, PDX1, NEUROD1, NEUROG3, SST,*

and *GIP*), many to levels approximating whole duodenal mucosa. The only exception to this was

CCK, which was significantly lower in AS when compared to G14 (Fig 3b). In addition, AS

increased expression of other secretory and ISC markers, including *ATOH1, MUC2,* and *LGR5,*

when compared to G14 and G2D12 (Supplementary Fig 3b). Interestingly, the use of AS

reduced expression of the enterocyte marker *ALPI* when compared to G14 and G2D12,

suggesting a possible role for FOXO1 inhibition in the induction of the secretory lineage

(Supplementary Fig 3b). Immunofluorescence staining revealed CHGA+ cells within a large

majority of individual enteroids (85%) (Fig 3c and 3d). By comparison, only 3.5% of enteroids

grown in G2D12 had CHGA+ cells (Fig 3d). Quantitative flow cytometric analysis revealed 5.2%

of all cells exposed to AS to be CHGA+, almost 50 times the number seen with G2D12 alone

(Fig 3e).

Treatment with Rimonabant/SP600125 or AS1842856 Promote Distinct Enteroid

Differentiation Trajectories

To better assess the transcriptomic distinctions between cells from the various culture

conditions, we utilized single cell RNA sequencing (scRNA-seq) on the 10X platform combined

with sample hashing, allowing us to assess the reproducibility of our findings from distinct

donors**.** Enteroids from three individuals were grown in G2D12, RSP or AS, for a total of nine

samples. After the 14-day protocol, enteroids were isolated and processed into a single-cell

suspension and their condition and patient source were marked using Hashtag antibodies prior

to pooling the samples together[43]. After labeling, cells were processed, and single-cell

transcriptomes were generated.

By using hashtag labeling, we were able to categorize cells into singlets (significant detection of

one antibody), doublets (significant detection of more than one antibody) or negative droplets

(no significant detection of any antibodies). Negative droplets were found to have a reduced

number of both detected genes and unique genes or unique molecular identifiers (UMIs),

supporting the idea that the majority of these datapoints do not correspond to an individual cell

(Supplementary Fig 4a). In contrast, both singlets and doublets had appreciable numbers of

both genes and UMIs detected. In addition, we found that they had similar UMIs, percentage of

mitochondrial RNA, which aids in identification of low-quality cells, and detected gene

distributions (Supplementary Fig 4a). However, upon dimensional reduction, we found that

singlets and doublets formed distinct groupings in both t-distributed stochastic neighbor

embedding (tSNE) visualization (Supplementary Fig 4b), as well as uniform manifold

approximation and projection (UMAP) visualization and clustering (Supplementary Figs 4b and

4c), indicating that although they had similar RNA content, singlets likely correspond to true

singlets, whereas doublets correspond to multiple cell types analyzed together in a single

droplet.

After removal of doublets, negative droplets, and low-quality cells (gene count < 200,

percentage of mitochondrial RNA > 25%), the resulting dataset consisted of 14,767 cells spread

across the three different culture conditions. UMAP visualization revealed distinct cell separation

between G2D12 and AS treatment with minimal overlap between the two populations (Fig 4a).

In contrast, enteroid treatment with RSP generated populations that overlapped with the two

other culture conditions (Fig 4a). To further characterize the developing populations within each

treatment, we performed Louvain clustering and identified 15 clusters within the combined

dataset (Supplementary Fig 4d). Of these 15 clusters, some were composed of cells sourced

from all three culture conditions, such as clusters 2, 4, 9, and 7, whereas other clusters were

primarily formed by cells from one treatment group, such as cluster 8 (AS) and cluster 3

(G2D12). To assist with manual annotation of each cluster, we calculated the top markers of

each Louvain cluster (Supplementary Fig 4e; Wilcoxon rank sum test, $p_{adj}<1.38\times10^{-21}$ for

displayed markers)[44]. By using these markers, as well as classical markers of various epithelial

populations[45,46], we broadly classified the data into six populations of epithelial cells, including

stem cells, proliferating progenitor cells, progenitor cells, enterocytes, goblet cells and

enteroendocrine cells (Fig 4b). ISCs were identified by expression of *LGR5, AXIN2* and

ASCL2[47]. Enterocytes were identified by the expression of *KRT20*, *FABP2*, and *FABP1*, goblet

cells by the expression of *MUC2, FCGBP,* and *GFI1*, and EE cells by expression of *CHGA*,

NEUROG3, *NEUROD1*, and *PAX4*[48,49] (Fig 4c). Finally, progenitor cells were identified as

having a mixture of stem cell markers, albeit at a lower level than the ISCs, while also

expressing mature enterocyte markers. We were able to distinguish proliferating cycling

progenitors from other progenitor cells by expression of the proliferation marker *MKI67*[50] (Fig 4b

and 4c); however, some progenitor markers were also lowly expressed in secretory cell

populations, suggesting that these groups likely also contain secretory progenitors.

To understand how different culture conditions impacted enteroid development and

differentiation, we first compared the proportions of the various cell types across each condition

(Fig 4d). We found that treatment with G2D12 primarily gave rise to cells of the absorptive

lineage, with 58% identified as enterocytes. The remaining cells exposed to G2D12 consisted of

stem cells (5.7%), proliferating progenitor cells (5.4%) and progenitor cells (30%). G2D12 did

not give rise to secretory cells, with almost no detection of goblet cells (0.6%) or EE cells

(0.2%). In contrast to G2D12, treatment with either RSP or AS led to a decrease in the

proportion of enterocytes (RSP: 17%, AS: 2.5%) and an increase in stem cells (RSP: 22%, AS:

42%), goblet cells (RSP: 4.6%, AS: 1.5%), and EE cells (RSP: 4.3%, AS: 18%), with significant

differences noted between G2D12 and AS conditions for stem cells, enterocytes, and EE cells.

To further characterize the differentiation path of each culture condition, we used scVelo to

model the RNA velocity and differentiation trajectory of each cell type[51]. Briefly, RNA velocity

analysis pipelines, such as scVelo, leverage the fact that scRNA-seq can distinguish between

un-spliced and spliced variants of each gene. Since recently transcribed RNA is present as an

un-spliced variant which is then spliced into its mature form before eventually being degraded, it

is possible to calculate the dynamic kinetics of each individual gene's expression across all cells

within the dataset based on the ratio of un-spliced to spliced RNA[51,52]. scVelo uses the

collection of gene kinetics to infer the direction and speed of cell state changes on a population

level. We applied this method to each culture condition and generated vector estimates of

cellular transition overlaid on each UMAP (Fig 4e). In all three conditions, the majority of vector

arrows initiate from a stem cell cluster in the lower half of the UMAP. In the G2D12 culture

condition, cells primarily move towards proliferating progenitors and progenitor cells, before

moving towards the enterocyte cluster. In contrast, in AS treated cells, the vector arrows

proceed through a completely different trajectory, with the majority of vectors moving towards

the goblet cell cluster and then into the EE cluster. Cells treated with RSP revealed vector

arrows in patterns that resembled both G2D12 and AS treated cells, suggesting it may promote

early differentiation of both secretory and absorptive cell types. Finally, a small proportion of

vector arrows suggested a "backwards" trajectory, with movement from a differentiated

enterocyte into a progenitor population. This behavior may be indicative of de-differentiation or

possibly movement towards cell death, as the cells lose features associated with mature cell

types[53,54]. Overall, these results indicate that all three culture conditions have distinct

differentiation potentials, with AS and RSP giving rise to an increased proportion of EE cells

compared to G2D12.

To verify that the EE cells identified in our enteroid system resemble those found *in situ*, we

compared our dataset with a reference gene set from isolated mouse EE cells[45]. This list

encompasses 77 genes, including conventional markers such as *CHGA*, *NEUROG3* and the

enzyme *TPH1* (Supplementary Fig 4f). An EE cell gene module score was calculated for each

cell in our scRNA dataset based on the number of shared features with the reference EE gene

set[45], which revealed exclusive marking of cells within our EE cell cluster (Fig 4f). In addition,

when looking across culture conditions, RSP and AS showed an increased effect size in relation

to G2D12, with AS having a larger effect size than RSP (Fig 4g). These results underscore

differences observed between the three culture conditions described in previous results and

further demonstrate that EE cells generated using these protocols resemble those found within

native tissue.

To understand the effect of each culture condition on the differentiation of individual

enteroendocrine subsets, we further clustered the EE cells previously identified in our scRNA-

seq dataset. Following sub-clustering, we identified a total of 471 EE cells, which consisted of

enteroendocrine progenitor cells, enterochromaffin cells and various hormone-producing cells

(Supplementary Fig 5a and 5b). We defined enteroendocrine progenitor cells by the expression

of *DLL1 and NEUROG3*, and the absence of expression of differentiated markers *NEUROD1,*

CHGA and *ARX*, the last of which specifically marks non-enterochromaffin EE cells[15,55]

(Supplementary Fig 5c). We found that RSP treatment gave rise to a greater proportion of

enteroendocrine progenitor cells compared to AS (RSP 50% vs AS 13%) (Supplementary Fig

5d). In contrast, AS treatment led to an increased frequency of mature EE cells compared to

RSP (RSP 50% vs AS 87%), with the majority consisting of enterochromaffin cells (RSP 40% vs

AS 62% of all cells) (Supplementary Fig 5d). To investigate the population of peptide hormone-

producing cells, we leveraged our scRNA-seq dataset to determine the expression of hormones

and related genes that have previously been attributed to various EE subsets[2,15,18,45,56]. Notably,

we found expression of both *TPH1*, associated with enterochromaffin cells, and *GHRL*,

associated with X cells, across both AS and RSP. *SST* was not differentially expressed between

AS and RSP, while *MLN*, associated with M cells, was identified predominantly in RSP-treated

cells, although these markers, along with *GAST*, associated with G cells, *GIP*, and *CCK*, were

expressed in only a small number of cells (Supplementary Fig 5e).

Treatment with AS1842856 Followed by Rimonabant and SP600215 Increases GIP

Expression in Human Duodenal Enteroids

Compared to RSP treated enteroids, AS treatment led to more robust EE differentiation, based

on overall induction of multiple EE cell markers and the larger fraction of CHGA+ cells (Figs 2-

4). Direct comparison of the two protocols revealed that AS exposure led to higher expression of

CHGA, NEUROD1, *NEUROG3*, and *SST* (Supplementary Fig 6a), consistent with the higher

percentage of CHGA+ cells (Figs 2 and 3). In contrast, we noted that RSP induced higher

expression of *GIP* compared to AS (Supplementary Fig 6a). Given these results, we next

hypothesized that the combination of Rim, AS, and SP (RASP) would further increase

expression of both CHGA and GIP. Exposure of enteroids to RASP for the full duration of the

differentiation protocol, however, was not compatible with viable enteroids, as evidenced by

their irregular morphology and extremely low RNA content (Supplementary Fig 6b and 6c).

We next tested the impact of adding RSP after exposure to AS, reasoning that AS treatment

would shunt a larger proportion of cells into the EE lineage, as indicated by the scRNA-seq data

(Figs 4d and 4g), with the later addition of RSP inducing *GIP* expression in more cells than RSP

alone. To identify the appropriate time for the addition of RSP, we performed a time-course

analysis of AS treated enteroids and found that multiple transcription factors required for EE

differentiation, including *NEUROD1* and *NEUROG3*, showed increased expression around the

fourth day of differentiation. Following this, both *CHGA* and *SST* had detectable transcript levels

by the sixth day of differentiation (Supplementary Fig 7a). Given these data, we next

hypothesized that AS treatment for six days followed by subsequent exposure to RSP would

lead to increased *GIP* expression compared to RSP alone. To test this, we utilized two

differentiation strategies: (1) switching from AS to RSP at day six of differentiation (AS→RSP) or

(2) adding RSP to AS at day six of differentiation (AS→RASP). Morphologically, AS→RASP

produced smaller enteroids compared to other conditions (Supplementary Fig 7b); however,

RNA concentrations were consistently above the minimum threshold of 10 ng/μL, suggesting

improved viability over exposure to RASP for the full duration of the differentiation protocol

(Supplementary Fig 7c). Enteroids treated with AS→RSP showed gene expression changes

that were similar to RSP alone, aside from *CHGA* and *SST* (Fig 5a). Enteroids treated with

AS→RASP showed 2-3-fold higher gene expression levels of most EE cell markers when

compared to AS, with the exception of *PDX1* and *CCK* (Fig 5a).

Furthermore, exposure to AS for the entire differentiation protocol, i.e., AS and AS→RASP,

decreased expression of *MUC2* and *ALPI* and increased expression of *LGR5* compared to

those that received RSP alone, for any amount of time (Supplementary Fig 8a). In terms of

other lineages, immunofluorescence staining showed that the enterocyte marker cytokeratin 20

(CK20) was expressed in all differentiation conditions, but with very little stain noted in

AS→RASP (Supplementary Fig 8b). MUC2 staining was only noted in RSP, AS and AS→RSP

enteroids while LYZ, similar to CK20, was present in all differentiation conditions except

AS→RASP (Supplementary Fig 8c and 8d).

We also evaluated the effect of our differentiation protocols on proliferation and apoptosis after

7 and 14 days of culture. As expected, at 7days GM supported cell proliferation, as evidenced

by a significantly higher fraction of EdU-positive (EdU+) cells (2.7%) after a two-hour pulse,

compared to all other conditions. There were no significant differences between DM, RSP and

AS, which ranged from 0.1-0.5% EdU+ cells. At day 14 of culture, no differences in the

percentage of EdU+ cells were detected between any groups (which ranged from 0.1-0.6%),

suggesting that GM conditions are not able to sustain ongoing proliferation indefinitely

(Supplementary Fig 8e). Analysis of apoptosis at 7 days showed GM had the lowest percentage

of Annexin V-positive cells (13%) compared to each differentiation condition, with RSP showing

the highest fraction at 29%. At day 14, despite modest differences between groups, levels of

apoptosis were fairly consistent ranging from 19-29% (Supplementary Fig 8f). Further, enteroids

exposed to RSP (including RSP, AS→RSP, and AS→RASP) appear to have more apoptotic

cells compared to DM and AS.

All four differentiation strategies induced a high fraction of CHGA+ enteroids, ranging from 79 to

88% (RSP, 79%; AS, 82%; AS→RSP, 81%; AS→RASP, 88%), as assessed by immunostaining

(Fig 5b and 5c). Quantitative flow cytometric analysis revealed 3.7% of all cells exposed to

AS→RSP to be CHGA+, which was between the 1.0% seen in RSP alone and the 6.1% seen in

AS alone. AS→RASP had the highest percentage of CHGA+ cells at 7.1%, more than 100 times

the number seen with G2D12 alone (Fig 5d). Taken together, these data reveal that exposure of

human enteroids to AS→RASP is the most effective way to induce EE cell differentiation, in

terms of expression of *CHGA*, *PAX4*, *NEUROD1*, and *NEUROG3*, as well as overall number of

CHGA+ cells.

Hormone Production and Secretion Mirror Gene Expression Changes in Duodenal

Enteroids

To assess hormone production and secretion during EE cell differentiation, we next assayed for

duodenal-associated hormones in response to the various differentiation conditions.

Immunofluorescence staining showed that SST was expressed similarly in all differentiation

conditions, aside from G2D12, with 42 to 53% of enteroids containing SST-positive cells (RSP,

45%; AS, 42%; AS→RSP, 53%; AS→RASP, 51%) (Fig 6a and 6b). 5HT was expressed in a

higher percentage of enteroids treated with AS throughout the entire differentiation period than

those that only received RSP (RSP, 44%; AS, 75%; AS→RSP, 59%; AS→RASP, 76%) (Fig 6c

and 6d). An induction in GIP-positive enteroids was detected in response to all differentiation

conditions, with the exceptions of G2D12 and AS→RASP (RSP, 66%; AS, 53%; AS→RSP,

56%; AS→RASP, 8.6%) (Fig 6e and 6f). Further, CCK-positive enteroids were predominantly

detected in response to RSP (29%) and AS→RSP (30%) compared to all other treatment

groups (AS, 4.5%; AS→RASP, 2.6%) (Fig 6g and 6h).

Finally, to assess hormone secretion, we assayed conditioned media from each differentiation

condition. Exposure to AS alone induced higher levels of 5HT secretion than all other

conditions, with AS→RSP conditioned media showing higher levels of 5HT when compared to

RSP and AS→RASP (Fig 7a). These patterns persisted when controlled for total DNA content,

with the exception of a modest difference between AS→RSP and AS→RASP (Supplementary

Fig 9a). Importantly, secretion of 5-HT was significantly increased when differentiated enteroids

were exposed to the adenylate cyclase activator forskolin (Fsk) (Fig 7b and Supplementary Fig

9b). Secretion of GIP was highest following exposure to AS→RSP, compared to RSP alone,

while AS and AS→RASP treated enteroids revealed no secretion of GIP (Fig 7c and

Supplementary Fig 9c). Secretion of GIP from enteroids exposed to AS→RSP was also

significantly increased after exposure to Fsk (Fig 7d and Supplementary Fig 9d). Overall, these

data show that RSP and AS, either alone or in combination, can induce protein expression and

secretion of multiple duodenal hormones (5HT, GIP, SST and CCK) from human enteroids, with

specific differences depending on exposure and/or timing of RSP and AS treatment.

Differentiation Media Induces EE Cell Differentiation in Human Rectoids

To evaluate the ability of our DM to induce EE cell differentiation outside of the duodenum, we exposed organoids derived from rectal ISCs (rectoids) to the G14 and G2D12 protocols (Supplementary Table 1). Rectoids treated with G14 maintained spherical structures, while those exposed to G2D12 had a mixture of spherical and budding structures (Fig 8a). In contrast to enteroids, rectoids exposed to G2D12 had significantly increased expression of the majority of EE markers, included *CHGA*, *PAX4*, *NEUROD1*, *NEUROG3*, *SST*, *GCG* and *PYY*, many of which were at or above the expression level observed in whole rectal mucosa (Fig 8b). Notably, rectoids exposed to G2D12 also had significantly higher expression of *ATOH1* and *MUC2*, decreased expression of *LGR5*, and no change in *CAII* expression, a marker for absorptive cells of the rectum[57,58], when compared to G14 (Supplementary Fig 10).

Immunofluorescence staining revealed CHGA+ cells within a majority of G2D12 rectoids (72%) (Fig 8c and 8d) and quantitative flow cytometric analysis revealed 4.7% of all cells exposed to G2D12 to be CHGA+ (Fig 8e). Additional immunofluorescence staining revealed that both GLP-1 and PYY were expressed in a large number of G2D12 rectoids (GLP-1, 61%; PYY, 40%) (Fig 9a-9d). We next assayed conditioned media from G14 and G2D12 rectoids to evaluate for hormone secretion and found significant levels of both GLP-1 and PYY in G2D12 rectoids compared to G14, which was increased by stimulation with Fsk (Fig 9e-9h, Supplementary Fig 11ab-11d). Overall, these data indicate that G2D12 is sufficient to induce EE cell differentiation of rectal ISCs.

DISCUSSION

Here, we establish that the addition of the small molecules rimonabant, SP600125 and AS1842856 leads to robust differentiation of human EE cells from duodenal ISCs without the

use of transgenic modification. This was assessed by EE marker gene expression, which

approached levels of the native duodenal mucosa, and protein expression, including

immunofluorescent staining, flow cytometric analysis, and hormone secretion assays.

Many of the studies establishing a role for Wnt, Notch, MAPK, and BMP signaling, as well as

short chain fatty acids and isoxazole-9, in EE lineage differentiation and hormone production

have been performed using the organoid culturing system, which approximates *in vivo* growth

and development due to its 3D nature[8,9,16,22-24]. However, there are significant shortcomings to

the analyses of intestinal organoid differentiation that our study has begun to address. First,

RNA expression does not always mirror protein expression, as evidenced by the presence of

GIP mRNA in AS→RASP enteroids (Fig 5a) with little protein expression noted on

immunofluorescence (Fig 6e and 6f) and no secreted protein seen on ELISA (Fig 7c). Further,

our study suggests that enteroids grown in G14 are not appropriate as the sole reference point

in differentiation experiments. For example, G2D12, but not G14, treatment induced expression

of *CHGA* mRNA (Fig 1b), but the comparison of enteroid mRNA expression levels to whole

duodenal mucosa revealed that G2D12 alone induces a very low level of *CHGA* compared to

native tissue. Moreover, immunodetection of CHGA expression revealed significant

heterogeneity between individual enteroids, even when cultured under the same experimental

conditions, as evidenced by only a small minority of enteroids and cells staining CHGA+ in

G2D12 and the variability in hormone expression using immunofluorescence seen in the other

differentiation conditions. These results highlight the limitations of current human EE

differentiation protocols.

Multiple human EE differentiation protocols using only small molecules have suggested that

removal of WNT3a from the base growth medium is critical for secretory lineage

differentiation[9,16]. Our results, in contrast, reveal that removal of WNT3a is detrimental to both

EE differentiation and long-term viability, as noted by a lack of *CHGA* expression at any time

during the differentiation protocol and a marked decrease in total RNA levels with time. This is

consistent with a previous study suggesting that maintaining high WNT3a concentrations during

differentiation was similar, if not better, at inducing *CHGA* expression than a reduction of

WNT3a[23]; however, that study only examined differentiation markers after two days of

differentiation. Additional studies have evaluated differentiation of human intestinal organoids

using reduced amounts of WNT3a with varying lengths of differentiation and found induction of

EE cell markers, when examined[23,24,59,60]. Regardless, our data indicate that high WNT3a

concentrations can promote and maintain differentiation of human intestinal organoids for at

least 12 days and that removal of WNT3a is not necessary for human EE cell differentiation.

We also show that treatment of human enteroids with rimonabant, an inverse agonist of CB1

signaling, in conjunction with a known JNK inhibitor, led to an increase in EE marker expression

and EE cell number, compared with G2D12 alone. We had similar findings when using a well-

described inhibitor of FOXO1, which has been previously implicated in beta cell differentiation[30],

but whose role in EE cell differentiation from primary human ISCs has not been closely

examined. These protocols showed high levels of human EE cell differentiation compared with

prior reports[9,16,23]. While transgenic over-expression of *NEUROG3* generated elevated levels of

human EE cells[14,18,19], our methods avoid the use of genetic techniques while maintaining EE

marker expression at levels at or above endogenous levels within the human duodenal mucosa.

Interestingly, both RSP and AS treatments also led to increased levels of *LGR5* expression,

suggesting some induction of stemness. It has been suggested that EE cells can act as

facultative stem cells in the setting of stress and injury[61], which in our organoid system may

result from chronic culture and/or loss of Matrigel integrity.

Regarding mechanism, it is not known what downstream changes occur after exposure to Rim.

Interestingly, GATA Binding Protein 4 (GATA4) is known to play roles in small intestine

formation and in the expression of CCK and GIP, as *Gata4* deletion *in vitro* and in mice leads to

reduced expression of both GIP and CCK[62-65], and GATA4 has been shown *in vitro* to bind to

the GIP promoter[66]. Similarly, both CCK and GIP have been shown to be regulated by the

endocannabinoid signaling[31-33]. Further, endocannabinoid signaling is known to inhibit

cAMP/protein kinase A signaling[67], which has been shown to play a role in GATA4 activation[68];

therefore, it is possible that reducing CB1 activity could lead to GATA4 activation in human

enteroids, leading to ISC differentiation and hormone production. However, *in vivo* murine

studies show that GATA4 is not expressed in mature EE and goblet cells[69,70], suggesting that

GATA4 may not be involved in mature EE cell function, but could be involved in EE cell

differentiation. In addition, Rim functions as a dual inhibitor of sterol O-acyltransferase 1

(SOAT1 or ACAT1) and SOAT2 (ACAT2)[71], though these play more of a role in the production

of steroid, not peptide, hormones[72].

It is notable that two recent scRNA-seq studies analyzing human and murine EE lineage

differentiation using inducible expression of neurogenin 3 did not identify JNK or FOXO1 as

potential regulatory factors[15,18], suggesting that they may work upstream of neurogenin 3. This

is supported by increased expression of *ATOH1*, a known secretory progenitor marker critical

for neurogenin 3 expression[73,74], in RSP and AS enteroids (Supplementary Figs 2c and 3b). It is

also possible that SP and AS are all having off target effects in our enteroid model. For

example, SP600125 did not alter PDX1 expression in our enteroid system. Therefore, it is

possible that JNK inhibition induces post-translational modifications on PDX1 or, that in human

intestinal cells, JNK inhibition does not affect PDX1. Regardless of the mechanism, the use of

RSP and AS induced differentiation of bona fide EE cells, as evidenced by our scRNA-seq data,

which showed the presence of unique populations of cells that expressed EE markers and that

were derived from populations of stem and progenitor cells.

Exposure of enteroids to combinations of Rim, SP, and AS led to the expression of specific

hormones, with RSP inducing GIP expression and AS inducing higher levels of 5HT expression.

Exposure to AS→RSP yielded similar levels of *GIP* mRNA and total GIP-positive enteroids, but

increased levels of GIP secretion, when compared to RSP alone; however, AS→RASP

seemingly inhibited GIP protein production and secretion. AS alone appears to be the most

potent inducer of 5HT secretion. Exposure to RSP, with or without AS, induced less 5HT

secretion when compared to AS alone. Taken together, these combinatorial data suggest that

AS exposure is a potent inducer of 5HT secretion while inhibiting GIP secretion, and exposure

to Rim and SP is a potent inducer of GIP secretion while leading to reduced 5HT secretion

when compared to AS treated enteroids. These data suggest that modulation of the

endocannabinoid signaling system, JNK, and FOXO1, as well as other potential targets, could

have multiple effects on EE function, controlling mRNA and protein production, as well as

secretion, of multiple hormones.

In comparison to enteroids, rectoids showed a much stronger induction of EE cells when

exposed to G2D12. It is clear that human small intestinal and colonic crypts demonstrate distinct

gene expression profiles, which includes genes involved in epithelial cell differentiation[75], so it is

likely that the requirements for EE cell lineage differentiation will be different between the

various regions. Additional experiments examining whether the mechanisms defined for

duodenal EE differentiation also regulate rectal EE differentiation need to be explored.

In summary, we have shown robust differentiation of human EE cells from duodenal and rectal

ISCs using Wnt3a-containing differentiation media, and, with respect to enteroids, the addition

of the small molecules rimonabant, SP600125, and AS1842856. These protocols improve upon

current methods of EE cell differentiation without the use of direct genetic alteration. These

studies also provide a platform for future experiments designed to identify endogenous factors

regulating EE differentiation, identifying the role and response of EE cells in human GI disease,

and further increasing EE cell numbers to potentially be used as personalized cell therapy.

METHODS

Isolation of human intestinal crypts

Tissues were procured as previously described[76]. In short, de-identified endoscopic and rectal

biopsies were collected from grossly unaffected tissues in adolescent/young adult patients

undergoing esophagogastroduodenoscopy and colonoscopy at Boston Children's Hospital for

gastrointestinal complaints and used for organoid derivation. Further, de-identified duodenal

resections were collected from adult patients undergoing pancreaticoduodenectomy at

Massachusetts General Hospital prior to chemotherapy or radiation therapy for pancreatic

carcinoma. Whole mucosal biopsies were generated from the duodenal resections using

endoscopic biopsy forceps and used for total RNA isolation. Age and sex of donors can be

found in Supplementary Table 2 but were unknown to researchers when experiments were

being performed. Only macroscopically normal-appearing tissue was used from patients without

a known gastrointestinal diagnosis. Each experiment was performed on at least three

independent organoid lines derived from adolescent/young adult biopsy samples. Informed

consent and developmentally appropriate assent were obtained at Boston Children's Hospital

from the donors' guardian and the donor, respectively. Informed consent was obtained at

Massachusetts General Hospital from the donors. All methods were approved and carried out in

accordance with the Institutional Review Boards of Boston Children's Hospital (Protocol number

IRB-P00000529) and Massachusetts General Hospital (Protocol number IRB-2003P001289).

Resections were briefly washed with pre-warmed DMEM/F12, after which the epithelial layer

was separated from the rest of the duodenum manually with sterilized surgical tools then taken

for RNA isolation. To isolate crypts, adolescent/young adult biopsies were digested in 2 mg/mL

of Collagenase Type I reconstituted in Hank's Balanced Salt Solution for 40 minutes at 37°C.

Samples were then agitated by pipetting followed by centrifugation at 500 x g for 5 minutes at

4°C. The supernatant was then removed, and crypts resuspended in 200-300 μL of Matrigel,

with 50μL being plated onto 4-6 wells of a 24-well plate and polymerized at 37°C.

Culturing of human duodenal and rectal organoids *in vitro*

Isolated crypts in Matrigel were grown in specific growth media (GM) (Supplementary Table 1)

based on the tissue of origin. The resulting organoids were passaged every 6-8 days as

needed, with media changes occurring every two days. To passage, Matrigel was mechanically

dissociated from the well and resuspended in 500 μL of Cell Recovery solution for 40-60

minutes at 4°C. To aid in separating the Matrigel and enteroids, the tubes are gently inverted

and then centrifuged at 500 x g for 5 minutes at 4°C. The supernatant was then removed, and

organoids resuspended in Matrigel, followed by mechanical disruption via a bent-tipped pipette.

Organoids were passaged at a 1:2 dilution, with 50 μL per well of a 24-well plate. After plating,

the organoids were incubated at 37°C for 10 minutes to allow the Matrigel to set. Once

complete, 500 μL of GM was added to each well.

For differentiation, organoids were passaged and grown in GM for two days, to allow for ISC

expansion, after which the organoids were transitioned to tissue-specific differentiation media

(DM) with additional small molecules added as described (Supplementary Table 1). Media was

changed every two days, with Tubastatin A being removed after the second day of

differentiation. Enteroid and rectoids were taken for analysis after 14 days.

Gene expression analysis

Total RNA was purified from individual wells using TRI®Reagent and the Direct-zol™ RNA kit,

following the manufacturer's protocol. RNA concentration was determined using a NanoDrop™

1000 spectrophotometer (Life Technologies). RNA was then reverse transcribed using the High-

Capacity cDNA Reverse Transcription Kit. Gene expression analysis was then performed by

Real Time quantitative PCR (qPCR) using a QuantStudio 6 Flex thermocycler (Thermo Fisher).

We used the following Taqman primers from Thermo Fisher (Supplementary Table 3). *18S*

transcripts were used as the internal control and data were expressed using the 2^{-ddCt} method

with Ct limit of 40. Fold change, unless otherwise stated, was compared to total RNA derived

from whole mucosal tissue biopsies, using duodenal mucosa with enteroids and rectal mucosa

with rectoids.

DNA isolation

To isolate organoid genomic DNA, 200 µL of 50 mM NaOH was added to a single well of a 24-

well plate and the Matrigel was mechanically dissociated. The samples were then transferred to

1.5 mL microcentrifuge tubes and placed in a 95°C heat block for 20 minutes. The tubes were

then vortexed, after which 25 µL of 1M Tris-HCl was added. The samples were then centrifuged

at 18,500 x g for 10 minutes. The DNA content of the supernatant was then assayed using a

NanoDrop™ 1000 spectrophotometer (Thermo Fisher).

Immunofluorescence

Immunofluorescence staining was performed as previously described with minor

modifications[77]. Organoids were grown in and isolated from Matrigel as noted above. 1-3 wells

from a 24-well plate were washed in 200 µL of 1X phosphate-buffered saline (PBS) and moved

in suspension to a 1.5 mL microcentrifuge tube. Each tube was centrifuged at 800 x g for 5

minutes at 4°C to pellet organoids. PBS was aspirated, and organoids were fixed in 200 µL of

4% paraformaldehyde (PFA) for 20 minutes on ice, shaking. Each tube was centrifuged again

as above, and PFA was aspirated. The organoids were then resuspended in 500 µL of 0.3%

Triton-X in PBS and moved to a 48-well plate for the remaining steps. The organoids were

permeabilized for 30 minutes at room temperature, shaking. Between each step, organoids

were allowed to settle to the bottom of each well, the plate was angled, and the solution

aspirated by careful pipetting. Organoids were then blocked with 5% bovine serum albumin

(BSA) in PBS for 1.5 hours at room temperature, shaking. This was followed by three five-

minute washes in 500 µL of PBS at room temperature, shaking.

Each well was then incubated in 200 µL of primary antibodies diluted in 5% BSA/0.1% Trition-X

in PBS at 4°C overnight. This was followed by five washes in 500 µL of 0.1% Triton-X in PBS for

15 minutes each at room temperature, shaking. 200 µL of secondary antibodies diluted in 0.1%

Triton-X in PBS were then added to each well and incubated for two hours at room temperature,

shaking. Organoids were then washed as above, then moved to new 1.5 mL centrifuge tubes.

During the last wash, 4′,6‑diamidino-2-phenylindole (DAPI) was added at a concentration of

1:1000 for nuclear staining. Organoids were then centrifuged at 1000 x g for 5 minutes at 4°C to

help remove as much PBS as possible. Slides were prepared by drawing three circles with a

hydrophobic pen. Enteroids were then resuspended in 20 µL of Prolong Gold Antifade mountant

and droplets placed within hydrophobic circles. The organoids were spread out to reduce

clumping, sealed with a coverslip, and allowed to dry overnight at room temperature. Slides

were stored at 4°C for future imaging. Images were acquired using a Nikon upright Eclipse 90i

microscope with a 20×/0.75 Plan-Apochromat objective and adjusted for brightness and

contrast in Fiji (v2.10/1.53c)[78].

Quantification of organoid immunofluorescence

This technique was adapted from a previously described method[79]. Immunofluorescent images

were acquired using an Invitrogen EVOS FL 2 Auto microscope (Thermo Fisher).

Representative images of stained organoids were taken at 2X magnification. The stitched

images were then processed in Fiji[78]. The color of the DAPI images was converted to 8-bit

grayscale and then the image was smoothed by applying a Gaussian Blur filter (radius = 4,

scaled units). Thresholding of the smoothed images was performed using manual adjustment to

achieve optimal separation of individual organoids. Watershed and Find Edges filters were then

applied to segment any clumped organoids. Post-segmentation analysis was performed to

outline and count individual organoids using Analyze Particles (size=4,000-100,000,

circularity=0.30-1.00, exclude on edges). Each image was then manually curated to exclude

debris and organoids exhibiting background fluorescence. Any remaining clumped organoids

were manually separated prior to quantification. The outlines generated from the DAPI images

were then applied to the corresponding images from the other fluorescent channels. Each color

image was converted to 8-bit grayscale and then the HiLo Lookup Table was applied. The

threshold gate for stained cells was found by manual adjustment of positively stained organoids

to achieve optimal representation. The threshold gate for each channel was then applied to

each experimental condition. The Mean metric was extracted with ROI manager (measure) and

compiled for analysis. Organoids with a Mean value of more than zero were considered positive.

Flow cytometry

Organoids were incubated in Cell Recovery solution for 40-60 minutes at 4°C to remove the

Matrigel and then centrifuged at 500 x g for 5 minutes at 4°C. To achieve single cell suspension,

organoids were then incubated in 500 µL of TrypLE Express at 37°C for 30 minutes and broken

up by repeated pipetting using a bent P1000 pipette tip. Each sample was then diluted in 800µL

of 20% fetal bovine serum (FBS) in Advanced DMEM/F12 and then centrifuged at 800 x g for 5

minutes at 4°C. To mark dead cells, each sample was then incubated in DAPI (1:1000) diluted

in 2% FBS/2 mM EDTA/calcium-free DMEM for 20 minutes at room temperature, then

centrifuged at 800 x g for 5 minutes at 4°C and washed in 2% FBS/2 mM EDTA/DMEM. Cells

were then incubated in 1% PFA for 15 minutes at room temperature, washed with 2% FBS/PBS

and then permeabilized in 0.2% Tween 20 in 2% FBS/PBS for 15 minutes at 37°C. Following

centrifugation, cells were resuspended in 0.1% Tween 20/2% FBS/2 mM EDTA in PBS with

PE/Alexa Fluor 647-conjugated CHGA, PE-conjugated mouse IgG1, K isotype, Alexa Fluor 647-

conjugated mouse IgG2b, K isotype, or with no antibody (the latter three acting as controls) for

30 minutes on ice. Cells were then washed in 0.1% Tween 20/2% FBS/2 mM EDTA in PBS,

filtered through a 37-micron mesh, and then analyzed on a BD LSRFortessa flow cytometer

using BD FACSDiva (v8.02) and FlowJo (v10.6.2). For gating strategy, please see

Supplementary Figure 12.

Cell proliferation

Enteroid cell proliferation was assessed using the Click-IT EdU Alexa Fluor 488 Flow Cytometry

Assay kit following the manufacturer's protocols. Briefly, enteroids were grown in Matrigel as

noted above and taken for analysis after 7 and 14 days. Enteroids were labeled following

incubation with 10 µM of EdU at 37°C for 2 hours and then incubated with TrypLE Express as

above to isolate single cells. All samples were then diluted in 800 µL of 20% FBS in Advanced

DMEM/F12 and then centrifuged at 800 x g for 5 minutes at 4°C. Fixation, permeabilization, and

the Click-iT reaction were performed in accordance with manufacturer's protocols before being

filtered through a 37-micron mesh, and then analyzed on a BD LSRFortessa flow cytometer

using BD FACSDiva and FlowJo.

Cell apoptosis

Apoptosis was assessed using the Dead Cell Apoptosis Kit with Annexin V Alexa Fluor 488 &

Propidium Iodide following the manufacturer's protocols. For these experiments, enteroids were

grown in Matrigel as noted above and taken for analysis after 7 and 14 days. To isolate single

cells, enteroids were incubated with TrypLE Express as above. All samples were then diluted in

800 µL of 20% FBS in Advanced DMEM/F12 and then centrifuged at 800 x g for 5 minutes at

4°C. Samples were then incubated with propidium iodide and annexin V per manufacturer's

protocol. Cells were filtered through a 37-micron mesh, and then analyzed on a BD

LSRFortessa flow cytometer using BD FACSDiva and FlowJo.

ELISA

Hormone quantification for 5HT, GIP, GLP-1, and PYY were performed using kits and following

the manufacturer's protocols. For these experiments, organoids were grown in 48-well plates to

aid in concentrating the hormone of interest. Conditioned media was taken on day 14 of

differentiation, 48 hours after last media change. Diprotin A (100µM), a dipeptidyl peptidase 4

inhibitor, was added daily to the media for the last two days of differentiation to prevent GIP and

GLP-1 degradation. For hormone induction studies, conditioned media was taken on day 13, 24

hours after last media change, after which forskolin (10 µM) was added to fresh media and

collected after 24 hours. As an additional control, differentiation media not exposed to cells was

also evaluated. This value was then subtracted from each experimental sample.

Sample preparation for single cell RNA sequencing

Enteroids were incubated in Cell Recovery solution for 40-60 minutes at 4°C to remove the

Matrigel and then centrifuged at 500 x g for 5 minutes at 4°C. To achieve single cell suspension,

organoids were then incubated in 500 µL of TrypLE Express at 37°C for 30 minutes and broken

up by repeated pipetting using a bent P1000 pipette tip. Each sample was then diluted in 800 µL

of 20% FBS in Advanced DMEM/F12, filtered through 37 µm mesh, and then centrifuged at 300

x g for 10 minutes at 4°C. Dead cells were then removed using MACS separation following the

manufacturer's protocol. Briefly, each sample was resuspended in 100 µL Dead Cell Removal

MicroBeads and incubated at room temperature for 15 minutes. Four hundred microliters of 1x

Binding Buffer were then added to each sample, after which the samples were applied to

prepared MACS columns placed in a MACS Separator. Flow-through containing live cells was

collected. This was followed by washing the columns four times with 1x Binding Buffer, which

were combined with the initial flow-through. These samples were then centrifuged at 800 x g for

5 minutes at 4°C.

Hashtag-labeling of sequencing samples

Single cells isolated after dead cell removal were labeled prior to sequencing with Biolegend

Totalseq-B hashtag antibodies, targeted against CD298 and β2-microglobulin, per the

manufacturer's protocol[43]. Briefly, each of the nine samples (three separate differentiation

conditions in three enteroid lines) were resuspended in 50 µL of Cell Staining Buffer and

blocked with 5 µL of Human TruStain FcX^{TM} blocking reagent for 10 minutes at 4°C. The

supernatant was then removed, and each sample resuspended in 50 µL of Cell Staining Buffer

containing 1 µg of a unique Totalseq-B hashtag antibody. Samples were incubated for 30

minutes at 4°C and then washed three times in 3.5 mL of Cell Staining Buffer. The samples

were then filtered through a 37 µm mesh, resuspended in 2% BSA in PBS at a concentration of

1500 cells/µL, and then pooled together. Of note, prior to the scRNA-seq experiment, antibody

binding was verified using fluorophore-conjugated antibodies with the same binding specificity.

Droplet-based single cell RNA sequencing and alignment

A total of 180,000 cells as one pool were input across four channels, with a recovery rate of

approximately 30,000 cells across all four channels. RNA libraries were prepared from single

cells by the Single Cell Core at Harvard Medical School, Boston, MA, using the 10X Chromium

Single Cell 3' Library Chip (v3.1) with dual indexed barcodes. Briefly, single cells were

partitioned into Gel Beads in Emulsion (GEMs) by the Chromium Controller. Prior to partitioning,

hashed cells were pooled together and "super-loaded" to reduce batch effects across

samples[43]. Once partitioned, cells were lysed and RNA was captured, reverse transcribed, and

amplified. cDNA was then enzymatically fragmented and sample indices were attached to both

ends of the fragment. Following library prep, the transcriptomes were sequenced by the

Molecular Biology Core Facilities at the Dana-Farber Cancer Institute on an Illumina NovaSeq

6000 sequencing platform. The pooled sample was split across two lanes on a NovaSeq S1

flow cell.

After sequencing, the files were processed, demultiplexed and aligned using 10X's Cell Ranger

software (v5.0.1). Briefly, raw BCL files were demultiplexed using the cellranger mkfastq

function, a wrapper of Illumina's bcl2fastq, with the filter dual index argument set to true.

Following the generation of the FASTQ files, the samples were aligned to the hg19 human

genome and features were counted using the cellranger count function. This function aligned

the UMIs of each individual cell to the related gene and counted the number of detected UMIs to

generate a feature-barcode matrix. In addition, it also measured the hashtag oligo counts for

each individual cell from the relevant hashtag antibodies that were used to label each condition.

Further, for analysis of RNA velocity, both un-spliced and spliced RNA variants were aligned

and counted using the velocyto.py package (v0.17.16)[52].

Single cell RNA sequencing analysis

Analysis of the scRNA-seq data was conducted using the Seurat package (v4.0.1), within the R

programming language (v4.0.3). Initially, cells were filtered to only include those with at least

one detectable hashtag antibody signal, and then hashtag counts were normalized, and the cell

samples were demultiplexed using the HTOdemux function built into Seurat. This function

ranked a cell as being positive for a specific hashtag signal if the detected signal was in the 99th

percentile or higher. Cells were then demultiplexed, with cells that had no positive signals being

labeled as negative cells, cells with only one positive signal being labeled by their respective

positive signal, and cells with two or more positive signals being labeled as doublets. Following

demultiplexing, the dataset was pre-processed to remove cells that were deemed of lower

quality, based on their number of unique genes (<200 unique genes) and the proportion of UMIs

that corresponded to mitochondrial RNA (>25% of UMIs)[44].

After the removal of doublet barcodes, negative barcodes and the remaining barcodes that were

deemed to be of lower quality, we were left with a combined dataset of 14,767 cells that was

used for further analysis. To better control for batch effects and to improve our ability to detect

diversity within our samples, we normalized the gene counts using regularized negative

binomial regression, using SCTransform (v0.3.2)[80]. Following normalization, we performed

dimensional reduction by running Principal Component Analysis (PCA) and, based on the elbow

plot generated, selected the first 20 principal components for defining the K nearest neighbors

and the UMAP plot. After dimensional reduction, 15 clusters were defined using Louvain

clustering (resolution=0.5). Differentially expressed genes were identified across each cluster

using the Wilcoxon rank sum test via the FindAllMarkers function built into Seurat. Only genes

that were expressed in at least 25% of cells and had a log-fold change of 0.25 were considered

in this analysis. These differential genes, as well as classical epithelial markers, were used to

broadly categorize the clusters into six epithelial cell types and include *LGR5, AXIN2* and

ASCL2 for intestinal stem cells, *KRT20*, *FABP2*, and *FABP1* for enterocytes, *MUC2, FCGBP,*

and *GFI1* for goblet cells, *CHGA*, *NEUROG3*, *NEUROD1* and *PAX4* for enteroendocrine cells,

CD44 for progenitor cells and *MKI67* for proliferating cells[47-49].

EE cells were then subsetted out using Seurat's subset function and further analyzed. Similar to

the parent dataset, gene counts were normalized using SCTransform and then dimensional

reduction and cluster assignment was calculated using the Seurat pipeline. Across 471 cells, a

total of 7 clusters were identified using Louvain clustering at a resolution of 0.8. Cluster-defining

features were calculated using a Wilcoxon rank sum test through the FindAllMarkers function,

with significant genes being defined as having been expressed in at least 5% of cells, with a log-

fold change of 0.25 minimum. These genes, alongside markers previously used to annotate EE

cell subsets, were used to assign identity to each cluster for downstream analysis. Using these

markers, we broadly categorized the clusters into enteroendocrine progenitor cells (*DLL1,*

NEUROG3), enterochromaffin cells (*CHGA, TPH1, NEUROD1*), and various peptide hormone

producing subsets including M/X cells (*ARX, MLN, GHRL*) and G/K/I cells (*ARX, GAST, GIP,*

CCK)[2,15,18,45,55,56]. Single cell gene expression of marker and hormone genes were visualized

using Seurat's FeaturePlot function[81].

Finally, enteroid EE cell identity was compared to the gene set signature from tissue EE cells

isolated from the murine small intestine[45]. To compare the gene signature, we calculated the

gene module score of each individual cell using the AddModuleScore function found in Seurat

v3 and v4, which calculates the score based on the enrichment of the specified gene set

compared to randomly selected genes with a comparable average gene expression[44]. The

AddModuleScore function calculates a scaled score between 0 to 1.

Trajectory analysis with scVelo

RNA velocity was calculated using the scVelo package (v0.2.3) with Scanpy (v1.7.1) on Python

(v3.7.10)[51]. To perform trajectory analysis, the un-spliced and spliced variant count matrix that

was previously calculated using velocyto (v0.17.15) was fused with an anndata object

containing the UMAP information and cluster identities defined in Seurat analysis. The

combined dataset was then processed using the scVelo pipeline: The ratio of un-spliced:spliced

RNA for each gene was normalized and filtered using the default settings. Afterwards, the first

and second moments were calculated for velocity estimation. Following moment calculation, the

dynamic model was used to calculate the RNA velocities. The dynamic model iteratively

estimates the parameters that best model the phase trajectory of each gene, therefore capturing

the most accurate, albeit more computationally intensive, estimate of the dynamics for each

gene. These approaches were used to graphically model the RNA velocity for each culture

condition.

Quantification and statistical analysis

All experiments were repeated using at least three different human organoid lines and

representative data from a single line is shown. For qPCR and flow cytometry studies, each

condition was performed using pooled enteroids from 3-5 wells, unless otherwise noted, with

each well acting as a single sample. Whole mucosal biopsies from duodenal resections and

whole mucosa rectal biopsies were combined from three different individuals to generate a

single reference sample for all enteroid and rectoid experiments.

Prior to statistical analysis, all qPCR data were transformed using $\log_{10}$. When analyzing only

two conditions, statistical significance was determined by unpaired, two-tailed Student's t test,

combined with the Holm-Sidak method to control for multiple comparisons. When analyzing

more than two conditions, statistical significance was determined by either one-way or two-way

ANOVA, followed by Tukey post hoc analysis. Specific conditions were excluded from statistical

analysis if the data from one or more samples was labeled as not detectable. To assess the

proportion of cell populations within our scRNAseq data, we used tidyverse (1.3.1), PNWColors

(0.1.0), and cowplot (1.1.1) to quantify and visualize our data. When analyzing the statistical

relationship between cell proportions identified using single cell RNA sequencing, the non-

parametric Kruskal-Wallis test was conducted, followed by Dunn's post-hoc analysis. We used

the Bonferroni correction to account for multiple hypothesis testing when performing the post-

hoc analysis. When comparing the distribution of gene module scores across the three different

culture conditions, effect size was used to compare large changes and was calculated as

Cohen's d. Statistical details for each experiment can be found within each figure legend.

Reagents

A list of all resources used in this study, including source and catalog number, can be found in

Supplementary Table 4.

Data Availability

The scRNA-seq data generated in this study have been deposited in the Gene Expression

Omnibus database under accession code GSE178342

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE178342). All other data generated in

this study are provided within the article and its Supplementary Information/Source Data file.

REFERENCES

1   Posovszky, C. Development and Anatomy of the Enteroendocrine System in Humans.
Endocr Dev **32**, 20-37, doi:10.1159/000475729 (2017).

2   Furness, J. B., Rivera, L. R., Cho, H. J., Bravo, D. M. & Callaghan, B. The gut as a
sensory organ. *Nat Rev Gastroenterol Hepatol* **10**, 729-740,
doi:10.1038/nrgastro.2013.180 (2013).

3   El-Salhy, M., Hausken, T., Gilja, O. H. & Hatlebakk, J. G. The possible role of
gastrointestinal endocrine cells in the pathophysiology of irritable bowel syndrome.
Expert Rev Gastroenterol Hepatol **11**, 139-148, doi:10.1080/17474124.2017.1269601
(2017).

4   Worthington, J. J., Reimann, F. & Gribble, F. M. Enteroendocrine cells-sensory sentinels
of the intestinal environment and orchestrators of mucosal immunity. *Mucosal Immunol*
11, 3-20, doi:10.1038/mi.2017.73 (2018).

5 Holst, J. J., Vilsboll, T. & Deacon, C. F. The incretin system and its role in type 2
diabetes mellitus. *Mol Cell Endocrinol* **297**, 127-136, doi:10.1016/j.mce.2008.08.012
(2009).

6 Yousefi, M., Li, L. & Lengner, C. J. Hierarchy and Plasticity in the Intestinal Stem Cell
Compartment. *Trends Cell Biol* **27**, 753-764, doi:10.1016/j.tcb.2017.06.006 (2017).

7 Yu, H. *et al.* The Contributions of Human Mini-Intestines to the Study of Intestinal
Physiology and Pathophysiology. *Annu Rev Physiol* **79**, 291-312, doi:10.1146/annurev-
physiol-021115-105211 (2017).

8 Fujii, M. *et al.* Human Intestinal Organoids Maintain Self-Renewal Capacity and Cellular
Diversity in Niche-Inspired Culture Condition. *Cell Stem Cell* **23**, 787-793 e786,
doi:10.1016/j.stem.2018.11.016 (2018).

9 Sato, T. *et al.* Long-term expansion of epithelial organoids from human colon, adenoma,
adenocarcinoma, and Barrett's epithelium. *Gastroenterology* **141**, 1762-1772,
doi:10.1053/j.gastro.2011.07.050 (2011).

10 Basak, O. *et al.* Induced Quiescence of Lgr5+ Stem Cells in Intestinal Organoids
Enables Differentiation of Hormone-Producing Enteroendocrine Cells. *Cell Stem Cell* **20**,
177-190 e174, doi:10.1016/j.stem.2016.11.001 (2017).

11 Sato, T. & Clevers, H. SnapShot: Growing Organoids from Stem Cells. *Cell* **161**, 1700-
1700 e1701, doi:10.1016/j.cell.2015.06.028 (2015).

12 Yin, X. *et al.* Niche-independent high-purity cultures of Lgr5+ intestinal stem cells and
their progeny. *Nat Methods* **11**, 106-112, doi:10.1038/nmeth.2737 (2014).

13 Almeqdadi, M., Mana, M. D., Roper, J. & Yilmaz, O. H. Gut organoids: mini-tissues in
culture to study intestinal physiology and disease. *Am J Physiol Cell Physiol* **317**, C405-
C419, doi:10.1152/ajpcell.00300.2017 (2019).

14 Sinagoga, K. L. *et al.* Deriving functional human enteroendocrine cells from pluripotent
stem cells. *Development* **145**, doi:10.1242/dev.165795 (2018).

15 Gehart, H. *et al.* Identification of Enteroendocrine Regulators by Real-Time Single-Cell
Differentiation Mapping. *Cell* **176**, 1158-1173 e1116, doi:10.1016/j.cell.2018.12.029
(2019).

16 Beumer, J. *et al.* Enteroendocrine cells switch hormone expression along the crypt-to-
villus BMP signalling gradient. *Nat Cell Biol* **20**, 909-916, doi:10.1038/s41556-018-0143-
y (2018).

17 Chen, C., Fang, R., Davis, C., Maravelias, C. & Sibley, E. Pdx1 inactivation restricted to
the intestinal epithelium in mice alters duodenal gene expression in enterocytes and
enteroendocrine cells. *Am J Physiol Gastrointest Liver Physiol* **297**, G1126-1137,
doi:10.1152/ajpgi.90586.2008 (2009).

18   Beumer, J. *et al.* High-Resolution mRNA and Secretome Atlas of Human
Enteroendocrine Cells. *Cell* **181**, 1291-1306 e1219, doi:10.1016/j.cell.2020.04.036
(2020).

19   Chang-Graham, A. L. *et al.* Human Intestinal Enteroids With Inducible Neurogenin-3
Expression as a Novel Model of Gut Hormone Secretion. *Cell Mol Gastroenterol Hepatol*
8, 209-229, doi:10.1016/j.jcmgh.2019.04.010 (2019).

20   Munera, J. O. *et al.* Differentiation of Human Pluripotent Stem Cells into Colonic
Organoids via Transient Activation of BMP Signaling. *Cell Stem Cell* **21**, 51-64 e56,
doi:10.1016/j.stem.2017.05.020 (2017).

21   McCracken, K. W. *et al.* Modelling human development and disease in pluripotent stem-
cell-derived gastric organoids. *Nature* **516**, 400-404, doi:10.1038/nature13863 (2014).

22   Petersen, N. *et al.* Generation of L cells in mouse and human small intestine organoids.
Diabetes **63**, 410-420, doi:10.2337/db13-0991 (2014).

23   VanDussen, K. L. *et al.* Development of an enhanced human gastrointestinal epithelial
culture system to facilitate patient-based assays. *Gut* **64**, 911-920, doi:10.1136/gutjnl-
2013-306651 (2015).

24   Tsakmaki, A., Fonseca Pedro, P., Pavlidis, P., Hayee, B. & Bewick, G. A. ISX-9
manipulates endocrine progenitor fate revealing conserved intestinal lineages in mouse
and human organoids. *Mol Metab* **34**, 157-173, doi:10.1016/j.molmet.2020.01.012
(2020).

25   Kousteni, S. FoxO1, the transcriptional chief of staff of energy metabolism. *Bone* **50**,
437-443, doi:10.1016/j.bone.2011.06.034 (2012).

26   Tothova, Z. & Gilliland, D. G. FoxO transcription factors and stem cell homeostasis:
insights from the hematopoietic system. *Cell Stem Cell* **1**, 140-152,
doi:10.1016/j.stem.2007.07.017 (2007).

27   Roy, S. A. *et al.* Dual regulatory role for phosphatase and tensin homolog in specification
of intestinal endocrine cell subtypes. *World J Gastroenterol* **18**, 1579-1589,
doi:10.3748/wjg.v18.i14.1579 (2012).

28   Garcia-Martinez, J. M., Chocarro-Calvo, A., De la Vieja, A. & Garcia-Jimenez, C. Insulin
drives glucose-dependent insulinotropic peptide expression via glucose-dependent
regulation of FoxO1 and LEF1/beta-catenin. *Biochim Biophys Acta* **1839**, 1141-1150,
doi:10.1016/j.bbagrm.2014.07.020 (2014).

29   Arriola, D. J., Mayo, S. L., Skarra, D. V., Benson, C. A. & Thackray, V. G. FOXO1
transcription factor inhibits luteinizing hormone beta gene expression in pituitary
gonadotrope cells. *J Biol Chem* **287**, 33424-33435, doi:10.1074/jbc.M112.362103
(2012).

30   Bouchi, R. *et al.* FOXO1 inhibition yields functional insulin-producing cells in human gut
organoid cultures. *Nat Commun* **5**, 4242, doi:10.1038/ncomms5242 (2014).

31   DiPatrizio, N. V. Endocannabinoids in the Gut. *Cannabis Cannabinoid Res* **1**, 67-77,
doi:10.1089/can.2016.0001 (2016).

32   Argueta, D. A., Perez, P. A., Makriyannis, A. & DiPatrizio, N. V. Cannabinoid CB1
Receptors Inhibit Gut-Brain Satiation Signaling in Diet-Induced Obesity. *Front Physiol*
10, 704, doi:10.3389/fphys.2019.00704 (2019).

33   Sathyapalan, T. *et al.* Effect of rimonabant and metformin on glucose-dependent
insulinotropic polypeptide and glucagon-like peptide-1 in obese women with polycystic
ovary   syndrome.   *Clin   Endocrinol   (Oxf)*   **72**,   423-425,   doi:10.1111/j.1365-
2265.2009.03643.x (2010).

34   Biteau, B., Hochmuth, C. E. & Jasper, H. JNK activity in somatic stem cells causes loss
of tissue homeostasis in the aging Drosophila gut. *Cell Stem Cell* **3**, 442-455,
doi:10.1016/j.stem.2008.07.024 (2008).

35   Sancho, R. *et al.* JNK signalling modulates intestinal homeostasis and tumourigenesis in
mice. *EMBO J* **28**, 1843-1854, doi:10.1038/emboj.2009.153 (2009).

36   Kaneto, H. *et al.* Oxidative stress and the JNK pathway in diabetes. *Curr Diabetes Rev*
1, 65-72 (2005).

37   Tang, C. *et al.* Glucose-Induced beta-Cell Dysfunction In Vivo: Evidence for a Causal
Role of C-jun N-terminal Kinase Pathway. *Endocrinology* **159**, 3643-3654,
doi:10.1210/en.2018-00566 (2018).

38   Mandic, A. D. *et al.* c-Jun N-terminal kinase 2 promotes enterocyte survival and goblet
cell differentiation in the inflamed intestine. *Mucosal Immunol* **10**, 1211-1223,
doi:10.1038/mi.2016.125 (2017).

39   Pagliuca, F. W. *et al.* Generation of functional human pancreatic beta cells in vitro. *Cell*
159, 428-439, doi:10.1016/j.cell.2014.09.040 (2014).

40   Harskamp, L. R., Gansevoort, R. T., van Goor, H. & Meijer, E. The epidermal growth
factor receptor pathway in chronic kidney diseases. *Nat Rev Nephrol* **12**, 496-506,
doi:10.1038/nrneph.2016.91 (2016).

41   Zhao, X., Mohan, R., Ozcan, S. & Tang, X. MicroRNA-30d induces insulin transcription
factor MafA and insulin production by targeting mitogen-activated protein 4 kinase 4
(MAP4K4)   in   pancreatic   beta-cells.   *J   Biol   Chem*   **287**,   31155-31164,
doi:10.1074/jbc.M112.362632 (2012).

42   Yu, F. *et al.* FoxO1 inhibition promotes differentiation of human embryonic stem cells
into insulin producing cells. *Exp Cell Res* **362**, 227-234, doi:10.1016/j.yexcr.2017.11.022
(2018).

43   Stoeckius, M. *et al.* Cell Hashing with barcoded antibodies enables multiplexing and
doublet detection for single cell genomics. *Genome Biol* **19**, 224, doi:10.1186/s13059-
018-1603-1 (2018).

44 Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902
e1821, doi:10.1016/j.cell.2019.05.031 (2019).

45 Haber, A. L. *et al.* A single-cell survey of the small intestinal epithelium. *Nature* **551**, 333-
339, doi:10.1038/nature24489 (2017).

46 Mead, B. E. *et al.* Harnessing single-cell genomics to improve the physiological fidelity of
organoid-derived cell types. *BMC Biol* **16**, 62, doi:10.1186/s12915-018-0527-2 (2018).

47 Barker, N. Adult intestinal stem cells: critical drivers of epithelial homeostasis and
regeneration. *Nat Rev Mol Cell Biol* **15**, 19-33, doi:10.1038/nrm3721 (2014).

48 van der Flier, L. G. & Clevers, H. Stem cells, self-renewal, and differentiation in the
intestinal epithelium. *Annu Rev Physiol* **71**, 241-260,
doi:10.1146/annurev.physiol.010908.163145 (2009).

49 Noah, T. K., Donahue, B. & Shroyer, N. F. Intestinal development and differentiation.
Exp Cell Res **317**, 2702-2710, doi:10.1016/j.yexcr.2011.09.006 (2011).

50 Basak, O. *et al.* Mapping early fate determination in Lgr5+ crypt stem cells using a novel
Ki67-RFP allele. *EMBO J* **33**, 2057-2068, doi:10.15252/embj.201488017 (2014).

51 Bergen, V., Lange, M., Peidli, S., Wolf, F. A. & Theis, F. J. Generalizing RNA velocity to
transient cell states through dynamical modeling. *Nat Biotechnol* **38**, 1408-1414,
doi:10.1038/s41587-020-0591-3 (2020).

52 La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494-498,
doi:10.1038/s41586-018-0414-6 (2018).

53 Asfaha, S. *et al.* Krt19(+)/Lgr5(-) Cells Are Radioresistant Cancer-Initiating Stem Cells in
the Colon and Intestine. *Cell Stem Cell* **16**, 627-638, doi:10.1016/j.stem.2015.04.013
(2015).

54 Tetteh, P. W. *et al.* Replacement of Lost Lgr5-Positive Stem Cells through Plasticity of
Their Enterocyte-Lineage Daughters. *Cell Stem Cell* **18**, 203-213,
doi:10.1016/j.stem.2016.01.001 (2016).

55 Beucher, A. *et al.* The homeodomain-containing transcription factors Arx and Pax4
control enteroendocrine subtype specification in mice. *PLoS One* **7**, e36449,
doi:10.1371/journal.pone.0036449 (2012).

56 Busslinger, G. A. *et al.* Human gastrointestinal epithelia of the esophagus, stomach, and
duodenum resolved at single-cell resolution. *Cell Rep* **34**, 108819,
doi:10.1016/j.celrep.2021.108819 (2021).

57 Bekku, S. *et al.* Carbonic anhydrase I and II as a differentiation marker of human and rat
colonic enterocytes. *Res Exp Med (Berl)* **198**, 175-185, doi:10.1007/s004330050101
(1998).

58   Gonzalez, L. M., Williamson, I., Piedrahita, J. A., Blikslager, A. T. & Magness, S. T. Cell
lineage identification and stem cell culture in a porcine model for the study of intestinal
epithelial regeneration. *PLoS One* **8**, e66465, doi:10.1371/journal.pone.0066465 (2013).

59   Dieterich, W., Neurath, M. F. & Zopf, Y. Intestinal ex vivo organoid culture reveals
altered programmed crypt stem cells in patients with celiac disease. *Sci Rep* **10**, 3535,
doi:10.1038/s41598-020-60521-5 (2020).

60   Mohammadi, S. *et al.* Assessing donor-to-donor variability in human intestinal organoid
cultures. *Stem Cell Reports*, doi:10.1016/j.stemcr.2021.07.016 (2021).

61   Yan, K. S. *et al.* Intestinal Enteroendocrine Lineage Cells Possess Homeostatic and
Injury-Inducible Stem Cell Activity. *Cell Stem Cell* **21**, 78-90 e76,
doi:10.1016/j.stem.2017.06.014 (2017).

62   Jepeal, L. I., Boylan, M. O. & Michael Wolfe, M. GATA-4 upregulates glucose-dependent
insulinotropic polypeptide expression in cells of pancreatic and intestinal lineage. *Mol*
Cell Endocrinol **287**, 20-29, doi:10.1016/j.mce.2008.01.024 (2008).

63   Walker, E. M., Thompson, C. A. & Battle, M. A. GATA4 and GATA6 regulate intestinal
epithelial cytodifferentiation during development. *Dev Biol* **392**, 283-294,
doi:10.1016/j.ydbio.2014.05.017 (2014).

64   Walker, E. M., Thompson, C. A., Kohlnhofer, B. M., Faber, M. L. & Battle, M. A.
Characterization of the developing small intestine in the absence of either GATA4 or
GATA6. *BMC Res Notes* **7**, 902, doi:10.1186/1756-0500-7-902 (2014).

65   Kohlnhofer, B. M., Thompson, C. A., Walker, E. M. & Battle, M. A. GATA4 regulates
epithelial cell proliferation to control intestinal growth and development in mice. *Cell Mol*
Gastroenterol Hepatol **2**, 189-209, doi:10.1016/j.jcmgh.2015.11.010 (2016).

66   Girard, R. *et al.* HNF4alpha is a novel regulator of intestinal glucose-dependent
insulinotropic polypeptide. *Sci Rep* **9**, 4200, doi:10.1038/s41598-019-41061-z (2019).

67   Zou, S. & Kumar, U. Cannabinoid Receptors and the Endocannabinoid System:
Signaling and Function in the Central Nervous System. *Int J Mol Sci* **19**,
doi:10.3390/ijms19030833 (2018).

68   Tremblay, J. J. & Viger, R. S. Transcription factor GATA-4 is activated by
phosphorylation of serine 261 via the cAMP/protein kinase a signaling pathway in
gonadal cells. *J Biol Chem* **278**, 22128-22135, doi:10.1074/jbc.M213149200 (2003).

69   Dusing, M. R. & Wiginton, D. A. Epithelial lineages of the small intestine have unique
patterns of GATA expression. *J Mol Histol* **36**, 15-24, doi:10.1007/s10735-004-2908-9
(2005).

70   Bosse, T. *et al.* Gata4 is essential for the maintenance of jejunal-ileal identities in the
adult mouse small intestine. *Mol Cell Biol* **26**, 9060-9070, doi:10.1128/MCB.00124-06
(2006).

71 Netherland, C. & Thewke, D. P. Rimonabant is a dual inhibitor of acyl CoA:cholesterol
acyltransferases 1 and 2. *Biochem Biophys Res Commun* **398**, 671-676,
doi:10.1016/j.bbrc.2010.06.134 (2010).

72 Ferraz-de-Souza, B. *et al.* Sterol O-acyltransferase 1 (SOAT1, ACAT) is a novel target of
steroidogenic factor-1 (SF-1, NR5A1, Ad4BP) in the human adrenal. *J Clin Endocrinol*
Metab **96**, E663-668, doi:10.1210/jc.2010-2021 (2011).

73 Shroyer, N. F. *et al.* Intestine-specific ablation of mouse atonal homolog 1 (Math1)
reveals a role in cellular homeostasis. *Gastroenterology* **132**, 2478-2488,
doi:10.1053/j.gastro.2007.03.047 (2007).

74 Yang, Q., Bermingham, N. A., Finegold, M. J. & Zoghbi, H. Y. Requirement of Math1 for
secretory cell lineage commitment in the mouse intestine. *Science* **294**, 2155-2158,
doi:10.1126/science.1065718 (2001).

75 Lewis, S. K. *et al.* DNA Methylation Analysis Validates Organoids as a Viable Model for
Studying Human Intestinal Aging. *Cell Mol Gastroenterol Hepatol* **9**, 527-541,
doi:10.1016/j.jcmgh.2019.11.013 (2020).

76 Kasendra, M. *et al.* Development of a primary human Small Intestine-on-a-Chip using
biopsy-derived organoids. *Sci Rep* **8**, 2871, doi:10.1038/s41598-018-21201-7 (2018).

77 Dekkers, J. F. *et al.* High-resolution 3D imaging of fixed and cleared organoids. *Nat*
Protoc **14**, 1756-1771, doi:10.1038/s41596-019-0160-8 (2019).

78 Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
Methods **9**, 676-682, doi:10.1038/nmeth.2019 (2012).

79 Borten, M. A., Bajikar, S. S., Sasaki, N., Clevers, H. & Janes, K. A. Automated brightfield
morphometry of 3D organoid populations by OrganoSeg. *Sci Rep* **8**, 5319,
doi:10.1038/s41598-017-18815-8 (2018).

80 Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-
seq data using regularized negative binomial regression. *Genome Biol* **20**, 296,
doi:10.1186/s13059-019-1874-1 (2019).

81 Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587
e3529, doi:10.1016/j.cell.2021.04.048 (2021).
1107

1108 ACKNOWLEDGEMENTS

1109 We thank members of the Breault laboratory, J. Majzoub, S. Bonner-Weir and J. Turner for

1110 helpful discussions. This research was supported by National Institute of Health (NIH) awards

1111 T32DK769937, K12DK09472109, the Juvenile Diabetes Research Foundation Advanced

1112 Postdoctoral Fellowship award 3-APF-2020-929-A-N, and Pediatric Endocrine Society

1121

AUTHOR CONTRIBUTIONS

D.Z. designed and performed experiments, analyzed data, prepared figures and co-wrote the

manuscript; E.S., P.M., and W.Q. performed experiments, analyzed data and edited the

manuscript; J.S.C. performed and analyzed the single cell RNA sequencing experiment; X.Y.

and E.P.S. performed cell culture experiments; M.S.S. performed flow cytometry experiments;

S.D. and S.H. assisted in qPCR and immunofluorescence experiments; J.M.K. provided

conceptual advice and reagents; D.L.C. and J.O.M. provided conceptual advice and edited the

manuscript, and D.T.B. directed the project and co-wrote the manuscript.

1130

COMPETING INTERESTS

The author J.M.K. holds equity in Frequency Therapeutics, a company that has an option to

license IP generated by J.M.K and that may benefit financially if the IP is licensed and further

validated. The interests of J.M.K. was reviewed and are subject to a management plan

overseen by their institutions in accordance with their conflict-of-interest policies. J.O.M. reports

compensation for consulting services with Cellarity and Hovione. The remaining authors declare

no competing interests. A patent was filed by D.Z. and D.T.B covering compositions and

methods for regulating enteroendocrine cell differentiation and uses thereof as described in the

manuscript.

1140

FIGURE LEGENDS

Figure 1. Base Differentiation Media Induces CHGA Expression in a Wnt-dependent Manner

(a) Representative light microscopy of enteroids (whole well) grown in growth media (GM) for 14

days (G14) or two days in GM followed by 12 days in differentiation media (DM, G2D12).

Specific culture schematic located above each panel, respectively. Scale bar = 1 mm.

(b) qPCR analysis of enteroendocrine (EE) markers of enteroids grown in either G14 or G2D12

compared to duodenal whole mucosa and normalized to *18S*. Dotted line denotes expression

level in whole duodenal mucosa. Representative experiment showing n = 3 wells for each

condition from a single enteroid line. *CHGA* = chromogranin A, *PAX4* = paired box 4, *PDX1* =

pancreatic and duodenal homeobox 1, *NEUROD1* = neuronal differentiation 1, *NEUROG3* =

neurogenin 3, *CCK* = cholecystokinin, *SST* = somatostatin, *GIP* = glucose-dependent

insulinotropic peptide, ND = not detectable in one or more samples.

(c and d) Representative immunofluorescence staining of CHGA (magenta) in enteroids (whole

well) treated with G2D12. Boxed portion in (c) shown magnified in (d). DNA (4′,6-diamidino-2-

phenylindole (DAPI) blue). Scale bars = 1 mm (c) and 50 µm (d).

(e) Left two panels: Representative flow cytometry plots of CHGA-positive (CHGA+) cells from

enteroids grown in either G14 or G2D12. Right panel: Percentage of CHGA+ cells per well.

Representative experiment showing n = 3 wells from each condition from single enteroid line.

(f) qPCR analysis of EE gene markers of enteroids grown in either G2D12 without Wnt (G2D12-

Wnt) or with Wnt (G2D12+Wnt) compared to whole mucosa and normalized to *18S*. Dotted line

denotes expression level in whole duodenal mucosa. *MUC2* = Mucin 2, *ALPI* = Intestinal

alkaline phosphatase. Representative experiment showing n = 3 wells from each condition from

a single enteroid line. ****$p < 0.0001$.

Bars show mean ± SEM; two-way ANOVA with Tukey correction for multiple comparisons (b,f);

two-tailed unpaired *t* test (e). Each experiment repeated with at least three different enteroid

lines. Specific conditions were excluded from statistical analysis if the data from one or more

samples was labeled as not detectable. Source data are provided as a Source Data file.

1168

Figure 2. Differentiation with Small Molecules Rimonabant and SP600125 Induces CHGA

Expression

(a) Representative light microscopy of enteroids (whole well) grown in growth media for two

days followed by 12 days in differentiation media with rimonabant and SP600125 (RSP). Culture

schematic located above panel. Scale bar = 1 mm.

(b) qPCR analysis of enteroendocrine markers of enteroids grown in G14, G2D12 or RSP

compared to duodenal whole mucosa and normalized to *18S*. Dotted line denotes expression

level in whole duodenal mucosa. Representative experiment showing n = 3 wells from each

condition from a single enteroid line. *CHGA* = chromogranin A, *PAX4* = paired box 4, *PDX1* =

pancreatic and duodenal homeobox 1, *NEUROD1* = neuronal differentiation 1, *NEUROG3* =

neurogenin 3, *CCK* = cholecystokinin, *SST* = somatostatin, *GIP* = glucose-dependent

insulinotropic peptide, ND = not detectable in one or more samples. **p = 0.0025; ***p = 0.0009

(*PDX1*), 0.0003 (*CCK*); ****p < 0.0001.

(c) Representative immunofluorescence staining of CHGA (magenta) in enteroids (whole well)

treated with G14, G2D12 and RSP. Boxed portion magnified in lower right corner. Nuclei (4',6-

diamidino-2-phenylindole (DAPI), blue). Scale bars = 1mm and 50µm (boxed portion).

(d) Percentage of enteroids with positive CHGA staining in G14, G2D12 and RSP treatments.

Table above graph shows the total number of enteroids examined per condition. Average

results are from three separate experiments from three different enteroid lines or passages.

****p < 0.0001.

(e) Left three panels: Representative flow cytometry plots of CHGA+ cells from enteroids grown

in G14, G2D12 or RSP. Right panel: Percentage of CHGA+ cells per well. Representative

experiment showing n = 3 wells from each condition from a single enteroid line. ****p < 0.0001.

Bars show mean ± SEM; two-way ANOVA with Tukey correction for multiple comparisons (b);

one-way ANOVA with Tukey correction for multiple comparisons (d,e). Each experiment

repeated with at least three different enteroid lines. Specific conditions were excluded from

statistical analysis if the data from one or more samples was labeled as not detectable. Source

data are provided as a Source Data file.

1197

Figure 3. Differentiation with Small Molecule AS1842856 Induces CHGA Expression

(a) Representative light microscopy of enteroids (whole well) grown in growth media for two

days followed by 12 days in differentiation media with AS1842856 (AS). Culture schematic

located above panel. Scale bar = 1 mm.

(b) qPCR analysis of enteroendocrine markers of enteroids grown in G14, G2D12 or AS

compared to whole duodenal mucosa and normalized to *18S*. Dotted line denotes expression

level in whole duodenal mucosa. Representative experiment showing n = 3 wells from each

condition from a single enteroid line. *CHGA* = chromogranin A, *PAX4* = paired box 4, *PDX1* =

pancreatic and duodenal homeobox 1, *NEUROD1* = neuronal differentiation 1, *NEUROG3* =

neurogenin 3, *CCK* = cholecystokinin, *SST* = somatostatin, *GIP* = glucose-dependent

insulinotropic peptide, ND = not detectable in one or more samples. *p = 0.038 (*NEUROG3*),

0.0130 (*CCK*); ***p = 0.0004 (*PDX1*, G14 to AS), 0.0009 (*PDX1*, G2D12 to AS); ****p < 0.0001.

(c) Representative immunofluorescence staining of CHGA (magenta) in enteroids (whole well)

treated with G14, G2D12 and AS. Boxed portion magnified in lower right corner. DNA (4′,6-

diamidino-2-phenylindole (DAPI), blue). Scale bars = 1 mm and 50 µm (boxed portion).

(d) Percentage of enteroids with positive CHGA staining in G14, G2D12 and AS treatments.

Table above graph shows the total number of enteroids examined per condition. Average

results are from three separate experiments from three different enteroid lines or passages.

****$p < 0.0001$.

(e) Left three panels: Representative flow cytometry plots of CHGA+ cells from enteroids grown

in G14, G2D12 or AS. Right panel: Percentage of CHGA+ cells per well. Representative

experiment showing n = 3 wells from each condition from a single enteroid line. ****$p < 0.0001$.

Bars show mean ± SEM; two-way ANOVA with Tukey correction for multiple comparisons (b);

one-way ANOVA with Tukey correction for multiple comparisons (d,e). Each experiment

repeated with at least three different enteroid lines. Specific conditions were excluded from

statistical analysis if the data from one or more samples was labeled as not detectable. Source

data are provided as a Source Data file.

1225

Figure 4. scRNA-seq Profiling of Enteroids Cultured with Rimonabant/SP600125 or AS1842856

(a) Uniform manifold approximation and projection (UMAP) visualization of 14,767 cells

summarizing enteroid differentiation from all samples, color labeled by culture condition.

(b) UMAP visualization from (a), color labeled by broad annotated cell identity, following Louvain

clustering.

(c) Dot plot of the average scaled expression (measured by average Pearson residual) of

canonical markers of various gut epithelial cell types, plotted against cluster identity.

(d) Proportional abundance of epithelial cell subsets by enteroid culture protocol. Each culture

condition consists of three different enteroid lines from distinct human donors, as denoted by

data point shape. Bars show mean ± SEM; Kruskal-Wallis test with Dunn's post-hoc analysis

displayed. P values were adjusted using Bonferroni correction for multiple comparisons. *$p =$

0.0219 (Intestinal Stem Cells and Enteroendocrine Cells), 0.0338 (Enterocytes).

(e) UMAP visualization of 14,767 cells divided by culture condition and colored by cell identity.

Trajectory analysis of each protocol was calculated using scVelo and the vector field was

overlaid on top of each UMAP. Arrows represent smoothed averages of the estimated cellular

differentiation trajectory, with arrow thickness corresponding to the "speed" of differentiation.

(f) UMAP visualization from (a), with individual cells colored by their EE cell module score. Each

score was scaled on a range from 0 to 1.

(g) Violin plots of the module score described in (f) split across culture condition. The effect size

between culture conditions was calculated as Cohen's d; \$\$ $0.8 < d < 1.2$. Source data are

provided as a Source Data file.

1247

Figure 5. Combinations of AS1842856 and Rimonabant/SP600125 Induce Different Levels of

Enteroendocrine Marker Expression

(a) qPCR analysis of enteroendocrine markers of enteroids grown in AS, AS for 6 days, followed

by RSP only for 6 days (AS→RSP) and AS for 6 days, followed by AS and RSP for 6 days

(AS→RASP) compared to enteroids grown in RSP and normalized to *18S*. Representative

experiment showing n = 3 wells from each condition from a single enteroid line. *CHGA* =

chromogranin A, *PAX4* = paired box 4, *PDX1* = pancreatic and duodenal homeobox 1,

NEUROD1 = neuronal differentiation 1, *NEUROG3* = neurogenin 3, *CCK* = cholecystokinin,

SST = somatostatin, *GIP* = glucose-dependent insulinotropic peptide. *p = 0.0424 (*PAX4*),

0.0123 (*SST*); **p = 0.039 (*CHGA*, RSP to AS→RSP), 0.0059 (*CHGA*, AS to AS→RSP), 0.0064

(*CHGA*, AS to AS→RASP), 0.0019 (*SST*), 0.0010 (*GIP*); ***p = 0.0006 (*NEUROD1*), 0.0004

(*NEUROG3*), 0.0003 (*CCK*), 0.0001 (*GIP*); ****p < 0.0001.

(b) Representative immunofluorescence staining of CHGA (magenta) in enteroids (whole well)

treated with G14, G2D12, RSP, AS, AS→RSP, and AS→RASP. DNA (4′,6-diamidino-2-

phenylindole (DAPI), blue). Scale bar = 1 mm.

(c) Percentage of enteroids with positive CHGA staining in G14, G2D12, AS, RSP, AS→RSP,

and AS→RASP treatments. Table above graph shows the total number of enteroids examined

per condition. Average results are from three separate experiments from three different enteroid

lines or passages. ****$p < 0.0001$.

(d) Left six panels: Representative flow cytometry plots of CHGA+ cells from enteroids grown in

G14, G2D12, RSP, AS, AS→RSP, and AS→RASP. Right panel: Percentage of CHGA+ cells

per well. Representative experiment showing n = 3 wells from each condition from a single

enteroid line. **$p < 0.0041$ (G14 to RSP), 0.0045 (G2D12 to RSP), 0.0020 (AS to AS→RASP);

****$p < 0.0001$.

Bars show mean ± SEM; two-way ANOVA with Tukey correction for multiple comparisons (a);

one-way ANOVA with Tukey correction for multiple comparisons (c,d). Each experiment

repeated with at least three different enteroid lines. Source data are provided as a Source Data

file.

1276

Figure 6. Multiple Differentiation Conditions Induce Hormone Production

(a) Representative immunofluorescence staining of somatostatin (SST, green) and

chromogranin A (CHGA, magenta) in enteroids treated with G14, G2D12, RSP, AS, AS→RSP,

and AS→RASP. DNA (4′,6-diamidino-2-phenylindole (DAPI), blue). Scale bar = 50µm.

(b) Percentage of enteroids with SST staining in G14, G2D12, RSP, AS, AS→RSP, and

AS→RASP treatments. Average results are from three different enteroid lines or passages. **p

= 0.0017 (G14 to RSP, G2D12 to RSP), 0.0029 (G14 to AS, G2D12 to AS); ***p = 0.0004 (G14

to AS→RSP, G2D12 to AS→RSP), 0.0006 (G14 to AS→RASP, G2D12 to AS→RASP).

(c) Representative immunofluorescence staining of serotonin (5HT, green) and CHGA

(magenta) in enteroids treated with G14, G2D12, RSP, AS, AS→RSP and AS→RASP. DNA

(DAPI, blue). Scale bar = 50 µm.

(d) Percentage of enteroids with 5HT staining in G14, G2D12, RSP, AS, AS→RSP and

AS→RASP treatments. Average results are from three different enteroid lines or passages. *p =

0.0183 (RSP to AS), 0.0134 (RSP to AS→RASP); **p = 0.0014; ***p = 0.0009; ****p < 0.0001.

(e) Representative immunofluorescence staining of glucose-dependent insulinotropic peptide

(GIP, green) and CHGA (magenta) in enteroids treated with G14, G2D12, RSP, AS, AS→RSP,

and AS→RASP. DNA (DAPI, blue). Scale bar = 50 µm.

(f) Percentage of enteroids with GIP staining in G14, G2D12, RSP, AS, AS→RSP, and

AS→RASP treatments. Average results are from three different enteroid lines or passages. ***p

= 0.0001; ****p < 0.0001.

(g) Representative immunofluorescence staining of cholecystokinin (CCK, green) and CHGA

(magenta) in enteroids treated with G14, G2D12, RSP, AS, AS→RSP, and AS→RASP. DNA

(DAPI, blue). Scale bar = 50 µm.

(h) Percentage of enteroids with CCK staining in G14, G2D12, RSP, AS, AS→RSP, and

AS→RASP treatments. Average results are from three different enteroid lines or passages. ***p

= 0.0002 (RSP to AS), 0.0001 (AS to AS→RSP); ****p < 0.0001.

Bars show mean ± SEM; one-way ANOVA with Tukey correction for multiple comparisons

(b,d,f,h). Tables above graphs show the total number of enteroids examined per condition. Each

experiment repeated with at least three different enteroid lines. Source data are provided as a

Source Data file.

1307

Figure 7. Differentiation Conditions Induce Hormone Secretion

(a) Serotonin (5HT) ELISA of conditioned media from the last two days of differentiation of

enteroids grown in G14, G2D12, RSP, AS, AS→RSP, and AS→RASP. Representative

experiment showing n = 3 wells from each condition from a single enteroid line. *p = 0.049; **p

= 0.0049 (RSP to AS→RSP), 0.0011 (AS to AS→RSP); ***p = 0.0005 (G14 to AS→RSP and

G2D12 to AS→RSP); ****p < 0.0001.

(b) 5HT ELISA of AS conditioned media collected after 24 hours on day 13 (solid bar) and after

24 hours with forskolin (Fsk) on day 14 (striped bar). Representative experiment showing n = 3

wells from each condition from a single enteroid line. **p = 0.0015.

(c) Glucose-dependent insulinotropic peptide (GIP) ELISA of conditioned media from the last

two days of differentiation of enteroids grown in G14, G2D12, RSP, AS, AS→RSP, and

AS→RASP. Representative experiment showing n = 3 wells from each condition from a single

enteroid line. **p = 0.0013 (AS to RSP), 0.0031 (RSP to AS→RSP); ***p = 0.0009 (G14 to RSP,

G2D12 to RSP, and RSP to AS→RASP); ****p < 0.0001.

(d) GIP ELISA of AS→RSP conditioned media collected after 24 hours on day 13 (solid bar) and

after 24 hours with Fsk on day 14 (striped bar). Representative experiment showing n = 3 wells

from each condition from a single enteroid line. **p = 0.0025.

Bars show mean ± SEM; one-way ANOVA with Tukey correction for multiple comparisons (a,c);

two-tailed unpaired *t* test (b,d). Dotted line at 2.6 ng/mL represents the lower limit of detection

for the 5HT ELISA (a,b). Dotted line at 4.2 pg/mL represents the lower limit of detection for the

GIP ELISA (c,d). Each experiment repeated with at least three different enteroid lines. Source

data are provided as a Source Data file.

1330

Figure 8. Induction of the Enteroendocrine Lineage in Rectoids

(a) Representative light microscopy of rectoids (whole well) grown in G14 or G2D12. Scale bar

= 1 mm.

(b) qPCR analysis of EE markers of rectoids grown in either G14 or G2D12 compared to whole

rectal mucosa (dotted line) and normalized to *18S*. Representative experiment showing n = 3

wells for each condition from a single rectoid line. *CHGA* = chromogranin A, *PAX4* = paired box

4, *NEUROD1* = neuronal differentiation 1, *NEUROG3* = neurogenin 3, *SST* = somatostatin,

GCG = glucagon, *PYY* = peptide YY, ND = not detectable in one or more samples. ****p <

0.0001.

(c) Representative immunofluorescence staining of CHGA (magenta) in rectoids (whole well)

treated with G14 and G2D12. DNA (4ʹ,6-diamidino-2-phenylindole (DAPI), blue). Scale bar = 1

mm.

(d) Percentage of rectoids with CHGA staining in G14 and G2D12 treatments. Average results

are from three different rectoid lines or passages. Table above graph shows the total number of

rectoids examined per condition.   **p < 0.0022.

(e) Left two panels: Representative flow cytometry plots of CHGA+ cells from rectoids grown in

G14 and G2D12. Right panel: Percentage of CHGA+ cells per well. Representative experiment

showing n = 3 wells from each condition from a single rectoid line. ***p < 0.0004.

Bars show mean ± SEM; two-way ANOVA with Tukey correction for multiple comparisons (b);

two-tailed unpaired *t* test (d,e). Each experiment repeated with at least three different rectoid

lines. Specific conditions were excluded from statistical analysis if the data from one or more

samples was labeled as not detectable. Source data are provided as a Source Data file.

1353

Figure 9. Hormone Production and Secretion in Rectoids

(a) Representative immunofluorescence staining of glucagon-like peptide-1 (GLP-1, green) and

CHGA (magenta) in rectoids treated with G14 and G2D12. DNA (4ʹ,6-diamidino-2-phenylindole

(DAPI), blue). Scale bar = 50 µm.

(b) Percentage of rectoids with GLP-1 staining in G14 and G2D12. Average results are from

three different rectoid lines or passages. ****p < 0.0001.

(c) Representative immunofluorescence staining of peptide YY (PYY, green) and CHGA

(magenta) in rectoids treated with G14 and G2D12. DNA (DAPI, blue). Scale bar = 50 µm.

(d) Percentage of rectoids with PYY staining in G14 and G2D12. Average results are from three

different rectoid lines or passages. *p < 0.0118.

(e) GLP-1 ELISA of conditioned media from the last two days of differentiation of rectoids grown

in G14 and G2D12. Representative experiment showing n = 3 wells from each condition from a

single rectoid line. *p < 0.0229.

(f) GLP-1 ELISA of G2D12 conditioned media collected after 24 hours on day 13 (solid bar) and

after 24 hours with forskolin (Fsk) on day 14 (striped bar). Representative experiment showing n

= 3 wells from each condition from a single rectoid line. **p < 0.0061.

(g) PYY ELISA of conditioned media from the last two days of differentiation of rectoids grown in

G14 and G2D12. Representative experiment showing n = 3 wells from each condition from a

single rectoid line. **p < 0.0056.

(h) PYY ELISA of G2D12 conditioned media collected after 24 hours on day 13 (solid bar) and

after 24 hours with forskolin (Fsk) on day 14 (striped bar). Representative experiment showing n

= 3 wells from each condition from a single rectoid line. ***p < 0.0006.

Bars show mean ± SEM; two-tailed unpaired *t* test (b,d,e,f,g,h). Dotted line at 25 pg/mL

represents the lower limit of detection for the GLP-1 ELISA (e,f). Dotted line at 7.3 pg/mL

represents the lower limit of detection for the PYY ELISA (g,h). Each experiment repeated with

at least three different rectoid lines. Tables above graphs show the total number of rectoids

examined per condition. Source data are provided as a Source Data file.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

a.

b.

c.

d.

Figure 8

Figure 9