Mechanical impact on the head has an antihypertensive effect

Authors:

Shuhei Murase[1,2,14], Naoyoshi Sakitani[1,14], Takahiro Maekawa[1], Daisuke Yoshino[3], Ayumu Konno[4], Hirokazu Hirai[4], Taku Saito[2], Sakae Tanaka[2], Keisuke Shinohara[5], Takuya Kishi[6], Yuki Yoshikawa[7], Takamasa Sakai[7], Makoto Ayaori[8], Hirohiko Inanami[9], Koji Tomiyasu[10], Toru Ogata[1,11], Atsushi Takashima[12], Masahiro Shinohara[1], Motoshi Nagao[1] and Yasuhiro Sawada[1,13]*

Affiliations:

[1]Department of Rehabilitation for Motor Functions, National Rehabilitation Center for Persons with Disabilities, Tokorozawa, Japan. [2]Department of Orthopaedic Surgery, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan. [3]Division of Advanced Applied Physics, Institute of Engineering, Tokyo University of Agriculture and Technology, Koganei, Japan. [4]Department of Neurophysiology & Neural Repair, Gunma University Graduate School of Medicine, Maebashi, Japan. [5]Department of Cardiovascular Medicine, Kyushu University Graduate School of Medical Sciences, Fukuoka, Japan. [6]Department of Cardiology, Graduate School of Medicine, International University of Health and Welfare, Okawa, Japan. [7]Department of Bioengineering, Graduate School of Engineering, The University of Tokyo, Tokyo, Japan.

[8]Tokorozawa Heart Center, Tokorozawa, Japan. [9]Inanami Spine & Joint Hospital/Iwai Orthopaedic

Medical Hospital, Iwai Medical Foundation, Tokyo, Japan. [10]Department of Internal Medicine,

[11]Center of Sports Science and Health Promotion, [12]Department of Assistive Technology,

[13]Department of Clinical Research, National Rehabilitation Center for Persons with Disabilities,

Tokorozawa, Japan.

[14]These authors contributed equally: Shuhei Murase, Naoyoshi Sakitani.

*Correspondence to Yasuhiro Sawada, Department of Clinical Research, National Rehabilitation

Center for Persons with Disabilities, 4-1 Namiki, Tokorozawa, Saitama 359-8555, Japan, Tel: +81-

4-2995-3100, Fax: +81-4-2995-0355, Email: ys454-ind@umin.ac.jp

Abstract

Nervous cell functions are known to be physiologically regulated by mechanical factors in the

brain. However, it remains unclear whether mechanical interventions can modulate the

pathophysiological processes underlying brain-related disorders and modify their

consequences. Here we show that passive head motion of hypertensive rats, which reproduces

mechanical accelerations generated at their heads during treadmill running at a moderate

velocity, decreases the expression of angiotensin II type 1 receptor (AT1R) in astrocytes in

their rostral ventrolateral medulla (RVLM). This decrease results in lowering their blood

pressure. Passive head motion generates interstitial fluid movement that is estimated to exert

shear stress with average magnitude of a few Pa on cells in rats' brainstem. Fluid shear stress

of a relevant magnitude decreases AT1R expression in cultured astrocytes, but not in neuronal

cells. Furthermore, in hypertensive rats, inhibition of movement of interstitial fluid by its

gelation with reactive polyethylene glycol injected into the RVLM eliminates the ability of

passive head motion to decrease their blood pressure and AT1R expression in RVLM

astrocytes. Consistent with these results from animal experiments, vertically oscillating chair

riding of hypertensive adult humans, which reproduces mechanical accelerations generated at

their heads during light jogging or fast walking, lowers their blood pressure. Our findings

indicate that moderate mechanical impact on the head has an antihypertensive effect by

modulating the function of RVLM astrocytes through interstitial fluid shear stress. We

anticipate mechanical regulation to underlie a variety of positive effects of physical exercise

on human health, particularly those related to brain functions.

Introduction

Hypertension, which is a major cause of stroke and cardiovascular diseases, is the biggest risk

factor for death worldwide[1]. Whereas numerous antihypertensive drugs have been developed and

used clinically, physical exercise is known to be effective for the treatment and prevention of

essential hypertension[2,3], which comprises the majority (>90%) of human hypertension[4]. Although

long-term regulation of blood pressure (BP) has been recognized to be largely dependent on sodium

excretion adjusting systems, which mainly involve kidney functions[5], elevated activity of the

sympathetic nervous system also importantly contributes to the development of hypertension[6-8].

Rostral ventrolateral medulla (RVLM), which is located in the brainstem, plays a critical role in

determining the basal activity of the sympathetic nervous system, and its functional integrity is

essential for the maintenance of basal vasomotor tone and regulation of BP[6,9]. Angiotensin II (Ang

II) is the major bioactive peptide of the renin-angiotensin system (RAS), and is known to regulate

BP as well as other biological processes such as cell growth/apoptosis/migration, inflammation, and

fibrosis[10]. The biological effects of Ang II are mediated by its interaction with two distinct high-

affinity G protein-coupled receptors, Ang II type 1 receptor (AT1R) and type 2 receptor. Of these

receptors, AT1R is responsible for most of the known physiological and pathophysiological

processes related to Ang II. Whereas RAS is involved in the functional regulation of various

"peripheral" organs and tissues such as kidney and vessels, it also regulates brain functions within

the blood-brain barrier, including the control and maintenance of sympathetic nerve activity and

cognitive ability[11]. In particular, the role of AT1R signaling in the RVLM in cardiovascular

regulation has been extensively studied and demonstrated. For example, the pressor/depressor

responses to Ang II and Ang II antagonists, injected into the RVLM have been reported to be

enhanced in spontaneously hypertensive rats (SHRs)[12,13]. We have previously demonstrated that

treadmill running at moderate velocities alleviates the sympathetic nerve activity, involving

attenuation of AT1R signaling in the RVLM of stroke-prone spontaneously hypertensive rats

(SHRSPs)[14], a substrain of SHRs that exhibit severer hypertension as compared with SHRs[15].

However, the details about the changes in AT1R signaling in the RVLM of these hypertensive rats

have yet to be elucidated. It remains unclear what type(s) of cells (e.g., neurons or astrocytes) are

primarily responsible for the altered AT1R signaling in the RVLM of SHRs or SHRSPs.

Furthermore, the causal relationship between the increased AT1R signal activity in the RVLM and

high BP in SHRs or SHRSPs in their steady state (i.e., apart from their responses to

pharmacological interventions) is left unrevealed.

AT1R has also been shown to play a vital role in regulating a variety of physiological or

pathological processes, including cellular responses to mechanical perturbations[16,17]. For example,

mechanical stretching of cardiac myocytes activates AT1R signaling[18], and fluid shear stress (FSS)

of average 1.5 Pa lowers AT1R expression in human vein endothelial cells[19]. Although intervening

the Ang II-AT1R system through pharmacological approaches, such as administration of

angiotensin-converting enzyme inhibitor or selective AT1R blocker, has been established as an

effective therapeutic strategy for hypertension[20], mechano-responsive attenuation of AT1R signaling

has not been clinically utilized as an antihypertensive measure.

Many of physical workouts, particularly aerobic exercises, involve vertical body motions,

which generate mechanical impacts (accelerations) on the head at the time of foot contact with the

ground (i.e., landing). The importance of mechanical loads is well established in the physiological

regulation of bones, the stiffest organ that only allows tiny deformation[21]. Osteocytes, the

mechanosensory cells embedded in bones[22], are assumed to undergo minimal deformations under

physiological conditions. We have reported that FSS on osteocytes derived from interstitial fluid

flow induced upon physical activity plays an important role in maintaining bone homeostasis[23].

Given that the brain is not a rigid organ, minimally deforming forces or stress distribution changes

in the brain during exercise or even activity of daily living (e.g., walking) may produce beneficial

effects. We have previously shown that in the prefrontal cortex (PFC) of rodents, moderate

mechanical impact-induced FSS modulates serotonin signaling in the neurons in situ[24]. Based on

these previous findings together with the distribution of interstitial fluid throughout the whole brain,

we hypothesized that moderate mechanical impact on the head might have antihypertensive effects

involving FSS-mediated modulation of AT1R signaling in the RVLM.

Results

Application of cyclical mechanical impact to the head by passive motion lowers the BP in

SHRSPs. To determine the effects of a mechanical intervention of a moderate intensity on BP, we

first sought to develop an experimental system that reproduces the impact exerted on the head

during rats' treadmill running at a modest velocity (20 m/minute), a typical experimental

intervention to test the effects of physical exercise on rats[25,26]. In a recent study, we observed that

treadmill running of rats (20 m/minute) generated 5-mm vertical oscillation of their heads with ~1.0

x g peak accelerations and 2-Hz frequency; therefore, we developed a "passive head motion"

(PHM) system to produce 2-Hz 5-mm vertical oscillation exerting 1.0 x g acceleration peaks at the

heads of rats[24]. In the current study, we examined the effects of mechanical impact on BP in

SHRSPs, using the PHM system. Similar to the antihypertensive effect of treadmill running on

SHRs or SHRSPs that we and others reported previously[14,25,27], application of PHM (30

minutes/day, 28 consecutive days; see Fig. 1a) significantly lowered their BP (Fig. 1b,c) as

compared to their controls, whereas HR was not significantly affected by PHM (Fig. 1d).

Anesthesia alone (daily 30 minutes) did not significantly alter the BP in SHRSPs (Extended Data

Fig. 1a), indicating that the antihypertensive effect resulted specifically from PHM. The anti-cardiac

hypertrophy effect of PHM on SHRSPs (Fig. 1e) as well as the lack of these PHM effects on control

normotensive rats (Wistar-Kyoto: WKY) (Fig. 1b,c,e) were also consistent with previous reports

describing treadmill running as an antihypertensive intervention for SHRs[27]. As was observed in our

treadmill running experiments[14], PHM decreased 24-hour urinary norepinephrine excretion of

SHRSPs (Fig. 1f). This suggests that PHM mitigates the sympathetic hyperactivity[28]. Collectively,

these results support our hypothesis that cyclical moderate mechanical impact on the head has an

antihypertensive effect.

PHM down-regulates AT1R expression in RVLM astrocytes of SHRSPs. We then looked into

the mechanism of how PHM alleviated the development of hypertension in SHRSPs. We previously

reported that down-regulation of AT1R signaling in the RVLM is responsible for the treadmill

running-induced sympathoinhibition in SHRSPs[14]. Given the mechanical regulation of AT1R

expression in endothelial cells[19], we examined whether PHM modulated AT1R expression in RVLM

neurons and astrocytes of SHRSPs. In our histochemical analysis, we defined neuronal nuclei

(NeuN)-positive cells as neurons[29] and glial fibrillary acidic protein (GFAP)-positive cells as

astrocytes[30]. PHM (30 minutes/day, 28 days) did not significantly change the relative population of

AT1R-expressing neurons and astrocytes in the RVLM of WKY rats (Fig. 1g). In contrast, 4-week

PHM significantly decreased the expression of AT1R in the astrocytes, but not in the neurons, of

SHRSPs' RVLM (Fig. 1h). Notably, AT1R expression in the RVLM neurons was comparable

between WKY rats and SHRSPs, either with and without PHM (Fig. 1i). In contrast, AT1R

expression was significantly higher in the RVLM astrocytes of SHRSPs without PHM (Fig. 1j,

column 3). PHM lowered the AT1R expression in the RVLM astrocytes of SHRSPs to the level

equivalent to that of WKY rats (Fig. 1j, columns 1, 2, and 4). Taken together, AT1R expression in

RVLM astrocytes appeared to be correlated with the antihypertensive effect of PHM on SHRSPs. In

line with this observation, 4-week treadmill running of SHRSPs also decreased the AT1R

expression in their RVLM astrocytes, but not neurons (Extended Data Fig. 1b−d).

PHM alleviates the sensitivity of RVLM in SHRSPs to Ang II or Ang II antagonist. We next

sought to examine whether the PHM-induced decrease in AT1R expression in the RVLM astrocytes

of SHRSPs (Fig. 1j, columns 3 and 4) was functionally relevant to the suppression of AT1R

signaling. To this end, we analyzed the pressor responses to Ang II injected into unilateral RVLM of

WKY rats and SHRSPs, either subjected to 4-week PHM or left sedentary under anesthesia (30

minutes/day, 28 days) (Fig. 2a). As we previously reported[14], SHRSPs without PHM exhibited

significantly greater pressor response to Ang II administered to RVLM than WKY rats (Fig. 2b,

compare between top and bottom of left panels; Fig. 2c, compare columns 1 and 3). Four-week

PHM ameliorated the pressor response to Ang II injected into the RVLM of SHRSPs, but not of

WKY rats (Fig. 2b, compare left and right; Fig. 2c, compare columns 1 vs. 2 and 3 vs. 4).

Furthermore, depressor responses to Ang II antagonist injected into unilateral RVLM[13] of WHY rats

and SHRSPs with and without 4-week PHM appeared to be approximate mirror images of the

pressor responses (Fig. 2d,e). These results support the functional relevance of the PHM-induced

decrease in AT1R expression in the RVLM astrocytes of SHRSPs (Fig. 1j).

To examine whether the increased AT1R expression in the RVLM astrocytes of SHRSPs was

associated with their development of hypertension, we manipulated AT1R signaling by introducing

exogenous expression of AT1R-associated protein (AGTRAP), which interacts with AT1R and

tempers Ang II-mediated signals by promoting AT1R internalization[31]. To this end, we used an

adeno-associated virus (AAV)-mediated gene delivery system[32]. AAV serotype 9 (AAV9) vectors

were injected locally to transduce RVLM cells (Fig. 3a and Extended Data Fig. 2a). To achieve

astrocyte- and neuron-specific gene expression, we used AAV9 vectors that harbored mouse GFAP

promoter (AAV-GFAP) and rat neuron-specific enolase (NSE) promoter (AAV-NSE), respectively

(Fig. 3a). Because these vectors contained a region encoding GFP and 2A sequence of porcine

teschovirus-1 (P2A; self-cleaving peptides[33]) (Fig. 3a), observation of green fluorescence allowed

us to identify the cells in which transgene was expressed (Fig. 3b,c and Extended Data Fig. 2a−e).

AAV-mediated expression of AGTRAP in astrocytes (Fig. 3b) but not in neurons (Fig. 3c) of

bilateral RVLMs in SHRSPs significantly lowered the BP as compared with their control SHRSPs

in which only GFP was virally expressed in RVLM astrocytes or neurons (Fig. 3d,e). Furthermore,

AAV-mediated expression of AGTRAP in astrocytes, but not neurons, of SHRSPs' bilateral RVLMs

decreased 24-hour urinary norepinephrine excretion (Fig. 3f). Injection of control AAV vector

(GFAP-control or NSE-control) did not significantly affect the BP of SHRSPs (Extended Data Fig.

2f). These results support the importance of AT1R signal intensity in RVLM astrocytes for SHRSPs'

development of hypertension and sympathetic hyperactivity, as well as the physiological relevance

of PHM-induced decrease in AT1R expression we observed in the RVLM astrocytes of SHRSPs

(Fig. 1j).

PHM generates low-amplitude pressure waves and induces interstitial fluid movement in rat

RVLM. We then sought to determine the physical effects that PHM produced in rat RVLM. To do

so, we analyzed local pressure changes using a telemetry pressure sensor (Fig. 4a) as we described

previously[24]. PHM generated pressure waves (changes) with ~1.2 mm Hg peak amplitude (Fig.

4b−d). Hydrostatic pressure of this magnitude (~1.6 cm H_2O) is unlikely to initiate mechano-

responsive signaling in cells[34]. Postulating an analogy to osteocytes embedded in bones, the

function of which is known to be modulated by interstitial fluid flow-derived shear stress[23], we have

demonstrated that minimal stress distribution changes generate interstitial fluid flow in the brain,

resulting in shear stress-mediated regulation of nervous cell functions[24].

To analyze the PHM-induced interstitial fluid movement in the RVLM, we injected an iodine-

based contrast agent (Isovist®) into the RVLM of anesthetized rats, and tracked its distribution with

sequential computed tomography (CT) (Fig. 4e) as we previously did to quantify the movement of

intramuscular interstitial fluid[35]. We found that PHM significantly promoted Isovist spreading in the

rostral-caudal and dorsal-ventral (*y*- and *z*-axes, Fig. 4f) directions (Fig. 4g,h). In contrast, PHM did

not significantly affect the left-right spreading (*x*-axis, Fig. 4f) of Isovist (Fig. 4g,h). As we

observed in rat PFC[24], PHM induced interstitial fluid movement in the brainstem in defined

directions, rather than in an isotropic manner. Our simulative calculation suggests that PHM

subjected RVLM cells to interstitial fluid flow-derived shear stress with an average magnitude of

0.59−2.64 Pa (Supplementary Table 1). FSS of this magnitude is known to modify the physiological

function of astrocytes[36], leading us to hypothesize that FSS derived from interstitial fluid movement

(Fig. 4e−h) mediated the PHM-induced decrease in AT1R expression in RVLM astrocytes (Fig. 1j).

FSS on astrocytes decreases AT1R expression in vitro. To test this hypothesis, we conducted in

vitro FSS experiments. Based on our simulation (Supplementary Table 1), we applied pulsatile FSS

with an average magnitude of 0.7 Pa to cultured primary astrocytes, which were prepared from

astrocyte-GFP mice[37] (Extended Data Fig. 3a), using a system we previously reported[23,24,35,38].

Quantitative polymerase chain reaction (qPCR) and immunostaining analyses revealed that FSS

application (0.5 Hz, 30 minutes) significantly decreased AT1R expression in astrocytes for at least

24 hours (Fig. 5a−c). In contrast, FSS application to Neuro2A cells, which exhibit neuronal

phenotypes and morphology[39,40], did not decrease AT1R expression (Extended Data Fig. 3b−d).

Consistent with the decreasing effect of FSS on AT1R expression, the binding of fluorescently

labeled Ang II to cultured astrocytes was significantly decreased by pre-exposure to 30-minute FSS

(Fig. 5d,e). The duration (>24 hours) of FSS effects on AT1R expression in astrocytes (Fig. 5) poses

a possibility of cumulative effects of FSS applied repeatedly at 24-hour intervals. Therefore, these

in vitro observations suggest that FSS-mediated persistent reduction of AT1R expression is involved

in the effects of daily PHM application on BP (Fig. 1b,c) and AT1R expression in the RVLM

astrocytes (Fig. 1j) in SHRSPs.

Hindrance of interstitial fluid movement by hydrogel introduction in RVLM eliminates the

ability of PHM to decrease the AT1R expression in RVLM astrocytes and the BP in SHRSPs.

To examine whether interstitial fluid movement in the RVLM mediated the effects of PHM on BP

and AT1R expression in RVLM astrocytes of SHRSPs, we modulated local interstitial fluid

dynamics. Following the procedure we used to hinder the interstitial fluid movement in mouse

PFC[24], we gelled interstitial fluid in situ and deprived its fluidity by microinjecting mutually

reactive polyethylene glycol (PEG) gel-precursor (pre-gel) solutions to RVLM (Fig. 6a). Injected

pre-gel solution spread over rat RVLM, and gelled the interstitial fluid in situ (Extended Data Fig.

4a).

Hydrogel introduction in bilateral RVLMs eliminated the ability of PHM to decrease BP (Fig.

6b, black and orange lines; Fig. 6c, columns 2 and 3), urinary norepinephrine excretion (Fig. 6d,

columns 2 and 3), and AT1R expression in the RVLM astrocytes (Fig. 6e, bottom 2 rows; Fig. 6f,

columns 2 and 3) in SHRSPs. In contrast, hydrogel introduction increased BP (Fig. 6b, blue and

orange lines; Fig. 6c, columns 1 and 3), norepinephrine excretion (Fig. 6d, columns 1 and 3), and

AT1R expression in the RVLM astrocytes (Fig. 6e, top 2 rows; Fig. 6f, columns 1 and 3) of

SHRSPs subjected to PHM. AT1R expression in the RVLM neurons of SHRSPs remained unaltered

irrespective of the combination of PHM and hydrogel introduction (Fig. 6e,g). These results suggest

that hydrogel introduction in the RVLM disrupts the mechanism mediating the PHM-induced

decrease in BP, norepinephrine excretion, and AT1R expression in the RVLM astrocytes of

SHRSPs.

We previously showed that gelation only inhibits the fluidity of the fluid but does not restrict

the diffusion of small molecules inside the gel[24,41]. Consistently, hydrogel introduction did not

apparently delay or attenuate the depressor response to Ang II antagonist injected to the RVLM

(Extended Data Fig. 4b−d), indicating rapid solute diffusivity through the hydrogels. As was the

case with mouse PFC[24], gelation by injecting PEG solution did not affect overall cell number or

apoptosis in RVLM (Extended Data Fig. 4e,f), and survival or apoptosis of RVLM astrocytes

(Extended Data Fig. 4g,h) and neurons (Extended Data Fig. 4i,j). Collectively, the loss of PHM

effects by hydrogel introduction in bilateral RVLMs of SHRSPs (Fig. 6b−f) is likely to result from

hydrogel-mediated alteration in interstitial fluid dynamics, rather than decreased cell viability

caused by impaired nutrient supply or removal of metabolic wastes.

Vertically oscillating chair riding (VOCR) lowers BP in hypertensive adult humans. The

results from our animal experiments reveal the antihypertensive effect of mechanical accelerations

generated at the head during treadmill running at a moderate velocity. This prompted us to test

whether mechanical impact on the head lowered BP in hypertensive humans. As we observed light

jogging or fast walking (locomotion at the velocity of 7 km/hour) typically produce ~2 Hz vertical

acceleration waves with an amplitude of ~1.0 x *g* at the person's head (Extended Data Fig. 5a, top),

we constructed a chair that could vertically oscillate at the frequency of 2 Hz (Extended Data Fig.

5b) and produce ~1.0 x *g* acceleration waves at the head of the occupant (Extended Data Fig. 5a,

bottom).

Given that previous reports regarding antihypertensive effects of aerobic exercise typically

recommend ≥3−4 days per week (frequency) and ≥30 minutes per session or day (duration)[3], we set

our regimen of VOCR as 3 days/week (Monday, Wednesday, and Friday unless needed to assign

otherwise for particular reasons such as public holidays) and 30 minutes/day. Our study of protocol

1, in which we simply compared the subjects' BP and HR before and after 4-week (12 times)

VOCR (Extended Data Fig. 5c), showed that VOCR decreased BP in hypertensive humans (Fig.

7a).

We then conducted a human study of protocol 2, in which we followed the changes in subjects'

BP and HR more minutely (Extended Data Fig. 5d). Encouraged by the positive results from the

study of protocol 1, we adopted the same VOCR regimen as to its frequency (3 days/week) and

duration (30 minutes/day). Participants were subjected to serial blood sampling to measure plasma

catecholamines (epinephrine, norepinephrine, and dopamine) and renin activity, and serum

aldosterone and C-reactive protein (CRP) before and after the intervention period (Extended Data

Fig. 5d). To conduct the 2nd blood sampling on the next day of the last bout of VOCR, the

intervention period was extended from 4 weeks (total 12 times, typically 26 days) to 4.5 weeks

(total 14 times, 30−31 days) because blood sampling could not be done during weekends at our

hospital. BP, both systolic and diastolic, and MAP immediately after the intervention period

significantly decreased as compared with those immediately before the intervention period (Fig.

7b). Furthermore, we also observed an antihypertensive effect of VOCR when we defined "BP of

the week" to reliably detect the trends by reducing the influences from interday BP variability (see

Methods) (Fig. 7c and Extended Data Fig. 6a). Notably, the post-intervention follow-up showed that

the BP-lowering effect apparently persisted for 4 weeks, but not 5 weeks, after the last bout of

VOCR (Fig. 7c). Similar to our animal study, we did not observe significant changes in HR by the

VOCR intervention (Fig. 7a−c). Significant differences were not detected in the blood levels of

catecholamines, aldosterone, renin activity, and CRP between before and after the VOCR

intervention (Extended Data Fig. 6b).

　　　Collectively, our studies of protocols 1 and 2 suggest that VOCR, which reproduces

mechanical impact exerted on the head during light jogging or fast walking, has an antihypertensive

effect in hypertensive humans. Importantly, in none of 21 subjects (Supplementary Table 2),

apparent adverse events, including motion sickness and low back pain, were observed or manifested

in relation to the VOCR intervention.

Discussion

Essential hypertension is defined as high BP in which secondary causes including renovascular

disease, renal failure, pheochromocytoma, and primary aldosteronism are not present. It accounts

for >90% of human cases of hypertension, resulting from combinations of multiple genetic and

environmental factors[4]. Recently, brain (dys)function has been implicated in the pathogenesis of

essential hypertension; however, the details of their molecular link remain unclear[42]. Physical

exercise is proven to be effective as a therapeutic/preventative measure for essential hypertension[3].

Although the antihypertensive effect of physical exercise has been shown to involve normalization

of sympathetic hyperactivity in the brain[43], it is still unclear whether exercise directly modulates

brain function. In this study, PHM, which reproduced mechanical accelerations generated at the

head during treadmill running, allowed us to dissect bodily activity-derived physical effects.

Whereas AT1R signaling in both neurons and astrocytes of the RVLM have been reported to be

involved in regulating BP[44,45], we observed that AT1R expression in RVLM astrocytes was

increased in SHRSPs as compared to that in WKY rats (Fig. 1j). In contrast, AT1R expression in

RVLM neurons was comparable between WKY rats and SHRSPs (Fig. 1i), although AT1R

expression in RVLM neurons has been shown to play an important role in other animal model(s) of

hypertension[44]. Together with the decreases in BP and urinary epinephrine excretion of SHRSPs in

which RVLM astrocytes were transduced with AGTRAP gene (Fig. 3d−f), the intensity of AT1R

signaling in RVLM astrocytes appears to be critically involved in the pathogenesis of hypertension

and sympathetic hyperactivity in SHRSPs.

Four-week PHM decreased urinary norepinephrine excretion and AT1R expression in RVLM

astrocytes of SHRSPs to the levels almost equivalent to those of WKY rats (Fig. 1f,j). However,

PHM only partially alleviated the development of hypertension in SHRSPs to the extent similar to

the antihypertensive effects of treadmill running we and others previously reported[14,25] (Fig. 1b,c).

Therefore, it is evident that factors other than AT1R signaling in RVLM astrocytes contribute to the

pathogenesis of essential hypertension.

AT1R expression in cultured astrocytes decreased upon FSS application (Fig. 5a−c). This was

consistent with our findings that PHM and treadmill running decreased in AT1R expression in

RLVM astrocytes of SHRSPs (Fig. 1j and Extended Data Fig. 1d). However, the AT1R expression

level in RVLM astrocytes was low in WKY rats even without PHM (Fig. 1j), and this may raise a

concern regarding the physiological relevance of our in vitro FSS experiments using cultured

astrocytes. Yet, it has been reported that cultured astrocytes typically exhibit increased "reactivity",

and do not fully recapitulate physiological astrocytes in vivo[46]. We suggest that the FSS-induced

decrease in AT1R expression in cultured astrocytes we observed represents physiological functions

of astrocytes, despite that their increased basal AT1R expression may relate to unphysiological

aspects of two-dimensional culture on stiff substrates (culture plastics). Cells in static culture are

exposed to a complete absence of FSS, which may not be physiologically realized in vivo. Previous

reports describe increased extracellular fluid in brains of hypertensive humans[47] and altered

dynamics of intracerebral interstitial fluid of SHRs[48]. Aberrant regulation of RVLM astrocytes'

function that relates to altered interstitial fluid movement-derived FSS may underlie the

pathogenesis of essential hypertension.

PHM did not significantly alter the AT1R expression in SHRSPs' RVLM neurons (Fig. 1i), and

FSS did not decrease AT1R expression in cultured Neuro2A cells (Extended Data Fig. 3b−d). Yet,

we do not suspect that these results represent the absence of sensitivity of neurons to FSS or other

type(s) of mechanical stimulation, particularly because we observed PHM- and FSS-induced

internalization of 5-HT$_{2A}$ receptor expressed in mouse PFC neurons and Neuro2A cells,

respectively[24]. Alternatively, we speculate that PHM and FSS may mitigate hyperexpression of

AT1R related to aforementioned pathological status of SHRSPs' RVLM astrocytes or

unphysiological nature of cultured astrocytes. In line with this notion, PHM did not significantly

affect the AT1R expression in RVLM astrocytes in normotensive WKY rats (Fig. 1j). Relevantly,

we and others have reported mechanical loading (in vivo)- or FSS (in vitro)- induced alleviation of

inflammatory processes[23,49], which can transcriptionally increase AT1R expression[50].

Whereas the amplitude of pressure waves that PHM generated in rat RVLM was tiny (Fig.

4b−d), the magnitude of interstitial fluid movement-derived shear stress in rat RVLM appeared

comparable with that of FSS previously reported for vascular endothelium[51,52], interstitium of

bone[21] and skeletal muscle[35] (Supplementary Table 1b). Consistent with the lack of strict cell

specificity in many of cellular responses to mechanical forces[53], FSS-induced decrease in AT1R

expression, which was reported in vascular cells[19], was also observed in cultured astrocytes. Taken

together, we speculate that there may be common homeostasis-regulatory mechanisms at the

cellular level that involve fluid flow-derived shear stress of ~0.5 to a few Pa.

Based on our hypothesis concerning the similarity in the pathogenesis of high BP between

human essential hypertension and SHRSPs, we conducted human studies in which we intended to

reproduce the mechanical impact on the head that lowered the BP in SHRSPs. Although the

mechanism behind the apparent antihypertensive effect of VOCR remains to be determined, the

significant role of interstitial fluid dynamics in the RVLM, which we demonstrated by our animal

experiments, might be shared between humans and rats or other animals (Extended Data Fig. 7).

Whereas plasma catecholamine levels were not significantly changed by the VOCR intervention

(Extended Data Fig. 6b), it is possible that urinary epinephrine measures collected over 24 hours in

our rat PHM experiments (Fig. 1f, 3f, and 6d) enhanced our ability to capture the sympathetic nerve

activity under "everyday-life" ambulatory conditions[54].

As the phrase "Exercise is Medicine" indicates, physical exercise is broadly useful to maintain

human health. Many of aerobic exercises, including walking and running, involve impact-

generating bodily actions creating sharp accelerations at the head upon foot contacting with the

ground. Therefore, their beneficial effects as therapeutic/preventative procedures for a variety of

brain function-related diseases and health disorders may rely at least partly on modest changes in

mechanical stress distribution in the brain, which may prompt optimal FSS on intracerebral nervous

cells. While we have recently demonstrated that brain function can be physiologically regulated by

mechanical forces[24], alterations in interstitial fluid movement-derived shear stress may underlie the

pathogenesis of various brain disorders, particularly those related to physical inactivity or aging.

Limitation of study. There are several limitations of this study. We were unable to test the response

of primary neurons, which were prepared from mouse cerebral cortex or hippocampus, to FSS of

relevant magnitudes (0.59−2.64 Pa, see Supplementary Table 1b) because of their easy detachment

from the substrates by FSS. As an alternative of cultured neuronal cells, we tested Neuro2A cells,

which stably adhered to the substrates through FSS of magnitudes up to ~1 Pa[24].

We did not comprehensively analyze the effects of PHM on brain functions, but focused on the

study of RVLM. PHM may modulate AT1R signaling in other brain regions that participate in the

regulation of sympathetic nerve activity, including the anteroventral third ventricle, paraventricular

nucleus of the hypothalamus, and nucleus tractus solitarii[6,9]. Nonetheless, elimination of PHM

effects on BP and urinary norepinephrine excretion by hydrogel introduction in RVLM (Fig. 6b−d)

supports the critical role for RVLM. Because hydrogel may exert yet unknown effects, experiments

of hydrogel introduction may not entirely prove the contribution of interstitial fluid movement. For

example, hydrogel introduction may alter the stiffness and elasticity of extracellular matrix, which

are known to affect the neurological physiology, pathology, and development[55]. Although further

studies are required to address these issues, our findings suggest that mechanical factors can be a

therapeutic target within the blood-brain barrier, the accessibility to which of antihypertensive drugs

appears variable and controversial[56].

Unlike the case of PHM in rats, VOCR of humans generates vertical accelerations at various

body parts in addition to the head. Therefore, we cannot preclude the possibility that the

antihypertensive effect of VOCR is mediated by mechanical regulation of tissues and organs other

than the brain. Furthermore, our clinical studies of protocol 1 and 2 are based on a small number of

subjects (Supplementary Table 2) with a fixed condition (frequency of 2 Hz, peak acceleration of

~1.0 x g, 30 minutes/day, 3 days/week, 12−14 rides), and need further analysis with a much larger

sample size and varying conditions to determine optimal VOCR. Still, application of vertical

mechanical impacts with moderate magnitudes is expected to be highly safe with minimal

possibility of adverse effects, providing a novel therapeutic/preventative strategy for physical

disorders including those that are resistant to conventional treatments such as drug administration.

Notably, our approach utilizing mechanical interventions may bring considerable benefits to those

who cannot receive benefits from exercise because of physical disabilities.

Data availability

All data are included in this article and its supplementary information files. Raw data are available

from the corresponding author upon reasonable request.

Acknowledgements

We thank K. Nakanishi, K. Hamamoto, and N. Kume for their consistent support. This work was in

part supported by Intramural Research Fund from the Japanese Ministry of Health, Labour and

Welfare; Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science

(15H01820, 15H04966, and 18H04088 to Y.S; 17H02127 and 18H03138 to T.O.; 19K06899 to

A.K.); MEXT-Supported Program for the Strategic Research Foundation at Private Universities,

2015−2019 from the Japanese Ministry of Education, Culture, Sports, Science and Technology

(S1511017 to Y.S. and T.O.); Brain Mapping by Integrated Neurotechnologies for Disease Studies

Author contributions

S.M. and N.S. conducted most of the experiments. Y.S. conceived the research, designed the study,

and led the project with help from M.N. and M.S. T.K. provided technical advice for all the

experiments involving measurement of cardiovascular parameters. S.M., N.S., T.K., M.S. and Y.S.

wrote the manuscript. T.M. and A.T. contributed to the design and construction of the machine for

PHM. D.Y. helped in vitro FSS experiments and carried out simulative calculation of in vivo FSS.

T.S. and Y.Y. developed and provided the PEG hydrogel system. A.K. and H.H. prepared and

provided AAV vectors. K.T., T.K., M.A., H.I., and T.O. contributed to the human studies. K.S., T.S.,

S.T. M.S., T.O., and M.N. provided technical, advisory and financial support.

Competing interests

431 The authors declare no competing interest. S.M., T.M., T.O., A.T., and Y.S. joined the application of

432 a patent for the vertically oscillating chair, which has been awarded in Japan (JP6592834) and filed

433 internationally (US16/616,935; EP18806753.2; CN201880033284.0; IN201927048891).

References

1.   Lim, S.S.*, et al.* A comparative risk assessment of burden of disease and injury attributable to
67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the
Global Burden of Disease Study 2010. *Lancet* **380**, 2224-2260 (2012).

2.   Hagberg, J.M., Park, J.J. & Brown, M.D. The role of exercise training in the treatment of
hypertension: an update. *Sports Med* **30**, 193-206 (2000).

3.   Pescatello, L.S., MacDonald, H.V., Lamberti, L. & Johnson, B.T. Exercise for hypertension: a
prescription update integrating existing recommendations with emerging research. *Curr*
Hypertens Rep **17**, 87 (2015).

4.   Carretero, O.A. & Oparil, S. Essential hypertension. Part I: definition and etiology. *Circulation*
101, 329-335 (2000).

5.   Guyton, A.C. Abnormal renal function and autoregulation in essential hypertension.
Hypertension **18**, III49-53 (1991).

6.   Guyenet, P.G. The sympathetic control of blood pressure. *Nat Rev Neurosci* **7**, 335-346 (2006).

7.   Grassi, G., Mark, A. & Esler, M. The sympathetic nervous system alterations in human
hypertension. *Circ Res* **116**, 976-990 (2015).

8.   Malpas, S.C. Sympathetic nervous system overactivity and its role in the development of
cardiovascular disease. *Physiol Rev* **90**, 513-557 (2010).

9.   Dampney, R.A. Functional organization of central pathways regulating the cardiovascular
system. *Physiol Rev* **74**, 323-364 (1994).

10.   Fyhrquist, F. & Saijonmaa, O. Renin-angiotensin system revisited. *J Intern Med* **264**, 224-236
(2008).

11.   Forrester, S.J.*, et al.* Angiotensin II signal transduction: an update on mechanisms of
physiology and pathophysiology. *Physiol Rev* **98**, 1627-1738 (2018).

12.   Ito, S., Komatsu, K., Tsukamoto, K., Kanmatsuse, K. & Sved, A.F. Ventrolateral medulla AT1
receptors support blood pressure in hypertensive rats. *Hypertension* **40**, 552-559 (2002).

13.   Muratani, H., Ferrario, C.M. & Averill, D.B. Ventrolateral medulla in spontaneously
hypertensive rats: role of angiotensin II. *Am J Physiol* **264**, R388-395 (1993).

14.   Kishi, T.*, et al.* Exercise training causes sympathoinhibition through antioxidant effect in the
rostral ventrolateral medulla of hypertensive rats. *Clin Exp Hypertens* **34**, 278-283 (2012).

15.   Nabika, T., Ohara, H., Kato, N. & Isomura, M. The stroke-prone spontaneously hypertensive
rat: still a useful model for post-GWAS genetic studies? *Hypertens Res* **35**, 477-484 (2012).

16.   Lu, X.*, et al.* Effects of local mechanical stimulation on coronary artery endothelial function
and angiotensin II type 1 receptor in pressure or flow-overload. *J Hypertens* **31**, 720-729
(2013).

17. Galie, P.A., Russell, M.W., Westfall, M.V. & Stegemann, J.P. Interstitial fluid flow and cyclic
strain differentially regulate cardiac fibroblast activation via AT1R and TGF-β1. *Exp Cell Res*
318, 75-84 (2012).
18. Zou, Y.*, et al.* Mechanical stress activates angiotensin II type 1 receptor without the
involvement of angiotensin II. *Nat Cell Biol* **6**, 499-506 (2004).
19. Ramkhelawon, B.*, et al.* Shear stress regulates angiotensin type 1 receptor expression in
endothelial cells. *Circ Res* **105**, 869-875 (2009).
20. Li, E.C., Heran, B.S. & Wright, J.M. Angiotensin converting enzyme (ACE) inhibitors versus
angiotensin receptor blockers for primary hypertension. *Cochrane Database Syst Rev*,
CD009096 (2014).
21. Weinbaum, S., Cowin, S.C. & Zeng, Y. A model for the excitation of osteocytes by mechanical
loading-induced bone fluid shear stresses. *J Biomech* **27**, 339-360 (1994).
22. Tatsumi, S.*, et al.* Targeted ablation of osteocytes induces osteoporosis with defective
mechanotransduction. *Cell Metab* **5**, 464-475 (2007).
23. Miyazaki, T.*, et al.* Mechanical regulation of bone homeostasis through p130Cas-mediated
alleviation of NF-κB activity. *Sci Adv* **5**, eaau7802 (2019).
24. Ryu, Y.*, et al.* Mechanical regulation underlies effects of exercise on serotonin-induced
signaling in the prefrontal cortex neurons. *iScience* **23**, 100874 (2020).
25. Minami, N.*, et al.* Effects of exercise and β-blocker on blood pressure and baroreflexes in
spontaneously hypertensive rats. *Am J Hypertens* **16**, 966-972 (2003).
26. Kim, S.E.*, et al.* Treadmill exercise prevents aging-induced failure of memory through an
increase in neurogenesis and suppression of apoptosis in rat hippocampus. *Exp Gerontol* **45**,
357-365 (2010).
27. Bertagnolli, M.*, et al.* Exercise training reduces sympathetic modulation on cardiovascular
system and cardiac oxidative stress in spontaneously hypertensive rats. *Am J Hypertens* **21**,
1188-1193 (2008).
28. Chidsey, C.A., Braunwald, E. & Morrow, A.G. Catecholamine excretion and cardiacstores of
norepinephrine in congestive heart failure. *Am J Med* **39**, 442-451 (1965).
29. Mullen, R.J., Buck, C.R. & Smith, A.M. NeuN, a neuronal specific nuclear protein in
vertebrates. *Development* **116**, 201-211 (1992).
30. Freeman, M.R. Specification and morphogenesis of astrocytes. *Science* **330**, 774-778 (2010).
31. Tamura, K.*, et al.* The physiology and pathophysiology of a novel angiotensin receptor-binding
protein ATRAP/Agtrap. *Curr Pharm Des* **19**, 3043-3048 (2013).
32. Huda, F.*, et al.* Distinct transduction profiles in the CNS via three injection routes of AAV9 and
the application to generation of a neurodegenerative mouse model. *Mol Ther Methods Clin Dev*
1, 14032 (2014).

33. Tang, X., *et al.* "Self-cleaving" 2A peptide from porcine teschovirus-1 mediates cleavage of dual fluorescent proteins in transgenic Eimeria tenella. *Vet Res* **47**, 68 (2016).

34. Tworkoski, E., Glucksberg, M.R. & Johnson, M. The effect of the rate of hydrostatic pressure depressurization on cells in culture. *PLoS One* **13**, e0189890 (2018).

35. Saitou, K., *et al.* Local cyclical compression modulates macrophage function in situ and alleviates immobilization-induced muscle atrophy. *Clin Sci (Lond)* **132**, 2147-2161 (2018).

36. Maneshi, M.M., *et al.* Mechanical stress activates NMDA receptors in the absence of agonists. *Sci Rep* **7**, 39610 (2017).

37. Gong, S., *et al.* A gene expression atlas of the central nervous system based on bacterial artificial chromosomes. *Nature* **425**, 917-925 (2003).

38. Yoshino, D., Sakamoto, N., Takahashi, K., Inoue, E. & Sato, M. Development of Novel Flow Chamber to Study Endothelial Cell Morphology: Effects of shear flow with uniform spatial gradient ondistribution of focal adhesion. *J Biomech Sci Eng* **8**, 233-243 (2013).

39. Goshima, Y., Ohsako, S. & Yamauchi, T. Overexpression of Ca^{2+}/calmodulin-dependent protein kinase II in Neuro2A and NG108-15 neuroblastoma cell lines promotes neurite outgrowth and growth cone motility. *J Neurosci* **13**, 559-567 (1993).

40. Yun, J., *et al.* Neuronal Per Arnt Sim (PAS) domain protein 4 (NPAS4) regulates neurite outgrowth and phosphorylation of synapsin I. *J Biol Chem* **288**, 2655-2664 (2013).

41. Fujiyabu, T., Toni, F., Li, X., Chung, U.I. & Sakai, T. Three cooperative diffusion coefficients describing dynamics of polymer gels. *Chem Commun (Camb)* **54**, 6784-6787 (2018).

42. Jennings, J.R., Muldoon, M.F. & Sved, A.F. Is the brain an early or late component of essential hypertension? *Am J Hypertens* (2020).

43. Joyner, M.J. & Green, D.J. Exercise protects the cardiovascular system: effects beyond traditional risk factors. *J Physiol* **587**, 5551-5558 (2009).

44. Jancovski, N., *et al.* Angiotensin type 1A receptor expression in C1 neurons of the rostral ventrolateral medulla contributes to the development of angiotensin-dependent hypertension. *Exp Physiol* **99**, 1597-1610 (2014).

45. Guo, F., *et al.* Astroglia are a possible cellular substrate of angiotensin(1-7) effects in the rostral ventrolateral medulla. *Cardiovasc Res* **87**, 578-584 (2010).

46. Liddelow, S.A. & Barres, B.A. Reactive astrocytes: production, function, and therapeutic potential. *Immunity* **46**, 957-967 (2017).

47. Maclullich, A.M., *et al.* Higher systolic blood pressure is associated with increased water diffusivity in normal-appearing white matter. *Stroke* **40**, 3869-3871 (2009).

48. Bedussi, B., *et al.* Enhanced interstitial fluid drainage in the hippocampus of spontaneously hypertensive rats. *Sci Rep* **7**, 744 (2017).

49. Hahn, C. & Schwartz, M.A. Mechanotransduction in vascular physiology and atherogenesis. *Nat Rev Mol Cell Biol* **10**, 53-62 (2009).

50. Elton, T.S. & Martin, M.M. Angiotensin II type 1 receptor gene regulation: transcriptional and
posttranscriptional mechanisms. *Hypertension* **49**, 953-961 (2007).

51. Galie, P.A., van Oosten, A., Chen, C.S. & Janmey, P.A. Application of multiple levels of fluid
shear stress to endothelial cells plated on polyacrylamide gels. *Lab Chip* **15**, 1205-1212 (2015).

52. Kadohama, T., Nishimura, K., Hoshino, Y., Sasajima, T. & Sumpio, B.E. Effects of different
types of fluid shear stress on endothelial cell proliferation and survival. *J Cell Physiol* **212**,
244-251 (2007).

53. Iskratsch, T., Wolfenson, H. & Sheetz, M.P. Appreciating force and shape-the rise of
mechanotransduction in cell biology. *Nat Rev Mol Cell Biol* **15**, 825-833 (2014).

54. Hughes, J.W., Watkins, L., Blumenthal, J.A., Kuhn, C. & Sherwood, A. Depression and anxiety
symptoms are related to increased 24-hour urinary norepinephrine excretion among healthy
middle-aged women. *J Psychosom Res* **57**, 353-358 (2004).

55. Li, S.C*., et al.* Tissue elasticity bridges cancer stem cells to the tumor microenvironment
through microRNAs: implications for a "watch-and-wait" approach to cancer. *Curr Stem Cell*
Res Ther **12**, 455-470 (2017).

56. Kishi, T., Hirooka, Y. & Sunagawa, K. Sympathoinhibition caused by orally administered
telmisartan through inhibition of the AT(1) receptor in the rostral ventrolateral medulla of
hypertensive rats. *Hypertens Res* **35**, 940-946 (2012).

57. Huang, N. & Bonn, D. Viscosity of a dense suspension in Couette flow. *J Fluid Mech* **590**, 497-
507 (2007).

58. Sugiyama, S*., et al.* Computational simulation of convection-enhanced drug delivery in the
non-human primate brainstem: a simple model predicting the drug distribution. *Neurol Res* **35**,
773-781 (2013).

59. Yao, W., Shen, Z. & Ding, G. Simulation of interstitial fluid flow in ligaments: comparison
among Stokes, Brinkman and Darcy models. *Int J Biol Sci* **9**, 1050-1056 (2013).

60. Tarbell, J.M. & Shi, Z.-D. Effect of the glycocalyx layer on transmission of interstitial flow
shear stress to embedded cells. *Biomech Model Mechanobiol* **12**, 111-121 (2013).

61. Sakai, K*., et al.* Overexpression of eNOS in NTS causes hypotension and bradycardia in vivo.
Hypertension **36**, 1023-1028 (2000).

62. Kishi, T*., et al.* Overexpression of eNOS in the RVLM causes hypotension and bradycardia via
GABA release. *Hypertension* **38**, 896-901 (2001).

63. Paxinos, G. & Watson, C. *The rat brain in stereotaxic coordinates* (Academic Press, 1998).

64. Hirooka, Y., Polson, J.W. & Dampney, R.A. Pressor and sympathoexcitatory effects of nitric
oxide in the rostral ventrolateral medulla. *J Hypertens* **14**, 1317-1324 (1996).

65. Shinohara, Y*., et al.* Effects of neutralizing antibody production on AAV-PHP.B-mediated
transduction of the mouse central nervous system. *Mol Neurobiol* **56**, 4203-4214 (2019).

66. Shinohara, Y., Ohtani, T., Konno, A. & Hirai, H. Viral vector-based evaluation of regulatory
regions in the neuron-specific enolase (NSE) promoter in mouse cerebellum in vivo.
Cerebellum **16**, 913-922 (2017).

67. Konno, A*., et al.* Mutant ataxin-3 with an abnormally expanded polyglutamine chain disrupts
dendritic development and metabotropic glutamate receptor signaling in mouse cerebellar
Purkinje cells. *Cerebellum* **13**, 29-41 (2014).

68. Matsuzaki, Y*., et al.* Neurotropic Properties of AAV-PHP.B are shared among diverse inbred
strains of mice. *Mol Ther* **27**, 700-704 (2019).

69. Sakitani, N*., et al.* Application of consistent massage-kike perturbations on mouse calves and
monitoring the resulting intramuscular pressure changes. *J Vis Exp* e59475 (2019).

70. Liu, C., De Luca, A., Rosso, A. & Talon, L. Darcy's law for yield stress fluids. *Phys Rev Lett*
122, 245502 (2019).

71. Schildge, S., Bohrer, C., Beck, K. & Schachtrup, C. Isolation and culture of mouse cortical
astrocytes. *J Vis Exp* e50079 (2013).

72. Umemura, S*., et al.* The Japanese Society of Hypertension guidelines for the management of
hypertension (JSH 2019). *Hypertens Res* **42**, 1235-1481 (2019).

Figures and figure legends

Fig. 1 | Application of cyclical mechanical impact to the head by passive motion lowers the BP

in SHRSPs, and AT1R expression in their RVLM astrocytes. a, Schematic representation of

experimental protocol for analysis of the effects of PHM on BP in rats. **b,c,** Time courses **(b)** and

values on Day 29 **(c)** of MAP of WKY rats and SHRSPs, subjected to either daily PHM or

anesthesia only (**b**: $P = 0.0814$ for Day15, $P = 0.0052$ for Day 22, and $P = 0.0046$ for Day 29. **c**: P

$= 0.9739$ for column 1 vs. 2 and $P = 0.0046$ for column 3 vs. 4. $n = 7$ rats for each group of WKY

and $n = 8$ rats for each group of SHRSP). **d,** HR values on Day 29 ($P = 0.9650$ for column 1 vs. 2

and $P = 0.2362$ for column 3 vs. 4. $n = 7$ rats for each group of WKY and $n = 8$ rats for each group

of SHRSP). **e,** Relative heart weight (heart weight / whole body weight) measured on Day 30 [$P =$

0.9866 for column 1 vs. 2 and $P = 0.0152$ for column 3 vs. 4. $n = 10$ rats for WKY, PHM (−); $n = 13$

rats for WKY, PHM (+); $n = 10$ rats for SHRSP, PHM (−); $n = 14$ rats for SHRSP, PHM (+)]. **f,**

Twenty-four-hour (Day 29 to Day 30) urinary norepinephrine excretion [$P = 0.9854$ for column 1

vs. 2 and $P = 0.0085$ for column 3 vs. 4. $n = 8$ rats for each group of WKY; $n = 16$ rats for SHRSP,

PHM (−); $n = 13$ rats for SHRSP, PHM (+)]. **g,h,** Micrographic images of anti-NeuN (blue), anti-

GFAP (green) and anti-AT1R (red) immunostaining of the RVLM of WKY rats **(g)** and SHRSPs

(h), either left sedentary (top) or subjected to PHM (bottom) under anesthesia (30 minutes/day, 28

days). Higher magnification images (center and right) refer to the areas indicated by dotted

rectangles in low magnification images (left). Arrows point to anti-AT1R immunosignals that

overlap with anti-GFAP, but not anti-NeuN, immunosignals in merged images. Scale bars, 50 μm.

Images are representative of three rats. **i,j,** Quantification of AT1R-positive neurons **(i)** and

astrocytes **(j)** in the RVLM of WKY rats and SHRSPs, either left sedentary or subjected to PHM.

Fifty NeuN-positive (NeuN$^+$) cells and one hundred GFAP-positive (GFAP$^+$) cells were analyzed

for each rat (**i**: $P = 0.7803$. **j**: $P = 0.9455$ for column 1 vs. 2, $P = 0.0004$ for column 1 vs. 3, and $P =$

0.0002 for column 3 vs. 4. $n = 3$ rats for each group). Data are presented as mean $\pm$ s.e.m. $^*P < 0.05$;

$^{**}P < 0.01$; $^{***}P < 0.001$; NS, not significant; one-way ANOVA with Tukey's post hoc multiple

comparisons test.

Extended Data Fig. 1 | Daily anesthesia alone does not affect BP, and treadmill running

decreases AT1R expression in the RVLM astrocytes in SHRSPs. a, Time courses of MAP of

SHRSPs either routinely kept in cages (no anesthesia) or subjected to daily anesthesia on the

platform of PHM machine without turning on its switch [PHM (−)] [$n = 6$ rats for no daily

anesthesia and $n = 8$ rats for PHM (−)]. **b,** Micrographic images anti-NeuN (blue), anti-GFAP

(green) and anti-AT1R (red) immunostaining of the RVLM of SHRSPs, either placed in the static

treadmill machine or subjected to treadmill running at the velocity of 20 m/minute (30 minutes/day,

28 days). Arrows point to anti-AT1R immunosignals that overlap with anti-GFAP, but not anti-

NeuN, immunosignals in merged images. Scale bar, 50 μm. Images are representative of three or

four rats. **c,d,** Quantification of AT1R-positive neurons **(c)** and astrocytes **(d)** in the RVLM of

SHRSPs with or without 4-week treadmill running. Fifty NeuN$^+$ cells and one hundred GFAP$^+$ cells

were analyzed for each rat [**c**: $P = 0.7056$. **d**: $P = 0.0048$. $n = 3$ rats for treadmill (−) and $n = 4$ rats

for treadmill (+)]. Data are presented as mean ± s.e.m. **$P < 0.01$; NS, not significant, unpaired

two-tailed Student's t-test.

Fig. 2 | PHM alleviates the sensitivity of RVLM in SHRSPs to Ang II or valsartan. a,

Schematic representation of experimental protocol for analysis of the effects of PHM on the

sensitivity to Ang II or valsartan injected into unilateral RVLM. Ang II (100 pmol) was injected into

the unilateral RVLM of WKY rats and SHRSPs, either left sedentary (daily anesthesia) or subjected

to PHM (30 minutes/day, 28 days), with their BP monitored under urethane anesthesia. Injection of

valsartan (100 pmol) into the RVLM was conducted at least 2 hours after the injection of Ang II. **b,**

Representative trajectories of BP (top in each panel) and MAP (bottom in each panel). Arrows point

to the time of the initiation of RVLM injection of Ang II. **c,** Quantification of MAP change caused

by Ang II injection [$P = 0.9876$ for column 1 vs. 2 and $P = 0.0003$ for column 3 vs. 4. $n = 5$ rats for

each group of WKY; $n = 7$ rats for SHRSP, PHM (−); $n = 8$ rats for SHRSP, PHM (+)]. **d,e,** Effects

of RVLM injection of valsartan (100 pmol) examined as in **(b,c)** [**e**: $P = 0.9953$ for column 1 vs. 2

and $P = 0.0099$ for column 3 vs. 4. $n = 3$ rats for WKY, PHM (−); $n = 5$ rats for WKY, PHM (+); n

$= 4$ rats for each group of SHRSP]. Data are presented as mean ± s.e.m. **$P < 0.01$; ***$P < 0.001$;

NS, not significant; one-way ANOVA with Tukey's post hoc multiple comparisons test.

Fig. 3 | AAV-mediated expression of AGTRAP in RVLM astrocytes, but not neurons, lowers

the BP in SHRSPs. a, Schematic representation of experimental protocol for analysis of the effects

of AAV-mediated transduction of RVLM astrocytes or neurons with AGTRAP gene. ITR; inverted

terminal repeat. **b,c,** Astrocyte- **(b)** and neuron- **(c)** specific transgene expression by RVLM

injection of AAV9 vectors. Micrographic images of GFP (green) and anti-GFAP **(b)** or anti-NeuN

(c) immunostaining (blue) of RVLM of SHRSPs 15 days after the injection of AAV9 vectors

indicated at the top of each panel. Scale bars, 50 μm. Images are representative of three rats. **d−f,**

Time courses **(d)** and values on Day 15 **(e)** of MAP (**d**: $P = 0.0229$ for Day 14. **e**: $P = 0.0229$ for

column 1 vs. 2 and $P = 0.6864$ for column 3 vs. 4. $n = 6$ rats for GFAP-control; $n = 7$ rats for

GFAP-AGTRAP; $n = 7$ rats for NSE-control; $n = 6$ rats for NSE-AGTRAP), and 24-hour urinary

norepinephrine excretion **(f)** ($P = 0.0497$ for column 1 vs. 2 and $P = 0.7455$ for column 3 vs. 4. $n =$

4 rats for each group) of SHRSPs subjected to RVLM injection of AAV9 vectors. Data are

presented as mean ± s.e.m. *$P < 0.05$; NS, not significant, one-way ANOVA with Tukey's post hoc

multiple comparisons test.

Extended Data Fig. 2

Extended Data Fig. 2 | AAV-mediated transduction of RVLM astrocytes or neurons in

SHRSPs. a, Schematic representation of injection of AAV9 vectors to RVLM. Micrographic image

is representative of three rats analyzed in Fig. 3b (15 days after the RVLM injection of GFAP-

AGTRAP vector). GFP-derived fluorescence indicates cells expressing the transgene. Scale bar, 1

mm. **b,c,** Efficiency and specificity of astrocyte-specific expression of the transgene. **(b)**

Micrographic images of GFP (green) and anti-NeuN immunostaining (blue) of RVLM of SHRSPs

analyzed in Fig. 3b. Scale bar, 50 μm. Images are representative of three rats. **(c)** Quantification of

the efficiency and specificity of transgene expression. The relative populations (%) of GFP- GFAP-

double positive (GFP$^+$ GFAP$^+$) or GFP- NeuN-double positive (GFP$^+$ NeuN$^+$) cells were calculated

by referring their numbers to those of GFP$^+$, GFAP$^+$ or NeuN$^+$ cells ($P < 0.0001$ for column 1 vs. 2

and $P < 0.0001$ for column 3 vs. 4. $n = 3$ rats for each group). **d,e,** Efficiency and specificity of

neuron-specific expression of the transgene. **(d)** Micrographic images of GFP (green) and anti-

GFAP immunostaining (blue) of RVLM of SHRSPs analyzed in Fig. 3c. Scale bar, 50 μm. Images

are representative of three rats. **(e)** The efficiency and specificity quantified as in **(c)** ($P < 0.0001$ for

column 1 vs. 2 and $P < 0.0001$ for column 3 vs. 4.). **f,** BP in SHRSPs injected with control vectors.

BP was measured and MAP was quantified as in Fig. 1b ($P = 0.9803$ for Week 12, $P = 0.5914$ for

Week 13, and $P = 0.4693$ for Week 14. $n = 6$ rats for no AAV injection; $n = 6$ rats for GFAP-control;

$n = 7$ rats for NSE-control). The data for blue line are identical with those demonstrated with blue

line in Extended Data Fig. 1a. Data are presented as mean ± s.e.m. ***$P < 0.001$; NS, not

significant; unpaired two-tailed Student's *t*-test **(c,e)** or one-way ANOVA with Tukey's post hoc

multiple comparisons test **(f)**.

Fig. 4

Fig. 4 | **PHM generates pressure waves of low amplitude, but facilitates interstitial fluid**

movement (flow) in rat RVLM. a, Schematic representation of pressure measurement in rat

RVLM. **b,** Representative pressure waves recorded in rat RVLM during sedentary condition and

PHM. Arrows point to the time of transition from inhalation to exhalation detected by simultaneous

respiration monitoring. Scale bar, 1 s. Images are representative of three independent experiments

with similar results. **c,** Respiration-unsynchronized pressure changes. Respiration-synchronized

pressure waves indicated by dotted rectangles in **(b)** are presented with high magnification. Right-

angled scale bar, 1 s / 1 mm Hg. Note that 2-Hz pressure waves indicated by a two-headed dotted

line arrow were specifically generated during PHM. **d,** Magnitude of PHM-specific and -unspecific

pressure changes unsynchronized with respiration. Peak to peak magnitudes indicated by two-

headed arrows in **(c)** were quantified ($P = 0.0089$. $n = 4$ rats for each group, 10 segments analyzed

for each rat). **e,** Schematic representation of experimental protocol for μCT analysis of Isovist

injected in rat RVLM. **f,** Definition of x-(left-right), y-(rostral-caudal), and z-(dorsal-ventral) axes

used in this study. **g,** Representative Isovist spreading presented on X-ray images. Isovist clusters

are indicated by green. Images are representative of three rats. A dotted line arrow indicates the

main direction of spreading in this sample. **h,** Quantification of Isovist spreading along each axis

(left chart: $P = 0.6666$. middle chart: $P = 0.0218$. right chart: $P = 0.0244$. $n = 3$ rats for each group).

Data are presented as mean ± s.e.m. $^*P < 0.05$; $^{**}P < 0.01$; NS, not significant, unpaired two-tailed

Student's t-test.

a

Property	Value
Pressure changes (ΔP; mmHg)	1.19
Viscosity (μ; mPa•s)	1–20 [#]
Spread distance along x-axis per each PHM cycle (Δx; μm)	–0.002
Spread distance along y-axis per each PHM cycle (Δy; μm)	0.039
Spread distance along z-axis per each PHM cycle (Δz; μm)	0.055
Velocity of interstitial fluid flow along x-axis ($u_{\infty,x}$; μm/s)	–0.004
Velocity of interstitial fluid flow along y-axis ($u_{\infty,y}$; μm/s)	0.079
Velocity of interstitial fluid flow along z-axis ($u_{\infty,z}$; μm/s)	0.111

b

FSS (τ_x) along x-axis at the cell surface:

$$\tau_x = \frac{\mu u_{\infty,x}}{\sqrt{K_{p,x}}}$$

$$K_{p,x} = \frac{\mu u_{\infty,x}\Delta x}{\Delta P}$$

, where $K_{p,x}$ is the Darcy permeability of brain tissue along x-axis.

The shear stresses along y- and z-axes can be calculated in a similar manner.

When the values listed in **a** are introduced in these equations, the magnitude of FSS is estimated as 0.59–2.64 Pa.

Supplementary Table 1 | Simulative calculation of the magnitude of PHM-generated FSS on

rat RVLM cells. a, Values referenced for simulative calculation of the magnitude of FSS that PHM

generated in rat RVLM. All referenced values except viscosity (marked with ♯) were drawn from

analyses with IBP measurement and Isovist-enhanced μCT scanning (Fig. 4d,h). The property of

interstitial fluid viscosity was referenced from previous studies[57-59]. **b,** Calculation of the magnitude

of FSS generated by PHM. FSS (τ) at the cell surface can be calculated as reported previously[60].

Fig. 5

Fig. 5 | FSS on cultured astrocytes decreases their AT1R expression and Ang II-binding

potential in vitro. a−c, AT1R expression in cultured astrocytes with or without exposure to FSS.

Astrocytes prepared from astrocyte-GFP mice, either left unexposed or exposed to pulsatile FSS

(average 0.7 Pa, 0.5 Hz, 30 minutes) were analyzed 6 and 24 hours after the cessation of 30-minute

FSS application. **(a)** AT1R mRNA expression levels normalized against GAPDH expression and

scaled as the mean value from control samples (cells left unexposed to FSS) set as 1 ($P = 0.0050$ for

column 1 vs. 2 and $P = 0.0011$ for column 1 vs. 3. $n = 6$). **(b)** Microscopic images of anti-AT1R

(red) and anti-GFP (green) immunostaining. Images are representative of three or four independent

experiments with similar results. Scale bar, 50 μm. **(c)** Relative population of AT1R- GFP-double

positive (AT1R$^+$ GFP$^+$) cells were quantified as a ratio to total GFP-positive (GFP$^+$) cells in each

sample [$P = 0.0021$ for column 1 vs. 2 and $P = 0.0002$ for column 1 vs. 3. >100 GFP-positive cells

were analyzed in each sample; $n = 4$ for FSS (−), $n = 3$ for 6 and 24 hours after FSS]. **d,e,** Effect of

FSS on astrocytes' Ang II-binding potential. Cultured astrocytes were either left unexposed or

exposed to FSS as in **(a−c)**. Six and twenty-four hours after the cessation of 30-minute FSS

application, cells were subjected to fluorescent Ang II binding assay. **(d)** Microscopic images

representative of three independent experiments with similar results. Scale bar, 50 μm. **(e)** GFP-

positive cells with punctate red fluorescence (TAMRA-Ang II-bound astrocytes) were quantified as

a ratio (%) to total GFP-positive cells in each sample ($P < 0.0001$ for column 1 vs. 2 and $P = 0.0001$

for column 1 vs. 3. 100 GFP-positive cells were analyzed in each sample; $n = 3$ for each group).

Data are presented as mean ± s.e.m. **$P < 0.01$, ***$P < 0.001$, one-way ANOVA with Tukey's post

hoc multiple comparisons test.

Extended Data Fig. 3 | Preparation of astrocyte-enriched primary culture, and lack of decreasing effect of FSS on AT1R expression in cultured neuronal cells. a, Representative micrographic image of anti-GFP immunostaining (green) and DAPI staining (blue) of astrocyte-enriched culture prepared from astrocyte-GFP mice. Scale bar, 50 μm. Note that most of the cells are GFP-positive. **b−d,** Effects of FSS on AT1R expression in Neuro2A cells analyzed as in Fig. 5a−c. **(b)** AT1R mRNA expression levels ($P = 0.2834$. $n = 4$). **(c)** Micrographic images representative of three independent experiments with similar results. Scale bar, 50 μm. **(d)** Relative population of AT1R- TUJ-1-double positive (AT1R$^+$ TUJ-1$^+$) cells quantified as a ratio to total TUJ-

749   1-positive (TUJ-1$^+$) cells in each sample ($P = 0.2308$. >100 TUJ-1-positive cells were analyzed in

750   each sample; $n = 3$). Data are presented as mean $\pm$ s.e.m. NS, not significant, one-way ANOVA with

751   Tukey's post hoc multiple comparisons test **(b)** or unpaired two-tailed Student's t-test **(d)**.

Fig. 6 | Hydrogel introduction eliminates decreasing effects of PHM on BP and AT1R

expression in the RVLM astrocytes in SHRSPs. a, Schematic representation of experimental

protocol for analysis of the effects of PHM with and without PEG hydrogel introduction in bilateral RVLMs of SHRSPs. PHM was applied daily for consecutive 28 days. **b−d,** Time courses **(b)** and values on Day 29 **(c)** of MAP, and 24-hour urinary norepinephrine excretion **(d)** of SHRSPs, subjected to various combinations of daily PHM application and hydrogel introduction in bilateral RVLMs. Note the absence of significant difference in BP **(b,c)** and urinary norepinephrine excretion **(d)** of SHRSPs with hydrogel introduced RVLMs [Gel (+)] between with and without PHM [**b**: $P = 0.0372$ for Day 29. **c**: $P = 0.0387$ for column 1 vs. 2, $P = 0.0372$ for column 1 vs. 3, and $P = 0.9959$ for column 2 vs. 3. $n = 7$ rats for Gel (−), PHM (+) and for Gel (+), PHM (+); $n = 6$ rats for Gel (+), PHM (−); **d**: $P = 0.0247$ for column 1 vs. 2, $P = 0.0307$ for column 1 vs. 3, and $P = 0.9920$ for column 2 vs. 3. $n = 5$ rats for each group]. **e**, Micrographic images of anti-GFAP (green), anti-AT1R (red), and anti-NeuN (blue) immunostaining of the RVLM in SHRSPs subjected to various combinations of hydrogel introduction in bilateral RVLMs and 4-week PHM application. Arrows point to anti-AT1R immunosignals that overlap with anti-GFAP, but not anti-NeuN, immunosignals in merged images. Scale bar, 50 μm. Images are representative of four rats. **f,g,** Quantification of AT1R-positive astrocytes **(f)** and neurons **(g)** in the RVLM. Note the absence of significant difference in the ratio of AT1R-positive astrocytes of SHRSPs with hydrogel introduced RVLMs [Gel (+)] between with and without PHM (**f**, columns 2 and 3). Fifty NeuN$^+$ cells and one hundred GFAP$^+$ cells were analyzed for each rat (**f**: $P < 0.0001$ for column 1 vs. 2, $P < 0.0001$ for column 1

773   vs. 3, and $P = 0.1597$ for column 2 vs. 3. **g**: $P = 0.5182$. $n = 4$ rats for each group). Data are

774   presented as mean $\pm$ s.e.m. *$P < 0.05$; ***$P < 0.001$; NS, not significant, one-way ANOVA with

775   Tukey's post hoc multiple comparisons test.

Extended Data Fig. 4

Extended Data Fig. 4 | Hydrogel introduction in the RVLM does not affect the depressor

response to valsartan or survival/apoptosis of RVLM astrocytes and neurons in SHRSPs. a,

Introduction of PEG hydrogel in rat RVLM. Twenty-four hours after the injection of control

ungelatable fluorescent PEG solution (left) or 1 week after the injection of pre-gel fluorescent PEG

solution (right), brainstem samples were prepared. Coronal-section images representative of three

rats with similar results are shown. Scale bar, 1 mm. **b,** Schematic representation of experimental

protocol. One week after the injection of pre-gel PEG solution or its ungelatable control, depressor

response was analyzed as in Fig. 2. **c,d,** Representative trajectories **(c)** and quantification **(d)** of BP

descent upon valsartan injection to the RVLM of SHRSPs with or without hydrogel introduction (**d**:

$P = 0.2342$. $n = 3$ rats for each group). **e−j,** Effects of hydrogel introduction on survival/apoptosis

of RVLM cells in SHRSPs. Rat RVLM sections prepared in the experiments for Fig. 6e−g [PHM

(+), Gel (−) and PHM (+), Gel (−)] were subjected to TUNEL assay **(e,f)**, or combinations of anti-

GFAP, anti-NeuN and anti-cleaved caspase-3 immunostaining **(g−j)**. DAPI-positive nuclei **(e)**,

GFAP- **(g)** or NeuN- **(i)** positive cells were counted, and the relative populations of cells doubly

positive for indicated combinations of TUNEL **(f)** and cleaved caspase-3 **(h,j)** were quantified. Each

value represents an average from five images of 1 x 1-mm area analyzed for each rat (**e**: $P = 0.6518$.

f: $P = 0.6943$. **g**: $P = 0.7938$. **h**: $P = 0.8722$. **i**: $P = 0.1679$. **j**: $P = 0.7247$. $n = 3$ rats for each group).

Data are presented as mean ± s.e.m. NS, not significant; unpaired two-tailed Student's *t*-test.

Fig. 7 | VOCR has an antihypertensive effect on hypertensive adult humans. a, BP and HR

"value of the day"s immediately before and after 4-week VOCR in the study of protocol 1 (SBP: P

$= 0.0018$. DBP: $P = 0.1509$. MAP: $P = 0.0459$. HR: $P = 0.3900$. $n = 5$). **b,** BP and HR "value of the

week"s immediately before and after 4.5-week VOCR in the study of protocol 2 (SBP: $P < 0.0001$.

DBP: $P = 0.0031$. MAP: $P = 0.0009$. HR: $P = 0.9817$. $n = 15$). **c,** Subject number-corresponding

trajectories and statistical analysis of BP and HR in the study of protocol 2. Gray-shaded rectangles

indicate the VOCR intervention periods (4.5 weeks). The colors and symbols of individual lines

correspond to those of individual subject numbers listed on the right, excluding subject 9 (see

Supplementary Table 2). Each "value of the week" was statistically compared with that of the week

immediately before the initiation of VOCR intervention (SBP: $P = 0.0293$ for Pre vs. Week 1, $P =$

0.00228 for Pre vs. Week 2, $P = 0.0013$ for Pre vs. Week 3, $P = 0.0035$ for Pre vs. Week 4, $P =$

0.0002 for Pre vs. Week 5, $P = 0.0078$ for Pre vs. Week 6, $P = 0.0035$ for Pre vs. Week 7, $P =$

0.0075 for Pre vs. Week 8, $P = 0.2132$ for Pre vs. Week 9, $P = 0.1314$ for Pre vs. Week 10, $P =$

0.0973 for Pre vs. Week 11, and $P = 0.3993$ for Pre vs. Week 12. DBP: $P = 0.3022$ for Pre vs. Week

1, $P = 0.0436$ for Pre vs. Week 2, $P = 0.0010$ for Pre vs. Week 3, $P = 0.0100$ for Pre vs. Week 4, $P =$

0.0006 for Pre vs. Week 5, $P = 0.0599$ for Pre vs. Week 6, $P = 0.0488$ for Pre vs. Week 7, $P =$

0.0096 for Pre vs. Week 8, $P = 0.9346$ for Pre vs. Week 9, $P = 0.7850$ for Pre vs. Week 10, $P =$

0.0769 for Pre vs. Week 11, and $P = 0.3137$ for Pre vs. Week 12. MAP: $P = 0.1075$ for Pre vs. Week

1, $P = 0.0132$ for Pre vs. Week 2, $P = 0.0008$ for Pre vs. Week 3, $P = 0.0063$ for Pre vs. Week 4, $P =$

0.0003 for Pre vs. Week 5, $P = 0.0251$ for Pre vs. Week 6, $P = 0.0136$ for Pre vs. Week 7, $P =$

0.0071 for Pre vs. Week 8, $P = 0.6795$ for Pre vs. Week 9, $P = 0.4295$ for Pre vs. Week 10, $P =$

0.0704 for Pre vs. Week 11, and $P = 0.3662$ for Pre vs. Week 12. HR: $P = 0.6287$ for Pre vs. Week

1, $P = 0.7840$ for Pre vs. Week 2, $P = 0.1573$ for Pre vs. Week 3, $P = 0.5380$ for Pre vs. Week 4, $P =$

0.7331 for Pre vs. Week 5, $P = 0.6995$ for Pre vs. Week 6, $P = 0.9110$ for Pre vs. Week 7, $P =$

0.9875 for Pre vs. Week 8, $P = 0.5866$ for Pre vs. Week 9, $P = 0.9566$ for Pre vs. Week 10, $P =$

0.6487 for Pre vs. Week 11, and $P = 0.9905$ for Pre vs. Week 12. $n = 15$ for Pre and Weeks 1 to 5, n

$= 13$ for Week 6, $n = 12$ for Week 7, $n = 11$ for Weeks 8 and 9, $n = 9$ for Week 10, $n = 7$ for Week

11, $n = 3$ for Week 12). *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$; NS, not significant, paired two-tailed

Student's t-test.

Extended Data Fig. 5

a

Light jogging
(7 km/hour)

VOCR

0.5 x *g*

0.5 s

b

c

Protocol 1

Informed consent with participants recruited via advertisement and their eligibility check

≥2 weeks

Measure home BP and HR ≥2 days per week (not recorded) ---▶ Two subjects were excluded because of lowered BP (SBP <130 mm Hg)

Measure and record home BP and HR ▶ BP and HR before intervention

Same day

First ride on vertically oscillating chair

4 weeks

Ten rides on vertically oscillating chair (3 days per week)

Last (twelfth) ride on vertically oscillating chair

Next day

Measure and record home BP and HR ▶ BP and HR after intervention

End of study

d

Protocol 2

Informed consent with participants recruited via advertisement and their eligibility check

≥2 weeks

Measure and record home BP and HR ≥3 days per week ---▶ Two subjects were excluded because of lowered BP (SBP <130 mm Hg)

Blood sampling ------▶ Blood data before intervention

≤1 week

First ride on vertically oscillating chair

4.5 weeks

Twelve rides on vertically oscillating chair (3 days per week)
Measure and record home BP and HR (≥3 days per week)

Last (fourteenth) ride on vertically oscillating chair

Next day

Blood sampling ------▶ Blood data after intervention

9 days to 8 weeks

Measure and record home BP and HR ≥3 days per week

End of study

Extended Data Fig. 5 | Vertically oscillating chair that reproduces mechanical impacts on the

head during light jogging, and protocols of clinical study to examine antihypertensive effect of

VOCR on hypertensive adult humans. a, Vertical accelerations generated at adult human head

during light jogging on a treadmill machine (velocity: 7 km/hour) and VOCR (frequency: 2 Hz).

The VOCR system was adjusted to produce ~1.0 x g vertical acceleration peaks. Right-angled scale

bar, 0.5 x g / 0.5 s. Images are representative of three independent experiments with similar results.

b, Photograph of the chair. **c,d,** Protocol 1 **(c)** and 2 **(d)**.

Extended Data Fig. 6

Extended Data Fig. 6 | Trajectories of BP and HR, and blood test values before and after the

intervention period in the study of protocol 2. a, Trajectories of "value of the week"s of BP and

HR. Gray-shaded rectangles indicate the VOCR intervention periods. The colors and symbols of

individual lines correspond to those of individual subjects (Fig. 7c). The black line with inverted

triangles indicates subject 9. **b,** Blood test values before and after the intervention period.

Significant change was not observed in any of tested parameters. NS, not significant; paired two-

tailed Student's *t*-test. (Aldosterone: $P = 0.6265$. Renin activity: $P = 0.3794$. Epinephrine: $P =$

0.5103. Norepinephrine: $P = 0.2653$. Dopamine: $P > 0.9999$. CRP: $P = 0.4412$.). A participant

(subject 18) showed a large increase in plasma renin activity after VOCR. We advised him to

consult his primary care physician, who ruled out the disqualifying conditions for this study (e.g.,

severe renal disease; see Methods) based on comprehensive evaluation. Therefore, we did not

exclude subject 18 from our statistical analysis of BP and HR.

Supplementary Table 2

Subject #	Sex / Age (years) / Body weight (kg) / Height (cm) / BMI	Period since diagnosis or self-recognition as hypertension / Health problems and diseases other than hypertension	Smoking / Alcohol (if yes, how often)	Current medication (dose per day)	Habitual exercise (times per week)	SBP/DBP (mm Hg) and HR (bpm) just before (left) and after (right) VOCR period (month of first - last bout of VOCR)
1	Male / 60 / 73 / 165.2 / 27.2	13 years / Hyperuricemia	No / Almost every day	Azilsartan · Amlodin (20 mg · 5 mg) Febuxostat (10 mg) Bisoprolol fumarate (5 mg)	Walking 90 minutes (3 or 4 times)	148/85 62 — 139/78 66 February - March
2	Female / 53 / 50 / 159 / 19.8	14 years / None	No / Almost every day	None	Sit-ups 2 x 30 times (every day)	135/90 70 — 127/81 69 March
3	Male / 37 / 113 / 186 / 32.7	19 years / None	No / Occasionally	None	Judo (once)	135/86 68 — 129/91 54 March
4	Female / 60 / 73 / 156 / 30	6 years / None	No / Occasionally	None	None	141/87 79 — 132/80 80 April - May
5	Female / 52 / 50 / 148 / 22.8	1 year / None	No / Occasionally	Azilsartan (20 mg) Amlodin (2.5 mg)	None	144/84 74 — 140/80 69 April - May
6	Female / 57 / 68 / 164 / 25.3	10 years / None	No / Occasionally	None	None	128/70 77 — 109/70 76 November - December
7	Male / 61 / 67 / 168 / 23.7	Uncertain (<1 year) / Diabetes mellitus Hyperlipidemia	No / No	Metformin hydrochloride (1000 mg) Rosuvastatin calcium (5 mg) Sitagliptin phosphate hydrate (50 mg)	Walking 120 minutes (once)	146/100 92 — 128/86 84 November - December
8	Male / 46 / 72 / 178 / 22.7	3 months / Allergic rhinitis	Yes / Occasionally	Montelukast sodium (10 mg) Ebastine (10 mg)	Walking 60 minutes (once) Stretching 30 minutes (once)	135/95 61 — 118/78 59 November - December
9	Male / 55 / 87 / 173 / 29.1	7 months / Diabetes mellitus	No / No	None	Karate (2 or 3 times)	156/89 74 — 145/94 67 November - December
10	Male / 70 / 62 / 166 / 22.5	6 months / None	No / Occasionally	None	Walking 40 minutes (4 times)	140/92 81 — 121/79 89 November - December
11	Female / 60 / 45 / 157 / 18.3	Uncertain (>1year) / Breast cancer (post-surgery)	No / Almost every day	Tamoxifen citrate (20 mg)	None	155/92 74 — 136/81 67 November - December
12	Male / 68 / 72 / 171 / 24.6	Uncertain (<1 year) / None	No / No	None	None	133/86 70 — 129/88 70 January - February
13	Female / 56 / 58 / 160 / 22.7	7 years / None	No / No	Telmisartan · Amlodipine (40 mg · 5 mg)	None	137/95 65 — 133/93 71 January - February
14	Male / 49 / 69 / 168 / 24.5	Uncertain (<1 year) / Diabetes mellitus	No / No	Sitagliptin phosphate hydrate (50 mg) Acetazolamide (250 mg)	None	140/90 95 — 124/77 83 January - February

Subject #	Sex / Age (years) / Body weight (kg) / Height (cm) / BMI	Period since diagnosis or self-recognition as hypertension / Health problems and diseases other than hypertension	Smoking / Alcohol (if yes, how often)	Current medication (dose per day)	Habitual exercise (times per week)	SBP/DBP (mm Hg) and HR (bpm) just before (left) and after (right) VOCR period (month of first - last bout of VOCR)
15	Male / 43 / 69 / 165 / 25.3	Uncertain (<1 year) / None	No / Occasionally	None	None	140/87 76 — 124/75 62 · August - September
16	Female / 48 / 45.3 / 157.2 / 18.3	15 years / None	No / Occasionally	None	Walking 60 minutes (6 times)	157/101 72 — 140/92 96 · October - November
17	Male / 65 / 60.1 / 167.1 / 21.5	8 years / None	Yes / Almost every day	None	None	140/93 59 — 134/89 56 · October
18	Male / 55 / 93.4 / 168.1 / 33.1	8 years / Reflex esophagitis Hyperlipidemia	No / Almost every day	Azilsartan (20 mg) Lansoprazole (15 mg) Atorvastatin calcium hydrate (10 mg) Benidipine hydrochloride (4 mg)	None	172/123 69 — 115/78 88 · November - December
19	Male / 41 / 92.3 / 183.2 / 27.5	2 years / None	No / Almost every day	None	Walking 60 minutes (3 times)	130/83 64 — 127/89 56 · January - February
20	Female / 65 / 48.7 / 158.6 / 19.4	1 year / Hyperlipidemia Lumbar spinal canal stenosis (post-surgery)	No / Almost every day	Amlodipine besilate (5 mg) Atorvastatin calcium hydrate (10 mg) Limaprost alfadex (15 µg) Loxoprofen sodium hydrate (120 mg) Rebamipide (200 mg)	Swimming 40 minutes (5 times)	136/87 70 — 118/75 67 · January - February
21	Male / 32 / 65.1 / 168.9 / 22.8	2 years / None	No / Almost every day	None	None	163/109 71 — 126/86 80 · January - February

Supplementary Table 2 | Information of subjects who participated in the human studies.

Subjects 1−5 were participants in the study of protocol 1, and subjects 6−21 participated in the

study of protocol 2. Subject 9 was excluded from our statistical analysis because of high serum CRP

value before VOCR (2.85 mg/dL), which made it difficult to rule out acute infection, a possible

disqualifier, at the time of initiation of VOCR, albeit the lack of specific complaint or local

symptom related to acute physical problem(s). His serum CRP after the VOCR period was within

the normal range (0.12 mg/dL).

Extended Data Fig. 7 | Schematic representation of the antihypertensive effects observed in

PHM of hypertensive rats and VOCR of hypertensive humans. The results from our animal

experiments indicate that cyclical mechanical impact on the head generates interstitial fluid

movement in the RVLM, leading to FSS-induced decrease in AT1R expression in astrocytes in situ,

and thereby ameliorates hypertension. Our studies also show that the VOCR of hypertensive adult

humans, which produces vertical accelerations at their heads, lowers their BP.

Methods

Animal experiments and human studies. Animals were housed under a 12/12 hour light-dark

cycle with controlled temperature (23−25°C), and treated with humane care under approval from

the Animal Care and Use Committee of National Rehabilitation Center for Persons with Disabilities

(approval number: 30-07). Male SHRSP/Izm and WKY/Izm rats were provided from the Disease

Model Cooperative Association (Kyoto, Japan) and astrocyte-GFP mice (Aldh1L1-GFP mice)[37]

were obtained from GENSAT (New York, NY), acclimated to the laboratory environments for at

least 1 week, randomly divided into experimental groups, and used for experiments.

All participants in our human studies provided written informed consent. The studies were

approved by the Ethics Committees of the Iwai Medical Foundation and the National Rehabilitation

Center for Persons with Disabilities (approval number: 30-01).

Chemicals and antibodies. All the chemicals were purchased from Sigma-Aldrich unless noted

otherwise. Primary antibodies and their dilution rates used in this study are as follows: mouse

monoclonal anti-GFAP (MAB360; Millipore, Billerica, MA) at 1:1,000; rabbit polyclonal anti-

GFAP (Z0334; Dako, Glostrup, Denmark) at 1:1,000; chicken polyclonal anti-GFAP (ab4674;

Abcam, Cambridge, UK) at 1:2,000; rabbit polyclonal anti-cleaved caspase-3 (9661; Cell Signaling

Technology, Danvers, MA) at 1:1,000; mouse monoclonal anti-NeuN (MAB377; Millipore) at

1:200; rabbit polyclonal anti-NeuN (ABN78; Millipore) at 1:1,000, rabbit polyclonal anti-AT1R

(HPA003596; Sigma-Aldrich, St. Louis, MO) at 1:200; rabbit polyclonal anti-AGTRAP

(HPA044120; Sigma-Aldrich) at 1:1000; rabbit polyclonal anti-GFP (598; MBL, Nagoya, Japan) at

1:2,000; chicken polyclonal anti-GFP (ab13970; Abcam) at 1:2,000; mouse monoclonal anti-TUJ-1

(ab78078; Abcam) at 1:1000. Secondary antibodies conjugated with Alexa Fluor 350, 488, 568,

633, and 647 (Thermo Fisher Scientific, Waltham, MA) were used at the dilution rate of 1:400. Cell

nuclei were stained with DAPI (D9542; Sigma-Aldrich).

PHM application to rats. Rats were subjected to PHM in a prone position using a platform that we

developed to move their heads up and down[24] (schematically represented in Fig. 1a). During PHM,

animals were kept anesthetized with 1.5% isoflurane except for the μCT study, in which we used

intraperitoneal injection of 2 mg/kg of midazolam (Sandoz, Basel, Switzerland), 2.5 mg/kg of

butorphanol (Meiji Seika, Tokyo, Japan) and 0.15 mg/kg of medetomidine (Kyoritsu Seiyaku,

Tokyo, Japan) for anesthesia. Body temperature of tested animals was maintained using a light

heater. The PHM system was set up to reproduce the head motion (5 mm, 2 Hz) of treadmill

running (20 m/minute) which made 1.0 x g vertical acceleration peaks at the head of rats

examined[24]. The control rats in PHM experiments were anesthetized likewise, and placed in a prone

position with their heads on the platform that was left static.

Treadmill running of rats. Rats were subjected to compulsive running using a belt drive treadmill

equipped with an electrical shock system (MK-680S; Muromachi, Tokyo, Japan). We habituated the

rats to the treadmill system by placing them in the machine several times without turning on the

treadmill belt during the acclimation period. The electrical stimulation was turned on only once or

twice during the first 5 minutes of the 30-minute treadmill running on the first day of the 4-week

treadmill running period. Thereafter, we did not need to turn on the electrical shock system to have

the animals keep running, perhaps because the velocities we employed (20 m/minute) were

moderate. The control rats in treadmill running experiments were placed on the belt daily for 30

minutes without turning on the treadmilling.

Measurement of BP and HR of rats by radio-telemetry. Rats were implanted with a telemetry

pressure probe (Millar, Houston, TX) in the abdominal aorta at 9−10 weeks of age, following the

surgical procedure described previously[61]. Rats were allowed to recover for at least 14 days before

the initiation of experimental interventions or analyses. During the periods of experimentation that

involved repeated BP and HR measurements over multiple weeks, BP and HR were monitored and

recorded for continuous 30 minutes every 7 days between 10:00 AM and 12:00 PM (noon) by a

multichannel amplifier and signal converter (LabChart 8; ADInstruments, Dunedin, New Zealand).

Mean arterial pressure (MAP) was calculated with a standard formula as follows: MAP = diastolic

BP (DBP) + 1/3 [systolic BP (SBP) – DBP].

Measurement of urinary norepinephrine excretion of rats. Urine excreted during the indicated

24-hour period was collected by means of a metabolic cage (KN-646; Natsume Seisakusho, Tokyo,

Japan), and stored at −80˚C until assayed. Excretion of urinary norepinephrine was calculated by

multiplying its concentration measured using an enzyme-linked immunosorbent assay (ELISA) kit

(KA1891; Abnova, Taipei, Taiwan) with the urine volume.

Microinjection into rat RVLM. Rats were anesthetized with intraperitoneal injection of

midazolam, butorphanol, and medetomidine except for Ang II or valsartan injection studies, in

which we used 1.2−1.4 g/kg of urethane (Sigma-Aldrich), and subjected to microinjection as

described previously[62]. In brief, a 25s-G microsyringe (Hamilton, Bonaduz, Switzerland) was

stereotaxically positioned on anesthetized rats after exposure of dorsal surface of medulla. The

needle placement was defined according to an atlas of the rat with stereotaxic coordinates[63];

anteroposterior angle: 18˚, 1.8 mm lateral to the calamus scriptorius, 3.5 mm ventral to the dorsal

surface of the medulla. The placement of the needle tip in RVLM was confirmed by ensuring the

pressure response to a test-dose injection of L-glutamate[12,64] (100 nL of 1 mmol/L in PBS).

Microinjection of various compounds or mediums was made through a needle reinserted at the

same coordinates with fixed infusion rates using a microsyringe pump instrument (KD scientific,

Holliston, MA). Except for experiments to analyze pressor or depressor responses, we held the

syringe for 5 minutes after the injection to avoid reflux, pulled out the needle carefully, and sutured

the skin. The volumes and rates of microinjection were as follows; Ang II (Auspep, Tullamarine,

Australia) and valsartan (Tocris Bioscience, Bristol, UK): 100 nL of 1 mmol/L in PBS at 0.1

μL/minute, AAV solutions: 300 nL at 0.03 μL/minute, PEG solutions: 1 μL at 0.1 μL/minute,

Isovist: 1 μL at 0.2 μL/minute.

Analysis of pressor/depressor responses. Rats implanted with a telemetry pressure probe were

anesthetized with urethane, and subjected to the analysis of pressor/depressor responses. Monitoring

the BP, we injected Ang II or valsartan (100 pmol) stereotaxically into the unilateral RVLM

following the microinjection procedures described above. The injection side (right or left) was

chosen randomly. When both pressor and depressor responses were analyzed, at least 2 hours

elapsed between the injections of Ang II and valsartan (Fig. 2a). The maximal MAP change elicited

from the baseline was statistically analyzed[14].

Production of AAV vectors. To obtain astrocyte- and neuron-specific transduction, we used AAV9

vectors that expressed a transgene under the control of mouse GFAP and rat NSE promoters,

respectively. The astrocyte-specific GFAP promoter consists of 0.6-kb hybrid fragments containing

ABC1D genomic regions upstream of the mouse *GFAP* gene[65]. The neuron-specific NSE promoter

is composed of the 1.2-kb genomic region upstream of the rat *NSE* gene[66]. Full-length rat AGTRAP

cDNA was synthesized (Eurofins Genomics, Tokyo, Japan) and inserted into plasmid pAAV-GFAP-

GFP-P2A-Cre-woodchuck hepatitis virus post-transcriptional regulatory element (WPRE)-simian

virus 40 polyadenylation signal (SV40pA) and pAAV-NSE-GFP-P2A-Cre-WPRE-SV40pA to

generate pAAV-GFAP-GFP-P2A-AGTRAP-WPRE-SV40pA and pAAV-NSE-GFP-P2A-

AGTRAP-WPRE-SV40pA. pAAV-GFAP-GFP-WPRE-SV40pA and pAAV-NSE-GFP-WPRE-

SV40pA were used for experimental controls. Recombinant single-strand AAV2/9 vectors were

produced by transfection of HEK293T cells (Thermo Fisher Scientific) with the respective pAAV

expression plasmid, pAAV2/9 and a helper plasmid (Stratagene, La Jolla, CA) as previously

described[67]. After harvesting conditioned medium, the viral particles were precipitated using

polyethylene glycol 8000 and iodixanol continuous gradient centrifugation as previously

described[68]. The genomic titer of purified AAV9 vectors was determined by qPCR that targeted the

WPRE sequence.

Measurement of pressure in rat RVLM. Intra-brainstem pressure (IBP) was measured using a

blood pressure telemeter by means of the procedure we previously reported[24,69]. The pressure sensor

was placed in rat RVLM using the same approach with the microinjection described above. During

IBP measurement, respiration was monitored using a respiration sensor attached to the tested rats.

Low-pass (50 Hz) filtered IBP waves were analyzed using LabChart 8 software (ADInstruments).

We observed ~0.5-Hz respiration-synchronized IBP changes with ~2.5 mm Hg magnitude (peak to

peak) as well as 2-Hz PHM-specific waves with ~1.2 mm Hg magnitude (peak to peak).

In vivo analysis of the distribution dynamics of interstitial fluid in rat RVLM using µCT.

Isovist (Bayer, Berlin, Germany) was stereotaxically microinjected into RVLM of anesthetized 12-

week-old male WKY rats following the procedure described above, and visualized using µCT

(inspeXio SMX-100CT, Shimadzu, Kyoto, Japan). After Isovist injection, rats were subjected to

two serial brain µCT scans between which PHM was either applied or left unapplied (kept

sedentary) for 30 minutes (Fig. 4e). µCT images were analyzed using software for 3D morphometry

(TRI/3D-BON-FCS64, RATOC System, Tokyo, Japan). Voxels with ≥1.02 times signal intensity as

compared with that of air was defined as Isovist cluster in rat RVLM.

Simulative calculation of the magnitude of FSS on the cells in rat RVLM during PHM. We

calculated interstitial fluid flow-derived FSS imposed on cells in rat RVLM by referring the results

from our μCT analysis (Fig. 4g,h) to the Henry Darcy's law, which defines the flux density of

penetrating fluid per unit time[70]). The velocity of interstitial fluid flow (u) is assumed to approach

the superficial velocity (u_∞) and zero ($u=0$) at the cell surface (i.e., a no-slip condition). Using these

two boundary conditions together with the Brinkman equation, FSS (τ) at the interstitial cell surface

can be obtained as described in Supplementary Table 1[60].

Cell culture. Primary cultures of astrocytes were prepared from the cortex of neonatal (2−3 days

old) astrocyte-GFP mouse[37] pups by physical dissociation as previously described[71]. Cells were

maintained in DMEM/F12 supplemented with 10% FBS (GE Healthcare Life Science,

Marlborough, MA), 100 I. U./mL penicillin and 100 μg/mL streptomycin at 37°C in a humidified

incubator (5% CO_2 and 95% air). Culture medium was replaced with fresh one every 3−4 days until

confluent. Astrocytes were detached with trypsin/EDTA (0.05% trypsin, 0.53 mM EDTA in PBS),

replated at a ratio of 1:10, and grown to approximately 80−90% confluence prior to use.

Mouse neuroblastoma-derived Neuro2A cells (provided from Dr. Yokota, Tokyo Medical and

Dental University, Japan), which exhibit neuronal phenotypes and morphology[39,40], were cultured in

DMEM (Wako, Japan) supplemented with 10% FBS, 100 I. U./mL penicillin and 100 μg/mL

streptomycin at 37°C in a 5% CO_2 incubator.

1007

Application of FSS to astrocytes or Neuro2A cells in culture. Astrocytes or Neuro2A cells grown

in a poly-D-lysine-coated 35-mm culture dish (Corning Life Sciences, Corning, NY) were exposed

to pulsatile FSS (average 0.7 Pa) for 30 minutes. As we previously reported[23,24,35], a parallel plate

flow-chamber and a roller pump (Masterflex; Cole-Parmer, Vernon Hills, IL) were used to apply

FSS. The flow-chamber, which was composed of a cell culture dish, a polycarbonate I/O unit, and a

silicone gasket, generated a 23-mm-long 10-mm-wide 0.5-mm-high flow channel. To maintain pH

and temperature of culture medium, we used a 5% CO_2-containing reservoir and a temperature-

controlled bath.

**Tissue preparation and immunostaining (immunohistochemical or immunocytochemical

analysis).** Rats were anesthetized with intraperitoneal injection of midazolam, butorphanol, and

medetomidine, and perfused transcardially with 4% paraformaldehyde (PFA; TAAB Laboratories

Equipment, Aldermaston, UK). The brainstems were excised and post-fixed with 4% PFA in PBS

overnight at 4°C. The tissues were cryoprotected by soaking in 20% sucrose/PBS for 24 hours and

in 30% sucrose/PBS for additional 24 hours at 4°C. Fixed brainstems were frozen in optimal cutting

temperature (OCT) compound (Sakura Finetek, Tokyo, Japan) and cut into 16 μm-thick coronal

sections using a cryostat (CM3050S; Leica Microsystems, Wetzlar, Germany). Sliced sections were

permeabilized and blocked with 0.1% Tween-20 and 4% donkey serum (Merck Millipore,

Burlington, MA) in Tris-buffered saline, blocked with, and stained by incubating with primary

antibodies at appropriate dilutions followed by their species-matched secondary antibodies. Cell

nuclei were stained with DAPI (Sigma-Aldrich). The slides were mounted with ProLong Gold

Antifade Reagent (Thermo Fisher Scientific) and images were captured with a BZ-9000 digital

microscope system (Keyence, Osaka, Japan).

For immunocytochemistry, cultured cells were fixed with 4% PFA in PBS for 20 minutes at

room temperature (RT) and permeabilized and blocked with 0.1% Triton X-100 and 10% FBS

(Thermo Fisher Scientific) in PBS for 30 minutes at RT. The cells were then incubated with primary

antibodies for 2 hours and then with secondary antibodies for 1 hour at RT.

1035

Quantitative PCR Analysis (reverse transcription and real-time PCR). 500 ng of total RNA

extracted from cell culture were subjected to reverse transcription, using ISOGEN II (NIPPON

GENE, Tokyo, Japan) and PrimeScript RT reagent Kit (TaKaRa, Kusatsu, Japan). The resulting

cDNA was subjected to real-time PCR analysis using glyceraldehyde-3-phosphate dehydrogenase

(GAPDH) as an internal control in Applied Biosystems 7500 Real Time PCR System with Power

SYBR Green PCR Master Mix (Thermo Fisher Scientific).

The primers (sense and antisense, respectively) were as follows: mouse *Agtr1a* (AT1R-encoding

gene); 5'-AAAGGCCAAGTCGCACTCAAG-3'

and 5'-TCCACCTCAGAACAAGACGCA-3',

mouse *Gapdh* (GAPDH-encoding gene); 5'-GCAAAGTGGAGATTGTTGCCAT-3' and

5'-CCTTGACTGTGCCGTTGAATTT-3', WPRE (for genomic titration of purified AAV9 vectors);

5'-CTGTTGGGCACTGACAATTC-3' and 5'-GAAGGGACGTAGCAGAAGGA-3'.

1048

Fluorescent Ang II binding assay. Six or twenty-four hours after the termination of FSS

application, cultured astrocytes were incubated with Ang II type 2 receptor inhibitor, PD123319

(10^{-6} mol/L in PBS; ab144564, Abcam), for 20 minutes, and then with tetramethylrhodamine

(TAMRA)-labeled Ang II (10^{-8} mol/L in PBS; AS-61181, AnaSpec, Fremont, CA) for 30 minutes.

After 3 times wash with PBS, samples were fixed and subjected to anti-GFP immunostaining to

strengthen the GFP-derived green fluorescence signals and corroborate our analysis on astrocytes

prepared from astrocyte-GFP mice as well as to secure the binding of fluorescent Ang II. Green and

red fluorescence was viewed with a fluorescence microscope (BZ-9000 HS, Keyence). Samples

from astrocytes left unexposed to FSS were prepared and viewed likewise, and provided an

experimental control.

1059

Hindrance of interstitial fluid movement (flow) by introduction of hydrogel in rat RVLM. Just

before use, a pre-mixture of polyethylene glycol (PEG) with functional groups (25 g/L in PBS) was

prepared from tetra-armed thiol-terminal (TetraPEG-SH) (Yuka-Sangyo, Tokyo, Japan) and

acrylate-terminal (Tetra-PEG-ACR) (JenKem Technology, TX, USA) PEG solutions as we

previously described[24]. Tetra-armed polyethylene glycol without functional groups (25 g/L in PBS)

was used as an ungelatable control. For the analysis of hydrogel distribution in rat RVLM, we used

Tetra-PEG-SH fluorescently labeled with a thiol-reactive probe (Merck KGaA, Darmstadt,

Germany). Microinjection of PEG solutions into rat RVLM was conducted as described above.

To specifically analyze the consequences of PHM and hydrogel introduction by minimizing

possible invasive influences of microinjection itself, we gave 1-week recovery time before the first

BP measurement, and then applied PHM to the rats (daily 30 minutes, 14 or 28 days). Immediately

subsequent to post-PHM 24-hour urine collection (Fig. 6a), rats were sacrificed by transcardial

infusion of PFA and subjected to histological analysis.

1073

Terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling (TUNEL) assay. Rat

RVLM sections were stained using a TUNEL kit (Biotium, Fremont, CA) according to the

manufacturer's protocols, counterstained with DAPI, and then viewed using a 20x objective with a

fluorescence microscope (BZ-9000 HS, Keyence). The nuclei of apoptotic cells were determined by

the existence of green fluorescent patches, and cell apoptosis was quantified by referring their

counts to total numbers of nuclei defined by DAPI staining.

1080

1081   **Measurement of accelerations at human head.** To measure the accelerations at the human head

1082   during treadmill running or VOCR, we fixed an accelerometer (NinjaScan-Light; Switchscience,

1083   Tokyo, Japan) on foreheads with a surgical tape. Vertical acceleration was evaluated using the

1084   software application provided from the manufacture.

1085

1086   **Blood sampling and measurement of parameters in plasma and serum of humans.** Blood

1087   sampling in the human study of protocol 2 was conducted between 12 PM and 3 PM. Participants

1088   were rested in a sitting position for at least 10 minutes before starting the sampling procedures.

1089   After plasma (for catecolamines and renin activity) and serum (for aldosterone and CRP) separation

1090   by centrifugation, we outsourced the measurement of parameters be tested (BML, Kawagoe, Japan).

1091

Design and participants of clinical study on antihypertensive effects of vertically oscillating

chair riding. We conducted single-arm clinical studies. The study of protocol 1 (Extended Data

Fig. 5c) was carried out at the affiliated health services facility of Iwai Medical Foundation (Iwai

Keiaien, Tokyo, Japan). The study of protocol 2 (Extended Data Fig. 5d) was carried out at the

National Rehabilitation Center for Persons with Disabilities Hospital.

Subjects were considered eligible if they were 20−75 years old of age and confirmed to have

130−160 mm Hg of SBP at the time of interview for informed consent and eligibility check.

Subjects with mental or psychological illnesses, history or presence of cardiovascular events,

history or presence of severe dysfunction/disorder of liver, kidney, lung, gastrointestinal tract, and

spine, or presence of acute injuries/diseases (e.g., recent traumas and infectious diseases) were

excluded with the exception of those who were given permission for participating in this study from

their primary care physicians. Whereas antihypertensive medication did not disqualify the subjects

(Supplementary Table 2), they were advised not to change their medication from at least one month

prior to the first bout of VOCR through the study period (i.e., up to 8 weeks after the last bout of

VOCR). At a certain (approximately fixed) time point in the morning (typically just before

breakfast), they conducted 3 consecutive measurements of BP (mm Hg) and HR (bpm) using an

automated upper arm-cuff sphygmomanometer, and recorded the values from all those

measurements. These procedures of BP measurement and recording accord with the Japanese

Society of Hypertension Guidelines for the management of hypertension (JSH2019)[72]. Subjects

were directed to start periodical (≥3 days per week) BP/HR measurements at least 2 weeks before

the initiation of the intervention (i.e., daily VOCR) and continue to measure BP/HR throughout the

study period using the same sphygmomanometers. Particularly at the study of protocol 2 (i.e., the

study at the National Rehabilitation Center for Persons with Disabilities), participants were advised

to record all the data of those measurements. Those whose BP lowered below the eligibility

requirement of the study (≥130 mm Hg of SBP) prior to the initiation of VOCR intervention were

eliminated from the study. Participants were directed to be rested and keep calm for at least 1

minute before starting to measure BP/HR. The mean BP/HR value from 3-time measurements was

defined as "value of the day", and used for statistical analysis. When BP and HR were measured

and recorded on ≥3 days during a particular week in the study of protocol 2, the mean of all the

"value of the day"s through the week was defined as "value of the week". For the participants who

agreed. periodical BP/HR measurement and recording (≥3 days per week) was extended up to 8

weeks after the last bout of VOCR.

1124

1125 **Statistical analysis.** All the quantitative data are presented as mean ± s.e.m. Parametric statistical

1126 analyses were conducted by paired or unpaired two-tailed Student's *t*-test for two-group

1127 comparison, and ANOVA with Tukey's post hoc test for multiple (≥3) group comparison, using

1128 Prism software (Version 8, GraphPad Software, San Diego, CA). Differences were considered as

1129 significant at P values below 0.05.