Prevailing role of mucosal immunoglobulions and B cells in teleost skin immune responses to bacterial infection

Xiao-Ting Zhang[1][¶], Yong-Yao Yu[1][¶], Hao-Yue Xu[1][¶], Zhen-Yu Huang[1], Xia Liu[1], Jia-Feng Cao[1],

Kai-Feng Meng[1], Zheng-Ben Wu[1], Guang-Kun Han[1], Meng-Ting Zhan[1], Li-Guo Ding[1], Wei-Guang

Kong[1], Nan Li[2], Fumio Takizawa[3], Sunyer J Oriol[4], Zhen Xu[1,5]*

1 Department of Aquatic Animal Medicine, College of Fisheries, Huazhong Agricultural University, Wuhan, Hubei, 430070, China. **2** State Key Laboratory of Freshwater Ecology and Biotechnology, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, Hubei, 430072, China. **3** Faculty of Marine Science and Technology, Fukui Prefectural University, Obama, Fukui 917-0003, Japan. **4** Department of Pathobiology, School of Veterinary Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, 19116, USA. **5** Laboratory for Marine Biology and Biotechnology, Qingdao National Laboratory for Marine Science and Technology, Qingdao, Shandong, 266071, China.

[¶]These authors contributed equally to this work.

* Corresponding author.

E-mail: zhenxu@mail.hzau.edu.cn

Abstract

The skin of vertebrates is the outermost organ of the body and serves as the first line of defense against external aggressions. In contrast to mammals, the skin of early vertebrates like fish, lacks keratinization and has evolved to operate as a mucosal surface. As such, it has been shown to contain a skin-associated lymphoid tissue (SALT) in the epidermis. Prior to the discovery of IgT as a key mucosal immunoglobulin (Ig) in fish, several studies reported IgM-specific titers in the skin mucus upon pathogenic infection or vaccination. In contrast, a recent report showed IgT being the main Ig responding in the skin upon parasitic challenge. Whether IgT is the main Ig isotype elicited in the skin against bacterial pathogens, and whether such responses are locally induced within the SALT are important questions that remain unanswered. Here we analyzed the immune response of trout skin upon infection with one of the most widespread fish skin bacterial pathogens, *Flavobacterium columnare* (*F. columnare*). To gain insight into the pathogenesis and progression of the infection in skin, a recombinant *F. columnare* expressing green fluorescent protein (GFP) was constructed and showed to successfully invade the skin mucosa. This pathogen induced strong skin innate immune and inflammatory responses at the initial phases of infection. More critically, we found that the skin mucus of fish having survived the infection contained significant IgT- but not IgM- or IgD-specific titers against the bacteria. Moreover, we demonstrate the local proliferation and production of IgT$^+$ B-cells and pathogen-specific IgT titers respectively within the SALT upon bacterial infection. Thus, our findings represent the first demonstration that IgT is the main Ig isotype induced by the skin mucosa upon bacterial infection, and that because of the large surface of the skin, its SALT probably represents a prominent IgT inductive site in fish.

Author summary

In contrast to mammalian skin, that of teleost fish is the largest mucosal surface containing a skin-associated lymphoid tissue (SALT). Thus far, IgT representing the prevalent immunoglobulin (Ig) in SALT have only been reported upon infection with a parasite. However, very little is known about the types of B cells and Igs responding to bacterial infection in the teleost skin mucosa, as well as the inductive or effector role of the SALT in such responses. To address these questions, we infected rainbow trout (*Oncorhynchus mykiss*) with *F. columnare* an important skin pathogen that causes significant losses both in wild and farmed fish populations. Remarkably, strong pathogen-specific IgT responses and accumulation of IgT$^+$ B cells were induced in the cutaneous mucus and skin epidermis respectively upon infection. Evidence for the skin being an IgT inductive site was shown by the finding of local IgT$^+$ B cells proliferation and pathogen-specific IgT tites in cultured trout skin explants. Thus, our findings demonstrate that IgT is the prevailing Ig isotype induced by the skin mucosa upon bacterial infection, and that the skin represents an IgT inductive site in fish.

Introduction

In contrast to five major Ig classes (IgM, IgG, IgA, IgD, and IgE) in mammals, only three Ig isotypes are present in teleosts, including IgM, IgT/IgZ, and IgD [1]. Teleost IgM is the most abundant Ig class that involved in systemic immunity upon infection or vaccination [2-5]. Secreted IgD (sIgD) has been identified and found coating a significant portion of the fish microbiota [6-8], however, its immune function against pathogens remains largely unknown. Teleost IgT has been demonstrated to play a specialized role in mucosal immunity akin to that of mammalian secretory IgA (sIgA) [6-10]. Similar to sIgA, sIgT plays a key function both in protecting fish mucosal sites against pathogens [7-12], as well as in promoting microbiota homeostasis at mucosal surfaces [13]. With regards to its role in pathogen control, we and others have shown that sIgT is the main Ig induced in several fish mucosal associated lymphoid tissues (MALTs) upon pathogen insult [7-12]. Thus far, pathogen-specific IgT responses have been documented in gut, skin, gill, nasal, buccal and pharyngeal mucosal areas [7-12]. However, in-depth studies assessing the local induction of IgT have only been performed in some of the fish MALTs, including those of the gills, nose and buccal and pharyngeal tissues [7, 8, 11, 12]. Moreover, such local responses have been assessed with the use of only one pathogen type, the parasite *Ichthyophthirius multifiliis* (Ich), a pathogen that is responsible for important losses to both wild and farmed populations of salmonid species.

At this point, nothing is known about the local induction of IgT responses in the skin of fish, by far the largest mucosal surface in these species. Moreover, pathogen-mediated IgT responses in that surface are restricted to one study that utilized the parasite Ich as the pathogen model [9], while it is unknown at this point whether bacterial pathogens are capable of inducing similar responses. Anatomically, the skin presents a similar structure in all vertebrates, which is generally composed of

two layers (epidermis and dermis) (S1 Fig). However, it should be noted that vertebrates of different taxonomic statuses have faced unique evolutionary pressures that have continuously shaped their own skin structures and components. These histological and componential differences in the skin of vertebrates are reflected also in their specific immune strategies [14]. It is worth noting that the term SALT was first used in reference to mammals [15] and that in these species, the main cellular constituents of the skin epidermis are specialized epithelial cells known as keratinocytes, Langerhans cells, and T cells [16]. Moreover, the skin dermis of mammals is home to a diverse array of specialized immune cells, including antigen-presenting dermal dendritic cells (DCs), T cells, B cells, and natural killer (NK) cells as well as mast cells, monocytes, and macrophages [17], which indicates that many immunological processes taking place in the dermis layer of the mammalian SALT (S1A Fig). Interestingly, teleost fish have also been shown to contain a SALT [2]. However, in contrast to mammalian skin, that of teleosts behaves as a mucosal surface since it lacks keratinization, harbors abundant mucus, and its living epithelial cells are in direct contact with the water environment where they live. While the fish dermis is mainly composed of dense connective tissue with a large amount of collagen fibers and is devoid for the most part of immune cells, its epidermis has been shown thus far to contain B cells, T cells, macrophages and granulocytes as the main immune cells (S1B Fig). Moreover, it is in the epidermis where significant immune responses have also been reported [18, 19]. IgT responses in the skin were first described by us, where we showed that IgT-specific responses against a parasite were chiefly detected in the skin mucus, whereas IgM-specific titers were for the most part observed in the serum [9]. Whether IgT and IgT[+] B cell responses are locally induced in the skin is an important question that remains to be addressed. In that regard, it is important to note that in mammals not all IgA responses are induced locally in all existing mucosal sites [20]. For

example, IgA found in some mucosal areas (e.g. female genital tract) and secretory glands (e.g. salivary glands and lacrimal glands) is derived from B cells that are locally induced in other mucosal sites (e.g. nasopharynx-associated lymphoid tissue (NALT) and gut-associated lymphoid tissue (GALT)) [21-23].

As stated above, only one report has described the induction of skin IgT-specific titers against a pathogen (Ich), and local IgT responses have only been assessed, with the use of the same pathogen, only in some fish mucosal sites, but not the skin. Prior to the discovery of the role of IgT in mucosal immunity, several studies in teleosts showed the induction of modest IgM-specific titers against several bacterial pathogens. More recently, a couple of studies were unable to detect IgM-specific titers in the skin mucus of rainbow trout exposed to *Flavobacterium pschrophilum* which prompted the authors to suggest that IgT might play a role against this bacterial pathogen. Thus, in this study we aimed to evaluate the specific contributions of the different fish Igs to skin bacterial pathogen in a representative teleost, and to assess whether such responses could be induced locally in the skin-associated lymphoid tissue (SALT). Here we choose *Flavobacterium columnare* (*F. columnare*), one of the most widespread fish skin bacterial pathogens. *F. columnare* is a long Gram-negative rod in the family *Flavobacteriaceae*, one of the main phyletic lines within the Bacteroidetes group from the domain Bacteria [24], which can survival in both fresh and brackish water throughout the world [25].This pathogen can cause columnaris disease that affects a variety of freshwater fishes, leading significant losses both in wild and farmed fish populations [26-32]. Columnaris disease is characterized by a pronounced erosion and necrosis of mucosal tissues including the fin, skin and gills and it can cause particularly high mortalities. Secreted enzymes, such as proteases and chondroitin sulfate lyases, have been potential *F. columnare* virulence factors, contributing to the

branchial and cutaneous necrosis [33, 34]. While we have shown in the past that *F. columnare* induces dominant IgT- and IgM-specific responses in the gill mucosa and serum of rainbow trout respectively, thus far, little is known with regards to the pathogenesis of columnaris in the fish skin as well as the innate and adaptive immune mechanisms induced by this pathogen in that surface upon infection.

To assess the mucosal immune responses induced in the skin by *F. columnare*, we infected rainbow trout with a newly constructed fluorescent version of this pathogen following a bath exposure strategy [35]. Our findings show that trout skin elicited strong innate and adaptive immune responses to *F. columnare*. More critically, we show that *F. columnare*-specific Ig responses in the skin mucus were mainly represented by the IgT isotype and that such responses were induced locally as shown by the local proliferation and accumulation of IgT$^+$ B cells. Moreover, when cultured in vitro, skin explants produced *F. columnare*-specific IgT titers. These data show for the first time that the use of a fluorescent version of *F. columnare* which was critical to further understand the pathogenesis of this important fish pathogen. More importantly, we demonstrate the previously undiscovered capacity of the fish skin of inducing potent mucosal IgT-specific responses against a bacterial pathogen and that such responses are induced locally upon mucosal infection.

Results

Pathological changes in trout skin after bacterial infection

To evaluate morphological changes in trout skin, we used a bath challenge model with *F. columnare*, which results in the infection of fish mucosal tissues, and elicits strong immune responses [36]. However, due to lack of specific antibody to this bacterial strain, a recombinant *F. columnare* strain containing a plasmid expressing green fluorescent protein (GFP) was constructed to gain insight into the pathogenesis and progression of the bacterial infection in fish skin (S2A Fig). Under the same culture conditions, the GFP-expressing and wild-type *F. columnare* strains grew at similar rate in Shieh medium (S2B Fig). Critically, both the GFP-expressing and wild-type *F. columnare* strains elicited very similar mortality rates when used at the same concentration (S2C Fig). Together, these results indicated that the growth and virulence of the GFP-expressing *F. columnare* (GFP-*F. columnare*) were not influenced by GFP or the plasmid. At 4 days post infection (dpi) with GFP-*F. columnare*, we observed that the classical symptoms of skin lessions, fin rot and gill necrosis appeared in the trout (Fig 1A). Importantly, high expression of *F. columnare*-specific 16S rRNA accumulated in the skin, gills, and fins of trout at 4 dpi (Fig 1B). Notably, by using fluorescence microscope, a large number of *F. columnare* with green fluorescence was observed in the fin, skin and gill tissues from 4-day infected fish, when compared to that of control fish (Fig 1C). In addition, the tissue homogenates of trout skin from control fish and 4-day infected fish were cultured on Shieh agar, and the yellow-green color bacterial colonies showed characteristic rhizoid shapes were detected only from homogenate samples of infected trout skin (red arrows). Moreover, the single colonies isolated from the plate were grown in pure culture, and displayed characteristic elongated rod-shaped *F. columnare* bacteria with green fluorescence (Fig 1D). By fluorescence microscopy, a

time series observation of GFP-*F. columnare* in skin tissue showed that much of the bacteria was found in the skin epithelium, and that bacteria accumulated in trout skin as early as 1 dpi (Fig 1E). Moreover, similar to these results, qRT-PCR analysis showed that the *F. columnare* load in trout skin increased significantly from 1 dpi and peaked at day 4 (Fig 1F), which further confirmed that the bacteria succeeded in invading the skin mucosal tissue.

In combination with paraffin sections and staining (hematoxylin & eosin or alcian blue, H & E / AB), morphological changes were observed in the skin epidermis. We observed a conspicuous decrease in both the thickness of the skin epidermis and the number of mucus cells in the infected trout skin (Fig 2A). Moreover, the thickness of the trout skin epidermis reduced significantly at 2 dpi, and it remained relatively stable for up to 14 days (Fig 2B). In addition, the significantly reduced number of mucus cells in the skin epidermis occurred at 1, 2, and 7 dpi (Fig 2C).

Bacterial infection elicits strong immune responses in trout skin

Using qRT-PCR, we detected the mRNA expression levels of 15 immune-related genes and cell markers, including cytokines (interleukin 8, IL8; interleukin 10, IL10; tumor necrosis factor α, TNFα; and interferon regulatory factor 4, IRF4), cathelicidin (CATH1), complement 3 (C3), poly Ig receptor (pIgR), and Ig heavy chain genes (IgT, IgM, and IgD) (Fig 2D; primers were shown in S1 Table). These studies indicated a strong immune response occurred in trout skin (Fig 2D), spleen (S3A Fig), and head kidney (S3B Fig) after challenge with *F. columnare*. Interestingly, significantly upregulated immune-related genes (e.g., IL10; inducible nitric oxide synthase, NOS2; SAA) were detected in trout skin as early as 1 day, while in trout head kidney (S3A Fig) and spleen (S3B Fig), immune responses were delayed upon challenge with *F. columnare*. It's worth noting that day 2 and day 14 were most correlated in terms of the strength of the skin immune responses. Taken the time point of

expression of the immune-related genes and significant histological responses from 2 dpi to 14 dpi,

we chose these two time points for subsequent high-throughput transcriptome sequencing analysis.

RNA-Seq libraries were constructed from nine samples that separately represented three groups

(Control, 2d, and 14d) which were sequenced on an Illumina platform. Upon *F. columnare* challenge,

we found significantly modified mRNA expression of 4291 genes (2074 genes upregulated and 2217

genes downregulated) at 2 dpi, and 784 genes (391 genes upregulated and 393 genes downredulated)

at 14 dpi, respectively (Fig 2E). More than 25% of DEGs indicated immune-related genes after

filtering by the immune gene library of *Oncorhynchus mykiss* (*O. mykiss*) (Fig 2F). Importantly,

among the immune-related genes, we found significantly modified mRNA expression of innate

immune genes (Fig 2G; e.g. interleukins, chemokines, antibacterial peptides, and mucins) and

adaptive immune genes (Fig 2H; e.g. genes associated with antigen processing and presentation, and

Igs) at both days 2 and 14. Gene ontology (GO) analysis showed that these DEGs were enriched for

genes involved in developmental processes, cell differentiation, signaling transduction, programmed

cell death, and immune response (Fig 2I).These data indicated that *F. columnare* invasion could

induce intense and long-term innate as well as adaptive immune responses. In order to validate the

DEGs identified by RNA-Seq, we chose 12 upregulated and 12 downregulated genes for qRT-PCR

confirmation. The result showed a significant correlation of the expression values determined by

RNA-Seq and qRT-PCR at each time point (S4 Fig), suggesting that the RNA-seq results performed

the same accuracy as that of qRT-PCR in the determination of gene expression in *vivo*.

Pathogen load decreased in fish skin after re-infection

Histopathology of the skin of the both infected and survivor groups showed no significant changes in

the epidermis compared to the control group (Fig 3A and 3B), and the number of mucus secreting

cells also recover to normal levels (Fig 3A and 3C). We then challenged the infected (28 dpi) and

survivor fish as well as uninfected control fish by bath with 5×10^6 CFU/ml of GFP-*F. columnare* for

4 h at 16 ℃. All fish were euthanized at 4 days post challenge to evaluate bacterial load in the skin

tissues from the three fish groups. Using fluorescence microscopy and qRT-PCR, we detected that

upon re-infection, the infected and survivor groups had markedly limited green fluorescence signal

and lower 16S rRNA expression of *F. columnare* compared with those of the control-challenged fish

(Fig 3D and 3E). These data strongly suggest that pre-exposure to *F. columnare* protects trout skin

from the bacterial re-infection, suggesting that mucosal adaptive immunity had been induced in trout

skin to fight pathogen invasion.

B cell responses and local proliferation in trout skin after bacterial infection

Our previous studies have demonstrated that IgT play a specialized role in skin mucosal immunity

against an aquatic parasite [25]. The decreases of *F. columnare* load in infected and survivor trout

skin after re-challenge with a higher dose of the same pathogen, suggesting whether IgT play an

important role in mucosal adaptive immune responses against this bacterial pathogen. To evaluate

the role of skin mucosal Igs in response to bacterial challenge, we detected Ig and B-cell responses in

trout skin infected with *F. columnare* by bath challenge. Using immunofluorescence analysis, we

found very few IgT$^+$ B cells as well as IgM$^+$ B cells in the skin of control fish (Fig 4A;

isotype-matched control antibodies, S5 Fig). However, fish infected after 28 days with *F. columnare*,

a moderate (~ 2-fold) but significant increase in the number of IgT$^+$ B cells was detected in the skin

(Fig 4A and 4B). Importantly, when compared to control fish, we observed a notable accumulation

of IgT$^+$ B cells in the skin of survivor fish. In stark contrast to this, there was no change in the

number of IgM$^+$ B cells whether in infected or survivor fish groups when compared with that of

control fish (Fig 4A and 4B).

To confirm whether the substantial increase of IgT$^+$ B cells in the survivor fish skin was due to

proliferative IgT$^+$ B cell in local skin mucosa or to the migration of these cells from other immune

tissues such as systemic lymphoid organs and other MALT, the *in vivo* proliferation of B cells in

control and survivor fish was detected by caudal vein injection of these fish with fluorescent EdU

(5-ethynyl-20-deoxyuridine). Using flow analysis, the percentage of proliferating IgT$^+$ B cells (EdU$^+$

IgT$^+$ B cells) showed a significant increase in the skin of survivor fish (~ 7.20%) when compared to

control fish (~ 3.45%), while there was no change in the percentage of IgM$^+$ B cells (EdU$^+$ IgM$^+$ B

cells) between control and survivor fish (Fig 4C and 4D). Interestingly, we found a notable rise of

EdU$^+$ IgM$^+$ B cells in the head kidney of survivor fish (~ 6.23%) compared to control fish (~ 2.89%)

while no differences in the proliferation of IgT$^+$ B cells were detected (Fig 4E and 4F).

Bacteria-specific immunoglobulin responses in trout skin

To further study the skin Ig responses in trout skin after *F. columnare* infection, we performed

SDS-PAGE and western blot analysis. Along with the significant increase of IgT$^+$ B cells observed

in the skin of infected and survivor fish, the IgT protein level in the skin mucus of the both groups

was ~ 8- and ~ 15-fold higher respectively, when compared with that of control fish, while the levels

of IgM and IgD did not differ (Fig 5A). In serum, about 2-fold and an approximately 5-fold increases

of IgT concentration were detected in infected and survivor fish, respectively. Conversely, the

concentration of IgM in serum was much higher than that of IgT and had approximately 4- and

3-fold increases in infected and survivor fish, respectively, when compared with those of control fish

(Fig 5E). In contrast, the IgD protein concentration did not change significantly in either the skin

mucus or serum of the same fish groups (Fig 5A and 5E).

The results of large increases in IgT$^+$ B cells and IgT protein levels in the skin of infected and

survivor fish suggested the generation of bacteria-specific IgT immune responses. To verify this

hypothesis, using a pull-down assay, we measured the capacity of skin mucus Igs binding to *F.*

columnare. Compared to control fish, we detected significantly higher (~ 5-fold) bacteria-specific

IgT binding up to 1/40 diluted skin mucus of infected and survivor fish, respectively (Fig 5B-5D).

While bacteria-specific IgT binding was detected in serum with only in the 1/10 dilution of the

survivor group (Fig 5F and 5H). In contrast, ~ 3-fold and ~ 2.4-fold higher bacteria-specific IgM

binding was observed in up to 1/100 serum dilutions both from infected and survivor fish,

respectively (Fig 5G and 5H).   However, bacteria-specific IgD binding is hardly detected in either

the skin mucus or the serum from infected and survivor fish (Fig 5A-5H).

To test further the local production of pathogen-specific Igs in trout mucosal and systemic

tissues respectively, we analyzed bacteria-specific Ig titers of culture medium derived from cultured

skin, head kidney, and spleen explants of control and survivor fish (Fig 6A-6F). We detected

bacteria-specific IgT binding up to the 1/100 dilution of medium of cultured skin explants from

survivor fish, while very low bacteria-specific IgM responses (Fig 6A and 6D) were detected (only in

the 1/2 dilution in the same medium). In contrast, dominant bacteria-specific IgM binding (up to the

1/10 dilution) was observed in the medium of survivor head kidney and spleen explants, while

similar (up to the 1/10 dilution) and lower (up to the 1/2 dilution) bacteria-specific IgT responses

were detected in the medium of survivor head kidney and spleen explants respectively (Fig 6B, 6E,

6C, and 6F). Interestingly, we could not detect any bacteria-specific IgD binding in either the

mucosal (skin) or systemic (head kidney and spleen) tissue explants from survivor fish (Fig 6A-6F).

pIgR in trout skin after bacterial infection

Previous studies have indicated that rainbow trout pIgR may take part in the skin mucosal immune

responses to parasitic infection by mediating the transepithelial transport of secretory Ig [9]. Thus,

we hypothesized that pIgR might also mediate the transport of IgT to the skin mucus during immune

responses to *F. columnare*. Here, using immunofluorescence analysis, we observed that some of the

skin epithelial cells from control fish stained by the anti-trout pIgR antibody (Fig 7A; isotype

matched control antibodies, S5B Fig). Importantly, the number of pIgR$^+$ cells significantly increased

in the epidermis of infected (~ 1.6-fold) and survivor fish (~ 2.4-fold) (Fig 7A and 7B) when

compared with those of control fish. Using qRT-PCR, the expression of pIgR was found to be

significantly upregulated in the skin of infected (~ 3-fold) and survivor fish (~ 6.6-fold), respectively,

when compared with that of control fish (Fig 7C). Moreover, the concentration of pIgR in skin

mucus post bacterial infection were also detected and calculated by western blot analysis. Our results

showed that the relative fold changes of pIgR in infected and survivor fish when compared to that of

control fish, were significantly increased (~ 1.7-fold for infected group and ~ 3.4-fold for survivor

group; Fig 7D and 7E).

Discussion

Fish skin, is the outermost organ of the organism's body, which is continuously exposed to a wide array of potentially harmful insults in the aquatic environment. This constant exposure to microbes has led to the evolution of a mucosal immune network in the teleost SALT, which ensures that adequate immune responses are mounted upon antigenic challenge while maintaining overall tissue and microbiota homeostasis [16, 37-39]. However, till now little is known with regards to the evolution of skin mucosal B cell and antibody responses to bacterial pathogens in early vertebrates. Most of the literature in that area for teleost fish predates the discovery of IgT as the key mucosal immunoglobulin in these species [10]. In that regard, some studies in several teleosts have reported low levels of pathogen-specific IgM titers upon bacterial infection [40, 41]. Moreover, whether the skin mucosa behaves as an inductive or effector site of mucosal Ig responses has never been investigated. Here, we report for the first time in a teleost that infection with *F. columnare*, a widespread fish bacterial pathogen, can elicit local skin proliferative IgT$^+$ B cell responses as well as significant bacterial-specific mucosal IgT titers.

Columnaris disease, caused by the Gram-negative bacterium *F. columnare*, severely impacts the global production of many fish species [33, 42]. As a mucosal bacterial pathogen, *F. columnare* infection results in skin epithelial erosion, gill necrosis and ulcers, with a high degree of mortality [33, 42]. To evaluate mucosal immune responses to *F. columnare*, we exposed fish to this pathogen following a well-established waterborne challenge model with a novel fluorescent (GFP) strain of *F. columnare* [45]. Importantly, GFP-*F. columnare* exhibited similar growth and virulence properties as the wild-type bacteria, and thus, this newly constructed GFP-*F. columnare* represents a new reagent that is expected to significantly advance the study of host- *F. columnare* interactions. At 4 dpi,

clinical signs were detected in the bacteria-infected rainbow trout, and qRT-PCR and

immunofluorescence further proved the successful invasion of GFP-*F. columnare* into rainbow trout.

It is worth point that the use of our unique GFP-*F. columnare* strain was extremely instrumental in

assessing the capacity of this pathogen to intrude into the skin epidermis of fish under the

experimental conditions tested. Importantly, upon infection, we observed significantly decreased

numbers of mucus cells and thickness of the epidermis in the trout skin. Thus, our results suggest

that *F. columnare*'s successful invasion relies on destroying the epidermal layer structure of fish skin,

which is in line with what has been shown in previous studies [44, 46]. However, the mechanisms of

bacterial infection still need to be further investigated. In agreement with the noteworthy

histopathological changes in trout skin, we found that 15 immune-related genes were significantly

upregulated in trout skin as early as 1 dpi when compared with those in the head kidney and spleen,

which indicates that the immune responses in trout skin are activated significantly earlier than in

systemic lymphoid tissues, thus supporting further the notion of the skin acting as the first line of

defense against bacterial infections in fish. Importantly, both innate and adaptive immune molecules

were induced in the skin upon *F. columnare* infection, including antibacterial peptides, cytokines,

chemokines, complement factors, and Igs. Thus, our results provide a general picture of the

predominant immune responses that take place in the trout skin after *F. columnare* infection. Of these

responses, it is worth pointing the increased transcript levels of cathelicidin 1 and proinflammatory

cytokines such as IL8 and TNFα found in trout skin immediately after *F. columnare* infection.

Moreover, the anti-inflammatory cytokine IL10 was also induced after bacterial infection. Combined

with previous studies [47-50], our results suggest that *F. columnare* infection induces a significant

inflammatory reaction as well as strong immune responses in the fish skin epidermis early upon

infection. In contrast, these types of responses occur in the dermis of mammals [16, 51], probably

reflecting the different physiological needs of fish and mammalian skin tissues with regards to the

water or terrestrial environments where these species live.

The use of GFP-*F. columnare* enabled us to find that the skin epithelium of trout previously

exposed and thus, immune to *F. columnare*, cannot get intruded upon re-exposure to this pathogen.

These findings suggested that local humoral immunity might be elicited in trout SALT thus

contributing to protection upon re-infection. Here we first showed significant increases in the

concentration of IgT but not IgM in the skin mucus of infected and survivor fish exposed to *F.*

columnare. These data correlated with the large accumulation of IgT$^+$ but not IgM$^+$ B cells in the

skin epidermis of the same animals. These results paralleled those obtained in a previous study where

fish surviving from a parasite infection exhibited large accumulations of IgT and IgT$^+$ B cells in the

skin epidermis [9]. It is worth mentioning that upon bacterial infection, several mammalian MALTs

have been shown to have similar significant accumulations of IgA$^+$ B cells [52-56]. In humans,

several studies have described a correlation between resistance to *Vibrio cholerae* infection and high

specific sIgA titers [57, 58]. Importantly, our findings show for the first time in fish the detection of

bacteria-specific IgT titers mainly in the skin mucus and to a much lesser degree in the serum of

survivor fish. Although the kinetics of IgM production in the skin following bacterial infection had

been investigated in some species [59-62], till now the role of sIgT in skin bacterial infections had

not been addressed, probably due to the lack of specific antibody reagents recognizing IgT [63, 64].

Moreover, we hardly found any bacteria-specific IgM titers in skin mucus of infected and survivor

fish, which were instead detected almost exclusively in the serum in very high levels. Therefore, we

can preliminarily conclude that skin IgT and IgT$^+$ B cell responses against *F. columnare* are

specifically confined in its mucosa, whereas IgM responses are overwhelmingly found in the serum, and thus of systemic nature. Interestingly, previous studies in trout with a similar bacterial pathogen (*Flavobacterium psychrophilum*) have shown that bath exposure of trout to this pathogen could not elicit bacteria-specific IgM responses in the trout skin mucus [65], as we have also found for *F. columnare*. Thus, the authors speculated that perhaps IgT is the immunoglobulin involved in skin responses against *F. pychrophilum*, a hypothesis that seems to be supported by our data, although that hypothesis remains to be investigated.

The accumulation of IgT[+] B cells and high bacteria-specific IgT titers observed in the skin mucosa from infected and survivor fish led us to hypothesize the local proliferation and induction of IgT[+] B cells and IgT antibody responses respectively upon bacterial infection. Confirming this hypothesis, we found strong local proliferative IgT[+] B cells responses in the teleost skin but not in the head kidney of the same fish, suggesting that the accumulation of IgT[+] B cells in their skin is due to local proliferation rather than migration of these cells from the systemic immune organs. We cannot rule out however the possibility that these IgT[+] B cells are induced in other mucosal surfaces, and upon migration to the SALT, they start proliferating. Future work will have to address the aforementioned possibility. Critically, the production of significant titers of bacteria-specific IgT in skin explant cultures confirmed that these IgT antibodies are produced locally and suggest the presence of specific plasma cells in the local skin mucosa. Overall, considering that *F. columnare* firstly invades and stays in the skin epidermis, it is unlikely that the inductive site is a systemic lymphoid tissue, thus our results imply that trout SALT is an inductive site of the observed IgT mucosal immune responses. We have previously shown similar results in the mucosa of gill and nose when exposed to pathogenic challenge [7, 8]. It is worth noting also that similar results have been

shown in mucosal inductive sites of some mammalian MALTS, including the GALT, NALT and

bronchus-associated lymphoid tissue (BALT) where a local mucosal response governed by a mucosal

Ig (sIgA) was strongly induced upon bacterial infection, and in contrast, the systemic response is

dominated by IgG/IgM [66-68].

A key feature of pIgR is the mediation of the transepithelial transport of secretory Igs into the

mucosal surfaces in both mammals and teleost [2, 69]. Previous studies have shown that the putative

trout secretory component (tSC) of pIgR is associated with mucosal IgT and IgM in gut [10], skin [9],

gills [7], and nasal mucus [8], contributing to the Ig response to parasite infection. Here, we found

that trout pIgR (tpIgR) was mainly expressed in the epithelial layer of the skin from control, infected,

and survivor fish. Critically, here we see a significant increase in the numbers of pIgR^{+} cells and

transcript levels of pIgR in the skin from bacterial infected trout when compared with that of control

fish. Interestingly, similar results have been reported in mammals, for example, the expression of

pIgR was found upregulated in germ-free mice implanted with *Bacteroides thetaiotaomicron* [70]. In

addition, pIgR expression can be upregulated in HT-29 cells by reovirus (double-stranded RNA) [71]

and bacteria (*Enterobacteriaceae*) with LPS on the surface [72]. Thus, our results indicate such

increases in the levels of pIgR upon pathogenic challenge is an evolutionary conserved feature from

fish to mammals which probably has the role to increase the transport rate of mucosal

immunoglobulins to mucosal surfaces upon pathogenic insult.

In conclusion, our study provides the first evidence of local B cell proliferation and IgT

secretion in teleost skin, which suggests that teleost SALT is an inductive site of mucosal B cell and

immunoglobulin responses. Based on our studies, we propose a model in which upon bacterial

infection, epidermis damage occur which leads to an up regulation of immune-related genes in the

skin as well as to processes of IgT$^+$ B cell activation and proliferation within the SALT, resulting into

the production of bacteria-specific IgT (Fig 8). Thus, from an evolutionary viewpoint, our results

reinforce the formerly stated view [73] that regardless of their phylogenetic origin and tissue

localization, specialized mucosal Igs (i.e., IgA, IgT) of all vertebrate MALT-containing surfaces

function under the guidance of primordially conserved principles. Moreover, our findings in this

phylogenetically primitive vertebrate strongly suggest a universal requirement for vertebrate

MALT-containing mucosal surfaces to use a dedicated Ig for the maintenance of homeostasis.

Additionally, since many fish pathogens enter the host through the skin mucosa, the knowledge

derived from our findings also has special relevance from a more practical perspective, as it may lead

to the development of fish vaccines that can effectively induce bacteria-specific IgT immune

responses in skin.

Materials and methods

Ethics statement

All experimental protocols were performed in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the Ministry of Science and Technology of China. They were approved by the Scientific Committee of Huazhong Agricultural University (permit number HZAUFI-2017-013). All efforts were made to minimize the suffering of the animals.

Fish

Rainbow trout (~ 5 g) were obtained from a fish farm in Shiyan (Hubei, China) and were maintained in aerated aquarium tanks with a water recirculation system including internal biofilters and thermostatic temperature control. Fish were acclimatized for at least 2 weeks at 16 ℃, and fed daily with commercial trout pellets at a rate of 1% biomass during the whole experiment periods. The feeding was terminated 2 days prior to sacrifice.

F. columnare culture and construction of GFP-*F. columnare* G$_4$ strain

The strain G$_4$ of *F. columnare* used in this study was provided by Professor Nie [74] and conserved at -80 ℃. *F. columnare* strains were grown at 28 ℃ in Shieh medium or on Shieh agar. The GFP-*F. columnare* G$_4$ strain was constructed by using the method previously described by Li et al [74]. Briefly, a fused fragment carrying P*ompA* of *F. johnsoniae* (*Flavobacterium johnsoniae*) and *gfp* was amplified by PCR using the primers pr44-Bam (CGCGGATCCGGCAGCGCATACCAAAGAACACTTAG) and pr38-Pst (GCTAGCTGCAGCAGATCTATTTGTATAGTTCATCCA) with pAS43 as template. The PCR product was digested with BamHI and PstI and cloned into the corresponding sites of pCP23 to

440 generate plasmid pCP-gfp. This plasmid was transferred into *Escherichia coli* (*E. coli*) S17-1 $^\lambda$ pir

441 and then was introduce into G_4 by conjugation. The GFP-labeled strain was screened on Shieh agar

442 containing 1 µg/ml tobramycin and 10 µg/ml tetracycline.

Growth curve determination

To compare the growth rates of GFP-*F. columnare* and wild-type *F. columnare* G_4 strain *in vitro*, both

strains were streaked on Shieh agar at 28 ℃ for 24 h. The colonies were then inoculated into Shieh

broth and incubated at 28 ℃ with shaking at 150 rpm. The optical density was measured at 600 nm

with 2ml cuvette using NanoPhotometer® NP80 (Implen, Germany). Until the OD_{600} reached 0.4,

500 µl of such early-exponential-phase were inoculated in 25 ml Shieh broth and incubated at 28 ℃

with shaking at 150 rpm. The OD_{600} of culture liquid from each group was measured every 1 h for 13

h.

Rainbow trout challenges

The method used for *F. columnare* infection were described previously [19] with slight modification.

Fish (~ 5 g per fish) were randomly chosen and exposed by immerse with wild-type or GFP-*F.*

columnare at a final concentration of 1×10^6 CFU/ml for 4 h at 16 ℃, and then migrated into the

aquarium containing new aquatic water. To detect the localization of *F. columnare* in trout, 30 fish

were infected with GFP-*F. columnare*, and tissue samples including skin, gill and fin were collected

at 1, 2, 4, 7 and 14 dpi. For the study of immune responses in trout, there were two types of

challenges were performed. In the first group (infected fish), 60 fish were challenged by wild-type

G_4 strain. Tissue samples including skin, head kidney, and spleen were collected at 1, 2, 4, 7, 14, and

28 dpi. Skin mucus and serum were taken 4 weeks later as infected group (fish survived 28 days after

one exposure). In the second group, another 60 fish were monthly exposed to wild-type *F. columnare*

for 75 days with the final concentration of 1×10^6 CFU/ml for 4 h at 16 ℃. Fish samples mentioned

above were taken 2 weeks after the last challenge as survivor group (fish survived 75 days after three

exposure). Both experiments were performed at least three independent times. As a control group

(mock infected), the same number of fish were maintained in similar tanks and were exposed to the

same culture medium without pathogens.

The detection of *F. columnare* in trout

To detect the *F. columnare* in rainbow trout after challenge, different tissue samples including skin,

gills and fins were collected, weighed and homogenized in tissue lysis buffer using steel beads and

shaking (60 HZ for 1 min) by TissueLyser II (Jingxin Technology). Then the DNA of different tissue

samples was extracted and amplified by PCR using the 16S rRNA specific primers of *F. columnare*.

The PCR products were extracted from a 2% agarose gel, and images were acquired using a Gel Doc

XR$^+$ (Bio-Rad, USA). Moreover, skin tissue samples were collected from both control and infected

fish and homogenized in 1 ml PBS (filtered with 0.22 μm) for coating plates, and the colonies were

observed using a fluorescence microscope.

RNA isolation and quantitative real-time PCR (qRT-PCR) analysis

Total RNA was isolated by TRIZol (Invitrogen) method as we previously described [45].

Approximately 2 μl of RNA (1000 ng in total) was reverse transcribed into cDNA with Hifair III 1st

Strand cDNA Synthesis SuperMix for qPCR (YEASEN) following the manufacturer's instructions.

And then, qRT-PCR was performed by with SYBR green qPCR mix (Monad) at 95˚C for 5 min, and

41 cycles of 95˚C 10 s, 58˚C 30 s. We calculated the relative fold changes by using the $2^{-\Delta\Delta Ct}$ method

and EF-1α was used as an internal control. The specific primers used for this experiment are list in

Table 1.

RNA-Seq libraries and differential expression analysis

Nine samples from rainbow trout was used for the RNA-Seq libraries construction and analyzed as

we previous described [76]. RNA-Seq data generated by Illumina paired-end sequencing was filtered

and only reads mapped to one site was used for further analysis. Genes with FDR $\leq$ 0.05 and |log2

(fold-change) | $\geq$ 1 were considered as the differentially expressed genes (DEGs).

Collection of serum and skin mucus

MS-222 was used to anesthetize the trout, and after that, the serum was obtained by

ultracentrifugation [10]. As for the skin mucus, we used the method described by Xu et al [9]. Briefly,

we transferred the scraping from the surface of skin into an Eppendorf tube, and then blew it

repeatedly through the injector. Subsequently, the skin mucus was collected by centrifuge. To

remove the skin bacteria, we centrifuged the previously treated skin mucus by 10,000 g

ultracentrifugation and then the skin mucus supernatant was obtained.

Western blot

Skin mucus and serum under non-reducing condition were separated by 4-15% SDS-PAGE Ready

Gel (Bio-Rad) according to previous research [7]. The samples were detected by western blot

analysis using primary anti-trout IgT (rabbit polyclonal Ab (pAb)), anti-trout IgM (mouse

monoclonal antibody (mAb)) or biotinylated anti-trout IgD (mouse mAb), respectively. The binding

of primary antibody was detected by secondary antibodies and observed by GE Amersham Imager

600 Imaging System (GE Healthcare).

Histology, light microscopy and immunofluorescence microscopy studies

The paraffin sections of tissues were obtained for histology or immunofluorescence study as previous report [76]. To detect the histological structure and mucus cells in skin, the sections were stained with H&E and AB. To observe the distribution of GFP-*F. columnare* in skin, gills and fins, the sections were stained with DAPI (4′, 6-diamidino-2-phenylindole; 1 μg/ml; Invitrogen). In the immunofluorescence study, polyclonal rabbit anti-trout IgT (0.5 μg/ml), monoclonal mouse anti-trout IgM (1 μg/ml), or polyclonal rabbit anti-trout pIgR antibody (0.5 μg/ml) were used to assess the IgT$^+$ and IgM$^+$ B cells or pIgR$^+$ cells in skin sections as previous study [9]. The rabbit and mouse IgG isotype antibodies were incubated at the same concentration as isotype controls. All images were gained by Olympus BX53 fluorescence microscope (Olympus), and processed by iVision-Mac scientific imaging processing software (Olympus).

Proliferation of B cells in the skin of trout

The proliferation of B cells in skin and head kidney were assessed in control and survivor fish by injection wih EdU (Invitrogen) as previously reported [7]. 24 hours later, the leucocytes from trout skin and head kidney were isolated for flow cytometry study using the existing methodology as described previously [7-9]. To detect IgT$^+$ or IgM$^+$ B cells, approximately 5×10^6 cells were incubated with anti-trout IgT (IgG 2b, 1 μg/ml) or mouse anti-trout IgM (IgG1, 1ug/ml), and the secondary antibodies were incubated subsequently. The EdU$^+$ cells were then stained according to the manufacturer's instructions (Click-iT EdU Alexa Fluor 647 Flow Cytometry Assay Kit, Invitrogen). Using a CytoFLEX flow cytometer (Beckman coulter) and FlowJo software (Tree Star), the percentages of EdU$^+$ cells within the IgM$^+$ or IgT$^+$ B cells populations were presented.

Tissue explants culture

Firstly, fish were treated with MS-222 for conveniently removing blood from the caudal vein. Then, the tissues of head kidney, spleen and skin (about 50 mg) were taken from each fish, and soaked with 75% ethanol 15 s to kill the surface bacteria. Thereafter, the tissues were submerged twice in PBS and moved to 400 μl prepared DMEM medium (Invitrogen). Lastly, tissues in the medium were cultured (5% CO_2, 17 ℃) for 7 days, and the tissue culture fluid was collected and stored until use.

Binding of trout immunoglobulins to *F. columnare*

Here, we measured the titers of *F. columnare*-specific Igs in fluid samples (skin mucus, serum and tissue explant medium) by pull-down assay as previously described [7]. Briefly, bacteria were incubated with fluid samples for 4 h at 4 ℃ and washed three times with PBS. Then the bound proteins were eluted with 2 × Laemmli Sample Buffer (Bio-Rad), and then analyzed by western blot as clarified above.

Statistical analysis

An unpaired Student's *t*-test (Excel version 11.0; Microsoft) and nonparametric Mann–Whitney *t*-test (Prism version 6.0; GraphPad) were used for analysis of differences between two groups. *P* values of 0.05 or less were considered statistically significant.

Acknowledgements

We thank Dr. J. Oriol Sunyer (University of Pennsylvania) for his generous gift of anti-trout IgM, anti-trout IgD, anti-trout IgT mAbs, anti-trout IgT and anti-trout pIgR pAbs.

   Figure Legends

   Fig 1. Invasion of *F. columnare* to the skin of rainbow trout.

   (A) The severe phenotype of rainbow trout was observed at 4 dpi with GFP-*F. columnare*. (B) The

   PCR products of *F. columnare* 16S rRNA gene of the fins, skin and gills tissues from three control

   and six infected fish at 4 dpi, were electrophoresed in 2% agarose and the bands were recorded using

   the gel imaging system. (C) Histological examination of GFP-*F. columnare* by fluorescence in trout

   fins, skin and gill paraffinic sections from infected fish at 4 dpi. Differential interference contrast

   (DIC) images show merged staining with *F. columnare* (green) and nuclei (blue). Scale bars, 20 μm.

   (D) The culture plates from trout skin of control fish (left, upper) and 4-day infected fish (right,

   upper), respectively. Red triangles indicate single colony of GFP-*F. columnare*. Colony image: a

   magnified view of circled colony from infected fish by fluorescence microscope (right, lower).

   Bacteria image: the observation of bacterial solution obtained by circled colony expansion (left,

   lower). (E) Localization of GFP-*F. cloumnare* in trout skin of control fish and infected fish at 1, 2, 4,

   7, and 14 dpi. Ep: epidermis, De: dermis, Sc: scale. Scale bars, 20 μm. (F) The relative expression of

   16S rRNA of *F. columnare* in infected fish versus control fish measured at days 1, 2, 4, 7, and 14

   post-infection in skin of rainbow trout ($n = 6$). Data are representative of three independent

   experiments (mean and s.e.m.).

   Fig 2. Pathological changes and immune responses in the skin of rainbow trout following *F.*

   ***columnare* infection.**

   (A) Histological examination by H & E and AB staining of skin from trout infected with *F.*

columnare at days 1, 2, 4, 7, and 14, as well as uninfected control fish ($n = 6$). The black lines with

double arrows and black triangle indicate the width of skin epidermis and the mucus secreting cells,

respectively. Ep: epidermis, De: dermis, Sc: scale, Me: melanophores. Scale bars, 20 μm. (B) The

width of skin epidermis of control and bacterial infected fish counted from A (upper) ($n = 6$). (C)

The number of mucus cells per millimeter in the epidermis of trout skin from control and bacterial

infected fish counted from A (lower) ($n = 6$). (D) Heat map illustrates results from qRT-PCR of

mRNA expression levels for selected immune markers in *F. columnare*-infected fish versus control

fish measured at 1, 2, 4, 7, and 14 dpi in the skin ($n = 6$). Color value: fold change. (E) Volcano plot

showed the overlap of genes upregulated or downregulated in the skin of rainbow trout at days 2 (left)

and 14 (right) infected with *F. columnare* versus control fish. Red spots: expression fold change of >

2 and FDR of < 0.05; Green bots: expression fold change of < 2 and FDR of < 0.05; Black spots: no

difference in expression. (F) Venn diagrams showed the statistical results of DEGs including

up-regulated and down-regulated genes and the percentage of immune genes after the DEGs filtered

by rainbow trout immune genes libraries in the trout skin at days 2 and 14 infected with *F. columnare*

versus control fish. (G and H) Representative innate (G) and adaptive (H) immune genes modulated

by *F. columnare* infection at 2 and 14 dpi ($n = 9$). (I) Biological processes that were significantly

altered of rainbow trout at days 2 and 14 infected with *F. columnare* versus control fish revealed by

RNA-Seq studies. Fold change differences between control and bacterial infected samples were

calculated using cutoff of two-fold. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ (unpaired Student's *t*-test).

Data are representative of three independent experiments (mean and s.e.m.).

Fig 3. Bacterial load of the skin from survivor fish upon *F. columnare* re-infection.

(A) Histological examination by H & E and AB staining of skin from fish infected with 28 days and

survivor fish ($n = 6$). The black lines with double arrows and black triangle indicate the width of skin

epidermis and the mucus secreting cells, respectively. Ep: epidermis, De: dermis, Sc: scale, Me:

melanophores. Scale bars, 20 μm. (B) The width of skin epidermis of control fish, infected fish and

survivor fish counted from A (upper) ($n = 6$). (C) The number of mucus cells per millimeter in the

epidermis of trout skin from control , infected and survivor fish counted from A (lower) ($n = 6$). (D)

DIC images of fluorescence analysis showed the GFP-*F. columnare* location and load in the skin

from control and control-challenge fish (left), 28-day infected and 28-day infected-challenge fish

(middle), survivor fish and survivor-challenge fish (right). These challenge fish were sampling at 4

days post re-infection with GFP-*F. columnare* ($n = 6$). Ep: epidermis, Sc: scale, De: dermis. Scale

bars, 20 μm. (E) The relative expression of 16S RNA of *F. columnare* in the trout skin tissue of

control, control-challenge, infected, infected-challenge, survivor and survivor-challenge group were

measured by qRT-PCR ($n = 6$ fish per group). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ (unpaired

Student's *t*-test). Data are representative of at least three independent experiments (mean and s.e.m.).

Fig 4. Accumulation and proliferation of IgT[+] B cells in the skin of rainbow trout following *F.*

***cloumnare* infection.**

(A) DIC images of immunofluorescence staining on trout skin paraffinic sections from control fish,

infected and survivor fish stained for IgT (green) and IgM (red); nuclei are stained with DAPI (blue)

(isotype-matched control antibody staining, S5A Fig). Ep: epidermis, Sc: scale, De: dermis. Scale

bars, 20 μm. (B) The number of IgT[+] and IgM[+] B cells in skin paraffin sections of control, infected

and survivor fish infected with *F. columnare* are counted ($n = 12$, original magnification, ×20). *$P <$

0.05, **$P < 0.01$, ***$P < 0.001$ (one-way ANOVA with Bonferroni correction). (C) Representative

flow cytometry dot plot showing proliferation of IgT$^+$ B cells and IgM$^+$ B cells in skin leucocytes of

control and survivor fish ($n = 12$). (D) Percentage of EdU$^+$ cells from the total skin IgT$^+$ or IgM$^+$

B-cell populations in control and survivor fish ($n = 12$). (E) Representative flow cytometry dot plot

showing proliferation of IgT$^+$ B cells and IgM$^+$ B cells in head kidney leucocytes of control and

survivor fish. (F) Percentage of EdU$^+$ cells from the total head kidney IgT$^+$ or IgM$^+$ B-cell

populations in control or survivor fish ($n = 12$). *$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$ (unpaired

Student's t-test) in D and F. Data are representative of at least three independent experiments (mean

and s.e.m.).

Fig 5. Ig responses in the skin mucus and serum from infected and survived fish.

(A) Concentration of IgT, IgM, and IgD in skin mucus from control, infected, and survivor fish ($n =$

12). (B) Western blot analysis of IgT-, IgM- and IgD-specific binding to *F. columnare* in skin mucus

(dilution 1:2) from infected and survivor fish. (C and D) IgT-, IgM- and IgD-specific binding to *F.*

columnare in dilutions of skin mucus from infected (C) and survivor (D) fish evaluated by

densitometric analysis of immunoblots and presented as relative values to those of control fish ($n =$

12-18). (E) Concentration of IgT, IgM, and IgD in serum from control, infected, and survivor fish (n

$= 12$). (F) Western blot analysis of IgT-, IgM- and IgD- specific binding to *F. columnare* in serum

(dilution 1:10) from infected and survivor fish. (G and H) IgT-, IgM- and IgD- specific binding to *F.*

columnare in dilutions of serum from infected (G) and survivor (H) fish, evaluated by densitometric

analysis of immunoblots and presented as relative values to those of control fish ($n = 12$-18). *$P <$

0.05, **$P < 0.01$, ***$P < 0.001$ (unpaired Student's t-test). Data are representative of at least three

independent experiments (mean and s.e.m.).

Fig 6. Local IgT-, IgM- and IgD-specific responses in skin explants of survivor fish.

(A-C) Skin, head kidney, and spleen explants (~ 50 mg each) from control and survivor fish were

cultured in medium (400 µl) for 7 days. Immunoblot analysis of IgT-, IgM- and IgD- specific

binding to *F. cloumnare* in the culture medium of skin (A), head kidney (B), and spleen (C) (dilution

1:2) from control and survivor fish. (D-F) IgT-, IgM- and IgD- specific binding to *F. cloumnare* in

dilutions of culture medium from skin (D), head kidney (E), and spleen (F) from control and survivor

fish, measured by densitometric analysis of immunoblots and presented as relative values to those of

control fish ($n = 12$). *$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$ (unpaired Student's *t*-test). Data are

representative of at least three independent experiments (mean and s.e.m.).

Fig 7. Accumulation of pIgR and pIgR⁺ cells in the skin of rainbow trout infected with *F. columnare*.

(A) DIC images of immunofluorescence staining on trout skin paraffinic sections from control fish,

infected, and survival fish, stained for pIgR (green) and nuclei with DAPI (blue) ($n = 12$)

(isotype-matched control antibodies for anti-pIgR in S5B Fig). Ep: epidermis, De: dermis, Sc: scale.

Scale bars, 20 µm. (B) The number of pIgR⁺ cells in skin paraffin sections of control, infected, and

survived fish are counted ($n = 12$, original magnification, ×20). (C) Relative mRNA expression of

pIgR in the skin of control fish, infected, and survival fish were detected by qRT-PCR ($n = 6$). (D)

Immunoblot analysis of pIgR concentration in skin mucus from control, infected and survivor fish

following *F. columnare* infection. (E) Relative concentration of pIgR from infected and survivor

trout skin mucus to that of control fish ($n = 12$). *$P < 0.05$, **$P < 0.01$ (unpaired Student's t-test).

Data are representative of at least three independent experiments (mean and s.e.m.).

Fig 8. Proposed model of local IgT and IgT$^+$ B cell induction in the skin after *F. columnare*

infection.

The presumed structure of skin displaying mucus layer, epithelia layer and dermis. The proposed

model contains two partitions: control (left) and infected/survivor (right). Induction of local IgT

responses in the trout skin based on our findings. On the left is the scheme of a typical skin structure

in control (naive) teleost fish. The number of IgT$^+$ B cells in control fish skin is low. IgT are

produced by IgT-secreting B cells and transported from the epithelium into the skin mucus layer via

pIgR. The secreted IgT coats the majority of microbiota in the skin surface. In the skin epidermis,

abundant mucus cells are present, and their production contains antibacterial molecules to protect

host against pathogens. Upon *F. columnare* invasion, their antigen (Ag) can be taken up by

antigen-presenting cells (APC) and presented to naive CD4$^+$ T cells. Then IgT$^+$ B cells are activated

by Ag-specific CD4$^+$ T cells and start to proliferate locally and differentiate into plasma cells (PCs)

in skin epithelial layer. These differentiated IgT-secreting cells can produce large amounts of

pathogen-specific IgT, which can be transported by pIgR into skin mucus layer where those

pathogen-specific IgT can specifically binding to the bacteria *F. columnare*. Alternatively, some IgT$^+$

plasma cells may differentiate further into memory IgT$^+$ B cells. When *F. columnare* invade the host

again, the memory IgT$^+$ B cells would directly proliferate and differentiate into plasma cells, and

then rapid produce specific-IgT to binding *F. columnare*.

667
668

669   Fig 1

670
671

672 Fig 2

675 **Fig 3**

F.columnare Nuclei

Fig 4

680 Fig 5

Fig 6

684   Fig 7

687   Fig 8

Control fish

Local immune response in trout skin to
F.columnare (Infercted/survival fish)

Mucus layer

Epidermis

B-cell
activation

Differentiation
to plasm cells

Ag
presentation

B-cell
proliferation

Differentiation
to memory
cells

Melanophore

	Epithelial cells		Commensal bacteria		Plasma cells		Cytokines & chemokines
	Squamous epithelial cells		Basal cells		Memory B cell		MHC/TCR/Ag complex
	Mucus cell		Granulocyte		pIgR		*Flavobacterium columnare*
	Macrophage	T	T cell		Cell surface IgT		Virulence factors
	Dendritic cell	B	IgT⁺ B cell		Polymeric IgT		Scale
		B	Naive B cell		Polymeric IgT with tSC		Melanophore

Supplemental information

S1 Fig. Physical and immunological barrier of the skin.

 Structures of human (left) and fish (right) skin. In mammal, there are three layers in skin including stratum corneum, stratum granulosum, and stratum spinosum in the epidermis. Tight junctions are formed in the stratum granulosum, and specialized dendritic cells (DCs) known as Langerhans cells and $CD8^+$ T cells reside between keratinocytes. Specialized immune cells including antigen-presenting dermal DCs, macrophages, neutrophil and $CD4^+$ T cells reside in the dermis. In contrast, there are live cells instead of stratum corneum with large abundant of mucus covering on the fish skin surface, and numerous microbiota living there. Interestingly, in the epidermis of teleost, various types of cells including immune cells such as B cells, T cells, DCs, macrophages, granulocytes and mucus cells have been identified. Differently, fish dermis is mainly composed of dense connective tissue with a large amount of collagen fibers.

 S2 Fig. Construction of GFP-*F. columnare* G$_4$ strain.

 (A) Map of the plasmid pCP23-gfp. (B) Growth curves for wild-type *F. columnare* and the GFP-*F.* *columnare*. (C) Survival curves of fish infected with wild-type G$_4$ and the GFP-*F. columnare* ($n = 30$). Data are representative of at least three independent experiments (mean and s.e.m.).

 S3 Fig. Immune responses in trout spleen and head kidney following *F. columnare* infection.

 (A and B) Heat map illustrates results from qRT-PCR of mRNAs for selected immune markers in *F.* *columnare*-infected fish versus control fish measured at 1, 2, 4, 7, and 14 dpi in trout head kidney (A)

and spleen (B) (*n* = 6). Data are representative of three different independent experiments (mean and

s.e.m.).

S4 Fig. Validation of transcriptomic data by qPCR.

(A-D) Transcriptomic differential expressed genes in experimental groups at 2 (A and B) and 14

days (C and D) post infection were randomly chosen and detected using qPCR to validate RNA-seq.

Positive numbers in the Y axis mean up-regulated, while negative values mean down-regulated (*n* =

6). Data are representative of three different independent experiments (mean and s.e.m.).

S5 Fig. Isotype control staining for anti-IgT, anti-IgM and anti-pIgR antibodies in trout skin

paraffin sections.

(A and B) Differential interference contrast images of skin paraffin sections from control fish, with

merged staining of isotype control antibodies for anti-trout IgT pAb (green) and anti-trout IgM mAbs

(red) (A); or for anti-trout pIgR pAb (green) (B). Nuclei were stained with DAPI (blue). Ep:

epidermis, Sc: scale, Me: melanophores. Scale bars, 20 μm. Data are representative of at least three

different independent experiments.

S1 Table. List of primers for real-time quantitative PCR amplifications.

S1 Fig

730   S2 Fig

734   S3 Fig

A

B

738   S4 Fig

742   S5 Fig

A

B

745 S1 Table

Gene	GenBank accession no.	Primer Sequence (5′-3′)	Tm (°C)	Amplicon Length (bp)
IMMUNOGLOBULINS AND THEIR RECEPTOR				
IgT	AY870265.1	F: TGTCCGTCCACATTCTC	58	169
		R: TTGCCTTTCTTGGTTTT		
IgM	EF438413.1	F: CTATGGAATAGGCTGGAT	58	137
		R: TGTTGCTGGAAGTGTCTC		
IgD	AY870261.1	F: CATCCTGAGTGGGTTCT	58	219
		R: GTAGGTGGTCCTTGTGC		
pIgR	XM_021599266.1	F: TGTTACACTCCGCATTCTC	58	253
		R: CAGGGCAGGTTTCTGATTT		
COMPLEMENT AND PATHWAY RELATED GENES				
C1R	NM_001124380.1	F: GCCGACTGCTACTACGA	58	240
		R: GGTGCCTATGTGCTCTATG		
C3	XM_021561577.1	F: CCTCACAACAAGAGTGCACATC	58	213
		R: CCAAGTGGGCAAACTCATCTCC		
C7-1	NM_001124618.1	F: AACGAGCATCTCCAATA	58	218
		R: TGCCACCTCTTTACTTC		
C1q-2	XM_021624859.1	F: TGAACAACCTGAACACCCC	58	176
		R: CAGCCTATTAGCCTGTAACTCC		
CYTOKINES				
Interleukin 8	NM_001124362.1	F: TGTCGTTGTGCTCCTGG	58	197
		R: CCTGACCGCTCTTGCTC		
Interleukin 10	NM_001245099.1	F: CACCGCCTTCTCCACCATC	58	101
		R: CCATAGCGTGACACCCCAC		
Interleukin 1β	NM_001124347.2	F: TGATGAATGAGGCTATGGA	58	216
		R: GATGGTGAAGGTGGTAAGG		
TNF α	XM_021578986.1	F: CAGAGCCTCAGGAAGAACT	58	180
		R: CAAGGGGTAGAAGAGCATA		
IRF 4	XM_021604510.1	F: CGCCCTACGGAGATAAAC	58	135
		R: TCAGCAGCACCTGGAGAC		
CCL 19	XM_021605155.1	F: GTTTCCCTCGCCACTTCAA	58	139
		R: GCCACCCACTTGCTCTTTG		
FLAVOBACTERIUM COLUMNARE				
16S rRNA	EU395796.1	F: GAGTGGCTAAGCGAAAGTGAT	58	146
		R: ACCTGACACCTCACGGCAC		
REFERENCE GENE				
EF-1α	XM_021571866.1	F: CAACGATATCCGTCGTGGCA	58	
		R: ACAGCGAAACGACCAAGAGG		
OTHERS				
CATH-1	NM_001124480.1	F: CTGGAGGCAAGCAACAAC	58	152
		R: CCCCCAAGACGAGAGACA		

Gene	Accession	Primer sequences (5'→3')	Tm	Size
CXCL10	XM_021622222.1	F: ACATCAACGGTCCTCATC R: ACACTTCTTCCCTTCTCC	58	216
NOS2	XM_021581479.1	F: GGCAGTCAAGAACCAACC R: GAGCACCAAACGCTAATT	58	254
HP	XM_021607150.1	F: CGGAGGAGGTTGGAAGC R: GCAGCAGAAGCCACAGC	58	221
SAA	XM_021607573.1	F: TTGTTCTGACCCTCGTTG R: CCTGGCAGCATCATAGTT	58	167
MUC-2	XM_021572851.1	F: GGAATCGTGGTGGACATG R: AGGCTAGGTAGCGAAAGG	58	177
PRDM1	XM_021592048.1	F: GGCAAGGAGGACCAGAG R: CCGGTTGTAGAGGCGTA	58	144
CD209	NM_001124633.1	F: GCCCTCTTTGTGCTGATT R: ACTCCCTACACTTCCTTA	58	214
MCOLN1	XM_021625006.1	F: TCATCAAGATTGGCATAG R: GGATTACATTAGGGAAGG	58	171
HSPA1S	XM_021567143.1	F: GATGGACAAGGCTCAGG R: CGTCAAGGAGGAGAAGGT	58	212
NPPC	XM_021601323.1	F: GGTGGCTTGTGGACTTAT R: GGTCTCGCAGCAATCGTA	58	182
ZNRF1	XM_021568721.1	F: ACGGCTACCAGGAAACAG R: TCATTGGACGCCACAGAC	58	166
col1a3	NM_001124206.1	F: GCAACAGCCGCTTCACA R: TTCGACGCCAAACTCCT	58	167
THBS2S	XM_021557617.1	F: ACATCAACGGTCCTCATC R: ACACTTCTTCCCTTCTCC	58	218
COL2A	XR_002469112.1	F: GTCTAAATGCTAACCCAT R: GTGTAGTGTTGATTCCCT	58	190
DCN	XM_021562199.1	F: TCGGAGCCTGATAACAAT R: ACAGATGCGGTGGTAGAG	58	117
CREB3	XM_021585854.1	F: TTGGATGTGGCTGGATTT R: TGAGGCGTGATTTGGTGT	58	251
ITGB1	XM_021620606.1	F: CAGCGAAGACCTGGATAA R: CAGAGTTTGTTGTTGGAGC	58	180
ANGPT1	XM_021587370.1	F: CCCACTATGACCGTTTCT R: ATCCCATTCAGTTTACCC	58	262
GAPDH	NM_001124209.1	F: GGCAGCGTTTGAGGAGATGA R: CATGGCGTCCAGTAGCTTGG	58	100
ENO	XM_021578347.1	F: TCAAGTCACCTGACGACCCT R: ACAGCCTGCTGGATACGC	58	214
ALDO	XM_021575432.1	F: GCATTGGTGGTGTCATCTTC R: TGCCTACAACCCAGCCTCT	58	102
LDH	XM_021564046.1	F: ACTCCAGCGTGCCCGTAT R: CGTGCTTCCAGTCCTCCTT	58	105

| PFKA | XM_021609920.1 | F: CGACACCCGTTCCACAAT | 58 | 103 |
| | | R: AGCCATCACAGCCTCCAC | | |

References

1. Fillatreau S, Six A, Magadan S, Castro R, Sunyer JO, Boudinot P. The astonishing diversity of Ig classes and B cell repertoires in teleost fish. Front Immunol. 2013;4:28. pmid: 23408183

2. Salinas I, Zhang YA, Sunyer JO. Mucosal immunoglobulins and B cells of teleost fish. Dev Comp Immunol. 2011;35(12):1346-1365. pmid: 22133710

3. Zapata A, Diez B, Cejalvo T, Gutiérrez-de Frías C, Cortés A. Ontogeny of the immune system of fish. Fish Shellfish Immunol. 2006;20(2):126-136. pmid: 15939627

4. Bromage ES, Kaattari IM, Zwollo P, Kaattari SL. Plasmablast and plasma cell production and distribution in trout immune tissues. J Immunol. 2004;173(12):7317-7323. pmid: 15585855

5. Solem ST, Stenvik J. Antibody repertoire development in teleosts-a review with emphasis on salmonids and *Gadus morhua* L. Dev Comp Immunol. 2006;30(1-2):57-76. pmid: 16084588

6. Perdiguero P, Martín-Martín A, Benedicenti O, Díaz-Rosales P, Morel E, Muñoz-Atienza E, et al. Teleost IgD$^+$IgM$^-$ B cells mount clonally expanded and mildly mutated intestinal IgD responses in the absence of lymphoid follicles. Cell Rep. 2019;29(13):4223-4235.e4225. pmid: 31875534

7. Xu Z, Takizawa F, Parra D, Gómez D, von Gersdorff Jørgensen L, LaPatra SE, et al. Mucosal immunoglobulins at respiratory surfaces mark an ancient association that predates the emergence of tetrapods. Nat Commun. 2016;7:10728. pmid: 26869478

8. Yu YY, Kong W, Yin YX, Dong F, Huang ZY, Yin GM, et al. Mucosal immunoglobulins protect the olfactory organ of teleost fish against parasitic infection. PLoS Pathog. 2018;14(11):e1007251. pmid: 30395648

9. Xu Z, Parra D, Gómez D, Salinas I, Zhang YA, von Gersdorff Jørgensen L, et al. Teleost skin, an ancient mucosal surface that elicits gut-like immune responses. Proc Natl Acad Sci U S A. 2013;110(32):13097-13102. pmid: 23884653

10. Zhang YA, Salinas I, Li J, Parra D, Bjork S, Xu Z, et al. IgT, a primitive immunoglobulin class specialized in mucosal immunity. Nat Immunol. 2010;11(9):827-835. pmid: 20676094

11. Yu YY, Kong WG, Xu HY, Huang ZY, Zhang XT, Ding LG, et al. Convergent evolution of mucosal immune responses at the buccal cavity of teleost fish. iScience. 2019;19:821-835. pmid: 31499339

12. Kong WG, Yu YY, Dong S, Huang ZY, Ding LG, Cao JF, et al. Pharyngeal immunity in early vertebrates provides functional and evolutionary insight into mucosal homeostasis. J Immunol. 2019;203(11):3054-3067. pmid: 31645417

13. Xu Z, Takizawa F, Casadei E, Shibasaki Y, Ding Y, Sauters TJC, et al. Specialization of mucosal immunoglobulins in pathogen control and microbiota homeostasis occurred early in vertebrate evolution. Sci Immunol. 2020;5(44). pmid: 32034088

14. Zhang J, Taniguchi T, Takita T, Ali AB. A study on the epidermal structure of

Periophthalmodon and *Periophthalmus* mudskippers with reference to their terrestrial adaptation. Ichthyological Research. 2003;50(4):310-317. DOI: 10.1007/s10228-003-0173-7

15. Streilein JW. Circuits and signals of the skin-associated lymphoid tissues (SALT). J Invest Dermatol. 1985;85(1 Suppl):10s-13s. pmid: 4008979

16. Kabashima K, Honda T, Ginhoux F, Egawa G. The immunological anatomy of the skin. Nat Rev Immunol. 2019;19(1):19-30. pmid: 30429578

17. Tong PL, Roediger B, Kolesnikoff N, Biro M, Tay SS, Jain R, et al. The skin immune atlas: three-dimensional analysis of cutaneous leukocyte subsets by multiphoton microscopy. J Invest Dermatol. 2015;135(1):84-93. pmid: 25007044

18. Ángeles Esteban M, Cerezuela R. 4 - Fish mucosal immunity: skin. In: Beck BH, Peatman E, editors. Mucosal Health in Aquaculture. San Diego: Academic Press. 2015. p. 67-92.

19. Ángeles Esteban M. An overview of the immunological defenses in fish skin. ISRN Immunology. 2012;2012. DOI: 10.5402/2012/853470

20. Brandtzaeg P. Mucosal immunity: induction, dissemination, and effector functions. Scand J Immunol. 2009;70(6):505-15. pmid: 19906191

21. Wright PF. Inductive/effector mechanisms for humoral immunity at mucosal sites. Am J Reprod Immunol. 2011;65(3):248-52. pmid: 21223423

22. Saitoh-Inagawa W, Hiroi T, Yanagita M, Iijima H, Uchio E, Ohno S, et al. Unique characteristics of lacrimal glands as a part of mucosal immune network: high frequency of IgA-committed B-1 cells and NK1.1$^+$ alphabeta T cells. Invest Ophthalmol Vis Sci. 2000;41(1):138-44. pmid: 10634613

23. Brandtzaeg P. Secretory immunity with special reference to the oral cavity. J Oral Microbiol. 2013;5. pmid: 23487566

24. Bernardet J-F, Segers P, Vancanneyt M, Berthe F, Kersters K, Vandamme PJIJoS, et al. Cutting a Gordian knot: emended classification and description of the genus *Flavobacterium*, emended description of the family *Flavobacteriaceae*, and proposal of *Flavobacterium hydatis* nom. nov. (basonym, *Cytophaga aquatilis* Strohl and Tait 1978). 1996;46(1):128-148.

25. Plumb J. Health maintenance and culture of microbial diseases of cultured fishes. Iowa State University Press, Ames. 1999.

26. Bernardet J-F, Bowman J. The prokaryotes: a handbook on the biology of bacteria. *Proteobacteria*: Delta and Epsilon Subclasses Deeply Rooted Bacteria. 2006;7:481-532.

27. Decostere A, Haesebrouck F, Devriese LA. Characterization of four *Flavobacterium columnare* (*Flexibacter columnaris*) strains isolated from tropical fish. Vet Microbiol. 1998;62(1):35-45. pmid: 9659690

28. Figueiredo HC, Klesius PH, Arias CR, Evans J, Shoemaker CA, Pereira DJ, Jr., et al. Isolation and characterization of strains of *Flavobacterium columnare* from Brazil. J Fish Dis. 2005;28(4):199-204. pmid: 15813861

29. Morley NJ, Lewis JW. Consequences of an outbreak of columnaris disease (*Flavobacterium columnare*) to the helminth fauna of perch (*Perca fluviatilis*) in the Queen Mary reservoir, south-east England. J Helminthol. 2010;84(2):186-192. pmid: 19735594

30. Řehulka J, Minařík B. Blood parameters in brook trout *Salvelinus fontinalis* (Mitchill, 1815), affected by columnaris disease. Aquaculture Research. 2007;38(11):1182-1197. DOI: 10.1111/j.1365-2109.2007.01786.x

31. Soto E, Mauel MJ, Karsi A, Lawrence ML. Genetic and virulence characterization of *Flavobacterium columnare* from channel catfish (*Ictalurus punctatus*). J Appl Microbiol. 2008;104(5):1302-1310. pmid: 18005349

32. Suomalainen LR, Bandilla M, Valtonen ET. Immunostimulants in prevention of columnaris disease of rainbow trout, *Oncorhynchus mykiss* (Walbaum). J Fish Dis. 2009;32(8):723-726. pmid: 19508505

33. Declercq AM, Haesebrouck F, Van den Broeck W, Bossier P, Decostere A. Columnaris disease in fish: a review with emphasis on bacterium-host interactions. Vet Res. 2013;44(1):27. pmid: 23617544

34. Li N, Zhu Y, LaFrentz BR, Evenhuis JP, Hunnicutt DW, Conrad RA, et al. The Type IX Secretion System Is Required for Virulence of the Fish Pathogen *Flavobacterium columnare*. Appl Environ Microbiol. 2017;83(23). pmid: 28939608

35. Moor K, Diard M, Sellin ME, Felmy B, Wotzka SY, Toska A, et al. High-avidity IgA protects the intestine by enchaining growing bacteria. Nature. 2017;544(7651):498-502. pmid: 28405025

36. Baldissera MD, Souza CF, Dias JB, Da Silva TO, Tavares GC, Valladao GMR, et al. Branchial bioenergetics dysfunction as a relevant pathophysiological mechanism in freshwater silver catfish (*Rhamdia quelen*) experimentally infected with *Flavobacterium columnare*. Microb Pathog. 2019;138:103817. pmid: 31672529

37. Byrd AL, Belkaid Y, Segre JA. The human skin microbiome. Nat Rev Microbiol. 2018;16(3):143-55. pmid: 29332945

38. Belkaid Y, Tamoutounour S. The influence of skin microorganisms on cutaneous immunity. Nat Rev Immunol. 2016;16(6):353-66. pmid: 27231051

39. Gomez D, Sunyer JO, Salinas I. The mucosal immune system of fish: the evolution of tolerating commensals while fighting pathogens. Fish Shellfish Immunol. 2013;35(6):1729-39. pmid: 24099804

40. Yin X, Mu L, Fu S, Wu L, Han K, Wu H, et al. Expression and characterization of Nile tilapia (*Oreochromis niloticus*) secretory and membrane-bound IgM in response to bacterial infection. Aquaculture. DOI: 10.1016/j.aquaculture.2019.03.058

41. Castro R, Abós B, González L, Granja AG, Tafalla C. Expansion and differentiation of IgM$^+$ B cells in the rainbow trout peritoneal cavity in response to different antigens. Dev Comp Immunol. 2017;70:119-27. pmid: 28088353

42. Declercq AM, Chiers K, Haesebrouck F, Van den Broeck W, Dewulf J, Cornelissen M, et al.

Gill infection model for columnaris disease in common carp and rainbow trout. J Aquat Anim Health. 2015;27(1):1-11. pmid: 25488182

43. Austin DA, Jordan EM, Austin B. Recovery of an unusual Gram-negative bacterium from ulcerated rainbow trout, *Oncorhynchus mykiss* (Walbaum), in Scotland. J Fish Dis. 2003;26(4):247-9. pmid: 12962233

44. Declercq AM, Chiers K, Van den Broeck W, Dewulf J, Eeckhaut V, Cornelissen M, et al. Interactions of highly and low virulent *Flavobacterium columnare* isolates with gill tissue in carp and rainbow trout. Vet Res. 2015;46:25. pmid: 25889257

45. Tongsri P, Meng K, Liu X, Wu Z, Yin G, Wang Q, et al. The predominant role of mucosal immunoglobulin IgT in the gills of rainbow trout (*Oncorhynchus mykiss*) after infection with *Flavobacterium columnare*. Fish Shellfish Immunol. 2020;99:654-62. pmid: 32001351

46. Xie HX, Nie P, Chang MX, Liu Y, Yao WJ. Gene cloning and functional analysis of glycosaminoglycan-degrading enzyme chondroitin AC lyase from *Flavobacterium columnare* G_4. Arch Microbiol. 2005;184(1):49-55. pmid: 16044262

47. Katzenback BA. Antimicrobial peptides as mediators of innate immunity in teleosts. Biology (Basel). 2015;4(4):607-39. pmid: 26426065

48. Wang TT, Song XH, Bao GM, Zhao LX, Yu X, Zhao J. Molecular characterization, expression analysis, and biological effects of interleukin-8 in grass carp *Ctenopharyngodon idellus*. Fish Shellfish Immunol. 2013;35(5):1421-32. pmid: 23994423.

49. Peng Y, Cai X, Zhang G, Wang J, Li Y, Wang Z, et al. Molecular characterization and expression of interleukin-10 and interleukin-22 in golden pompano (*Trachinotus ovatus*) in response to *Streptococcus agalactiae* stimulus. Fish Shellfish Immunol. 2017;65:244-55. pmid: 28442416.

50. Hong S, Li R, Xu Q, Secombes CJ, Wang T. Two types of TNF-α exist in teleost fish: phylogeny, expression, and bioactivity analysis of type-II TNF-α3 in rainbow trout *Oncorhynchus mykiss*. J Immunol. 2013;191(12):5959-72. pmid: 24244011

51. Bose M, Farnia P. Proinflammatory cytokines can significantly induce human mononuclear phagocytes to produce nitric oxide by a cell maturation-dependent process. Immunol Lett. 1995;48(1):59-64. pmid: 8847093

52. Tamura S, Iwasaki T, Thompson AH, Asanuma H, Chen Z, Suzuki Y, et al. Antibody-forming cells in the nasal-associated lymphoid tissue during primary influenza virus infection. J Gen Virol. 1998;79 (Pt 2):291-299. pmid: 9472613

53. Fagarasan S. Intestinal IgA synthesis: a primitive form of adaptive immunity that regulates microbial communities in the gut. Curr Top Microbiol Immunol. 2006;308:137-153. pmid: 16922089

54. Cerutti A, Rescigno M. The biology of intestinal immunoglobulin A responses. Immunity. 2008;28(6):740-750. pmid: 18549797

55. Onodera T, Takahashi Y, Yokoi Y, Ato M, Kodama Y, Hachimura S, et al. Memory B cells in the

lung participate in protective humoral immune responses to pulmonary influenza virus reinfection. Proc Natl Acad Sci U S A. 2012;109(7):2485-2490. pmid: 22308386

56. Milligan GN, Bernstein DI. Generation of humoral immune responses against herpes simplex virus type 2 in the murine female genital tract. Virology. 1995;206(1):234.

57. Uddin T, Harris JB, Bhuiyan TR, Shirin T, Uddin MI, Khan AI, et al. Mucosal immunologic responses in cholera patients in Bangladesh. Clin Vaccine Immunol. 2011;18(3):506-12. pmid: 21248157

58. Johnson RA, Uddin T, Aktar A, Mohasin M, Alam MM, Chowdhury F, et al. Comparison of immune responses to the O-specific polysaccharide and lipopolysaccharide of *Vibrio cholerae* O1 in Bangladeshi adult patients with cholera. Clin Vaccine Immunol. 2012;19(11):1712-21. pmid: 22993410

59. Zilberg D, Klesius PH. Quantification of immunoglobulin in the serum and mucus of channel catfish at different ages and following infection with *Edwardsiella ictaluri*. Vet Immunol Immunopathol. 1997;58(2):171-180. pmid: 9336885

60. Coscia MR, Simoniello P, Giacomelli S, Oreste U, Motta CM. Investigation of immunoglobulins in skin of the Antarctic teleost *Trematomus bernacchii*. Fish Shellfish Immunol. 2014;39(2):206-214. pmid: 24821425

61. Guardiola FA, Cuesta A, Arizcun M, Meseguer J, Esteban MA. Comparative skin mucus and serum humoral defence mechanisms in the teleost gilthead seabream (*Sparus aurata*). Fish Shellfish Immunol. 2014 Feb;36(2):545-51. pmid: 24412437

62. Valdenegro-Vega VA, Crosbie P, Vincent B, Cain KD, Nowak BF. Effect of immunization route on mucosal and systemic immune response in Atlantic salmon (*Salmo salar*). Vet Immunol Immunopathol. 2013 Jan 15;151(1-2):113-23. pmid: 23206403

63. Piazzon MC, Galindo-Villegas J, Pereiro P, Estensoro I, Calduch-Giner JA, Gómez-Casado E, Novoa B, Mulero V, Sitjà-Bobadilla A, Pérez-Sánchez J. Differential modulation of IgT and IgM upon parasitic, bacterial, viral, and dietary challenges in a perciform fish. Front Immunol. 2016 Dec 27;7:637. pmid: 28082977

64. Xu J, Yu Y, Huang Z, Dong S, Luo Y, Yu W, Yin Y, Li H, Liu Y, Zhou X, Xu Z. Immunoglobulin (Ig) heavy chain gene locus and immune responses upon parasitic, bacterial and fungal infection in loach, *Misgurnus anguillicaudatus*. Fish Shellfish Immunol. 2019 Mar;86:1139-1150. pmid: 30599252

65. Makesh M, Sudheesh PS, Cain KD. Systemic and mucosal immune response of rainbow trout to immunization with an attenuated *Flavobacterium psychrophilum* vaccine strain by different routes. Fish Shellfish Immunol. 2015;44(1):156-63. pmid: 25687393

66. Woof JM, Kerr MA. The function of immunoglobulin A in immunity. J Pathol. 2006;208(2):270-282. pmid: 16362985

67. Lamm ME. Interaction of antigens and antibodies at mucosal surfaces. Annu Rev Microbiol. 1997;51:311-340. pmid: 9343353

68. Randall TD. Bronchus-associated lymphoid tissue (BALT) structure and function. Adv
Immunol. 2010;107:187-241. pmid: 21034975

69. Mostov KE. Transepithelial transport of immunoglobulins. Annu Rev Immunol. 1994;12:63-84.
pmid: 8011293

70. Hooper LV, Wong MH, Thelin A, Hansson L, Falk PG, Gordon JI. Molecular analysis of
commensal host-microbial relationships in the intestine. Science. 2001;291(5505):881-884.
pmid: 11157169

71. Pal K, Kaetzel CS, Brundage K, Cunningham CA, Cuff CF. Regulation of polymeric
immunoglobulin receptor expression by reovirus. J Gen Virol. 2005;86(Pt 8):2347-57. pmid:
16033983.

72. Bruno ME, Rogier EW, Frantz AL, Stefka AT, Thompson SN, Kaetzel CS. Regulation of the
polymeric immunoglobulin receptor in intestinal epithelial cells by *Enterobacteriaceae*:
implications for mucosal homeostasis. Immunol Invest. 2010;39(4-5):356-82. pmid: 20450283.

73. Sunyer JO. Fishing for mammalian paradigms in the teleost immune system. Nat Immunol.
2013;14(4):320-6. pmid: 23507645

74. Li N, Qin T, Zhang XL, Huang B, Liu ZX, Xie HX, et al. Gene deletion strategy to examine the
involvement of the two chondroitin lyases in *Flavobacterium columnare* virulence. Appl
Environ Microbiol. 2015;81(21):7394-7402. pmid: 26253667

75. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic
Acids Res. 2001;29(9):e45. pmid: 11328886

76. Zhang X, Ding L, Yu Y, Kong W, Yin Y, Huang Z, et al. The change of teleost skin commensal
microbiota is associated with skin mucosal transcriptomic responses during parasitic infection
by *Ichthyophthirius multifillis*. Front Immunol. 2018;9:2972. pmid: 30619329