Identification of microRNAs that stabilize p53 in HPV-positive cancer cells

Gustavo Martínez-Noël [a], Patricia Szajner [a], Rebecca E. Kramer [a], Kathleen A. Boyland [a], Asma Sheikh [a], Jennifer A. Smith [a,b], and Peter M. Howley [a#]

[a]Department of Immunology, Harvard Medical School, 77 Avenue Louis

Pasteur, Boston MA 02115

[b]ICCB-Longwood Screening Facility, Harvard Medical School, 250 Longwood Avenue,

Boston MA 02115

Running Title: HPV p53 miRNA Stabilization Screen

Abstract word count: 250

Text word count: 6030

#Corresponding author: Peter M. Howley , peter_howley@hms.harvard.edu

Gustavo Martínez-Noël and Patricia Szajner contributed equally to this work.

Abstract

Etiologically, 5% of all cancers worldwide are caused by the high-risk human papillomaviruses (hrHPVs). These viruses encode two oncoproteins (E6 and E7) whose expression is required for cancer initiation and maintenance. Among their cellular targets are the p53 and the retinoblastoma tumor suppressor proteins. Inhibition of the hrHPV E6-mediated ubiquitylation of p53 through the E6AP ubiquitin ligase results in the stabilization of p53, leading to cellular apoptosis. We utilized a live cell high throughput screen to determine whether exogenous microRNA (miRNA) transfection had the ability to stabilize p53 in hrHPV-positive cervical cancer cells expressing a p53-fluorescent protein as an *in vivo* reporter of p53 stability. Among the miRNAs whose transfection resulted in the greatest p53 stabilization was 375-3p that has previously been reported to stabilize p53 in HeLa cells, providing validation of the screen. The top 32 miRNAs in addition to 375-3p were further assessed using a second cell-based p53 stability reporter system as well as in non-reporter HeLa cells to examine their effects on endogenous p53 protein levels, resulting in the identification of 23 miRNAs whose transfection increased p53 levels in HeLa cells. While a few miRNAs that stabilized p53 led to decreases in E6AP protein levels, all targeted HPV oncoprotein expression. We further examined subsets of these miRNAs for their abilities to induce apoptosis and determined whether it was p53-mediated. The introduction of specific miRNAs revealed surprisingly heterogeneous responses in different cell lines. Nonetheless, some of the miRNAs described here have potential as therapeutics for treating HPV-positive cancers.

Importance

Human papillomaviruses cause approximately 5% of all cancers worldwide and encode genes that contribute to both the initiation and maintenance of these cancers. The viral oncoprotein E6 is expressed in all HPV-positive cancers and functions by targeting the degradation of p53 through the engagement of the cellular ubiquitin ligase E6AP. Inhibiting the degradation of p53 leads to apoptosis in HPV-positive cancer cells. Using a high throughput live cell assay we identified several miRNAs whose transfection stabilize p53 in HPV-positive cells. These miRNAs have the potential to be used in the treatment of HPV-positive cancers.

Introduction

Cervical cancer is the second leading cause of cancer deaths among women worldwide, with approximately 500,000 new cases diagnosed and 275,000 deaths each year. Virtually all cases of cervical cancer are attributable to infection by HPVs. There are over 200 different HPV types and a subset of these are linked to anogenital tract infections, 14 of which are referred to as hrHPV because they are associated with lesions that can progress to cancer (1). These same hrHPVs are also associated with other genital tract cancers (penile, vaginal, vulvar, and anal), as well as an increasing number of oropharyngeal cancers. Indeed, approximately 5% of all cancers worldwide are caused by hrHPVs (2).

Despite the success of the virus like particle (VLP)-based vaccine that has the potential to prevent millions of new HPV-associated cancers, there is the need for effective therapies to treat cancers that will develop in individuals who have not been vaccinated

and those already infected with hrHPVs. In this regard, hrHPVs provide a number of

potential viral targets for specific HPV antiviral therapies, including the viral encoded

oncoproteins E6 and E7, the viral E1 helicase, and the viral E2 regulatory protein.

The hrHPVs associated with cervical cancer encode two oncoproteins, E6 and E7,

that are invariably expressed in HPV-positive cancers and target the important cellular

growth regulatory proteins p53 and pRb, respectively (1). These cancers are dependent on

the continued expression of E6 and E7 and silencing these oncoproteins results in

apoptosis and senescence (3, 4). The E6 proteins of the hrHPV types promote the ubiquitin-

dependent degradation of the p53 protein (5) through its binding to the E3 ubiquitin ligase

E6AP (also known as UBE3A), which ubiquitylates p53 (6-8). In promoting the ubiquitin-

dependent proteolysis of p53, E6 counters E7-mediated replication stress signals that

activate and stabilize p53 (9). Thus, inhibition of the E6/E6AP-mediated ubiquitylation of

p53 is pro-apoptotic and provides a validated target for HPV-positive cancers. For example,

direct targeting of E6 induces apoptosis in HPV-positive cancer cells (10). Although the

hrHPV E6 proteins have other cellular activities that have been described in the literature,

we have focused on the E6/E6AP pathway as a potential therapeutic target for HPV-

positive cancers, as well as precancerous lesions.

MicroRNAs (miRNAs/miRs) are small non-coding RNAs (approximately 22

nucleotides in length) that significantly impact and regulate many essential biological

pathways. miRNAs negatively regulate gene expression either by translational repression

or cleavage of target mRNAs. Many miRNAs have been shown to play roles in cancer,

sometimes as oncogenes and other times as tumor suppressors, making them suitable as

therapeutic targets. miRNAs themselves can be used as cancer therapeutics as they may

affect one or more pathways essential for the replication and survival of specific cancer

cells. There have been a number of studies focused on HPV and miRNAs in recent years.

These studies have examined whether HPVs, like many other viruses, encode viral miRNAs

(11) and whether the HPV oncoproteins regulate the expression of cellular miRNAs (12).

Other studies have proposed miRNAs as potential biomarkers of hrHPV infections (13).

There have also been studies that examined the effects of miRNA expression in various

cancers including cervical cancers (For review see (14)). It was with this mindset that we

conducted a screen to identify miRNAs whose transfection into HPV-positive cancer cells

stabilize p53.

Here we present a high throughput screen using a cell-based platform to assay p53

stability in hrHPV E6-expressing HeLa cells that identified miRNAs whose transfection

increased p53 protein levels. Several of these miRNAs caused apoptosis and/or p21

induction when transfected into hrHPV-positive cells. The goal of this work was to find

miRNAs that could be used as therapeutic tools for the treatment of hrHPV-associated

diseases, regardless of whether these miRNAs are physiological regulators of p53 stability.

To our knowledge, this is the first study to systematically exam miRNAs for their abilities to

stabilize p53 in hrHPV-positive cancer cells. We also provide insights into the mechanisms

by which some of these miRNAs function in stabilizing p53 and promoting apoptosis.

Results

A high throughput screen identified transfected miRNAs that

stabilize p53 in HeLa cells.

To identify miRNAs that stabilize p53 and induce apoptosis when transfected into

hrHPV-positive cancer cells, we performed a high throughput screen in HPV18-positive

HeLa cells harboring a reporter for p53 protein stability. The reporter was originally

designed by the Elledge lab for their "global protein stability system"(15, 16), and

expresses a bicistronic mRNA that encodes an EGFP-p53 fusion protein and the red

fluorescent protein DsRed as reference (Figure 1A). The stability of the EGFP-p53 fusion

protein is determined by the less stable protein, in this case p53, with a half-life less than

30 minutes in HeLa cells due to E6/E6AP-mediated proteolysis (5, 17). To detect changes

in p53 stability, we monitored the EGFP/DsRed fluorescence ratio by laser scanning

cytometry. In our screen, for the fusion protein we used a p53 gene encoding the R273C

mutant, which is also efficiently targeted by E6/E6AP for ubiquitin-dependent proteolysis

(18). Because this mutant is defective in DNA binding (19, 20) and probably in the p53

mitochondrial pro-apoptotic pathways (21), it does not add an extra pro-apoptotic

stimulus to the eventual stabilization of the endogenous p53 during the course of the assay.

Furthermore, p53(R273C) can interfere with p53 transcriptionally driven apoptosis (22),

thus facilitating the acquisition of data for p53 stabilizers that might otherwise kill all cells.

The results of this screen, which assayed 885 miRNA mimics, are shown in Figure 1B and

Table S1. The top hit, 375-3p, has been previously reported to stabilize p53 in HPV-positive

cells (23) providing validation for this screen. Of note, since all the miRNAs used in this

study are human, we omitted the use of the prefix hsa-miR in the miRNAs names. We

selected for further testing the miRNAs that produced the next 32 highest EGFP/DsRed

fluorescence ratios (Table 1).

To confirm the results obtained in the screen, we tested the selected miRNAs in

HeLa cells containing a different bicistronic reporter vector that has been successfully used

in a small molecule screen to find p53 stabilizers in HeLa cells (24). In this vector,

p53(R273C) is fused to mRuby, a bright monomeric red fluorescent protein, and the

reference monomeric green fluorescent protein SGFP2 is fused to H2B (Figure 2A). For this

experiment, we determined the percentage of H2B-SGFP2 expressing cells that also

expressed mRuby-p53 (R273C) using flow cytometry. Two randomly selected miRNAs

(498-5p and 665), which were negative in the initial screen, and the miRNA Ctrl. 1 were

used as negative controls, along with 375-3p used as a positive control. As shown in Figure

2B, 25 of the 32 originally selected miRNAs were confirmed as significantly stabilizing

mRuby-p53 (R273C) in this reporter HeLa cell line.

Validation of candidate miRNAs in HeLa cells

Twenty seven miRNAs were next tested for their ability to stabilize endogenous p53

in HeLa cells by western blotting (Figure 3 and Table S2), including 3 miRNAs, 17-5p, 153-

3p, and 592 that produced a moderate increase in the levels of the reporter p53 . While

transfection of most of the miRNAs resulted in some degree of p53 stabilization compared

to the mock and negative controls, four miRNAs (219-5p, 106a-3p, 592, and 508-3p) had

little or no effect on endogenous p53 levels. The other miRNAs ranged from those that

produced a mild to moderate increase in p53 levels (511-5p, 519d-3p, 631, 374a-3p, 153-

3p, 17-5p, 373-3p, 210-3p, and 302c-3p) to those that caused stronger p53 stabilization

(519c-3p, 943, 374b-5p, 148b-5p, 519b-3p, 93-3p, 520d-3p, 520c-3p, 520b-3p, 519a-3p,

302a-3p, 302b-3p, 509-3p, and 1287-5p). We also examined the effects of the miRNAs on

E6AP and HPV18 E6 protein levels (Figure 4A-F and Table S2). Although most miRNAs had

little or no effect on E6AP levels, miRNAs 148b-5p, 374a-3p, and 374b-5p decreased the

levels of E6AP to values comparable to those observed in cells transfected with the positive

control 375-3p, which has been previously shown to target the E6AP mRNA (23).

Transfection of a few miRNAs, most notably 93-3p and 1287-5p, resulted in elevated levels

of E6AP. In contrast, transfection of most of the miRNAs tested in this experiment led to a

reduction in E6 protein levels. While there was no correlation between the protein levels of

E6 and E6AP ($r(35) = 0.02098$, $p = 0.9019$) or between E6AP and p53 ($r(35) = -0.07389$, p

$= 0.6638$), there was a significant negative correlation between the levels of E6 and p53

($r(35) = -0.6288$, $p<0.0001$) (Figure 4G), suggesting that a decrease in E6 expression plays

a significant role in the stabilization of p53 by several of these miRNAs in HeLa cells.

Examination of these data suggest that, under the conditions used in this experiment, a

~50% reduction in E6 protein levels appears necessary for p53 stabilization in HeLa cells.

Sequence analysis of the validated miRNAs showed a significant enrichment of miRNAs

sharing a common seed sequence (5'-AAGUGC-3') that defines the 302-3p/372-3p/373-

3p/519-3p/520-3p family of miRNAs (from now on referred as 302/519 miRNA family)

(Figure 5). Based on these results, we selected miRNAs 148b, 374a, 374b, 1287, and five

members of the 302/519 family for further analysis.

Transfection of selected miRNAs increase p53 in different

hrHPV-positive cell lines.

To examine whether there was any HPV specificity to the stabilization of p53 by

these miRNAs, we examined their effects on the stability of p53 in other hrHPV-positive

cancer cell lines. We transfected three cervical cancer cell lines, SiHa (HPV16), ME-180

(HPV68), and MS751 (HPV45), as well as two head and neck squamous carcinoma cell

lines, UM-SCC-47 and UM-SCC-104 (both HPV16), with the selected miRNAs and examined

p53, E6AP, and actin protein levels by western blot. As shown in Figure 6A-E, the

transfection of these miRNAs increased the protein levels of p53 in all HPV-positive cell

lines, indicating this effect is conserved in other HPV-associated cervical cancer cell lines

independent of the hrHPV type. However, the relative magnitude of the p53 stabilization by

individual miRNAs varied in the five HPV-positive cell lines, likely reflecting differences

between the cell lines and/or HPV types. 519d-3p consistently produced the lowest

increase in p53 protein levels compared to the other 302/519 miRNA family members

used in these experiments. These miRNAs share considerable sequence homology

throughout and the same seed sequence with exception of 519d-3p, whose seed sequence

is displaced one position by the addition of a C at its 5' end (Figure 5). This suggests that a

good pairing of the seed sequence of these miRNAs with their target mRNA is important for

the ability of the members of 302/519 miRNA family to induce accumulation of p53 when

transfected into hrHPV-positive cancer cells.

To evaluate the involvement of the E6/E6AP pathway in the stabilization of p53 by

these miRNAs, we examined their effects on the levels of p53 and E6AP in HPV-negative

human U2OS cells (Figure 6F), an osteosarcoma-derived cell line that harbors wt p53

under the control of Mdm2. Transfection of miRNAs 375-3p and 148b-5p increased the

levels of p53 in U2OS cells, indicating that these two miRNAs can affect p53 protein levels

independent of HPV oncoprotein expression. In contrast, transfection of the miRNAs from

the 302/519 family and 1287-5p had minimal or no effect on the levels of p53 in U2OS

cells, suggesting that the effect of these miRNAs on p53 stability is dependent upon the

expression of hrHPV E6. Of note, transfection of miRNAs 375-3p, 148b-5p and to a lesser

degree 374a-3p and 374b-5p, decreased the levels of E6AP in each of the cell lines

analyzed.

Several miRNAs decrease expression of E6/E7.

Since the increase of p53 in HeLa cells transfected with the miRNAs identified in this

study negatively correlated to the levels of E6 (Figure 4G), we hypothesized that the ability

of these miRNAs to stabilize p53 in HPV-positive cancer cells might be mediated, at least in

part, by a negative effect on E6 expression. This could be a consequence of destabilizing the

E6 protein or decreasing its expression, either by targeting the E6/E7 mRNAs for

degradation, preventing their translation, or decreasing viral transcription. We observed

that transfection of these miRNAs into HeLa cells, particularly members of the 302/519

family and 1287-5p, decreased both E6 and E7 protein levels similarly (Figure 7A). This is

consistent with the idea that these miRNAs affect the expression of the viral E6/E7 mRNAs

since the hrHPV E6 and E7 genes are expressed from a single promoter located in the viral

LCR. This effect on the level of the E6/E7 mRNA in HeLa cells was confirmed by qRT-PCR

using primers that detect all HPV E6/E7 transcripts (Figure 7B). Interestingly, all miRNAs

tested, including 375-3p and 148b-5p that increased p53 protein levels in the absence of

hrHPV proteins (Figure 6F), also decreased the levels of the E6/E7 transcripts. Although

498-5p reduced the level of the E6/E7 mRNAs in HeLa cells, consistent with the reduced

level of E6 protein shown in Figures 3, 4 and 7A, we did not observe an increase in p53

protein levels (Figures 3 and 6). While 375-3p was as efficient in repressing the expression

of the E6/E7 mRNAs in HPV16 SiHa cells as in HPV18 HeLa cells, most of the tested

miRNAs were less efficient decreasing the levels of the E6/E7 transcripts in SiHa cells, even

though they were efficiently transfected as determined by siGlo RISC-free transfection

control (data not shown). In addition to 375-3p, only 519b-3p and 1287-5p showed

comparable effects on the E6/E7 transcripts in both HPV-positive cell lines. Surprisingly,

148b-5p, which stabilized p53 in SiHa cells, produced an increase in the levels of the E6/E7

mRNAs but we could not confirm whether E6 protein levels were affected in this cell line

(e.g. by preventing translation) because we were unable to detect HPV16 E6 using any of

the commercially available anti-HPV16 E6 antibodies.

To determine whether these miRNAs were targeting the untranslated regions (UTR)

of the HPV16 E6/E7 mRNAs, we examined their effect in 1321 cells, a human keratinocyte

cell line that expresses HPV16 E6/E7 under the control of the human β-actin promoter and

from a mRNA containing the 3'UTR of the human β-actin mRNA (25). In this cell line, the

302/519 miRNA family members (302a-3p, 373-3p, 519b-3p, 519d-3p, and 520c-3p) and

1287-5p failed to increase the p53 protein levels, suggesting that their effect involves

actions over the promoter activity of the HPV16 LCR and/or the 3'UTR of the E6/E7

transcripts (Figure 7C). In contrast, 375-3p, 148b-5p, 374a-3p, and 374b-5p transfection

resulted in an increase in p53 protein levels, indicating it is likely through a different

mechanism than the miRNAs of the 302/519 family and 1287-5p. Considering that the

E6/E7 transcripts in the different hrHPV-positive cell lines tested have different 3'UTRs,

depending on the viral sequence inserted in the host cell and the insertion site, it is more

likely that the miRNAs of the 302/519 family and 1287-5p affect E6 and E7 protein levels

by acting on the viral LCR promoter activity. In line with this hypothesis, 1287-5p and the

302/519 family members efficiently repressed the HPV16 LCR and HPV18 LCR promoter

activity in a reporter system in the HPV-negative cervical cancer cells C33A (Figure 7D),

Selected miRNAs induce apoptosis and/or p21 expression.

Stabilization of p53 is expected to be pro-apoptotic in hrHPV-positive cells. We

therefore tested whether miRNAs that stabilized p53 in HeLa cells induce apoptosis. Flow

cytometric analysis of HeLa cells transfected with different miRNAs and stained with the

apoptosis marker Annexin V revealed an increase in the fraction of apoptotic cells when

compared to untreated or mock transfected cells (Figure 8A). Of note, miRNAs Ctrl.1 and

498-5p, which did not increase p53 levels in hrHPV-positive cells (Figures 3 and 6), did

increase Annexin V staining upon transfection into HeLa cells. In contrast, there was a

correlation between induction of apoptosis and stabilization of p53 in HeLa cells

transfected with the 302/519 family members (Figure 3). In this family, 519d-3p, which

produced the lowest increase in p53 levels in HeLa cells, was also the least efficient in

inducing apoptosis.

We next examined the ability of a subset of these miRNAs to induce apoptosis in

SiHa cells and two HPV-negative cancer cell lines, C33A and U2OS, using Annexin V staining

(Figure 8B). Transfection with the miRNA 20a-3p was utilized as a negative control, since it

does not induce apoptosis in cervical cancer cells (26). The 1287-5p miRNA was also

included in these experiments as it was found to stabilize p53 (Figures 3 and 6) and to

efficiently decrease E6/E7 mRNAs expression in both HeLa and SiHa cells (Figure 7B).

Similar to their effect in HeLa cells, 148b-5p and 1287-5p induced apoptosis in SiHa cells

whereas 375-3p and 519b-3p had very little or no effect. While all of these miRNAs

increased the levels of p53 when transfected into SiHa cells (Figure 6A), 148b-5p and

1287-5p led to the highest p53 levels. Both 148b-5p and 1287-5p stimulated apoptosis in

C33A cells, an HPV-negative cervical cancer cell line that already expresses high levels of an

inactive mutant form of p53 (27), indicating that these miRNAs can induce apoptosis

independently of p53.

Since 519b-3p has been reported to induce senescence in HeLa cells through p21

expression, which is partially dependent of p53 (28), we tested whether other members of

the 302/519 family have a similar effect on p21 protein levels and if it also occurs in

HPV16-positive SiHa cells. Since 375-3p has been shown to increase p21 protein levels in

both HeLa and SiHa cells (23), it was included as the positive control. As shown in Figure

8C, the miRNAs of the 302/519 family tested in this experiment increased p21 protein

levels in both cell lines with the exception of 519d-3p, which did not increase p21 levels in

SiHa cells. Interestingly, 498-5p led to an increase in p21 protein levels in SiHa cells;

however this is likely to be at least partially p53 independent since this miRNA failed to

induce a marked p53 accumulation under similar conditions in these cells (Figure 6A).

miRNAs of the 302/519 family and 1287-5p induce p53-

dependent apoptosis in HeLa cells.

We next examined if induction of apoptosis by several representative miRNAs is

p53-dependent in HeLa cells. We performed Annexin V staining of HeLa cells co-

transfected with the miRNAs and an siRNA targeting p53 or its C911 variant (29) to

address possible off-target effects of the siRNA (Figure 9A). As expected, the positive

control 375-3p increased the percentage of Annexin V-stained cells while 20a-3p had no

significant effect when compared to untreated cells. Surprisingly, the apoptotic effect of

375-3p was independent of p53 since knock-down of p53 had little effect on the percentage

of apoptotic cells in HeLa cells co-transfected with this miRNA. The effect of 498-5p on

apoptosis is also p53-independent and consistent with the inability of this miRNA to

stabilize p53 when transfected into hrHPV-positive cell lines (Figures 3 and 6). Similarly,

the apoptotic effects of miRNAs 148b-5p and 374a-3p are also independent of p53, as co-

transfection of these miRNAs with either the siRNA against p53 or its C911 version have

similar effects on the percentage of apoptotic cells. This indicates that, in these cases, the

effect observed when p53 is knocked down can be attributed to off-target effects of the

siRNA against p53. In contrast, the numbers of apoptotic cells observed with 519b-3p (as a

representative of the 302/519 family of miRNAs) and 1287-5p were reduced when co-

transfected with the p53 siRNA but not with its C911 version, suggesting that the apoptotic

effect of these miRNAs is largely dependent on p53 in HeLa cells. Similar results were

observed when using PARP cleavage as a marker for apoptosis in western blot analysis (Fig

9B).

Discussion

In this study we identified several miRNAs that increased p53 protein levels when

transfected in HeLa cells. Some of the miRNA hits from our high throughput screen (148b-

5p, 302a-3p, 373-3p, and 374b-5p) have been previously reported to have decreased

expression in cervical cancer cells and tissues compared to the controls used in each of the

studies (26, 30-32). Another positive miRNA from the screen, 1287-5p, has been found to

have no change in its expression level in cervical cancer cells, but is sequestered away from

its targets by the circular RNA circSLC26A4, which is overexpressed in hrHPV-positive

cancer cells (33). Together these findings suggest that interfering with the functions of

some of the miRNAs identified in this work might contribute to the establishment of HPV

infections or/and progression of hrHPV-infected cells towards cancer. Consistent with that,

several of the miRNAs identified in our screen, namely 148b-5p, 302a-3p, 374b-5p, or

519b-3p, have been shown to slow cell growth and induce apoptosis in hrHPV-positive

cervical cancer cells (28, 31, 32, 34, 35). Taken together, these previous findings validate

this screening platform as a reliable tool to identify miRNAs that affect the cell growth and

survival of hrHPV-positive cells, as expected from miRNAs whose transfection stabilize

p53.

Since the ubiquitin ligase activity of E6AP/E6 is responsible for the efficient

proteolysis of p53 in hrHPV-positive cervical cancer cells, we determined whether

transfection of any of the miRNAs identified in this screen altered the levels of E6AP in

HeLa cells. Interestingly, only three of the 27 miRNAs analyzed in HeLa cells, 148b-5p,

374a-3p, and 374b-5p, led to a decrease in E6AP protein levels comparable to the decrease

produced by 375-3p. Although we did not determine whether they directly bind to the

E6AP mRNA, target prediction analysis using the miRDB database (www.mirdb.org) (36,

37) revealed that 148b-5p and 374b-5p each have three sites complementary to their seed

sequences within the 3'UTR of the E6AP mRNA. Remarkably, these four miRNAs (148b-5p,

374a-3p, 374b-5p, and 375-3p) also led to decreased E6 expression. This simultaneous

decrease of both E6AP and E6 may be required to stabilize p53 in HeLa cells, since a small

amount of E6AP is sufficient to efficiently promote p53 degradation in this cell line

(38). The need to suppress the very efficient E6/E6AP-mediated degradation of p53 to

increase the levels of this protein in hrHPV-positive cancer cells is exemplified by miRNAs

375-3p and 1287-5p. Both miRNAs stabilized p53 in absence of E6 when transfected into

U2OS cells, however, they also efficently decrease the levels of E6 when transfected into

HeLa and SiHa cells, suggesting that this is a necessary condition to stabilize p53 in these

cells. Moreover, in this study we haven't found any miRNA that stabilize p53 in hrHPV-

cancer cells that do not significantly decrease the E6 protein level too. In contrast,

transfection of 498-5p, which also resulted in decreased levels of E6 in these cells, did not

stabilize p53 in HeLa cells, suggesting that the remaining E6 protein was sufficient to target

p53 for degradation. In line with this idea, in our study, miRNAs that failed to reduce E6

protein levels to approximately half of its original amount or beyond also failed to increase

p53 levels in HeLa cells, suggesting that there is a considerable excess of E6 in these cells

that must be overcome to stabilize p53. High levels of expression of E6 could therefore be a

limiting factor for the use of some miRNAs such as 498-5p as therapeutics, at least in some

hrHPV-associated cancers.

While several of the transfected miRNAs were found to induce apoptosis, the

mechanisms by which this occurred varied. The apoptotic induction in HeLa cells for three

of the miRNAs, 148b-5p, 374a-3p, and 375-3p, was independent of p53 stabilization. In

contrast, transfection of 1287-5p induced apoptosis in HeLa cells in a p53-dependent

manner. The miRNA 1287-5p was also found to efficiently induce apoptosis in U2OS cells

without a noticeable increase in p53 levels and to increase the number of apoptotic C33A

cells, which do not express wt p53, indicating that this miRNA promotes cell death

independent of p53 stabilization in these HPV-negative cells. Therefore, 1287-5p can

induce apoptosis via different pathways depending on the cell type. This is also exemplified

by Ctrl. 1 whose transfection stabilized p53 only in U2OS cells. The other miRNA that we

found to induce apoptosis in HeLa cells in a p53-dependent manner upon transfection was

519b-3p, a member of the 302/519 family. Abdelmohsen and colleagues have previously

reported that 519-3p transfection in HeLa cells increases p21 levels in a partly p53-

independent manner, resulting in growth inhibition and cell survival (28). There are a

number of technical differences between our study and that of Abdelmohsen et al. which

could account for the discordant results, including different miRNA concentrations and end

points. However, we both found that 519b-3p transfection did not induce apoptosis in SiHa

cells, but led to an increase in p53 and p21 protein levels. It has been reported that p53

must accumulate to a threshold level to induce apoptosis (39). Thus, it is feasible that

under the experimental conditions used in our study 519b-3p did not increase p53 levels to

the threshold necessary to trigger apoptosis in SiHa cells. In addition, we observed that

transfection of all miRNAs belonging to the 302/519 family, with the exception of 519d-3p

in SiHa cells, increased p21 protein levels in hrHPV-positive cells indicating this is a family

characteristic rather than a feature of 519b-3p alone.

Throughout this study, 519d-3p has been less efficient than the other members of

the 302/519 family of miRNAs in promoting increased p53 and p21 protein levels in the

hrHPV-positive cell lines, as well as in repressing expression from the HPV16 and HPV18

LCRs, and inducing apoptosis in HeLa cells. In addition to a few base substitutions in the 3'

end sequences, a difference between 519d-3p and the other miRNAs of this family included

in this study is the addition of a C at its 5'end. This additional nucleotide produces a

displacement of the miRNA sequences, including its seed sequence, which is 5'-AAAGUG-3'

instead of 5'- AAGUGC-3', the most common seed sequence between the members of the

302/519 family. Since a seed sequence match is very important for miRNA target

recognition (40, 41), this change in 519d-3p's seed sequence is expected to impact its

functions when compared to the other members of the 302/519 family. The data presented

in this study supports this hypothesis and indicate that the correct matching of the seed

sequence 5'-AAGUGC-3' with their target mRNAs is an important factor mediating the

effects of this family of miRNAs on p53 and p21 protein levels as well as on cell growth and

survival of hrHPV-positive cancer cells.

Since their discovery in 1993 (42, 43), the interest in miRNAs continues to increase.

Research in miRNAs is being translated into diagnostics as miRNAs are considered

sensitive and specific biomarkers. In contrast, their use as therapeutics have lagged with

only a few clinical trials currently underway (reviewed in (44, 45)). This may be due in part

to the lack of single target specificity characteristic of miRNAs. This multiplicity of targets

makes miRNAs potentially challenging as therapeutics because of the possibility of

unexpected and/or undesirable side effects. However, this property makes them an

attractive tool for cancer treatment since miRNAs can simultaneously regulate multiple

cellular pathways by modulating the expression of several genes in a coordinated fashion.

The utility of miRNAs as therapeutic tools will be improved by a more complete

understanding of how they function in the cellular networks in which they are embedded.

In this study, we identified several miRNAs that stabilize p53 in hrHPV-cancer cells and

induce apoptosis in HeLa cells in a p53-dependent or in p53-independent manner. Despite

the heterogenicity of the responses to different miRNAs, decreasing E6 protein levels

seems to be required to increase p53 protein levels in HeLa cells since all miRNAs found to

stabilize p53 also affected the expression of E6. Further research into the mechanisms

involved in the stabilization of p53 and/or induction of apoptosis by these miRNAs in

hrHPV-positive cells will improve our understanding of the hrHPV-host interactions and

contribute to the identification of new potential therapeutic targets for hrHPV infections

and cancers.

Materials and methods

Plasmids

To create a lentiviral reporter vector to monitor the protein levels of p53 *in vivo* in

hrHPV cancer cells, the p53 open reading frame (ORF) carrying the mutation R273C was

first amplified by PCR using the primers hp53-1 (5'-CACCATGGAGGAGCCGCAGTCAGATCC-

3') and hp53-8 (5'-GGATCCTCAGTCTGAGTCAGGCCCTTCTGTCTTG-3') and cloned into the

pENTR/D-TOPO vector (ThermoFisher) generating the entry vector pENTR p53(R273C)

(p6140). The p53(R273C) ORF was then recombined from the entry vector into a lentiviral

vector containing a GPS (<u>G</u>lobal <u>P</u>rotein <u>S</u>tability) cassette (gift from S. Elledge) using the

GATEWAY cloning system (ThermoFisher), resulting in the vector pHAGE-P CMVt N-RIG2

p53(R273C) (p7987). The GPS reporter cassette expresses a bicistronic mRNA encoding

EGFP fusion proteins and the red fluorescent protein DsRed as reference (15, 16, 46, 47).

The reporter vector pHAGE-N CMVt N-RIG3 p53(R273C) (p7709), which contains a variant

of the GPS reporter cassette, expresses p53(R273C) fused to the monomeric red

fluorescent protein mRuby and a monomeric green fluorescent protein (SGFP2) fused to a

histone 2B (H2BC11) as reference (38).

Cell lines

HeLa, HeLa RIG2 p53(R273C), HeLa RIG3 p53(R273C), Siha (ATCC® HTB-35™),

C33A (ATCC® HTB-31™), U2OS (ATCC® HTB-96™), ME-180 (ATCC® HTB-33™), MS-751

(ATCC® HTB-34™), UM-SCC-47 and UM-SCC-104 (EMD-Millipore), and human

keratinocytes immortalized by HPV16 E6 and E7 expressed from the β-actin promoter

(p1321) (25) referred to here as 1321 cells were grown in Dulbecco's modified Eagle's

medium (DMEM) supplemented with 10% (v/v) fetal bovine serum at 37°C 5% CO_2.

miRNA screen

For the primary miRNA screen, HeLa RIG2 p53(R273C) cells were reverse

transfected with the Ambion Human Pre-miR miRNA Mimic Library (2009) based on

miRBase release 13.0. in a one-target, one-well format using 384-well microtiter plates in a

high throughput format. The Z'-factor (48) was calculated for each assay plate and was

consistently > 0.6, indicating assay robustness. siRNA buffer (1×) (Horizon #B-002000-UB-

100) was aliquoted into wells, miRNA mimics were added so that the final concentration

was 40 nM/well, and DharmaFECT1/OptiMEM was dispensed into wells. While the

miRNA/lipid mix was allowed to complex, cells were trypsinized, counted, and

resuspended to reach a plating density of 600 cells/well. Cells were seeded on top of the

miRNA/lipid mixture, briefly centrifuged, and incubated for 72 hours, a time that we

determined to be optimal for p53 stabilization for small miRNAs. The cells were then

equilibrated to room temperature (~15 min) before analysis. Using TTP LabTech's Acumen

eX3 laser scanning cytometer, the total EGFP and DsRed intensities in each well were

quantitated. Internal controls were present in Columns 13 and 14 of each library plate

(Pre-miR miRNA Precursor Molecules Negative Control #1 - Ambion AM17110; Pre-miR

miRNA Precursor Molecules Negative Control #2 – Ambion AM17111; PLK1 SMARTpool

Dharmacon M-003290-01). In addition, siRNAs against E6AP (UBE3A, Dharmacon D-

005137-04) and non-targeting siRNAs (Dharmacon D-001210-01 and D-001210-02) were

added to each plate. The miRNA mimics were screened in triplicate. High throughput

libraries and screening capability was provided by the ICCB-Longwood Screening Facility

at Harvard Medical School.

miRNA transfections

All cells were reverse transfected with miRNAs (Table 2) using RNAiMax according

to manufacturer's instructions (ThermoFisher Scientific). More specifically, 750 µl of

suspended cells were added to 250 µl of a transfection mix containing 2 µl of Lipofectamine

RNAiMax and different miRNAs for a final concentration of 10nM and plated in 12-well

plates. miRNA transfection efficiency in different cell lines was confirmed using siGlo Red

Transfection Indicator (Horizon, D-001630-02-05).

Confirmation assay

458   HeLa cells were reverse transfected with 10 nM of the indicated miRNAs using

459   Lipofectamine RNAiMAX according to manufacturer's instructions. 72h after transfection

460   the cells were trypsinized, harvested in 1X flow cytometry buffer (1X PBS, 1mM EDTA, 2%

461   FBS) and analyzed for green and red fluorescence by flow cytometry using a BD

462   FACSymphony instrument (BD-Biosciences).

Western blots and antibodies

After incubation for 72h, transfected cells were washed once with 1X PBS, lysed

directly in wells with 100-150µl of SDS Lysis buffer (62.5mM Tris-Cl, pH 6.8, 2% SDS) and

sonicated (amplitude of 35%; 10 pulses with 2 seconds ON and 0.5 seconds OFF). The

lysates were centrifuged for 2 minutes and protein concentrations quantitated by BCA

Protein Assay (Pierce). Lysates (10 µg of protein/well) were run on 20-well 4-12% NuPage

BisTris midi gels in 1X MES buffer (ThermoFisher Scientific) and transferred to

polyvinylidene difluoride (PVDF) membranes using 25 mM Tris-HCl pH 7.6, 192 mM

glycine, and 10% methanol. Membranes were blocked in 5% nonfat dried milk in Tris-

buffered saline [pH 7.4], 0.05% Tween 20 (TBST) and then incubated with primary

antibodies as follows: mouse monoclonal anti-E6AP (clone E4) at 1:1000 dilution (sc-

166689 from Santa Cruz Biotechnology); rabbit monoclonal anti-cleaved PARP (Asp214)

(D64E10 XP® #5625 from Cell Signaling Technologies); mouse monoclonal anti-HPV18-E6

(clone G-7) at a 1:250 dilution (sc-365089 from Santa Cruz Biotechnology); mouse

monoclonal anti-p21 Waf1/Cip1/CDKN1A (clone 187) at a 1:1000 dilution (sc-817 from

Santa Cruz Biotechnology); and mouse monoclonal anti-HPV18-E7 (clone F-7) at a 1:250

dilution (sc-365035 from Santa Cruz Biotechnology). Membranes were washed in TBST

and incubated with horseradish peroxidase (HRP)-conjugated anti-mouse antibody at a

1:1,000 dilution (R1005 from Kindle Biosciences). The primary antibodies against β-Actin

and P53 were directly conjugated to HRP and were used as follows: mouse monoclonal

anti-β-Actin-HRP at a 1:50,000 dilution (A3854 from Millipore Sigma); and goat polyclonal

anti-p53-HRP at a 1:3,000 dilution (HAF1355 from R&D systems). All blots were developed

using KwikQuant Ultra Digital-ECL™ Substrate Solution and processed using KwikQuant™

Imager (Kindle Bioscience LLC). Western blots were quantified using the Image Studio Lite

program (LI-COR Biotechnologies).

RNA isolation and qRT-PCR.

At 72h post-transfection, total RNA was isolated using the Nucleospin RNA kit

(Takara Bio) according to the manufacturer's instructions. Samples were quantified using

NanoDrop™ Lite Spectrophotometer and equal amounts of RNA (500ng) were reverse

transcribed using the high-capacity cDNA reverse transcription kit (Applied Biosystems).

Quantitative real-time PCR (qRT-PCR) was performed in an Applied Biosystems ABI 7500

fast sequence detection system using TaqMan Fast Advanced Master Mix (ThermoFisher

Scientific), and the following TaqMan custom gene expression assays for HPV18-E6E7:

HPV18-E6E7 FWD 5'-CAACCGAGCACGACAGGAA-3'; HPV18-E6E7 PROBE 5'-

AATATTAAGTATGCATGGACCTAAGGCAACATTGCAA-3'; and HPV18-E6E7 REV 5'-

CTCGTCGGGCTGGTAAATGTT-3' and for HPV16-E6E7: HPV16-E6-FWD 5'-

AGGAGCGACCCGGAAAGT-3'; HPV16-E6-PROBE 5'-

ACCACAGTTATGCACAGAGCTGCAAACAA-3'; and HPV16-E6-REV 5'-

CACGTCGCAGTAACTGTTGCTT-3'. All reactions were normalized using a TaqMan gene

expression assay for PPIA (Hs99999904_m1 ThermoFisher Scientific).

Luciferase assays

C33A cells were transfected in 12-well plates with 10 nM of miRNAs as described

above. After 24h, the miRNA transfection mix was removed and cells were transfected with

0.5 µg of either plasmid pGL4-LCRHPV16 (p6239, HPV16 nt 7000-85) (49) or pGL4-

LCRHPV18 (p5194, HPV18 nt 6943-105) (50), containing the firefly luciferase ORF under

the control of the HPV16 and HPV18 LCRs respectively, together with 0.5µg of a plasmid

expressing the renilla luciferase under the control of the human β-actin promoter using

Lipofectamine 3000 (ThermoFisher Scientific) according to manufacturer's protocol. At

48h post-transfection of plasmids (72 hours after miRNA transfection), cells were lysed in

the plate with 250 µl of 1x Passive Lysis Buffer and luciferase activity was determined

using the Dual-Luciferase Reporter Assay System (Promega) according to the

manufacturer's instructions. Luciferase intensities were measured using a SpectraMax L

luminescence microplate reader (Molecular Devices). Firefly luciferase readings were

normalized to the respective renilla luciferase readings.

Apoptosis assays

HeLa cells were reverse transfected with 10 nM of the different miRNAs either alone

or combined with 20 nM of siRNAs directed to p53 (Dharmacon, D-003329-26: 5'-

GCUUCGAGAUGUUCCGAGA-3') or its C911 version (sense, 5'-GCUUCGAGUACUUCCGAGA-

UU-3') using Lipofectamine RNAiMAX according to manufacturer's instructions. After

incubation for 72h, attached cells were trypsinized and combined with all non-adherent

cells in the supernatant and PBS washes. The cells were stained with Annexin V using the

Dead Cell Apoptosis Kit with Annexin V Alexa Fluor™ 488 & Propidium Iodide (PI)

(ThermoFisher Scientific #V13241 and #V13245). Stained cells were analyzed by flow

cytometry using a BD FACSCanto flow cytometer (BD Biosciences).

Figures and statistics

528   All graphics and statistical analyses were conducted using Prism 8 (GraphPad).

529   When the values corresponding to mock transfections were used to normalize other values

530   obtained in an experiment, the mock values were not included in the corresponding

531   graphic since invariably their mean is equal to one and their standard deviation is zero.

532   Figures were assembled using Affinity Photo (Affinity).

Table 1. miRNAs selected from primary screen

Rank	miRNA	Mature_Sequence	EGFP / DsRed (Mean, n = 3)	EGFP / DsRed (SD, n =3)
1	hsa-miR-375-3p	UUUGUUCGUUCGGCUCGCGUGA	0.12298	0.01761
2	hsa-miR-219a-5p	UGAUUGUCCAAACGCAAUUCU	0.07703	0.00559
3	hsa-miR-511-5p	GUGUCUUUUGCUCUGCAGUCA	0.07371	0.00962
4	hsa-miR-519c-3p	AAAGUGCAUCUUUUUAGAGGAU	0.07180	0.01690
5	hsa-miR-943	CUGACUGUUGCCGUCCUCCAG	0.06086	0.02073
6	hsa-miR-200c-5p	CGUCUUACCCAGCAGUGUUUGG	0.05781	0.00790
7	hsa-miR-944	AAAUUAUUGUACAUCGGAUGAG	0.05766	0.03304
8	hsa-miR-106a-3p	CUGCAAUGUAAGCACUUCUUAC	0.05210	0.00779
9	hsa-miR-374b-5p	AUAUAAUACAACCUGCUAAGUG	0.05065	0.00389
10	hsa-miR-148b-5p	AAGUUCUGUUAUACACUCAGGC	0.05013	0.00438
11	hsa-miR-519d-3p	CAAAGUGCCUCCCUUUAGAGUG	0.04929	0.01507
12	hsa-miR-631	AGACCUGGCCCAGACCUCAGC	0.04839	0.01369
13	hsa-miR-592	UUGUGUCAAUAUGCGAUGAUGU	0.04690	0.01220
14	hsa-miR-519b-3p	AAAGUGCAUCCUUUUAGAGGUU	0.04442	0.01903
15	hsa-miR-93-3p	ACUGCUGAGCUAGCACUUCCCG	0.04441	0.00884
16	hsa-miR-340-5p	UUAUAAAGCAAUGAGACUGAUU	0.04336	0.00383
17	hsa-miR-508-3p	UGAUUGUAGCCUUUUGGAGUAGA	0.04083	0.00853
18	hsa-miR-374a-3p	UUAUAAUACAACCUGAUAAGUG	0.03978	0.00887
19	hsa-miR-520d-3p	AAAGUGCUUCUCUUUGGUGGGU	0.03790	0.01428
20	hsa-miR-520c-3p	AAAGUGCUUCCUUUUAGAGGGU	0.03290	0.01176
21	hsa-miR-520b-3p	AAAGUGCUUCCUUUUAGAGGG	0.03226	0.01171
22	hsa-miR-153-3p	UUGCAUAGUCACAAAAGUGAUC	0.03096	0.00799
23	hsa-miR-17-5p	CAAAGUGCUUACAGUGCAGGUAG	0.02945	0.01485
24	hsa-miR-519a-3p	AAAGUGCAUCCUUUUAGAGUGU	0.02878	0.00725
25	hsa-miR-1287-5p	UGCUGGAUCAGUGGUUCGAGUC	0.02854	0.00375
26	hsa-miR-373-3p	GAAGUGCUUCGAUUUUGGGGUGU	0.02800	0.00787
27	hsa-miR-302a-3p	UAAGUGCUUCCAUGUUUUGGUGA	0.02791	0.00701
28	hsa-miR-302b-3p	UAAGUGCUUCCAUGUUUUAGUAG	0.02749	0.01175
29	hsa-miR-509-3p	UGAUUGGUACGUCUGUGGGUAG	0.02568	0.00559
30	hsa-miR-210-3p	CUGUGCGUGUGACAGCGGCUGA	0.02560	0.00996
31	hsa-miR-302c-3p	UAAGUGCUUCCAUGUUUCAGUGG	0.02376	0.00492
32	hsa-let-7a-2-3p	CUGUACAGCCUCCUAGCUUUCC	0.02275	0.01039
33	hsa-miR-224-5p	CAAGUCACUAGUGGUUCCGUU	0.02230	0.00997

544 **Table 2. miRNAs used in this study**

Description	Vendor	catalog number
miRIDIAN microRNA Human hsa-miR-375 - Mimic, 2 nmol	Dharmacon	C-300682-05-0002
miRIDIAN microRNA Human hsa-miR-219a-5p - Mimic, 2 nmol	Dharmacon	C-300576-05-0002
miRIDIAN microRNA Human hsa-miR-511-5p - Mimic, 2 nmol	Dharmacon	C-300752-03-0002
miRIDIAN microRNA Human hsa-miR-519c-3p - Mimic, 2 nmol	Dharmacon	C-300786-03-0002
miRIDIAN microRNA Human hsa-miR-943 - Mimic, 2 nmol	Dharmacon	C-301275-01-0002
miRIDIAN microRNA Human hsa-miR-200c-5p - Mimic, 2 nmol	Dharmacon	C-301070-01-0002
miRIDIAN microRNA Human hsa-miR-944 - Mimic, 2 nmol	Dharmacon	C-301276-01-0002
miRIDIAN microRNA Human hsa-miR-106a-3p - Mimic, 2 nmol	Dharmacon	C-301159-01-0002
miRIDIAN microRNA Human hsa-miR-374b-5p - Mimic, 2 nmol	Dharmacon	C-301250-01-0002
miRIDIAN microRNA Human hsa-miR-148b-5p - Mimic, 2 nmol	Dharmacon	C-301087-01-0002
miRIDIAN microRNA Human hsa-miR-519d-3p - Mimic, 2 nmol	Dharmacon	C-300812-05-0002
miRIDIAN microRNA Human hsa-miR-631 - Mimic, 2 nmol	Dharmacon	C-300957-01-0002
miRIDIAN microRNA Human hsa-miR-592 - Mimic, 2 nmol	Dharmacon	C-300916-01-0002
miRIDIAN microRNA Human hsa-miR-519b-3p - Mimic, 2 nmol	Dharmacon	C-300791-05-0002
miRIDIAN microRNA Human hsa-miR-93-3p - Mimic, 2 nmol	Dharmacon	C-301207-01-0002
miRIDIAN microRNA Human hsa-miR-340-5p - Mimic, 2 nmol	Dharmacon	C-301081-01-0002
miRIDIAN microRNA Human hsa-miR-508-3p - Mimic, 2 nmol	Dharmacon	C-300848-05-0002
miRIDIAN microRNA Human hsa-miR-374a-5p - Mimic, 2 nmol	Dharmacon	C-300681-05-0002
miRIDIAN microRNA Human hsa-miR-520d-3p - Mimic, 2 nmol	Dharmacon	C-300816-05-0002
miRIDIAN microRNA Human hsa-miR-520c-3p - Mimic, 2 nmol	Dharmacon	C-300803-05-0002
miRIDIAN microRNA Human hsa-miR-520b - Mimic, 2 nmol	Dharmacon	C-300797-03-0002
miRIDIAN microRNA Human hsa-miR-153-3p - Mimic, 2 nmol	Dharmacon	C-300615-07-0002
miRIDIAN microRNA Human hsa-miR-519a-3p - Mimic, 2 nmol	Dharmacon	C-300836-05-0002
miRIDIAN microRNA Human hsa-miR-1287-5p - Mimic, 2 nmol	Dharmacon	C-301341-00-0002
miRIDIAN microRNA Human hsa-miR-373-3p - Mimic, 2 nmol	Dharmacon	C-300680-03-0002
miRIDIAN microRNA Human hsa-miR-302a-3p - Mimic, 2 nmol	Dharmacon	C-300653-05-0002
miRIDIAN microRNA Human hsa-miR-302b-3p - Mimic, 2 nmol	Dharmacon	C-300669-05-0002
miRIDIAN microRNA Human hsa-miR-509-3p - Mimic, 2 nmol	Dharmacon	C-300849-07-0002
miRIDIAN microRNA Human hsa-miR-210-3p - Mimic, 2 nmol	Dharmacon	C-300565-03-0002
miRIDIAN microRNA Human hsa-miR-302c-3p - Mimic, 2 nmol	Dharmacon	C-300671-05-0002
miRIDIAN microRNA Human hsa-let-7a-2-3p - Mimic, 2 nmol	Dharmacon	C-301297-00-0002
miRIDIAN microRNA Human hsa-miR-224-5p - Mimic, 2 nmol	Dharmacon	C-300581-07-0002
miRIDIAN microRNA Human hsa-miR-302d-3p - Mimic, 2 nmol	Dharmacon	C-300672-05-0002
miRIDIAN microRNA Human hsa-miR-520e - Mimic, 2 nmol	Dharmacon	C-300772-03-0002
miRIDIAN microRNA Human hsa-miR-520a-3p - Mimic, 2 nmol	Dharmacon	C-300788-03-0002
miRIDIAN microRNA hsa-miR-20a-3p mimic, 2 nmol	Dharmacon	C-301022-01-0002
miRIDIAN microRNA Human hsa-miR-498 - Mimic, 2 nmol	Dharmacon	C-300771-03-0002
miRIDIAN microRNA Human hsa-miR-665 - Mimic, 2 nmol	Dharmacon	C-301246-01-0003
miRIDIAN microRNA Mimic Negative Control #1, 5 nmol	Dharmacon	CN-001000-01-05

Acknowledgements

548   We thank other members of the Howley laboratory for helpful discussions and

549 suggestions. All plasmids generated in the Howley laboratory that are described in this

550 manuscript are available through Addgene. GMN, JAS and PMH are co-inventors on a patent

551 filed by Harvard on the therapeutic use of miRNAs in hrHPV positive cancers. This work

552 has been supported by NIH grant R35CA197262 to P.M.H.

Bibliography

1.   Schiller JT, Lowy DR. 2021. Papillomaviruses, p 45-67. *In* Knipe DM, Howley PM

(ed), Fields Virology, 7th ed, vol 2. Wolkers Kluwers, Philadelphia.

2.   de Martel C, Plummer M, Vignat J, Franceschi S. 2017. Worldwide burden of cancer

attributable to HPV by site, country and HPV type. Int J Cancer 141:664-670.

3.   Goodwin EC, DiMaio D. 2000. Repression of human papillomavirus oncogenes in

HeLa cervical carcinoma cells causes the orderly reactivation of dormant tumor

suppressor pathways. Proc Natl Acad Sci U S A 97:12513-8.

4.   Wells SI, Francis DA, Karpova AY, Dowhanick JJ, Benson JD, Howley PM. 2000.

Papillomavirus E2 induces senescence in HPV positive cells via pRB- and p21^{CIP}-

dependent pathways. EMBO J 19:5762-5771.

5.   Scheffner M, Werness BA, Huibregtse JM, Levine AJ, Howley PM. 1990. The E6

oncoprotein encoded by human papillomavirus types 16 and 18 promotes the

degradation of p53. Cell 63:1129-36.

6.   Huibregtse JM, Scheffner M, Howley PM. 1993. Cloning and expression of the cDNA

for E6-AP: A protein that mediates the interaction of the human papillomavirus E6

oncoprotein with p53. Mol Cell Biol 13:775-784.

7.   Huibregtse JM, Scheffner M, Howley PM. 1993. Localization of the E6-AP regions that

direct HPV E6 binding, association with p53, and ubiquination of associated

proteins. Mol Cell Biol 13:4918-4927.

8.   Scheffner M, Huibregtse JM, Vierstra RD, Howley PM. 1993. The HPV-16 E6 and E6-

AP complex functions as a ubiquitin-protein ligase in the ubiquination of p53. Cell

75:495-505.

9.   Park RB, Androphy EJ. 2002. Genetic analysis of high-risk e6 in episomal

maintenance of human papillomavirus genomes in primary human keratinocytes. J

Virol 76:11359-64.

10.   Butz K, Ristriani T, Hengstermann A, Denk C, Scheffner M, Hoppe-Seyler F. 2003.

siRNA targeting of the viral E6 oncogene efficiently kills human papillomavirus-

positive cancer cells. Oncogene 22:5938-3945.

11.   Chirayil R, Kincaid RP, Dahlke C, Kuny CV, Dalken N, Spohn M, Lawson B, Grundhoff

A, Sullivan CS. 2018. Identification of virus-encoded microRNAs in divergent

Papillomaviruses. PLoS Pathog 14:e1007156.

12.   Wald AI, Hoskins EE, Wells SI, Ferris RL, Khan SA. 2011. Alteration of microRNA

profiles in squamous cell carcinoma of the head and neck cell lines by human

papillomavirus. Head Neck 33:504-12.

13.   Wang X, Wang HK, Li Y, Hafner M, Banerjee NS, Tang S, Briskin D, Meyers C, Chow

590   LT, Xie X, Tuschl T, Zheng ZM. 2014. microRNAs are biomarkers of oncogenic human

591   papillomavirus infections. Proc Natl Acad Sci U S A 111:4262-7.

592   14.   Tornesello ML, Faraonio R, Buonaguro L, Annunziata C, Starita N, Cerasuolo A,

593   Pezzuto F, Tornesello AL, Buonaguro FM. 2020. The Role of microRNAs, Long Non-

594   coding RNAs, and Circular RNAs in Cervical Cancer. Front Oncol 10:150.

595   15.   Yen HC, Elledge SJ. 2008. Identification of SCF ubiquitin ligase substrates by global

596   protein stability profiling. Science 322:923-9.

597   16.   Yen HC, Xu Q, Chou DM, Zhao Z, Elledge SJ. 2008. Global protein stability profiling in

598   mammalian cells. Science 322:918-23.

599   17.   Hubbert NL, Sedman SA, Schiller JT. 1992. Human papillomavirus type 16 E6

600   increases the degradation rate of p53 in human keratinocytes. Journal of virology

601   66:6237-6241.

602   18.   Scheffner M, Takahashi T, Huibregtse JM, Minna JD, Howley PM. 1992. Interaction of

603   the human papillomavirus type 16 E6 oncoprotein with wild-type and mutant

604   human p53 proteins. J Virol 66:5100-5.

605   19.   Kamaraj B, Bogaerts A. 2015. Structure and Function of p53-DNA Complexes with

606   Inactivation and Rescue Mutations: A Molecular Dynamics Simulation Study. PLoS

607   One 10:e0134638.

608   20.   Olotu FA, Soliman MES. 2019. Dynamic perspectives into the mechanisms of

609   mutation-induced p53-DNA binding loss and inactivation using active perturbation

610   theory: Structural and molecular insights toward the design of potent reactivators in

611   cancer therapy. J Cell Biochem 120:951-966.

612   21.   Tomita Y, Marchenko N, Erster S, Nemajerova A, Dehner A, Klein C, Pan H, Kessler H,

613   Pancoska P, Moll UM. 2006. WT p53, but not tumor-derived mutants, bind to Bcl2

614   via the DNA binding domain and induce mitochondrial permeabilization. J Biol

615   Chem 281:8600-6.

616   22.   Gencel-Augusto J, Lozano G. 2020. p53 tetramerization: at the center of the

617   dominant-negative effect of mutant p53. Genes Dev 34:1128-1146.

618   23.   Jung HM, Phillips BL, Chan EK. 2014. miR-375 activates p21 and suppresses

619   telomerase activity by coordinately regulating HPV E6/E7, E6AP, CIP2A, and 14-3-

620   3zeta. Mol Cancer 13:80.

621   24.   Martinez-Noel G, Vieira VC, Szajner P, Lilienthal EM, Kramer RE, Boyland KA, Smith

622   JA, Howley PM. 2021. Live cell, image-based high-throughput screen to quantitate

623   p53 stabilization and viability in human papillomavirus positive cancer cells.

624   Virology 560:96-109.

625   25.   Munger K, Phelps WC, Bubb V, Howley PM, Schlegel R. 1989. The E6 and E7 genes of

626   the human papillomavirus type 16 together are necessary and sufficient for

627   transformation of primary human keratinocytes. J Virol 63:4417-21.

628   26.   Liu X. 2018. Up-regulation of miR-20a by HPV16 E6 exerts growth-promoting effects

629   by targeting PDCD6 in cervical carcinoma cells. Biomed Pharmacother 102:996-

630   1002.

631   27.   Scheffner M, Munger K, Byrne JC, Howley PM. 1991. The state of the p53 and

632   retinoblastoma genes in human cervical carcinoma cell lines. Proc Natl Acad Sci U S

633   A 88:5523-7.

28.   Abdelmohsen K, Srikantan S, Tominaga K, Kang MJ, Yaniv Y, Martindale JL, Yang X,

Park SS, Becker KG, Subramanian M, Maudsley S, Lal A, Gorospe M. 2012. Growth

inhibition by miR-519 via multiple p21-inducing pathways. Mol Cell Biol 32:2530-

48.

29.   Buehler E, Chen YC, Martin S. 2012. C911: A bench-level control for sequence

specific siRNA off-target effects. PLoS One 7:e51942.

30.   Jiang Y, Hou R, Li S, Li S, Dang G. 2018. MicroRNA-302 inhibits cell migration and

invasion in cervical cancer by targeting DCUN1D1. Exp Ther Med 16:1000-1008.

31.   Li GC, Cao XY, Li YN, Qiu YY, Li YN, Liu XJ, Sun XX. 2018. MicroRNA-374b inhibits

cervical cancer cell proliferation and induces apoptosis through the p38/ERK

signaling pathway by binding to JAM-2. J Cell Physiol 233:7379-7390.

32.   Mou Z, Xu X, Dong M, Xu J. 2016. MicroRNA-148b Acts as a Tumor Suppressor in

Cervical Cancer by Inducing G1/S-Phase Cell Cycle Arrest and Apoptosis in a

Caspase-3-Dependent Manner. Med Sci Monit 22:2809-15.

33.   Ji F, Du R, Chen T, Zhang M, Zhu Y, Luo X, Ding Y. 2020. Circular RNA circSLC26A4

Accelerates Cervical Cancer Progression via miR-1287-5p/HOXA7 Axis. Mol Ther

Nucleic Acids 19:413-420.

34.   Abdelmohsen K, Kim MM, Srikantan S, Mercken EM, Brennan SE, Wilson GM, Cabo R,

Gorospe M. 2010. miR-519 suppresses tumor growth by reducing HuR levels. Cell

Cycle 9:1354-9.

35.   Cai N, Wang YD, Zheng PS. 2013. The microRNA-302-367 cluster suppresses the

proliferation of cervical carcinoma cells through the novel target AKT1. Rna 19:85-

95.

36.   Chen Y, Wang X. 2020. miRDB: an online database for prediction of functional

microRNA targets. Nucleic Acids Res 48:D127-D131.

37.   Liu W, Wang X. 2019. Prediction of functional microRNA targets by integrative

modeling of microRNA binding and target expression data. Genome Biol 20:18.

38.   Kuhnle S, Martinez-Noel G, Leclere F, Hayes SD, Harper JW, Howley PM. 2018.

Angelman syndrome-associated point mutations in the Zn(2+)-binding N-terminal

(AZUL) domain of UBE3A ubiquitin ligase inhibit binding to the proteasome. J Biol

Chem 293:18387-18399.

39.   Kracikova M, Akiri G, George A, Sachidanandam R, Aaronson SA. 2013. A threshold

mechanism mediates p53 cell fate decision between growth arrest and apoptosis.

Cell Death Differ 20:576-88.

40.   Bartel DP. 2009. MicroRNAs: target recognition and regulatory functions. Cell

136:215-33.

41.   Shukla GC, Singh J, Barik S. 2011. MicroRNAs: Processing, Maturation, Target

Recognition and Regulatory Functions. Mol Cell Pharmacol 3:83-92.

42.   Lee RC, Feinbaum RL, Ambros V. 1993. The C. elegans heterochronic gene lin-4

encodes small RNAs with antisense complementarity to lin-14. Cell 75:843-54.

43.   Wightman B, Ha I, Ruvkun G. 1993. Posttranscriptional regulation of the

heterochronic gene lin-14 by lin-4 mediates temporal pattern formation in C.

elegans. Cell 75:855-62.

44.   Bonneau E, Neveu B, Kostantin E, Tsongalis GJ, De Guire V. 2019. How close are

miRNAs from clinical practice? A perspective on the diagnostic and therapeutic

market. Ejifcc 30:114-127.

45.   Hanna J, Hossain GS, Kocerha J. 2019. The Potential for microRNA Therapeutics and

681   Clinical Research. Front Genet 10:478.

682   46.   Emanuele MJ, Elia AE, Xu Q, Thoma CR, Izhar L, Leng Y, Guo A, Chen YN, Rush J, Hsu

683   PW, Yen HC, Elledge SJ. 2011. Global identification of modular cullin-RING ligase

684   substrates. Cell 147:459-74.

685   47.   Lin HC, Yeh CW, Chen YF, Lee TT, Hsieh PY, Rusnac DV, Lin SY, Elledge SJ, Zheng N,

686   Yen HS. 2018. C-Terminal End-Directed Protein Elimination by CRL2 Ubiquitin

687   Ligases. Mol Cell 70:602-613 e3.

688   48.   Zhang JH, Chung TD, Oldenburg KR. 1999. A Simple Statistical Parameter for Use in

689   Evaluation and Validation of High Throughput Screening Assays. J Biomol Screen

690   4:67-73.

691   49.   Ottinger M, Smith JA, Schweiger MR, Robbins D, Powell ML, You J, Howley PM. 2009.

692   Cell-type specific transcriptional activities among different papillomavirus long

693   control regions and their regulation by E2. Virology 395:161-71.

694   50.   Schweiger MR, Ottinger M, You J, Howley PM. 2007. Brd4-independent

695   transcriptional repression function of the papillomavirus e2 proteins. J Virol

696   81:9612-22.

Figure captions

700   **Figure 1. Primary screen.**

A. Reporter cassette of pHAGE-P CMVt RIG2 p53(R273C). The expressed open reading

frames are shown with the red and green arrows. CMVp: CMV promoter, IRES:

Encephalomyocarditis virus internal ribosomal entry site**. B.** Dot plot showing the effect of

the different miRNAs on individual experimental wells during the primary screen. Bars

indicate the means of triplicate experiments and the error bars show one standard

deviation. The split Y axis has two different scales to help with the visualization of the data.

Figure 2. p53 stabilization in HeLa cells expressing the reporter pHAGE-P CMVt RIG3

p53(R273C).

A. Reporter cassette in pHAGE-P CMVt RIG3 p53(R273C) (24). The expressed fusion

proteins are indicated with the red and green arrows. CMVp: CMV promoter, IRES:

Encephalomyocarditis virus internal ribosomal entry site, H2B: histone 2B (H2BC11), SG

linker: serine-glycine linker. **B.** 72 h after transfection cells were harvested and the

percentage of cells expressing mRuby-p53(R273C) was determined by flow cytometry.

Columns indicate the means of three independent experiments and the error bars show

one standard deviation. Variability between the means was tested using the one-way

ANOVA test ($F_{(40, 82)}$ = 84.19 p<0.0001). Then, each sample was compared to non-treated

cells using the "Dunnett's multiple comparisons test" (* = p<0.01).

Figure 3. Effect of miRNAs transfection on p53, E6AP, and E6 protein levels in HeLa

cells.

Cells were lysed 72 h post-transfection and the lysates were resolved by SDS-PAGE. Protein

levels of p53, E6AP, E6, and actin were visualized by western blotting. **A, C, and E.** Western

blot results from a single experiment. **B, D, and F.** Quantification of p53 from western blot

experiments. Values obtained for p53 were normalized to the amount of actin in the same

sample and then all values were divided by the corresponding value obtained with the

mock transfection in the same blot. Circles represent values from independent experiments

and the bars show the mean ± standard deviation. To determine significant differences

between the samples in an experiment we used the "one-way ANOVA" test (A. $F_{(16, 34)}$=8.271 $p<0.0001$, n=3; B. $F_{(14, 30)}$=17.76 $p<0.0001$, n=3; C. $F_{(3, 12)}$=52.63 $p<0.0001$,

34)=8.271 p<0.0001, n=3; B. F(14, 30)=17.76 p<0.0001, n=3; C. F(3, 12)=52.63 p<0.0001,

n=4). We then compared each sample with the cells transfected with the Ctrl.1 miRNA

using the "Dunnett's multiple comparisons test" (* = $p<0.01$).

Figure 4. Quantification of the effect of miRNAs transfection on E6AP and E6 protein

levels in HeLa cells.

Quantification of E6AP (**A-C**) and E6 (**D-F**) protein levels from western blot experiments.

The values for E6AP and E6 were normalized to the amount of actin in the same sample,

then all values were divided by the corresponding value of the mock transfection in the

same blot. Circles show the values of independent experiments and the bars indicate the

mean ± standard deviation. The significance of differences between the samples means

was determined using the one-way ANOVA test (A. E6AP protein levels $F_{(16, 34)}$=9.540

$p<0.0001$, n=3; B. E6AP protein levels $F_{(14, 30)}$=1.218 $p<0.3135$, n=3; C. E6AP protein

levels $F_{(3, 12)}$=11.16 $p<0.0009$, n=4 D. E6 protein levels $F_{(16, 34)}$=8.672 $p<0.0001$, n=3; E.

E6 protein levels $F_{(14, 30)}$=18.15 $p<0.0001$, n=3; F. E6 protein levels $F_{(3, 12)}$=55.35

$p<0.0001$, n=4). Each sample was then compared to the cells transfected with the Ctrl.1

miRNA using the "Dunnett's multiple comparisons test" (* = $p<0.01$). **G.** The protein levels

of p53, and E6 calculated from the western blot experiments were used to analyze whether

there is a significant correlation between the levels of these proteins after transfection of

HeLa cells with the different miRNAs. This was done by calculating the Pearson's

correlation coefficient (r) and a two tailed p value. **H.** Heat map summarizing the effect of

the miRNAs on the protein levels of E6, E6AP, and p53 relative to the cells transfected with

Ctrl.1 in the corresponding gel after averaging the quantifications of the western blot

experiments showed in Figures 3 and 4, and Table S2).

Figure 5. Alignment of members of the 302/519 family of miRNAs.

Several of the miRNAs identified in our screen have the same seed sequence indicated with

bold letters and underlined. The sequences of the miRNAs were aligned based on the most

common seed sequence found in this family framed with a dashed blue line. miRNAs of this

family present between the 32 miRNAs selected from the primary screen are indicated

with black letters.

Figure 6. Effect of transfection of selected miRNAs on p53 and E6AP protein levels in

different cell lines.

72 h after transfection cells were lysed and the protein extracts were resolved by SDS-

PAGE, and the p53, E6AP, and actin proteins levels were determined by western blotting.

Figure 7. Effect of selected miRNAs on HPV E6 and E7 expression.

Cells were harvested 72 h post-transfection and then used for the following experiments. **A.**

Western blot showing the effect of the selected miRNAs on E6, E7, and actin protein levels

in HeLa cells. **B.** Expression of the E6/E7 mRNAs after transfection of the selected miRNAs

was quantitated by QRT-PCR. Measures from each experiment were normalized to the

value of the corresponding mock transfection. Differences between samples were analyzed

by the "one-way ANOVA" test (HeLa $F_{(12, 39)}$ = 42.32 p<0.0001, n=4; SiHa $F_{(12, 26)}$ =

41.38 p<0.0001, n=3) , then each sample was compared to the corresponding non-treated

cells using the "Dunnett's multiple comparisons test" (* = p<0.01). **C.** Western blot showing

the effect of transfecting the indicated miRNAs into 1321 cells on the p53, E6AP, and actin

proteins levels after 72 h incubation. **D.** Effect of transfection of selected miRNAs on the

promoter activity of the HPV16 and HPV18 LCRs using a luciferase reporter assay in C33A

cells. Values from each experiment were normalized to the value obtained for the

corresponding mock transfection. Differences between samples were analyzed by the "one-

way ANOVA" test (HPV 16 LCR, $F_{(12, 38)}$ = 34.91 p<0.0001, n=4; HPV 18 LCR, $F_{(12, 39)}$ =

23.47 p<0.0001, n=4), then each sample was compared to the corresponding non-treated

cells using the "Dunnett's multiple comparisons test" (* = p<0.01). In B and D, circles

represent values from independent experiments and the bars show the mean ± standard

deviation.

Figure 8. Transfection of selected miRNAs induces apoptosis in HeLa cells.

A. Cells were harvested 72 h post-transfection and stained with Annexin V as an indicator

of apoptosis. The percentage of cells stained with Annexin V was determined by flow

cytometry. Variability between samples was first analyzed with the one-way ANOVA test

($F_{(12,26)}$=48.04 p<0.0001, n=3). Then each sample was compared to the non-treated cells

with the "Dunnett's multiple comparisons test" (* = p<0.01). Circles represent values from

independent experiments and the bars show the mean ± standard deviation. **B.** Cells were

transfected with the indicated miRNAs and 72 h later the cells were harvested and stained

with Annexin V as a marker for apoptosis. The percentage of Annexin V positive cells was

quantitated by flow cytometry. Variability between samples for each cell line was analyzed

with the one-way ANOVA test (SiHa $F_{(6,21)}$=45.17 $p<0.0001$, n=4; C33A $F_{(6,21)}$=85.96

$p<0.0001$, n=3; U2OS $F_{(6,21)}$=104.07 $p<0.0001$, n=4). Each sample was then compared to

the non-treated cells with the "Dunnett's multiple comparisons test" (* = $p<0.01$). Circles

represent values from independent experiments and the bars show the mean ± standard

deviation. **C.** Effect of the indicated miRNAs on the p21 and actin proteins levels HeLa and

SiHa cells visualized by western blotting 72 h post transfection.

Figure 9. p53 dependence of apoptotic effect of selected miRNAs in HeLa cells.

A. Cells were transfected with the indicated miRNAs either alone or together with a siRNA

against p53 or its C911 variant. After incubation for 72 h the cells were harvested and

stained with Annexin V as a marker for apoptosis. The percentage of Annexin V stained

cells was determined by flow cytometry. The values obtained for the three transfections

with each miRNA were first analyzed with the one-way ANOVA test (375-3p $F_{(2, 6)}$ = 3.075

$p=0.1204$; 498-5p $F_{(2, 6)}$ = 1.914 $p=0.2276$; 20a-3p $F_{(2, 6)}$ = 0.6553 $p=0.5528$; 148b-5p

$F_{(2, 6)}$ = 23.57 $p=0.0014$; 374a-3p $F_{(2, 6)}$ = 2.681 $p=0.1472$; 519b-3p $F_{(2, 6)}$ = 17.71

$p=0.0030$; 1287-5p $F_{(2, 6)}$ = 48.00 $p=0.0002$). For all samples n=3. Then, the values from

samples transfected with the same miRNA were compared with each other using the

"Tukey's multiple comparisons test" (* = $p<0.01$). **B.** 72 h post-transfection cells were lysed

and the p53, cleaved PARP, and GAPDH proteins levels were determined by SDS-PAGE and

western blotting.

A.

B.

A.

B.

302a-3p	U**AAGUGC**UUCCAUGUUUUGGUGA
302b-3p	U**AAGUGC**UUCCAUGUUUUAGUAG
302c-3p	U**AAGUGC**UUCCAUGUUUCAGUGG
302d-3p	U**AAGUGC**UUCCAUGUUUGAGUGU
372-3p	A**AAGUGC**UGCGACAUUUGAGCGU
373-3p	G**AAGUGC**UUCGAUUUUGGGGUGU
519a-3p	A**AAGUGC**AUCCUUUUAGAGUGU
519b-3p	A**AAGUGC**AUCCUUUUAGAGGUU
519c-3p	A**AAGUGC**AUCUUUUUAGAGGAU
519d-3p	C**AAAGUG**CCUCCCUUUAGAGUG
519e-3p	A**AGUGCC**UCCUUUUAGAGUGUUU
520a-3p	A**AGUGC**UUCCCUUUGGACUGU
520b-3p	A**AAGUGC**UUCCUUUUAGAGGG
520c-3p	A**AAGUGC**UUCCUUUUAGAGGGU
520d-3p	A**AAGUGC**UUCUCUUUGGUGGGU
520e-3p	A**AAGUGC**UUCCUUUUUGAGGG
520f-3p	A**AGUGCU**UUCCUUUUAGAGGGUU
520g-3p	AC**AAAGU**GCUUCCCUUUAGAGUGU

A.

B.