Identification of genetic heterogeneity in a South Indian family affected by primary open angle glaucoma.

Mohd Hussain Shah[1*], Manojkumar Kumaran[2,5*], Prakash Chermakani[1,6] Mohideen Abdul Kader[3], R. Ramakrishnan[3], Subbiah. R. Krishnadas[4], Bharanidharan Devarajan[2@], and Periasamy Sundaresan[1@]

[1]Department of Genetics, Aravind Medical Research Foundation, Madurai, Tamil Nadu, India

[2]Department of Bioinformatics, Aravind Medical Research Foundation, Madurai, Tamil Nadu, India

[3]Glaucoma Clinic, Aravind Eye Hospital, Tirunelveli, Tamil Nadu, India

[4]Glaucoma Clinic, Aravind Eye Hospital, Madurai, Tamil Nadu, India

[5]School of Chemical and Biotechnology, SASTRA (Deemed to be University), Thanjavur, Tamil Nadu, India

[6]Department of Molecular Biology, Alagappa University, Karaikudi, Tamil Nadu, India

[*]First two authors contributed equally to this work

[@]Corresponding Authors.

@ Corresponding Authors: Bharanidharan, Devarajan E-Mail: bharani@aravind.org Telephone: 91-452-4356550 Extn.444

Periasamy, Sundaresan E-Mail: sundar@aravind.org Telephone: 91-4356550 Extn.423

Abstract

Primary open angle glaucoma (POAG) is a complex hereditary disorder; larger pedigree analysis of POAG positive family history has led to the discovery of many candidate genes associated with disease pathogenesis. However, only 5-6% of POAG can explain these genetic factors. In the present study, we recruited a single south Indian family with 84 family members over five-generation with an Egyptian origin had a positive family history of POAG from Kayalpatanam, Tamil Nadu, India. All the study participants had a comprehensive ocular evaluation (of the 84, 19 study subjects were diagnosed as POAG). Sanger sequencing of the candidate genes associated with POAG (*MYOC, OPTN and TBK1*) showed no genetic variation in the POAG affected family members. Therefore, we performed whole exome sequencing (WES) for 16 samples including (9 POAG and 7 unaffected controls). Samples were selected based on the inheritance pattern observed in the large pedigree of the South Indian family. On the whole, this study identified six pathogenic variants and all the variants showed consistent co-segregation with POAG affected individuals. Among the six, a novel pathogenic variant in Rho guanine nucleotide exchange factors Gene 40 (ARHGEF40) and a genetic variant in Retinitis Pigmentosa GTPase regulator-interacting protein1 (RPGRIP1) showed significant association with glaucoma phenotype suggested that the genetic complexity in POAG requires NGS platform to understand the factors underlying disease pathogenesis.

1. Introduction

Glaucoma is typically characterized by a progressive degeneration of optic nerve, which causes irreversible blindness. It is the second leading cause of global blindness after cataract [1]. Primary open-angle glaucoma (POAG) is a subset of glaucoma majorly associated with loss of retinoganglion cells and their axons triggers permanent vision loss with an apprehensive exponential growth affecting around 60.5 million people worldwide. Due to the exponential increase in the global aging population, it is estimated that 80 million people will be affected by POAG by the end of 2020 [2] and the count could be expected to rise 111.8 million people by 2040. Which has an inexplicable impact on Asian and African peoples [3]. Asia alone accounted for approximately 60% of the global glaucoma, whereas the Africa population represented (13%) is the second largest proportion of glaucoma cases in the world. To date in India, it is estimated that 12 million people have been affected by glaucoma and this number is expected to increase by 16 million by the end of 2020 [4, 5].

POAG is associated with several external risk factor including advanced age, central corneal thickness, myopia, steroid responsiveness and elevated intraocular pressure (IOP) [6]. However, these risk factors do not capture the full spectrum of the disease. Though, the positive family history is also one of the risk factors for POAG. Genetic characterization of the POAG Positive family history, is useful for the identification of POAG-candidate genes (MYOC, OPTN, & TBK1) [6-8] that are capable of causing POAG. However, these candidate genes were discovered through large pedigrees with a positive family history of glaucoma. In addition, many studies have shown that POAG development is associated with various genetic risk factors, including genetic variants in CDKN2B-AS [9-12] CAV1/CAV2 [13], TMCO1 [14]

AFAP1 [15], TXNRD2, FOXC1/GMDS, ATXN2 [16], FNDC3B [17, 18], GAS [14], PMM2 [19], TGFBR3 [20], and SIX1/SIX6 [10, 11]. Due to its genetic heterogeneity and familial clustering nature of consistent genetic inheritance, necessitated an extensive molecular characterization to identify the factors responsible for the genetic predisposition in POAG affected individuals.

Our previous report, suggested that genetic screening of known candidate genes *(MYOC, OPTN* and *TBK1)* in a single large South Indian family with POAG did not detect the genetic risk factors underlying the pathogenesis of the disease [21]. Therefore, this study aims to perform whole exome sequencing (WES), to identify the potential genetic risk factors associated with the positive POAG family history of the five-generation south Indian family.

2. Materials and Methods

2.1 POAG study subjects

The study was approved by the Institutional Review Board at the Aravind Eye Care System, Madurai, Tamil Nadu, India (IRB2011008BAS). This research adhered the tenets of the Declaration of Helsinki. All the study subjects were recruited and clinically evaluated as previously described [21]. Briefly, an ophthalmic examination was conducted for 240 subjects during a field trip to Kayalpatanam, for this current study 84 members were recruited from a single family of a large South Indian family of five generations with a positive history of POAG.

2.2 DNA isolation and quantification

5 ml of peripheral blood was collected in an anticoagulant (EDTA) vacutainer from each study subjects. The genomic DNA was extracted using a salting-out precipitation method [22] and the concentration of the DNA samples was quantified using Qubit fluorometer.

2.3 Whole exome sequencing

Samples were subjected for WES using the Agilent's SureSelect Human All Exon V6 kit. WES was performed in Centogene, Germany. The DNA libraries have been sequenced to mean >150X coverage on an Illumina HiSeq 4000 platform. The coding regions of all 20,000 genes were captured with RNA baits approximately 60MB (targeting >99% of regions in CCDS, RefSeq, and Gencode databases) for generating 300 bp paired-end reads. The DNA library of each sample was constructed by shearing (3 μg) of intact genomic DNA into 300-350 bp DNA fragments using Covaris. The quality of the fragmented DNA was checked using Bioanalyzer 2100. Then the fragmented DNA was end-repaired and dA-tailed using SureSelect XT library kit, followed by the ligation of paired-end adapters and the DNA library was enriched through PCR amplification using adaptor specific primers. Each step was followed by purification of the AMPure XP beads. Always, the quality of the DNA library was assessed using a bioanalyzer. After enrichment, the DNA libraries were pooled in equal quantity to give ~500 ng and hybridised with biotinylated RNA bait, after hybridization the biotinylated probes were captured using streptavidin-coated beads and finally the captured DNA library was amplified with 8-bp indexing primers and cleaned using AMPure XP beads for massive parallel paired end sequencing.

2.4 Data Analysis

We developed an automated pipeline (Supplementary Figure.1) for the identification of pathogenic variant from WES data using UNIX script (https://github.com/bharani-lab/WES-pipelines/tree/master/Script). First, raw reads (FASTQ file) were processed to remove the adapter and low-quality sequences using

Cutadapt. The processed reads were further aligned against the human genome build GRCh37 using BWA-mem version 0.7.12. GATK version 4.1.0. was used to identify single-nucleotide variant (SNV) and small Insertion and Deletion (InDel) followed by annotation using ANNOVAR [23]. Variants were prioritized by setting the minor allele frequency (MAF) less than or equal to 0.5% in 1000genome, ESP, ExAC and genomeAD. Next, all exonic variants, including non-synonymous, frameshift and splice variants, were filtered. For pathogenic variant identification, the non-synonymous variants were subsequently filtered by a two-step process; firstly, variants were selected with the conservation score >2.5 (GERP score) and CADD score greater than 10; secondly, the variants should be predicted to be deleterious with at least three prediction tools among the five (Polyphen2, SIFT, Mutation Taster, FATHMM and LRT). To avoid the mapping errors, all the variants were manually checked with IGV viewer. All predicted deleterious variants were further filtered based on their presence in at least more than three affected individuals in the pedigree. Finally, the variants were sorted out by their presence in number of affected individuals in the family members and their association with glaucoma phenotype. We used VarElect software [24] to sort the genes based on their direct or indirect association with a glaucoma disease.

We performed pathway and gene ontology analysis using DAVID for all the genes identified in the final set of variants. A gene network was created using Cytoscape with the enriched pathways and biological processes.

2.5 Sanger sequencing for the validation of novel variant

The novel variant of the ARHGEF40 gene was validated by Sanger sequencing. For PCR amplification, following gene specific primers were used (FW-5′-CTGAGCTGACGCCTGAACTT-3′);

(RV-5′-GCCGTGGGTACTGAGAAAG- 3′). 100 ng of genomic DNA template was PCR amplified with 1X Taq buffer, 200 μM DNTP's, 5 Pm of forward and reverse primer and 1 U of Taq DNA polymerase (Sigma Aldrich) under the PCR conditions with initial denaturation 10 minutes at 96°C, followed by 37 cycles at 96°C for 30 seconds, annealing at 58°C for 30 seconds, and final extension at 72°C for 5 mins (Applied Biosystems-Invitrogen). After amplification, the PCR amplicon was purified with QIA quick PCR purification kit (Bio Basic Inc.,) and cyclic PCR was performed for bi-directional sequencing using (3130 Genetic Analyser; Applied Biosystems) and the results were compared with the reference sequence of ARHGEF40 gene using NCBI-BLAST program and the chromatogram was analysed using Chromas lite (2.1) software.

3. Results

3.1 Clinical Evaluation of patients

A total of 84 family members were recruited from a single large south Indian family of five generation with a positive family history of POAG after a comprehensive ophthalmic screening of 240 family members in Kayalpatanam (as shown in the Supplementary Figure 2). Clinical assessment and complex pedigree analysis revealed that 19 of the 84 samples had been diagnosed with POAG. The clinical features of all 19 POAG affected individuals were discussed in detail [21]. Only 9 out of 19 clinically diagnosed POAG individuals were subjected for Whole exome sequencing

3.2 Exome Sequencing and Variant Filtering

Samples for the WES were selected solely on the basis of the POAG inheritance pattern observed in the pedigree of the South Indian family, which included nine POAG cases belonging to the generation (II- 2,5,15; III- 2,3,19,32 and IV- 26,27) and seven unaffected controls (II- 5; III- 4,16,34,41 and IV- 11,28) (Supplementary Figure 2).

WES was carried out using Agilent SureSelect Human All Exon V6 kit and the DNA libraries have been sequenced to the Illumina HiSeq 4000 platform with an average coverage depth of ~150x. The raw data were processed and analysed to identify the pathogenic variants (as indicated in the methods section of Figure 1). Approximately 60,000 variants (SNV and InDel) were identified in the exome of each patient's aligned to the human reference genome build GRCh37.

3.3 Pathogenic variants

WES results revealed that pathogenic variants were identified in the six of the sixteen samples were co-segregated among the POAG affected family members (Supplementary Figure 2). Over all, we found six pathogenic variants (RPGRIP1; ARHGEF40; OR11H12; OR4K14; RNASE13 and OR11G2) (5 non-synonymous and one frameshift variant, respectively); all of which showed consistent co-segregation over generations in POAG affected individuals. Of the six pathogenic variants, RPGRIP1 alone showed direct association with glaucoma (Table.1). A heterozygous variant with MAF less than 0.01 in Retinitis Pigmentosa GTPase regulator-interacting protein1 (RPGRIP1) (c. G3719A; p. G1240E) had a deleterious effect exhibit highest phenotype score (Varelect score 8.35). Moreover, this variant also co-segregated in six POAG cases, which may affect the function of RPGR protein complex. In contrast, the RPGRIP1 variant (G1240E) is relatively common in most populations especially in Africans [25]. A novel variant in Rho guanine nucleotide exchange factors Gene 40 ARHGEF40 gene (c. A1295G; p. Q432R), which was also segregated within the POAG affected individuals in the family. In addition, the novel variant of the ARHGEF40 gene was validated (c. A1295G; p. Q432R) in other family members through Sanger sequencing represented the genetic association in eight POAGs diagnosed and two unaffected family members (Supplementary Figure 2). Though the

other four genetic variants of exome sequencing found consistent co-segregation OR11G2 (c.847delC; p. H282fs), OR4K14 (c. A355G; p. M119V), RNASE13 (c.C338T; p.S113F) and OR11H12 (c.T719G; p.V240G), but it did not show any association with glaucoma. In addition to these, this study also identified 54 pathogenic variants encoded by 51 different genes, but these genes are not co-segregated across the affected individuals. Among the 54 pathogenic variants, 52 were missense and 2 were Indel variants with frame-shifting coding regions as shown in Table 2. Based on the glaucoma clinical phenotype, the pathogenic variants, that are inconsistently segregating was also further prioritized in the POAG affected individual. The genetic variant in RPGR gene gene (c.G3430A; p.V1144I) was found in four POAG family members belongs to the generation (III-3;III-2;II-2;II-15) displayed it may affects its protein partner RPGRIP1 in the RPGR proteasome complex [26] and, eventually, PLK4 gene mutation in (c.C2513T; p.T838I) has adverse impact on microcephaly, failure of growth, and retinopathy [27]. PLK4 mutations microcephaly, failure of growth, and retinopathy. Surprisingly, six of the 54 non-segregating pathogenic variants were identified as novel variants. Of which, the genetic variant in the neural cell adhesion molecule 1 (NCAM1) gene (c. A1841 T; p. D614V) was further confirmed by Sanger sequencing, since the genetic mutation in NCAM1 gene alters the optic nerve function which is further associated with increase in intraocular pressure[28].

3.4 Functional network analysis

A functional network has been developed for all the pathogenic variants identified in POAG affected individuals, to investigate the pathways and biological processes involved in glaucoma pathogenesis. Initially, DAVID database was used to integrate all genes with KEGG pathways and Gene Ontology (GO) process. A total of 57 genes

were significantly enhanced by three pathways, and 17 GO biological processes (P < 0.01). These pathways include Focal adhesion, ECM-receptor interaction and PI3K-Akt signalling pathway. Further Gene-functional network (Figure. 2), analysis revealed that NCAM1, LAMB4 and PDGFRA genes connected all the three pathways to the other GO process. Of these genes, NCAM1 was shown to be linked to RPGRIP1 and ARHGEF40 with pathogenic variants through RPGR protein interaction and GO processes of positive regulation of GTPase activity and visual perception.

4. **Discussion**

Studies of larger pedigree in POAG diagnosed families led to the discovery of mutation in MYOC, OPTN and TBK1genes. Ophthalmic examination of a single family in South India with 84 family members over five generations with Egyptian heritage had a positive POAG family history revealed no mutation for the primary candidate genes (MYOC, OPTN and TBK1) associated with POAG [21].

Hence WES of 16 samples including (9 POAG and 7 unaffected controls) of the 84 family members displayed a consistent co-segregation of six pathogenic genes ARHGEF40, RPGRIP1, OR4K14, RNASE13, OR11H12, and OR11G2 in six POAG samples. No pathogenic variants have been identified in 3 of the 9 POAGs and the remaining 7 unaffected individuals. Furthermore, candidate genes for the 3 individuals diagnosed with POAG can be identified through either deep intronic or whole genome sequencing. All the pathogenic variants identified from WES was further prioritized based on the glaucoma phenotype using an insilico VarElect phenotype sorting tool. Among the six co-segregating pathogenic variants, only two (ARHGEF40 and RPGRIP1) showed association with glaucoma. One of the key interesting fact is all six

pathogenic variants were present in chromosome 14q, which had previously been reported to have potential POAG loci [10, 29, 30].

The pathogenic variant in retinitis pigmentosa GTPase regulator-interacting protein 1 (RPGRIP1) gene is observed with the highest phenotype score in six POAG cases, suggesting that it may have prominent role in POAG disease pathogenicity. Ferná ndez-Martı́ nez et al., has shown that the heterozygous non-synonymous variants in C2 domain of RPGRIP1gene might cause the various forms of glaucoma including POAG [25]. In addition, it has also demonstrated that RPGRIP1 interaction with NPHP4 protein was shown to play a key role in glaucoma pathogenesis [25]. In this study, four POAG cases were found to have a heterozygous pathogenic missense variant in the RPGR gene. In contrast to this study homozygous or compound heterozygous variants detected in RPGRIP1 are also associated to photoreceptor dystrophies [31, 32]. Interestingly, we observed a pathogenic variant in RPGR gene, which is existed in four POAG cases. RPGRIP1 and its interacting partner RPGR, have been shown to express in human retina and also outside of the retina [26, 33-35] may regulate cilia genesis, maintenance, and function mainly through signalling pathways [36]. Luo et al., 2014 has reported that the primary cilia of trabecular meshwork (TM) mediates intraocular pressure regulation through signalling pathway in the eye, and further highlighted that defect in the signalling pathway leads to Lowe syndrome that developed congenital glaucoma at birth [37]. RPGR and its protein partners play an important role in actin cytoskeleton remodelling of cilia through these signalling pathways by activating the small GTPase, RhoA [38].

This research also identified a novel pathogenic variant in ARGHEF40 gene and this variant was further confirmed in all the affected family members using Sanger sequencing (Supplementary Figure 3). Study show that Rho guanine nucleotide

exchange factors Gene Family protein (ARHGEF12) has been implicated as a risk factor of glaucoma by increasing intraocular pressure through RhoA/RhoA kinase pathway [39]. Furthermore, the activation of the Rho/ROCK pathway results in trabecular meshwork (TM) contraction, and the inhibition of this pathway would aggravate relaxation of TM with a consequent increase in outflow facility and, thereby, decrease intraocular pressure [40]. In the present study, we speculate that ARGHEF40 variant may affect the RhoA signalling through RPGRIP1 and its interacting partner RPGR in actin cytoskeleton remodelling of trabecular meshwork (TM) cilia, which may subsequently increase the intraocular pressure.

The pathogenic variants detected in other genes have not been reported to be directly associated with POAG. Therefore, we constructed a network of genes using GO and pathway enrichment. We have shown three pathways Focal adhesion, ECM-receptor interaction and PI3K-Akt signalling pathway to be associated with the pathogenesis of POAG. Furthermore, the highlighted genes ARHGEF40, RPGRIP1 and RPGR were enriched through visual perception and positive regulation of GTPase activity. Intriguingly, the genes NCAM1, HSP1 and PDGFRA including ARHGEF40 and RPGR in the biological process of positive regulation of GTPase activity is prioritized as top pathogenic variants based on the phenotype score. A study has shown that neural cell adhesion molecule (NCAM) participate in the optic nerve changes associated with elevated intraocular pressure [28].

5. **Conclusion**

Overall, this study presented a panel of pathogenic variants in multiple genes, and their possible association with POAG pathogenesis in the five generation South Indian family. Thus, our findings strongly suggested that WES of the five-generation South Indian family showed extreme genetic heterogeneity of POAG within the family and

the identified pathogenic variants showed continuous co-segregation among POAG affected individuals. Pathway analysis also displayed the association of the candidate genes involved in POAG pathogenesis but it requires larger sample size to ensure the authentic association of these identified genetic variants in POAG affected individuals. Even though our study reported rare pathogenic variants in multiple genes and the pathways associated with the large family in POAG affected individuals, epigenetic changes and copy number variations may be explored to understand the incredible complexity of the POAG pathogenesis.

Funding details:

This research was supported by Indian Council of Medical Research, Government of India (2012-0383/F1)

Acknowledgments

We thank all the patients and their families for their participation in this study. We are grateful to the Indian Council of Medical Research (ICMR) for their financial support. We are also thankful to Mr. Saravanan, Mrs. Kalarani and Mrs. Muthu Selvi for helping us in recruiting the samples and maintaining the clinical data in our lab.

Ethics approval and consent to participate

The study adhered to the tenets of the Declaration of Helsinki, and ethics committee approval was obtained from the Institutional Review Board of the Aravind Eye Care System (IRB2011008BAS). All study participants read and signed informed consent after explaining the nature and possible significances of the study.

Declaration of interest

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of this article.

Author's contributions

Mohd Hussain Shah done the sample preparation, sanger sequencing and wrote the manuscript. Manojkumar Kumaran analysed the whole exome sequencing data and wrote the manuscript. Prakash chermakani wrote the manuscript. Periasamy Sundaresan and Bharanidharan Devarajan designed the work and co-wrote the manuscript. Mohideen Abdul Kader, R Ramakrishnan and Subbiah R Krishnadas assisted in clinical diagnosis and sample collection. All authors read and approved the final manuscript.

Availability of data and material

The pipeline lines used for analysis in this study and a detailed tutorial is openly available for the public at (https://github.com/bharani-lab/Wole-Exome-Analysis-Pipeline).

We have submitted the data that support the finding of this study to SRA project ID PRJNA555016. Data can be accessed upon request

Supportive/Supplementary Materials

Figure Legends

Figure 1. Work Flow for variant prioritization

Figure 2. Functional network enriched with pathways and Gene Ontology (GO) on genes identified with pathogenic variants. Hexagon represents the gene; rectangle represent pathways and Diamond represents GO.

Table Legends

Table.1. List of the Pathogenic variants co- segregated with phenotype. Varlect score with symbol † represents the direct association with glaucoma phenotypes and ‡ represent the indirect association. * represent the Novel variant.

Table 2. List of pathogenic variants. Varlect score with symbol † represents the direct association with glaucoma phenotypes, and ‡ represent the indirect association. * represent the Novel variant.

Supplementary Figure Legends

Supplementary Figure 1. Modular Pipeline

Supplementary Figure 2. Pedigree from south India Family. Family members diagnosed with POAG are shaded with black.

Supplementary Figure 3. Pedigree of selected family members from large south India family as shown in supplementary figure 1 (A). Sanger sequencing results of novel variant c.A1295G in ARGEF40 gene (marked with down arrow). The variant is detected in the family members II-2, III-1, III-2 and III-3 (B).

Reference

[1] S. Resnikoff, D. Pascolini, D. Etya'ale, I. Kocur, R. Pararajasegaram, G. Pokharel, S. Mariotti, Global data on visual impairment in the year 2002, Bulletin of the World Health Organization 82 (2004) 844-51.

[2] H. Quigley, A.T. Broman, The number of people with glaucoma worldwide in 2010 and 2020, The British journal of ophthalmology 90 (2006) 262-7.

[3] Y.-C. Tham, X. Li, T.Y. Wong, H. Quigley, T. Aung, C.-y. Cheng, Global Prevalence of Glaucoma and Projections of Glaucoma Burden through 2040 A Systematic Review and Meta-Analysis, Ophthalmology 121 (2014).

[4] B. Thylefors, A.D. Négrel, R. Pararajasegaram, Y. Dadzie, Global Data on Blindness, Bulletin of the World Health Organization 73 (1995) 115-21.

[5] L. Vijaya, R. George, M. Baskaran, H. Arvind, P. Sriram, R. S Ve, G.

Kumaramanicakvel, C. McCarty, Prevalence of Primary Open-angle Glaucoma in an Urban South Indian Population and Comparison with a Rural Population The Chennai Glaucoma Study, Ophthalmology 115 (2008) 648-654.e1.

[6] J. Fingert, Primary open-angle glaucoma genes, Eye (London, England) 25 (2011) 587-95.

[7] T. Rezaie, A. Child, R. Hitchings, G. Brice, L. Miller, M. Coca-Prados, E. Héon, T. Krupin, R. Ritch, D. Kreutzer, R. Crick, M. Sarfarazi, Adult-Onset Primary Open-Angle Glaucoma Caused by Mutations in Optineurin, Science (New York, N.Y.) 295 (2002) 1077-9.

[8] E. Stone, J. Fingert, W. Alward, T. Nguyen, J. Polansky, S. Sunden, D. Nishimura, A. Clark, A. Nystuen, B. Nichols, D. Mackey, R. Ritch, J. Kalenak, E. Craven, V. Sheffield, Identification of a Gene That Causes Primary Open Angle Glaucoma, Science (New York, N.Y.) 275 (1997) 668-70.

[9] M. Nakano, Y. Ikeda, Y. Tokuda, M. Fuwa, N. Omi, M. Ueno, K. Imai, H. Adachi, M. Kageyama, K. Mori, S. Kinoshita, K. Tashiro, Common Variants in CDKN2B-AS1 Associated with Optic-Nerve Vulnerability of Glaucoma Identified by Genome-Wide Association Studies in Japanese, PloS one 7 (2012) e33389.

[10] W. Osman, S.-K. Low, A. Takahashi, M. Kubo, Y. Nakamura, A genome-wide association study in the Japanese population confirms 9p21 and 14q23 as susceptibility loci for primary open angle glaucoma, (2018).

[11] W. Ramdas, D. Rizopoulos, R. Wolfs, A. Hofman, P. Jong, J. Vingerling, N. Jansonius, Defining Glaucomatous Optic Neuropathy from a Continuous Measure of

Optic Nerve Damage – The Optimal Cut-off Point for Risk-factor Analysis in Population-based Epidemiology, Ophthalmic epidemiology 18 (2011) 211-6.

[12] J. Wiggs, The Cell and Molecular Biology of Complex Forms of Glaucoma: Updates on Genetic, Environmental, and Epigenetic Risk Factors, Investigative ophthalmology & visual science 53 (2012) 2467-9.

[13] G. Thorleifsson, G. Walters, A. Hewitt, G. Masson, A. Helgason, A. Dewan, A. Sigurdsson, A. Jonasdottir, S. Gudjonsson, K. Magnusson, H. Stefansson, D. Lam, P. Tam, G. Gudmundsdottir, L. Southgate, K. Burdon, M. Gottfredsdottir, M. Aldred, P. Mitchell, K. Stefansson, Common variants near CAV1 and CAV2 are associated with primary open-angle glaucoma, Nature genetics 42 (2010) 906-9.

[14] L. Koolwijk, W. Ramdas, K. Ikram, N. Jansonius, F. Pasutto, P. Hys, S. Macgregor, S. Janssen, A. Hewitt, A. Viswanathan, J. ten Brink, S. Hosseini, N. Amin, D. Despriet, J. Willemse-Assink, R. Kramer, F. Rivadeneira, M. Struchalin, Y. Aulchenko, C. Duijn, Common Genetic Determinants of Intraocular Pressure and Primary Open-Angle Glaucoma,   (2016).

[15] P. Gharahkhani, K. Burdon, R. Fogarty, S. Sharma, A. Hewitt, S. Martin, M. Law, K. Cremin, J. Cooke Bailey, S. Loomis, L. Pasquale, J. Haines, M. Hauser, A. Viswanathan, P. McGuffin, F. Topouzis, P. Foster, S. Graham, R. Casson, J. Craig, Common variants near ABCA1, AFAP1 and GMDS confer risk of primary open-angle glaucoma, Nature genetics 46 (2014).

[16] J. Cooke Bailey, S. Loomis, J. Kang, R.R. Allingham, P. Gharahkhani, C.C. Khor, K. Burdon, H. Aschard, D. Chasman, R. Igo, P. Hysi, C. Glastonbury, A. Ashley-Koch, M. Brilliant, A. Brown, D. Budenz, A. Buil, P. Chen, H. Choi, J.

Wiggs, Genome-wide association analysis identifies TXNRD2, ATXN2 and FOXC1 as susceptibility loci for primary open-angle glaucoma, Nature genetics 48 (2016).

[17] P. Hysi, C.-y. Cheng, H. Springelkamp, S. Macgregor, J. Cooke Bailey, R. Wojciechowski, V. Vitart, A. Nag, A. Hewitt, R. Hoehn, C. Venturini, A. Mirshahi, W. Ramdas, G. Thorleifsson, E. Vithana, C.C. Khor, A. Stefansson, J. Liao, J. Haines, T. Aung, Genome-wide analysis of multiethnic cohorts identifies new loci influencing intraocular pressure and susceptibility to glaucoma, Nature genetics 46 (2014).

[18] Y. Liu, V. Vitart, K. Burdon, C.C. Khor, Y. Bykhovskaya, A. Mirshahi, H. Aw, D. Koehn, H. Pg, W. Ramdas, T. Zeller, V. En, C. Bk, W.T. Tay, E.S. Tai, C.-y. Cheng, J. Liu, J.N. Foo, S.-M. Saw, T.Y. Wong, Genome-wide association analyses identify multiple loci associated with central corneal thickness and keratoconus, Nature Genetics (2013).

[19] Y. Chen, Y. Lin, E. Vithana, L. Jia, X. Zuo, T.Y. Wong, L.J. Chen, X. Zhu, P. Tam, B. Gong, S. Qian, Z. Li, X. Liu, M. Baskaran, Q. Luo, C. Guzman, C. Leung, X. Li, W. Cao, Z. Yang, Common variants near ABCA1 and in PMM2 are associated with primary open-angle glaucoma, Nature genetics 46 (2014).

[20] Z. Li, R.R. Allingham, M. Nakano, L. Jia, Y. Chen, Y. Ikeda, M. Baskaran, L.-J. Chen, C. Kee, D. Garway-Heath, S. Sripriya, N. Fuse, K. Abu-Amero, C. Huang, P. Namburi, K. Burdon, S. Perera, P. Gharahkhani, Y. Lin, E. Vithana, A common variant near TGFBR3 is associated with primary open angle glaucoma, Human Molecular Genetics 24 (2015) 3880-3892.

[21] M. abdul kader, P. Namburi, S. Ramugade, R. Ramakrishnan, S. Krishnadas, B. Roos, S. Periasamy, A. Robin, J. Fingert, Clinical and genetic characterization of a

large primary open angle glaucoma pedigree, Ophthalmic Genetics 38 (2016) 1-4.

[22] S.A. Miller, D. Dykes, H. Polesky, Miller SA, Dykes DD, Polesky HF. A simple salting-out procedure for extracting DNA from nucleated cells. Nucleic Acids Res 16: 1215, Nucleic acids research 16 (1988) 1215.

[23] J. Wang, X. Liu, Y. Zhong, Rho/Rho-associated kinase pathway in glaucoma (Review), International journal of oncology 43 (2013).

[24] G. Stelzer, I. Plaschkes, D. Oz-Levi, A. Alkelai, T. Olender, S. Zimmerman, M. Twik, F. Belinky, S. Fishilevich, R. Nudel, Y. Guan-Golan, D. Warshawsky, D. Dahary, A. Kohn, Y. Mazor, S. Kaplan, T. Stein, H. Baris, N. Rappaport, D. Lancet, VarElect: The phenotype-based variation prioritizer of the GeneCards Suite, BMC Genomics 17 (2016).

[25] L. Fernández-Martínez, S. Letteboer, C. Mardin, N. Weisschuh, E. Gramer, B. Weber, B. Rautenstrauss, P. Ferreira, F. Kruse, A. Reis, R. Roepman, F. Pasutto, Evidence for RPGRIP1 gene as risk factor for primary open angle glaucoma, European journal of human genetics : EJHG 19 (2011) 445-51.

[26] R. Roepman, N. Bernoud-Hubac, D. Schick, A. Maugeri, W. Berger, H.-H. Ropers, F. Cremers, P. Ferreira, The retinitis pigmentosa GTPase regulator (RPGR) interacts with novel transport-like proteins in the outer segments of rod photoreceptors, Human Molecular Genetics 9 (2000).

[27] C.-A. Martin, I. Ahmad, A. Klingseisen, M. Hussain, L. Bicknell, A. Leitch, G. Nürnberg, M. Toliat, J. Murray, D. Hunt, Z. Ali, S. Tinschert, J. Ding, C. Keith, M. Harley, P. Heyn, R. Müller, I. Hoffmann, A. Jackson, Mutations in PLK4, encoding a

master regulator of centriole biogenesis, cause microcephaly, growth failure and retinopathy, Nature Genetics 46 (2014).

[28] C. Ricard, J. Pena, M.R. Hernandez, Differential expression of neural cell adhesion molecule isoforms in normal and glaucomatous human optic nerve heads, Brain research. Molecular brain research 74 (2000) 69-82.

[29] B.J. Fan, D. Wang, L. Pasquale, J. Haines, J. Wiggs, Genetic Variants Associated with Optic Nerve Vertical Cup-to-Disc Ratio Are Risk Factors for Primary Open Angle Glaucoma in a US Caucasian Population, Investigative ophthalmology & visual science 52 (2011) 1788-92.

[30] J. Wiggs, R.R. Allingham, A. Hossain, J. Kern, J. Auguste, E. Delbono, B. Broomer, F. Graham, M. Hauser, M. Pericak-Vance, J. Haines, Genome-wide scan for adult onset primary open angle glaucoma, Human molecular genetics 9 (2000) 1109-17.

[31] J. Booij, R. Florijn, J. ten Brink, W. Loves, F. Meire, M. van Schooneveld, P. Jong, A. Bergen, Identification of mutations in the AIPL1, CRB1, GUCY2D, RPE65, and RPGRIP1 genes in patients with juvenile retinitis pigmentosa, Journal of medical genetics 42 (2005) e67.

[32] T. Dryja, S. Adams, J. Grimsby, T. McGee, D.H. Hong, T. Li, S. Andréasson, E. Berson, Null RPGRIP1 alleles in patients with Leber congenital amaurosis, American journal of human genetics 68 (2001) 1295-8.

[33] P. Castagnet, T. Mavlyutov, Y. Cai, F. Zhong, P. Ferreira, RPGRIP1s with distinct neuronal localization and biochemical properties associate selectively with

RanBP2 in amacrine neurons, Human molecular genetics 12 (2003) 1847-63.

[34] P. Ferreira, Identification of Novel Murine- and Human-Specific RPGRIP1 Splice Variants with Distinct Expression Profiles and Subcellular Localization, Investigative ophthalmology & visual science 46 (2005) 1882-90.

[35] T. Mavlyutov, H. Zhao, P. Ferreira, Species-specific subcellular localization of RPGR and RPGRIP isoforms: Implications for the phenotypic variability of congenital retinopathies among species, Human molecular genetics 11 (2002) 1899-907.

[36] S. Patnaik, R. Kotapati Raghupathy, X. Zhang, D. Mansfield, X. Shu, The Role of RPGR and Its Interacting Proteins in Ciliopathies, Journal of Ophthalmology 2015 (2015) 1-10.

[37] N. Luo, M. Conwell, X. Chen, C. Insinna, C. Westlake, L. Cantor, C. Wells, R. Weinreb, T. Corson, D. Spandau, K. Joos, C. Iomini, A. Obukhov, Y. Sun, Primary cilia signaling mediates intraocular pressure sensation, Proceedings of the National Academy of Sciences of the United States of America 111 (2014).

[38] M. Gakovic, X. Shu, I. Kasioulis, S. Carpanini, I. Moraga, A. Wright, The role of RPGR in cilia formation and actin stability, Human molecular genetics 20 (2011) 4840-50.

[39] H. Abiko, S. Fujiwara, K. Ohashi, R. Hiatari, T. Mashiko, N. Sakamoto, M. Sato, K. Mizuno, Rho-guanine nucleotide exchange factors involved in cyclic stretch-induced reorientation of vascular endothelial cells, Journal of cell science 128 (2015).

[40] K. Wang, M. Li, Wang K, Li M, Hakonarson HANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. Nuc Acids Res 38(16): e164, Nucleic acids research 38 (2010) e164.

Chromosome Position	Accession number	Nucleotide changes	Gene Name	Amino acid change	dbSNP	Varlect	Number of cases (sample ID)
14:21816432	NM_020366.3	c.G3719A	RPGRIP1	p.G1240E	rs34725281	8.35†	6 (III-3;III-2;II-2;III-32;IV-26;IV-27)
14:21550588	NM_001278529.2	c.A1295G	ARHGEF40*	p.Q432R	.	1.59‡	6 (III-3;III-2;II-2;III-32;IV-26;IV-27)
14:20666340	NM_001005503.1	c.847delC	OR11G2	p.H282fs	rs528205284	0.99‡	6 (III-3;III-2;II-2;III-32;IV-26;IV-27)
14:20482998	NM_001004712.1	c.A355G	OR4K14	p.M119V	rs7157076	0.95‡	6 (III-3;III-2;II-2;III-32;IV-26;IV-27)
14:21502110	NM_001012264.4	c.C338T	RNASE13	p.S113F	rs114504351	0.71‡	6 (III-3;III-2;II-2;III-32;IV-26;IV-27)
14:19378312	NM_001013354.1	c.T719G	OR11H12	p.V240G	rs61969158	0.22‡	6 (III-3;III-2;II-2;III-32;IV-26;IV-27)

Table.1. List of the Pathogenic variants co- segregated with phenotype. Varlect score with symbol † represents the direct association with glaucoma phenotypes and ‡ represent the indirect association. * represent the Novel variant.

Chromosome Position	Accession number	Nucleotide changes	Gene Name	Amino acid change	dbSNP	Varlect	Number of cases (sample ID)	Number of controls (sample ID)
X:38144822	NM_001034853.2	c.G3430A	RPGR	p.V1144I	rs12688514	9.02†	4 (III-3;III-2;II-2;II-15)	1 (III-41)
4:128816154	NM_001190799.2	c.C2513T	PLK4	p.T838I	rs557954721	6.01†	6 (III-3;III-2;III-32;IV-26;IV-27;II-15)	1 (III-4)
14:45605338	NM_001308133.2	c.C104T	FANCM*	p.P35L	.	4.84‡	6 (III-3;III-2;III-32;IV-26;IV-27;II-15)	1 (III-4)
6:56394545	NM_015548.5	c.A9427G	DST	p.N3143D	rs530170321	4.2‡	6 (III-3;III-2;II-5;II-15;III-32;IV-27)	2 (III-4;III-16)
13:101759922	NM_001350750.2	c.A2408G	NALCN	p.Y803C	rs549182297	4.11†	8 (III-3;III-2;II-2;II-5;III-19;III-32;IV-27;II-15)	3 (III-41;III-4IV-10)
1:145534254	NM_001303041.1	c.A1330G	ITGA10	p.T444A	rs782732004	3.98‡	3 (III-3;III-2;II-2)	1 (III-4)
1:145541806	NM_001303041.1	c.T2900C	ITGA10*	p.L967P	.	3.98‡	3 (III-3;III-2;II-2)	1 (III-4)
2:179528378	NM_001267550.2	c.G36508A	TTN	p.E12170K	rs2163008	3.78‡	4 (III-3;III-2;II-2;III-32)	1 (IV-28)
19:55350963	NM_001281971.2	c.509_510insCCCGGAGCTCCTATGACATGTA	KIR2DS4	p.S151fs	rs551456772	3.56†	5 (III-3;II-5;III-32;IV-26;IV-27)	2 (IV-11;III-4)
4:55147769	NM_001347827.2	c.C2345T	PDGFRA	p.T782M	rs2291591	3.56†	4 (III-32;IV-26;IV-27;II-15)	1 (IV-28)

4:141483476	NM_021833.5	c.C680T	UCP1	p.T227I	rs148598275	*3.51‡*	6 (III-3;III-2;III-32;IV-26;IV-27;II-15)	1 (III-4)
1:175046835	NM_022093.2	c.C281T	TNN	p.T94M	rs41266080	*3.23‡*	4 (III-3;III-2;II-5;II-15)	2 (III-41;III-34)
3:141526640	NM_139209.2	c.G1204A	GRK7	p.D402N	rs150840377	*3.21‡*	8 (III-3;III-2;II-5;III-32;IV-26;IV-27;III-19;II-15)	4 (III-34;IV-11;IV-28;III-41)
5:33951693	NM_001012509.4	c.G1122C	SLC45A2	p.L374F	rs16891982	*3.17‡*	5 (III-2;III-32;IV-26;IV-27;II-15)	1 (III-4)
11:113126641	NM_000615.7	c.A1841T	NCAM1*	p.D614V	.	3.16†	6 (III-3;III-2;III-32;IV-26;IV-27;II-15)	2 (IV-28;III-4)
19:45853924	NM_177417.3	c.C1298T	KLC3*	p.S433F	.	*3.05‡*	6 (III-3;III-2;II-5;IV-27;III-32;IV-26)	3 (IV-28;III-4;III-16)
19:50752298	NM_024729.3	c.T1360G	MYH14	p.W454G	rs572234218	*2.94‡*	6 (III-3;III-2;II-5;III-32;IV-26;IV-27)	3 (IV-28;III-16;III-4)
1:209791929	NM_001318046.2	c.C2777A	LAMB3	p.A926D	rs2076222	*2.6‡*	7 (III-3;III-2;II-5;III-32;IV-26;IV-27;III-19)	3 (IV-11;III-4;IV-10)
10:100189242	NM_001322492.1	c.C808T	HPS1	p.P270S	rs34533614	*2.55‡*	3 (III-3;III-2;II-2)	1 (III-4)
10:100202987	NM_000195.5	c.T11C	HPS1	p.V4A	rs58548334	2.55†	6 (III-3;III-2;III-32;IV-26;V-27;II-15)	3 (III-41;III-34;IV-28)
10:97192237	NM_001034957.1	c.C173T	SORBS1	p.P58L	rs200179325	*2.49‡*	6 (II-5;III-32;IV-26;IV-27;II-15;III-19)	1 (IV-10)

7:107746432	NM_001350531.2	c.C700T	LAMB4	p.H234Y	rs2074749	2.47‡	8 (III-3;III-2;II-5;III-32;IV-26;IV-27;II-15;III-19)	2 (III-41;III-16)
10:90530612	NM_001289967.1	c.G683C	LIPN	p.G228A	rs201135817	2.33‡	9 (III-3;III-2;II-2;II-5;II-15;III-32;IV-26)	4 (III-16;IV-28;IV-27;III-41;III-19;III-4)
3:130361856	NM_001102608.2	c.G5216A	COL6A6	p.R1739Q	rs16830494	2.17‡	6 (III-3;III-2;II-5;III-32;IV-26;IV-27)	3 (III-4;III-16;;IV-28)
19:44571252	NM_013361.6	c.1506dup	ZNF223	p.R424fs	rs562593501	2.05‡	7 (III-3;III-2;II-5;III-32;IV-26;II-15;III-19)	3 (IV-11;III-4;III-16)
5:90016871	NM_032119.4	c.G9743A	ADGRV1	p.G3248D	rs16869032	2.05†	8 (III-3;III-2;II-5;III-32;IV-26;II-15;III-19)	4 (III-16;IV-28;IV-27;III-41;III-34)
10:73434888	NM_001171930.2	c.G1469C	CDH23	p.G490A	rs1227049	2.05†	6 (III-3;III-2;II-5;III-32;IV-27;III-19)	2 (III-34;III-16;)
10:71160787	NM_001322367.1	c.C2554G	HK1*	p.P852A	.	2.05†	6 (III-3;III-32;IV-26;II-15;III-2;II-5)	2 (III-16;III-4)
5:96117554	NM_001040458.3	c.C2290T	ERAP1*	p.P764S	.	1.95‡	7 (III-3;III-2;II-2;II-5;IV-26;IV-27;III-19)	0 ()
3:182788862	NM_001293273.1	c.A335G	MCCC1	p.E112G	rs142629318	1.83‡	6 (III-3;III-32;IV-26;IV-27;II-15;II-5)	2 (III-16;III-4)
3:179408072	NM_003940.3	c.A338G	USP13	p.N113S	rs771971543	1.78‡	7 (III-3;III-2;II-5;III-32;IV-26;IV-27;II-15)	3 (III-4;III-16;IV-28)

4:141483476	NM_021833.5	c.C680T	TMEM63B	p.R82P	rs371238478	1.69‡	6 (III-3;II-5;III-32;IV-26;IV-27;II-15)	3 (III-4;IV-10;III-16)
19:40363916	NM_003890.2	c.G14726A	FCGBP	p.T1975M	rs372872173	1.66‡	6 (III-3;III-2;II-5;IV-27;III-32;IV-26)	3 (IV-28;III-4;III-16)
19:40366240	NM_003890.2	c.C13994T	FCGBP	p.R4909H	rs77005739	1.66‡	3 (III-3;II-2;IV-27)	1 (IV-28)
3:141526640	NM_139209.2	c.G1204A	FCGBP	p.P4665L	rs62106922	1.66‡	3 (III-3;II-2;IV-27)	1 (IV-28)
5:96117554	NM_001040458.3	c.C2290T	EIF2AK3	p.S136C	rs867529	1.62†	7 (III-3;III-2;II-5;III-32;IV-27;II-15;III-19;)	2 (III-4;III-16)
2:179528378	NM_001267550.2	c.G36508A	TIGD4	p.C204F	rs576908904	1.5‡	6 (III-3;III-2;II-5;III-32;IV-26;IV-27;IV-10)	3 (III-16;IV-28)
15:82934639	NM_001322400	c.G941A	GOLGA6L10	p.R314H	rs200928526	1.564‡	7 (III-3;III-2;II-5;III-32;IV-26;IV-27;II-15)	3 (III-4;III-16;IV-28)
X:38144822	NM_001034853.2	c.G3430A	PEX5L	p.F173C	rs141827659	1.54‡	7 (III-3;III-2;II-5;III-32;IV-26;IV-27;II-15)	3 (III-4;III-16;IV-28)
17:39622068	NM_001017402.1	c.C665A	KRT32	p.S222Y	rs2071561	1.1†	6 (III-3;III-2;III-32;IV-26;IV-27;II-15)	1 (III-4)
11:103006630	NM_001080463.2	c.G2527A	DYNC2H1	p.A843T	rs548461924	0.97‡	6 (III-3;III-2;III-32;IV-26;IV-27;II-15)	2 (III-4;IV-28)
4:128816154	NM_001190799.2	c.C2513T	PDZD3	p.Q226K	rs147147532	0.85‡	6 (III-3;III-2;III-32;IV-26;IV-27;II-15)	0
3:179529649	NM_001349404.2	c.T518G	OR51E1	p.A156T	rs202113356	0.84‡	6 (III-3;III-2;III-32;II-15;IV-26;IV-27)	3 (III-4;III-16;IV-28)

11:119058667	NM_001168468.2	c.C676A	OR11H2	p.P269S	rs2815979	0.82‡	6 (III-3;III-2;III-32;IV-26;IV-27;II-15)	0 (III-4;IV-28)
21:33735605	NM_014825.3	c.T1369A	URB1	p.S457T	rs148292685	0.81‡	7 (III-3;III-2;II-5;III-32;IV-26;IV-27;II-15)	3 (III-4;III-16;IV-28)
11:30974115	NM_001350255.1	c.G1449T	DCDC1	p.R483S	rs158633	0.78‡	6 (III-3;III-32;IV-26;IV-27;II-15;III-19)	2 (III-41;III-34)
11:31123752	NM_001350255.1	c.A812G	DCDC1	p.D271G	rs183555899	0.78‡	6 (III-3;III-32;IV-26;IV-27;II-15;III-19)	2 (III-41;III-34)
19:49969382	NM_001145396.2	c.G1627C	ALDH16A1	p.A543P	rs555667637	0.42‡	6 (III-3|III-2|II-5|IV-27|III-32|IV-26|)	3 (III-4_C|III-16_C|IV-28_C|)
2:88913273	NM_004836.7	c.C407G	EFHC1	p.R353W	rs527295360	0.42‡	6 (III-3;III-32;IV-26;IV-27;II-15;II-5)	2 (III-4;III-16)
1:209791929	NM_001318046.2	c.C2777A	KLHL24	p.D197A	rs116961268	0.42‡	7 (III-3;III-2;II-5;III-32;IV-26;IV-27;II-15)	3 (III-4;IV-28;III-16)
19:50752298	NM_024729.3	c.T1360G	MAGIX	p.P221L	rs781930221	0.28‡	7 (III-2;II-2;II-5;III-32;IV-26;IV-27;III-19)	3 (III-4;III-34;III-16)
5:33951693	NM_001012509.4	c.G1122C	PIEZO1	p.I2265V	rs1803382	0.24‡	6 (III-3;III-2;III-32;IV-26;IV-27;II-15)	1 (III-4)
10:90530612	NM_001289967.1	c.G683C	LARP1B*	p.I178F	.	0.21‡	7 (III-3;III-2;II-5;III-32;IV-26;IV-27;II-15)	3 (III-4;III-16;IV-28)
6:44103070	NM_001318792.1	c.G245C	SSTR1	p.T390delinsTLX	rs775405351	0.18†	6 (III-3;III-2;II-5;IV-27;II-15;III-32)	2 (IV-28;III-16)

Table 2. List of pathogenic variants. Varlect score with symbol † represents the direct association with glaucoma phenotypes, and ‡ represent the indirect association. * represent the Novel variant.