Probing Causality of the Brainstem-Hypothalamic Murine Models of Sleep-Wake Regulation

Authors: Fatemeh Bahari[1,2]*, Myles W. Billard[1,2], John Kimbugwe[1,2], Carlos Curay[1], Glenn D.R. Watson[1,3], Kevin D. Alloway[1,3], Bruce J. Gluckman[1,2,4,5]*

Affiliations:

[1]Center for Neural Engineering, Pennsylvania State University, University Park, PA 16802, USA

[2]Department of Engineering Science and Mechanics, Pennsylvania State University, University Park, PA 16802, USA

[3]Neural and Behavioral Sciences, College of Medicine, Pennsylvania State University, Hershey, PA 17033, USA

[4]Department of Neurosurgery, College of Medicine, Pennsylvania State University, Hershey, PA 17033, USA

[5]Department of Bioengineering, Pennsylvania State University, University Park, PA 16802, USA

*Corresponding authors email addresses: Fatemeh Bahari, fa.bahari92@gmail.com, Bruce J. Gluckman BruceGluckman@psu.edu

Abstract

Sleep-wake regulation is thought to be governed by interactions among several nuclei in midbrain, pons, and hypothalamic regions. Determination of the causal role of these nuclei in state transitions requires simultaneous measurements from the nuclei with sufficient spatial and temporal resolution. We obtained long-term experimental single- and multi-unit measurements simultaneously from multiple nuclei of the putative hypothalamic and brainstem sleep-wake regulatory network in freely behaving rats. Cortical and hippocampal activity, along with head acceleration were also acquired to assess behavioral state. Here, we confirm that the general activity profile of the recorded sleep-wake regulatory nuclei is similar to the patterns presented previously in brief recordings of individual nuclei in head-fixed animals. However, we found that

these activity profiles when studied with respect to cortical and behavioral signs of state transitions differ significantly from previous reports. Our findings pose fundamental questions about the neural mechanisms that maintain specific states and the neural interactions that lead to the emergence of sleep-wake states.

Keywords: Sleep-wake regulatory network, flip-flop models of sleep, single-unit recordings, brainstem, hypothalamic area, freely behaving rodents

1. Introduction

Sleep-wake states are thought to be driven by basal forebrain, brainstem, and hypothalamic circuits (ECONOMO, 1930; Moruzzi and Magoun, 1949; Saper et al., 2005, 2010; Datta and MacLean, 2007; Weber et al., 2015; Xu et al., 2015; Sherin et al., 2018). These states are predominantly characterized from electroencephalogram (EEG) and electromyogram (EMG) recordings, with transitions between states described as discrete mechanisms (Saper et al., 2001, 2010; Datta and Siwek, 2002; Luppi et al., 2006; Weber and Dan, 2016). Models of underlying network mechanisms of sleep-wake regulation have attempted to describe and reproduce these discrete transitions (Tamakawa et al., 2006; Fulcher et al., 2008; Phillips and Robinson, 2008; Diniz Behn and Booth, 2010; Rempe et al., 2010; Fleshner et al., 2011). To do so, these models often invoke co-inhibitory dynamics of sleep-wake regulatory cell groups in configurations analogous to electrical flip-flops (Saper et al., 2001; Lu et al., 2006; Weber and Dan, 2016). However, observations of cell group activity and involvement are derived from experiments conducted on head-fixed (Boissard et al., 2002; Verret et al., 2003; Takahashi et al., 2010; Sakai, 2011; Boucetta et al., 2014) or lightly anesthetized animals (Aston-Jones et al., 2001; Sherin et al., 2018), via short recordings, and often within novel environments (Datta and Siwek, 2002; Lu, 2006; Alam et al., 2014; Weber et al., 2015; Xu et al., 2015; Bjorness et al., 2016). These conditions inherently modulate sleep-wake patterns (Sigl-Glöckner and Seibt, 2019). Further, short recording sessions provide too few state transitions to accurately

determine the relationship between firing rate and state transitions. These recording sessions are often 2-6 hours long amounting to approximately 6-20 sleep cycles (Takahashi et al., 2009, 2010; Sakai, 2018). Following the Poisson distribution laws and a typical bin size of 10 ms for firing rate calculations, a few hundred state transitions are required to detect a statistically significant change in firing rates relevant to the within-state activities of these neurons.

To investigate the neural basis of sleep-wake transitions in normal conditions, we chronically recorded single- and multi-unit activity simultaneously from multiple hypothalamic and brainstem sleep-wake regulatory nuclei in freely behaving rats. Cortical and hippocampal activity, along with head acceleration were also acquired to assess behavioral state. The dataset comprises 100-300 sleep-wake cycles for each neuron, which provides a sufficient number of state transitions to detect a statistically significant change in firing rate. Our findings address current hypotheses regarding the causal role of the represented brainstem and hypothalamic cell groups in initiating naturally emerging transitions between states of vigilance.

2. Methods

All animal work was approved by and performed under the oversight of the Institutional Animal Care and Use Committee at the Pennsylvania State University.

To study the sleep-wake regulatory network as well as to identify sleep-wake transitions we used the systemDrive (Billard et al., 2018): a customizable micro-wire multitrode microdrive for targeting multiple cell groups along non-parallel non-co-localized trajectories. Animals were implanted with the systemDrive's micro-wire multitrodes targeting several sleep-wake regulatory cell-groups. The systemDrive houses additional screw electrodes for electrocorticography (ECoG), and fixed depth electrodes for hippocampal local field potential (LFP) recordings. Following implantation, animals were returned to their home-cages and long-term continuous recordings started after ample post-operation recovery time (7 days).

80   In accordance with common microdrive system use, we advanced electrodes along the dorsal-

81   ventral axis of each targeted cell group and recorded neuronal activity of the area for continuous

82   periods of 3 to 10 days. The procedure was repeated until the entire depth of the target was

83   covered. Single- and/or multi-unit activity, when present, was extracted for all channels in each

84   target.

85   2.1.   Animal Surgery and Care:

Surgical and implantation techniques are previously described (Billard et al., 2018). Briefly, male

and female Long Evans rats weighing between 275-350 grams were implanted for continuous

recordings that lasted for a duration of 2-6 weeks.

Recording electrodes include micro-wire bundles for monitoring brainstem neuronal activity, as

well as hippocampal LFP pairs and ECoG screws for determining the EEG and behavioral state.

For observation of sleep-wake regulatory network (SWRN) dynamics, we recorded from five

brainstem structures with target coordinates referenced to intraural line (IA) according to the

Paxinos-Watson rat brain atlas:

Dorsal Raphe (DR) [AP: +1.5 mm, ML: 0 mm, DV: -3.6 mm, 21.33^{o} or -21.33^{o}]

PeduncloPontine Tegmentum (PPT) [AP: +0.5 mm, ML: ±2 mm, DV: -3.6 mm]

LateroDorsal Tegmentum (LDT) [AP: +0.36 mm, ML: ±0.8 mm, DV: -3 mm, 13.65^{o} or -13.65^{o}]

VentroLateral Preoptic Area (VLPO) [AP: +8.76 mm, ML: ±1 mm, DV: -0.75 mm, 15.52^{o}]

Locus Coerleus (LC) [AP: -0.72 mm, ML: +1.3 mm, DV: -3.2 mm]

A number of these nuclei are located near sensitive structures such as the sagittal sinus and

lateral ventricles. Therefore, trajectories were chosen to avoid these sensitive structures and

allow enough room on the cranium for all electrodes.

For hippocampal LFPs, two pairs of custom-made 50 µm iridium oxide deposited micro-reaction chamber (µRC) electrodes formed from gold coated stainless-steel wire (Shanmugasundaram and Gluckman, 2017) with ends staggered by 250-300 µm were implanted bilaterally in the dorsal hippocampus at coordinates [AP: +5 mm, ML: ±3 mm, DV: -2.5 mm from the cortex]. For ECoG, four stainless steel screws (one-eighth inch 18-8 stainless steel 0-80 cortical screws, McMaster-Carr) were implanted at coordinates [AP: +7.5 mm, ML: ±4 mm], [AP: +3 mm, ML: -4 mm], and [AP: +2.5 mm, ML: +3 mm]. Two additional stainless screws were implanted at [AP: +11 mm, ML: ±3 mm] for ground and reference.

After completing surgery, animals were returned to their individual home-cages for recovery with free access to food and water. Following recovery, the animals were cabled for continuous video and EEG monitoring. Every 3 to 10 days the animals were briefly anesthetized to advance the micro-wire electrodes incrementally between recording sessions and until they covered the entire dorsal-ventral length of the structures of interest.

2.2. Histological Techniques:

At the completion of recordings animals were sacrificed and their brains were histologically processed to examine the electrode tracks with respect to the targeted brain structures. Details and results of the histological methods and analysis are previously described (Billard et al., 2018).

2.3. Data Acquisition:

All rats were housed individually in custom-made plexiglass cages with dimensions 9" (w) x 15" (d) x 24" (h). At the first electrode driving session, each animal received a differential or referential digitizing head-stage amplifier with 3-axis accelerometer (RHD2116 or RHD2132, Intan Technologies, Los Angeles, CA) which can directly connect to an electrode interface board on the systemDrive. The amplifier was connected to a data acquisition board (RHD2000 Evaluation System, Intan Technologies) via 3' lightweight serial-peripheral interface

(SPI) cable (Intan Technologies) and in-house adapted low-torque commutator (SRA-73540-12,

Moog Inc.).

Hippocampal LFP, ECoG, as well as single and multi-unit signals were simultaneously acquired

at 30,000 samples per second (SPS) and high-pass filtered at 1 Hz. Head acceleration

measurements were sampled at a rate of 7500 Hz. A 12 hour light/dark cycle was maintained

with the lights on between 6 am and 6 pm. Video monitoring, at 3 frames per second, began on

the day of implantation and continued within each recording session. Video files were stored in

one-hour-long files, while biopotentials and head acceleration data were stored in 5 to 15-

minute-long files for further analysis.

To measure single- and multi-unit activity we used the microdrive arrangements within the

systemDrive. For each animal, electrodes aimed at sleep-wake regulatory cell-groups in the

brainstem and/or hypothalamus were first advanced after the recovery period.

During this and subsequent driving sessions, the animal was maintained under anesthesia with

0.5%-2% isoflurane gas and the head-mount was opened to access the systemDrive. We

maintained the recordings throughout the driving session. Each electrode bundle was advanced

incrementally (about 25μm) and the signal was monitored until we observed new single-unit

activity. For electrode trajectories that passed through the superior and inferior colliculi or other

sensory-responsive regions, we tested auditory and visual stimuli to evoke responses that

would confirm that the electrode was advancing through the appropriate brain structures.

2.4. Data Analysis:

147   All analyses were performed using custom-written MATLAB (Mathworks) and Labview (National

148   Instruments) scripts. Data used for analysis in this manuscript are available in an online

149   repository.

2.4.1. Sleep and Behavioral Scoring:

Hippocampal LFP, ECoG, and accelerometer time-series were down-sampled to 1000 samples per second and reformatted into 1 hour-long blocks of binary data within a custom Labview script. The raw data were then band-pass filtered at 1-125 Hz for LFP, 1-55 Hz for ECoG, and 2-100 Hz for head acceleration.

The SOV was marked using an adaptation of the semi-automatic Linear Discriminant Analysis (LDA) previously described (Sunderam et al., 2007; Sedigh-Sarvestani et al., 2014). Briefly, the SOV was classified into five states: Non-rapid eye movement (NREM) sleep, rapid eye movement (REM) sleep, an intermediate state between NREM and REM (IS), awake (wake$_{wb}$), and active exploration with non-coherent theta rhythm (wake$_\theta$). The frequency band specifications for NREM, REM, and awake states are as described in (Sunderam et al., 2007; Sedigh-Sarvestani et al., 2014). The IS was characterized by appearance of weak bursts of theta-band activity in the presence of identifiable delta band activity (GOTTESMANN, 1996; Boucetta et al., 2014). IS termination is defined by a large reduction of power in the lower delta band and a sharp increase in the theta band power (Fig. S1A). During the IS, the head acceleration power is low with no evidence of the short spikes characteristic of muscle twitches during REM.

For each animal, 4-6 hours of video-EEG data within 1 day were randomly selected and manually scored for SOV and SOV transition time. These data were then used to train the LDA. The remaining 18-20 hours of data were set aside as out-of-sample data (test set). For both training and test sets, features were computed, within 2 second long overlapping windows, from EEG spectral power in frequency bands 0.5-4, 4-8, 8-12, 12-25, and 25-80 Hz, plus coherence measurements (to distinguish REM bouts from IS and wake$_\theta$) and head acceleration. Head acceleration measurements were used for more accurate SOV scoring (Sunderam et al., 2007). The features were generated using causal filters and updated every second at the end of

175　computation window. Therefore, all the scores are based on information up to the time leading

176　to the state transition and *not* after that.

2.4.2. Spike Scoring:

We used Intan Technologies differential (RHD2216) and referential (RHD2132) preamplifiers to

acquire electrophysiological signals. The default pass-band for the filters in these preamplifiers

is between 0.1 Hz and 7.5 kHz. A 16-bit analog to digital converter then samples the signal from

the AC-coupled amplifiers. At 7.5 kHz, the cut-off frequency in the analog low-pass filter is much

higher than what is used conventionally (e.g. 3.5 kHz in (Datta and Siwek, 2002; Sakai, 2011)).

Data segments were filtered with a causal band-pass filter with 250 Hz to 7.5 kHz pass-band,

and then thresholded to detect single-unit activity. Units were detected as instances of threshold

crossings where thresholds were specified as multiples of the standard deviation of the filtered

background signals. The background signals represent separate epochs of time that are free of

distinguishable spiking activity and are used to calculate the mean and variance of the

background distribution. We detected units with a threshold of 5-7 standard deviations. When

spike events were detected, the waveform of each action potential was extracted from the data.

2.4.3. Statistical Analysis:

The core analysis of this work is to characterize the state-dependent firing rates and the peri-

state-transition time firing rates. These measurements are calculated from the firing times of

individual neurons and averaged over all like states or state-transition types. With just a mean-

firing rate description, these measurement statistics should follow Poisson statistics, such that

given a firing rate FR, the probability $P_\lambda(k)$ of observing the number of k spikes measured within

a time bin τ with respect to the transition time summed over N observed transitions, as:

$$P_\lambda(k) = \frac{\lambda^k e^{-\lambda}}{k!}, \qquad \lambda = FR \times N \times \tau$$

Critical in our analysis is to determine for each neuron when the instantaneous firing rate

$FI = k/(N \times \tau)$ is significantly different from the pre-transition state mean firing rate, (see for

example Fig. S3). We define the time t_{sig} when the observed k first falls within the upper or

lower 1% of $P_\lambda(k)$.

During the course of our experiments, the activity of each neuron at each target location was

measured over hundreds of sleep-wake cycles $N \in (100,300)$. Coupled with within-state and

instantaneous firing rates typically much less than 20 per second, the large ensemble state

transitions enables us to have sub-second resolution (τ) in our statistical analysis, which is

crucial for causality determination.

3. Results

We successfully acquired simultaneous measurements from up to four separate sleep-wake

regulatory nuclei (SWRN) in each rat. The success rates of chronic systemDrive targeting and

recording from multiple (n = 12) rats are shown in Table S1. Cumulative recording duration,

number of recording sessions, and number of single units exceeding a threshold of 7 standard

deviations are presented for each animal. Data presented in this manuscript pertain to the SOV-

dependent subset of neurons presented in Table S1.

Recording single units in deep brain structures across multiple days in freely moving rats is

extremely challenging. The dataset described here was previously summarized in (Billard et al.,

2018). In this previous work we confirmed the location of recorded targets using histological

analysis (as shown in Fig. 6 in (Billard et al., 2018)). Further, to confirm long-term stability of our

unit recordings we presented 1) measurements of signal-to-noise ratio for recordings from

SWRN (Fig. 7 in (Billard et al., 2018)), 2) unit waveforms simultaneously extracted from up to 3

SWRN that were continuously recorded over 4-7 days from a subset of animals (Fig. 8 in

(Billard et al., 2018)), and 3) examples of simultaneously recorded units from PPT, LDT, and DR

over 6 continuous days (Fig. 9 in (Billard et al., 2018)). We therefore confirmed that unit

detection is not disrupted during normal animal behavior and that recorded units remained

stable throughout the duration of the recording periods with no significant deterioration of

amplitude and signal-to-noise ratio.

We recorded single-unit activity with time-matched hippocampal, cortical, and head motion data

scored for sleep-wake states. The dataset corresponds to 569 total days of recording spread

over 12 animals. The SOV transition probabilities over all states are shown in Fig. S1D. The

time-of-day dependence of the different SOVs are illustrated in Fig. S2A. This is likely the

longest dataset acquired from freely behaving animals in their home cage environment for which

the analysis includes identification of the intermediate state (IS). Although previous reports

identify IS as the transitory state immediately prior to REM (GOTTESMANN, 1996; Gervasoni,

2004; Boucetta et al., 2014), we found that about 20% of the transitions out of IS continue into

Wake (Fig. S1C&D). Additionally, the IS bouts in our recordings are typically 35-60 s long (Fig.

S2B) which is much longer than previously reported.

3.1.   Mean state-dependent neuronal firing rate and activity profiles of the
brainstem-hypothalamic SWRN

In the brainstem-hypothalamic model of sleep-wake regulation interactions between cholinergic

PPT/LDT, monoaminergic LC/DR, and GABAergic VLPO lead to different SOVs (Saper et al.,

2010). Cumulatively, we recorded 855 single units across multiple regions and many continuous

days (3-10 days) from 12 freely behaving animals all in their home-cage environment. A range

of 100-300 complete sleep cycles were recorded from each neuron. About half (n = 428) of the

identified single-units recorded were SOV-dependent Of the n=428 SOV-dependent neurons,

152 were REM-active, 15 were NREM active, 142 were wake-active, 60 were wake and REM

active, and 41 were active in all SOVs except REM (denoted as REM-off). These neurons were

distributed throughout multiple brain regions: We found Wake-active neurons in PPT, LDT, and

DR; NREM-active neurons in VLPO; REM-active neurons in PPT and LDT. We also identified a

group of REM-off neurons located in vlPAG. The anatomical distribution of the state-dependent

and state-indifferent neurons is detailed in Table 1.

Cell Type	Number of recorded neurons (%)	Recording Sites/Target				
		PPT	LDT	DR	VLPO	LC
Wake-active	83 (8.2)	21	0	62	0	0
Burst	20 (2.3)	0	0	20	0	0
Wake$_\theta$-active	39 (4.6)	19	20	0	0	0
NREM-active	12 (1.4)	0	0	0	12	0
Burst	3 (0.3)	0	0	0	3	0
IS-active	18 (2.1)	0	0	0	18-MPA	0
REM-active	152 (17.8)	89	63	0	0	0
Wake-REM	60 (7)	30	30	0	0	0
REM-off	41 (4.8)	0	0	41-vlPAG	0	0
State Indifferent	427 (49.9)	295	198	270	88	4

Table 1. Anatomical distribution of the state-dependent and state-indifferent neurons in

brainstem and hypothalamus. Listed are the number of neurons recorded in the indicated

areas. In addition to the targeted brain regions, PPT, LDT, DR, VLPO, and LC, we measured state-

dependent activity from adjacent structures. These include vlPAG from a subset of animals with

electrodes targeting DR and neurons within the posterior region of LH. Abbreviations: PPT,

pedunculopontine tegmentum, LDT, laterodosral tegmentum, DR, dorsal raphe, vlPAG,

ventrolateral periacquaductal gray, VLPO, ventrolaterolateral preoptic area, MPA, medial

preoptic area, LC, locus coerleus.

3.1.1. Wake-active neurons

One hundred and forty two of the 428 state-dependent neurons (33%) were marked by having a

higher discharge rate during wake periods than during sleep bouts. Of these, 20 – located in DR

– displayed burst-like behavior with short inter-spike-interval (ISI) and relatively low firing rates

(Fig.1). These neurons were specifically active during wake and not wake$_\theta$ bouts.

Figure 1. Wake-active neurons in DR with bursting behavior. We identified 20 neurons in DR that were mostly active during wake bouts and off during NREM and REM sleep (A) and (D). These neurons, whose average waveform is demonstrated by the solid black line in (B) – standard deviation is shown in light gray shading – had bursting behavior with very short inter-spike intervals (C). The gray and orange bars (D) indicate dark and light periods respectively. We did not find any consistent or significant difference in the neurons' firing rate during light vs dark periods.

The remaining wake-active neurons in DR were divided into three subgroups based on their average waveforms and ISIs: Most of the neurons (n=36) showed a broad asymmetric biphasic spike waveform with large positive peak and long spike duration (Fig. 2A&B). The descending portion of the waveform displayed a notch and was then followed by a smaller amplitude and longer negative component. The second type of cells (n=16) displayed a similar biphasic pattern however their spike waveforms were more symmetric and with a much less pronounced notch (Fig. 2C&D). The third type (n=10) displayed a biphasic spike waveform with a smaller positive

peak which descended into a large amplitude negative trough (Fig. 2E&F). These waveforms

are consistent with previous reports demonstrating wake-active DR neurons with various

average waveforms both in anesthetized and un-anesthetized (head-fixed) animals (Urbain et

al., 2006; Sakai, 2011).

In 8 out of 12 rats we identified wake-active neurons in the PPT areas. These neurons displayed

a triphasic spike waveform with mean spike duration of 1.36 ± 0.14 ms (Fig. 2G&H). We also

identified a group of neurons in PPT (10 out of 12 rats, Fig. 2I&J) and LDT (3 out of 7 rats, Fig.

2K&L) that were only active during wake$_\theta$ bouts. Based on their average waveform, ISI, and

state-dependency they potentially belong to the group of previously reported wake-active

neurons in PPT and LDT areas (Datta and Siwek, 2002; Saper et al., 2010).

Figure 2. Wake and wake$_\theta$ active neurons. We identified 4 types of wake-active neurons in DR

and 2 types of wake$_\theta$ active neurons in PPT and LDT. The average waveform and firing rates of

the wake-active neurons (B, D, F, H) is similar to previous reports (Sakai, 2011). The gray and

orange bars (A, C, E, G, I, K) indicate dark and light periods respectively. Although the neurons

had higher firing rate in the dark vs light period, given the standard deviation, the differences

were not statistically significant.

3.1.2. NREM-active neurons

We identified 15 out of the 428 state-dependent neurons (1.7%), specifically in the VLPO

region, with higher discharge rates during NREM than during REM or wake bouts. The activity

of these neurons was continuously monitored for at least 24 hours in each animal (n = 5). These

neurons had a biphasic spike waveform with a small amplitude but long negative trough (Fig.

3A&B).

Figure 3. Sleep and wake-REM active neurons. We identified 15 neurons in the VLPO that were

predominantly active during NREM (A, B). The average waveform (B) of these neurons is

consistent with previous report (Takahashi et al., 2009; Alam et al., 2014). Similarly the activity

pattern (D) and waveform (C) of the REM-active neurons in PPT (grouped as PPT$_a$ and PPT$_b$) and

LDT (grouped as LDT$_a$ and LDT$_b$) were consistent with findings of Datta et al (Datta and Siwek,

2002). We also identified neurons in the LDT and PPT areas that were active during REM, wake,

and wake$_\theta$ bouts (E). The orange bars (A, C, E) indicate light periods. The other shades of color

(i.e. gray, brown, white) indicate the dark period for different neurons. We did not find any

consistent or significant difference in the neuron's firing rate during light vs dark periods.

3.1.3. REM-active neurons
We identified 152 out of the 428 state-dependent neurons as REM-active. These neurons were

located either in PPT (89 neurons, n=10 animals) or in LDT (63 neurons, n=7 animals) with

triphasic sharp waveforms. Based on their ISI, average waveform, and location of clusters in the

principle component space during spike sorting we subdivided these neurons into 2 groups for

each region (Fig. 3C&D).

3.1.4. Wake-REM-active neurons
A subset of recorded neurons (7%) from PPT and LDT were active both during REM and awake

bouts (Fig. 3E&F). These neurons were co-localized with a portion of REM-active neurons in

PPT and LDT. Therefore, differences in their activity profiles do not stem from their specific

location.

3.1.5. REM-off neurons
In 4 animals, we identified a number of neurons in and around the vlPAG region that were active

during all SOVs except REM. The firing rate of these neurons during wake$_\theta$ was significantly

higher than during REM (Fig. 4). Therefore, we hypothesize that although active during wake$_\theta$

bouts, these neurons do not code for theta rhythm or other cortical and behavioral signs of

REM.

Figure 4. REM-off neurons. In 4 animals we found neurons potentially located in vlPAG that

were active during all sleep states except IS and REM (A). These neurons did not demonstrate

any theta-dependent behavior because although they are active during wake$_\theta$, they were

significantly less active during REM (high theta activity). The gray and orange bars (A) indicate

dark and light periods respectively. We did not find any consistent or significant difference in

the neuron's firing rate during light vs dark periods. The average waveform of these neurons

was triphasic and similar to the cholinergic REM-active neurons in PPT and LDT.

3.1.6. IS-active neurons

None of the identified state-dependent neurons active during NREM, REM, or wake states were

active during the IS. However, in 4 animals we identified a group of neurons potentially within

the medial preoptic area (MPA) dorsal to the VLPO that were active during REM and IS (Fig. 5).

Although their firing rate was slightly lower in IS bouts that ended in REM than the ones that

ended in wake, the difference was not statistically significant.

The IS emerges during transitions out of NREM with clear behavioral and cortical signs (see

methods, Fig. S1) that, when reported (GOTTESMANN, 1996; Gervasoni, 2004; Boucetta et al.,

2014), are associated with human NREM-II. Although in most reports IS appears to occur only

between NREM and REM states, we found that 20% of the time IS was directly followed by

wake bouts (Fig. S1C&D). Additionally, the median duration of IS in our animals was ~45s (Fig.

S2B).

The identification of neurons whose firing rate is significantly less during NREM and only

increases during transitions out of NREM into IS or REM indicates that there is a distinct

circuitry that drives IS.   That the IS-active neurons are active during IS and REM states, but not

wake$_\theta$ bouts indicates that IS-active neurons are not required for coding of theta band activity.

Figure 5. Identified neurons in the preoptic areas are active during the intermediate state. (A)
Stereotaxic coordinates of the regions with IS-active neurons, recorded in 4 animals, are
indicated by the yellow dots. For visual clarity purposes, the coronal histologic section shown
here is from 1 of the 4 animals chronically implanted with this target. The yellow dots indicate
the electrode locations in all 4 animals when the IS-active neurons where monitored. **(B)**The IS-
active neurons are active during REM and IS bouts and increase their firing rate slightly upon
transitions from IS to REM. However they are off during wake and wake$_\theta$ bouts. This indicates
that 1) IS is a distinct state of vigilance with its own corresponding neuronal activity and 2) the
IS-active neurons do not code for theta-rhythm activity. **(C)** Average waveform.

The mean firing rate of each identified SOV-dependent neuron during each SOV is summarized

in Fig. 6. The firing rates for each neuronal group (rows) during different SOVs (columns) are

separated based on light (orange dots) and dark cycles (blue dots). For each neuronal group

the firing rates are normalized to the maximum observed firing rate of that group across all

SOVs (M_{FR}, shown in black dots). The neuronal groups are separated by a combination of

anatomical location and SOV-dependence, regardless of differences in their waveform patterns.

For example all the identified PPT REM-active neurons (Fig. 3 C&D) are grouped together. We

found that although different in their waveforms, the neurons had similar activity profiles (i.e.

mean firing rate, and light/dark patterns).

The data presented here pertains to the state-dependent neurons in the targeted SWRN. We

aimed to investigate the original flip-flop model of sleep-wake regulation which identifies certain

regions as only active during certain SOVs (i.e. NREM-active neurons in VLPO) (Saper et al.,

2001; Diniz Behn and Booth, 2010). Thus, although most of the targeted regions contain

multiple populations of neurons with diverse state-firing dependency, the neuronal activity

presented here specifically includes NREM-active subpopulation of VLPO, REM and wake-REM

active subpopulations of PPT and LDT, wake-active subpopulation of DR, REM-off

subpopulation of vlPAG, and IS-active neurons that are potentially located in medial region of

the preoptic area.

We found that the population mean and general activity profile of these state-dependent

neurons were similar to previous reports (Datta and Siwek, 2002; Sakai, 2011, 2018; Alam et

al., 2014; Boucetta et al., 2014). We observed some level of variability in mean firing rates for

each neuronal group (indicated by green stars in Fig. 6) during light or dark periods. However,

the variance of the firing rate distributions, specially the M_{FR} (black dots) was much lower

compared to what is reported in previous studies. The statistics of these distributions are highly

affected by how long and over how many SOVs each neuron's activity is monitored. Because

we have such a large sample population for each state, and for each neuron, we have a longer

(and more stable) observation window over the behavior of the neuron which inherently reduces

the variance. This can in turn explain the differences in distribution width compared to previous

work.

Of the represented neuronal groups, VLPO is one of the most challenging structures for

continuous stable recordings. This explains the dearth of reports of continuous electrode

recordings, and the fact that such recordings, when available, are performed in short bouts

(Szymusiak et al., 1998) or when the animal's movements are restricted (Takahashi et al., 2009;

Alam et al., 2014). To our knowledge, we are the first to report consistent multi-day continuous

observations of VLPO neuronal activity. The large number of NREM bouts wherein VLPO

neurons were highly active enabled calculations of firing rate with low margin of error and

provided sufficient power to compare neuronal activity over different SOVs.

We found a slight difference in the mean state-dependent firing rate of a subset of neuronal

groups during light versus dark periods. These included the wake-REM active neurons of PPT

and LDT and the REM-off neurons of vlPAG. However, this circadian difference was neither

statistically significant nor consistent in all groups. This is in line with our previous work showing

no clear difference in neurons' ISI and average waveform in dark versus light periods (Fig. 9 in

(Billard et al., 2018)).

Figure 6. **Mean and standard deviation of the firing rates of all recorded neurons in brainstem-hypothalamic sleep-wake regulatory network.** Firing rates of state-dependent neurons whose recording sites were histologically confirmed pooled from all recording days across all animals. The neurons are separated based on their location and activity dependence. The firing rate of each neuron is marked by orange for the light period and blue for the dark period. For each neuron the firing rate is normalized to the maximum firing rate (black dots) of the group across all states of vigilance (M_{FR}). Although the firing rates in all population were somewhat consistent, we found higher variability in subset of neuronal populations (green stars). In PPT and LDT we found a variety of neurons that were REM-active (RA), Wake and REM active (WRA), or Wake active (WA). VLPO, ventrolateral preoptical area; vlPAG, ventrolateral periacquaductal gray; PPT, pedunculopontine tegmentum; LDT, laterodorsal tegmentum; MPA, medial preoptic area; DR, dorsal raphe.

3.2. Investigation of causality of neuronal activity in emergence of cortical signs of SOV

Our aim was to investigate the role of state-dependent neurons, within the conventional brainstem-hypothalamic flip flop model of sleep-wake regulation, in the emergence of SOVs and SOV transitions. There is a tendency to implicitly interpret the term 'XXX-active neuron' (i.e. REM-active) as a neuron whose activity is causal of initiating the 'XXX' state (i.e. REM state). An alternative role denoted in the literature is that such neurons are involved in the maintenance of the state in which they are active (i.e. (Grace et al., 2014; Xu et al., 2015; Weber and Dan, 2016; Sakai, 2018)). We therefore investigated whether changes in firing rate could be causal in state initiation – as indicated by significantly increasing firing rate ahead of the observed transition time.

Neuronal firing rates over many sleep cycles (~100-300) were derived from simultaneous measurements of single-unit activity from a subset of the SWRN. For each SOV transition (from state A to B), we collected the distribution of firing rates while in state A. The firing rate was calculated from spike-times away from state transitions by at least 3 seconds. Procedurally, we determined SOV transition times (t_{trans}) as the time when EEG and behavioral signs of state B first emerged. Such signs emerge from brain regions downstream from the SWRN, and therefore t_{trans} as defined must be delayed from transitions in activity state of the SWRN. Changes in firing rate of a particular neuron or neuron type that follow t_{trans} cannot be causal for the transition, while those that precede it can but are not guaranteed to be so.

Following the Poisson process statistics (see Methods) we determined the time-point that the instantaneous state-conditioned firing rate during state B differed significantly from the firing rate in state A. The time-point is denoted as t_{sig}. The large ensemble of firing rates for each neuron during each transition allows for sub-second resolution determination of t_{sig}.

If the significant change in the state-conditioned firing rate occurred prior to transition time

$t_{sig} < t_{trans}$, then it is counted as negative t_{sig}. Positive t_{sig} indicates that the state-conditioned firing

rate changed after transition time, $t_{sig} > t_{trans}$. Therefore, a positive t_{sig} implies that the changes in

firing of that neuron type **cannot** be causal for the state transition. In addition, our sleep scoring

was done with 1s discretization. Therefore, there may be as much as an average of 0.5s delay

in marking t_{trans} from the actual change in signs. This means that an even more conservative

scoring to which changes in neuronal firing could be causal would be $t_{sig} < t_{trans}$-0.5.

The t_{sig} values for the 428 identified state-dependent neurons are marked by blue dots in Fig. 7.

The gray shading indicates the positive t_{sig} and therefore non-causal relationships. The white

shading indicates the negative t_{sig} where the change in firing rate occurred prior to state

transition. For each cell group and state transition, dashed gray lines indicate the representative

average state-dependent firing rate as a function of time during state transitions. We found that

the t_{sig} values for individual cells are tightly distributed within each cell group for each state

transition. This indicates the consistency of neuronal activity during state transitions.

Cell groups whose mean firing rates increase during state transitions (e.g. REM-active PPT or

LDT neurons during NREM to REM transitions) can either turn on to cause the state ($t_{sig} < 0$,

white shading in Fig. 7), which means high discharge rates prior to emergence of cortical signs

of a SOV, or can be turned on as a consequence of state transition ($t_{sig} > 0$, gray shading in Fig.

7). Among all cell groups with increasing firing rate during state transitions, only NREM-active

neurons of VLPO and REM-off neurons of vlPAG during REM to NREM transitions had negative

t_{sig}. The discharge rate of all other neurons increases *after* the appearance of EEG and

behavioral signs of SOV. Therefore, the increase in their discharge rate does not mean that they

are causing the state change, only that their behavior is a consequence of the state change.

Likewise, cell groups whose mean firing rate decreases during state transitions (e.g. wake-

active DR neurons during wake to NREM transitions) can either be responsible for the

emergence of the state ($t_{sig} < 0$, white shading in Fig. 7), or be inhibited as a consequence of the

state ($t_{sig} > 0$, gray shading in Fig. 7). We found that all the cell groups that decreased their firing

rates during state transitions did so prior to transition time ($t_{sig} < 0$), which could allow a new

SOV to emerge. We anticipate that such inactivation is sourced from changes in the activity of

the presynaptic source.

The NREM-active neurons of VLPO are the only population with negative t_{sig} values during all

state transitions. Specifically, during transitions into NREM the activity of the VLPO neurons

increases significantly with respect to the state-conditioned average firing rate prior to the state

transition. These neurons also decrease their firing rate significantly prior to transitions out of

NREM. Therefore, the causality hypothesis (Szymusiak et al., 1998; Saper et al., 2005, 2010)

holds for the NREM-active neurons of VLPO.

Figure 7. The t_{sig} for all neurons with respect to cortically defined state transitions. Each

column indicates a state transition, where the transition is marked by time zero. The state

before and after the transitions are marked by the white and gray backgrounds, respectively.

The t_{sig} for each neuron is indicated by a blue dot. Representative average firing rates during

transition are shown by the dashed gray lines. The firing rates for DR, PPT/LDT$_{RA}$, PPT/LDT$_{WRA}$

groups are normalized to half height of the plot box. Other firing rates are normalized to the full

height of the plot box. Notice that for every neuronal group and transition, the t_{sig} values are

tightly distributed. This indicates that the activity of these neurons is extremely consistent. The

gray shading indicates the positive t_{sig} and not causal relationships while the white areas are

negative t_{sig} indicating that the neuron changed its behavior prior to the appearance of cortical

signs of a state. VLPO, ventrolateral preoptical area; vlPAG, ventrolateral periacquaductal gray;

PPT, pedunculopontine tegmentum; LDT, laterodorsal tegmentum; MPA, medial preoptic area;

DR, dorsal raphe.

491 3.3. Neuronal activity in preoptic area and vlPAG demonstrate distinct
492 dynamics during IS

Because we found that IS seems to be characteristic of ending of NREM and often precedes

REM, we utilized the t_{sig} values to investigate the behavior of REM and NREM active neurons

during IS transitions. During NREM to IS transitions, the discharge rates of both the solely

NREM-active neurons of VLPO and the REM-off neurons of vlPAG decreased prior to transition

into IS ($t_{sig} < 0$). The behavior of NREM-active neurons implies a level of causality, but whether

the decrease in firing rate is programmed internally or caused by other factors remains to be

investigated. In contrast, the discharge rate of the REM-active as well as REM-wake-active

neurons of PPT and LDT does not change during NREM to IS transitions.

The IS-active neurons remained active with no substantial change in their firing rate during IS to

REM transitions. However, we observed a statistically significant decrease in their firing rate

during IS to wake transitions ($t_{sig} < 0$). Similarly these neurons stopped firing prior to transitions

out of REM to NREM or wake ($t_{sig} < 0$). These indicate that IS-active neurons might be coding

for transitions out of IS (or REM).

The activity patterns of the NREM-active neurons in VLPO, REM-off neurons in vlPAG, and IS-
active neurons in the preoptic area provide another line of evidence that IS is a distinct state of
vigilance rather than only a transitory dynamic between NREM and REM.

4. Discussion

We developed a dataset including 569 cumulative days of SOV-scored single- and multi-unit
measurements from multiple cell-groups in the sleep-wake regulatory network (SWRN). These
measurements are from 12 freely behaving rodents, with each rat recorded for over 30 day
periods, which consisted of multiple 3-10 day long segments of continuous recordings. The
dataset (previously presented in (Billard et al., 2018)) includes unit activity from up to five
noncoplanar, noncollinear targets of the SWRN (PPT, LDT, DR, vlPAG, VLPO, and MPA), along
with ECoG and LFP recordings of cortex and hippocampus in freely behaving rats.

Current theory about wake-sleep regulation posits that there are wake-promoting and NREM-
promoting cell groups, and that these inhibit one another. This theory would predict changes in
the wake-promoting groups before awakening from NREM sleep, and in the NREM-promoting
neurons prior to transitions into NREM sleep. Our dataset complements previous work testing
this theory (Aston-Jones et al., 2001; Datta and MacLean, 2007; Saper et al., 2010; Takahashi
et al., 2010; Sakai, 2011; Alam et al., 2014; Boucetta et al., 2014; Van Dort et al., 2015; Cox et
al., 2016; Weber and Dan, 2016), but is characterized by several significant distinctions.

First, our simultaneous measurements of hippocampal LFP, ECoG, head acceleration, and
neuronal activity from multiple cell-groups in freely behaving animals over long (3-10 day)
continuous recording periods in the animals' home cages represent a novel "systems
neuroscience" attempt aimed at characterizing the dynamics of the brainstem-hypothalamic
SWRN system described by Saper et al. (Saper et al., 2001, 2010). Second, the data include
continuous and simultaneous measurements from multiple cell groups of SWRN (including

NREM-active VLPO neurons) over many (100-300) consecutive sleep-cycles without introduction of any restrictions to affect animal behavior. Third, our approach to sleep scoring resulted in continuous and temporally precise determination of SOV and SOV transition times. Fourth, time-resolved analysis of the SOV and SOV transition times as well as acquisition of sufficient sleep-wake cycles provide suitable statistical power to probe causality between neuronal firing rates and emergence of cortical signs of states. Finally, we identified a subset of neurons, potentially located in or close to the medial preoptic area (MPA) that code for the intermediate state (IS, (GOTTESMANN, 1996; Boucetta et al., 2014)). We therefore characterized IS as a distinct sleep state from NREM that dominantly transitions to REM.

Much of the research investigating the neuronal activity of the cell-groups in the SWRN is performed within conditions that inherently modulate the animals' sleep cycles and/or limit the time-resolution required for determination of the relationship between firing rate and state transitions. Almost all sleep studies available today are performed with head-fixed animals (Takahashi et al., 2006, 2010; Sakai, 2012, 2018), or in novel environments such as isolation boxes, and provide short recordings of the sleep cycles (Szymusiak et al., 1998; Datta and Siwek, 2002; Takahashi et al., 2006; Sakai, 2011, 2012). While such methods avoid the technical challenges associated with simultaneously recording the hippocampal, cortical, and behavioral indices of different sleep states as wells as state-dependent neuronal activity, they complicate the analysis of SWRN dynamics.

Deviations from normal home-cage environments change the animals' normal behavior and sleep-wake cycles. Short-term recordings from restricted animals eliminate the effect of exogenous (light-dark cycles) and endogenous circadian cycles on sleep dynamics. Furthermore, to detect a statistically significant change in neuronal firing rates during state transitions, with sub-second temporal resolution, many sleep cycles are required. Finally, conventional SOV scoring techniques assign a score to non-overlapping time-windows

(Takahashi et al., 2009, 2010; Bastianini et al., 2017; Sakai, 2018; Grieger et al., 2021; Huffman et al., 2021) with fixed length and predetermined boundaries. These fixed windows prevent accurate assessment of the SOV dynamics and identification of the true onset of a state.

4.1. Implications of the intermediate state for sleep-wake regulation

Gottesmann was the first to identify an intermediate state in Wistar rats that occurred between NREM and REM, and which exhibited both cortical spindles (a sign of NREM) and hippocampal theta rhythm (a sign of REM) (Gottesmann, 1964) . Gottesmann's finding of IS was further confirmed both in later rat experiments (Gandolfo et al., 1990; Neckelmann et al., 1994; GOTTESMANN, 1996) and observations of human sleep (Lairy, 1966). More recently this state has appeared in a small subset of publications focused on sleep-wake regulation in healthy (Gervasoni, 2004; Boucetta et al., 2014) and diseased brain (Andrade et al., 2017; Pitkänen et al., 2019). We also identified IS in our animals and found distinct temporal, spectral, and neuronal patterns associated with it.

The conventional/routine sleep-scoring analyses assume state transitions are discrete and easily classified by fast changes in EEG features, and in behavioral and physiological activity represented by EMG power, head acceleration, heart rate, and/or breathing. Therefore, research has focused on identifying brain regions involved in triggering discrete transitions between these states. However, the existence of a state with clear spectral and neuronal characteristics that combines and expands upon the features of both NREM and REM in the transition from NREM to REM or to wakefulness poses new questions regarding the mechanisms underlying the onset of any SOV.

The IS-active neurons reported here remained active during IS to REM transitions. However, these neurons turned off prior to transitions out of IS and into wake. Therefore, we hypothesize that the activity of these neurons determines whether transitions out of IS progress into REM or

wake. Based on our histological analysis, these neurons might be located in the medial regions

of the preoptic area.

The technical challenges in measuring from a number of SWRN indicate the value of SWRN

mathematical models (see for example (Tamakawa et al., 2006; Fleshner et al., 2011)) for

understanding the underlying mechanisms of sleep-wake regulation. However, the models and

the experiments they are based on have generally not considered the existence of the IS or the

unique sets of IS-active neurons. Implementation of the IS as a distinct SOV with its own state-

active neurons will enable the models to provide mechanistic insight into differentiation between

transitions to wake versus transitions to REM.

4.2. Implications of correlative vs. causal activity in the brainstem-
hypothalamic SWRN

We used time-resolved transition-based analysis to probe the causality of the SWRN dynamics

in transitions between states of vigilance. With the exception of NREM-active VLPO neurons,

increases in state-conditioned mean firing rate always followed the emergence of the state

which is determined from cortical and behavioral signs. A change in neuronal activity that

follows the SOV cannot be programming it.

Previous work including brief recordings (Datta and Siwek, 2002) and lesion experiments (Saper

et al., 2001) has indicated that PPT neurons are involved in both the induction and the

maintenance of REM from NREM sleep and are a precursor to the signs of REM (Datta and

MacLean, 2007). We too observed increases in PPT activity during REM. However, we

observed increased PPT activity only *after* emergence of cortical signs of REM. This questions

the hypothesized role of cholinergic REM-on PPT neurons in REM induction. We suspect that

this mismatch is due to two factors. First, the spectral and temporal features used in our sleep-

scoring are generated via causal filters and updated every second at the end of computation

window. Therefore, all the scores are based on information up to the time leading to the marked

state transition and *not* after that. This is in contrast to fixed length windows used in

conventional sleep-scoring that are susceptible to missing the correct transition time, or hand-

scored versions that use a-causal potentially biased methods to estimate transition times.

Second, for each neuron we collected a large ensemble of firing rates computed over many

sleep-wake cycles. The large ensemble size enables sub-second resolution in determining peri-

transition-time changes in firing rate (see Methods).

We observed a similar behavior for other identified SWRN nuclei. Neurons in REM-active LDT,

Wake/REM-active PPT and LDT, Wake-active DR, Wake-active PPT, and REM-off vlPAG cell

groups all show a significant change in their firing rate only *after* their corresponding state

transitions.

The cell groups involved in sleep-wake regulation are functionally and spatially heterogeneous

and extend across large neural territories. Many aspects of how such a large number of variable

cell groups interact to govern rapid ultradian transitions within a sleep cycle (i.e. NREM to REM)

as well as much slower homeostatic and circadian processes are still unknown. Acute

disruptions of the interactions within the SWRN via anatagonist injections or optogenetic

inhibition are shown to cause acute loss of the corresponding SOV. However, chronic ablation

of the basal forebrain cholinergic neurons (Kaur et al., 2008), the tuberomammillary

histaminergic neurons (Gerashchenko et al., 2004), the LC and pontine cholinergic neurons

(Webster and Jones, 1988; Shouse and Siegel, 1992; Lu, 2006), or combinations of these

structures (Blanco-Centurion et al., 2007) have minimal effects on the regulation of that SOV.

Further, almost all of the SWRN contain intermingled cell types that are active during more than

one SOV. For example in the basal forebrain, glutamatergic, cholinergic, and parvalbumin (PV)-

expressing GABAergic neurons are shown to be active during both REM and wake, but their

optogenetic activation promotes only wakefulness (Xu et al., 2015). A routine based on acute

study of stimulus (lesion/optogenetics)-response (i.e. changes to SOV length and occurrence)

dynamics might not be fitting for investigation of a naturally occurring spontaneous behavior

such as sleep-wake transition and cycling. Therefore, although internal/external stimuli might

cause transitions into a state, it does not necessarily mean that the stimulated neurons are

causal for the emergence of the state in normal conditions.

The *in vivo* experimental approach described here and in (Billard et al., 2018) allows for

simultaneous recording of neuronal activity in multiple deep brain structures throughout many

days in freely behaving animals. As evidenced by the findings reported here, this approach has

the potential to uncover new insights on mechanisms underlying the dynamics of sleep-wake

regulation.

5. Funding sources

639 This work was supported by the National Institutes of Health (NIH) grant (No. R01EB019804)

640 and a doctoral Academic Computing Fellowship from the Pennsylvania State University to F.B.

641 Declaration of interest: none.

6. Appendix

Animal # (Sex)	Days Recorded [# of Sessions]	PPT Left	PPT Right	LDT Left	LDT Right	DR	LC	VLPO Right
1 (M)	41 [13]	(red)	(red)			47 (23)		
2 (M)	37 [12]	42 (21)	33 (13)			43 (17)		
3 (M)	57 [9]	29 (19)	29(8)			18 (10)		
4 (M)	73 [12]	7 (1)	(red)			18 (7)		
5 (M)	66 [10]	20 (9)			14 (4)	27 (10)		
6 (M)	43 [8]	24 (20)			29 (22)	32 (11)		26 (10)
7 (F)	39 [7]		6 (4)	19 (12)		3 (2)		9 (4)
8 (F)	54 [9]	K			47 (28)	44 (31)		(red)
9 (M)	57 [9]	8 (6)			43 (21)	23 (3)		13 (5)
10 (M)	34 [8]		42 (27)	K		15 (9)		16 (6)
11 (M)	29 [8]		25 (17)	30 (19)		(red)		16 (5)
12 (M)	39 [6]		30 (14)	16 (7)		K	4 (0)	8 (3)
Total Neuron Count SOV-Dependent Neurons ()		295 (159)		198 (113)		270 (123)	4 (0)	88 (33)

(green)	(orange)	(red)
Record & Histological validation	Record from target successfully	Record but no neuron crossing threshold (>7 std)

Table S1. Detailed recording information from brainstem-hypothalamic structures. The cells include information about which targets were recorded from successfully as well as the number of well-defined SOV-dependent and total units counted over the entire course of recordings for each axis and animal. The final row is the sum of SOV and total units over all animals in the cohort. Animals 1-5 had three Microdrive axes, animals 6-11 had a total of four microdrive axes, and animal 12 had a total of five microdrive axes. Formatting of cells refer to the following conditions: recorded units with histological validation that the electrodes hit the target (green); recorded units with no histological validation (orange). For this group the state-dependency of the unit-recordings was established based on EEG and behavioral measures; recorded activity but no units crossed the 7 SD threshold (red). Some implanted microwire bundles were kinked (K) during the Microdrive implant procedure or later driving sessions, which prevented electrodes from driving. Sessions were continuous recording periods between electrode-driving sessions. Recording sessions typically lasted between 3 and 10 days.

6.1. Characterization of the intermediate state

Figure S1. Time-frequency characteristics of the intermediate state. The IS was observed in almost all NREM to REM transitions. **(A)** In an example hippocampal LFP recording in one animal, the power spectrum during NREM shows lower frequency activity in the delta range (high delta peak in red in B). The transition to REM is not discrete. As the brain exits NREM, power increases in the theta range appear simultaneous with the persisting peaks in the delta band (theta and delta peaks in magenta in B). The IS onset and end are marked by the solid magenta lines. The REM onset is then marked by a clear consistent peak in the theta band (high theta peak in green in B). Example time-series traces for NREM (red panel, ECoG), IS (magenta panel, LFP), and REM (green panel, LFP) are shown in the bottom panels. All vertical and horizontal scale bars are 100 mV and 1 second; respectively. Power is calculated over 8 second windows with 4 second overlap. **(B)** The simultaneous delta and theta activity in the IS

is highlighted in the power plot averaged over 20 second long time-series for each state. Power

is calculated from the same hippocampal LFP recording as in A over 2 second long windows

with 1 second overlap. **(C)** Scatter plot of the first two linear canonical discriminants highlight the

three main groups. Each dot corresponds to a 2 second window for which the spectral features

were calculated. For clarity, the dataset presented here is from 48 hour of recording in one

normal animal. After the clustering analysis, within each cluster, the 95% most-inclusive contour

was chosen as the boundary of that cluster. We define the transitions between states as

trajectories connecting the clusters in the canonical space. Almost all transitions from NREM to

REM (yellow) or to Wake (red) seem to first dwell in the IS region (black dots). **(D)** Shown in the

upper panel are the possible transitions between SOVs. The majority of transitions from NREM

to REM passed the IS region, therefore direct transitions (much fewer) were marked by a red

dashed line. The transition probabilities between SOVs, across all animals, are shown in the

bottom panel. The majority of transitions out of IS state end in REM (78%), whereas only 22% of

those transitions end in Wake.

Figure S2. IS patterns and duration across the light/dark cycle for all rats. (A) Hourly

percentages spent in NREM (red), IS (magenta), REM (green), and Wake (blue) are shown for

all rats. The majority of the light period (yellow) is spent in sleep and the majority of the dark

period (gray) is spent in Wake. **(B)** Distribution of the duration of the IS pooled from all animals

both during light (blue) and dark (orange) periods. Regardless of the time of day, the

intermediate state lasts for a considerable amount of time that should be detectable in the

current sleep-scoring algorithms. **(C)** Mean and standard deviation of percent of day spent in IS

for light and dark period. Difference in the light/dark cycle was calculated using a one-sided

unequal variance t-test with alpha=0.05. Although animals exhibit an increase in daily IS time

during the light cycle, it is not significant. **(D)** IS bouts pooled across all rats (n=13) were scored

as either IS → REM or IS → Wake. Hazard rates were calculated using competing groups.

700 Competing groups analysis assumes one event precludes the other. λ_{REM} is on average 4 times

701 larger than λ_{Wake}. This ratio indicates the relative frequency of REM and Wake transition types in

702 the data set. The λ_{REM} first grows and then quickly decays, with peaks at IS duration of 30 and

703 70 seconds. λ_{Wake} remains comparatively flat with a small peak at 80.

6.2. Characterization of neural activity during state transitions

Our main goal is to understand the role of brainstem and hypothalamic neurons in the

emergence of sleep-wake states and the transitions between them. By clarifying the temporal

relationship between spontaneous neuronal activity and state transitions in freely behaving

animals, we can elucidate the role of specific cell-groups in the initiation or maintenance of a

SOV. Therefore, we acquired simultaneous measurements of single- and multi-unit activity from

a subset of the sleep-wake regulatory cell-groups as well as hippocampal LFP and ECoG.

As described in the methods, SOV and state transition times were defined from the

hippocampal, cortical, and behavioral signs of the state, using spectral features of the

hippocampal LFP, ECoG, and head acceleration.

We first examined the activity of each identified neuron during all states of vigilance. We

extracted the firing rates for the identified neurons and validated their location via histological

analysis. An example of these processing and analysis steps for an identified REM-active

neuron in the PPT cell-group is shown in Fig. S3. The data shown are from one animal, over all

IS to REM instances (n = 168) in one 5-day recording session. The histological analysis

(Fig. S3A) confirmed that the electrode was in PPT. Likewise, the neuron's waveform (Fig. S3B)

and inter-spike interval histogram (ISIH) is consistent with cholinergic cells of the PPT (Fig.

S3C) (Satoh and Fibiger, 1986; Datta and Siwek, 2002; Grace et al., 2014; Van Dort et al.,

2015).

Every detected spike of this particular neuron during every IS to REM transition was marked

(raster plot in Fig. S3D upper panel) and the average state-conditioned firing rate, as a function

of time with respect to transition time, was computed. The average state conditioned firing rate

(over all REM to IS incidents) for this particular PPT neuron during both the IS and REM is

shown in black traces in the bottom panel of Fig. S3D. Consistent with a REM-active neuron

(one whose activity is high during REM) the average firing rate is nearly constant for times

positive with respect to the transition time, which is consistent with the raster-plot having

consistent firing in the green-shaded regions (Fig. S3D upper panel). Note that the variance in

firing rate goes up for large time periods because there are fewer very long REM periods over

which it is calculated.

To calculate changes in firing rate during state transitions, we used Poisson statistics. The pre-

transition average firing rate over many trials formed the reference distribution. An example of

the mean of the average firing rate distribution during the all the ISs is indicated by the solid

dashed line in the lower panel of Fig. S3D. We then used the cumulative distribution function

with 1%-99% confidence intervals to find the post-transition instantaneous firing rate outside of

the confidence bounds. The time point associated with the significantly different instantaneous

firing rate was then determined as t_{sig}. The orange dashed line in the lower panel of Fig. S3D

indicated the t_{sig} for this specific neuron.

In addition to the state-conditioned firing rates, we simultaneously compute and report the

probability of being in a state, P(state). The green and magenta lines in the bottom panel of

Fig. S1D indicate the P(IS) and P(REM). By construction, according to the homeostatic sleep

drive, once the dynamics transition into a state, the probability of staying in that state

monotonically decreases over time. The P(state) shown in Fig. S3D is an example of this

phenomenon, where the P(REM) monotonically decreases during time spent in REM.

The procedural steps detailed in Fig. S3 for one PPT neuron were implemented to collect the

state-conditioned firing rates and P(state) for all identified neurons as well as for all periods

before and after every allowed transition type.

Figure S3. Activity analysis of a representative REM-active PPT neuron. **(A)** NADPH-stained coronal section of the brain show the cholinergic population in the PPT as well as the electrode track in the PPT. **(B)** Average waveform of the identified action potential for one PPT neuron over one 5-day long driving session. **(C)** The inter-spike interval (ISI) of the neuron. **(D)** Peri-REM firing rate of a PPT neuron in one animal. The raster plot is ranked according to the duration of REM bouts and with respect to time since transition into REM. The firing activity of the neuron is indicated by the black markers. In a raster plot format they each indicate every time an action potential is recorded. We calculated the probability of each state given time from state onset; indicated by the dashed lines in the bottom panel (magenta for IS; P(IS) and green for REM; P(REM)). On average, the PPT neuron fires much more infrequently during the IS bout, and only significantly increases its activity during REM (green area). The significance is

calculated based on the Poisson statistics, where the reference firing rate distribution is the

average firing rate of the prior state (here, IS) over many trials indicated by the solid orange line.

The time point when the instantaneous firing rate falls out of the 1%-99% confidence bounds is

marked as t_{sig} marked here by the orange dashed line. As P(REM) decays, so does the average

firing rate. The average firing rate here indicates the neuronal activity conditioned on state and

is calculated in 1 second bins. The probability traces are rescaled from [0,1] to [0,10] to match

the scale of the average firing rate.

References

Alam MA, Kumar S, McGinty D, Alam MN, Szymusiak R (2014) Neuronal activity in the preoptic hypothalamus during sleep deprivation and recovery sleep. J Neurophysiol 111:287–299.

Andrade P, Nissinen J, Pitkänen A (2017) Generalized Seizures after Experimental Traumatic Brain Injury Occur at the Transition from Slow-Wave to Rapid Eye Movement Sleep. J Neurotrauma.

Aston-Jones G, Chen S, Zhu Y, Oshinsky ML (2001) A neural circuit for circadian regulation of arousal. Nat Neurosci 4:732–738.

Bastianini S, Alvente S, Berteotti C, Lo Martire V, Silvani A, Swoap SJ, Valli A, Zoccoli G, Cohen G (2017) Accurate discrimination of the wake-sleep states of mice using non-invasive whole-body plethysmography. Sci Rep 7.

Billard MW, Bahari F, Kimbugwe J, Alloway KD, Gluckman BJ (2018) The systemdrive: A multisite, multiregion microdrive with independent drive axis angling for chronic multimodal systems neuroscience recordings in freely behaving animals. eNeuro 5:ENEURO.0261-18.2018.

Bjorness TE, Dale N, Mettlach G, Sonneborn A, Sahin B, Fienberg AA, Yanagisawa M, Bibb JA, Greene RW (2016) An Adenosine-Mediated Glial-Neuronal Circuit for Homeostatic Sleep. J Neurosci 36:3709–3721 Available at: http://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.3906-15.2016.

Blanco-Centurion C, Gerashchenko D, Shiromani PJ (2007) Effects of Saporin-Induced Lesions of Three Arousal Populations on Daily Levels of Sleep and Wake. J Neurosci 27:14041–14048 Available at: http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.3217-07.2007.

Boissard R, Gervasoni D, Schmidt MH, Barbagli B, Fort P, Luppi P-H (2002) The rat ponto-medullary network responsible for paradoxical sleep onset and maintenance: a combined microinjection and functional neuroanatomical study. Eur J Neurosci 16:1959–1973 Available at: http://doi.wiley.com/10.1046/j.1460-9568.2002.02257.x.

Boucetta S, Cisse Y, Mainville L, Morales M, Jones BE (2014) Discharge Profiles across the Sleep-Waking Cycle of Identified Cholinergic, GABAergic, and Glutamatergic Neurons in the Pontomesencephalic Tegmentum of the Rat. J Neurosci 34:4708–4727.

Cox J, Pinto L, Dan Y (2016) Calcium imaging of sleep–wake related neuronal activity in the dorsal pons. Nat Commun 7:10763.

Datta S, MacLean RR (2007) Neurobiological mechanisms for the regulation of mammalian sleep–wake behavior: reinterpretation of historical evidence and inclusion of contemporary cellular and molecular evidence. Neurosci Biobehav Rev 31:775–824.

Datta S, Siwek DF (2002) Single cell activity patterns of pedunculopontine tegmentum neurons across the sleep-wake cycle in the freely moving rats. J Neurosci Res 70:611–621.

Diniz Behn CG, Booth V (2010) Simulating Microinjection Experiments in a Novel Model of the Rat Sleep-Wake Regulatory Network. J Neurophysiol 103:1937–1953 Available at: http://jn.physiology.org/content/103/4/1937.abstract.

ECONOMO C V (1930) SLEEP AS A PROBLEM OF LOCALIZATION. J Nerv Ment Dis 71:249–259 Available at: https://insights.ovid.com/crossref?an=00005053-193003000-00001.

Fleshner M, Booth V, Forger DB, Diniz Behn CG (2011) Circadian regulation of sleep–wake behaviour in
nocturnal rats requires multiple signals from suprachiasmatic nucleus. Philos Trans R Soc A Math
Phys Eng Sci 369:3855–3883 Available at:
https://royalsocietypublishing.org/doi/10.1098/rsta.2011.0085.

Fulcher BD, Phillips AJK, Robinson PA (2008) Modeling the impact of impulsive stimuli on sleep-wake
dynamics. Phys Rev E 78:51920 Available at:
https://link.aps.org/doi/10.1103/PhysRevE.78.051920.

Gandolfo G, Romettino S, Gottesmann C, Van Luijtelaar G, Coenen A (1990) Genetically epileptic rats
show a pronounced intermediate stage of sleep. Physiol Behav 47:213–215.

Gerashchenko D, Chou TC, Blanco-Centurion CA, Saper CB, Shiromani PJ (2004) Effects of Lesions of the
Histaminergic Tuberomammillary Nucleus on Spontaneous Sleep in Rats. Sleep 27:1275–1281
Available at: https://academic.oup.com/sleep/article-lookup/doi/10.1093/sleep/27.7.1275.

Gervasoni D (2004) Global Forebrain Dynamics Predict Rat Behavioral States and Their Transitions. J
Neurosci 24:11137–11147 Available at:
http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.3524-04.2004.

Gottesmann C (1964) Données sur l'activité corticale au cours du sommeil profond chez le rat. CR Soc
Biol(Paris) 158:1829–1834.

GOTTESMANN C (1996) The Transition from Slow-wave Sleep to Paradoxical Sleep: Evolving Facts and
Concepts of the Neurophysiological Processes Underlying the Intermediate Stage of Sleep.
Neurosci Biobehav Rev 20:367–387 Available at:
https://linkinghub.elsevier.com/retrieve/pii/0149763495000550.

Grace KP, Vanstone LE, Horner RL (2014) Endogenous Cholinergic Input to the Pontine REM Sleep
Generator Is Not Required for REM Sleep to Occur. J Neurosci 34:14198–14209.

Grieger N, Schwabedal JTC, Wendel S, Ritze Y, Bialonski S (2021) Automated scoring of pre-REM sleep in
mice with deep learning. Sci Rep 11.

Huffman D, Ajwad A, Yaghouby F, O'Hara BF, Sunderam S (2021) A real-time sleep scoring framework for
closed-loop sleep manipulation in mice. J Sleep Res 30.

Kaur S, Junek A, Black MA, Semba K (2008) Effects of Ibotenate and 192IgG-Saporin Lesions of the
Nucleus Basalis Magnocellularis/Substantia Innominata on Spontaneous Sleep and Wake States
and on Recovery Sleep after Sleep Deprivation in Rats. J Neurosci.

Lairy GC (1966) Données récentes sur la physiologie et la physiopathologie de l'activité onirique.
Données EEG chez le malade mental. IV World Congr Psychiatry 1:189–197.

Lu J (2006) Identification of Wake-Active Dopaminergic Neurons in the Ventral Periaqueductal Gray
Matter. J Neurosci.

Lu J, Sherman D, Devor M, Saper CB (2006) A putative flip–flop switch for control of REM sleep. Nature
441:589–594.

Luppi P-H, Gervasoni D, Verret L, Goutagny R, Peyron C, Salvert D, Leger L, Fort P (2006) Paradoxical
(REM) sleep genesis: the switch from an aminergic–cholinergic to a GABAergic–glutamatergic
hypothesis. J Physiol 100:271–283.

Moruzzi G, Magoun HW (1949) Brain stem reticular formation and activation of the EEG.
Electroencephalogr Clin Neurophysiol.

Neckelmann D, Olsen OE, Fagerland S, Ursin R (1994) The reliability and functional validity of visual and
semiautomatic sleep/wake scoring in the Moll-Wistar rat. Sleep.

Phillips AJK, Robinson PA (2008) Sleep deprivation in a quantitative physiologically based model of the
ascending arousal system. J Theor Biol 255:413–423 Available at:
https://linkinghub.elsevier.com/retrieve/pii/S002251930800444X.

Pitkänen A, Ekolle Ndode-Ekane X, Lapinlampi N, Puhakka N (2019) Epilepsy biomarkers – Toward
etiology and pathology specificity. Neurobiol Dis.

Rempe MJ, Best J, Terman D (2010) A mathematical model of the sleep/wake cycle. J Math Biol 60:615–
644.

Sakai K (2011) Sleep-waking discharge profiles of dorsal raphe nucleus neurons in mice. Neuroscience
197:200–224.

Sakai K (2012) Discharge properties of presumed cholinergic and noncholinergic laterodorsal tegmental
neurons related to cortical activation in non-anesthetized mice. Neuroscience 224.

Sakai K (2018) Single unit activity of periaqueductal gray and deep mesencephalic nucleus neurons
involved in sleep stage switching in the mouse. Eur J Neurosci 47:1110–1126.

Saper CB, Chou TC, Scammell TE (2001) The sleep switch: Hypothalamic control of sleep and
wakefulness. Trends Neurosci.

Saper CB, Fuller PM, Pedersen NP, Lu J, Scammell TE (2010) Sleep State Switching. Neuron.

Saper CB, Scammell TE, Lu J (2005) Hypothalamic regulation of sleep and circadian rhythms. Nature
437:1257–1263.

Satoh K, Fibiger HC (1986) Cholinergic neurons of the laterodorsal tegmental nucleus: Efferent and
afferent connections. J Comp Neurol.

Sedigh-Sarvestani M, Thuku GI, Sunderam S, Parkar A, Weinstein SL, Schiff SJ, Gluckman BJ (2014) Rapid
eye movement sleep and hippocampal theta oscillations precede seizure onset in the tetanus toxin
model of temporal lobe epilepsy. J Neurosci 34:1105–1114 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/24453303.

Shanmugasundaram B, Gluckman BJ (2017) Micro-reaction chamber microelectrodes especially for
neural and biointerfaces.

Sherin JE, Elmquist JK, Torrealba F, Saper CB (2018) Innervation of Histaminergic Tuberomammillary
Neurons by GABAergic and Galaninergic Neurons in the Ventrolateral Preoptic Nucleus of the Rat. J
Neurosci.

Shouse MN, Siegel JM (1992) Pontine regulation of REM sleep components in cats: integrity of the
pedunculopontine tegmentum (PPT) is important for phasic events but unnecessary for atonia
during REM sleep. Brain Res 571:50–63 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/1611494.

Sigl-Glöckner J, Seibt J (2019) Peeking into the sleeping brain: Using in vivo imaging in rodents to

886 understand the relationship between sleep and cognition. J Neurosci Methods.

887 Sunderam S, Chernyy N, Peixoto N, Mason JP, Weinstein SL, Schiff SJ, Gluckman BJ (2007) Improved
888  sleep-wake and behavior discrimination using MEMS accelerometers. J Neurosci Methods
889  163:373–383.

890 Szymusiak R, Alam N, Steininger TL, McGinty D (1998) Sleep-waking discharge patterns of ventrolateral
891  preoptic/anterior hypothalamic neurons in rats. Brain Res 803:178–188.

892 Takahashi K, Kayama Y, Lin JS, Sakai K (2010) Locus coeruleus neuronal activity during the sleep-waking
893  cycle in mice. Neuroscience 169:1115–1126.

894 Takahashi K, Lin JS, Sakai K (2006) Neuronal activity of histaminergic tuberomammillary neurons during
895  wake-sleep states in the mouse. J Neurosci 26.

896 Takahashi K, Lin JS, Sakai K (2009) Characterization and mapping of sleep-waking specific neurons in the
897  basal forebrain and preoptic hypothalamus in mice. Neuroscience.

898 Tamakawa Y, Karashima A, Koyama Y, Katayama N, Nakao M (2006) A quartet neural system model
899  orchestrating sleep and wakefulness mechanisms. J Neurophysiol 95:2055–2069.

900 Urbain N, Creamer K, Debonnel G (2006) Electrophysiological diversity of the dorsal raphe cells across
901  the sleep-wake cycle of the rat. J Physiol 573:679–695 Available at:
902  http://doi.wiley.com/10.1113/jphysiol.2006.108514.

903 Van Dort CJ, Zachs DP, Kenny JD, Zheng S, Goldblum RR, Gelwan NA, Ramos DM, Nolan MA, Wang K,
904  Weng F-J, Lin Y, Wilson MA, Brown EN (2015) Optogenetic activation of cholinergic neurons in the
905  PPT or LDT induces REM sleep. Proc Natl Acad Sci.

906 Verret L, Goutagny R, Fort P, Cagnon L, Salvert D, Léger L, Boissard R, Salin P, Peyron C, Luppi PH (2003)
907  A role of melanin-concentrating hormone producing neurons in the central regulation of
908  paradoxical sleep. BMC Neurosci.

909 Weber F, Chung S, Beier KT, Xu M, Luo L, Dan Y (2015) Control of REM sleep by ventral medulla
910  GABAergic neurons. Nature.

911 Weber F, Dan Y (2016) Circuit-based interrogation of sleep control. Nature.

912 Webster HH, Jones BE (1988) Neurotoxic lesions of the dorsolateral pontomesencephalic tegmentum-
913  cholinergic cell area in the cat. II. Effects upon sleep-waking states. Brain Res.

914 Xu M, Chung S, Zhang S, Zhong P, Ma C, Chang W-C, Weissbourd B, Sakai N, Luo L, Nishino S, Dan Y
915  (2015) Basal forebrain circuit for sleep-wake control. Nat Neurosci 18:1641–1647 Available at:
916  http://www.ncbi.nlm.nih.gov/pubmed/26457552.