Deciphering how interneuron specific 3 cells control oriens lacunosum-moleculare cells to contribute to circuit function

Alexandre Guet-McCreight[1,2] and, Frances K Skinner[1,3]

[1]*Krembil Brain Institute, University Health Network, Toronto, ON, Canada*
[2]*Department of Physiology, University of Toronto, Toronto, ON, Canada*
[3]*Departments of Medicine (Neurology) and Physiology, University of Toronto, Toronto, ON, Canada*

Running title: Interneuron specific cells sharpen recruitment

Corresponding Author:
Alexandre Guet-McCreight
alexandre.guet.mccreight@mail.utoronto.ca
University of Toronto

Number of pages: 30

Number of figures: 8

Number of tables: 0

Number of words Abstract: 247

Number of words New & Noteworthy: 75

Number of words Introduction: 1193

Number of words Discussion: 2537

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgements: This research was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC): Discovery Grant (RGPIN - 2016 - 06182) to FKS and Graduate Scholarship (CGSD2 - 504375 - 2017) to AGM. We thank L. Topolnik for reading and providing feedback on this manuscript. We would also like to thank the reviewers for their various comments that helped us make significant improvements in the presentation of our work.

Abstract

The wide diversity of inhibitory cells across the brain makes them suitable to contribute to network dynamics in specialized fashions. However, the contributions of a particular inhibitory cell type in a behaving animal are challenging to untangle as one needs to both record cellular activities and identify the cell type being recorded. Thus, using computational modeling and theory to predict and hypothesize cell-specific contributions is desirable. Here, we examine potential contributions of interneuron-specific 3 (I-S3) cells - an inhibitory interneuron found in CA1 hippocampus that only targets other inhibitory interneurons - during simulated theta rhythms. We use previously developed multi-compartment models of oriens lacunosum-moleculare (OLM) cells, the main target of I-S3 cells, and explore how I-S3 cell inputs during *in vitro* and *in vivo* scenarios contribute to theta. We find that I-S3 cells suppress OLM cell spiking, rather than engender its spiking via post-inhibitory rebound mechanisms, and contribute to theta frequency spike resonance during simulated *in vivo* scenarios. To elicit recruitment similar to *in vitro* experiments, inclusion of disinhibited pyramidal cell inputs is necessary, implying that I-S3 cell firing broadens the window for pyramidal cell disinhibition. Using *in vivo* virtual networks, we show that I-S3 cells contribute to a sharpening of OLM cell recruitment at theta frequencies. Further, shifting the timing of I-S3 cell spiking due to external modulation shifts the timing of the OLM cell firing and thus disinhibitory windows. We propose a specialized contribution of I-S3 cells to create temporally precise coordination of modulation pathways.

New & Noteworthy

How information is processed across different brain structures is an important question that relates to the different functions that the brain performs. Using modelling and theoretical analyses, we show that an inhibitory cell type that only inhibits other inhibitory cells can broaden the window for disinhibition of excitatory cells, manage input pathway switching, and modulate inhibitory cell spiking. This work contributes to knowledge of how coordination between sensory and memory consolidation information can be attained.

Introduction

Across the brain there is a variety of different types of excitatory and inhibitory neurons that control how information is processed (1–7). This diversity spans morphological, electrophysiological and molecular aspects, and examination of specific inhibitory cell types shows that there are distinct neuronal subclasses that can be mapped to function and behaviour (8). However, identifying a cell type goes beyond characterization of any single genetic marker making such mappings a difficult endeavour. Explorations of inhibitory cells have additional challenges since they are typically smaller in size and are rarer than their excitatory cell counterparts, making them more difficult to record from *in vivo*. Even though there are conceptual and technical challenges in classifying cell types, it is clear, for example, that brain diseases can be specific in the cell types that they affect (9).

The CA1 hippocampus, a brain area associated with memory formation (10), stands out in the field by the vast amount of experimental literature characterizing its inhibitory cell types (2, 4, 6, 11). In particular, there are four known types of interneuron-specific (I-S) cells - inhibitory cells that are specialized to primarily target other interneurons, and not pyramidal (PYR) cells (12–18). Of them, we here focus on one type, the interneuron-specific 3 (I-S3) cell type via consideration of its primary cellular targets. I-S3 cells express vasoactive intestinal polypeptide (VIP) and calretinin (CR), and their primary targets are oriens lacunosum moleculare (OLM) cells (12, 15), a cell type that expresses somatostatin (SOM) and inhibits the distal apical dendrites of PYR cells in CA1 (19, 20). PYR cells represent by far the largest proportion of neurons in the CA1 network (2, 6). Inputs carrying "sensory-related information" from entorhinal cortex synapse onto their distal dendrites, and inputs carrying "retrieval-related information" from CA3 synapse onto their proximal dendrites (4, 21). The OLM cells have been shown to gate sensory encoding via the inhibition of PYR cell distal dendrites (20, 22), and I-S3 cells are thus well placed for input-specific information gating in the CA1 area. This dis-inhibitory circuitry (i.e., I-S3 cell inhibition of OLM cells causing disinhibition of PYR cells) is not necessarily unique to the hippocampus since strikingly similar circuitries with VIP+ and SOM+ cell types have also been reported across several different areas of cortex (17). Generally speaking, activation of VIP+ cells *in vivo* across different cortical areas has been associated with improved performances in a variety of different learning and memory paradigms as well as the facilitation of synaptic potentiation in PYR cells (17).

From *in vitro* studies, it has been shown that I-S3 cells might exert control over OLM cells via post-inhibitory rebound (PIR) mechanisms (15). However, it is possible that inhibitory input from I-S3 cells could suppress OLM cell firing or I-S3 cells may indirectly control OLM cell firing via other cellular pathways. Further, it is unknown how and whether *in vitro* observations would translate to *in vivo* settings, and targeting and dissecting out the contributions of I-S3 cell inputs to OLM cell spiking *in vivo* would be technically difficult and challenging. To begin to understand the inhibitory control that I-S3 cells exert over OLM cells, it is first important to understand the various input integration mechanisms that govern OLM cell dendritic physiology.

In addition to having highly active dendritic properties (23–25), OLM cells exhibit prominent sag currents due to

hyperpolarization-activated cation channels (h-channels) (26–28), which can also promote PIR spiking (29). H-

channels are also thought to contribute to spike resonance properties in OLM cells *in vivo* at theta frequencies, but

when tested *in vitro* using dynamic-clamp experimental techniques, OLM cells phase-locked well to theta frequency-

modulated inputs, independently of whether h-channels were blocked (30). However, since dynamic-clamp

experiments are largely limited to somatic injections, possible contributions of dendritic h-channels were not

assessed. Previous computational modelling work did explore OLM cell dendritic h-channel contributions to spike

resonance in which OLM cell models with synapses distributed onto their somatic and dendritic compartments to

generate simplified *in vivo*-like states were used (31). It was found that there is a shift in spike resonance from high to

low theta frequencies if, respectively, dendritic h-channels are present or not (31). Subsequently, a "neuron-to-

model matched" (NMM) approach was used to create new OLM cell models in which there was a tight integration

of experimental data with the multi-compartment model building, so as to determine whether dendritic h-channels

are in fact present or absent (27). These new OLM models created using this NMM approach strongly suggested

that h-channels must be present on OLM cell dendrites.

In parallel, in previous work using I-S3 cell models (32), we had developed an approach for performing high-resolution

parameter searches to find excitatory and inhibitory synaptic input parameter combinations that could generate *in*

vivo-like states, and thus create virtual networks from a particular cell's perspective. We took advantage of this

approach to do the same for these new OLM cell models where we demonstrated that h-channel dendritic currents are

consistently enhanced during *in vivo*-levels of synaptic bombardment (33). Importantly, I-S3 cell inputs are a

major source of inhibitory synaptic inputs to OLM cells (13). Overall, these model developments provide an

opportunity to do a detailed *in silico* exploration of the potential contributions of I-S3 cells in a behaving animal via

its control over OLM cells during theta activities, given all of our previous findings on OLM cell dendritic h-

channels (27, 31), advancements in generating detailed *in vivo* virtual network simulations (32, 33), and discovery of

the activation patterns of I-S3 cells during animal behaviour (16).

In this study, we use computational modelling and theoretical analyses to examine how I-S3 cells manifest control

over OLM cells so as to decipher how these specialized cell types contribute to function. In the strategy

implemented in this work, we use the new OLM cell models (27) with synaptic inputs from I-S3 cells as well as

other inputs (33), and perform simulations with these models under *in vitro* and *in vivo*-like contexts (see **Fig 1**

illustration), as motivated by experimental data specific to OLM and I-S3 cells. Our explorations in the *in vitro*

context are based on observations from Tyan et al. (15) which showed that I-S3 cells could preferentially control the

timing of OLM cell spiking at 5 and 10 Hz (theta frequencies) but not at non-theta frequencies. Our explorations in

the *in vivo* context are based on results from Luo et al. (16) demonstrating that I-S3 cells are activated with a delay

relative to theta-run epochs and spike during the rising to peak phases of theta rhythms. Our models allow a focus

on biophysical details of OLM cells and we use theoretical phase response curve (PRC) analyses from *in vitro*

aspects to motivate and understand mechanisms of entrainment and spiking resonance *in vivo* in considering
specialized contributions of I-S3 cells. In the end, our modeling study predicts that I-S3 cells contribute to function by
suppressing spiking in OLM cells and broadening the window for the synaptic disinhibition of pyramidal cells.
Overall, our work shows that close interfacing of computational and experimental studies can help us make progress
toward the challenging endeavour of mapping specialized cell types to function and behaviour.

Materials and Methods

121 Associated code for all of the simulations that are presented in this work can be found at the following link:
122 `https://github.com/FKSkinnerLab/OLM_IS3control`.

OLM cell models

The models upon which the present work is based have been previously published. Detailed descriptions of these
models with equations and parameter values can be accessed from Guet-McCreight and Skinner (33) and Sekulic et al.
(27). Here we provide a brief description:

Two detailed multi-compartment OLM cell models, cell 1 and cell 2, were developed in NEURON (34) using
a "neuron-to-model matched" (NMM) approach (27). That is, the morphology, passive properties and h-channel
characteristics in each model were specific to cell 1 or 2 as measured in the biological OLM cell, and other
biophysical currents were optimized using electrophysiological data specific to cell 1 or 2. As expected, model
parameter values, such as the biophysical channel conductances, differed between cell 1 and cell 2. Both cell
models were constructed using morphological and electrophysiology data from intermediate CA1 mouse
hippocampus. Code for the models is available from `https://github.com/FKSkinnerLab/OLMng`. Ion
channel mechanisms include: hyperpolarization-activated cation channels (*H*), transient sodium (*NaT*), fast and
slow delayed rectifier potassium (*Kdrf*, *Kdrs*), A-type potassium (*A*), M-type potassium (*M*), T- and L-type
calcium (*CaT*, *CaL*), and calcium-dependent potassium (*KCa*) channels. All of these mechanisms are distributed
throughout the cell, with *CaT*, *CaL*, and *KCa* being inserted only in dendritic compartments, the rest inserted in
somatic and dendritic compartments, and *NaT*, *Kdrf*, and *Kdrs* also being inserted in axonal compartments. We refer
to currents generated by the different ion channel mechanisms with subscripts.

These two OLM cell models developed using an NMM approach are the most up-to-date OLM cell models currently
available (27), and optimized fits to the electrophysiological data support the inclusion of h-channels on the dendrites
of OLM cells. Although each of these models represent OLM cells as captured by mimicking the electrophysiological
recording outputs from the particular biological OLM cell, they differ in their detailed morphologies and conductance
values for each of the various ion channel mechanisms. As such, they encapsulate cellular heterogeneity as is known
to exist (35). For example, cell 1 has a maximal h-channel conductance of 0.1063 pS/μm^2 whereas it is 0.0608

pS/µm² in cell 2. All parameter values can be found in Guet-McCreight and Skinner (33) and Sekulic et al. (27).

Throughout the paper we use both models so as to consider whether these differences would affect our results.

Synapse models

For the synapse model we use NEURON's built-in Exp2Syn function. The input populations to OLM cells include:

I-S3 cell inputs, GABAergic long-range projecting inputs from medial septum (MS), bistratified cell inputs (BIS), and

local PYR cell inputs [I-S3 cells & MS: see Chamberland et al. (13); BIS & PYR cells: see Leão et al. (20)]. In the

absence of specific spatial constraints, these inputs are distributed randomly across all dendritic compartments.

OLM cells also receive inputs locally from long-range projecting VIP+ cells in CA1 (18), though because these

cells are silent during theta rhythms, we did not include them in this study. In previous work, we had performed

optimizations across dendritic compartments so as to estimate the synaptic weights for each input type that

matched the experimentally recorded synaptic current amplitudes measured at the soma, also taking into

consideration measured synaptic kinetics [see Table 2 in Guet-McCreight and Skinner (33)]. In doing this, we

obtained increasing values with distance from soma (cell 1: G_{PYR} = 0.00020 to 0.00082 µS; G_{MS} = 0.00024 to

0.00132 µS; G_{I-S3} = 0.00018 to 0.00068 µS; G_{BIS} = 0.00021 to 0.00100 µS; cell 2: G_{PYR} = 0.00020 to 0.00120

µS; G_{MS} = 0.00024 to 0.00132 µS; G_{I-S3} = 0.00018 to 0.00120 µS; G_{BIS} = 0.00021 to 0.00100 µS).

Consideration of other inputs and further details can be found in Guet-McCreight and Skinner (33).

In vitro model explorations

In vitro setup. We use our models to consider an *in vitro* context as used by Tyan et al. (15) of hippocampal slice

recordings. In these experiments, although excitatory and inhibitory synapses were not specifically blocked during

intracellular OLM cell recordings, cells are relatively silent compared to *in vivo* recordings, and so we do not include

any additional synaptic bombardment (see **Fig. 1A**). To model this context, we apply various holding currents to the

somatic compartment and rhythmic synaptic input activations on dendritic compartments, which is analogous to the

optogenetic experiments performed by Tyan et al. (15). More specifically, we use a similar protocol in our

computational OLM cell models where we first depolarize the cell enough (cell 1: I_{inj} = 49 pA; cell 2: I_{inj} = 45 pA)

to attain a spike rate of 7.25 Hz. We then activate select input populations at different frequencies in our models.

The synaptic locations for the inputs are chosen randomly across the dendritic arbour of OLM cells (**Fig. 2**) using our

previously developed techniques (33). We look at a variety of different input schemes to gauge possible contributions

of different network components, but we mainly focus our explorations on the contributions from I-S3 cell inputs.

When PYR cell inputs are included together with inhibitory ones, they are activated with a delay that has been observed

experimentally (see section below).

Phase response curve (PRC) analyses. The control of OLM cells by I-S3 cells has been suggested to be due to

PIR mechanisms (15). That is, inhibitory inputs would activate biophysical currents such as I_H in OLM cells which

could lead to spiking. Considering this, we are in a position to take advantage of our detailed OLM cell models that

were developed in a "neuron-to-model matched" (NMM) fashion (27) regarding its morphology, passive properties,

and biophysical currents. These are currently the most biophysically, detailed OLM cell models and using them

serves as a biological proxy for exploring PIR mechanisms in a comprehensive fashion. We do this by turning to

phase response curve (PRC) analyses, which have been used to study morphologically detailed neuron models in the

past (36). If PIR mechanisms are in play, then one would expect to observe that I-S3 cells could cause phase

advances and not phase delays of OLM cell spiking.

To compute PRCs as shown in **Figs. 3B,C, 4A**, we compute the phase response ($\Delta\Phi$) for perturbations given at

different phases of the cycle. We obtain the interspike interval of the two spikes preceding the perturbation ($T0$)

and the interspike interval between the last spike preceding the perturbation and the first spike following the

perturbation ($T1$). We calculate the phase response as follows:

$$\Delta\Phi = \frac{T1 - T0}{T0} \times 100 \qquad (1)$$

In this sense, the phase response is normalized to 1 and converted to a percent. A negative value means a phase

advance, or a shortening of the interspike interval. A positive value means a phase delay, or a lengthening of the

interspike interval. We compute the phase responses where the zero perturbation is defined as the time when the spike

crosses a threshold of -20 mV during its upstroke.

To compute the change in biophysical currents (the same calculation used for each current type; see **Figs. 3D,F,**

4B,C) caused by the perturbation, we obtain the peak current amplitude generated in the period from the 2nd

last spike preceding the perturbation to the perturbation time ($I0$), and the peak current amplitude generated in the

period from the perturbation time to the 2nd spike following the perturbation ($I1$). Percent change in peak current is

calculated as follows:

$$\Delta I = \frac{I1 - I0}{I0} \times 100 \qquad (2)$$

As such, positive values indicate percent increases in peak current amplitude, and negative values indicate percent

decreases in peak current amplitude.

In vivo model exploration

***In vivo* setup**. It is known that the large amount of synaptic bombardment in the behaving animal *in vivo* changes

the response profile of individual neurons (37). In previous work using I-S3 cell models (32), we had

developed techniques for performing high-resolution parameter searches in parallel on the Neuroscience Gateway

(NSG) for high-performance computing (38) to find synaptic input parameter combinations (i.e. excitatory and

inhibitory numbers of synapses and spike rates) that could generate *in vivo*-like (IVL) states for the given cell

model. IVL states were characterized by metrics that ensured, for example, large enough fluctuations of

subthreshold membrane potential activities and interspike interval coefficient of variations. In Guet-McCreight

and Skinner (33), we used similar techniques to obtain IVL states for OLM cells. We found that IVL states

resulted in synaptic parameter values for cell 1 of 1268 excitatory synapses firing at 1.6 Hz and 1254 inhibitory

synapses firing at 8.7 Hz, and for cell 2, 1503 excitatory synapses firing at 1.5 Hz and 1532 inhibitory synapses

firing at 8 Hz, distributed throughout the dendritic tree.

Spiking resonance

Setup for *in vitro* spiking resonance. To examine spiking resonance in *in vitro* states, we generate 50 different

baseline spike rates (f_B's) in our OLM models by applying a range of holding currents from 30-152.5 pA (cell 1)

and 22-144.5 pA (cell 2). Note that plots of the frequency-current relationships of these models have been reported

previously (33). This elicits a range of f_B's from about 1-35 Hz in both models. Specifically, for cell 1, we obtain a

f_B range of 1.14-35.72 Hz, and a range of 0.95-34.80 Hz for cell 2, for the holding current ranges used. Each of these

models is then subjected to a range of inputs at different input frequencies (0.5 - 30 Hz) to determine spike resonance

frequencies.

Setup for *in vivo* spiking resonance. To examine spiking resonance in *in vivo* states, we also generate 50 different

f_B's in our OLM models. However, this is done by changing the random seed that controls placement of synapses and

presynaptic spike times rather than by changing the holding current since *in vivo* one is not injecting tonic current

into the soma. This generates different f_B's (across random seeds), though not with as wide of a range as changing the

holding current in the *in vitro* state. We obtain 50 f_B's by using 50 different random seeds. Specifically, for cell 1, we

obtain a f_B range of 11.82-15.83 Hz, and a range of 9.34-13.57 Hz for cell 2. Each of these models is then subjected

to a range of inputs at different input frequencies (0.5 - 30 Hz) to determine spike resonance frequencies.

Spiking resonance calculation. For each of the models, we compute the power spectral density (PSD) of the

spike train (1's for spikes and 0's for no spikes, using a -20mV threshold for spikes) without modulatory inputs

(i.e., at f_B). The PSD is computed in python using the welch function available as part of the scipy module:

`signal.welch(signal, fs=1/dt, scaling='density', nperseg=20000)`. We then apply a

series of different input frequencies (f_I) using different input populations, and the PSD is computed following each

input frequency.

The different input populations used are: 30 frequency-modulated I-S3 cell inputs, or 30 frequency-modulated PYR

cell inputs, or 30 frequency-modulated inputs from each input populations (i.e., I-S3, MS, BIS, and PYR cells). For

the latter, the start time delays from each other are based on their relative timing during theta rhythms (16, 39, 40).

Specifically, these start times (phases) relative to a 125 ms theta cycle are the following: I-S3, 45°; MS, 225°; BIS,

270°; PYR, 270 °. We note that the choice to use 30 I-S3 cell synapses is because at this number of synapses, I-S3 cell

synaptic activation can exhibit a sufficient level of synaptic control over OLM cell spiking at 5 Hz in both models (see

Fig. 2). The choice to use the same numbers of PYR cell, MS cell, and BIS cell synapses is simply to keep the numbers

equivalent in the absence of appropriate experimental data obtained in this context to constrain these numbers.

To gauge spike frequency resonance, we use a ratio to baseline measurement (i.e., 'baseline ratio' or δPSD), which

is computed by dividing the PSD at the f_I by the PSD at the f_I in the corresponding baseline trace (i.e. without the

modulatory inputs):

$$\delta PSD = \frac{PSD_{f_{I_{modulated}}}}{PSD_{f_{I_{baseline}}}} \tag{3}$$

In other words, it is a measurement of how much the PSD at the f_I changes relative to baseline once modulatory inputs

are added. A value of 1 would therefore indicate that there is no change in PSD, values less than 1 indicate a decrease

in PSD, and values greater than 1 indicate an increase in PSD. The magnitude of this metric therefore indicates the

effectiveness of synaptic inputs at entraining spiking to a prescribed input frequency. Spike resonance (f_R) is defined

as the f_I generating the largest baseline ratio.

In vivo theta rhythm exploration setup

We use our models to consider an *in vivo* context as found in (16). We run simulations using the full IVL scenarios +

8 Hz modulatory inputs from all input population (I-S3, MS, BIS, and PYR cells). As done in Luo et al. (16), we

add a small amount of noise in all of the theta-timed inputs, as this is both more realistic than having perfectly-timed

inputs, and enhances theta recruitment.

Results

Focusing on I-S3 cell inhibitory control over OLM cells, we perform detailed computational explorations of OLM cell models

considering *in vitro* and *in vivo* contexts. These explorations represent virtual network scenarios as the OLM cell is

either depolarized with holding current injections (*in vitro* state) or virtually embedded in a network where it is receiving

much synaptic bombardments (*in vivo* state). In **Fig. 1** we show an example of output in *in vitro* (A) and *in vivo* (B)

contexts along with the biophysical currents that underlie the outputs.

I-S3 cell control over OLM cell spiking – reproducing in vitro experimental observations

In vitro, optogenetic activation of CR+ cells, which includes I-S3 cells, can preferentially control the timing of

OLM cell spiking at 5 and 10 Hz, but not 1 and 20 Hz. Tyan et al. (15) considered that this was achieved through

PIR spiking mechanisms. However, because other inhibitory synapses and excitatory synapses were not blocked in

these experiments, it is unclear whether this is due to network effects or to intrinsic, biophysical OLM cell ion channel

properties that promote PIR spiking.

We first note that following negative long-square, high amplitude current injections, OLM cells

are capable of exhibiting rebound spikes (27). Our OLM cell models also exhibit this feature

(Cell 1 threshold: -220 pA; Cell threshold 2: -230 pA), primarily through activation of I_H and

subsequent activation of I_{Na} (**OSF Figure 1** on **https://osf.io/28r4s/**). However, it is

unclear if synaptic inhibition alone is sufficient to trigger this same phenomenon. As well, it is

unknown whether and how findings obtained *in vitro* would translate to *in vivo* contributions in a behaving animal.

I-S3 cell input populations alone cannot entrain OLM cell spiking as observed experimentally.   There
are several ways in which I-S3 cells could manifest their control over OLM cells. If it is primarily through PIR
spiking, this implies that inhibitory perturbations from I-S3 cells to OLM cells essentially speed up the onset of
an OLM cell spike – a phase advance due to intrinsic OLM cell properties such as I_H, I_{CaT}, and I_M that would
interact with the incoming inhibitory perturbations. There are other ways in which I-S3 cells could influence OLM
cells in the hippocampal circuitry. I-S3 cells could be controlling the timing of OLM cell spiking by suppressing
and delaying OLM cell spiking. This would require OLM cells to be spiking faster than the frequency at which
I-S3 cells are activated to exert their control. During the *in vitro* experimental recordings, the average spike rate of
the OLM cells is 7.3 Hz and OLM cells could be recruited to spike at 10 Hz photo-activation patterns. Thus,
simple suppression by I-S3 cells seems unlikely to be the case.

Let us start by examining how I-S3 cell synaptic inputs alone could control OLM cell output. Using optimized synaptic
parameters of I-S3 cell inputs to our detailed OLM cell models, we explore a range of different numbers of synapses
when I-S3 cells are activated at different frequencies as was done in Tyan et al. (15). An example of 30 I-S3 cell
synapses distributed on one of the OLM cell models is shown in **Fig. 2A**.

We apply I-S3 cell synaptic inputs alone with rhythmic spiking at 1 Hz, 5 Hz, 8 Hz, 10 Hz, and 20 Hz and using
0 to 60 synapses. The resulting output from one of the OLM cell models (cell 1) is shown in **Fig. 2A**. Output for
cell 2 is similar and is provided in **OSF Figure 2** on **https://osf.io/28r4s/**. In looking at the spike traces
across both cell 1 and cell 2, we observe that entrainment by the I-S3 cell synapses occurs primarily at 5 Hz when
there is a large enough number of frequency-modulated synapses. However, at higher frequencies, I-S3 cell inputs
only lead to spiking on every other cycle. For example, at 10 Hz stimulation with 30 I-S3 cell synapses, the OLM cell
models spike at 5 Hz. This is in stark contrast with the experimental results that show that OLM cells spike at 10 Hz
when receiving 10 Hz inhibition from CR+ cells (15). These modeling results show that I-S3 cell synaptic inputs
delay the spiking of OLM cells on each cycle, which is contrary to what one would expect if PIR mechanisms in
OLM cells were the main way by which I-S3 cells were exerting their control of OLM cell firing. We also tested the
inclusion of inhibitory input from other populations onto OLM cells [e.g., MS cell inputs, which were reported by
Chamberland et al. (13) to have comparatively larger IPSC amplitudes than I-S3 cell inputs], as well as clustering I-
S3 cell synapses on distal or proximal dendrites [i.e., since I-S3 cell IPSC amplitudes from Chamberland et al.
(13) suggest preferred distal localization of I-S3 cell synapses]. These manipulations did not lead to better spike
entrainment at 10 Hz frequencies (not shown). Further, these explorations suggest that any other purely
inhibitory pathway, e.g., a 'I-S3 to BIS to OLM cell' pathway, would also lead to spiking delays, and not be able
to reproduce the experimental observations of Tyan et al. (15). We note that IPSP magnitudes (**Fig. 2D**) are
comparable to those generated experimentally during wide-field optogenetic stimulation of CR+ cells [i.e., Fig
8C1-C4 in Tyan et al. (15)], likely because the I-S3 cell input conductances in our models were optimized to capture

the amplitudes of I-S3 to OLM cell connections reported by Chamberland et al. (13). However, this blown-up

trace (**Fig. 2D**) also highlights that the full measured amplitudes of the IPSPs are dependent on their timing

relative to the after-spike hyperpolarization portion of the trace, which can occlude the IPSP almost completely.

I-S3 cell inputs together with PYR cell inputs can entrain OLM cell spiking as observed experimentally.

To reproduce the *in vitro* experiments, it would seem to be the case that I-S3 cells do not simply suppress and delay

inhibitory cell spiking without PIR spiking. However, since excitatory synapses were not blocked in the *in vitro*

experiments, it is possible that photoactivation of I-S3 cells might have the general effect of disinhibiting PYR cells,

which could then further phase lock the spiking of OLM cells through recurrent excitation (i.e., a 'I-S3 to OLM to

PYR to OLM cell' pathway). There are additional ways in which this could occur. For example, it is possible that

inhibition of BIS cells by I-S3 cell activation could contribute towards disinhibiting PYR cells, which could

further augment recurrent excitation to OLM cells in an 'I-S3 to BIS to PYR to OLM cell' pathway.

Thus, since inhibitory inputs alone are not able to entrain OLM cell spiking in a manner that is consistent with the

in vitro experiments, we include PYR cell inputs in a disinhibitory manner to determine whether we could achieve

a match with experiment. As we are not explicitly modeling PYR cells, we include frequency-modulated PYR cell

synaptic inputs with a delay relative to inhibitory inputs. Also, we add only a few PYR cell synapses (i.e., enough

to generate spiking) on the OLM cell models to determine whether we could obtain the level of spike entrainment

previously seen in experiment (15). This is shown in **Fig. 2B** for cell 1 (cell 2 in **OSF Figure 2** on

https://osf.io/28r4s/) where it can be seen that one-to-one entrainment is possible at 5 and 10 Hz as

observed experimentally, but now also at 20 Hz, which is unlike what is seen experimentally in Tyan et al. (15).

The inclusion of frequency-modulated inputs from both I-S3 and PYR cells is 'virtual', meaning that we are

not explicitly modelling I-S3 and PYR cells in a circuit configuration with the OLM cell models. That is, in our

virtual network simulations here, we include PYR cells inputs at the various frequencies, assuming that they

reliably spike at the given input frequency, whether it be 5 or 20 Hz and so on, as there is no explicit modeling

of tri-synaptic connectivity between I-S3 cells, OLM cells, and PYR cells. While this is a clear limitation, the

alternative to simulate full microcircuits brings about many other aspects to consider and introduces other

limitations and caveats (see Discussion). That being said, a possible interpretation of our simulation results is

that, experimentally, photo-activation of CR+ cells at 20 Hz may not lead to disinhibited PYR cell spiking at 20

Hz, as modelled here with our virtual networks - at least not enough to excite OLM cells to spike at these higher

frequencies. Additionally, wide-field optogenetic stimulation of CR+ cells in Tyan et al. (15) was often capable

of generating 2-3 OLM cell spikes. This effect was not captured when scaling up the number of I-S3 cell inputs

alone (**Fig. 2A**) or with a minimal number of PYR cell inputs (**Fig. 2B**). In scaling up the number of PYR cell

inputs on its own, however, we find that this effect can be replicated (**Fig. 2C & F**). As such, our simulations

suggest that wide-field activation of CR+ cells, that includes I-S3 cells, may disinhibit a large enough number of

PYR cells to cause a robust recruitment of OLM cells to spike.

Insights from PRC analyses with biophysical correlates

PRCs are used in studies of oscillatory systems such as neuronal spiking (41, 42) to indicate the change in the cycle
period as a function of the phase at which a perturbation is received, thus predicting its response to rhythmic
inputs - that is, its entrainment at various frequencies. Thus, we here use PRCs to uncover whether there are
any scenarios under which I-S3 cells could cause phase advances of OLM cell spiking and thus indicative of PIR
mechanisms. In the previous section where we focused on replicating *in vitro* experiments (15), we did not find
support for PIR mechanisms contributing to I-S3 cell control of OLM cell firing.

Inhibitory perturbations rarely elicit PIR spiking in OLM cell models. In **Fig. 3**, we illustrate theoretical
PRC explorations. **Fig. 3A** shows the spiking frequency of cell 1 for a given amount of somatically injected tonic
current. At 30 pA injected current, as indicated by the dashed line, the cell fires at about 1 Hz. The PRCs generated
for the cell firing at this frequency are given in **Fig. 3B & 3C**, in response to different positive and negative 3ms
square current pulses at the soma, and different numbers of I-S3 cell synapses spread throughout the dendrites,
respectively. Both negative current pulses with magnitudes up to -500 pA (**Fig. 3B**) and up to 60 inhibitory synaptic
inputs (**Fig. 3C**) could generate phase advances at early phases of the interspike interval. In comparison, much
smaller amplitude positive current pulses caused large phase advances at later phases of the interspike interval (**Fig.**
3B), and similar PRC shapes could be obtained with PYR cell synapses, but with conductance values that are much
smaller than optimized PYR synapse values (not shown). Increasing the inhibitory perturbation past 30 synapses had
little effect on further enhancing phase advances (**Fig. 3C**) but was accompanied with a continuous increase in I_H response
(**Fig. 3D**). The dashed lines in **Fig. 3C** show that cell 1 can either show a phase advance (yellow) or a phase delay
(red) of the subsequent spike if the cell is perturbed at the given phases indicated along the x-axis.

In **Fig. 3E** we show an overlay of the model voltage traces without perturbation (dashed grey), when there is a phase
advance (yellow) or a phase delay (red) using a perturbation of 30 inhibitory synapses. As can be seen in **Fig. 3C**,
perturbations for this scenario (OLM cell spiking around 1 Hz, close to its rheobase) mostly lead to phase delays,
and any possible phase advances are small. **Fig. 3F** shows the changes in all of the biophysical currents in cell 1
due to the phase-dependent perturbations from 30 inhibitory synapses. It is clear that I_H is the main contributor
when there is a phase advance given its change relative to all of the other biophysical currents. This change is as an
increase which makes sense for I_H contributing to producing a phase advance since it is an inward current. I_{CaT}, and
I_{Kdrs} also are strong contributors here, but showing decreases for either phase advance or delay scenarios.

We compute the PRCs at four different baseline spike rates (rheobase, 4 Hz, 7.25 Hz, 15 Hz) in the OLM cell models
(see **Fig. 4A**) using a perturbation of 30 inhibitory synapses. From this, it is clear that phase delays are almost always
present in response to inhibitory perturbations, with the smallest phase delay occurring when the perturbation is

given during the spike refractory period, and the largest phase delay occurring when the perturbation is just
before spike threshold - this aspect is visualized in the voltage traces shown in **Fig. 3E**. Only at the lowest
baseline spike rate (rheobase) can phase advances (negative values) occur when the perturbation is near 40% of
the inter-spike interval, and these small phase advancements only occur with cell 1, and not with cell 2. This
exposes the biophysical heterogeneity in these two OLM cell models, each of which were developed using an NMM
approach (see Methods). In particular, given that I_H constitutes the main current increase with the perturbation, and
we know that the h-channel conductance is larger in cell 1 relative to cell 2, we can say that this is what underlies
the small phase advancement that occurs in cell 1. **Fig. 4B** shows the I_H changes for the four different baseline spike
rates and one can see the larger I_H for cell 1 as compared to cell 2. It is important to note that the various channel
conductances in the cell 1 and cell 2 OLM cell models were optimized to the specific electrophysiological data, and
not arbitrarily set to different values. However, to verify that variations in I_H in cell 1 or 2 models do not affect these
results, we adjust the h-channel conductance by $\pm 25\%$ and find little change in the PRCs (**OSF Figure 3** on
https://osf.io/28r4s/). Unlike I_H, most currents exhibited perturbation phase invariance at rheobase (**Fig.**
3F), and stronger current changes near spike threshold (i.e., 100% phase) when the cells were spiking at higher
frequencies, such as with I_M for cell 1 and cell 2 (**Fig. 4C** and see **OSF Figure 4** on **https://osf.io/28r4s/**
for all current changes).

In examining all of perturbation phase-dependent amplitude shifts of the different currents due to the various ion
channel mechanisms (shown in **OSF Figure 4** on **https://osf.io/28r4s/**), we observe the following: We
obtain phase-dependent increases in I_H (cyan), and I_{CaL} (yellow), and decreases in I_M (purple), which should all
enhance the likelihood of PIR spiking (i.e. more inward current and less outward current). We note that maximal I_{CaT}
(grey) is decreased following the inhibitory perturbation, regardless of the perturbation phase, but is steadily decreased
less at later phases of the interspike interval for base firing rates of 4 and 7.25 Hz. A phase-dependent increase, albeit
a small one, in I_{K_A} (blue), however, would counteract rebound firing since I_{K_A} is known to suppress PIR spiking (29).
It should be noted that contributions from I_{K_A} are large in these models, which could partially account for the lack of
inhibitory perturbation-dependent phase advances. Other currents (I_{Kdrf}: orange, I_{Kdrs}: green, I_{KCa}: red, I_{Na}: pink, I_L:
brown) do not show appreciable inhibitory perturbation phase-dependent changes in maximal magnitude. Comparing
relative changes across the baseline firing rates and perturbation-dependent changes, enhancements in contributions
from I_{K_A}, I_{Kdrf}, I_{Kdrs} and I_{KCa} all contribute towards the prevention of phase advances from occurring more broadly
across higher spike rates. Since I_{K_A}'s phase-dependence near rheobase is flat compared to its phase-dependence
at higher spike rates, this may also account for why some phase advances are permitted near rheobase. Though
there is a difference in PRCs across the two models, mainly at the rheobase baseline spike rate, the ion channel
current responses are consistent for both OLM cell models. Altogether, these results suggest that inhibitory inputs
alone, via I-S3 cells, are unlikely to be sufficient to elicit PIR mechanisms in OLM cells. Rather, the modeling
predicts that I-S3 cell activation suppresses OLM cell spiking (i.e., causes phase delays).

I-S3 cell inhibitory inputs and spike resonance in OLM cells

In vitro spike resonance at theta frequencies occurs for excitatory but not inhibitory inputs. From our PRC

analyses and simulations of the *in vitro* experiments of Tyan et al. (15), it seems clear that I-S3 cells likely do not

control OLM cell firing through PIR mechanisms since small phase advances were obtained in a very limited

fashion (rheobase for cell 1 only). We have so far limited our explorations to a few input frequencies and

baseline spike rates. From these results it is clear that inhibitory modulation depends on the baseline firing rate of

the OLM cell (f_B), as well as the frequency of the incoming frequency-modulated synaptic inputs (f_I), though the

precise relationship between the two is unclear. Let us now explore a wide range of baseline firing rates in OLM

cells and apply a wide range of frequency-modulated inputs to determine the scenarios under which preferred firing

(i.e., spike resonance) could occur, and whether it includes theta (3-12 Hz) frequencies. We do this by

incrementally changing the holding current to elicit different f_B, while at the same time submitting the model at

each holding current to a barrage of different input frequencies. We note that this range of baseline spike rates

includes the range of spike rates that have been reported for OLM cells *in vivo* across different behavioral and

network states (43, 44). In this way we can establish a 'ground-truth' regarding preferred output responses of

OLM cells in the face of various input frequencies, i.e., an OLM cell spiking resonance, given different baseline

spike rates in an *in vitro* context.

As noted earlier, the *in vitro* context simply means that the OLM cells are not bombarded with excitatory and

inhibitory synaptic inputs. In this context, we apply the various holding currents to the somatic compartment (see **Fig.**

1A), and rhythmic synaptic input activations are applied to dendritic compartments. In **Fig 5**, we show the spike

resonant frequencies (f_R) of cell 1 when receiving inhibitory inputs from I-S3 cells (**Fig. 5A-D**) or excitatory inputs

from PYR cells (**Fig. 5E-H**). Results from inputs across all populations (i.e., including MS and BIS inputs) are

shown in **OSF figure 5** on **https://osf.io/28r4s/**, as well as for both cell 1 and cell 2. When receiving only

I-S3 cell inhibitory inputs (**Fig. 5A**), f_R is largely dependent on f_B (**Fig. 5B**; cell 1 f_R: 15.37 ± 8.76 Hz; cell 2 f_R:

15.23 ± 8.88 Hz). That is, if the OLM cell is firing at a higher rate, then f_R is also higher. This can be seen more

clearly in **Fig. 5C**, which also highlights that f_R can also occasionally be larger than f_B, despite having spike rates

at f_R that are always suppressed compared to f_B (**Fig. 5D**; i.e., consistent with the phase delays due to inhibitory

perturbations reported in previous sections). One explanation for this is that when the baseline ratio is enhanced in

those cases, the spikes that are being suppressed are spikes that occasionally fall in phase with the modulatory inputs.

In this sense, the modulatory inhibitory inputs are keeping the spiking entrained through suppression. This can happen

regardless of whether f_R is greater than or lesser than f_B, and depends on how the intrinsic spike train aligns with the

modulatory input spike trains.

Overall, these results make sense given the findings from the previous section showing that I-S3 cell synapses alone

rarely lead to phase advances. Specifically, recruitment of OLM cell spiking to certain inhibitory input frequencies is

largely baseline spike rate-dependent because inhibitory perturbations mostly just cause phase delays. For example, if
the baseline spike rate is much smaller than the input frequency, inhibitory perturbations will only cause suppression
of spikes, and the PSD at the input frequency is more likely to drop. Likewise, if the baseline spike rate is much larger
than the input frequency, then inhibitory perturbations will have minimal effects on spiking, as they may only suppress
a small fraction of spikes. Thus, the spike resonant frequency is largest when the baseline spike frequency is near the
inhibitory input frequency. These tests were also performed using MS inputs alone (not shown), which have larger
IPSC amplitudes, and similar results are obtained.

In contrast, OLM cell spike resonance due to excitatory PYR cell inputs (**Fig. 5E**) is consistently in the 2-15 Hz input
frequency range, regardless of baseline spike frequency (**Fig. 5F-G**; cell 1 f_R: 7.8 ± 4.28 Hz; cell 2 f_R: 4.38 ± 2.48
Hz). However, the magnitude of the baseline ratio at f_R is dependent on baseline spike frequency, with larger baseline
spike frequencies having smaller baseline ratios at the f_R (**Fig. 5F**). In all cases the presence of excitatory inputs causes
increases in spike rates, which is consistent with phase advances (**Fig. 5H**). Similar findings are seen when including
all input populations (see **OSF Figure 5** on **https://osf.io/28r4s/**), suggesting that any theta frequency (3-12
Hz) spike resonance *in vitro* is largely dictated by excitatory inputs when they are present and not by inhibitory inputs
alone. We note that using a lower number of PYR cell synapses could affect these results, but not likely in a drastic
way given that OLM cell entrainment is strong even when using a small number of PYR cell synapses (see **Fig. 2C**).

In vivo, theta frequency spike resonance can occur with inhibitory but not excitatory inputs. We have
previously simulated synaptic bombardment conditions on OLM cells to generate *in vivo*-like (IVL) states. This
includes a random spread of synaptic locations from different input populations leading to irregular firing - an
illustration is shown in **Fig. 1B**. We now investigate I-S3 cell control over OLM cells during IVL states. Since
we have already investigated I-S3 cell control over OLM cells in an *in vitro* context, we are able to compare whether
or not the same level of control over OLM cell spiking might apply *in vivo*. We use different random seeds to
obtain a range of f_B's, and we note that the range of frequencies is not as expansive as *in vitro* as the level of control is
limited (i.e., we are not simply injecting tonic somatic currents).

We perform the same spike resonance tests as in **Fig. 5**, but now under this IVL state (Cell 1: **Fig. 6A,E**). We see
across all input population types (I-S3 cells and PYR cells) that there is a rightward shift in the resonant frequency
distributions, relative to *in vitro*, towards higher values where more if not all f_R values are larger than the f_B values
(**Fig. 6B,F**; cell 1 $f_{R, I-S3}$: 15.92 ± 5.84 Hz; cell 2 $f_{R, I-S3}$: 18.07 ± 7.74 Hz; cell 1 $f_{R, PYR}$: 27.42 ± 3.43 Hz; cell 2
$f_{R, PYR}$: 25.6 ± 5.31 Hz). That is, even when considering the smaller range of the baseline spike rates, the resonant
frequencies are shifted towards higher input frequency ranges (**Fig. 6C,G**). We note that, as in the *in vitro* case, there
is no change in the suppressing effects of inhibitory inputs (i.e., PIR spiking remains absent) or the excitable effects of
PYR cell inputs on spike rates (**Fig. 6D,H**). The shift towards higher f_R values than in the *in vitro* case (i.e., **Fig. 5C,G**
vs. **Fig. 6C,G**) is possibly because of different aspects associated with the IVL state, including irregular spiking
patterns (i.e., spiking is no longer periodic; **Fig. 1B**), as well as a reduction in sensitivity due to a decrease in input

resistance. In other words, irregular spike patterns at baseline can allow larger variabilities in f_R values depending on
how spike times align with incoming modulatory excitatory and inhibitory inputs. As well, decreased input
resistance can increase the required magnitude of excitatory and inhibitory currents needed to modulate the model
cell firings. This can be brought about by higher input frequency modulation, thus explaining the increased f_R
values. We note again that cell 2 exhibited similar f_R profiles to cell 1, and similar to the *in vitro* case, inclusion of
all input populations as modulatory inputs exhibited similar f_R profiles to having just PYR cell inputs (**OSF Figure 6**
on **https://osf.io/28r4s/**).

When considering the amplitudes of the baseline ratios, the IVL magnitudes (**Fig. 6B,F**) are considerably smaller than
the *in vitro* baseline ratios (**Fig. 5B,F**). This observation could be due to an increase in currents in the model once it is
put into an IVL state, which generates a lower input resistance and decreased sensitivity to rhythmically-timed inputs.
In other words, higher input frequencies will be necessary to elicit spike resonance since there is a decrease in
sensitivity. We note that we have previously established that the addition of synaptic bombardment leads to decreases
in sensitivity to additional inputs (32), so it is not surprising that spiking resonance in OLM cells will be different
during IVL states relative to *in vitro* states.

The results between the two OLM cell models are qualitatively similar, despite differences in morphologies and
intrinsic parameters. One potential reason for this is the presence of dendritic I_H, and dendritic Na^+ and K^+ channels
in both models, which would promote similar integration of synaptic inputs and action potential propagation and thus
generate qualitatively similar results across both models. As well, our simulation results are quite similar between
using I-S3 cell inputs or MS inputs (not shown), the latter of which have larger IPSC amplitudes (13). Indeed, there
would not be a strict differentiation of incoming inhibitory inputs as modelled in these virtual networks.

Simulating I-S3 cell inputs to OLM cells in theta-run epochs during behaviour

In previous work, we found that I-S3 cells are activated with a delay relative to theta-run epochs and spike during the
rising to peak phases of theta rhythms (16). As well, it was predicted from modelling work that the timing of this
phasic preference would be modulated by inputs from entorhinal cortex (EC; rising phase) and CA3 (peak phase).
Thus, we investigate the effects of a ramp-up of I-S3 cell activation (i.e., simulating a delay in I-S3 cell activation)
on a per theta cycle basis during an ongoing theta rhythm (**Fig. 7**, which shows results for cell 1). We run
simulations using the full IVL scenarios + 8 Hz modulatory inputs from all input population (I-S3, MS, BIS, and
PYR cells). As schematized in **Fig. 7A**, starting at 2 s into the simulations until the end of the simulation (10 s),
we add 7 I-S3 cell input spikes per cycle, which essentially ramps up the inhibition from I-S3 cells gradually. To
consider possible shifts in balances of inputs from CA3 and EC, which can present a possible dis-inhibitory circuitry
for switching between sensory inputs and memory consolidation inputs, we also explore shifts in the timing of I-S3
cell inputs alone (**Fig. 7A**) or together with a shift in the timing of PYR cell inputs (**Fig. 7E**). Note that synaptic
location sites are chosen randomly for each addition, as described previously (32). The rationale behind these

simulations is that the majority of I-S3 cell activation occurs near the ends of theta-run epochs, with a delay

relative to the timing of activation of other neuron types, and so we want to see the effect of a ramp-up of I-S3 cell

inputs to OLM cell spiking during a behavioural context with ongoing theta rhythms. Here we use five different

random seeds for IVL synapses, where IVL synapse locations and spike times are re-randomized (e.g., in **Fig. 1B**).

Ramp-up of I-S3 cell inputs during theta sharpens and modulates OLM cell recruitment. In the

simulations where we shift the timing of I-S3 cell inputs alone, the theta-timing of the OLM cell models do not

change across any of the conditions (i.e., no ramp-up - black, stronger EC inputs - red, even EC/CA3 inputs -

green, or stronger CA3 inputs - blue; **Fig. 7B**). We do see a moderate decrease in the spike rate (**Fig. 7C**), and an

increase in the 8 Hz power in the PSD (**Fig. 7D**) across all conditions compared to baseline. More specifically, the

enhancement in the PSD is strongest when the timing of the I-S3 cell inputs is shifted towards the peak, corresponding

with stronger inputs from CA3 (**Fig. 7D**, inset). This appears to be because inhibition occurring following the peak of

theta is the most out-of-phase with the trough of theta, which is when the OLM cell models spike, due to excitation

from local PYR cells inputs (**Fig. 7B**). Moreover, this also corresponds to the time at which the I-S3 cell inputs will,

on average, be most out-of-phase with the OLM cell spike refractory period, which can allow a stronger I_H-mediated

response to inhibition (**Fig. 3**). As such, I-S3 cell inputs alone can sharpen OLM cell recruitment at theta frequencies

by suppressing spikes that are out-of-phase with the theta-timing of OLM cell spiking. We note that when tested across

five different random seeds, these same results are generated consistently (i.e., the effect of ramped up I-S3 cell inputs

causing spike suppression, and the strongest 8 Hz power when modulated by CA3).

In the simulations where we shift the timing of I-S3 cell inputs together with PYR cell inputs (**Fig. 7E**), the theta-timing

of the OLM cell models are shifted by the same phasic amount (**Fig. 7F**). The interpretation in these simulations is

that a shift in the timing of I-S3 cells due to inputs from either EC or CA3 will shift the dis-inhibitory window for

PYR cells, and as such, their phasic timing will shift by the same amount. In these simulations we show that this

network effect would also shift the timing of OLM cell spiking. Moreover, since the out-of-phase timing is relative to

the phasic timing of when the OLM cell models are spiking, we see a similar PSD magnitude at 8 Hz regardless of

the direction in which I-S3 and PYR cell inputs are shifted (**Fig. 7H**). As well, the spike rates are again moderately

decreased when compared to baseline (**Fig. 7G**). Again, when tested across five different random seeds, the same

results were generated consistently, and similar results were obtained when testing cell 2 (**OSF Figure 7** on

https://osf.io/28r4s/).

Discussion

Mapping identified inhibitory cell types to function and behaviour is a challenging endeavour. To address this, we

took advantage of our previous studies interfacing experiment with modeling work of inhibitory cell types in the

hippocampus. We focused on how specialized interneuron-selective cells, I-S3 cells, might exert their influence over

OLM cells, which have a demonstrated gating control between incoming 'sensory' and ongoing 'memory' information

flow. We used two OLM cell models with parameters fit according to experimental data obtained from the same cell,

populations of synapses with parameters specific to cell types that synapse onto OLM cells, PRC theoretical

analyses, knowledge of I-S3 cell firing *in vivo*, as well as a methodology for generating *in vivo*-like states.

How do I-S3 cells control OLM cell spiking in vivo?

From the *in vitro* experimental observation of Tyan et al. (15) in which it was suggested that PIR mechanisms

contributed to control of OLM cell spiking by I-S3 cell inhibitory inputs, we considered other possibilities. In

simulated *in vitro* scenarios, we found that I-S3 cell-mediated disinhibition of pyramidal cells seemed most likely

to be the way in which I-S3 cells exert their influence on OLM cell spike timing. I-S3 cell-mediated suppression

of OLM cells would be part of enabling this contribution, but *not* PIR mechanisms, and BIS cells could also

contribute in these interactions. Thus, based on our explorations, we predict that the influence that I-S3 cells would

have on OLM cells would be through spike suppression followed by enhanced excitation of OLM cells due to PYR

cell disinhibition.

Interestingly, when simulating *in vivo* states in OLM cells, only inhibitory inputs, including those from I-S3 cells,

could lead to a spike resonance at theta frequencies. This was not the case during simulation of *in vitro* states

though - there it was excitatory and not inhibitory inputs that could lead to theta frequency spike resonances. In

previous modeling work, it was found that OLM cell spike resonance with inhibitory inputs in simplified, simulated

in vivo states could occur at high or low theta frequencies depending on whether dendritic h-channels were present

or not, respectively (31). Here, we used our NMM OLM cell models that were shown to necessarily have dendritic

h-channels (27), a different metric for gauging spike resonance (i.e., the maximal baseline ratio), a series of different

baseline spike rates [i.e., instead of just 2.5 Hz, as in Sekulić and Skinner (31)], and synaptic parameters

estimated from actual input populations measured experimentally. Based on the results here, we predict that theta

frequency spike resonance could occur in OLM cells *in vivo* with incoming inhibitory I-S3 cell inputs, as well as

from other rhythmic inhibitory inputs like MS and BIS cell inputs.

From previous work, we know that I-S3 cells exhibit a delay relative to theta-run epochs, with a phasic preference

towards the rising/peak phases of theta (16). I-S3 cell activation patterns during running and immobility are also

highly variable and even PYR cell-like when compared to other CA1 interneuron types (45), possibly due to mixed

modulatory excitation from both CA3 and EC (16), which have delayed activation patterns relative to one another

(46). Simulating a ramp-up of I-S3 cell spiking in our virtual networks, we found that we could obtain a

sharpening of the timing of OLM cells during theta rhythms, which could contribute to shaping pyramidal cell place

fields (47). In fact, silencing SOM+ cells increases PYR cell burst spiking during place field traversals in awake

mice (47), and also un-gates synaptic activation along pyramidal cell apical dendrites (48). In line with this,

optogenetic silencing of VIP+ cells, which promotes activation of SOM+ cells, dampens both the reshaping of

PYR cell place fields and learning of reward site locations (49). Coincidentally, I-S3 cell activation *in vivo* is

significantly less correlated with BIS cell activation than it is with basket cell activation, and even less correlated

with OLM cell activation, indicating a suppression of BIS cells and an even heavier suppression of OLM cells when

I-S3 cells are activated (45). Previous modeling has also shown that OLM cells, via network pathways that

include BIS cells, were key to LFP signal robustness of ongoing intrinsic theta rhythms (50). Thus, I-S3 cell

contributions, via OLM and BIS cells, could be essential for the existence of robust theta rhythms.

Based on all of this, our proposition for I-S3 cell contributions is illustrated in **Fig**. **8** in a series of steps. We predict that

I-S3 cells cause phase delays in OLM cell spiking by spike suppression, as opposed to PIR-mediated phase advances

(step 1). This would lead to PYR cell spiking due to disinhibition at a particular phase of theta rhythms (step 2), which

would subsequently dictate the timing of the OLM cell spiking through excitation (step 3), leading to re-gating of

inputs at distal dendrites (step 4).

Experimental Investigations

The predictions from our computational studies lead to several suggestions for experimental investigation. One precise

prediction is that 30 I-S3 cell synapses were necessary to elicit inhibitory control over OLM cell spiking. If the

variance-mean analysis estimation of three release sites by Tyan et al. (15) is correct, this would imply activation of at

least 10 presynaptic I-S3 cells. Results from Tyan et al. (15) can also be re-tested by performing frequency-modulated

optogenetic stimulation of CR+ cells while recording from OLM cells in the presence of excitatory synaptic blockers.

If OLM cell frequency-modulated spiking control is then lost, the interpretation would be that the frequency

modulation was due to feedforward disinhibition-driven excitation onto OLM cells. Additionally, instead of recording

from OLM cells during these tests, one could record from PYR cells to verify frequencies at which I-S3 cell

activation might lead to disinhibited spiking in PYR cells, if at all.

Another suggestion would be to perform closed-loop feedback stimulation with the theta-filtered LFP as has been done

previously (21), but with photo-activation of CR+ cells at specific phases of theta. One could stimulate CR+ cells at

phases closer to the peak versus the trough of theta, to assess whether the phasic timing of CR+ cells has an effect

on the encoding and retrieval of information during a learning test such as the T-maze task. Our results predict an

un-gating of pyramidal cells during theta phases that follow the activation of CR+ cells. On average (but with a large

variance), I-S3 cells spike near the rising to peak phases of theta (16) which follows excitation from EC and precedes

excitation from CA3. As such, if stimulated to spike at earlier phases, integration of inputs from EC by pyramidal

cells could be favoured to promote sensory encoding. Likewise, if stimulated to spike at later phases, integration of

inputs from CA3 by pyramidal cells could be favoured to promote retrieval of information.

Further, it is now clear that the location of OLM cells along the dorsoventral axis of the hippocampus matters in terms

of whether one records high or low theta rhythms (22, 51, 52), possibly due to biophysical and connectivity differences

with OLM cells in these different locations. As already noted above, modeling studies have suggested that whether

h-channels are present or not in the dendrites of OLM cells could 'control' the theta frequency spike resonance. Here,

we used OLM cell models with dendritic h-channels developed from intermediate CA1 mouse hippocampal data (27),

and our PRC explorations indicated that current changes with inhibitory perturbations were mostly due to h-channels.

Theoretical and Modeling Considerations

Our use of theoretical PRCs here is of a pulse-coupled type (53) where we used excitatory and inhibitory synaptic inputs to

consider phase-locked entrainment of OLM cell models. PRCs could also be used to predict synchronous output, and for example,

Type II PRCs imply synchronization with excitation (54, 55). We focused on the ability of our model OLM cells to exhibit

phase advances or delays using inhibitory perturbations for I-S3 cell input considerations. Other studies have used

synaptic perturbations with PRCs and with varying levels of neuron model complexity (36, 56).

Here, we have detailed multi-compartment models with a full suite of biophysical channel types, and since we

wanted to situate our explorations in a realistic biological setting, our inhibitory perturbations took the form of

dendritically distributed inhibitory synaptic inputs on the OLM cell models with synaptic features estimated from

experiment. These PRCs serve as an approximation for how OLM cells would respond to repeated 'periodic

forcing' (41), and they show that incoming inhibitory inputs mostly serve to slow down the firing of OLM cells.

Moreover, by also examining the underlying changes in the biophysical currents of OLM cells, we found that h-

channel changes dominate, but they are not sufficient to lead a phase advance with PIR mechanisms rather than a phase

delay. Previous work has highlighted that T-type calcium and persistent sodium channels in deep cerebellar nucleus cell

models can also underlie PIR spiking following long square negative current injections (57). In slight contrast, PIR

spiking following long square negative current injections in our OLM cell models (OSF Figure 1) were initiated

primarily by h-currents and then shaped by sodium currents, followed by T-type calcium currents, as well as several

outward currents.

Another interesting aspect to consider are the more complex dynamical regime changes that occur *in vivo* during

different awake and sleep cycles as well as behavioral states. While we have modelled *in vivo* states quite simply as

high levels of AMPA-mediated synaptic bombardment, more complex neuromodulatory phenomena are likely to be

governing state changes more closely associated with behavior. For example, experimental cholinergic modulation

has previously been shown to switch the PRC in L2/3 pyramidal neurons from type II to type I (58), which would

have synchronization consequences, and cholinergic inputs are known to be cell-specific (59). Other types of inputs

to OLM cells have been reported previously, including serotonergic receptors (60), metabotropic glutamate

receptors (61, 62), cholinergic receptors (63–65), additional nuances in NMDA/AMPA/Kainate receptors (66, 67),

and TRPV1 receptors (68). However, in the absence of particular constraints, we opted to not include them at this

time. It is unclear if a cholinergic-associated down-regulation of M-type current in our OLM cell models would

enhance the small inhibitory synapse-driven phase advances that we report in our *in vitro* PRC analyses. However,

given the phase invariance of non-h-type currents to inhibitory perturbations during rheobase spiking, we would not

anticipate a large effect. Certainly, a more complex picture may exist in the *in vivo* state however.

To allow a focus on cellular details, we carried out virtual network explorations using detailed OLM cell models.

Because we used models developed using an NMM approach, where a select number of neuronal model fits were

closely matched with experimental recordings (27), we did not explore a full range of parameters to exhaustively

assess the robustness of our findings. This can be done if we used the populations of 'acceptable' models that are

generated for each optimization (27), but would require a considerable amount of computing resources to fully

explore. The use of a virtual network allowed us to directly compare and translate *in vitro* aspects to *in vivo* to come

up with our proposed specialized contribution of I-S3 cells via OLM cell control. Since *in vivo* recordings of

specialized cell types is highly challenging, computational studies to develop hypotheses, predict and guide

experimental studies are strongly needed. However, a virtual network is limited as it does not directly model all

of the interacting network effects that can occur consequentially as a result of changes in the activation of the cell

type of interest. At the same time, the large variety of neuron types and subclass interactions makes the

consideration of interacting effects incredibly complex. Despite these interactions not being modelled explicitly,

our approach using the detailed, biophysical OLM cell models offers the benefit of being able to dissect out putative

mechanisms without having to build a full-blown circuit model initially, and has led to explicit suggestions for

experimental investigation. Further, we note that recent computational (69, 70) and experimental (45) studies have

begun to disentangle the co-activity dynamics of disinhibitory neurons with different neuron types in relation to

behaviour. We would also like to note that our interpretations are in line with previous network modelling showing

that VIP/SOM connectivity is sufficient for switching circuit activity between two processing modes where synaptic

inputs in pyramidal cells are either integrated or suppressed (69, 71, 72).

While the use of virtual networks has limitations, building and using full microcircuit models requires other

considerations with subsequent limitations. Two CA1 microcircuit models by Turi et al (49) and Bezaire et al.

(39) have been built. The Turi et al model (49) consists of 150 cells that includes PYR cells and 6 types of

interneurons including I-S3 cells. It is supportive of a disinhibitory role for I-S3 cells in encoding reward

locations, and indeed, the disinhibitory effects of VIP+ cell activation on PYR cell spiking has been broadly

reported in the literature (17). The Bezaire et al model (39) is a full-scale CA1 microcircuit model that consists

of 338,740 cells including 8 types of interneurons, and shows the importance of some cell types over others in

the production of theta rhythms. However, I-S3 cell types were not part of the Bezaire et al model. While both

microcircuit models include OLM cells, the OLM cell models within them are not detailed in morphology (4

compartments) or biophysical currents relative to the developed OLM cell models used in the work here. Of

particular note, no h-channels were included in the dendrites of the OLM cell models used in these
microcircuit models. These are all reasonable limitations given the respective goals of these microcircuit
models. In fact, there is an explosion of parameters at the circuit level (e.g., different cell types, connectivity
parameters & strengths, network size, etc.) that usually necessitates a reduction in the resolution of analysis at the
single cell level (e.g., dendritic compartments, ion channel types, etc.). A circuit model would also require a
further expansion of high-performance computing resources towards simulating the full circuit model rather than
focusing on the single-cell details and parameter spaces that we have focused on in this work. While our OLM cell
models were developed carefully using an approach that directly matched the model development with the same
biological cell (27), models at this level of detail do not currently exist for the many other interneuron types in the
CA1. The huge endeavour of achieving this level of detailed models for many interneuron types in a full-scale CA1
microcircuit has been undertaken (73), although I-S3 cells have not yet been included.

Summary & Conclusions

This work provides a deep understanding into how I-S3 cells could contribute to circuit function *in vivo*. This was
accomplished by studying inputs to detailed OLM cells models and dissecting out how I-S3 cell inputs may
contribute. By analyzing these factors carefully in biophysical computational models, we were able to generate
predictions into the mechanisms through which I-S3 cells can effectively control OLM cell spiking and precisely
coordinate modulation pathways. We also note that our interpretations are in line with previous network modelling
showing that VIP/SOM connectivity is sufficient for switching circuit activity between two processing modes where
synaptic inputs in pyramidal cells are either integrated or suppressed (69, 71, 72).

References
697

1.   **Tremblay R**, **Lee S**, **Rudy B**. GABAergic Interneurons in the Neocortex: From Cellular Properties
to Circuits. *Neuron* 91: 260–292, 2016. doi: 10.1016/j.neuron.2016.06.033.
2.   **Bezaire MJ**, **Soltesz I**. Quantitative assessment of CA1 local circuits: Knowledge base for
interneuron-pyramidal cell connectivity. *Hippocampus* 23: 751–785, 2013. doi: 10.1002/hipo.22141.
3.   **Cembrowski MS**, **Spruston N**. Heterogeneity within classical cell types is the rule: lessons from
hippocampal pyramidal neurons. *Nature Reviews Neuroscience* 20: 193, 2019. doi: 10.1038/s41583-019-
0125-5.
4.   **Klausberger T**, **Somogyi P**. Neuronal Diversity and Temporal Dynamics: The Unity of
Hippocampal Circuit Operations. *Science* 321: 53–57, 2008. doi: 10.1126/science.1149381.
5.   **Markram H**, **Toledo-Rodriguez M**, **Wang Y**, **Gupta A**, **Silberberg G**, **Wu C**. Interneurons of the
neocortical inhibitory system. *Nature Reviews Neuroscience* 5: 793, 2004. doi: 10.1038/nrn1519.
6.   **Pelkey KA**, **Chittajallu R**, **Craig MT**, **Tricoire L**, **Wester JC**, **McBain CJ**. Hippocampal
GABAergic Inhibitory Interneurons. *Physiological Reviews* 97: 1619–1747, 2017. doi:
10.1152/physrev.00007.2017.
7.   **Ascoli GA**, **Alonso-Nanclares L**, **Anderson SA**, **Barrionuevo G**, **Benavides-Piccione R**,
Burkhalter A, **Buzsáki G**, **Cauli B**, **DeFelipe J**, **Fairén A**, **Feldmeyer D**, **Fishell G**, **Fregnac Y**, **Freund**
TF, **Gardner D**, **Gardner EP**, **Goldberg JH**, **Helmstaedter M**, **Hestrin S**, **Karube F**, **Kisvárday ZF**,
Lambolez B, **Lewis DA**, **Marin O**, **Markram H**, **Muñoz A**, **Packer A**, **Petersen CCH**, **Rockland KS**,
Rossier J, **Rudy B**, **Somogyi P**, **Staiger JF**, **Tamas G**, **Thomson AM**, **Toledo-Rodriguez M**, **Wang Y**,
West DC, **Yuste R**, **The Petilla Interneuron Nomenclature Group (PING)**. Petilla terminology:
nomenclature of features of GABAergic interneurons of the cerebral cortex. *Nature Reviews Neuroscience*
9: 557–568, 2008. doi: 10.1038/nrn2402.
8.   **Kepecs A**, **Fishell G**. Interneuron cell types are fit to function. *Nature* 505: 318–326, 2014. doi:
10.1038/nature12983.
9.   **Zeng H**, **Sanes JR**. Neuronal cell-type classification: challenges, opportunities and the path
forward. *Nature Reviews Neuroscience* 18: 530, 2017. doi: 10.1038/nrn.2017.85.
10.   **Andersen P**, **Morris R**, **Amaral D**, **Bliss T**, **O'Keefe J**. *The Hippocampus Book*. Oxford, New
York: Oxford University Press, 2006.
11.   **Harris KD**, **Hochgerner H**, **Skene NG**, **Magno L**, **Katona L**, **Gonzales CB**, **Somogyi P**, **Kessaris**
N, **Linnarsson S**, **Hjerling-Leffler J**. Classes and continua of hippocampal CA1 inhibitory neurons
revealed by single-cell transcriptomics. *PLOS Biology* 16: e2006387, 2018. doi:
10.1371/journal.pbio.2006387.
12.   **Acsády L**, **Görcs TJ**, **Freund TF**. Different populations of vasoactive intestinal polypeptide-
immunoreactive interneurons are specialized to control pyramidal cells or interneurons in the hippocampus.
Neuroscience 73: 317–334, 1996. doi: 10.1016/0306-4522(95)00609-5.
13.   **Chamberland S**, **Salesse C**, **Topolnik D**, **Topolnik L**. Synapse-Specific Inhibitory Control of
Hippocampal Feedback Inhibitory Circuit. *Front Cell Neurosci* 4, 2010. doi: 10.3389/fncel.2010.00130.
14.   **Chamberland S**, **Topolnik L**. Inhibitory control of hippocampal inhibitory neurons. *Front*
Neurosci 6, 2012. doi: 10.3389/fnins.2012.00165.
15.   **Tyan L**, **Chamberland S**, **Magnin E**, **Camiré O**, **Francavilla R**, **David LS**, **Deisseroth K**,
Topolnik L. Dendritic Inhibition Provided by Interneuron-Specific Cells Controls the Firing Rate and
Timing of the Hippocampal Feedback Inhibitory Circuitry. *J Neurosci* 34: 4534–4547, 2014. doi:
10.1523/JNEUROSCI.3813-13.2014.
16.   **Luo X**, **Guet-McCreight A**, **Villette V**, **Francavilla R**, **Marino B**, **Chamberland S**, **Skinner FK**,
Topolnik L. Synaptic Mechanisms Underlying the Network State-Dependent Recruitment of VIP-
Expressing Interneurons in the CA1 Hippocampus. *Cerebral Cortex* 30: 3667–3685, 2020. doi:
10.1093/cercor/bhz334.
17.   **Guet-McCreight A**, **Skinner FK**, **Topolnik L**. Common Principles in Functional Organization of

VIP/Calretinin Cell-Driven Disinhibitory Circuits Across Cortical Areas. *Front Neural Circuits* 14, 2020. doi: 10.3389/fncir.2020.00032.

18. **Francavilla R**, **Villette V**, **Luo X**, **Chamberland S**, **Muñoz-Pino E**, **Camiré O**, **Wagner K**, **Kis V**, **Somogyi P**, **Topolnik L**. Connectivity and network state-dependent recruitment of long-range VIP-GABAergic neurons in the mouse hippocampus. *Nature Communications* 9: 5043, 2018. doi: 10.1038/s41467-018-07162-5.

19. **Maccaferri G**, **McBain CJ**. Passive propagation of LTD to stratum oriens-alveus inhibitory neurons modulates the temporoammonic input to the hippocampal CA1 region. *Neuron* 15: 137–145, 1995. doi: 10.1016/0896-6273(95)90071-3.

20. **Leão RN**, **Mikulovic S**, **Leão KE**, **Munguba H**, **Gezelius H**, **Enjin A**, **Patra K**, **Eriksson A**, **Loew LM**, **Tort ABL**, **Kullander K**. OLM interneurons differentially modulate CA3 and entorhinal inputs to hippocampal CA1 neurons. *Nature Neuroscience* 15: 1524–1530, 2012. doi: 10.1038/nn.3235.

21. **Siegle JH**, **Wilson MA**. Enhancement of encoding and retrieval functions through theta phase-specific manipulation of hippocampus. *eLife* 3: e03061, 2014. doi: 10.7554/eLife.03061.

22. **Siwani S**, **França ASC**, **Mikulovic S**, **Reis A**, **Hilscher MM**, **Edwards SJ**, **Leão RN**, **Tort ABL**, **Kullander K**. OLMα2 Cells Bidirectionally Modulate Learning. *Neuron* 99: 404-412.e3, 2018. doi: 10.1016/j.neuron.2018.06.022.

23. **Martina M**, **Vida I**, **Jonas P**. Distal Initiation and Active Propagation of Action Potentials in Interneuron Dendrites. *Science* 287: 295–300, 2000. doi: 10.1126/science.287.5451.295.

24. **Topolnik L**, **Chamberland S**, **Pelletier J-G**, **Ran I**, **Lacaille J-C**. Activity-Dependent Compartmentalized Regulation of Dendritic Ca2+ Signaling in Hippocampal Interneurons. *J Neurosci* 29: 4658–4663, 2009. doi: 10.1523/JNEUROSCI.0493-09.2009.

25. **Maccaferri G**. Stratum oriens horizontal interneurone diversity and hippocampal network dynamics. *The Journal of Physiology* 562: 73–80, 2005. doi: https://doi.org/10.1113/jphysiol.2004.077081.

26. **Maccaferri G**, **McBain CJ**. The hyperpolarization-activated current (Ih) and its contribution to pacemaker activity in rat CA1 hippocampal stratum oriens-alveus interneurones. *The Journal of Physiology* 497: 119–130, 1996. doi: 10.1113/jphysiol.1996.sp021754.

27. **Sekulic V**, **Yi F**, **Garrett T**, **Guet-McCreight A**, **Lawrence JJ**, **Skinner FK**. Integration of within-cell experimental data with multi-compartment modeling predicts h-channel densities and distributions in hippocampal OLM cells. *Front Cell Neurosci* 14, 2020. doi: 10.3389/fncel.2020.00277.

28. **Zemankovics R**, **Káli S**, **Paulsen O**, **Freund TF**, **Hájos N**. Differences in subthreshold resonance of hippocampal pyramidal cells and interneurons: the role of h-current and passive membrane characteristics. *The Journal of Physiology* 588: 2109–2132, 2010. doi: 10.1113/jphysiol.2009.185975.

29. **Ascoli GA**, **Gasparini S**, **Medinilla V**, **Migliore M**. Local Control of Postinhibitory Rebound Spiking in CA1 Pyramidal Neuron Dendrites. *J Neurosci* 30: 6434–6442, 2010. doi: 10.1523/JNEUROSCI.4066-09.2010.

30. **Kispersky TJ**, **Fernandez FR**, **Economo MN**, **White JA**. Spike Resonance Properties in Hippocampal O-LM Cells Are Dependent on Refractory Dynamics. *J Neurosci* 32: 3637–3651, 2012. doi: 10.1523/JNEUROSCI.1361-11.2012.

31. **Sekulić V**, **Skinner FK**. Computational models of O-LM cells are recruited by low or high theta frequency inputs depending on h-channel distributions. *eLife Sciences* 6: e22962, 2017. doi: 10.7554/eLife.22962.

32. **Guet-McCreight A**, **Skinner FK**. Using computational models to predict in vivo synaptic inputs to interneuron specific 3 (IS3) cells of CA1 hippocampus that also allow their recruitment during rhythmic states. *PLOS ONE* 14: e0209429, 2019. doi: 10.1371/journal.pone.0209429.

33. **Guet-McCreight A**, **Skinner FK**. Computationally going where experiments cannot: a dynamical assessment of dendritic ion channel currents during in vivo-like states. *F1000Res* 9: 180, 2020. doi: 10.12688/f1000research.22584.2.

34. **Carnevale NT**, **Hines ML**. *The NEURON Book*. 1 edition. Cambridge, UK; New York: Cambridge University Press, 2006.

35. **Marder E**. Variability, compensation, and modulation in neurons and circuits. *PNAS* 108: 15542–
15548, 2011. doi: 10.1073/pnas.1010674108.
36. **Schultheiss NW**, **Edgerton JR**, **Jaeger D**. Phase Response Curve Analysis of a Full Morphological
Globus Pallidus Neuron Model Reveals Distinct Perisomatic and Dendritic Modes of Synaptic Integration. *J*
Neurosci 30: 2767–2782, 2010. doi: 10.1523/JNEUROSCI.3959-09.2010.
37. **Destexhe A**, **Rudolph M**, **Paré D**. The high-conductance state of neocortical neurons in vivo. *Nat*
Rev Neurosci 4: 739–751, 2003. doi: 10.1038/nrn1198.
38. **Sivagnanam S**, **Majumdar A**, **Yoshimoto K**, **Astakhov V**, **B A**, **Martone M**, **Carnevale NT**.
Introducing The Neuroscience Gateway [Online]. http://ceur-ws.org/Vol-993/paper10.pdf.
39. **Bezaire MJ**, **Raikov I**, **Burk K**, **Vyas D**, **Soltesz I**. Interneuronal mechanisms of hippocampal
theta oscillations in a full-scale model of the rodent CA1 circuit. *eLife Sciences* 5: e18566, 2016. doi:
10.7554/eLife.18566.
40. **Hangya B**, **Borhegyi Z**, **Szilágyi N**, **Freund TF**, **Varga V**. GABAergic Neurons of the Medial
Septum Lead the Hippocampal Network during Theta Activity. *J Neurosci* 29: 8094–8102, 2009. doi:
10.1523/JNEUROSCI.5665-08.2009.
41. **Rinzel J**, **Ermentrout GB**. Analysis of neural excitability and oscillations [Online]. In: *Methods in*
Neuronal Modeling, edited by Koch C, Segev I. MIT Press, p. 135–169.
http://dl.acm.org/citation.cfm?id=94605.94613 [31 Jul. 2019].
42. **Schultheiss NW**, **Prinz AA**, **Butera RJ**, editors. *Phase Response Curves in Neuroscience: Theory,*
Experiment, and Analysis. 2012 edition. New York: Springer, 2011.
43. **Katona L**, **Lapray D**, **Viney TJ**, **Oulhaj A**, **Borhegyi Z**, **Micklem BR**, **Klausberger T**, **Somogyi**
P. Sleep and Movement Differentiates Actions of Two Types of Somatostatin-Expressing GABAergic
Interneuron in Rat Hippocampus. *Neuron* 82: 872–886, 2014. doi: 10.1016/j.neuron.2014.04.007.
44. **Varga C**, **Golshani P**, **Soltesz I**. Frequency-invariant temporal ordering of interneuronal discharges
during hippocampal oscillations in awake mice. *PNAS* 109: E2726–E2734, 2012. doi:
10.1073/pnas.1210929109.
45. **Geiller T**, **Vancura B**, **Terada S**, **Troullinou E**, **Chavlis S**, **Tsagkatakis G**, **Tsakalides P**, **Ócsai**
K, **Poirazi P**, **Rózsa BJ**, **Losonczy A**. Large-Scale 3D Two-Photon Imaging of Molecularly Identified CA1
Interneuron Dynamics in Behaving Mice. *Neuron* 108: 968-983.e9, 2020. doi:
10.1016/j.neuron.2020.09.013.
46. **Buzsáki G**, **Mizuseki K**. The log-dynamic brain: how skewed distributions affect network
operations. *Nature Reviews Neuroscience* 15: 264–278, 2014. doi: 10.1038/nrn3687.
47. **Royer S**, **Zemelman BV**, **Losonczy A**, **Kim J**, **Chance F**, **Magee JC**, **Buzsáki G**. Control of
timing, rate and bursts of hippocampal place cells by dendritic and somatic inhibition. *Nature Neuroscience*
15: 769–775, 2012. doi: 10.1038/nn.3077.
48. **Lovett-Barron M**, **Turi GF**, **Kaifosh P**, **Lee PH**, **Bolze F**, **Sun X-H**, **Nicoud J-F**, **Zemelman BV**,
Sternson SM, **Losonczy A**. Regulation of neuronal input transformations by tunable dendritic inhibition.
Nature Neuroscience 15: 423–430, 2012. doi: 10.1038/nn.3024.
49. **Turi GF**, **Li W-K**, **Chavlis S**, **Pandi I**, **O'Hare J**, **Priestley JB**, **Grosmark AD**, **Liao Z**, **Ladow**
M, **Zhang JF**, **Zemelman BV**, **Poirazi P**, **Losonczy A**. Vasoactive Intestinal Polypeptide-Expressing
Interneurons in the Hippocampus Support Goal-Oriented Spatial Learning. *Neuron* 101: 1150-1165.e8,
2019. doi: 10.1016/j.neuron.2019.01.009.
50. **Chatzikalymniou AP**, **Skinner FK**. Deciphering the Contribution of Oriens-
Lacunosum/Moleculare (OLM) Cells to Intrinsic θ Rhythms Using Biophysical Local Field Potential (LFP)
Models. *eNeuro* 5, 2018. doi: 10.1523/ENEURO.0146-18.2018.
51. **Hilscher MM**, **Nogueira I**, **Mikulovic S**, **Kullander K**, **Leão RN**, **Leão KE**. Chrna2-OLM
interneurons display different membrane properties and h-current magnitude depending on dorsoventral
location. *Hippocampus* 29: 1224–1237, 2019. doi: https://doi.org/10.1002/hipo.23134.
52. **Mikulovic S**, **Restrepo CE**, **Siwani S**, **Bauer P**, **Pupe S**, **Tort ABL**, **Kullander K**, **Leão RN**.
Ventral hippocampal OLM cells control type 2 theta oscillations and response to predator odor. *Nature*

Communications 9: 3638, 2018. doi: 10.1038/s41467-018-05907-w.

53.   **Canavier CC**, **Achuthan S**. Pulse coupled oscillators. *Scholarpedia* 2: 1331, 2007. doi: 10.4249/scholarpedia.1331.

54.   **Hansel D**, **Mato G**, **Meunier C**. Synchrony in Excitatory Neural Networks. *Neural Computation* 7: 307–337, 1995. doi: 10.1162/neco.1995.7.2.307.

55.   **Stiefel KM**, **Ermentrout GB**. Neurons as oscillators. *Journal of Neurophysiology* 116: 2950–2960, 2016. doi: 10.1152/jn.00525.2015.

56.   **Stiefel KM**, **Gutkin BS**, **Sejnowski TJ**. The effects of cholinergic neuromodulation on neuronal phase-response curves of modeled cortical neurons. *J Comput Neurosci* 26: 289–301, 2009. doi: 10.1007/s10827-008-0111-9.

57.   **Steuber V**, **Schultheiss NW**, **Silver RA**, **De Schutter E**, **Jaeger D**. Determinants of synaptic integration and heterogeneity in rebound firing explored with data-driven models of deep cerebellar nucleus cells. *J Comput Neurosci* 30: 633–658, 2011. doi: 10.1007/s10827-010-0282-z.

58.   **Stiefel KM**, **Gutkin BS**, **Sejnowski TJ**. Cholinergic Neuromodulation Changes Phase Response Curve Shape and Type in Cortical Pyramidal Neurons. *PLOS ONE* 3: e3947, 2008. doi: 10.1371/journal.pone.0003947.

59.   **Lawrence JJ**. Cholinergic control of GABA release: emerging parallels between neocortex and hippocampus. *Trends in Neurosciences* 31: 317–327, 2008. doi: 10.1016/j.tins.2008.03.008.

60.   **Böhm C**, **Pangalos M**, **Schmitz D**, **Winterer J**. Serotonin Attenuates Feedback Excitation onto O-LM Interneurons. *Cereb Cortex* 25: 4572–4583, 2015. doi: 10.1093/cercor/bhv098.

61.   **Govindaiah G**, **Kang Y-J**, **Lewis HES**, **Chung L**, **Clement EM**, **Greenfield LJ**, **Garcia-Rill E**, **Lee S-H**. Group I metabotropic glutamate receptors generate two types of intrinsic membrane oscillations in hippocampal oriens/alveus interneurons. *Neuropharmacology* 139: 150–162, 2018. doi: 10.1016/j.neuropharm.2018.06.035.

62.   **Losonczy A**, **Somogyi P**, **Nusser Z**. Reduction of Excitatory Postsynaptic Responses by Persistently Active Metabotropic Glutamate Receptors in the Hippocampus. *Journal of Neurophysiology* 89: 1910–1919, 2003. doi: 10.1152/jn.00842.2002.

63.   **Haam J**, **Zhou J**, **Cui G**, **Yakel JL**. Septal cholinergic neurons gate hippocampal output to entorhinal cortex via oriens lacunosum moleculare interneurons. *PNAS* 115: E1886–E1895, 2018. doi: 10.1073/pnas.1712538115.

64.   **Lawrence JJ**, **Grinspan ZM**, **Statland JM**, **McBain CJ**. Muscarinic receptor activation tunes mouse stratum oriens interneurones to amplify spike reliability. *The Journal of Physiology* 571: 555–562, 2006. doi: 10.1113/jphysiol.2005.103218.

65.   **Schmid LC**, **Mittag M**, **Poll S**, **Steffen J**, **Wagner J**, **Geis H-R**, **Schwarz I**, **Schmidt B**, **Schwarz MK**, **Remy S**, **Fuhrmann M**. Dysfunction of Somatostatin-Positive Interneurons Associated with Memory Deficits in an Alzheimer's Disease Model. *Neuron* 92: 114–125, 2016. doi: 10.1016/j.neuron.2016.08.034.

66.   **Hájos N**, **Freund TF**, **Mody I**. Comparison of single NMDA receptor channels recorded on hippocampal principal cells and oriens/alveus interneurons projecting to stratum lacunosum-molecular (O-LM cells). *Acta Biologica Hungarica* 53: 465–472, 2002. doi: 10.1556/ABiol.53.2002.4.7.

67.   **Oren I**, **Nissen W**, **Kullmann DM**, **Somogyi P**, **Lamsa KP**. Role of Ionotropic Glutamate Receptors in Long-Term Potentiation in Rat Hippocampal CA1 Oriens-Lacunosum Molecular Interneurons. *J Neurosci* 29: 939–950, 2009. doi: 10.1523/JNEUROSCI.3251-08.2009.

68.   **Hurtado-Zavala JI**, **Ramachandran B**, **Ahmed S**, **Halder R**, **Bolleyer C**, **Awasthi A**, **Stahlberg MA**, **Wagener RJ**, **Anderson K**, **Drenan RM**, **Lester HA**, **Miwa JM**, **Staiger JF**, **Fischer A**, **Dean C**. TRPV1 regulates excitatory innervation of OLM neurons in the hippocampus. *Nature Communications* 8: 15878, 2017. doi: 10.1038/ncomms15878.

69.   **Hertäg L**, **Sprekeler H**. Amplifying the redistribution of somato-dendritic inhibition by the interplay of three interneuron types. *PLOS Computational Biology* 15: e1006999, 2019. doi: 10.1371/journal.pcbi.1006999.

70.   **Hertäg L**, **Sprekeler H**. Learning prediction error neurons in a canonical interneuron circuit. *eLife*

9: e57541, 2020. doi: 10.7554/eLife.57541.
71.   **Wang X-J**, **Tegnér J**, **Constantinidis C**, **Goldman-Rakic PS**. Division of labor among distinct
subtypes of inhibitory neurons in a cortical microcircuit of working memory. *PNAS* 101: 1368–1373, 2004.
doi: 10.1073/pnas.0305337101.
72.   **Yang GR**, **Murray JD**, **Wang X-J**. A dendritic disinhibitory circuit mechanism for pathway-
specific gating. *Nat Commun* 7: 1–14, 2016. doi: 10.1038/ncomms12815.
73.   **Ecker A**, **Romani A**, **Sáray S**, **Káli S**, **Migliore M**, **Falck J**, **Lange S**, **Mercer A**, **Thomson AM**,
Muller E, **Reimann MW**, **Ramaswamy S**. Data-driven integration of hippocampal CA1 synaptic
physiology in silico. *Hippocampus* 30: 1129–1145, 2020. doi: https://doi.org/10.1002/hipo.23220.
74.   **Alonso LM**, **Marder E**. Visualization of currents in neural models with similar behavior and
different conductance densities. *eLife* 8: e42722, 2019. doi: 10.7554/eLife.42722.

Figure 1: OLM cell model output in *in vitro* and *in vivo* contexts. A: Schematic demonstrating the *in vitro* context where DC current is injected directly into the OLM cell model somatic compartment (top) such that spiking is generated. Currentscape plots (74) are shown below, where in each currentscape plot, the top trace is the voltage trace (y-axis scale bar = -50 mV [horizontal dashed line] to -20 mV), the filled-in black traces above and below the coloured plots are the total outward and inward currents respectively (dotted lines = ± 0.5 pA, ± 5 pA, ± 50 pA), and the coloured plots (see colour references in the legend on the right) show the percent contributions of each individual outward (top half of the plot) and inward (bottom half of the plot) current to the total outward and inward currents. B: Schematic demonstrating the *in vivo* context where the OLM cell model dendritic compartments are bombarded with synaptic inputs from different excitatory (PYR) and inhibitory (I-S3, BIS: Bistratified cell inputs, MS: Medial Septal inputs) presynaptic populations (top) such that the voltage dynamics exhibit characteristics that are typical of a high-conductance state. As in panel A, the currentscape plot is shown below.

Figure 2: I-S3 cell inputs alone cannot entrain OLM cell spiking at frequencies higher than their baseline spike rate. A: Example synaptic locations (left) of 30 I-S3 cell inputs onto an OLM cell model (note that synaptic locations can be overlayed). Simulated voltage traces (right) in the 9-10s simulation times across several frequency-modulated I-S3 cell input frequencies (left to right) and numbers of I-S3 cell synapses (top to bottom). Blue arrows indicate times when I-S3 cell synapses are activated. For each voltage trace axis, the two notches indicate voltages of -50 mV and 0 mV. B: Example synaptic locations (left) of 30 I-S3 cell inputs and 3 PYR cell inputs onto an OLM cell model. Simulated voltage traces (right; as in A) but with inclusion of 3 PYR cell inputs (timing indicated by the yellow arrows). Note that this number of PYR cell inputs is fixed across simulations. The times at which the 3 PYR cell inputs are activated are indicated by the yellow arrows and start with a delay of 78.125 ms relative to I-S3 cell inputs, consistent with their relative timing during theta rhythms (4, 16). C: Example synaptic locations (left) of 30 PYR cell inputs onto an OLM cell model. Simulated voltage traces (right; as in A) but with a scaling up of PYR cell inputs alone instead of I-S3 cell inputs alone. D-F: Blown up versions of the traces highlighted by the stars in A-C, respectively. In all cases, the models are given somatically injected current to generate spiking at 7.25 Hz (cell 1: 49 pA; cell 2: 45 pA) to generate the mean baseline OLM cell spike rates reported by Tyan et al. (15). This figure displays results for cell 1. Results for cell 2 are similar (**OSF Figure 2**).

Figure 3: PRC exploration illustration. A: Relationship between holding current and resulting spike rates in cell 1. Dashed line indicates the holding current level used for generating baseline spiking that is used to examine phase responses in panels B-F. B: Resulting PRCs due to large negative and small positive 3ms square current pulses (-500, -300, -100, 10, 30, 50 pA) being moved from earlier (0%) to later (100%) phases of an inter-spike interval cycle. More positive phase shifts indicate larger phase delays. C: Resulting PRCs due to inhibitory perturbation (6 to 60 I-S3 cell synapses) being moved from earlier (0%) to later (100%) phases of an inter-spike interval cycle. Example perturbation phases at which an advance and a delay can be seen are indicated by the yellow and red dashed lines, respectively. D: Perturbation phase-dependent changes in peak I_H amplitude for different numbers of I-S3 cell synapses. E: Voltage traces at baseline without perturbation (dashed), with an advance-causing perturbation of 30 I-S3 cell synapses (yellow), and with a delay-causing perturbation of 30 I-S3 cell synapses (red). Colors correspond with the perturbation phases highlighted by the colored dashed lines in C and F. F: Perturbation phase-dependent changes (30 I-S3 cell synapses) in peak current amplitude for each ion channel present in the model. Currents are colored as follows (same order as in legend): I_{KA} (blue), I_{Kdrf} (orange), I_{Kdrs} (green), I_{KCa} (red), I_M (purple), I_L (brown), I_{Na} (pink), I_{CaT} (grey), I_{CaL} (yellow), and I_H (cyan). The measurements shown here are extracted from current traces obtained from the first dendritic compartment adjacent to the soma since calcium channels are not present in the somatic compartments.

Figure 4: *I_H-modulated phase advances only occur when spiking is near rheobase*. A: PRCs for cell 1 (solid) and cell 2 (dashed), during rheobase, 4 Hz, 7.25 Hz, and 15 Hz OLM cell spiking, using a perturbation of 30 I-S3 cell synapses. B: Perturbation phase-dependent changes in peak I_H amplitude for cell 1 (solid) and cell 2 (dashed). C: Perturbation phase-dependent changes in peak I_M amplitude for cell 1 (solid) and cell 2 (dashed). It is clear that inhibitory perturbations alone mostly just cause phase delays in OLM cell models.

Figure 5: Cell 1 *in vitro* spike resonant frequencies due to inhibitory inputs, but not excitatory inputs, are dependent on the baseline spike rate. A: Example synaptic locations of 30 I-S3 cell modulatory inputs during somatic current injections to generate different f_B. B: Baseline ratios computed across different I-S3 cell modulation frequencies, f_I, and different holding currents (as shown by the range of different baseline frequencies, f_B, indicated by the colorbar). The colored dots and histograms plotted above the traces indicate the peak/resonant frequencies and distributions across f_B values, respectively. The dashed black line indicates a baseline ratio of 1 (note that the y-axis scale is large, so this line appears very close to 0). Values larger/smaller than 1 indicate an increase/decrease in the PSD at f_I. C: Resonant frequencies, f_R, plotted against f_B. Dots in blue/red areas indicate simulations where f_R is greater/lesser than f_B. D: Spike rate at the f_R plotted against f_B. Dots in blue/red areas indicate simulations where the f_R spike rate is increased/decreased compared to f_B (i.e. consistent with phase advances/delays). E-H: Same as in A-D, but with 30 PYR cell modulatory inputs instead.

Figure 6: Cell 1 *in vivo* spike resonant frequencies are in theta frequency ranges only for inhibitory inputs. A: Example synaptic locations of IVL synapses (top) and 30 I-S3 cell modulatory inputs (bottom). B: Baseline ratios computed across different I-S3 cell modulation frequencies, f_I, and different randomized IVL states (as shown by the f_B range, indicated by the colorbar). The colored dots and histograms plotted above the traces indicate the peak/resonant frequencies and distributions across f_B values, respectively. The dashed black line indicates a baseline ratio of 1. Values larger/smaller than 1 indicate an increase/decrease in the PSD at f_I. C: Resonant frequencies, f_R, plotted against f_B. Dots in blue/red areas indicate simulations where f_R is greater/lesser than f_B. D: Spike rate at the f_R plotted against f_B. Dots in blue/red areas indicate simulations where the f_R spike rate is increased/decreased compared to f_B (i.e. consistent with phase advances/delays). Note that axis scales are chosen to match the scales in **Fig. 5C-D & G-H** for comparison purposes. E-H: Same as in A-D, but with 30 PYR cell modulatory inputs instead.

984

985 **Figure 7: Ramping up I-S3 cell inputs can sharpen and modulate OLM cell recruitment (cell 1).** A: Schematic

986 of simulations where, on top of IVL inputs, we add theta-timed inputs, a ramp-up of I-S3 cell inputs as each

987 simulation progresses (depicted in the bottom-left subplots) as well as either CA3 (+15.625°) or EC (-15.625°) -

988 modulated shifts in the timing of different input populations. In this first case we impose a shift in the timing of I-S3

989 cell inputs alone as I-S3 cell inputs are ramped up. B: Plots of the simulated OLM cell model voltage traces averaged

990 across theta cycles. From top to bottom, these plots show the model simulations without a ramp-up (black), with an I-

991 S3 cell ramp-up near the rising phase (red; i.e. stronger CA3 inputs), with an I-S3 cell ramp-up between the rising

992 and peak phases (green; i.e. even inputs from CA3 and EC), as well as with an I-S3 cell ramp-up near the peak

993 phase (blue; i.e. stronger EC inputs), while it is being modulated by 8 Hz frequency inputs from all input

994 population types. To generate these average traces, we take the voltage traces (1,000 to 10,000 ms), split them each

995 into their 72 theta cycles (i.e. 9000 ms/125 ms = 72 cycles), and then compute the average 125 ms theta cycle

996 traces. Shaded areas show the amount of standard deviation above or below the mean. C: Resulting suppression

997 on the spike rate of the OLM cell model throughout the simulations. D: The PSD of the voltage traces before and

998 after applying ramp-ups of I-S3 cell inputs. The inset plot shows the magnified PSD between 7 to 9 Hz. E: In the

999 second case we impose shifts in the timing of both I-S3 cell inputs as well as PYR cell inputs as I-S3 cell inputs are

1000 ramped up. F-H: Same as in B-D but with shifts in both I-S3 and PYR cell inputs, as described in E.

1001

Figure 8: I-S3 cells are fit to suppress OLM cell spiking and disinhibit pyramidal cells.
In our proposed mechanism, I-S3 cells control OLM cell spiking through spike suppression (step 1), leading to disinhibition of PYR cell distal dendrites (step 2), enabling excitation of OLM cells (step 3), and subsequent re-gating of PYR cell distal dendrites via OLM cell-mediated inhibition.