

Title: Investigation of COVID-19 comorbidities reveals genes and pathways coincident with the SARS-CoV-2 viral disease.

Authors:

Mary E. Dolan^{1*,2}, David P. Hill^{1,2}, Gaurab Mukherjee², Monica S. McAndrews², Elissa J. Chesler², Judith A. Blake²

¹ These authors contributed equally and should be considered co-first authors

* Corresponding author mary.dolan@jax.org

² The Jackson Laboratory, 600 Main St, Bar Harbor, ME 04609, USA

Abstract:

The emergence of the SARS-CoV-2 virus and subsequent COVID-19 pandemic initiated intense research into the mechanisms of action for this virus. It was quickly noted that COVID-19 presents more seriously in conjunction with other human disease conditions such as hypertension, diabetes, and lung diseases. We conducted a bioinformatics analysis of COVID-19 comorbidity-associated gene sets, identifying genes and pathways shared among the comorbidities, and evaluated current knowledge about these genes and pathways as related to current information about SARS-CoV-2 infection. We performed our analysis using GeneWeaver (GW), Reactome, and several biomedical ontologies to represent and compare common COVID-19 comorbidities. Phenotypic analysis of shared genes revealed significant enrichment for immune system phenotypes and for cardiovascular-related phenotypes, which might point to alleles and phenotypes in mouse models that could be evaluated for clues to COVID-19 severity. Through pathway analysis, we identified enriched pathways shared by comorbidity datasets and datasets associated with SARS-CoV-2 infection.

Introduction

COVID-19 emerged as a global pandemic through the first half of 2020¹. The severity of the disease varies from asymptomatic to lethal with a case mortality rate in the 20 most affected countries ranging between 1 and 15.3% (<https://coronavirus.jhu.edu/data/mortality> ; retrieved 24 July 2020). Severe disease shows manifestations of both acute respiratory distress syndrome (ARDS) and cytokine release syndrome (CRS)^{2,3}. In pediatric patients, a blood vessel inflammatory pathology similar to Kawasaki disease is sometimes present⁴. All of these presentations have common elements of abnormality of inflammatory responses and manifestations of vascular defects such as thrombosis, which may be causally related^{5,6,7,8,9}.

Since the emergence and global transmission of the SARS-CoV-2 virus, many studies have reported that patients with certain underlying medical conditions have especially severe responses to the coronavirus infection¹⁰. Some of the identified comorbidities that lead to severe disease are Cardiovascular Disease, Diabetes, Hepatitis, Lung Disease, and Kidney Disease^{11,12,13,14,15,16,17}.

Understanding what makes some patients suffer from severe COVID-19 is an ongoing puzzle that is being investigated from both the virus and host perspectives¹⁸. We hypothesize that by exploring the underlying genetic basis of comorbidities associated with severe disease, we can identify putative host genes and pathways that are responsible for or contribute to the severity. Identification of these genes and pathways can serve as a gateway for further investigation into understanding how the host responds to the virus and for potential therapeutic strategies to interfere with a severe outcome.

We interrogated gene sets that are associated with the five previously mentioned underlying comorbidities to determine gene products that are shared among them. We identified several pathways and phenotypes in common, including those that are associated with severe COVID-19 pathology. All of the comorbid diseases have been and continue to be actively studied, now in the additional context of response to SARS-CoV-2 infection^{19,20,21,22,23}. In particular, the laboratory mouse has been extensively utilized as an animal model to study these conditions²⁴. As a result, mouse strains carrying mutations in shared genes or genes in shared pathways, and engineered to be capable of being infected by the virus, can present useful starting points for investigating the biological basis of disease severity²⁵.

We report here on investigations of the host genetics and genomics of a set of comorbidity conditions. We include data identifying the shared pathways and cellular mechanisms associated with these diseases and correlate these data with recent studies of the genetic basis of the COVID-19 to identify elements that are

shared among comorbidities and the host response to the disease. Our results suggest specific directions of future study to understand the genetic foundation of severe COVID-19.

Materials and Methods

Gene Sets Used for Analysis

All gene sets used in our analysis are publicly available from the GeneWeaver resource

(www.geneweaver.org)²⁶. Genes associated with Cardiovascular Disease, Diabetes, Hepatitis, Lung Disease were derived from gene sets associated with MeSH terms that relate to these comorbidities. The gene set for Kidney Disease was derived from the union of genes associated with Proteinuria Hematuria, Elevated Serum Creatinine, Increased Blood Urea Nitrogen and Decreased Glomerular Filtration Rate in the Human Phenotype Ontology (HPO)²⁷. The genes in the MeSH and HPO gene sets and associated metadata (indicating their association with COVID-19 and citations supporting the association) were incorporated into GW and used for analysis. The comorbidity-related gene sets are shown in Table 1.

To create gene sets that are directly related to SARS-CoV-2 infection (COVID-19), we identified several relevant reports^{5,28,29} and captured gene sets available from these studies into the GeneWeaver environment. Details are shown in Table 2.

Gene Set Comparison

To identify genes that were shared by all five comorbidities or four out of five comorbidities, we used the GeneWeaver 'Combine GeneSets' tool. To visualize the intersection of comorbidity gene sets graphically, we used the GeneWeaver 'HiSim graph' tool. To create a hierarchical view of interleukin pathways and the genes that are shared among comorbidities we used the HiSim graph tool at the GeneWeaver resource with homology excluded.

Functional Analysis of Gene Sets

To evaluate the shared biology of the common genes, we interrogated the genes that were shared among comorbidities in two ways: 1) we conducted a phenotype enrichment analysis using the VLAD enrichment tool³⁰ and 2) we conducted a pathway enrichment analysis using the Reactome Knowledgebase resource³¹.

Mammalian Phenotype Enrichment Analysis

To investigate details of the phenotypes associated with the shared genes, we took advantage of the mammalian phenotype data available from the Mouse Genome Informatics site (www.informatics.jax.org). The Mouse Genome Database (MGD) group captures phenotypic data using the Mammalian Phenotype Ontology (MP), a computable ontological structure, that can be queried and used for phenotypic enrichment analysis³². MGD also integrates these murine data into the context of human disease data based on orthology and gene expression^{32,24}. This integrated resource allowed us to exploit the associations of mouse genes and their phenotypes for enrichment purposes using VLAD, and gives us an entry into identification of potential mouse models for future study³³.

Mouse orthologs for the shared human genes were identified using data available from the Alliance of Genome Resources (Alliance) (www.alliancegenome.org/)³⁴ using the Alliance release 3.1 stringent mouse-human orthology set. If a human gene symbol matched more than one mouse marker, that gene was not included in the analysis. This resulted in the following fourteen human genes being excluded from the analysis: *AGTR1*, *CCL2*, *CFH*, *CYP2D6*, *CYP3A4*, *GSTM1*, *GSTP1*, *HAMP*, *HLA-B*, *HLA-DRB1*, *IFNA1*, *MMP1*, *SERPINA1*, and *TIMP2*. We did not identify mouse orthologs for two human genes (*CXCL8*, *HLA-DQB1*). Excluding these 16 from the initial 123 genes that were shared among four of five comorbidities, left us with 107 remaining mouse orthologs which we used in the VLAD analysis. All but one of the 107 mouse genes (*H2-Ea*, a polymorphic pseudogene; i.e. a coding gene in some strains and a pseudogene in others) had annotations to MP. The VLAD phenotype analysis was run on 20 August 2020 using annotation data from 20 August 2020, ontology data from 9 July 2020 and default parameters.

Reactome Pathway Enrichment Analysis

For pathway enrichment, we submitted the gene lists to the Reactome 'Analyze Gene List' enrichment tool (<https://reactome.org/PathwayBrowser/#TOOL=AT>) based on Reactome version 72. The analysis was performed on 18 May 2020. Results were downloaded using the 'Pathway Analysis Results' and 'Analysis Report' functionality at Reactome.

Pathway enrichment analysis was also performed for six COVID-19-related gene sets shown in Table 2. We selected up-regulated (\log_2 fold change ≥ 2) genes in host transcriptional response to SARS-CoV-2 in three cell cultures: human A549 lung alveolar cells (102 genes), Calu3 human lung adenocarcinoma epithelial cells (333 genes), normal human bronchial epithelium (NHBE) cells (14 genes); genes that are up-regulated in post-mortem lung samples from COVID-19-positive patients relative to biopsied healthy lung tissue from uninfected individuals (586 genes)²⁸; immune-response and angiogenesis-related genes that are up-regulated in lungs from patients who died from COVID-19 (114 genes)⁵; genes that are overexpressed in severe compared to mild cases of COVID-19 (4 genes)²⁹. We corrected for any symbols that were out of date and again used the Reactome Pathway analysis tool. The analysis was performed on 10 August 2020.

Results

COVID-19 comorbidities share associated genes

To test our hypothesis that comorbidities associated with COVID-19 severity have common underlying molecular bases, we chose five comorbidities that have been reported in the literature as closely associated with poorer disease outcome: Kidney Disease, Liver Disease, Diabetes, Lung Disease and Cardiovascular Disease. We searched the GeneWeaver Data repository for gene sets associated with these comorbidities and identified gene sets from MeSH and HPO that we used in our analyses (Table 1).

To identify genes that were shared among the five comorbidity gene sets, we used the ‘Combine GeneSets’ tool to create a matrix of genes and sets in which they were contained. We tabulated the number of gene sets that contained each gene and determined that eight genes were present in all five sets: *APOA1*, *APOE*, *B2M*, *CTLA4*, *F2*, *F5*, *HMOX1* and *STAT3*; 123 genes were common to at least four out of five comorbidity sets (Table 3).

Genes shared among COVID-19 comorbidities are enriched for phenotypes corresponding to immune system processes and circulatory system biology

We tested the functional significance of the genes shared among the five comorbidities by performing two different kinds of enrichment analysis on our gene sets. First, we identified the mouse orthologs of the human

genes and performed a phenotype enrichment using the Visual Annotation Display tool (VLAD)

(<http://proto.informatics.jax.org/prototypes/vlad/>). For genes shared among all five comorbidities, VLAD enrichment analysis identified 762 significantly enriched ($p < 0.05$) mammalian phenotypes (supplemental table 1). The most significantly enriched terms fall into three general categories: T-cell related phenotypes, inflammation or infection related phenotypes, and cardiovascular phenotypes including blood clotting. Table 4 shows that of the eight shared genes, several were annotated to each of the significantly enriched phenotypes.

We repeated our phenotype enrichment analysis using genes that are co-annotated to four of the five comorbidities associated with COVID-19. When we examined the shared genes among comorbidity sets, we found that 123 genes were shared among four out of five comorbidities. Phenotype enrichment analysis performed with the 107 one-to-one mouse orthologs of these human genes was consistent with our analysis of the eight genes that were conserved in all five comorbidities. The increase in gene number resulted in an increase in the number of significantly enriched mammalian phenotype terms ($p < 0.05$) with 3232 terms included in the enrichment analysis (supplemental table 2). VLAD analysis showed that the major areas of the ontology with the most highly significant enrichment were, as in the analysis for the eight genes shared by all the comorbidities, in inflammatory response and infection, leukocyte biology and blood vessel morphology. Abnormal blood coagulation was no longer in the most highly significant group of phenotypes, but was significantly enriched ($p = 1.52 \times 10^{-11}$).

Similar to our results for the eight genes shared among all five comorbidities, the mouse orthologs of the 123 genes shared in four out of five comorbidities showed many genes associated with each of the most significant phenotypes (Table 5).

Pathway analysis enrichment includes cytokine signaling pathways, blood coagulation and plasma lipoprotein metabolism

In addition to our phenotype analysis, we were also interested in investigating whether the genes shared among comorbidities were enriched for specific biological pathways. To answer this question, we used our human gene sets and the Reactome Knowledgebase in an enrichment analysis for biological pathways.

Reactome is a manually curated resource that captures information about reactions, their relationships and the genes and chemicals that play a role in those reactions³⁵.

We interrogated the Reactome Knowledgebase using the eight genes that were shared among all five comorbidities and identified 103 pathways/subpathways that were significantly enriched (FDR=<0.05, supplemental table 5). Reactome pathways are organized into a hierarchical format where grouping pathways are subcategorized into more specific pathways which in turn are eventually represented by individual reactions. Reactome captures information about not only the genes and molecules that act in a pathway but also those that are acted upon, thus casting a wide net for genes that are included in an analysis. The 25 most significantly enriched pathways grouped under several parent pathways are shown in Figure 1. Two genes, *APOA1* and *APOE*, are shared among several plasma lipoprotein assembly, remodeling and clearance pathways. Three genes, *APOA1*, *F2* and *F5*, are found in the hemostasis pathway, all are included in platelet activation and the latter two in blood clotting. Five of the eight genes shared among the comorbidities were involved in immune system pathways: *B2M*, *HMOX1*, *CTLA4*, *STAT3* and *F2*. Of these five genes, three are included in cytokine signaling: *B2M*, *HMOX1* and *STAT3*. Other informative pathways showed that *APOA1*, *APOE*, *F2* are in GPCR downstream signaling, and *F5*, *APOA1*, *APOE* are in vesicle-mediated transport.

We repeated the pathway enrichment analysis with the 123 genes that were shared by four out of five comorbidities. We identified 172 pathways that were significantly enriched (FDR=<0.05, supplemental table 6). These results supported and confirmed the results we obtained with the eight genes that were shared among all five comorbidities. Although with lower significance, enriched pathways include 'common pathway of fibrin clot formation' (FDR=5.9x10e-3; four genes), 'platelet degranulation' (FDR=6.0x10e-6; thirteen genes) and 'plasma lipoprotein assembly remodeling and clearance' (FDR=0.034; five genes). Immune signaling pathways and particularly interleukin signaling pathways were frequent in our enrichment results (Figure 2). The downstream GPCR signaling pathway and the retinoid/vitamin pathways were no longer significantly enriched.

We compared our pathway enrichment results with our phenotype enrichment results for the eight genes conserved among all five comorbidities to determine if there was consistency between the results obtained

from independently curated resources: MGI phenotype and Reactome. Like the pathway analysis, our phenotype enrichment analysis also revealed lipoprotein phenotypes for significant enrichment, for example 'abnormal circulating lipoprotein level' ($p=1.81 \times 10^{-2}$). Phenotype analysis also revealed 'abnormal blood coagulation' and 'decreased platelet aggregation' ($p=5.81 \times 10^{-8}$ and $p=2.64 \times 10^{-2}$ respectively) in common with the pathway analysis. Our results from the analyses of the 123 genes conserved in four out of five comorbidities were also consistent. The pathway analysis revealed that the 25 most significant pathways were pathways related to the immune system, in particular there was concordance with the results from the eight genes, identifying pathways related to inflammatory responses, interferon and interleukin signaling. In total, the results of all of our analyses show that five comorbidities associated with severe COVID-19 share common physiological aspects including cytokine signaling, blood clotting, and plasma lipoprotein biochemistry.

STAT3 is shared among interleukin signaling pathways that are enriched in COVID-19 comorbidities

To further investigate whether there is a common molecular basis for the interleukin signaling pathway enrichment we examined the Reactome enrichment results using the 123 genes conserved in four out of five comorbidities for interleukin signaling pathways (supplemental table 3). We created gene sets of the shared genes that were in Reactome interleukin pathways that are significantly enriched ($FDR < 0.05$). We used the HiSim graph tool in the GeneWeaver resource to create a graphical view of the genes that are found in the sets. The gene conserved among the largest number of sets is *STAT3* which is found in nine of the eleven significantly enriched interleukin signaling pathways. *IL12B* is shared among four signaling pathways (data not shown).

Genes associated with SARS-CoV-2 infection response and genes shared among COVID-19 comorbidities identify common cytokine signaling pathways and hemostasis

Since we had identified pathways that were common to COVID-19-associated comorbidities, we investigated whether these pathways were also associated with the COVID-19 itself. To answer this question, we created gene sets directly associated with SARS-CoV-2 infection from published literature (Table 2). GeneWeaver

Gene Set GS398287, represents four plasma cytokines that are significantly elevated in patients with severe disease versus patients with mild disease²⁹. GS398329 represents 114 genes associated with angiogenesis or inflammation that were upregulated in COVID-19 postmortem samples⁵. Two gene sets, GS398539 and GS398534, of 119 and 333 genes respectively represent genes that are upregulated in two distinct lung adenocarcinoma epithelial cells infected with SARS-CoV-2; GS398533 with 14 genes represents genes that are upregulated in normal human bronchial epithelium cells infected with SARS-CoV-2; GS398334 represents 587 genes upregulated in post-mortem COVID-19 samples²⁸. We ran Reactome pathway enrichment analysis on each of these sets and determined the enriched pathways that were shared with those identified in the comorbidity analyses (supplemental tables 7-12). Unsurprisingly, GS398329 and GS398287, which were preselected for genes involved in the immune response, were enriched for immune response pathways. GS398329, preselected to be associated with angiogenesis, also showed significant enrichment for the comorbidity pathways associated with platelet biology. All of the data sets showed significant enrichment for signaling mediated by interleukin-4, -10 and -13 (Table 6).

Identification of potential mouse models to study comorbidities and COVID-19 severities

The results of our phenotype analysis using mouse orthologs of shared human genes shows that phenotypic enrichment is consistent with the pathway enrichment using the human genes and is also consistent with pathologies associated with severe COVID-19: blood coagulation, inflammation and cardiovascular pathologies^{36,37,38,39,40,41}. Since mice provide an attractive genetic system for disease modeling, we investigated the phenotypes associated with each of these genes in further detail. Figure 3 shows each of the eight genes shared by all five comorbidities and the phenotype categories that were enriched in this set. Yellow highlighting indicates that mutations in the mouse gene have been annotated to a phenotype of a category that is enriched in the eight shared genes. For example homozygous mice of the genotype *Ctla4*^{tm1Shr}/*Ctla4*^{tm1Shr} display multiple phenotypes that are shared with severe COVID-19: abnormal lung inflammation, abnormal cytokine secretion (interferon secretion) and autoimmune response⁴². *Hmox1*^{tm1Mlee}/*Hmox1*^{tm1Mlee} mice are another example with cardiovascular, immune and liver system phenotypes (<http://www.informatics.jax.org/allele/MGI:2429784>).

Discussion and Summary

COVID-19 is a global health concern. The disease is complex and varies in severity from asymptomatic to lethal⁴³. As our understanding of the disease has progressed, a number of comorbidities associated with the disease have been identified that lead to greater severity. The goal of our work is to identify underlying genetic factors that might explain the mechanism of why certain comorbidities lead to more severe disease. To this end, we studied genetic features of five comorbidities that are associated with severe COVID-19:

Cardiovascular Disease, Diabetes, Hepatitis, Lung Disease, and Kidney Disease. We identified common genes that were associated with each of the comorbidities and the pathways and phenotypes with which they are associated. We compared the results of the comorbidity analysis with genes that were directly associated with SARS-CoV-2 and showed that they shared common pathways involved in the immune response and platelet biology. Our results are encouraging in that these areas of physiology have also been correlated with severe disease. Here we discuss our results in the context of COVID-19 severity.

Our analysis of genes shared among both comorbidities and SARS-CoV-2 infection identified several interleukin signaling pathways that were enriched in both categories. Interleukin-4/-13 and interleukin-10 signaling shared enrichment among the largest number of gene sets we examined, and interleukin-6, interleukin-12 and interleukin-2 shared enrichment between at least one comorbidity set and a set of genes upregulated in patients who died from COVID-19. STAT3 positively regulates the transcription of IL-6, which controls inflammation⁴⁴ and is a downstream signaling player in the IL-6 pathway through the IL6ST protein^{45,46}. IL-12 is produced in response to infection and signals through the JAK-STAT pathway, including STAT3, to induce the proliferation of NK cells and T cells. These cells in turn trigger cytokine signaling including interferon gamma⁴⁷.

One mechanism proposed for the severity of COVID-19 is the coincidence of severe acute respiratory distress triggered by a cytokine related syndrome triggered by the angiotensin signaling pathway⁴⁸. An interesting aspect of this proposal is its action through STAT3, one of the genes that we also found conserved in the comorbidities we studied. Targeting the JAK-STAT pathway has been proposed as a therapeutic approach to

COVID-19². Our results show that *STAT3* is conserved in all five comorbidities we analyzed. This supports the hypothesis that a promising therapeutic strategy to combat severe COVID-19 compounded by preexisting comorbidities may be to target common JAK-STAT pathways.

In addition to immune signaling pathways, we also saw shared enrichment for pathways involved in platelet biology. Platelets are the cells that are responsible for blood clotting⁴⁹. Abnormal clotting has been observed in severe COVID-19 patients and it has been suggested as a complication that leads to more severe disease^{50,51}. Magro *et al.* reported that the abnormal clotting observed in severe COVID-19 patients correlated with activation of the complement pathway⁵². Our results show that *HMOX1*, *APOA1*, *APOE* and two members of the coagulation cascade, *F2* and *F5*, are shared among all five comorbidities we examined. In mice, *Hmox1* deficiency leads to coagulation defects and results in arterial damage due to oxidative stress⁵³. *APOA1* is released during platelet degranulation as part of the platelet secretory granule⁵⁴ (Reactome:R-HSA-482770). *APOA1* levels have also been shown to be significantly decreased in severe COVID-19 patients⁵⁵. *APOE* is a gene that encodes a lipid binding protein involved in cholesterol metabolism⁵⁶. Preliminary evidence suggests that the e4 allele of *APOE* may lead to a higher risk of deep vein thrombosis and the same allele also predicts severe COVID-19^{57,58}. *F2* and *F4* are both involved in the formation of a fibrin clot (Reactome:R-HSA-140877). The complement pathway and coagulation cascade have been shown to interact, tying together inflammation and hemostasis⁵⁹. Additionally in a proteomic study of proteins differentially expressed in the serum of 28 severe COVID-19 patients compared with non-severe COVID-19 patients, Shen *et al.* report that 50 of 93 differentially regulated proteins fall into three categories one of which is platelet degranulation⁵⁵. These results suggest that one of the factors contributing to severe disease in patients with any of the five comorbidities may be due to an underlying genetic mechanism that acts through the hemostatic pathway.

Our results show that genes that are shared among five comorbidities associated with severe COVID-19 identify pathways that are consistent with the pathologies associated with the disease. In our analysis we excluded mouse orthologs that did not correlate 1:1 with human genes to avoid potential skewing of the enrichment analysis by having multiple paralogs over-represented. Despite this, and the exclusion of several potentially important immune system genes such as some histocompatibility genes, our results show that

analysis using mouse orthologs of the shared genes also identifies phenotypes that are consistent with disease pathology. The laboratory mouse provides a tractable system to study the effects of genetic foundations of the comorbidities and severe disease. As mentioned above mice carrying *Ctla4*^{tm1Shr}/*Ctla4*^{tm1Shr} and *Hmox1*^{tm1Mlee}/*Hmox1*^{tm1Mlee} homozygous mutations display multiple phenotypes consistent with severe COVID-19 pathology. Mice engineered for mutations in these genes crossed with mice engineered to carry the human ACE-2 SARS-CoV-2 viral receptor, ICR-Tg(Ace2-ACE2)1Cqin/J, would be a starting point to explore the underlying genetic variants related to comorbidities interact with viral infection⁶⁰. Humanized mice, like the *ApoE*^{tm3(APOE*4)Mae} strain, which carries the human E3 variant mentioned above as being implicated in both severe disease and an underlying thrombosis pathology, could be used in conjunction with ICR-Tg(Ace2-ACE2)1Cqin/J to study the effects of the human variant on viral infection. A comprehensive resource for using the mouse as a model system for COVID-19 research is maintained by the Mouse Genome Informatics Group [<http://www.informatics.jax.org/mgihome/other/coronavirus.shtml>].

In this study we have used a bioinformatics approach to interrogate genes associated with five COVID-19 comorbidities that correlate with severe disease. Using genes that have been annotated to these comorbidities in the MeSH or HPO resource we have shown that genes are shared among the comorbidities and that shared genes are enriched for pathways that could be the genetic basis for the pathologies observed with severe COVID-19, specifically our results suggest that the interrelated pathways of hemostasis and inflammation may be key players in understanding the severity of comorbidities with COVID-19^{61,62}. Our studies provide a gateway to understand how host genetics interacts with and influences the consequences of viral infection. Our knowledge about COVID-19 continues to grow at a rapid rate and future work will entail the examination of additional comorbidities, more specific comorbidities, a wider survey of genes beyond our initial seed set from MeSH and HPO. As we learn more about correlations between individual comorbidities and disease pathologies, we may be able to identify specific pathway/comorbidity combinations that can be used to inform us about treatment decisions. Our work also provides an entry point into an experimental system using the laboratory mouse to manipulate host genetics and to study its subsequent effect on the pathology of viral infection.

Data availability

All gene sets generated during and analyzed during the current study are based on data published in peer-reviewed papers, are available in the public GeneWeaver repository [www.geneweaver.org] and are accessible using the gene set identifiers given in the text (e.g. GS398287). Results data generated during this study are included in this published article and its supplementary files.

References

1. Wang, C., Horby, P. W., Hayden, F. G. & Gao, G. F. A novel coronavirus outbreak of global health concern. *Lancet* **395**, 470–473 (2020).
2. Luo, W. *et al.* Targeting JAK-STAT Signaling to Control Cytokine Release Syndrome in COVID-19. *Trends Pharmacol. Sci.* **41**, 531–543 (2020).
3. Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **395**, 497–506 (2020).
4. Xu, S., Chen, M. & Weng, J. COVID-19 and Kawasaki disease in children. *Pharmacol. Res.* **159**, 104951 (2020).
5. Ackermann, M. *et al.* Pulmonary Vascular Endothelialitis, Thrombosis, and Angiogenesis in Covid-19. *N. Engl. J. Med.* **383**, 120–128 (2020).
6. Terpos, E. *et al.* Hematological findings and complications of COVID-19. *Am. J. Hematol.* **95**, 834–847 (2020).
7. Gauthier, J. & Turtle, C. J. Insights into cytokine release syndrome and neurotoxicity after CD19-specific CAR-T cell therapy. *Curr Res Transl Med* **66**, 50–52 (2018).
8. Pearce, L., Davidson, S. M. & Yellon, D. M. The cytokine storm of COVID-19: a spotlight on prevention and protection. *Expert Opin. Ther. Targets* 1–8 (2020) doi:10.1080/14728222.2020.1783243.
9. Obi, A. T. *et al.* Empirical systemic anticoagulation is associated with decreased venous thromboembolism in critically ill influenza A H1N1 acute respiratory distress syndrome patients. *J Vasc Surg Venous Lymphat Disord* **7**, 317–324 (2019).
10. Richardson, S. *et al.* Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA* **323**, 2052–2059 (2020).
11. Cheng, Y. *et al.* Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney Int* **97**, 829–838 (2020).
12. CDC COVID-19 Response Team. Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions Among Patients with Coronavirus Disease 2019 - United States, February 12-March 28, 2020. *MMWR Morb. Mortal. Wkly. Rep.* **69**, 382–386 (2020).
13. Chen, T. *et al.* Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ* **368**, (2020).
14. Guan, W. *et al.* Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. *Eur Respir J* **55**, (2020).
15. Emami, A., Javanmardi, F., Pirbonyeh, N. & Akbari, A. Prevalence of Underlying Diseases in Hospitalized Patients with COVID-19: a Systematic Review and Meta-Analysis. *Arch Acad Emerg Med* **8**, (2020).
16. Korean Society of Infectious Diseases and Korea Centers for Disease Control and Prevention. Analysis on 54 Mortality Cases of Coronavirus Disease 2019 in the Republic of Korea from January 19 to March 10, 2020. *J. Korean Med. Sci.* **35**, e132 (2020).

17. Rodriguez-Morales, A. J. *et al.* Clinical, laboratory and imaging features of COVID-19: A systematic review and meta-analysis. *Travel Med Infect Dis* **34**, 101623 (2020).
18. Koutsakos, M. & Kedzierska, K. A race to determine what drives COVID-19 severity. *Nature* **583**, 366–368 (2020).
19. Muniyappa, R. & Gubbi, S. COVID-19 pandemic, coronaviruses, and diabetes mellitus. *Am. J. Physiol. Endocrinol. Metab.* **318**, E736–E741 (2020).
20. Zhu, L. *et al.* Association of Blood Glucose Control and Outcomes in Patients with COVID-19 and Pre-existing Type 2 Diabetes. *Cell Metab.* **31**, 1068-1077.e3 (2020).
21. Tuitou, E., Duchi, S. & Natsheh, H. A new nanovesicular system for nasal drug administration. *Int J Pharm* **580**, 119243 (2020).
22. South, A. M., Diz, D. I. & Chappell, M. C. COVID-19, ACE2, and the cardiovascular consequences. *Am. J. Physiol. Heart Circ. Physiol.* **318**, H1084–H1090 (2020).
23. Bansal, M. Cardiovascular disease and COVID-19. *Diabetes Metab Syndr* **14**, 247–250 (2020).
24. Bult, C. J. *et al.* Mouse Genome Database (MGD) 2019. *Nucleic Acids Res.* **47**, D801–D806 (2019).
25. Körner, R. W., Majjouti, M., Alcazar, M. A. A. & Mahabir, E. Of Mice and Men: The Coronavirus MHV and Mouse Models as a Translational Approach to Understand SARS-CoV-2. *Viruses* **12**, (2020).
26. Baker, E., Bubier, J. A., Reynolds, T., Langston, M. A. & Chesler, E. J. GeneWeaver: data driven alignment of cross-species genomics in biology and disease. *Nucleic Acids Res.* **44**, D555-559 (2016).
27. Köhler, S. *et al.* Expansion of the Human Phenotype Ontology (HPO) knowledge base and resources. *Nucleic Acids Res.* **47**, D1018–D1027 (2019).
28. Blanco-Melo, D. *et al.* Imbalanced Host Response to SARS-CoV-2 Drives Development of COVID-19. *Cell* **181**, 1036-1045.e9 (2020).
29. Chen, G. *et al.* Clinical and immunological features of severe and moderate coronavirus disease 2019. *J. Clin. Invest.* **130**, 2620–2629 (2020).
30. Richardson, J. E. & Bult, C. J. Visual annotation display (VLAD): a tool for finding functional themes in lists of genes. *Mamm. Genome* **26**, 567–573 (2015).
31. Beninato, M. & Spencer, R. F. The cholinergic innervation of the rat substantia nigra: a light and electron microscopic immunohistochemical study. *Exp Brain Res* **72**, 178–184 (1988).
32. Smith, C. L. *et al.* Mouse Genome Database (MGD)-2018: knowledgebase for the laboratory mouse. *Nucleic Acids Res.* **46**, D836–D842 (2018).
33. Eppig, J. T. *et al.* Mouse Genome Database: From sequence to phenotypes and disease models. *Genesis* **53**, 458–473 (2015).
34. Alliance of Genome Resources Consortium. Alliance of Genome Resources Portal: unified model organism research platform. *Nucleic Acids Res.* **48**, D650–D658 (2020).
35. Jassal, B. *et al.* The reactome pathway knowledgebase. *Nucleic Acids Res.* **48**, D498–D503 (2020).
36. Spiezia, L. *et al.* COVID-19-Related Severe Hypercoagulability in Patients Admitted to Intensive Care Unit for Acute Respiratory Failure. *Thromb. Haemost.* **120**, 998–1000 (2020).
37. Lin, J. *et al.* COVID-19 and coagulation dysfunction in adults: A systematic review and meta-analysis. *J. Med. Virol.* (2020) doi:10.1002/jmv.26346.
38. Tay, M. Z., Poh, C. M., Rénia, L., MacAry, P. A. & Ng, L. F. P. The trinity of COVID-19: immunity, inflammation and intervention. *Nat. Rev. Immunol.* **20**, 363–374 (2020).
39. Merad, M. & Martin, J. C. Pathological inflammation in patients with COVID-19: a key role for monocytes and macrophages. *Nat. Rev. Immunol.* **20**, 355–362 (2020).
40. Clerkin, K. J. *et al.* COVID-19 and Cardiovascular Disease. *Circulation* **141**, 1648–1655 (2020).

41. Fried, J. A. *et al.* The Variety of Cardiovascular Presentations of COVID-19. *Circulation* **141**, 1930–1936 (2020).
42. Tivol, E. A. *et al.* Loss of CTLA-4 leads to massive lymphoproliferation and fatal multiorgan tissue destruction, revealing a critical negative regulatory role of CTLA-4. *Immunity* **3**, 541–547 (1995).
43. Velavan, T. P. & Meyer, C. G. The COVID-19 epidemic. *Trop. Med. Int. Health* **25**, 278–280 (2020).
44. Tanaka, T., Narazaki, M. & Kishimoto, T. IL-6 in inflammation, immunity, and disease. *Cold Spring Harb Perspect Biol* **6**, a016295 (2014).
45. Breslow, E., Co, R. T., Hanna, P. & Laborde, T. Influence of neurophysin residues 1-8 on the optical activity of neurophysin-peptide complexes. Direct evidence that the 1-8 sequence alters the environment of bound peptide. *Int. J. Pept. Protein Res.* **34**, 21–27 (1989).
46. Greenhill, C. J. *et al.* IL-6 trans-signaling modulates TLR4-dependent inflammatory responses via STAT3. *J. Immunol.* **186**, 1199–1208 (2011).
47. Trinchieri, G. Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nat. Rev. Immunol.* **3**, 133–146 (2003).
48. Hirano, T. & Murakami, M. COVID-19: A New Virus, but a Familiar Receptor and Cytokine Release Syndrome. *Immunity* **52**, 731–733 (2020).
49. Holinstat, M. Normal platelet function. *Cancer Metastasis Rev.* **36**, 195–198 (2017).
50. Helms, J. *et al.* High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study. *Intensive Care Med* **46**, 1089–1098 (2020).
51. Al-Samkari, H. *et al.* COVID-19 and coagulation: bleeding and thrombotic manifestations of SARS-CoV-2 infection. *Blood* **136**, 489–500 (2020).
52. Magro, C. *et al.* Complement associated microvascular injury and thrombosis in the pathogenesis of severe COVID-19 infection: A report of five cases. *Transl Res* **220**, 1–13 (2020).
53. True, A. L. *et al.* Heme oxygenase-1 deficiency accelerates formation of arterial thrombosis through oxidative damage to the endothelium, which is rescued by inhaled carbon monoxide. *Circ. Res.* **101**, 893–901 (2007).
54. Coppinger, J. A. *et al.* Characterization of the proteins released from activated platelets leads to localization of novel platelet proteins in human atherosclerotic lesions. *Blood* **103**, 2096–2104 (2004).
55. Shen, B. *et al.* Proteomic and Metabolomic Characterization of COVID-19 Patient Sera. *Cell* **182**, 59-72.e15 (2020).
56. Martínez-Martínez, A. B. *et al.* Beyond the CNS: The many peripheral roles of APOE. *Neurobiol. Dis.* **138**, 104809 (2020).
57. Zhu, S., Wang, Z., Wu, X., Shu, Y. & Lu, D. Apolipoprotein E polymorphism is associated with lower extremity deep venous thrombosis: color-flow Doppler ultrasound evaluation. *Lipids Health Dis* **13**, 21 (2014).
58. Kuo, C.-L. *et al.* APOE e4 genotype predicts severe COVID-19 in the UK Biobank community cohort. *J. Gerontol. A Biol. Sci. Med. Sci.* (2020) doi:10.1093/gerona/glaa131.
59. Oikonomopoulou, K., Ricklin, D., Ward, P. A. & Lambris, J. D. Interactions between coagulation and complement--their role in inflammation. *Semin Immunopathol* **34**, 151–165 (2012).
60. Bao, L. *et al.* The pathogenicity of SARS-CoV-2 in hACE2 transgenic mice. *Nature* **583**, 830–833 (2020).
61. Verhamme, P. & Hoylaerts, M. F. Hemostasis and inflammation: two of a kind? *Thromb J* **7**, 15 (2009).
62. Koupenova, M., Clancy, L., Corkrey, H. A. & Freedman, J. E. Circulating Platelets as Mediators of Immunity, Inflammation, and Thrombosis. *Circ. Res.* **122**, 337–351 (2018).

Acknowledgements

This work was funded by NIH grants to the Mouse Genome Database (NHGRI U41 HG000330), the Jackson Laboratory Center for Precision Genetics (OD U54 OD020351) and GeneWeaver (NIAID RO1 AA18776). The authors would like to thank Dr. Peter D'Eustachio and Dr. Laurens Wilming for their critical reading of the manuscript.

Author contributions

DPH and JAB conceived the study and were in charge of overall direction and planning. DPH and MED carried out the implementation and performed the computations. GM, MSM, and EJC contributed to the interpretation of the results. DPH took the lead in writing the manuscript with support from MED. All authors discussed the results and contributed to the final manuscript.

Competing interests

All authors declare that they have no competing interests.

Author affiliations

Mary E. Dolan, David P. Hill, Gaurab Mukherjee, Monica S. McAndrews, Elissa J. Chesler, Judith A Blake
The Jackson Laboratory, Bar Harbor, ME 04609, USA

Materials & Correspondence

Correspondence and material requests should be addressed to Mary E. Dolan.

Tables

Table 1: SARS-CoV-2 (COVID-19) comorbidity-related gene sets

Table 2: SARS-CoV-2 (COVID-19) Gene Sets

Table 3. Genes shared by COVID-19 comorbidities. This table shows the genes that were annotated to four out of five comorbidities that are associated with COVID-19 severity. Genes in **bold** are annotated to all five comorbidities.

Table 4. Significantly enriched phenotype categories. Top significantly enriched phenotype categories identified by VLAD analysis for the set of eight genes shared by all five comorbidities.

Table 5. Significantly enriched phenotype categories. Top significantly enriched phenotype categories identified by VLAD analysis for the set of 123 genes shared by four of five comorbidities.

Table 6. Comparison of pathway enrichment false discovery rates (FDR). FDRs for pathways with FDR ≤ 0.05 (shown in light red) for at least one of the comorbidity sets and for at least one of the COVID-19 related gene sets are shown.

Table 1: SARS-CoV-2 (COVID-19) comorbidity-related gene sets

Gene Set Identifier	Gene count	Gene Set Title	Gene Set Description/metadata
GS380453	219	Kidney Disease	The Boolean Algebra tool was used to find the Union of 5 sets: GS380448, GS380449, GS380450, GS380451, GS380452. Kidney disease has been reported as a comorbidity factor in the disease (COVID-19) caused by SARS-CoV-2 infection: PMID:32247631, PMID:32233161, PMID:32232218.
GS380473	267	Hepatitis-MeSH:D006505	This gene set represents the genes in GS237945 queried on April 16, 2020 which was derived from the gene2mesh procedure. GS237945 was updated on 2019-01-07. Liver disease has been reported as a comorbidity factor in the disease (COVID-19) caused by SARS-CoV-2 infection: PMID: 32179124, PMID:32233161.
GS380478	2550	Cardiovascular Diseases-MeSH:D002318	This gene set represents the genes in GS235830 queried on April 17, 2020 which was derived from the gene2mesh procedure. GS235830 was updated on 2019-01-07. Cardiovascular disease has been reported as a comorbidity factor in the disease (COVID-19) caused by SARS-CoV-2 infection: PMID: 32179124, PMID:32267833, PMID:32232218, PMID:32217556.
GS380470	1003	Diabetes Mellitus, type 2-MeSH:D003924	This gene set represents the genes in GS242930 queried on April 16, 2020 which was derived from the gene2mesh procedure. GS242930 was updated on 2019-01-07. Diabetes has been reported as a comorbidity factor in the disease (COVID-19) caused by SARS-CoV-2 infection: PMID: 32179124, PMID:32233161, PMID:32267833, PMID:32232218, PMID:32217650, PMID:32345579.
GS380479	1927	Lung Diseases-MeSH:D008171	This gene set represents the genes in GS245016 queried on April 16, 2020 which was derived from the gene2mesh procedure. GS245016 was updated on 2019-01-07. Lung (pulmonological) disease has been reported as a comorbidity factor in the disease (COVID-19) caused by SARS-CoV-2 infection: PMID:32233161.

Table 2: SARS-CoV-2 (COVID-19) Gene Sets

Gene Set Identifier	Gene Count	Gene Set Title	Gene Set Description/metadata
GS398329	119	Up regulated angiogenesis and inflammation genes in lungs from patients who died from COVID-19	This gene set describes genes that are up-regulated in lungs from patients who died from COVID-19. COVID-19 is the disease caused by SARS-CoV-2 virus. Note that this expression analysis includes only the angiogenesis-associated and inflammation-associated genes available on NanoString panels. The authors define up-regulated as those genes that show a (FDR) of ≤ 0.05 . These data are from the publication (angiogenesis) and supplementary (inflammation) materials associated with the publication. PMID:32437596
GS398539	102	Up regulated genes in host transcriptional response to SARS-CoV-2 in Human adenocarcinomic alveolar basal epithelial (A549) cells	This gene set describes genes that are up-regulated by the host transcriptional response to SARS-CoV-2 infection in human adenocarcinomic alveolar basal epithelial (A549) cells. COVID-19 is the disease caused by SARS-CoV-2 virus. We define up-regulated as those genes that show a (log 2 fold change) of ≥ 2 . These data are from the supplementary materials associated with the publication. PMID:32416070
GS398534	333	Up regulated genes in host transcriptional response to SARS-CoV-2 in Human lung adenocarcinoma epithelial (Calu3) cells	This gene set describes genes that are up-regulated by the host transcriptional response to SARS-CoV-2 infection in human lung adenocarcinoma epithelial cells derived from pleural effusion (Calu3). COVID-19 is the disease caused by SARS-CoV-2 virus. We define up-regulated as those genes that show a (log 2 fold change) of ≥ 2 . These data are from the supplementary materials associated with the publication. PMID:32416070
GS398533	14	Up regulated genes in host transcriptional response to SARS-CoV-2 in normal human bronchial epithelium (NHBE) cells	This gene set describes genes that are up-regulated by the host transcriptional response to SARS-CoV-2 infection in normal human bronchial epithelium cells (NHBE). COVID-19 is the disease caused by SARS-CoV-2 virus. We define up-regulated as those genes that show a (log 2 fold change) of ≥ 2 . These data are from the supplementary materials associated with the publication. PMID:32416070
GS398334	587	Up regulated genes in post-mortem lung samples from COVID-19-positive patients	This gene set describes genes that are up-regulated in post-mortem lung samples from COVID-19-positive patients relative to biopsied healthy lung tissue from uninfected individuals. COVID-19 is the disease caused by SARS-CoV-2 virus. We define up-regulated as those genes that show a (log 2 fold change) of ≥ 2 . These data are from the supplementary materials associated with the

			publication. Note: the following HGNC id is part of this data set but was not recognized HGNC:13378. PMID:32416070
GS398287	4	Genes that are overexpressed in severe compared to mild cases of Covid-19	People with severe cases of Covid-19 express these proteins at significantly higher levels than people with mild cases of Covid-19. Data from Figure 2 of the paper: plasma cytokine levels in patients with COVID-19. PMID:32217835

<i>ACE</i>	<i>CHI3L1</i>	<i>FGA</i>	<i>IL10</i>	<i>MIF</i>	<i>SERPINA1</i>	<i>VDR</i>
<i>ADIPOQ</i>	<i>CTGF</i>	<i>FN1</i>	<i>IL12A</i>	<i>MMP1</i>	<i>SLC6A4</i>	<i>VEGFA</i>
<i>AGT</i>	<i>CTLA4</i>	<i>FOXC2</i>	<i>IL12B</i>	<i>MMP2</i>	<i>SOCS1</i>	<i>XRCC1</i>
<i>AGTR1</i>	<i>CTNNB1</i>	<i>GSTM1</i>	<i>IL15</i>	<i>MMP3</i>	<i>SOCS3</i>	
<i>AKT1</i>	<i>CX3CL1</i>	<i>GSTP1</i>	<i>IL18</i>	<i>MMP9</i>	<i>SOD2</i>	
<i>APOA1</i>	<i>CXCL10</i>	<i>HAMP</i>	<i>IL1A</i>	<i>MPO</i>	<i>SPP1</i>	
<i>APOE</i>	<i>CXCL12</i>	<i>HFE</i>	<i>IL1B</i>	<i>MTHFR</i>	<i>STAT3</i>	
<i>AR</i>	<i>CXCL8</i>	<i>HGF</i>	<i>IL1RN</i>	<i>MTOR</i>	<i>TERT</i>	
<i>ARG1</i>	<i>CYP2D6</i>	<i>HLA-B</i>	<i>IL2</i>	<i>NFE2L2</i>	<i>TGFB1</i>	
<i>B2M</i>	<i>CYP2E1</i>	<i>HLA-DQA1</i>	<i>IL4</i>	<i>NFKB1</i>	<i>THBD</i>	
<i>BCL2</i>	<i>CYP3A4</i>	<i>HLA-DQB1</i>	<i>IL4R</i>	<i>NOS2</i>	<i>TIMP1</i>	
<i>C3</i>	<i>EGF</i>	<i>HLA-DRA</i>	<i>IL6</i>	<i>NOS3</i>	<i>TIMP2</i>	
<i>CCL2</i>	<i>EGFR</i>	<i>HLA-DRB1</i>	<i>IRS1</i>	<i>NOTCH2</i>	<i>TLR2</i>	
<i>CCL5</i>	<i>ENG</i>	<i>HMOX1</i>	<i>ITGB3</i>	<i>NPPB</i>	<i>TLR4</i>	
<i>CCR2</i>	<i>EPO</i>	<i>HSPA5</i>	<i>LCAT</i>	<i>PIK3CA</i>	<i>TLR9</i>	
<i>CCR5</i>	<i>ESR1</i>	<i>ICAM1</i>	<i>LDLR</i>	<i>PON1</i>	<i>TNF</i>	
<i>CD40</i>	<i>F2</i>	<i>IFNA1</i>	<i>LEP</i>	<i>PRKDC</i>	<i>TNFRSF10B</i>	
<i>CDKN1A</i>	<i>F5</i>	<i>IFNG</i>	<i>LTA</i>	<i>PTGS2</i>	<i>TNFRSF1A</i>	
<i>CDKN2A</i>	<i>FAS</i>	<i>IGF1</i>	<i>MAPK1</i>	<i>PTPN22</i>	<i>TNFRSF1B</i>	
<i>CFH</i>	<i>FASLG</i>	<i>IGF2</i>	<i>MBL2</i>	<i>SCARB1</i>	<i>TP53</i>	

Table 3. Genes shared by COVID-19 comorbidities. This table shows the genes that were annotated to four out of five comorbidities that are associated with COVID-19 severity. Genes in **bold** are annotated to all five comorbidities.

Phenotype Category	Gene Count	P-value range
T Cell Related	5	1.6x10e-06 to 3.9x10e-08
Autoimmune Response	5	1.7x10e-6
Interferon gamma secretion	4	3.7x10e-6
Lung Inflammation	4	1.7x10e-6
Susceptibility to infection	5	8.9x10e-7
Blood Coagulation	5	5.8xe10-8
Cardiovascular physiology	7	1.6xe10-6

Table 4. Significantly enriched phenotype categories. Top significantly enriched phenotype categories identified by VLAD analysis, showing how many of the eight genes shared among all five comorbidities are annotated to each phenotype category. T Cell related phenotypes included ‘increased CD-4 positive, alpha beta T cell number’ (p=3.9x10e-8) and ‘increased T-helper Cell number’ (p=1.6x10e-06). (Complete list of enriched phenotypes available supplemental table 1)

Phenotype Category	Gene Count	P-value range
Innate immunity	49	3.1x10e-43
Inflammatory response	66	7.0x10e-43
Leukocyte Physiology	65	5.7x10e-42
Cytokine levels	43	1.9x10e-41
Susceptibility to infection	48	9.3x10e-40
Mononuclear cell morphology	68	1.1x10-e39
Phagocyte morphology	52	3.8x10-e39
Leukocyte number	69	2.0x10e-40
Leukocyte physiology	66	8.2x10e-39
Injury response	54	2.3x10e-49
Blood vessel morphology	60	1.4x10e-39
Professional antigen presenting cell morphology	50	4.5x10e-40

Table 5. Significantly enriched phenotype categories. Top significantly enriched phenotype categories identified by VLAD analysis, showing the number of genes from the set of 107 mouse orthologs shared among four out of five comorbidities annotated to each phenotype category. (Complete list of enriched phenotypes available supplemental table 2)

pw_id	pw_name	comorbid8	comorbid123	A549	Calu3	NHBE	postmortem	PMID:32437596	PMID:32217835	score
R-HSA-168256	Immune System	3.133E-02	5.773E-15	8.450E-02	1.832E-14	1.485E-06	2.376E-14	1.343E-14	1.259E-02	7
R-HSA-1280215	Cytokine Signaling in Immune system	2.870E-02	5.773E-15	9.169E-03	1.832E-14	1.422E-11	2.376E-14	1.343E-14	5.990E-04	8
R-HSA-449147	Signaling by Interleukins	3.473E-02	5.773E-15	3.453E-04	6.084E-14	1.093E-10	4.014E-09	1.343E-14	4.840E-05	8
R-HSA-6783783	Interleukin-10 signaling	7.900E-02	5.773E-15	3.953E-11	1.832E-14	1.144E-14	7.138E-11	1.343E-14	1.340E-05	7
R-HSA-6785807	Interleukin-4 and Interleukin-13 signaling	1.153E-02	5.773E-15	5.224E-05	6.880E-09	1.334E-05	2.228E-03	1.343E-14	4.840E-05	8
R-HSA-6783589	Interleukin-6 family signaling	3.473E-02	8.082E-02	2.852E-01	2.485E-01	6.745E-02	6.313E-01	2.274E-02	1.601E-02	3
R-HSA-1059683	Interleukin-6 signaling	3.220E-02	3.448E-02	2.852E-01	2.526E-01	6.745E-02	6.313E-01	5.788E-03		3
R-HSA-913531	Interferon Signaling	5.346E-02	5.773E-15	4.843E-01	1.832E-14	1.829E-02	2.376E-14	9.975E-02		4
R-HSA-909733	Interferon alpha/beta signaling		5.773E-15		1.832E-14	1.982E-03	2.376E-14	7.299E-02		4
R-HSA-877300	Interferon gamma signaling	3.473E-02	5.773E-15	2.852E-01	4.546E-13		1.970E-08	5.761E-01		4
R-HSA-168249	Innate Immune System	3.662E-01	4.366E-02	2.852E-01	4.483E-01	8.976E-01	3.656E-12	4.336E-02		3
R-HSA-109582	Hemostasis	4.045E-02	3.400E-03	2.852E-01	9.355E-01		6.313E-01	2.762E-09		3
R-HSA-76002	Platelet activation, signaling and aggregation	1.903E-02	3.484E-05	2.852E-01	8.186E-01		6.313E-01	2.125E-11		3
R-HSA-76005	Response to elevated platelet cytosolic Ca2+	3.133E-02	9.286E-07	2.852E-01	4.692E-01		6.313E-01	1.848E-09		3
R-HSA-114608	Platelet degranulation	3.133E-02	4.837E-07	2.852E-01	6.213E-01		6.313E-01	1.059E-09		3
R-HSA-76009	Platelet Aggregation (Plug Formation)	4.940E-02	2.912E-02	2.852E-01	8.100E-01		6.313E-01	1.544E-02		3

Table 6. Comparison of pathway enrichment false discovery rates (FDR) for pathways with FDR ≤ 0.05 (shown in light red) for at least one of the comorbidity sets and for at least one of the COVID-19 related gene sets: comorbid8 column displays FDR for comorbidity set for all five comorbidities; comorbid123 for comorbidity set for four of the five comorbidities; A549 for up regulated genes in host transcriptional response to SARS-CoV-2 in human A549 lung alveolar cells; Calu3 for up regulated genes in Calu3 human lung adenocarcinoma epithelial cells; NHBE for up regulated genes in normal human bronchial epithelium (NHBE) cells; postmortem for up regulated genes in post-mortem lung samples from COVID-19-positive patients; PMID:32437596 for up regulated angiogenesis and inflammation genes in lungs from patients who died from COVID-19; PMID:32217835 for genes that are overexpressed in severe compared to mild cases of Covid-19. The score column gives the number of sets with FDR meeting our criteria. Shaded in yellow are the pathways significantly enriched for both the comorbidity sets and several COVID-19 sets based on highest scores. The full table displaying 28 pathways meeting our criteria is given in supplemental material. (Complete list of enriched pathways available supplemental table 4; enriched pathway details for each gene set available supplemental tables 5-12)

Figures

Figure 1. The top 25 most significantly enriched pathways involving the eight genes shared among all five comorbidities. Pathways that are similar or directly related in the Reactome knowledgebase are color coded. Yellow: lipoprotein-related processes; peach: scavenger receptor pathways; blue: blood clotting; green; retinoid-related pathways; grey; signaling through STAT3.

Figure 2. The top 25 most significantly enriched pathways involving the 123 genes shared among four out of five comorbidities. Pathways that are similar or directly related in the Reactome knowledgebase are color coded. Yellow: adaptive immune system pathways; blue: interleukin signaling pathways; green; interferon signaling pathways.

Figure 3. Phenotype Enrichment for the Eight Genes Shared by All Five Comorbidities. Cells shaded in yellow indicate that there is a mouse model in the MGI resource that has been studied for the specific phenotype.

Pathway name	FDR	Genes in Pathway
Regulation of Insulin-like Growth Factor (IGF) transport and uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs)	1.560E-04	<i>F5; APOA1; APOE; F2</i>
Chylomicron remodeling	1.066E-03	<i>APOA1; APOE</i>
Chylomicron assembly	1.066E-03	<i>APOA1; APOE</i>
HDL remodeling	1.066E-03	<i>APOA1; APOE</i>
Post-translational protein phosphorylation	1.477E-03	<i>F5; APOA1; APOE</i>
Plasma lipoprotein assembly	1.779E-03	<i>APOA1; APOE</i>
Scavenging by Class A Receptors	1.779E-03	<i>APOA1; APOE</i>
Common Pathway of Fibrin Clot Formation	2.072E-03	<i>F5; F2</i>
Plasma lipoprotein remodeling	3.715E-03	<i>APOA1; APOE</i>
Plasma lipoprotein clearance	3.715E-03	<i>APOA1; APOE</i>
Formation of Fibrin Clot (Clotting Cascade)	4.854E-03	<i>F5; F2</i>
Retinoid metabolism and transport	4.934E-03	<i>APOA1; APOE</i>
Metabolism of fat-soluble vitamins	5.375E-03	<i>APOA1; APOE</i>
Metabolism of proteins	6.453E-03	<i>F5; B2M; APOA1; APOE; STAT3;</i>
Platelet activation, signaling and aggregation	6.453E-03	<i>F5; APOA1; F2</i>
Plasma lipoprotein assembly, remodeling, and clearance	9.811E-03	<i>APOA1; APOE</i>
Amyloid fiber formation	1.123E-02	<i>B2M; APOA1</i>
Defective ABCA1 causes Tangier disease	1.123E-02	<i>APOA1</i>
Visual phototransduction	1.473E-02	<i>APOA1; APOE</i>
MET activates STAT3	1.473E-02	<i>STAT3</i>
Signalling to STAT3	1.473E-02	<i>STAT3</i>
Interleukin-4 and Interleukin-13 signaling	1.683E-02	<i>HMOX1; STAT3</i>
PTK6 Activates STAT3	1.683E-02	<i>STAT3</i>
Platelet degranulation	1.753E-02	<i>F5; APOA1</i>
Binding and Uptake of Ligands by Scavenger Receptors	1.753E-02	<i>APOA1; APOE</i>

Figure 1. The top 25 most significantly enriched pathways involving the eight genes shared among all five comorbidities. Pathways that are similar or directly related in the Reactome knowledgebase are color coded. Yellow: lipoprotein-related processes; peach: scavenger receptor pathways; blue: blood clotting; green; retinoid-related pathways; grey; signaling through STAT3.

Pathway name	FDR	Gene Count
Translocation of ZAP-70 to Immunological synapse	4.770E-15	5
Interleukin-10 signaling	4.770E-15	21
Phosphorylation of CD3 and TCR zeta chains	4.770E-15	5
Interleukin-4 and Interleukin-13 signaling	4.770E-15	33
Antigen Presentation: Folding, assembly and peptide loading of class I MHC	4.770E-15	3
Endosomal/Vacuolar pathway	4.770E-15	2
Interferon gamma signaling	4.770E-15	10
Interferon alpha/beta signaling	4.770E-15	4
Class I MHC mediated antigen processing & presentation	4.770E-15	8
Signaling by Interleukins	4.770E-15	49
Cytokine Signaling in Immune system	4.770E-15	62
Costimulation by the CD28 family	4.770E-15	8
ER-Phagosome pathway	4.770E-15	4
Adaptive Immune System	4.770E-15	21
Interferon Signaling	4.770E-15	10
Immunoregulatory interactions between a Lymphoid and a non-Lymphoid cell	4.770E-15	5
Immune System	4.770E-15	81
Antigen processing-Cross presentation	4.770E-15	4
PD-1 signaling	4.770E-15	4
Generation of second messenger molecules	4.770E-15	4
Downstream TCR signaling	2.840E-12	6
TCR signaling	5.490E-12	7
MHC class II antigen presentation	7.010E-09	4
Extra-nuclear estrogen signaling	2.730E-08	12
Regulation of Insulin-like Growth Factor (IGF) transport and uptake by Insulin-like Growth Factor Binding Proteins (IGFBPs)	6.850E-08	15

Figure 2. The top 25 most significantly enriched pathways involving the 123 genes shared among four out of five comorbidities. Pathways that are similar or directly related in the Reactome knowledgebase are color coded. Yellow: adaptive immune system pathways; blue: interleukin signaling pathways; green; interferon signaling pathways.

	<i>APOA1</i>	<i>APOE</i>	<i>B2M</i>	<i>CTLA4</i>	<i>F2</i>	<i>F5</i>	<i>HMOX1</i>	<i>STAT3</i>
T cell related		X	X	X			X	X
Autoimmune response		X	X	X	X		X	
Interferon secretion		X		X			X	X
Lung inflammation		X		X			X	X
Susceptibility to infection		X	X		X		X	X
Blood coagulation		X			X	X	X	X
Abnormal cardiovascular system physiology	X	X		X	X	X	X	X

Figure 3. Phenotype Enrichment for the Eight Genes Shared by All Five Comorbidities. Cells shaded in yellow indicate that there is a mouse model in the MGI resource that has been studied for the specific phenotype.