Title: Host evolutionary history and ecological context modulate the adaptive potential of the

microbiome

Authors: Lucas P. Henry[1,2,3], Michael Fernandez[1,2], Andrew Webb[1,2], Julien F. Ayroles[1,2]

Affiliations:

[1]Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton NJ, USA.

[2]Ecology and Evolutionary Biology Department, Princeton University, Princeton NJ, USA.

[3]Center for Genomics and Systems Biology, New York University, New York NY, USA.

*Correspondence to: lph260@nyu.edu or jayroles@princeton.edu

Abstract:

Can the microbiome serve as a reservoir of adaptive potential for hosts? To address this

question, we leveraged ~150 generations of experimental evolution in *Drosophila melanogaster*

in a stressful, high-sugar (HS) diet. We performed a fully reciprocal transplant experiment using

the control and evolved HS bacteria. If the microbiome confers benefits to hosts, then transplant

recipients should gain fitness benefits compared to controls. Interestingly, we found that benefits

do exist, but mismatches between fly evolution and microbiome exerted fitness costs by slowing

development and reducing fecundity, especially in the stressful HS diet. The dominant HS

bacteria (*Acetobacter pasteurianus*) uniquely encoded several genes to enable uric acid

degradation, mediating the toxic effects of uric acid accumulation due to the HS diet for flies.

Our study demonstrates that host genotype x microbiome x environment interactions have

substantial effects on host phenotype, highlighting how host evolution and ecological context

together shape the adaptive potential of the microbiome.

Competing interests: The authors declare no competing interests.

INTRODUCTION

The microbiome has profound effects on many aspects of organismal biology. From modulating physiology [1] to pathogen protection [2] to social interactions [3], the microbiome is frequently implicated as a key regulator of many host phenotypes--but can the microbiome also influence host adaptation? For hosts where microbes are tightly controlled and faithfully transmitted across generations, microbes can have clear influence on host evolution [4–6]. For example, the acquisition of nutrition-provisioning symbionts enables the ecological expansion of sap-feeding insects [7]. For many eukaryotes, though, their microbiomes are environmentally acquired or through other weakly controlled transmission mechanisms [5]. Weak control over microbial transmission disrupts the links between the microbiome and host over generations. Without these links, the microbiome may have limited effects on host evolution [4, 6, 8]. Microbiomes may still have the potential to influence host evolution because environmentally acquired microbes must survive and adapt to external environments. In doing so, microbiomes may acquire beneficial traits that hosts can leverage. Thus, microbiomes have the potential to precede and facilitate rapid host adaptation, but this process is poorly understood.

Environmentally acquired microbiomes appear to facilitate rapid host adaptation in several systems. For example, bean bugs acquire pesticide resistance from their soil-dwelling *Burkholderia* symbiont [9]. The gut microbiome of woodrats facilitates survival on toxic plant diets [10]. In plants, drought-adapted soil communities can increase fitness in drought conditions [11]. These studies suggest that microbes can adapt and evolve to stressful conditions in natural environments and buffer the stressor for the host, but much about the evolution of these interactions remains unclear—for example, we do not know to what extent these microbial effects depend on host genotypes. Host genetic variation may influence the responsiveness to microbial variation such that some combinations of host and microbiome differentially affect host phenotypes within species. The effects on host phenotype will also likely depend on the environmental context, where mismatches between the environment, host genotype, and microbiome may negatively affect host fitness. In other words, host genotype x microbial genotype x environment (G_H x G_M x E) interactions may shape how host-microbiome systems evolve. Such complex G_H x G_M x E interactions are likely to be common, and yet their contribution in shaping host phenotypic variation and adaptation are underexplored. As the study of microbiomes moves beyond descriptive attributes, experimentally assessing the interactions between host and microbial evolution are a key research priority for better understanding the interplay in host-microbiome interactions [12, 13].

The fruit fly, *Drosophila melanogaster*, has emerged as an excellent model to study host-
microbe interactions. Flies have relatively simple microbiomes, often composed of fewer than 10
bacterial species from acetic acid and lactic acid bacteria families with effects on many different
fly traits [14]. The microbiome is largely environmentally acquired through feeding, defecation,
and egg smearing during oviposition [15, 16]. After hatching, larvae acquire their microbes
through a combination of microbial persistence and colonization and form associations over the
lifespan of the fly [17–19]. Many fly-associated microbes can be individually cultured and
reconstituted in various combinations to inoculate sterile flies [14, 20]. With the feasibility of
microbiome manipulations and rich genetic resources, *Drosophila* is an ideal model to study the
interplay between host genetic, microbiome, and environment (i.e., G_H x G_M x E interactions) in
shaping host phenotypes that underlie rapid host adaptation.
In this study, to investigate whether the microbiome can serve as a reservoir of adaptive
potential for hosts, we studied the contribution of the microbiome in the context of experimental
evolution. Experimental evolution has a long history of providing fundamental insights into
adaptive processes, but has largely neglected the contribution of the microbiome [6, 21, 22].
One particularly powerful approach is the Evolve and Resequence (E&R) framework to measure
the genomic response to selection [23]. E&R experiments typically begin with large, outbred
populations which are then subjected to a selective regime, and in parallel, control populations
are maintained in a benign, non-selective environment. After several generations, the control
and evolved populations are then sequenced to identify the genomic response to selection.
Although traditionally E&R experiments have focused on the organism targeted by selection
(e.g., yeast, flies, mice, etc.), the E&R approach can also identify changes in the microbiome
associated with the evolved populations [22]. However, to determine whether any changes in
the microbiome over generations of selection actually contributed to host adaptation, it is
necessary to transplant the microbiome between control and evolved populations. This
approach holds great potential to inform how the microbiome shapes host phenotypes and
contributes to host adaptation.
Here, we leverage ~150 generations of fly and microbial adaptation to a nutritionally
stressful, high sugar (HS) diet. HS diets exert stress on flies, and flies exhibit similar obesity-like
phenotypes as humans with elevated triglycerides, insulin resistance, and shortened lifespans
[24–27]. In turn, the different bacteria in the fly microbiome have a range of effects on fly
phenotypes, shaping life history traits [28, 29] and metabolism [28, 30], and these microbiome
effects on host phenotype can depend on host genetics [31, 32]. Together, these findings
suggest that the microbiome could potentially contribute to host adaptation to the HS diet.

Our goal was to determine whether transplanting the adapted microbiome into non-
adapted fly genotype would confer adaptive traits in the stressful HS diet. Through a fully
reciprocal host x microbiome x diet transplant experiment (Fig. 1), we evaluate how each
component contributes to fitness associated traits in flies. The power of our approach is to
compare the benefit of adapted microbes relative to non-adapted microbes under different host
genetic and environmental contexts. Our results point to the important role microbiomes may
play in influencing host adaptation but also highlight the complexities of environmental context
dependence on host-microbiome interactions.
107
MATERIALS AND METHODS
Experimental design
We first identified if the microbiome responded to selection from experimental evolution
to the HS diet. Then, to test if the microbiome can transfer adaptive traits, we performed a
reciprocal microbiome transplant experiment (Fig. 1). All host genotypes x microbiome x diet
interactions were generated in the reciprocal transplant. Both genotypes (C and HS flies) were
inoculated with microbiomes (C, HS) or kept sterile and raised in both diets (C and HS) for a
total of 12 treatments. This experimental design allowed us to evaluate how host x microbiome x
environment interactions influence the transfer of adaptive phenotypes.
117
Fly populations
All fly populations were maintained at 25ºC with 12 hour light:dark cycles. The base
ancestral population was derived by round-robin crossing a subset of the Global Diversity lines
[33], which were then maintained at large population size (>10,000 flies) and allowed to
interbreed freely for ~50 generations before HS selection began. The control (C) diet was
composed of 8% glucose, 8% yeast, 1.2% agar, with 0.04% phosphoric acid and 0.4% propionic
acid as preservatives. Three replicate populations for HS selection were derived from the base
population and maintained on the same diet, but with an additional 12% sucrose and at lower
population size (>5,000 flies/cage). The control population was the ancestral population
maintained at large population size (>10,000 flies) in a parallel on the C diet during the entirety
of the HS selection. The approach used here allowed for natural selection on HS diets; we did
not guide or select for particular traits, which is a common approach in experimental evolution
studies in _Drosophila_. We consider the two populations, C and HS, to be two different
genotypes, though note they are outbred and retain genetic diversity.
132

Identifying the microbiome response to selection in the HS diet

We first identified how the microbiome changed between C and HS flies through 16S

rRNA amplicon sequencing. In brief, DNA was extracted from single, age-matched (7-10 days

post eclosion adults) using Zymo Quick-DNA extraction kit (Zymo D3012). The 16S V1-V2

region was amplified, pooled, and digested with BstZ17I (NEB R3954) to deplete Wolbachia

reads [34]. Libraries were sequenced using 300 bp paired-end reads using the Illumina MiSeq

platform at the Princeton University Genomics Core. We used QIIME2 v2019.4 [35] to process

reads, cluster into amplicon sequence variants (ASVs) with DADA2, and assign taxonomy with

Greengenes classifier trimmed to the 16S rRNA V1-V2 regions. Data was imported into

phyloseq for visualization [36]. Flies with fewer than 500 reads/individual were discarded before

the analyses.

To identify differentially abundant bacteria between the C and HS flies, we used ANCOM

(analysis of composition of microbiomes) [37] implemented in QIIME2. We cultured the most

differentially abundant bacteria from HS flies, *A. pasteurianus*, and the significantly differentially

abundant bacteria within the *Acetobacter* genus and enriched in C flies, *A. indonesiensis*. We

confirmed 16S rRNA sequence identity using Sanger sequencing. For simplicity, we refer to *A.*

indonesiensis as C *Acetobacter* and *A. pasteurianus* as HS *Acetobacter*.

Transplanting microbiomes to test for adaptive significance

We first performed both a complete microbiome transplant. We then performed a

focused transplant of the dominant C and HS *Acetobacter* strains identified through ANCOM to

measure the impact of host genetic, microbial, and environment interactions on fitness-

associated phenotypes.

First, to transplant the complete microbiome, we collected frass for ~ 5 hours from 300 C

and HS donor flies (age matched 7-10 days post eclosion). Flies were placed in sterile 50 ml

tubes, lined with a plastic sheet. Frass was then washed off the plastic using sterile water and

filtered through fine mesh to remove particulates and any eggs. Preliminary tests suggested that

the HS microbiome had much lower bacterial abundance, and so the C microbiome was

resuspended in 6 ml sterile water and 3 ml for the HS microbiome. 50 µl of the frass wash was

used to inoculate axenic eggs in sterile diets following standard protocols [20], and flies were

reared in a biosafety cabinet at 25ºC and 12:12 light:dark cycle through the lifespan of the

recipient flies. At 7-10 days post eclosion, recipient flies were stored at -80ºC and DNA

subsequently extracted from individuals. The microbiome of recipients was assessed through

16S rRNA V1-V2 profiling as described above and rarified to 500 reads/fly. To assess the

fidelity of the microbiome transplant, we tested for the effects of host genotype and microbiome
on Bray-Curtis dissimilarity for each diet separately using the ADONIS2 implementation of
PERMANOVA in vegan [38].

To perform the *Acetobacter* transplant, both *Acetobacter* strains were cultured in liquid
MRS, density was normalized to OD_{600} = 0.1, and 50 µl of bacteria (or PBS for sterile
treatments) was used to inoculate axenic eggs in sterile diets [20]. For each treatment, 4-12
replicate populations were maintained. Flies were then reared in a biosafety cabinet at 25ºC and
12:12 light:dark cycle through the lifespan of the recipient flies. For the *Acetobacter* transplant,
we measured two fitness-associated phenotypes: developmental time and fecundity.

To measure developmental time, we counted the number of new pupae every 6 hours
during their daytime (i.e., 8am, 2pm, 8pm) until all treatments began eclosing (4-12
replicates/treatment, see Supp. Table 1 for sample sizes and replication). For developmental
time, we used the survival [39] and survminer [40] packages in R to analyze development data.
We used Cox proportional hazards to identify how interactions between fly genotype,
microbiome, and diet influence development time.

For fecundity, we counted the number of eggs laid by individual 8-10 day old females
(N=644 total, see Supp. Table 2 for sample sizes). Females were placed in a 24-well plate with
oviposition media. Oviposition media was the same as the fly food media, but with yeast extract,
which makes the diet transparent and easier to count eggs. Females laid eggs on both C and
HS media; media did not affect egg lay (Mann-Whitney-Wilcoxon test, W=50964, p=0.09), and
data were pooled from both oviposition media for analysis. To minimize plate effects, two
treatments were assayed on each plate, and treatments were randomly assigned across plates.
Females laid eggs for ~4-6 hrs at 25ºC during the late afternoon (2pm-8pm) to capture similar
circadian rhythms in egg laying across all treatments. We used a zero-inflated Poisson
distribution implemented in glmmTMB [41] testing for the effects of interactions between fly
genotype, *Acetobacter* strain, and diet with oviposition plate (N=29 plates) as the random effect.
193
Characterizing the genomic differences between C and HS Acetobacter
To further characterize the mechanisms underlying the fitness-associated benefits from
the C and HS *Acetobacter* strains, we performed whole genome sequencing on these bacteria.
Libraries were prepared following the manufacturer instructions using the Illumina PCR-free
Library Prep kit, and then 150-bp paired end reads were sequenced on an Illumina NovaSeq
platform. Genomes were assembled using SPAdes [42] with --isolate flag and annotated using
the Rapid Annotation using Subsystem Technology (RAST) server [43].

We first compared average nucleotide identity (ANI) between the C and HS *Acetobacter*

using OrthoANIu [44]. OrthoANIu computationally fragments bacterial genomes into 1020 bp

(contigs <1020 bp are discarded), performs reciprocal BLASTn searches for all fragments, and

then computes ANI for all reciprocal BLASTn hits. Then to identify functional changes, we

manually compared the RAST annotations to identify pathways unique to either C or HS

Acetobacter. We also generated the RAST annotations from two other abundant bacteria in the

C microbiome and compared them with the HS *Acetobacter* annotations: *Algoriella xinjiangensis*

(NCBI accession FOUZ01000040.1) and *Acetobacter persici* (NCBI accession

JOPC00000000.1). To confirm the differences between C and HS *Acetobacter* strains, we

identified orthologs between the two *Acetobacter* genomes using OrthoFinder v2.5.2 [45]. Then

to gain insight into the function of the putative genes, we performed BLASTP v2.10.1+ searches

against the NCBI non-redundant protein sequence database [46].

The genomic analysis performed above suggested that the HS *Acetobacter* genome

encoded several pathways for urea degradation, but were not present in the C *Acetobacter*

genome. To validate this observation, we measured the ability of the *Acetobacter* strains to

degrade uric acid in fly food. Uric acid was added at 50 µmol concentration to 30 µl of C and HS

diet. Then 10 µl of C or HS Acetobacter (or sterile PBS control), normalized to OD_{600}=0.1, were

added to the diet and incubated for 4 days at 25ºC to degrade the uric acid. Uric acid

concentration was measured using the Amplex Red Uric Acid/Uricase kit (ThermoFisher

A22181) following the manufacturer instructions. ANOVA was used to analyze the uric acid

degradation by bacteria. We tested for the effects of bacteria (C or HS *Acetobacter* strain), diet,

and the interaction between bacteria and diet.

RESULTS

Fly microbiome responds to HS selection

First, we examined if the microbiome differed between C and HS flies following ~150

generations of adaptation to the C or HS diets (Fig. 2A). HS flies were dominated by

Acetobacter pasteurianus (86.7% +/- 1.8 SE) and unclassified Acetobacteraceae strain (5.1%

+/- 1.8 SE), while C flies were more diverse with *Algoriella xinjiangensis* (37.3% +/- 5.3 SE),

Acetobacter persici (16.9% +/-2.6 SE), and *Acetobacter indonesiensis* (14.8% +/- 2.1 SE). The

most differentially abundant bacteria were *A. pasteurianus* and Acetobacteraceae strain for the

HS flies, while *A. xinjiangensis* and *A. indonesiensis* were associated with C flies (Fig. 2B,

Supp. Table 3). These bacteria are unique to either C or HS flies and found across individual

flies, demonstrating that the HS diet selected for a different fly microbiome, dominated by *A.*

pasteurianaus.

<u>*Transplanting Acetobacter influences fitness-related traits*</u>

To test if transferring the microbiome could transfer adaptive phenotypes to naive hosts,

we performed a fully reciprocal, host genotype x microbiome x diet transplant experiment. We

first transplanted a complete microbiome. On the C diet, the difference in C and HS

microbiomes were maintained in recipients (Fig. 2C, Supp. Table 4, PERMANOVA, R^2 = 0.38, p

= 0.001). However, the microbiome was not successfully transplanted for flies on the HS diet, as

the variance explained by microbiome was much lower in recipients (Fig. 2C, Supp. Table 5,

PERMANOVA, R^2 = 0.04, p = 0.004). This suggests that the HS diet was highly selective and as

we discuss later, likely limits bacterial survival.

Instead, given the simplicity of the microbiome associated with HS flies, we performed

another microbial transplant using the *Acetobacter* strains identified as differentially abundant

between C and HS flies: *A. indonesiensis* (C *Acetobacter*) and *A. pasteurianus* (HS

Acetobacter). While not the complete bacterial community from the C flies, we reasoned that

because both community complexity (i.e., single dominant species in the HS microbiome versus

several in the C microbiome) and phylogenetic diversity (i.e., single dominant *Acetobacter* in the

HS microbiome versus two *Acetobacter* species, *Algoriella*, *Commensalibacter*, etc. in the C

microbiome) differ, using only C *Acetobacter* still provides an ecologically relevant substitute,

while controlling for differences in diversity between the C and HS microbiome. We then

measured higher order, fitness-associated traits (developmental timing and fecundity), which

reflect cumulative adaptations to the multitude of selective pressures in the HS diet.

Development was shaped by interactions between fly genotype, *Acetobacter*, and diet

(Fig. 3A, Supp. Table 6). On the C diet, fly genotype influenced the response to the *Acetobacter*

transplant. HS flies with the C *Acetobacter* developed more rapidly (avg. 140.8 hrs +/- 0.54 SE)

than with the HS *Acetobacter* (avg. 147.6 hrs +/- 0.60 SE). There was no effect for the C fly.

However, on the HS diet, the HS *Acetobacter* accelerated development (HS *Acetobacter* avg.

176.5 hrs +/- .50 SE, C *Acetobacter* avg. 186.8 hrs +/-0.60 SE). The three-way interaction

between fly genotype, *Acetobacter*, and diet significantly affected development time (log-

likelihood =-58500, X^2 = 47.625, p < 0.0001).

Fecundity responded to interactions between fly genotype, *Acetobacter*, and diet (Fig.

3B, Supp. Table 7). On average, treatments were more fecund on the C diet, while treatments

on HS diets laid 54.3% fewer eggs (C avg. 13.0 eggs +/-0.6 SE, HS, avg. 5.9 eggs +/- 0.4 SE).

Sterile treatments substantially lowered fecundity, only on the HS diet (+microbes avg. 7.45

eggs +/- 0.5 SE, sterile avg. 1.67 eggs +/- 0.4 SE). Notably, the response to *Acetobacter*

treatment depended on fly genotype for the HS diet. On the HS diet, HS flies with HS

Acetobacter (avg. 8.4 eggs +/-1.4 SE) were almost twice as fecund as flies with the C

Acetobacter (avg. 4.5 eggs +/- 1.0 SE). The interaction between fly genotype, microbiome, and

diet significantly affected fecundity (interaction X^2 = 16.4, df = 2, p=0.0003).

HS Acetobacter uniquely encodes uric acid degradation pathway

To better understand how the HS *Acetobacter* may benefit flies in the HS diet, we

sequenced and compared the genomes of the C and HS *Acetobacter* strains. ANI between the

strains was 73.68% (Supp. Table 8). Annotation by RAST showed that strains shared a majority

of predicted functions (>90% for both strains, Supp. Fig. 1). One key difference between the two

genomes was that HS *Acetobacter* encodes for several genes that degrade uric acid, but C

Acetobacter does not (Supp. Table 9). Further, the other dominant bacteria in the C microbiome

(*Algoriella xinjiangensis* and *Acetobacter persici*) also lack uric acid degradation genes (Supp.

Table 9). Indeed, we found that interactions between *Acetobacter* and the diet shape uric acid

degradation (Fig. 4, Supp. Table 10). Only on the HS diet does HS *Acetobacter* reduce the

concentration of uric acid by ~50% compared to C *Acetobacter*, with only negligible changes to

uric acid concentration on the C diet.

DISCUSSION

Here, we explored how the microbiome influences host fitness in stressful environments.

Because all combinations of G_H x G_M x E interactions were produced in the transplant

experiment, the relative benefits of C and HS *Acetobacter* across the different host genetic

(selected or unselected) and environmental conditions (HS or control diet) could be evaluated.

Through reciprocal host x microbiome x diet transplant experiments, we show that G_H x G_M x E

interactions shape host fitness by modifying developmental timing and fecundity. The HS

Acetobacter genome encoded uric acid degradation genes and, *in vitro*, lowered uric acid on the

HS diet, while the C *Acetobacter* did not. Our results highlight the complex interactions

generated by the microbiome, and we discuss important future directions to further dissect

these relationships.

Acetobacter dominates HS microbiome

The HS microbiome was dominated by a single species of bacteria, *Acetobacter*
pasteurianus (Fig. 2). *Acetobacter* species are common in flies, both in laboratory settings and
wild environments [18, 47–51]. While *Acetobacter* are typically sugar specialists [52], one major
axis of variation separating the C and HS microbiome was the ability to degrade uric acid. We
did not find an effect of C or HS *Acetobacter* strains in shaping triglyceride accumulation (Supp.
Fig. 2); triglyceride accumulation is often a benefit associated with *Acetobacter* in *Drosophila*
[28, 31]. In the context of *Drosophila* biology, uric acid accumulation is commonly associated
with high sugar diets in flies [53]. The accumulation of uric acid may also exacerbate the
deleterious effects of HS diet, altering oxidative stress and inflammation that can promote
obesogenic phenotypes [54, 55]. Uric acid is normally excreted by flies, but high concentrations
slow development [56] and shorten lifespan [53, 57]. Not all *Acetobacter* species can degrade
uric acid, including the C *Acetobacter*, but the ability to degrade is more common in fly-
environment than in other free-living *Acetobacter* species [55]. Here, comparative analyses
between the C and HS *Acetobacter* genomes showed that HS *Acetobacter* encoded for several
genes in the uric acid degradation pathway, while the C *Acetobacter* genome did not. Uric acid
degradation may be the key fitness benefit of the HS *Acetobacter* over the C *Acetobacter* in the
HS diet.
The fitness advantage of uric acid degradation was likely direct for the HS *Acetobacter*,
but indirect for the fly. Over the course of experimental evolution, both fly and microbiome were
exposed to the HS diet, and in turn, the HS microbiome was dominated by *A. pasteurianus*,
which could degrade uric acid. *Acetobacter* species that degrade uric acid can outcompete
Acetobacter species that cannot in the fly environment [55]. For the HS *Acetobacter*, uric acid
degradation may underlie its persistence in the stressful HS diet, and thus indirectly benefiting
flies through reductions of uric acid in the diet. The benefit of uric acid degradation may be
especially important during development. Larvae have limited endogenous abilities to detoxify
uric acid, and the primary mechanism of uric acid tolerance is by decreasing exposure through
decreased uptake or increased excretion [56, 58, 59]. For adults, it is unclear if HS *Acetobacter*
also degrades uric acid in the fly gut or simply benefiting from the reduction in the diet. While
these questions require further characterization of fly physiology, uric acid degradation by HS
Acetobacter is similar to a handful of examples of environmentally acquired microbiomes that
facilitate adaptation through detoxifying the environment, such as insecticide resistance in bean
bugs [9] or terpene detoxification in woodrats [10]. Our results contribute to a growing body of
literature that suggests that the microbiome can encode novel functions that hosts can leverage
the microbiome during adaptation to stressful environments [6].

Adaptive potential of the microbiome depends on evolutionary and ecological context

Many have predicted that the microbiome could accelerate local adaptation in the host,

facilitating phenotypic changes to buffer local stressors [6, 7, 60, 61]. If so, transplanting the

locally adaptive microbiomes should confer adaptive phenotypes to naive individuals. However,

the benefit may depend on interactions between host genotype and environment. G_H x G_M x E

interactions have the potential to facilitate or impede evolution in challenging environments [12,

13]. Through combining experimental evolution with reciprocal transplants, our data allowed us

to test if and how ecological context (i.e., diet) and evolutionary context (i.e., fly genotype)

shape the adaptive potential of the microbiome.

To shape adaptive potential, the microbiome must contribute to host phenotypic

variation, not unlike traditional (host) additive genetic variance [6]. The G_H x G_M x E interaction

for development illustrates how diet (the ecological context), when matched with host genotype

(the evolutionary context) may increase the fitness benefit of locally adaptive microbiomes (Fig.

3A). The HS *Acetobacter* accelerated development on the HS diet (a match between microbe

and diet, independent of fly genotype) and slowed development compared to the C *Acetobacter*

on the C diet for only the HS fly (a mismatch between microbe and environment, but dependent

on fly genotype). The mismatch only mattered for the HS fly; C fly development was similar

between the C and HS *Acetobacter* on the C diet. However, the G_H x G_M x E interaction that

shaped fecundity is suggestive of how genetic interactions may impede the transfer of adaptive

potential. The naive host, C flies, did not receive a fecundity increase from the HS *Acetobacter*

in the HS diet, but HS flies did (Fig. 3C). The presence of G_H x G_M x E interactions for both

development and fecundity suggests that the microbiome contributed to fly evolution. HS flies

responded differently to the *Acetobacter* variation from the C flies, and more so, responsiveness

depended on the diet. Because host genotypes may vary in their ability to leverage beneficial

microbes in stressful environments, genetic interactions may limit the benefits of locally adaptive

microbiomes. Understanding these G_H x G_M interactions will be especially important in using

evolution to engineer beneficial microbiomes to solve pressing problems in public health and

agriculture [6, 12, 13, 62]. While longitudinal sampling of the evolutionary trajectory is necessary

to determine to what extent the HS fly evolved or selected for beneficial *Acetobacter* traits, our

results clearly demonstrate how the ecological context and host evolution can shape the

adaptive potential of the microbiome.

Research priorities

Here, we leveraged host and microbial evolution from ~150 generations of adaptation to
a stressful diet through the E&R approach. Such experiments have illuminated many important
findings about the role of demography, population genetics, and genetic architecture in shaping
adaptive traits [23, 63]. However, missing from this framework has been the role of the
microbiome. In a recent survey of ten E&R experiments, the microbiome also frequently
responded to experimental evolution, particularly in traits associated with metabolism [22].
Taken together with our reciprocal transplant experiment, our data also highlight how the
microbiome may contribute to host adaptation – effects previously unseen when focusing on the
Drosophila genome. Recent work on the *Drosophila* microbiome linked to natural variation in
wild populations suggested that a single inoculation of two different bacteria linked to different
life-history strategies, can lead to rapid phenotypic change and genetic differentiation in one fly
population [29, 64]. Our current study advances these findings by comparing differences in
ecologically relevant *Acetobacter* evolution to differences in fly evolution, providing one of the
first experimental dissections of host-microbiome interactions in the E&R context. Through
reciprocal transplants, we identified how interactions between fly genotype, microbiome, and
diet shape fitness related traits.

The G_H x G_M x E interactions observed here highlight how evolution in both the host and
microbiome occurred during adaptation to the HS diet. In particular, the ability of HS
Acetobacter to degrade uric acid in the HS diet reflects the potential for diffuse coevolution [65].
The HS diet likely led to an increase in uric acid production by flies, and then the *Acetobacter*
that could exploit the new niche increased, which the HS flies benefited from the reduction in
toxicity. Ultimately, in our system and perhaps many others, the evolutionary benefits between
host and microbiome are intertwined. Future work utilizing experimental evolution will
demonstrate how the joint evolution of host and microbiome shapes the response to selection,
providing fundamental insights into how natural selection operates on complex systems.
394

ACKNOWLEDGEMENTS
We thank members of the Ayroles lab for helpful feedback on this work. LPH was supported by
NSF-GRFP under grant DGE1656466 and National Institutes of Health (NIH) grant R35-
GM124881 to JFA.

DATA AVAILABILITY

Genome sequencing will be uploaded to NCBI [upon acceptance]. Phenotypic data will be

deposited in Dryad [upon acceptance]. Code used to analyze data will be posted on github

[upon acceptance].

FIGURES

406
407
Figure 1: Experimental design to leverage ~150 generations of fly adaptation to the high sugar

(HS) diet to study the adaptive potential of the microbiome. The conceptualized evolutionary

trajectory shows control (C) flies and microbiome in red and HS flies and microbiome in blue. By

the end of experimental evolution, the C fly population (red) has a different microbiome than the

HS fly population (blue). After the ~150 generations of selection, we performed a fully reciprocal

host x microbiome x diet transplant experiment for all combinations of fly genotypes (C or HS),

microbes (sterile, C, or HS), and diets (C, or HS). Together, this approach allows for the

assessment of the benefit of adapted relative to non-adapted microbes in both host genetic and

environmental contexts.

428
Figure 2: The microbiome in HS flies is different from C flies. A) Relative abundance of bacteria
determined using 16S rRNA sequences. Each bar represents an individual fly (N=38 C flies, 21
HS flies). The C fly microbiome is enriched for *A. indonesiensis*, *A. persici*, and *Algoriella*
xinjiangensis. The HS fly microbiome is enriched for *A. pasteurianus* and unclassified
Acetobacteraceae. B) ANCOM results for differentially abundant bacteria between C and HS
flies. Each point represents a different ASV, colored by whether significantly enriched for C (red)
or HS flies (blue). Labels denote the taxonomic classification for significant ASVs and are
highlighted to match the color from the relative abundance in 2A. C) PCoA plots based on Bray-
Curtis dissimilarity for recipients in the whole microbiome transplant. Plot is faceted by diet, and
colors represent the different combination of host genotype, microbiome, and diet. The
difference between C and HS microbiomes was maintained for recipients on the C diet, but not
on the HS diet.

Figure 3: Effects of the *Acetobacter* transplant on fitness-related traits. A) Hazard ratio plot for

Acetobacter effects on development time. Treatment is shown with the fly diagram, where fly

color represents fly genotype, circle represents *Acetobacter* (no circle = sterile), and box

represents diet. The full treatment is written out (i.e., $C_H+C_M+C_E$ = C fly with C microbiome on

the C diet) with the total number of replicates (N_{TUBE}) and pupae (N_{PUP}) phenotyped across

replicates shown below. Boxplots show hazard ratios and 95% confidence intervals for each

treatment relative to $C_H+0_M+C_E$. Mismatches between diet and microbiome slowed

development. In the C diet, HS flies with C *Acetobacter* developed faster than the HS flies with

HS *Acetobacter*. However, on the HS diet, C *Acetobacter* slowed, while HS *Acetobacter*

accelerated development. B) Least square means (LSM) from fecundity, modeled with a zero-

inflated Poisson distribution show fly genotype x *Acetobacter* x diet interactions shaped

fecundity. Plots are faceted by fly diet, where colors represent *Acetobacter* treatment, and the

number of age-matched females assayed below the fly diagram. Error bars show standard error

for LSMs. For the HS fly genotype on the HS diet, the HS *Acetobacter* increased fecundity

relative to the C *Acetobacter*, suggesting a fecundity benefit with the adapted microbiome.

485
486 Figure 4: Uric acid degradation by HS *Acetobacter*. Boxplots show uric acid in the *in vitro* diet
487 assay. Boxplots are colored by diet with points colored by *Acetobacter* treatment. Only in the HS
488 diet, HS *Acetobacter* degraded significantly more uric acid than C *Acetobacter*.

REFERENCES

1. Sommer F, Bäckhed F. The gut microbiota—masters of host development and physiology.

Nat Rev Microbiol 2013; **11**: 227.

2. Selosse M-A, Bessis A, Pozo MJ. Microbial priming of plant and animal immunity:

symbionts as developmental signals. *Trends Microbiol* 2014; **22**: 607–613.

3. Vuong HE, Yano JM, Fung TC, Hsiao EY. The Microbiome and Host Behavior. *Annu Rev*

Neurosci 2017; **40**: 21–49.

4. Moran NA, Sloan DB. The hologenome concept: Helpful or hollow? *PLoS Biol* 2015; **13**:

e1002311.

5. Koskella B, Hall LJ, Metcalf CJE. The microbiome beyond the horizon of ecological and

evolutionary theory. *Nat Ecol Evol* 2017; **1**: 1606–1615.

6. Henry LP, Bruijning M, Forsberg SKG, Ayroles JF. The microbiome extends host

evolutionary potential. *Nat Commun* 2021; **12**: 5141.

7. Sudakaran S, Kost C, Kaltenpoth M. Symbiont Acquisition and Replacement as a Source of

Ecological Innovation. *Trends Microbiol* 2017; **25**: 375–390.

8. Douglas AE, Werren JH. Holes in the Hologenome: Why Host-Microbe Symbioses Are Not

Holobionts. *MBio* 2016; **7**: e02099.

9. Kikuchi Y, Hayatsu M, Hosokawa T, Nagayama A, Tago K, Fukatsu T. Symbiont-mediated

insecticide resistance. *Proc Natl Acad Sci U S A* 2012; **109**: 8618–8622.

10. Kohl KD, Dearing MD. The Woodrat Gut Microbiota as an Experimental System for

Understanding Microbial Metabolism of Dietary Toxins. *Front Microbiol* 2016; **7**: 1165.

11. Lau JA, Lennon JT. Rapid responses of soil microorganisms improve plant fitness in novel

environments. *Proc Natl Acad Sci U S A* 2012; **109**: 14058–14062.

12. Mueller UG, Sachs JL. Engineering Microbiomes to Improve Plant and Animal Health.

Trends Microbiol 2015; **23**: 606–617.

13. Busby PE, Soman C, Wagner MR, Friesen ML, Kremer J, Bennett A, et al. Research

priorities for harnessing plant microbiomes in sustainable agriculture. *PLoS Biol* 2017; **15**:

e2001793.

14. Douglas AE. The Drosophila model for microbiome research. *Lab Anim* 2018; **47**: 157–

164.

15. Wong AC-N, Luo Y, Jing X, Franzenburg S, Bost A, Douglas AE. The host as the driver of

the microbiota in the gut and external environment of Drosophila melanogaster. *Appl*

Environ Microbiol 2015; **81**: 6232–6240.

16. Blum JE, Fischer CN, Miles J, Handelsman J. Frequent replenishment sustains the

beneficial microbiome of Drosophila melanogaster. *MBio* 2013; **4**: e00860–13.

17. Bakula M. The persistence of a microbial flora during postembryogenesis of Drosophila

melanogaster. *J Invertebr Pathol* 1969; **14**: 365–374.

18. Broderick NA, Lemaitre B. Gut-associated microbes of Drosophila melanogaster. *Gut*

Microbes 2012; **3**.

19. Obadia B, Güvener ZT, Zhang V, Ceja-Navarro JA, Brodie EL, Ja WW, et al. Probabilistic

Invasion Underlies Natural Gut Microbiome Stability. *Curr Biol* 2017; **27**: 1999–2006.e8.

20. Koyle ML, Veloz M, Judd AM, Wong ACN, Newell PD, Douglas AE, et al. Rearing the Fruit

Fly Drosophila melanogaster Under Axenic and Gnotobiotic Conditions. 2016; e54219.

21. Hoang KL, Morran LT, Gerardo NM. Experimental Evolution as an Underutilized Tool for

Studying Beneficial Animal–Microbe Interactions. *Front Microbiol* 2016; **07**.

22. Henry LP, Ayroles JF. Meta-analysis suggests the microbiome responds to Evolve and

Resequence experiments in Drosophila melanogaster. *BMC Microbiol* 2021; **21**: 108.

23. Long A, Liti G, Luptak A, Tenaillon O. Elucidating the molecular architecture of adaptation

via evolve and resequence experiments. *Nat Rev Genet* 2015; **16**: 567–582.

24. Musselman LP, Fink JL, Narzinski K, Ramachandran PV, Hathiramani SS, Cagan RL, et al.

A high-sugar diet produces obesity and insulin resistance in wild-type Drosophila. *Dis*

Model Mech 2011; **4**: 842–849.

566 25. Na J, Musselman LP, Pendse J, Baranski TJ, Bodmer R, Ocorr K, et al. A Drosophila

567   model of high sugar diet-induced cardiomyopathy. *PLoS Genet* 2013; **9**: e1003175.

568 26. Rovenko BM, Kubrak OI, Gospodaryov DV, Perkhulyn NV, Yurkevych IS, Sanz A, et al.

569   High sucrose consumption promotes obesity whereas its low consumption induces

570   oxidative stress in Drosophila melanogaster. *J Insect Physiol* 2015; **79**: 42–54.

571 27. Pallares LF, Lea AJ, Han C, Filippova EV, Andolfatto P. Diet unmasks genetic variants that

572   regulate lifespan in outbred Drosophila. *bioRxiv* 2021.

573 28. Newell PD, Douglas AE. Interspecies Interactions Determine the Impact of the Gut

574   Microbiota on Nutrient Allocation in Drosophila melanogaster. *Appl Environ Microbiol* 2013;

575   **80**: 788–796.

576 29. Walters AW, Hughes RC, Call TB, Walker CJ, Wilcox H, Petersen SC, et al. The microbiota

577   influences the Drosophila melanogaster life history strategy. *Mol Ecol* 2020; **29**: 639–653.

578 30. Shin SC, Kim S-H, You H, Kim B, Kim AC, Lee K-A, et al. Drosophila microbiome

579   modulates host developmental and metabolic homeostasis via insulin signaling. *Science*

580   2011; **334**: 670–674.

581 31. Chaston JM, Dobson AJ, Newell PD, Douglas AE. Host Genetic Control of the Microbiota

582   Mediates the Drosophila Nutritional Phenotype. *Appl Environ Microbiol* 2016; **82**: 671–679.

583 32. Dobson AJ, Chaston JM, Newell PD, Donahue L, Hermann SL, Sannino DR, et al. Host

584   genetic determinants of microbiota-dependent nutrition revealed by genome-wide analysis

585   of Drosophila melanogaster. *Nat Commun* 2015; **6**: 7296.

586 33. Grenier JK, Roman Arguello J, Moreira MC, Gottipati S, Mohammed J, Hackett SR, et al.

587   Global Diversity Lines–A Five-Continent Reference Panel of Sequenced Drosophila

588   melanogaster Strains. *G3: Genes, Genomes, Genetics* 2015; **5**: 593–603.

589 34. Simhadri RK, Fast EM, Guo R, Schultz MJ, Vaisman N, Ortiz L, et al. The Gut Commensal

590   Microbiome of Drosophila melanogaster Is Modified by the Endosymbiont Wolbachia.

591   *mSphere* 2017; **2**.

35. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al.

Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2.

Nat Biotechnol 2019; **37**: 852–857.

36. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and

graphics of microbiome census data. *PLoS One* 2013; **8**: e61217.

37. Mandal S, Van Treuren W, White RA, Eggesbø M, Knight R, Peddada SD. Analysis of

composition of microbiomes: a novel method for studying microbial composition. *Microb*

Ecol Health Dis 2015; **26**: 27663.

38. Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, et al. Vegan

community ecology package: ordination methods, diversity analysis and other functions for

community and vegetation ecologists. *R package ver* 2015; 2–3.

39. Therneau TM, Lumley T. Package 'survival'. *R Top Doc* 2015; **128**: 112.

40. Kassambara A. Kosinski M. survminer: Drawing survival curves using 'ggplot2,' 2018. *URL*

https://CRAN R-project org/package= survminer R package version 0 4 2018; **3**.

41. Magnusson A, Skaug H, Nielsen A, Berg C, Kristensen K, Maechler M, et al. glmmTMB:

generalized linear mixed models using template model builder. *R package version 0 1*

2017; **3**.

42. Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes De Novo

Assembler. *Curr Protoc Bioinformatics* 2020; **70**: e102.

43. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST Server:

rapid annotations using subsystems technology. *BMC Genomics* 2008; **9**: 75.

44. Yoon S-H, Ha S-M, Lim J, Kwon S, Chun J. A large-scale evaluation of algorithms to

calculate average nucleotide identity. *Antonie Van Leeuwenhoek* 2017; **110**: 1281–1286.

45. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative

genomics. *Genome Biol* 2019; **20**: 238.

46. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J*

Mol Biol 1990; **215**: 403–410.

47. Chandler JA, Lang JM, Bhatnagar S, Eisen JA, Kopp A. Bacterial communities of diverse

Drosophila species: ecological context of a host-microbe model system. *PLoS Genet* 2011;

7: e1002272.

48. Staubach F, Baines JF, Künzel S, Bik EM, Petrov DA. Host species and environmental

effects on bacterial communities associated with Drosophila in the laboratory and in the

natural environment. *PLoS One* 2013; **8**: e70749.

49. Adair KL, Wilson M, Bost A, Douglas AE. Microbial community assembly in wild populations

of the fruit fly Drosophila melanogaster. *ISME J* 2018; **12**: 959–972.

50. Wang Y, Kapun M, Waidele L, Kuenzel S, Bergland AO, Staubach F. Common structuring

principles of the Drosophila melanogaster microbiome on a continental scale and between

host and substrate. *Environ Microbiol Rep* 2020; **12**: 220–228.

51. Henry LP, Ayroles JF. Drosophila melanogaster microbiome is shaped by strict filtering and

neutrality along a latitudinal cline. *Mol Ecol* 2022; **31**: 5861–5871.

52. Kersters K, Lisdiyanti P, Komagata K, Swings J. The Family Acetobacteraceae: The

Genera Acetobacter, Acidomonas, Asaia, Gluconacetobacter, Gluconobacter, and Kozakia.

In: Dworkin M, Falkow S, Rosenberg E, Schleifer K-H, Stackebrandt E (eds). *The*

Prokaryotes: Volume 5: Proteobacteria: Alpha and Beta Subclasses. 2006. Springer New

York, New York, NY, pp 163–200.

53. van Dam E, van Leeuwen LAG, Dos Santos E, James J, Best L, Lennicke C, et al. Sugar-

Induced Obesity and Insulin Resistance Are Uncoupled from Shortened Survival in

Drosophila. *Cell Metab* 2020; **31**: 710–725.e7.

54. Sautin YY, Johnson RJ. Uric acid: the oxidant-antioxidant paradox. *Nucleosides*

Nucleotides Nucleic Acids 2008; **27**: 608–619.

55. Winans NJ, Walter A, Chouaia B, Chaston JM, Douglas AE, Newell PD. A genomic

investigation of ecological differentiation between free-living and Drosophila-associated

bacteria. *Mol Ecol* 2017.

56. Henry Y, Renault D, Colinet H. Hormesis-like effect of mild larval crowding on

thermotolerance in Drosophila flies. *J Exp Biol* 2018; **221**.

57. Lang S, Hilsabeck TA, Wilson KA, Sharma A, Bose N, Brackman DJ, et al. A conserved

role of the insulin-like signaling pathway in diet-dependent uric acid pathologies in

Drosophila melanogaster. *PLOS Genetics* . 2019. , **15**: e1008318

58. Pierce VA, Mueller LD, Gibbs AG. Osmoregulation in Drosophila melanogaster selected for

urea tolerance. *J Exp Biol* 1999; **202**: 2349–2358.

59. Etienne R, Fortunat K, Pierce V. Mechanisms of urea tolerance in urea-adapted

populations of Drosophila melanogaster. *J Exp Biol* 2001; **204**: 2699–2707.

60. Alberdi A, Aizpurua O, Bohmann K, Zepeda-Mendoza ML, Gilbert MTP. Do vertebrate gut

metagenomes confer rapid ecological adaptation? *Trends Ecol Evol* 2016; **31**: 689–699.

61. Carrier TJ, Reitzel AM. The Hologenome Across Environments and the Implications of a

Host-Associated Microbial Repertoire. *Front Microbiol* 2017; **8**: 802.

62. Mueller EA, Wisnoski NI, Peralta AL, Lennon JT. Microbial rescue effects: How

microbiomes can save hosts from extinction. *Funct Ecol* 2020; 33.

63. Schlötterer C, Kofler R, Versace E, Tobler R, Franssen SU. Combining experimental

evolution with next-generation sequencing: a powerful tool to study adaptation from

standing genetic variation. *Heredity* 2015; **114**: 431–440.

64. Rudman SM, Greenblum S, Hughes RC, Rajpurohit S, Kiratli O, Lowder DB, et al.

Microbiome composition shapes rapid genomic adaptation of Drosophila melanogaster.

Proc Natl Acad Sci U S A 2019; **116**: 20025–20032.

65. Koskella B, Bergelson J. The study of host-microbiome (co)evolution across levels of

selection. *Philos Trans R Soc Lond B Biol Sci* 2020; **375**: 20190604.

668
669

Supp. Fig. 1: Summary of the RAST annotation for the C and HS Acetobacter. A) Venn diagram showing the number of functions unique to C Aceto (106/1133 total), unique to HS Aceto (66/1093 total), or shared (1027). B) Number of predicted functions per category colored by either unique to C Acetobacter (red), unique to HS Acetobacter (blue), or shared (grey). In general, both genomes encoded similar functions, but small differences are observed across the different categories.

Supp. Fig. 1: Metabolic effects of the *Acetobacter* transplant. Triglyceride levels were assessed from pools of three, age-matched (7-10 days old) females. Tryglycerides were measured using colorimetric assay test (Sigma TR0100-1KT) and then normalized to protein using a Bradford assay (BioRad 5000205). A) Boxplot for triglycerides (TAG) normalized to protein. TAG amount reflects fat storage in *Drosophila*. Each point represents a pool of three age-matched female flies. TAG levels were mean centered to control for each batch assayed. Interactions between fly genotype and diet shaped TAG levels ($F_{1,78.1}$ = 9.00, p=0.004), but *Acetobacter* had no effect. B) LSM from the mixed model shows the significant interaction between fly genotype and diet. Error bars show standard error for LSM. TAG levels responded to diet for HS flies, but not for C flies.

SUPPLEMENTARY TABLES

Supp. Table 1: Sample size and replication for development measured during the *Acetobacter* transplant experiment.

Fly genotype	Microbiome	Diet	No. tubes	Avg larvae/tube	SE avg larvae/tube
control	sterile	control	8	79.2	7.88
control	sterile	high sugar	5	55.2	11.1
control	control	control	4	92.2	10.1
control	control	high sugar	8	87.2	8.37
control	high sugar	control	5	85.2	5.89
control	high sugar	high sugar	11	78.3	4.6
high sugar	sterile	control	9	95.7	7.01
high sugar	sterile	high sugar	5	72.8	10.8
high sugar	control	control	11	84.7	9.44
high sugar	control	high sugar	6	108	9.21
high sugar	high sugar	control	12	87.3	10.9
high sugar	high sugar	high sugar	6	87	6.93

Supp. Table 2: Sample sizes for fecundity in the *Acetobacter* transplant experiment .

Fly genotype	Microbiome	Diet	N Females
control	sterile	control	44
control	sterile	high sugar	46
control	control	control	48
control	control	high sugar	71
control	high sugar	control	59
control	high sugar	high sugar	81
high sugar	sterile	control	45
high sugar	sterile	high sugar	45
high sugar	control	control	61
high sugar	control	high sugar	46
high sugar	high sugar	control	59
high sugar	high sugar	high sugar	59

Supp. Table 3: ANCOM results for differentially abundant ASVs from C and HS fly microbiomes. Taxonomy was characterized to the lowest level an assignment could be made, and most ASVs were classified to species level. CLR=centered log ratio, W=test statistic, Sig=significant difference between groups (N.S. is nonsignificant).

ASV ID	CLR	W	Sig	Taxonomy
d84f3e1a7dbb97bc4695b954c8a2d2b2	4.30129864	25	HS fly	*Acetobacteraceae*
fb59adcbdd838dc7853f7388afb62b5a	7.82830283	25	HS fly	*Acetobacter pasteurianus*
9329896cd08f88a88296d921369cd8ea	-3.4609509	20	C fly	*Algoriella xinjiangensis*
f0d11f9f8ba452d5485a79e0b31b538f	-3.6539534	20	C fly	*Acetobacter indonesiensis*
3f2be3d57ac38f60b0b9c56fa5857f63	-2.7057494	18	N.S.	*Algoriella xinjiangensis*
70e118808cdc81b5b2670837c7833d8d	-3.3688051	18	N.S.	*Acetobacter persici*
dd4826376c1de49b8687c4f882d3bcf3	-2.1106948	17	N.S.	*Enterobacteriaceae*
606aa8c7117f801dc5ccc8c2a78c68c3	-1.7000214	16	N.S.	*Commensalibacter*
1c735b8a08dcbd5ca158987733234114	-1.6719837	15	N.S.	*Algoriella xinjiangensis*
da74ac9885f0aa2d29ec4338b46aa051	2.00310762	13	N.S.	*Orbaceae*
4043baca9447b9860efabab56603efc0	0.79635173	10	N.S.	*Acetobacter species*
61fd19eda2dee9c93613432491556a99	0.69392446	10	N.S.	*Alphaproteobacteria*
7070daee4850cf801719eab99ce88a82	0.96974134	10	N.S.	*Commensalibacter*
b3ccbf1d4fc85411f948429591f371cf	0.59968784	10	N.S.	*Pseudomonas*
ca7279bfdcb977287193c567f96b025c	0.75112887	10	N.S.	*Acetobacter indonesiensis*
0a04cded1e993516d5cdc851bfe07180	-1.6048832	9	N.S.	*Algoriella xinjiangensis*
6e4961fe480153e97ebdcf8fca4913fb	0.99515931	9	N.S.	*Providencia*
8dc187c9810636b7199c030caaaf0694	0.58697562	9	N.S.	*Morganella*
9f9bb5e1a63c5a0820d1571839534a2f	0.62165362	9	N.S.	*Commensalibacter*
677d95c5a0bc1c25435861fe262cb487	0.50699766	8	N.S.	*Alphaproteobacteria*
4af26492bbe6d0d4980bc82c2f08f32d	0.3940931	7	N.S.	*Providencia*
ad8b60b9428bc76f3b50e428f4f5a29f	0.41862229	7	N.S.	*Stenotrophomonas*
f72576c61dde7b9b6ad683dea71b105f	0.06411428	7	N.S.	*Enterobacteriaceae*
3c3c65e801e07d1be83bfa6d962b5627	-0.3759816	6	N.S.	*Commensalibacter*
681de033774ada405895dd81abc7593f	-0.5750105	6	N.S.	*Providencia rettgeri*
97b76e52dafa224603888fb56ef0b831	-0.3031253	5	N.S.	*Commensalibacter*

Supp. Table 4: Comparison of microbial community composition (Bray-Curtis dissimilarity) for recipients on C diets using PERMANOVA with 999 permutations.

Category	Df	SS	R^2	F	P value
Fly genotype	1	0.683	0.021	5.220	0.007
Microbiome	1	12.415	0.380	94.938	0.001
Fly genotype x Microbiome	1	0.506	0.015	3.871	0.018
Residual	146	19.092	0.584		
Total	149	32.695			

Supp. Table 5: Comparison of microbial community composition for recipients on HS diets using PERMANOVA with 999 permutations.

Category	Df	SS	R^2	F	P value
Fly genotype	1	1.827	0.065	8.5721	0.001
Microbiome	1	1.159	0.041	5.439	0.003
Fly genotype x Microbiome	1	1.261	0.045	5.915	0.004
Residual	112	23.869	0.849		
Total	115	28.116			

Supp. Table 6A: Analysis of deviance for Cox model with interactions between fly genotype, microbiome, and diet for developmental timing in the *Acetobacter* transplant.

Category	Log-likelihood	X^2	Df	*P* value
Null	-60640			
Fly genotype	-60584	111.642	1	<2.20E-16***
Acetobacter	-60517	134.874	2	<2.20E-16***
Diet	-58732	3568.585	1	<2.20E-16***
Fly x Acetobacter	-58675	115.244	2	<2.20E-16***
Fly x Diet	-58658	34.288	1	4.75E-09***
Acetobacter x Diet	-58523	268.613	2	<2.20E-16***
Fly x Acetobacter x Diet	-58500	47.625	2	4.55E-11***

Significance codes: '***' <0.001; '**' <0.01; '*' <0.05

Supp. Table 6B: Model selection statistics for developmental timing in the *Acetobacter* transplant. Model 1 is without interactions, while Model 2 includes all two and three-way interactions.

Model	Log-likelihood	X^2	X^2 Df	P(>|Chi|)
Model 1	-58732			
Model 2	-58500	465.77	7	<2.20E-16***

Significance codes: '***' <0.001; '**' <0.01; '*' <0.05

Supp. Table 7A: Fixed effects for fecundity in the *Acetobacter* transplant. Significance was evaluated using Type III Wald F tests with Kenward-Roger degrees of freedom.

Category	X^2	Df	*P value*
Intercept	637.8446	1	<2.20E-16***
Fly genotype	4.0333	1	0.0446106*
Acetobacter	39.7839	2	2.30E-09***
Diet	53.9716	1	2.03E-13***
Fly x Acetobacter	7.787	2	0.0203736*
Fly x Diet	8.4003	1	0.0037517**
Acetobacter x Diet	14.4492	2	0.0007285***
Fly x Acetobacter x Diet	16.3387	2	0.0002832***

Significance codes: *'***' <0.001; '**' <0.01; '*' <0.05*

Supp. Table 7B: Model selection statistics for fecundity in the *Acetobacter* transplant. Model 1 is without interactions, while Model 2 includes all two and three-way interactions.

Model	Df	AIC	BIC	logLik	deviance	X^2	X^2 Df	*P value*
Model 1	7	6333.5	6365	-3159.8	6319.5			
Model 2	14	6296.8	6359.8	-3134.4	6268.8	50.753	7	1.03E-08***

Significance codes: *'***' <0.001; '**' <0.01; '*' <0.05*

Supp. Table 8: Average nucleotide identity (ANI) between C and HS *Acetobacter* strains.

Metric	Value
ANI value (%)	73.68
C Acetobacter genome length (bp)	3,436,380
HS Acetobacter genome length (bp)	2,895,396
Average aligned length (bp)	1,055,729
C Acetobacter coverage (%)	30.72
HS Acetobacter coverage (%)	37.10

Supp. Table 9: Uric acid degradation presence/absence in the most abundant bacteria from the C and HS microbiomes. Annotation was performed using RAST, and each were compared to the HS *Acetobacter* (Y= present, N= not present). Only HS *Acetobacter* encoded for uric acid degradation pathways.

Role	HS Aceto	C Aceto	A. persici	Algoriella
Urea ABC transporter, ATPase protein UrtD	Y	N	N	N
Urea ABC transporter, ATPase protein UrtE	Y	N	N	N
Urea ABC transporter, permease protein UrtB	Y	N	N	N
Urea ABC transporter, permease protein UrtC	Y	N	N	N
Urease accessory protein UreE	Y	N	N	N
Urease accessory protein UreF	Y	N	N	N
Urease accessory protein UreG	Y	N	N	N
Urease alpha subunit (EC 3.5.1.5)	Y	N	N	N
Urease beta subunit (EC 3.5.1.5)	Y	N	N	N
Urease gamma subunit (EC 3.5.1.5)	Y	N	N	N

Supp. Table 10: ANOVA results for uric acid degradation by *Acetobacters*.

Category	Sum Sq	Df	F value	Pr(>F)
Intercept	12504.2	1	1188.4315	<2.2E-16***
Acetobacter	374.6	2	17.8035	1.82E-05***
Diet	1.5	1	0.1442	0.7075
Acetobacter x Diet	810.9	2	38.534	3.22E-08***
Residuals	252.5	24		

Significance codes: '***' <0.001; '**' <0.01; '*' <0.05