Dorsal raphe nucleus to anterior cingulate cortex 5-HTergic neural circuit modulates

consolation and sociability

Lai-Fu Li[1,2,*], Li-Zi Zhang[1,*], Zhi-Xiong He[1], Yu-Ting Zhang[1], Huan Ma[1], Yu-Feng Xun[1], Wei

Yuan[1,3], Wen-Juan Hou[1], Yi-Tong Li[1], Zi-Jian Lv[1], Rui Jia[1], Fa-Dao Tai[1,#]

[1]Institute of Brain and Behavioral Sciences, College of Life Sciences, Shaanxi Normal University, Xi'an,

710062, China

[2]College of Life Sciences, Nanyang Normal University, Nanyang, 473061, China

[3]Provincial Key Laboratory of Acupuncture and Medications, Shaanxi University of Chinese Medicine,

Xianyang 712046, China

Running title: DR-ACC 5-HTergic neural circuit modulate consolation

*The first two authors contributed equally to this article.

#Corresponding authors: Fa-Dao Tai, PhD, Professor, College of Life Sciences, Shaanxi Normal

University, Xi'an, 710062, China. E-mail: taifadao@snnu.edu.cn

ABSTRACT

Consolation is a common response to the distress of others in humans and some social animals,

but the neural mechanisms underlying this behavior are not well characterized. By using socially

monogamous mandarin voles, we found that optogenetic or chemogenetic inhibition of

5-HTergic neurons in the dorsal raphe nucleus (DR) or optogenetic inhibition of 5-HT terminals

in the anterior cingulate cortex (ACC) significantly decreased allogrooming time in the

consolation test and reduced sociability in the three-chamber test. The release of 5-HT within the

ACC and the activity of DR neurons were significantly increased during allogrooming, sniffing

and social approaching. Finally, we found that the activation of 5-HT1A receptors in the ACC

was sufficient to reverse consolation and sociability deficits induced by the chemogenetic

inhibition of 5-HTergic neurons in the DR. Our study provided first direct evidence that

DR-ACC 5-HTergic neural circuit is implicated in consolation-like behaviors and sociability.

Keywords: Consolation; Empathy; Anterior cingulate cortex; Dorsal raphe nuclei; Serotonin;

Fiber photometry

INTRODUCTION

Consolation behavior, which is referred to as an increase in affiliative contact toward a distressed individual by an uninvolved bystander, is an important component of the social capabilities of humans (de Waal and Preston, 2017; Field et al., 2009). Impaired consolation has been frequently observed in many psychiatric diseases, such as depression, autism, and schizophrenia (Young et al., 2015). According to de Waal's multilevel conceptualization of empathy (de Waal, 2008; de Waal and Preston, 2017), consolation represent an intermediate level of empathy (the primary level of "emotional contagion", the more complex level of "consolation", and the most elaborate level of "perspective taking and targeted helping"), which has long been assumed to exist in species possessing complex cognitive functions, such as humans, apes, dolphins and elephants (Perez-Manrique and Gomila, 2018); however, recent studies have indicated that it also exists in some socially lived rodents, such as prairie voles (Burkett et al., 2016), mandarin voles (Li et al., 2019) and rats (Knapska et al., 2010).

Currently, studies of the neural mechanisms underlying consolation and other forms of empathy have primarily focused on the oxytocin systems (Burkett et al., 2016; Li et al., 2019). However, as a complex social behavior, consolation may require the coordinated actions of numerous neuromodulators and neurotransmitters. Serotonin (5-HT) is an evolutionarily ancient neurotransmitter that has long been implicated in a variety of emotional disorders (Faye et al., 2020; Garcia-Garcia et al., 2018; Meneses and Liy-Salmeron, 2012). According to recent studies, 5-HT transmission is also involved in a series of social behaviors such as social interaction (Walsh et al., 2018), social reward and aggression (Dolen et al., 2013). Regarding empathy, a recent study revealed an association between salivary 5-HT levels and the empathic abilities of people (Matsunaga et al., 2017); a polymorphism in the promoter region of the serotonin transporter gene has been linked to individual differences in empathy (Gyurak et al., 2013); and MDMA (±3,4-methylenedioxymethamphetamine, better known as the recreational drug ''ecstasy''), which is well known to stimulate a feeling of closeness and empathy in its users (Carlyle et al., 2019), had been confirmed to robustly increase the release of 5-HT in an

activity-independent manner (Heifets and Malenka, 2016). In animal studies, Kim, et al. found

that microinjection of 5-HT into the anterior cingulate cortex (ACC) impairs vicarious fear and

alters the regularity of neural oscillations in mice (Kim et al., 2014). Our recent study indicated

that 5-HT1A receptors within the ACC are involved in consolation deficits induced by chronic

social defeat stress in mandarin voles (Li et al., 2020). However, to our knowledge, direct

evidence for an association between 5-HT and consolation has yet to be obtained.

Dorsal raphe nucleus (DR) is a main source of 5-HT neurons and provide 70% of 5-HTergic

projections in the forebrain (Fu et al., 2010; Luo et al., 2015). DR 5-HTergic neurons form dense,

broad and bidirectional neural connections with a broad range of forebrain and limbic structures,

including the ACC (Celada et al., 2013; Charnay and Leger, 2010), which is a central hub for

various types of empathy. Therefore, direct modulation of the DR→ACC 5-HTergic circuit to

investigate its function role in consolation-like behaviors is interesting and meaningful.

The released 5-HT binds to pre- and postsynaptic receptors. To date, at least 14 different

5-HT receptor subtypes have been identified in the brain (Artigas, 2013). Among which, 5HT1AR

and 5HT2AR are the two main subtypes that are expressed at high levels in the prefrontal cortex

(Carhart-Harris and Nutt, 2017; Santana and Artigas, 2017). The distribution, signaling pathways

and functions of these two receptors are substantially different, and both receptors play critical

roles in modulating cortical activity and neural oscillations (Celada et al., 2013). Previous studies

indicated that 5HT2AR gene single nucleotide polymorphisms are associated with

empathy-related social communication abilities (Gong et al., 2015), and a 5HT2AR agonist

increases emotional empathic ability (Dolder et al., 2017). However, in animal studies by Kim, et

al., blockade of serotonin receptors in the ACC did not affect observational fear responses in

mice (Kim et al., 2014). Clearly, the specific functions of 5HT1AR and 5HT2AR in empathy-like

behaviors still require further examination.

The mandarin vole (*Microtus mandarinus*) is a socially monogamous rodent that is widely

distributed across China (He et al., 2019). As shown in our previous studies, this species is

capable of displaying consolation-like behaviors upon exposure to a distressed partner (Li et al.,

2020; Li et al., 2019). In the present study, we first investigated the function of DR→ACC

5-HTergic circuits in consolation-like behavior using optogenetic and chemogenetic approaches.

To provide more direct evidence, we then monitored ACC 5-HT release and DR neuron activities

during this behavior by using *in vivo* fiber photometry. Finally, we used chemogenetics plus

pharmacological approaches to investigate which types of 5-HT receptors in the ACC are

involved in consolation-like behaviors in mandarin voles. In order to investigate any potential

sex differences during these processes, both male and female subjects were included in our study.

As consolation is in general a pro-social behavior, some social behaviors were investigated

during the tests.

RESULTS

Optogenetic inhibition of DR 5-HT neurons in the DR→ACC neural circuit impaired

consolation and reduced sociability

We first determined the 5-HTergic projection relationship between the DR and ACC in mandarin

voles. For this experiment, the retrograde tracer CTB was injected into the ACC, followed by

immunofluorescence staining of the DR sections with TPH2, a marker of 5-HTergic neurons. A

substantial number of TPH2+ neurons colocalized with CTB, indicating the presence of DR-ACC

5-HTergic projections (Figure 1—figure supplement 1). We then used a novel dual-virus

optogenetics approach to explore the function of the DR-ACC 5-HTergic circuit in

consolation-like behaviors and sociability, where double-floxed AAV-DIO-ChR2-mCherry

(DIO-ChR2) or AAV-DIO-eNpHR3.0-mCherry (DIO-eNpHR3.0) was injected into the DR and

retro-AAVs containing the TPH2 promoter and Cre element (rAAV(Retro)-TPH2-Cre) were

injected into the ACC (Figure 1A). This virus strategy ensures that opsins (excitatory CHR2 or

inhibitory eNpHR3.0) are mainly expressed within the DR-ACC 5-HTergic circuit.

Immunohistochemical staining showed that more than 75% of mCherry-labeled neurons

expressed TPH2 in both male and female voles, and more than 80% of TPH2+ cells coexpressed

mCherry (Figure 1—figure supplement 2). In the electrophysiological study, we found that the

DR neurons reliably responded to pulses of 473 (activation)/593 (inhibition) nm light stimuli (Figure 1D-E). These results indicate the viability of this virus strategy.

To test whether modulation of DR 5-HT neuron activity alters consolation and sociability, five weeks after the virus injection, optic fibers were implanted above the DR (Figure 1A-B). The virus expression sites and the optic fiber placement schematics are shown in Figure 1—figure supplement 3A. According to Burkett's and our previous results (Burkett et al., 2016; Li et al., 2019), the time spent grooming their stressed partners (allogrooming) is an important indicator of consolation-like behaviors. We then exposed the subjects to their electric-shocked partners and recorded the time spent allogrooming, chasing (closely following) and selfgrooming (consolation test; for details, please refer to the Methods section). The three-chamber test was used to assay the sociability along with open-field anxiety and locomotion (Figure 1B). In the CHR2-expressing animals, there was no conclusive evidence that optogenetic activation of 5-HTergic neurons affected their time spent allogrooming, chasing and self-grooming in the 10-min consolation test (Figure 1G, Figure 1—figure supplement 4A-D), the social interaction ratio in the 3-chamber test (Figure 1I), and behavioral performance in the open-field test (Figure 1K, Figure 1—figure supplement 4E).

However, in the eNPHR3.0-expressing animals, there was extremely strong evidence that optogenetic inhibition of 5-HT neurons in the DR reduced the time spent allogrooming (Figure 1H) and chasing (Figure 1—figure supplement 4B), but had little effect on the control behavior self-grooming (Figure 1—figure supplement 4D). Following experiments indicated that the light-inhibition effects significantly reduced within 24 h (Figure 1—figure supplement 5). In the 3-chamber test, light inhibition also significantly reduced the social preference ratio (Figure 1J). In the open-field test, the results were inconclusive, which suggests that the data for the time spent in the central area and the total distance travelled are equally likely under H0 and H1 (Figure 1L, Figure 1—figure supplement 4F).

The above results indicate that although we have no conclusive results for optogenetic activation, optogenetic inhibition of 5-HT neurons within the DR→ACC 5-HTergic circuit

impaired consolation-like behavior and sociability in mandarin voles.

Figure 1. Optogenetic bidirectional modulation of 5-HT neuron in the DR in the DR-ACC

neural circuit. (A): Schematic of optogenetic manipulation; (B): timeline of experiments; (C):

immunohistological image showing virus expression in the DR (c1, × 100) and amplified images

in the left box showing the mCherry, TPH2 and the colocaliztion of the two ('c2-c4', × 200); (D):

electrophysiological recordings model; (E & F): representative traces from electrophysiological

recordings showing photostimulation (E) and photoinhibition of a 5-HT neuron (F); (G-L):

quantification of allogrooming time in the consolation test (G & H); social preference ratio in the

three-chamber test (I & J) and time spent in the central area in the open-field test (K & L) during

optogenetic modulations. Data are presented as mean ± SE, $n = 7$–8 in each group, **$P < 0.01$. For

'G-H', independent samples t tests along with Bayesian independent samples t-test; for 'I-L',

two-way repeated measures ANOVA along with two-way repeated Bayesian measures ANOVA

(light as within subject factor). For raw data in this figure, please refer to Figure 1—source data 1.

For detailed quantifications, please refer to Supplementary 2. ACC: anterior cingulate cortex;

CTR: control; DR: dorsal raphe nucleus; TPH2: tryptophan hydroxylase 2.

Optogenetic inhibition of ACC 5-HT terminals in the DR→ACC neural circuit similarly

impaired consolation and reduced sociability

Modulation of 5-HT neurons may affect other neurons in the DR, and thus confound behavioral

performance. In subsequent experiments, we placed optic fibers in the ACC and investigated

whether the direct modulation of 5-HT terminals in this region would exert the same effects

(Figure 2A). Similarly, we found that optogenetic activation of DR-ACC 5-HT terminals did not

significantly alter behavioral performance in the consolation test (Figure 2C, Figure 2—figure

supplement 1A, 1C) and open-field test (Figure 2G, Figure 2—figure supplement 1E). In the

3-chamber test, two-way repeated measures ANOVA revealed a moderate 'light × group'

interaction in females ($F_{(1,14)} = 6.297$, $P = 0.025$, $BF_{incl} = 4.186$). Subsequent *post hoc* comparison

results showed that light stimulation slightly increased the social preference ratio in

CHR2-expressing females but not in mCherry-expressing females (ChR2: off *vs.* on, $P = 0.033$;

mCherry: off *vs.* on, $P = 0.975$; Figure 2E). The 'light × group' interaction was not evidently

observed in males ($F_{(1,14)} = 3.556$, $P = 0.08$, $BF_{incl} = 1.239$).

Optogenetic inhibition of DR-ACC 5-HT terminals significantly reduced the time spent

allogrooming and chasing in the consolation test (Figure 2D, Figure 2—figure supplement 1B)

and the social preference ratio in the 3-chamber test (Figure 2F). Similarly, the inhibition had

little effect on the control behavior of selfgrooming (Figure 2—figure supplement 1D) and

behavioral performance in open-field test (Figure 2H, Figure 2—figure supplement 1F) and the

inhibitory effect lasted no more than 24 h (Figure 2—figure supplement 2). The above results

indicated that optogenetic modulation of ACC 5-HT terminals have nearly identical effects as

modulation of DR 5-HT neurons.

Figure 2. Optogenetic bidirectional modulation of 5-HT terminals within the ACC in the

DR-ACC neural circuit. (A): Schematic of optogenetic manipulation; (B): timeline of experiments;

(C-H): quantification of time spent in allogrooming in the consolation test (C & D), sociability in

the three-chamber test (E & F) and time spent in the central area in the open-field test (G & H)

during optogenetic modulations. Data are presented as mean ± SE, *n* = 7–8 in each group, *P* <

0.05, **P* < 0.01. For C-D, independent samples *t* tests along with Bayesian independent samples

t-test; for E-H, two-way repeated measures ANOVA along with two-way repeated Bayesian

measures ANOVA (light as within subject factor). For raw data in this figure, please refer to

Figure 2—source data 1. For detailed quantifications, please refer to Supplementary 2. ACC:

anterior cingulate cortex; DR: dorsal raphe nucleus; CTR: control.

Chemogenetic inhibition of DR 5-HT neurons in the DR→ACC neural circuit impaired

consolation and reduced sociability

To confirm the above optogenetic results over a longer time frame, we used a chemogenetic

approach to selectively express 'Gq-DREADD' or 'Gi-DREADD' in DR 5-HT neurons by injecting

AAV-DIO-hM4Dq-mCherry (Gq-DREADD) or AAV-DIO-hM4Di-mCherry (Gi-DREADD) into

the DR and rAAV(Retro)-TPH2-Cre into the ACC (Figure 3A). Immunohistochemical staining

revealed that more than 65% of TPH2 labeled neurons were infected by mCherry virus and more

than 60% of mCherry cells coexpressed TPH2 (Figure 3—figure supplement 1). The virus

expression sites schematics were shown in Figure 1—figure supplement 3B. To determine

whether the ligand CNO can activate or inhibit DR 5-HT neurons, whole-cell current-clamp

recordings were performed. The results showed that the addition of 10 μM CNO remarkably

increased the number of action potentials in the Gq-DREADD-transfected neurons (Figure 3D).

In contrast, CNO caused a significantly decrease in the number of spikes (Figure 3E) and

increased the spike rheobase during current step injections in Gi-DREADD-transfected neurons

(Figure 3F). These results indicate the specificity and viability of this virus strategy.

In subsequent behavioral studies, we found CNO (1 mg/kg, i.p.) treated

Gi-DREADD-expressing voles showed reduced grooming toward their shocked partners in the

consolation test (Figure 3G) and decreased sociability in the 3-chamber test (Figure 3H). The

CNO treated males also showed a trend of spending less time in the open-field test, but the

statistics were inconclusive (Figure 3I; treatment: $F_{(1,19)}$ = 4.305, P = 0.052, BF(incl) = 4.438; Group:

$F_{(2,19)}$ = 2.064, P = 0.155, BF(incl) = 0.384; treatment × Group: $F_{(1,19)}$ =0.164, P = 0.850, BF(incl) =

0.300). The treatment had no detectable effects on the total distance travelled in the open-field

test (Figure 3—figure supplement 2C) and some control behaviors in the consolation test (Figure

3—figure supplement 2A&B). In Gq-DREADD and virus control subjects, CNO treatment had no

detectable effects on the behavioral performance in all the test (Figure 3).

Based on the results of the chemogenetic and optogenetic experiments described above, we concluded that inhibition of DR-ACC 5-HTergic circuit activity was sufficient to impair consolation and sociability in mandarin voles.

Figure 3. Chemogenetic modulation of DR 5-HT neuron activities in the DR-ACC neural

circuit. (A): Schematic of chemogenetical manipulations; (B): timeline of experiments; (C):

immunohistological image showing virus expression in the DR (c1, × 100) and amplified images

in the left white box showing the mCherry, TPH2 and the colocaliztion of the two (c2-c4, × 200);

(D): representative trace from a Gq-DREADD neuron; (E): representative trace from a

Gi-DREADD-transfected neuron; (F): quantification of spike rheobase and spike numbers under

current step injections in Gi-DREADD-transfected neurons (n = 5 neurons, *P < 0.05); (G-I):

quantification of allogrooming time in the consolation test (G); social preference ratio in the

three-chamber test (H) and time spent in the central area in the open-field test (I) in male (upper

panels) and female (down panels) voles. Data are presented as mean ± SE, n = 7–8 in each group,

*P < 0.05, **P < 0.01 compared with vehicle control. For 'F', one tailed paired t test along with one

tailed Bayesian paired samples t-test; for 'G-I', two-way repeated measures ANOVA along with

two-way repeated Bayesian measures ANOVA. For raw data in this figure, please refer to Figure

3—source data 1. For detailed quantifications, please refer to Supplementary 2. ACC: anterior

cingulate cortex; DR: dorsal raphe nucleus; TPH2: tryptophan hydroxylase 2; M: male; F: female.

DR 5-HT neuron activity increased during allogrooming and social approaching

If the DR-ACC 5-HTergic circuit is involved in consolation and sociability, ACC projecting DR

5-HT neurons may change their activity during the corresponding behaviors. To verify this idea,

we performed photometric recording in freely moving animals by injecting Cre-inducible

AAV-DIO-GCaMp6, a genetically encoded fluorescent Ca^{2+} sensor, into the DR and

rAAV(Retro)-TPH2-Cre into the ACC (Figure 4A). Ten days later, fibers were implanted above

the DR injection sites. Post hoc histological analysis revealed that more than 80% $GCaMp6^+$ cells

overlapped with $TPH2^+$ expressing cells (Figure 4—figure supplement 1). The virus expression

sites and the optic fiber placements schematics were shown in Figure 1—figure supplement 3C.

During the consolation test, we found that the activity of DR 5-HT neurons was reliably

increased during allogrooming, social approaching and sniffing in both sexes (Figure 4E1-E2,

F1-F2 and G1-G2). DR 5-HT neurons showed the highest response during allogrooming which

may indicate the specific role of 5-HT in consolation-like behaviors (Figure 4—figure supplement

2A-2B). Interesting, the increased Ca^{2+} in DR 5-HT neurons often preceded these behaviors

which may indicate the motivational function of 5HT (Arakawa, 2020; Yagishita, 2020). When

aligning the fluorescence changes to the end of these social bouts, we found that the activity of

5-HT neurons decreased accordingly before withdrawal from allogrooming and approaching

(Figure 4—figure supplement 3A-3B). Withdrawing from sniffing also showed a similar trend

but was not statistically significant (Figure 4—figure supplement 3C; pre *vs*. post: male: $t_{(5)} = 1.611$,

$P = 0.084$, BF+0 = 1.589 with median posterior $\delta = 0.528$, 95%CI = [0.041, 1.363]; female: $t_{(5)} = 1.724$, P

$= 0.073$, BF+0 = 1.778 with median posterior $\delta = 0.557$, 95%CI = [0.046, 1.415]). The increased activity

of DR 5-HT neurons did not simply reflect movement initiation as no fluorescence changes were

observed when aligning to running and selfgrooming (Figure 4H1-H2, I1-I2). Furthermore, no

significant changes in fluorescence signals were detected in DR 5-HT neurons of GFP-expressing

control voles during all the behaviors, confirming that the GCaMp6 signals genuinely indicated

neuronal activity and did not simply reflecting motion artifacts (Figure 4E3-E4, F3-F4, G3-G4,

H3-H4 and I3-I4).

The above results provide direct evidence that DR 5-HT neurons within the DR-ACC

5-HTergic circuit are recruited in consolation-like behavior and sociability in mandarin voles.

Figure 4. Fiber photometry recording DR 5-HT neural dynamics during the consolation test.

(A): Schematic diagrams depicting the virus injection and recording sites; (B): histology showing

the expression of GCaMP6 (left) and GFP control (right) in the DR (× 200); (C): experimental

timeline for photometry experiments; (D): representative fluorescence changes of GCaMP6 (red

line) and GFP (blue line) during photometry recordings; (E1-I1): representative peri-event plot of

GCaMP6 fluorescence signals aligned to onsets of various behavior (for all peri-event plots, the

red line denotes the mean signals of 4-6 bouts of behaviors, whereas the red shaded region

denotes the SE); (E2-I2): quantification of change in GCaMP6 fluorescence signals before and

after the events (*n* =6 in each group); (E3-I3): representative peri-event plot of GFP signals

aligned to onsets of various behavioral events (for all peri-event plots, the blue line denotes the

mean signals of 4-6 bouts of behaviors, whereas the blue shaded region denotes the SE); (E4-I4):

quantification of change in GFP fluorescence signals before and after the events (n = 5 in each

group). Data are presented as mean ± SE, *P < 0.05, **P < 0.01, Paired samples t-test along with

Bayesian paired samples t-test. For raw data in this figure, please refer to Figure 4—source data 1.

For detailed quantifications, please refer to Supplementary 2. ACC: anterior cingulate cortex;

TPH2: tryptophan hydroxylase 2; Allo: allogrooming; Sni: sniffing; App: approaching; Gro:

selfgrooming; Run: running.

ACC 5-HT release increased during allogrooming, social approach and sniffing

To further establish the potential relevance of 5-HT in consolation-like behaviors, we directly

detected the dynamics of 5-HT within the ACC using a newly developed GPCR-activation-based

5-HT sensor along with photometric recording (Wan, *et al* 2020). To do this, we injected AAV

encoding the fluorescent 5-HT sensor (AAV-5HTsensor 2.1) unilaterally into the ACC and

implanted optic fibers above the injection site ten days later (Figure 5A). The virus expression

sites and the optic fiber placements schematics were shown in Figure 1—figure supplement 3D.

Control animals were injected with AAVs encoding eGFP. Post hoc histological verification

indicated expression of both the 5-HT sensor and eGFP within the ACC (Figure 5B).

Next, we measured the dynamics of endogenous 5-HT during the consolation test.

Consistent with the above fiber photometry of Ca^{2+} signals in DR 5-HT neurons, we found

reliably increased fluctuations in fluorescence within the ACC in conjunction with allogrooming,

social approaching and sniffing, but not observed with self-grooming and running (Figure 5, left

columns). Similarly, the peek 5-HT signals during allogrooming were larger than during the

other behaviors (Figure 4—figure supplement 2C-2D), and the release of 5-HT significantly

dropped before the end of allogrooming, approaching and sniffing (Figure 5—figure supplement

1). In control animals that expressed GFP in the ACC, we observed little fluorescence change

during any of these behaviors (Figure 5E3-E4, F3-F4, G3-G4, H3-H4 and I3-I4).

To validate the selectivity of this 5-HT sensor in mandarin voles, in another set of animals

expressing the 5-HT sensor, a 5-HT receptor antagonist metergoline (Met) was injected (i.p., 4

mg/kg) before conducting the fiber photometry experiments. We found that treating voles with

Met completely blocked the fluorescence changes of the 5-HT sensor during allogrooming

(Figure 5—figure supplement 2), validating the viability and specificity of this 5-HT sensors.

 The above results provide further evidence that 5-HT released within the ACC may play an

important role in consolation-like behavior and sociability in mandarin voles.

Figure 5. Fiber photometry recording dynamics of 5-HT within the ACC during the

consolation test. (A): Schematic diagrams depicting the virus injection and recording sites; (B):

histology showing the expression of 5-HT sensor (left) and GFP control (right) within the ACC (×

200); (C): experimental timeline for photometry experiments; (D): representative fluorescence

changes of 5-HT sensor (red line) and GFP (blue line) during photometry recordings; (E1-I1):

representative peri-event plot of 5-HT fluorescence signals aligned to onsets of various behavior

(for all peri-event plots, the red line denotes the mean signals of 4-6 bouts of behaviors, whereas

the red shaded region denotes the SE); (E2-I2): quantification of change in 5-HT fluorescence

signals before and after the events (n = 6 in each group); (E3-I3): representative peri-event plot of

GFP signals aligned to onsets of various behavioral events (for all peri-event plots, the blue line

denotes the mean signals of 4-6 bouts of behaviors, whereas the blue shaded region denotes the

SE); (E4-I4): quantification of change in GFP fluorescence signals before and after the events (n =

5 in each group). Data are presented as mean ± SE, *P < 0.05, **P < 0.01; paired samples t-test

along with Bayesian paired samples t-test. For raw data in this figure, please refer to Figure

5—source data 1. For detailed quantifications, please refer to Supplementary 2. ACC: anterior

cingulate cortex; M: motor cortex; TPH2: tryptophan hydroxylase 2; Allo: allogrooming; Sni:

sniffing; App: approaching; Gro: selfgrooming; Run: running.

Serotonin in the ACC mediated consolation-like behaviors through 5-HT1A receptors

The most abundant 5-HT receptors expressed in the mPFC are 5-HT1AR and 5-HT2AR

(Carhart-Harris and Nutt, 2017). Increasing evidence indicates that these two types of receptors

exert opposite effects within the mPFC (Celada et al., 2013; Puig and Gulledge, 2011). In our

previous study, we indicated that ACC administration of a 5HT1AR agonist (8-OH-DPAT)

rescued the impaired consolation and sociability induced by social defeat (Li et al., 2020). To

further examine which receptor mediates consolation-like behaviors within the ACC, we directly

infused 8-OH-DPAT (200 nL/side; 1.5 mg/mL) or a 5-HT2AR antagonist (MDL 100907, 200

nL/side; 1 mg/mL) into the ACC along with chemogenetic inhibition of DR 5-HT neurons

before conducting behavioral assays (Figure 6A & B). Not surprisingly, CNO elicited deficits in

allogrooming and sociability (Figure 6C, D, F, G: left two bars). Pretreatment with 8-OH-DPAT

significantly reduced the impact of CNO (Figure 6C, D: right two bars; see also the comparison

results of 'Vehicle+Saline - Vehicle+CNO' *vs.* '8-OH-DPAT+Saline - 8-OH-DPAT+CNO'; Figure

6—figure supplement 1A, B), whereas MDL 100907 had no such effect (Figure 5F, G: right two

bars; Figure 6—figure supplement 1C, D). Neither the drugs nor viruses had any detectable

effect on the control behavior of self-grooming or behavioral performance in the open-field test

(Figure 6E-H, Figure 6—figure supplement 2).

Figure 6. Intra-ACC injection of 8-OH-DPAT reduced sociability deficits induced by

chemogenetic inhibition of DR 5-HT neurons in the DR→ACC neural circuit. (A): Schematic

representation of ACC infusion sites and virus strategy; (B): timeline of experimental design;

(C-E): effect of a 5-HT1AR agonist 8-OH-DPAT on allogrooming time in the consolation test (C),

social preference ratio in the three-chamber test (D), and time spent in the central area in the

open-field test (E) in male (left panels) and female (right panels) voles; (F-H): effect of a 5-HT2AR

antagonist (MDL 100907) on allogrooming time in the consolation test (F), social preference ratio

in the three-chamber test (G) and time spent in the central area in the open-field test (H) in male

(left panels) and female (right panels) voles. Data are presented as mean ± SE, n = 5–6 in each

group; two-way ANOVA along with two-way Bayesian ANOVA; *P < 0.05, **P < 0.01 for

'vehicle + saline' *vs.* 'vehicle + CNO'; ##P < 0.01 for 'MDL 100907 + CNO' *vs.* 'vehicle + saline'.

For raw data in this figure, please refer to Figure 6—source data 1. For detailed quantifications,

please refer to Supplementary 2. Anta: antangonist; ACC: anterior cingulate cortex; DR: dorsal

raphe nucleus.

DISCUSSION

In the present study, we demonstrated a crucial role for the DR→ACC 5-HTergic neural circuit

in the regulation of consolation-like behaviors and sociability in mandarin voles for the first time.

Our major findings are listed below. First, inhibition of DR 5-HT neurons or their terminals in

the ACC decreased allogrooming behavior and reduced sociability. Second, DR 5-HT neuron

activity and ACC 5-HT release increased during allogrooming, social approaching and sniffing.

Third, direct activation of ACC 5-HT1A receptors was sufficient to ameliorate deficits in

consolation and sociability induced by chemogenetic inhibition of DR 5-HT neurons.

We found optogenetic inhibition of DR 5-HT neurons (Figure 1) or ACC 5-HT terminals

(Figure 2) significantly decreased intimate behaviors (allogrooming and chasing) toward their

distressed partners and reduced sociability. Chemogenetic inhibition of DR 5-HT neurons

produced similar results (Figure 3). Consistent with our results, Walsh et al found that

optogenetic inhibition of DR 5-HT neurons reduced social interactions in the three-chamber test

(Walsh et al., 2018). As consolation is in general a pro-social behavior, it is difficult to determine

whether the reduced allogrooming is due to an overall decrease in sociability. However, in the

fiber photometry studies, we found both DR 5-HT neurons and ACC 5-HT release showed the

highest responses during allogrooming, which may indicate the specific role of 5-HT in

consolation (Figure 4—figure supplement 2).

Another open question is how this process occurs, namely, what are the neural mechanisms

underlying this process? The neocortical excitatory/inhibitory (E/I) balance hypothesis may

help address this question. The hypothesis indicates that an increase in the cortical cellular E/I

balance, for example through increased activity in excitatory neurons or a reduction in inhibitory

neuron function, is the common etiology and final pathway for some psychiatric diseases, such

as autism and schizophrenia (Bozzi et al., 2018; Vattikuti and Chow, 2010). This hypothesis has

recently been verified in mice, as optogenetic excitation of glutamatergic neurons in the mPFC

elicits a profound impairment in sociability, while compensatory excitation of inhibitory neurons

in this region partially rescues the social deficits caused by an increase in the E/I balance (Yizhar

et al., 2011). 5-HT tends to inhibit prefrontal pyramidal activity (Puig and Gulledge, 2011; Tian et

al., 2017). For example, according to Puig et al., electrical stimulation of the DR inhibits

approximately two-thirds of pyramidal neurons in the mPFC (Puig et al., 2005). Therefore, a

reasonable hypothesis is that inhibition of the DR-ACC 5-HTergic neural circuit may increase the

E/I balance, which ultimately leads to abnormal social behaviors in mandarin voles. Clearly, this

hypothesis should be verified in electrophysiological studies in the future.

In contrast to inhibition manipulations, we found activation of the DR-ACC 5-HTergic

neural circuit did not elicit corresponding increases in allogrooming and sociability (Figures 1-3).

One possible explanation is that there are various 5-HT receptors expressed in the ACC and

some have opposite functional effects (Puig and Gulledge, 2011; Tian et al., 2017). For example,

5-HT1AR coupled to the Gi family of G proteins induces hyperpolarization of pyramidal

neurons, whereas 5-HT2AR coupled to Gq proteins induces depolarization in the same neurons

(Carhart-Harris and Nutt, 2017). In addition, these two types of receptors are expressed in both

pyramidal and GABAergic neurons of the mPFC (Santana et al., 2004). Therefore, the net

behavioral effects of 5-HT release must result from the combined effects of all these receptors.

The other possible explanation is a ceiling effect, which means that the enhancement of

consolation and sociability are difficult to accomplish in these normal animals. This may occur in

situations where levels of consolation and sociability are low such as in the stressed subjects in

our previous study (Li et al., 2020) or genetically less sociable subjects as in Walsh's study (Walsh

et al., 2018). However, in a previous study by Kim et al., the direct administration of 5-HT into

the ACC impaired observational fear learning, which is an indicator of emotional contagion (Kim

et al., 2014). Species differences (mice *vs.* voles), methodological differences (pharmacology *vs.*

optogenetics and chemogenetics) or different behavioral indicators (consolation *vs.* emotional

contagion) may account for the discrepancies. Furthermore, optogenetic or chemogenetic

activation of DR 5-HT neurons is also distinct from pharmacological intervention with MDMA,

which robustly induces 5-HT releases in the whole brain and enhances closeness and empathy in

its users in human studies (Carlyle et al., 2019; Heifets and Malenka, 2016). Nevertheless, at least

some effects of activation were visible. For example, optogenetic activation of ACC 5-HT

terminals increased sociability in CHR2-expressing females (Figure 2E). Clearly, the effects of

activation of the DR-ACC 5-HTergic neural circuit on sociability and empathy still require

further in-depth study.

Our results show that neither activation nor inhibition the DR-ACC 5-HTergic neural circuit

exerts any significant effects on some control behaviors (Figure 1—figure supplement 4; Figure

2—figure supplement 1; Figure 3—figure supplement 2), i.e., self-grooming in the consolation

test or behavioral performance in the open-field test, which is consistent with pervious results

(for self-grooming, please refer to (Correia et al., 2017); for the open-field test, please refer to

(Walsh et al., 2018)). However, inconsistent results have also been reported. For example,

Ohmura et al. found that optogenetic activation of 5-HT neurons in the median raphe nucleus

enhanced anxiety-like behaviors in mice (Ohmura et al., 2014). In the study by Correia et al.,

activation of 5-HTergic neurons in the DR affected behavior in the open-field test, but not anxiety

(Correia et al., 2017). Different stimulation protocols or manipulations of different groups of

5-HT neurons may account for these discrepancies.

Our results obtained using *in vivo* fiber photometry indicated an increase in DR 5-HT

neuron activity during allogrooming, sniffing and social approaching (Figure 4). This result is

consistent with previous studies showing increases in the activity of 5-HTergic neurons in the

DR during nonaggressive social interactions (Li et al., 2016; Walsh et al., 2018). Furthermore,

using the highly sensitive 5-HT fluorescent sensors developed by Wan et al., (Wan et al, 2020),

we provided the first evidence that allogrooming, sniffing and social approaching elicit robust

5-HT release in the ACC (Figure 5). Self-grooming and running did not induce robust 5-HT

release within the ACC, which may indicate the behavioral relevance of the 5-HT sensors.

Interestingly, we found that the fluorescence changes in GCaMP6 and 5-HT sensors usually

precede allogrooming and social approaching and decreased accordingly before withdrawing

from them (Figure 4—figure supplement 3, Figure 5—figure supplement 1). These results

indicate that 5-HT may be involved in some motivational aspects of these social behaviors, which

is consistent with previous findings (Arakawa, 2020; Yagishita, 2020).

Although both 5-HT1AR and 5-HT2AR are expressed at high levels in the mPFC and work

together to modulate cortical network activity (Celada et al., 2013), we found only the 5-HT1AR

receptor agonist significantly reduced the consolation and sociability deficits induced by

chemogenetic inhibition of 5-HTergic neurons in the DR (Figure 6). This result is consistent with

our previous finding that the administration of WAY-100635 (a 5-HT1AR antagonist) into the

ACC attenuates consolation and sociability in mandarin voles (Li et al., 2020). 5-HT1AR within

the mPFC has frequently been reported to exert antidepressant and anxiolytic effects (Artigas,

2013; Fukumoto et al., 2020; Fukumoto et al., 2014; Wang et al., 2019), but our findings clearly

indicate that this type of 5-HT receptor within the ACC is also involved in regulating some social

behaviors. Clearly, this effect should be verified in many other species in future studies.

Administration of the 5-HT2AR antagonist into the ACC did not exert an obvious effect on the

social deficits induced by the chemogenetic inhibition of DR 5-HT neurons (Figure 5F-H). Thus,

5-HT2AR within the ACC may not play a major role in consolation and sociability. However,

this conclusion should be interpreted very cautiously because 5-HT2AR expression within the

mPFC shows clear rostral-to-caudal gradients in mice (Weber and Andrade, 2010) and we were

unable to determine its functional role in this dimension in the present study. Furthermore, the

effects of 5-HT in the ACC on consolation and sociability probably involve other subtypes of

5-HT receptors, such as 5-HT1BR and 5-HT3R, which require further examination.

In conclusion, our findings establish the importance of the DR-ACC 5-HTergic neural circuit

in consolation-like behaviors and sociability in mandarin voles. One major limitation of our

study is that although our virus strategies were generally successful, a few of the infected

neurons were indeed TPH2-negative, which may challenge the specificity of our manipulations

and confound our results. However, the fiber photometry and pharmacological studies mitigate

these concerns. Considering the widespread innervation of 5-HT terminals and abundant

distribution of 5-HT receptors in the whole brain, detailed knowledge of cellular and circuit

mechanisms of the 5-HTergic system, will not only improve our understanding of its

complicated features and functions but also have implications for the development of novel

therapies for the treatment of prevalent neuropsychiatric disorders, such as depression, autism,

and schizophrenia. Additionally, although both male and female subjects were included in this

study, sexually dimorphic effects were rarely observed. This finding may provide additional

evidence of cooperative evolution to adapt to environmental challenges, particularly in species

that adapt monogamous relationships and require cooperative breeding.

497 **METHODS AND MATERIALS**

498 **Key Resources Table**

Reagent type	Source	Identifier	Additional information
Antibodies			
Anti-TPH2 goat polyclonal antibody	Abcam	ab121013	1/500
Donkey anti-goat secondary antibody conjugated with Dylight 488	Jackson	705-545-147	1/300
Donkey anti-goat secondary antibody conjugated with TRITC	Jackson	705-025-003	1/300
DAPI	Boster	AR1177	Original fluid
Drugs and compounds			
8-OH-DPAT	Sigma	H8520	Injection site : ACC; volume: 200 nL/side; concentration: 1.5 mg/mL
MDL 100907	Tocris Bioscience	Cat#4173	Injection site : ACC; volume: 200 nL/side; concentration: 1 mg/mL
Metergoline	MedChem Express	Cat#HY-B1033/CS-4551	4mg/kg, i.p.
CNO	BrainVTA	Cat# CNO-02	1 mg/kg, i.p.
Cholera Toxin Subunit B (CTB)594	Thermo Fisher	Cat#C34777	Injection site : ACC ; volume : 400 nL/side ; expression time : 10 days
Optogenetic virus			
rAAV(Retro)-TPH2-Cre-WPRE-pA	BrainVTA	R-396-K181127	Injection site : ACC ; volume : 300 nL/side ; expression time : 5 wks
rAAV-Ef1α-DIO-ChR2-mC	BrainVTA	9-2-K190827	Injection site : DR;

herry-WPRE-pA			volume : 500 nL ; expression time : 5 wks
rAAV-Ef1α-DIO-eNpHR3.0-mCherry-WPRE-pA	BrainVTA	rAAV9-7-5-1	Injection site : DR; volume : 500 nL ; expression time : 5 wks
rAAV-Ef1α-DIO-mCherry-WPRE-pA	BrainVTA	9-13-K190430	Injection site : DR; volume : 500 nL ; expression time : 5 wks
Chemogenetic virus			
rAAV(Retro)-TPH2-Cre-WPRE-pA	BrainVTA	R-396-K181127	Injection site : ACC ; volume : 300 nL/side ; expression time : 5 wks
rAAV-Ef1α-DIO-hM4Di(Gi)-mCherry-WPRE-pA	BrainVTA	9-43-K190521	Injection site : DR; volume : 500 nL ; expression time : 5 wks
rAAV-Ef1α-DIO-hM4Dq(Gq)-mCherry-WPRE-pA	BrainVTA	9-42-K190530	Injection site : DR; volume : 500 nL ; expression time : 5 wks
Fiber photometry virus			
rAAV-hSyn-5HT2.1-WPRE-hGHpA (5-HT sensor)	BrainVTA	9-1826-K190620	Injection site : ACC ; volume : 400 Nl ; expression time : 3 wks
rAAV-hSyn-EGFP-WPRE-hGH-pA	BrainVTA	PT-1990	Injection site : ACC ; volume : 400 nL; expression time : 3 wks
rAAV-EF1α-DIO-GCaMp6-WPRE-hGH pA	BrainVTA	PT-0071	Injection site : DR; volume : 500 nL ; expression time : 4 wks
rAAV-hSyn-DIO-EGFP-WPRE-hGH-PA	BrainVTA	9-1103-K200409	Injection site : DR; volume : 500 nL; expression time : 4 wks
rAAV(Retro)-TPH2-Cre-WPRE-pA	BrainVTA	R-396-K181127	Injection site : ACC ; volume : 300 nL/side ; expression time : 4 wks

Animals

The mandarin voles used in this study were laboratory bred strains (F2-F3) whose ancestors derived from a wild population from Lingbao city (Henan, China). The voles were weaned on postnatal day 21, socially housed in same-sex in each polycarbonate cage (44×22×16 cm) and housed on a 12-h light/dark cycle with food and water ad libitum. Voles used for experiments were about 70-90 days old at the time. All breeding, housing, and experimental procedures were in accordance with Chinese guidelines for the care and use of laboratory animals and were approved by the Animal Care and Use Committee of Shaanxi Normal University.

Stereotaxic surgery and virus infusions

The kind of virus, total injection volumes and the expression time were listed in Key Resources Table. For surgery, voles (about 50 days) were anesthetized with 1.5%-3.0% isoflurane and placed in a stereotaxic instrument. Next, thirty-three-gauge syringe needles (Hamilton) were used for virus delivery. The injection rates were set at 50 nL/min. After each injection, the needle was left in the brain for another 5 min before being slowly withdrawn in order to prevent the virus from leaking out. The bregma coordinates for the virus injection were as follows: ACC: A/P: 1.6, M/L: 0.5, D/V: 1.6; DR: AP: -4.3; DV: -3.3, ML: +1.2, with a 20 angle toward the midline in the coronal plane.

Microinjection and drugs

The 5-HT1AR agonist 8-OH-DPAT was prepared in saline with a final concentration of 1.5 mg/mL. The 5-HT2AR antagonist MDL 100907 was prepared in 0.01 M PBS (adjust PH value with 0.1 M HCl to 6.4) with a final concentration of 1 mg/mL. All the microinjections were administered 30 min before the behavioral test. The speed of injection was 0.1 µL/min, and the total injection volume were 0.2 µL per side for all the drugs. The injector tips remained in situ for another 2 min for drug diffusions. The dose and timing of drug administration were chosen based on previous studies with 8-OH-DPAT (Cooper et al., 2008; Li et al., 2020) and MDL 100907

(Ishii et al., 2015; Pockros et al., 2011), which adjusted according to preliminary studies. In

chemogenetic studies, CNO (1 mg/kg) was dissolved in saline and delivered intraperitoneally 30

min before the behavioral test.

Optogenetic studies

For optogenetic activation, ferrules were connected (by patch cords) to a 473 nm laser diode

through a FC/PC adaptor and a fiber optic rotary joint. The output parameters were: 10 ms, 20

Hz, 8 s on and 2 s off cycle, ～10 mW for terminal stimulation, ～5 mW for somatic stimulation.

For optogenetic inhibition, ferrules were connected to a 593 nm laser diode. The output

parameters were: 10 ms, 20 Hz, constant, ～10 mW for both terminal and somatic inhibition. The

optogenetic stimulation parameters were chosen and slightly modified based on previous

studies (Garcia-Garcia et al., 2018; Walsh et al., 2018; Zhao et al., 2011)

Behavioral assays

Generally, all the behavioral experiments were performed under dim light during the dark phase

of the light-dark cycle. In all tests, all groups of experimental voles were randomly selected and

the observers were blinded to the treatments. During the test, if the subject (not the stimulus

voles) showed very rare or no movements, we think these individuals did not show normal

activity, and their data are not appropriate to include in the following analysis. For optogenetic

and fiber photometry studies, all subjects were allowed to habituate to patch cords for at least

three days and allowed 30 min acclimation to the connection before the experiment started.

Consolation test

The consolation test was performed as previously described (Burkett et al., 2016). Briefly, five

days before the experiment, age-matched adult male and female voles were cohoused together to

promote the formation of a pair bond (Yu et al., 2012). On the testing day, the subjects' partners

were gently transferred in a cup to a sound-proof electric shock chamber and then subjected to

10 rounds of moderately strong foot shocks (3 s, 0.8 mA, 2 min intertrial intervals). At the same

time, the test voles were left undisturbed in their home cages. After the separation, the pairs

were reunited, and the behavior of the subjects was recorded using a video camera for 10 min in

the test room. The digital videos were viewed and quantified using J Watcher software

(http://www.jwatcher.ucla.edu/). According to previous studies (Burkett et al., 2016; Li et al.,

2019), allogrooming is designated as behavioral indicators of consolation, which defined as head

contact with the body or head of their partner, accompanied by a rhythmic head movement. The

duration of chasing (following the shocked partners) and selfgrooming were also collected and

analyzed.

For optogenetic tests, it designed as a between-subjects test in which both groups (mCherry

or opsin) received laser stimulation. The lasting effects of optogenetic inhibition were measured

24 h after the original test, i.e., consolation test was conducted once again without light

stimulation in eNPHR 3.0 animals one day later.

<u>Three-chamber test</u>

The three-chamber test was used to assess the sociability of a subject (Horie et al., 2019; Walsh et

al., 2018). The apparatus consisted of a rectangular box with three separate chambers (20 × 40 ×

20 cm each). One side of each chamber contained a circular metal wire cage (stimulus animal

cage, 11 cm high and 9 cm in diameter). One day before the test, all subjects were habituated to

the arena for 5 min and an age- and sex-matched unfamiliar stimulus voles also habituated to the

wire cages for 5 min. On the testing day, the test vole was placed in the center chamber and a

'stimulus' vole (housed in a same room with the test voles) was randomly placed into one of the

wire cages. After 2 min, the partitioning walls between the chambers were removed and the test

voles were allowed to explore freely for a 10 min session. The time spent exploring each chamber

was automatically recorded using a video tracking system (Shanghai Xinruan, China). Sociability

was calculated as follows: (time spent in the stimulus vole side − time spent in the empty

side)/(time spent in the stimulus vole side + time spent in the empty side).

For optogenetic tests, voles received four epochs of light beginning with a light OFF baseline

epoch (OFF-ON-OFF-ON). Each epoch lasted for 5 min and the total duration was 20 min.

<u>Open-field test</u>

The anxiety level and locomotor function of the subjects were assessed by using the open-field

test (Flanigan et al., 2020; Walsh et al., 2018). Briefly, a square open field (50 cm × 50 cm × 25 cm)

was virtually subdivided into 16 even square. The four central squares were designated as

central area. At the beginning of the test, the test vole was placed into the center area facing

away from the experimenter. Behavior was recorded for 5 min. Outcome measures were distance

traveled, frequency entries and time spent in the central area.

For optogenetic tests, voles were tested twice in the same day in both light-off and light-on

conditions with at least 2 h between sessions (within-subjects design).

Fiber photometry

The fiber photometry recording set-up (ThinkerTech, Nanjing, China) was generated and used as

previously described (Feng et al., 2019; Yuan et al., 2019). Briefly, the emission light was

generated by a 480 LED, reflected with a dichroic mirror and delivered to the brain in order to

excite the GCaMP6m or the 5-HT sensor. The emission light passed through another band pass

filter, into a CMOS detector (Thorlabs, Inc. DCC3240M) and finally recorded by a Labview

program (TDMSViewer, ThinkerTech, Nanjing, China).

On the test day, voles were mildly anesthetized with isoflurane and connected to a

muti-mode optic fiber patch cord (ThinkerTech Nanjing Bioscience®, NA: 0.37, OD: 200 µm)

which the other end connected to fiber photometry apparatus. After 30 min habituation, the

voles were subjected to consolation test as described above (exposure to their shocked partners)

and the behavioral recording consisted of allogrooming, sniffing, social approaching,

self-grooming and running. During the test, if a specific behavior occurred no more than four

times within 1 h, the record was replicated in the following day.

Fiber photometry signals were processed with custom-written MATLAB software. Briefly,

all the data were segmented based on the behavioral events and baseline phase. The change in

fluorescence ($\Delta F/F$) was calculated as $(F–F0)/F0$, where F0 represents the baseline fluorescence

signal averaged over a 10 s-long control time window. We first segmented the data based on the

behavioral events. Then, we calculated the average 5-HT and calcium signals in both the pre-

and post-phases (2/8 s). The time window was determined according to the duration of a

behavior bout. The response elicited during a behavior was calculated as the average $\Delta F/F$

during all trials of a particular behavior. The peak response during a behavior was calculated as

the maximum $\Delta F/F$ during the behavior minus F0.

For specificity verification of 5-HT sensors in Figure 5—figure supplement 2, a 5-HT

receptor antagonist metergoline (Met) was injected (i.p., 4 mg/kg) 20 min before the fiber

photometry experiments. The timing and dosing were chosen according to previous studies

(Onasanwo et al., 2016; Wan, et al 2020).

In vitro **electrophysiological recordings**

To verify the optogenetic and chemogenetic manipulations, we performed *in vitro* whole-cell

patch-clamp recordings from DR neurons. Neurons expressing ChR2, eNpHR3.0, hM3Dq and

hM4Di were visually identified by fluorescence of mCherry. The voles were anesthetized with

isoflurane. Brains were quickly dissected and 300 µm coronal slices containing the DRN were

prepared in a chamber filled with ACSF (32-34℃) using vibratome (Campden 7000 smz). The

recordings were obtained using a Multiclamp 700B amplifier, filtered at 5 kHz and sampled at 10

kHz with Digidata 1440A. Clampex 10.5 was used for analysis.

Current-clamp recordings were performed to measure evoked action potentials. For

photoactivation and photoinhibition, the light protocols used during behavioral tests were

delivered through a 200 µm optical fiber close to recorded neurons. For CNO activation,

spontaneous firing of action potentials in the cell was recorded in current clamp mode at -60 mV.

After 5 min of recording, the slices were perfused with 10 µM CNO. The total recording time for

each cell was 10 min. For CNO inhibition, we applied currents in steps of 25 pA, ranging from 0

pA to 250 pA. Neurons were allowed to recover for 10 min then perfused with 10 µM CNO. The

same current procedure was performed. Afterwards, the CNO was removed by washes with

ASCF and cells were recorded for another 10 min.

Data analysis

All data are represented as the mean ± SE. All data were assessed for normality using a

one-sample Kolmogorov-Smirnov test, and the Levene's test was used to confirm homogeneity

of variance. Comparisons between two groups were performed by unpaired or paired t tests.

Comparisons among three or more groups of different animals were performed

using one-way ANOVA. Two-way ANOVAs or two-way repeated measures ANOVAs were

used to compare multiple groups under multiple testing conditions when appropriate. Post hoc

comparisons were conducted using Tukey. To confirm the significance and calculate the effect

size, Bayesian t test (paired or unpaired) and Bayesian ANOVA were conducted for all the

comparisons using default priors. For more detailed information about the Bayesian test, please

refer to Dr. Keysers's work (Keysers et al., 2020). The data were analyzed using JASP 14.0 and are

presented as the mean ± SE. The significant level was set at $P < 0.05$. The detailed analysis

method in each figure and the statistical quantifications are presented in Supplementary 2.

ACKNOWLEDGMENTS AND DISCLOSURES

This research was supported by the National Natural Science Foundation of China (31970424,

31670421, 31372213 and 31772473), the Natural Science Basic Research Plan in Shaanxin Province

of China (2018JM3032) and the Fundamental Research Funds for Central University

(GK201903059). We thank Yu-Ying Yang, Xin Zhang and Yi-Xin Feng for assistance in

conducting experiments and caring for voles. The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

Prof. F.D.T. designed the study; L.F.L. conducted the majorities of experiments and wrote the

original draft; L.Z.Z participated in the electrophysiological experiment; Z.X.H. and R.J.

discussed the results and provided constructive comments; Y.T.Z participated in the fiber

photometry studies. H.M., Y.F.X, W.Y., W.J.H., Z.J.L. and L.Y.T. participated in the behavior

study and helped to collect and analyze the data. All authors contributed to and have approved

the final manuscript.

APPENDIX

Supplementary 1. Supplemental figures.

Supplementary 2. Analysis methods and the statistical quantifications

REFERENCE

Arakawa, H., 2020. Somatosensorimotor and Odor Modification, Along with Serotonergic

Processes Underlying the Social Deficits in BTBR T+ Itpr3(tf)/J and BALB/cJ Mouse

Models of Autism. Neuroscience 445, 144-162.

Artigas, F., 2013. Serotonin receptors involved in antidepressant effects. Pharmacology &

therapeutics 137, 119-131.

Bozzi, Y., Provenzano, G., Casarosa, S., 2018. Neurobiological bases of autism-epilepsy

comorbidity: a focus on excitation/inhibition imbalance. The European journal of

neuroscience 47, 534-548.

Burkett, J.P., Andari, E., Johnson, Z.V., Curry, D.C., de Waal, F.B., Young, L.J., 2016.

Oxytocin-dependent consolation behavior in rodents. Science (New York, N.Y.) 351,

375-378.

Carhart-Harris, R.L., Nutt, D.J., 2017. Serotonin and brain function: a tale of two receptors.

Journal of psychopharmacology (Oxford, England) 31, 1091-1120.

Carlyle, M., Stevens, T., Fawaz, L., Marsh, B., Kosmider, S., Morgan, C.J., 2019. Greater empathy

in MDMA users. Journal of psychopharmacology (Oxford, England) 33, 295-304.

Celada, P., Puig, M.V., Artigas, F., 2013. Serotonin modulation of cortical neurons and networks.

Frontiers in integrative neuroscience 7, 25.

Charnay, Y., Leger, L., 2010. Brain serotonergic circuitries. Dialogues in clinical neuroscience 12,

471-487.

Cooper, M.A., McIntyre, K.E., Huhman, K.L., 2008. Activation of 5-HT1A autoreceptors in the

dorsal raphe nucleus reduces the behavioral consequences of social defeat.

Psychoneuroendocrinology 33, 1236-1247.

Correia, P.A., Lottem, E., Banerjee, D., Machado, A.S., Carey, M.R., Mainen, Z.F., 2017. Transient

inhibition and long-term facilitation of locomotion by phasic optogenetic activation of

serotonin neurons. eLife 6.

de Waal, F.B., 2008. Putting the altruism back into altruism: the evolution of empathy. Annual

review of psychology 59, 279-300.

de Waal, F.B.M., Preston, S.D., 2017. Mammalian empathy: behavioural manifestations and

neural basis. Nature reviews. Neuroscience 18, 498-509.

Dolder, P.C., Grunblatt, E., Muller, F., Borgwardt, S.J., Liechti, M.E., 2017. A Single Dose of LSD

Does Not Alter Gene Expression of the Serotonin 2A Receptor Gene (HTR2A) or Early

Growth Response Genes (EGR1-3) in Healthy Subjects. Frontiers in pharmacology 8, 423.

Dolen, G., Darvishzadeh, A., Huang, K.W., Malenka, R.C., 2013. Social reward requires

coordinated activity of nucleus accumbens oxytocin and serotonin. Nature 501, 179-184.

Faye, C., Hen, R., Guiard, B.P., Denny, C.A., Gardier, A.M., Mendez-David, I., David, D.J., 2020.

Rapid Anxiolytic Effects of RS67333, a Serotonin Type 4 Receptor Agonist, and Diazepam,

a Benzodiazepine, Are Mediated by Projections From the Prefrontal Cortex to the Dorsal

Raphe Nucleus. Biological psychiatry 87, 514-525.

Feng, J., Zhang, C., Lischinsky, J.E., Jing, M., Zhou, J., Wang, H., Zhang, Y., Dong, A., Wu, Z., Wu,

H., Chen, W., Zhang, P., Zou, J., Hires, S.A., Zhu, J.J., Cui, G., Lin, D., Du, J., Li, Y., 2019.

A Genetically Encoded Fluorescent Sensor for Rapid and Specific In Vivo Detection of

Norepinephrine. Neuron 102, 745-761 e748.

Field, T., Diego, M., Hernandez-Reif, M., 2009. Depressed mothers' infants are less responsive to

faces and voices. Infant behavior & development 32, 239-244.

Flanigan, M.E., Aleyasin, H., Li, L., Burnett, C.J., Chan, K.L., LeClair, K.B., Lucas, E.K.,

Matikainen-Ankney, B., Durand-de Cuttoli, R., Takahashi, A., Menard, C., Pfau, M.L.,

Golden, S.A., Bouchard, S., Calipari, E.S., Nestler, E.J., DiLeone, R.J., Yamanaka, A.,

Huntley, G.W., Clem, R.L., Russo, S.J., 2020. Orexin signaling in GABAergic lateral

habenula neurons modulates aggressive behavior in male mice. Nature neuroscience.

Fu, W., Le Maitre, E., Fabre, V., Bernard, J.F., David Xu, Z.Q., Hokfelt, T., 2010. Chemical

neuroanatomy of the dorsal raphe nucleus and adjacent structures of the mouse brain.

The Journal of comparative neurology 518, 3464-3494.

Fukumoto, K., Fogaça, M.V., Liu, R.J., Duman, C.H., Li, X.Y., Chaki, S., Duman, R.S., 2020.

Medial PFC AMPA receptor and BDNF signaling are required for the rapid and

sustained antidepressant-like effects of 5-HT(1A) receptor stimulation.

Neuropsychopharmacology : official publication of the American College of

Neuropsychopharmacology 45, 1725-1734.

Fukumoto, K., Iijima, M., Chaki, S., 2014. Serotonin-1A receptor stimulation mediates effects of a

metabotropic glutamate 2/3 receptor antagonist,

2S-2-amino-2-(1S,2S-2-carboxycycloprop-1-yl)-3-(xanth-9-yl)propanoic acid (LY341495),

and an N-methyl-D-aspartate receptor antagonist, ketamine, in the novelty-suppressed

feeding test. Psychopharmacology 231, 2291-2298.

Garcia-Garcia, A.L., Canetta, S., Stujenske, J.M., Burghardt, N.S., Ansorge, M.S., Dranovsky, A.,

Leonardo, E.D., 2018. Serotonin inputs to the dorsal BNST modulate anxiety in a 5-HT1A

receptor-dependent manner. Molecular psychiatry 23, 1990-1997.

Gong, P., Liu, J., Blue, P.R., Li, S., Zhou, X., 2015. Serotonin receptor gene (HTR2A) T102C

polymorphism modulates individuals' perspective taking ability and autistic-like traits.

Frontiers in human neuroscience 9, 575.

Gyurak, A., Haase, C.M., Sze, J., Goodkind, M.S., Coppola, G., Lane, J., Miller, B.L., Levenson,

R.W., 2013. The effect of the serotonin transporter polymorphism (5-HTTLPR) on

empathic and self-conscious emotional reactivity. Emotion (Washington, D.C.) 13, 25-35.

Heifets, B.D., Malenka, R.C., 2016. MDMA as a Probe and Treatment for Social Behaviors. Cell

166, 269-272.

Horie, K., Inoue, K., Suzuki, S., Adachi, S., Yada, S., Hirayama, T., Hidema, S., Young, L.J.,

Nishimori, K., 2019. Oxytocin receptor knockout prairie voles generated by

CRISPR/Cas9 editing show reduced preference for social novelty and exaggerated

repetitive behaviors. Hormones and behavior 111, 60-69.

Ishii, H., Ohara, S., Tobler, P.N., Tsutsui, K., Iijima, T., 2015. Dopaminergic and serotonergic

modulation of anterior insular and orbitofrontal cortex function in risky decision making.

Neuroscience research 92, 53-61.

Keysers, C., Gazzola, V., Wagenmakers, E.J., 2020. Using Bayes factor hypothesis testing in

neuroscience to establish evidence of absence. Nature neuroscience 23, 788-799.

Kim, B.S., Lee, J., Bang, M., Seo, B.A., Khalid, A., Jung, M.W., Jeon, D., 2014. Differential

regulation of observational fear and neural oscillations by serotonin and dopamine in the

mouse anterior cingulate cortex. Psychopharmacology 231, 4371-4381.

Knapska, E., Mikosz, M., Werka, T., Maren, S., 2010. Social modulation of learning in rats.

Learning & memory (Cold Spring Harbor, N.Y.) 17, 35-42.

Li, L.F., Yuan, W., He, Z.X., Ma, H., Xun, Y.F., Meng, L.R., Zhu, S.J., Wang, L.M., Zhang, J., Cai,

W.Q., Zhang, X.N., Guo, Q.Q., Lian, Z.M., Jia, R., Tai, F.D., 2020. Reduced Consolation

Behaviors in Physically Stressed Mandarin Voles: Involvement of Oxytocin, Dopamine

D2, and Serotonin 1A Receptors Within the Anterior Cingulate Cortex. The international

journal of neuropsychopharmacology 23, 511-523.

Li, L.F., Yuan, W., He, Z.X., Wang, L.M., Jing, X.Y., Zhang, J., Yang, Y., Guo, Q.Q., Zhang, X.N.,

Cai, W.Q., Hou, W.J., Jia, R., Tai, F.D., 2019. Involvement of oxytocin and GABA in

consolation behavior elicited by socially defeated individuals in mandarin voles.

Psychoneuroendocrinology 103, 14-24.

Li, Y., Zhong, W., Wang, D., Feng, Q., Liu, Z., Zhou, J., Jia, C., Hu, F., Zeng, J., Guo, Q., Fu, L.,

Luo, M., 2016. Serotonin neurons in the dorsal raphe nucleus encode reward signals.

Nature communications 7, 10503.

Luo, M., Zhou, J., Liu, Z., 2015. Reward processing by the dorsal raphe nucleus: 5-HT and

beyond. Learning & memory (Cold Spring Harbor, N.Y.) 22, 452-460.

Matsunaga, M., Ishii, K., Ohtsubo, Y., Noguchi, Y., Ochi, M., Yamasue, H., 2017. Association

between salivary serotonin and the social sharing of happiness. PloS one 12, e0180391.

Meneses, A., Liy-Salmeron, G., 2012. Serotonin and emotion, learning and memory. Reviews in

the neurosciences 23, 543-553.

Ohmura, Y., Tanaka, K.F., Tsunematsu, T., Yamanaka, A., Yoshioka, M., 2014. Optogenetic

activation of serotonergic neurons enhances anxiety-like behaviour in mice. The

international journal of neuropsychopharmacology 17, 1777-1783.

Onasanwo, S.A., Faborode, S.O., Ilenre, K.O., 2016. Antidepressant-like Potentials of Buchholzia

Coriacea Seed Extract: Involvement of Monoaminergic and Cholinergic Systems, and

Neuronal Density in the Hippocampus of Adult Mice. Nigerian journal of physiological

sciences : official publication of the Physiological Society of Nigeria 31, 93-99.

Perez-Manrique, A., Gomila, A., 2018. The comparative study of empathy: sympathetic concern

and empathic perspective-taking in non-human animals. Biological reviews of the

Cambridge Philosophical Society 93, 248-269.

Pockros, L.A., Pentkowski, N.S., Swinford, S.E., Neisewander, J.L., 2011. Blockade of 5-HT2A

receptors in the medial prefrontal cortex attenuates reinstatement of cue-elicited

cocaine-seeking behavior in rats. Psychopharmacology 213, 307-320.

Puig, M.V., Artigas, F., Celada, P., 2005. Modulation of the activity of pyramidal neurons in rat

prefrontal cortex by raphe stimulation in vivo: involvement of serotonin and GABA.

Cerebral cortex (New York, N.Y. : 1991) 15, 1-14.

Puig, M.V., Gulledge, A.T., 2011. Serotonin and prefrontal cortex function: neurons, networks,

and circuits. Molecular neurobiology 44, 449-464.

Santana, N., Artigas, F., 2017. Laminar and Cellular Distribution of Monoamine Receptors in Rat

Medial Prefrontal Cortex. Frontiers in neuroanatomy 11, 87.

Santana, N., Bortolozzi, A., Serrats, J., Mengod, G., Artigas, F., 2004. Expression of serotonin1A

and serotonin2A receptors in pyramidal and GABAergic neurons of the rat prefrontal

cortex. Cerebral cortex (New York, N.Y. : 1991) 14, 1100-1109.

Tian, Z., Yamanaka, M., Bernabucci, M., Zhao, M.G., Zhuo, M., 2017. Characterization of

serotonin-induced inhibition of excitatory synaptic transmission in the anterior cingulate

cortex. Molecular brain 10, 21.

Vattikuti, S., Chow, C.C., 2010. A computational model for cerebral cortical dysfunction in

autism spectrum disorders. Biological psychiatry 67, 672-678.

Walsh, J.J., Christoffel, D.J., Heifets, B.D., Ben-Dor, G.A., Selimbeyoglu, A., Hung, L.W.,

Deisseroth, K., Malenka, R.C., 2018. 5-HT release in nucleus accumbens rescues social

deficits in mouse autism model. Nature 560, 589-594.

Wan JX, Peng WL, Li XL, Qian TG, Song K, Zeng JZ, et al. A genetically encoded GRAB sensor

for measuring serotonin dynamics in vivo. *Biorxiv*. 2020; 38:569-580; posted February 25,

2020; doi: 10.1101/2020.02.24.962282.

Wang, L., Zhu, Z., Hou, W., Zhang, X., He, Z., Yuan, W., Yang, Y., Zhang, S., Jia, R., Tai, F., 2019.

Serotonin Signaling Trough Prelimbic 5-HT1A Receptors Modulates CSDS-Induced

Behavioral Changes in Adult Female Voles. The international journal of

neuropsychopharmacology 22, 208-220.

Weber, E.T., Andrade, R., 2010. Htr2a Gene and 5-HT(2A) Receptor Expression in the Cerebral

Cortex Studied Using Genetically Modified Mice. Frontiers in neuroscience 4.

Yagishita, S., 2020. Transient and sustained effects of dopamine and serotonin signaling in

motivation-related behavior. Psychiatry and clinical neurosciences 74, 91-98.

Yizhar, O., Fenno, L.E., Prigge, M., Schneider, F., Davidson, T.J., O'Shea, D.J., Sohal, V.S., Goshen,

I., Finkelstein, J., Paz, J.T., Stehfest, K., Fudim, R., Ramakrishnan, C., Huguenard, J.R.,

Hegemann, P., Deisseroth, K., 2011. Neocortical excitation/inhibition balance in

information processing and social dysfunction. Nature 477, 171-178.

Young, K.S., Parsons, C.E., Stein, A., Kringelbach, M.L., 2015. Motion and emotion: depression

reduces psychomotor performance and alters affective movements in caregiving

interactions. Frontiers in behavioral neuroscience 9, 26.

Yu, P., An, S., Tai, F., Zhang, X., He, F., Wang, J., An, X., Wu, R., 2012. The effects of neonatal

paternal deprivation on pair bonding, NAcc dopamine receptor mRNA expression and

serum corticosterone in mandarin voles. Hormones and behavior 61, 669-677.

Yuan, Y., Wu, W., Chen, M., Cai, F., Fan, C., Shen, W., Sun, W., Hu, J., 2019. Reward Inhibits

Paraventricular CRH Neurons to Relieve Stress. Current biology : CB 29, 1243-1251

e1244.

Zhao, S., Ting, J.T., Atallah, H.E., Qiu, L., Tan, J., Gloss, B., Augustine, G.J., Deisseroth, K., Luo,

M., Graybiel, A.M., Feng, G., 2011. Cell type-specific channelrhodopsin-2 transgenic mice

for optogenetic dissection of neural circuitry function. Nature methods 8, 745-752.

    SUPPLEMENTARY 1. SUPPLEMENTAL FIGURES.

834   **Figure 1—figure supplement 1. The histology of CTB injecting into the right ACC of male (up**

835   **row) and female voles (down row).** The right panels showed colocalization of TPH2$^+$ neurons

836   (green) and CTB (red) in the DR (× 200). M: male; F: female; ACC: anterior cingulate cortex; DR:

837   dorsal raphe nucleus; TPH2: tryptophan hydroxylase 2; Aq: aqueduct.

Figure 1—figure supplement 2. Immunohistological images showing colocalization of opsins

(mCherry, red), TPH2+ neurons (green) and DAPI (blue) in the DR of males (A) and (B)

females voles. (a1 & b1): merged image of DAPI and mCherry (× 100); (a2-a4, b2-b4): amplified

images in the left box showing the mCherry, TPH2 and the colocaliztion of mCherry and TPH2

(× 200); (a5, b5): quantification rates of mCherry neurons colabled with TPH2 (left pies), and

TPH2 neurons colabled with mCherry (right pies), n = 3 in each sex. DR: dorsal raphe nucleus;

TPH2: tryptophan hydroxylase 2.

Figure 1—figure supplement 3. Schematics depict virus spread (shades) and optic fiber placements (lines) for recording and functional manipulation experiments, related to figures 1, 3, 4 and 5 (A, B C and D, respectively).

Figure 1—figure supplement 4. Effect of bidirectional optogenetic modulation of DR 5-HT

neuron activities in the DR-ACC neural circuit on some control behaviors. Quantification of

chasing time (A & B) and grooming time (C & D) in the consolation test; and the distance

traveled in the open-field test (E & F). Data are presented as mean ± SE, *n* = 7–8 in each group,

**P* < 0.01. For A-D, Independent samples *t* test and Bayesian samples t-test; for E-F, two-way

repeated ANOVA and two-way Bayesian Repeated Measures ANOVA (light as within subject

factors). ACC: anterior cingulate cortex; CTR: control; DR: dorsal raphe nucleus.

Figure 1—figure supplement 5. **Effect of optogenetic inhibition of DR 5-HT neurons in the**

DR-ACC neural circuit does not elicit long-lasting effects (< 24 h) on allogrooming and

chasing behavior in the consolation test. The data are derived from Figure 1H. (A): Time

spent in allogrooming; (B) allogrooming frequency; (C): time spent in chasing and chasing

frequency (D). Data are presented as mean ± SE, n = 8 in each group; Paired sample t-test and

Bayesian Paired samples t-test; **P < 0.01. ACC: anterior cingulate cortex; DR: dorsal raphe

nucleus; CTR: control.

Figure 2—figure supplement 1. Effect of bidirectional optogenetic modulation of ACC 5-HT

terminals in the DR-ACC neural circuit on some control behaviors. Quantification of chasing

time (A & B) and grooming time (C & D) in the consolation test and the distance traveled in the

open-field test (E & F). Data are presented as mean ± SE, *n* = 7–8 in each group, **P < 0.01. For

A-D, Independent samples *t* test and Bayesian samples t-test; for E-F, two-way repeated

ANOVA and two-way Bayesian Repeated Measures ANOVA (light as within subject factors).

ACC: anterior cingulate cortex; DR: dorsal raphe nucleus; CTR: control.

Figure 2—figure supplement 2. **Effect of optogenetic inhibition of ACC 5-HT terminals in**

the DR-ACC neural circuit does not elicit long-lasting effects (< 24 h) on allogrooming and

chasing behaviors in the consolation test. The data are derived from Figure 2D. (A): Time

spent in allogrooming; (B) allogrooming frequency; (C): time spent in chasing; (D): chasing

frequency. Data are presented as mean ± SE, *n* = 7–8 in each group; Paired sample t-test and

Bayesian Paired samples t-test; **P < 0.01. ACC: anterior cingulate cortex; DR: dorsal raphe

nucleus; CTR: control; Allo: allogrooming.

Figure 3—figure supplement 1. Immunohistological image showing colocalization of

DREADD (mCherry, red), TPH2+ neurons (green) and DAPI (blue) in the DR of male (A)

and (B) female voles. (a1 & b1): merged image of DAPI and mCherry (× 100); (a2-a4, b2-b4):

amplified images in the left box showing the mCherry, TPH2 and the colocaliztion of mCherry

and TPH2 (× 200); (a5, b5): quantification rates of mCherry neurons colabled with TPH2 (left

pies), and TPH2 neurons colabled with mCherry (right pies), *n* = 3 in each sex. TPH2:

tryptophan hydroxylase 2.

Figure 3—figure supplement 2. Effect of chemogenetic modulation of DR 5-HT neuron

activities in the DR-ACC neural circuit on some control behaviors. Quantification of chasing

time (A) and grooming time (B) in the consolation test, and distance traveled in the open-field

test (C). Data are presented as mean ± SE, n = 7–8 in each group; two-way repeated measures

ANOVA along with two-way Bayesian repeated measures ANOVA. ACC: anterior cingulate

cortex; DR: dorsal raphe nucleus; M: male; F: female; OFT: open-field test.

Figure 4—figure supplement 1. Representative viral infection images of GCaMp6. (a1):

Immunohistological image showing GCaMP6 expression in the DR (× 100); (a2-a4): amplified

images in the left box showing the GCaMP6 (green), TPH2 (red) and the colocaliztion (yellow)

of the two (× 200); (a5): quantification rates of GCaMP6 neurons colabled with TPH2 (left pie),

and TPH2 neurons colabled with GCaMP6 (right pie), n = 3; (b1): immunohistological image

showing GFP expression in the DR (× 100); (b2-b4): amplified images in the left box showing

the GFP (green), TPH2 (red) and the colocaliztion (yellow) of the two (× 200); (b5):

quantification rates of GFP neurons colabled with TPH2 (left pie), and TPH2 neurons colabled

with GFP (right pie), n = 3.

Figure 4—figure supplement 2. Peak GCaMP6 (A & B) and 5-HT sensor (C & D)

fluorescence values during a few of behaviors in both male and female voles. Data are

presented as mean ± SE, *n* = 6 in each group. Groups not sharing the same letter significantly

differ from each other; one-way ANOVA along with one-way Bayesian ANOVA. Allo:

allogrooming; Sni: sniffing; App: social approaching; Gro: selfgrooming; Run: running.

Figure 4—figure supplement 3. GCaMP6 fluorescent signals align to the end of some

behaviors. (A1, B1, C1): Representative peri-event plot of GCaMP6 fluorescence signals

aligned to the end of allogrooming, social approaching and sniffing (for all peri-event plots, the

red line denotes the mean signals of 4-6 bouts of behaviors, whereas the red shaded region

denotes the SE); (A2, B2, C2): quantification of change in GCaMP6 fluorescence signals. Data

are presented as mean ± SE, *P < 0.05, **P < 0.01, n = 6 in each group; paired samples t test

along with Bayesian paired samples t-Test. ALLO: allogrooming; APP: approaching; SNI:

sniffing.

Figure 5—figure supplement 1. 5-HT sensor fluorescent signals align to the end of some behaviors. (A1, B1, C1): Representative peri-event plot of 5-HT sensor fluorescence signals aligned to the end of allogrooming, social approaching and sniffing (for all peri-event plots, the blue line denotes the mean signals of 4-6 bouts of behaviors, whereas the blue shaded region denotes the SE); (A2, B2, C2): quantification of change in 5-HT sensor fluorescence signals. Data are presented as mean ± SE, *$P < 0.05$, **$P < 0.01$, $n = 6$ in each group; simple Paired t test along with Bayesian Paired Samples t-Test. ALLO: allogrooming; APP: approaching; SNI: sniffing.

Figure 5—figure supplement 2. 5-HT sensor fluoresce changes during allogrooming could

be blocked by Met. (A): Schematic diagrams; (B): combined individual 5-HT sensor

fluorescence traces during allogrooming in response to Met (blue) or Veh (red); (C, D):

quantification of 5-HT sensor fluorescence signals during allogrooming in response to Met or

Veh in male (C) and female subjects. Data are presented as mean ± SE, **$P < 0.01$, $n = 6$ in each

group; two-way Repeated measures ANOVA and two-way Bayesian repeated measures

ANOVA. Met: metergoline, a 5-HT receptor antagonist; Veh: vehicle.

Figure 6—figure supplement 1. Comparisons of 'Vehicle+Saline - Vehicle+CNO' vs

'8-OH-DPAT+Saline - 8-OH-DPAT+CNO' (A & B) and 'Vehicle+Saline - Vehicle+CNO' vs

'MDL 100907+Saline - MDL 100907+CNO' (C & D). Independent samples *t*-test and Bayesian

independent samples *t*-test. *N* = 5-6 in each group, *P < 0.05.

Figure 6—figure supplement 2. Chemogenetic inhibition of DR 5-HT neuron in the

DR-ACC neural circuit along with intra-ACC injection of 5-HT1AR agonist (8-OH-DPAT) or

5-HT2AR antagonist (MDL 100907) had no significant effect on control behaviors of

selfgrooming (A, C) and distance traveled in the open-field test (B, D). Two-way ANOVA

and two-way Bayesian ANOVA. *N* = 5-6 in each group. ACC: anterior cingulate cortex; DR:

dorsal raphe nucleus; M: male; F: female.