Enhanced functionality of low-affinity CD19 CAR T-cells is associated with activation priming and a

polyfunctional cytokine phenotype

Ilaria M. Michelozzi[1], Eduardo Gomez Castaneda[1], Ruben V.C. Pohle[1], Ferran Cardoso Rodriguez[2],

Jahangir Sufi[2], Pau Puigdevall Costa[3], Meera Subramaniyam[1], Si Wei Wu[1], Aleks Guvenel[1], Sara

Ghorashian[4], Martin A. Pule[5], Christopher J. Tape[2], Sergi Castellano[3,6*], Persis J. Amrolia[1,7*] and Alice

Giustacchini[1].

1. Molecular and Cellular Immunology Section, UCL Great Ormond Street Institute of Child Health,
London, UK
11
12 2. Cell Communication Lab, Department of Oncology, University College London Cancer Institute,
13 London, UK
14
15 3. Genetics and Genomic Medicine Programme, Great Ormond Street Institute of Child Health,
16 University College London, London, UK
17
18 4. Department of Bone Marrow Transplant, Great Ormond Street Children's Hospital, London, UK.
19
20 5. Cancer Institute, University College London, London, UK
21
22 6. UCL Genomics, Great Ormond Street Institute of Child Health, University College London, London,
23 UK
24
25 7. Departments of Bone Marrow Transplant and Haematology, Great Ormond Street Hospital for
26 Children, London, UK
27

*These authors (S.C. and P.J.A.) contributed equally to this work

Corresponding author: Alice Giustacchini (a.giustacchini@ucl.ac.uk)

Abstract

We recently described a low-affinity second-generation CD19 chimeric antigen receptor (CAR, CAT)

that showed enhanced expansion, cytotoxicity, and anti-tumour efficacy compared to the high-affinity

(FMC63 based) CAR used in tisagenlecleucel, in pre-clinical models. Furthermore, CAT demonstrated

an excellent toxicity profile, enhanced *in vivo* expansion, and long-term persistence in a Phase I clinical

study. To understand the molecular mechanisms behind the improved properties of CAT CAR T-cells,

we performed a systematic *in vitro* characterization of the transcriptomic (RNA-seq) and protein (CyTOF) changes occurring in T-cells expressing low-affinity *vs* high-affinity CD19 CARs following stimulation with CD19-expressing cells. We demonstrate that CAT CAR T-cells show enhanced activation to CD19 stimulation and a distinct transcriptomic/protein profile with increased cytokine polyfunctionality post-stimulation compared with FMC63 CAR T-cells. Our results suggest that the enhanced functionality of low-affinity CAR T-cells may be sustained by the establishment of a self-reinforcing circuit activated through cytokines polyfunctional crosstalk.

Research Article

Introduction

T-cells genetically engineered to express CD19 chimeric antigen receptors (CAR T-cells) have shown remarkable efficacy in relapsed/refractory (r/r) B-cell malignancies leading to clinical licensing for r/r B-cell acute lymphoblastic leukaemia (B-ALL) and Non-Hodgkin Lymphoma.[1] Despite this success, several safety and efficacy hurdles remain[2]. CAR T-cells can trigger potent immune responses leading to transient but potentially life-threatening inflammatory events, such as cytokine release syndrome (CRS) and neurotoxicity[3,4]. CAR T-cell engraftment and expansion is a prerequisite for efficacy. Not all CD19 CAR T-cell products persist *in vivo* and long-term immune surveillance appears critical for preventing disease relapse, particularly in ALL. Thus, the design of versatile CAR T-cells, capable of balancing safety and efficacy, is contingent on our understanding of the molecular mechanisms underlying CAR T-cell function. The engagement of CARs to their cognate antigens results in the activation of CAR T-cells and promotes their rapid expansion as well as their differentiation into distinct T-cell subsets, mediating tumour cytolysis (effector cells) and providing long-lasting protection (memory cells). As tumour cell recognition by CAR T-cells relies on the binding of the CAR's single-chain variable fragment (scFv) to its epitope, fine-tuning the affinity of CARs to their antigens has become a strategy to modulate the strength of CAR T-cell responses[5].

The affinity of CARs is determined by their binding kinetics and the rates at which they associate to- and dissociate from their targets. The optimal affinity of a CAR is likely to vary depending on a number of factors, including the CAR design, the CAR expression levels and the antigen density on the target cells[6,7]. Chimeric immunoreceptors have an activation ceiling above which increasing the binding affinity does not improve T-cell activation but can rather result in T-cell exhaustion[8]. In contrast, by reducing CAR scFv affinity, the strength of the T-cell signal can be modulated, so that CAR T-cells discriminate different levels of antigen expression. CARs exhibiting slower antigen-association rates

to ErbB2, EGFR and CD123 targets showed reduced activation in response to low antigen

concentrations, favouring differential targeting of tumour cells overexpressing the target *vs* normal

tissue expressing the same target at physiological levels[8-10]. We have recently described a novel low-

affinity CD19 CAR (CAT), with epitope, structure and stability similar to the widely used FMC63, but

characterized by faster rates of antigen dissociation, leading to an overall 40-fold reduction of its

affinity[6,11,12]. Pre-clinical testing of CAT CAR T-cells has revealed greater antigen-specific cytotoxicity,

higher proliferation and more potent *in vivo* anti-tumour activity when compared to FMC63. In a Phase

I clinical trial in patients with high-risk treatment-refractory paediatric B-ALL, CAT CAR T-cells resulted

in lower toxicity in terms of severe CRS and displayed greater expansion than that reported for the

FMC63 based tisagenlecleucel as well as excellent persistence[12]. In this regard, while CAT CAR T-cells

have been designed to prevent excessive activation and reduce CRS and neurotoxicity risks, in line

with other recently developed low-affinity CD19 CARs[13], enhancements in their cytotoxicity,

expansion and persistence were unexpected[7] and the underlying molecular mechanisms to these

functional phenotypes remain unknown.

The interaction between T-cell receptor (TCR) and the peptide-MHC complex offers however some

insight on how immunoreceptors' affinity can dramatically influence T-cell functions[14]. Faster target

off-rates in TCRs allow a single peptide-MHC complex to serially trigger several TCRs, resulting in

amplified and sustained T-cell activation[14]. Similar to TCRs, low-affinity CARs may facilitate serial killing

of target tumour cells, leading to enhanced activation and decreased exhaustion. Importantly,

reduction of TCR avidity has been shown to also affect T-cell fate, with low-avidity TCRs promoting the

expansion of self-renewing memory T-cell subsets[15]. Little is known about whether fine-tuning of CARs

affinity influences T-cell fate in a similar fashion to TCRs, however, memory T-cell subsets are crucial

for the establishment of long-term protection upon adoptive T-cell transfer, and their expansion in

CAR T-cell products would be highly attractive[16,17]. Herein we perform a systematic *in vitro*

characterization of the molecular and biochemical changes occurring in CAR T-cells, in response to a

low-affinity CD19 CAR (CAT) as compared to a high-affinity one (FMC63). By combining bulk RNA

sequencing with single-cell mass cytometry analyses (cytometry by time of flight, CyTOF), we show

that the expression of the CAT CAR induces a memory-like activation priming in response to low

concentrations of CD19-expressing B-cells found in the manufacture product. Upon antigen

stimulation, we identified distinct molecular features downstream of CAT CAR activation responsible

for enhancing CAT CAR T-cell functional response.

Results

Generation and quality assessments of low- (CAT) and high- (FMC63) affinity CD19 CARs from healthy donors

To dissect the molecular mechanisms behind the functional differences observed between low- (CAT) and high- (FMC63) affinity CD19 CAR T-cells[12], we interrogated transcript and protein expression readouts from T-cells lentivirally (LV) transduced with CD19 CARs differing only in their scFv (Supplementary Fig. 1a), before and after antigenic stimulation. In particular, we performed bulk transcriptomic analyses (RNA-seq) to identify CAR T-cell distinct gene expression signatures and mass cytometry analyses (CyTOF) to model at a single-cell resolution, differences in their downstream signalling. RNA-seq and CyTOF readouts from untransduced (UNTR) controls and T-cells LV transduced to express CAT or FMC63 CD19 CARs from healthy donors (HD1-HD8, Supplementary Table 1) were compared at baseline and following stimulation with CD19+ ALL cell line NALM6 (unstimulated and stimulated conditions, respectively), as schematized in Fig. 1a. Importantly, we ruled out significant differences in the transduction of the 2 CARs by fluorescence activated cell sorting (FACS), assessing the percentage of mCherry+ T-cells (LV fluorescent marker) across donors and experimental conditions. These ranged between 11.50 - 93.80% (median 48.55%) in FMC63 and 13.40 - 93.60% (median 43.60%) in CAT (Fig. 1b, top) and were thus comparable among individual HDs. Similarly, we excluded major differences in transgene expression between the 2 CARs by measuring mCherry mean fluorescent intensity (MFI), a proxy for CARs expression, in transduced T-cells (mCherry+) (Fig. 1b, top). These ranged between 8,277 – 21,578 (median 11,189) and 8,187 – 28,451 (median 10,843) in FMC63 and CAT, respectively, (Fig. 1b, top) and were comparable among individual donors. We also confirmed that both levels of LV transduction and transgene expression intensities remained comparable between constructs upon antigen stimulation (Fig. 1b, bottom). Finally, because the CD4/CD8 composition in CAR T-cell products can affect their anti-tumour activity *in vivo*[18], we compared the CD4:CD8 ratios in CAT and FMC63 CAR T-cells. While no difference was observed in the absence of antigen stimulation, upon NALM6 co-culture we found a slight but statistically significant reduction of CD4:CD8 ratio in CAT as compared to FMC63 (Fig. 1c). This variation was largely attributable to a relative decrease of the CD8+ fraction (and increase of CD4+) in FMC63 CAR T-cells after antigen stimulation (Fig. 1c and Supplementary Fig. 1b) as indicated by the similar frequencies observed in CAT and UNTR T-cells (Fig. 1c). These assessments confirmed that our CAR T-cell products were suitable to investigate the molecular features of CAT and FMC63 CAR T-cells.

Low-affinity CD19 CAT CARs have higher activation priming in absence of antigen stimulation

We next performed bulk RNA-seq of FACS-sorted UNTR T-cells (CD3+) and FMC63 or CAT CAR T-cells (CD3+mCherry+) expanded for 4 days with or without stimulation with NALM6 at 2 time points (T1, 3h and T2, 24 h). Transcriptomic results confirmed that mCherry was an accurate proxy for CAR transgene expression levels, as shown by the significant positive correlation between the MFI of mCherry by FACS and the normalized RNA-seq counts aligning to the scFv region of each of the 2 CARs (Fig. 2a) or to the exogenous sequence of mCherry (Supplementary Fig. 2a). Similarly, the proportion of CD4 and CD8 T-cells detected by FACS was in line with the *CD4* and *CD8* mRNA levels (Supplementary Fig. 2b and Fig. 2b). Principal component analysis (PCA) on the 500 most variable expressed genes (top 100 genes shown in Supplementary Fig. 2c), distributed samples according to a T-cell activation gradient (PC1, from UNTR to CAR activated samples), with the majority of variance in gene expression across experimental conditions being explained by CD19-mediated CAR activation. As expected, UNTR T-cells not expressing any CARs were mostly unaffected by antigen stimulation (Fig. 2c). Similarly, PCA on protein expression from CyTOF, based on earth mover's distance (EMD) scores (a sensitive measure of multivariate changes in protein levels)[19], and in the same experimental conditions and timepoints than the gene expression, followed a similar gradient of sample activation, shifting from UNTR samples to antigen stimulated CAR T-cells on PC1 (Fig. 2d and Supplementary Fig. 3a). Indeed, the 20 most variable genes upregulated upon stimulation with NALM6 in both FMC63 and CAT CAR T-cells include genes involved in T-cell activation (*IL2RA*, *GZMB*) and proliferation (*PCNA*, *LDHA*), which are expressed at very low levels in control T-cells (Fig. 2e and Supplementary Fig. 2c). Consistent with this, the highest protein expression variation is from markers of T-cell activation (CD25, NFAT1, HLA-DR) and proliferation (pRB) (Supplementary Fig. 3b). Of note in both RNA and protein analyses unstimulated CAR T-cells show an intermediate RNA/protein expression profile between UNTR and stimulated CAR T-cells. This suggests that CAR expression on its own in absence of antigen stimulation induces basal T-cell activation (Fig. 2c-e). Importantly, unstimulated CAT samples appear closer to antigen stimulated CAR T-cells than unstimulated FMC63, as well as having less overlap with UNTR samples.

To gain further insights into the intermediate activation state observed in unstimulated CAT, we fuzzy clustered genes by their gene expression and classified individual samples into more than 1 cluster to resolve intermediate cell states and trajectories (Fig. 2f and Supplementary Fig. 2d)[20]. We identified 2 clusters, highly enriched for either inactive T-cells (cluster 1, which includes UNTR samples) or antigen-activated CAR T-cells (cluster 2, which includes stimulated CAR T samples) (Fig. 2f). Notably, the

probability of unstimulated CAT samples of belonging to cluster 2 (activated CAR T-cells) was
substantially higher than that of unstimulated FMC63 (4/6 HDs in CAT *vs* 0/6 HDs inFMC63), further
evidencing the functional proximity between unstimulated CAT and activated CAR T-cells. In this
regard, using gene set enrichment analyses (GSEA) on the Hallmark collection, we confirmed that
while antigen stimulated CAT and FMC63 CAR T-cells have similar enrichment for most of the gene
sets involved in immune functions and cell proliferation, in the absence of antigen stimulation CAT are
uniquely enriched for T-cell activation pathways (Myc Targets V1, Hedghog signalling, Inflammatory
response, Interferon α response) (Fig. 2g). Altogether, these results support that unstimulated CAT
CAR T-cells have stronger T-cell activation priming prior to antigenic stimulation.

Low-affinity CD19 CAT CAR activation priming is associated with and may be driven by residual
CD19-expressing B-cells

To investigate whether the introduction of high- or low-affinity CD19 CARs can differentially affect
gene expression patterns in T-cells, we directly compared the transcriptome of FMC63 and CAT CAR
T-cells, in absence of antigen stimulation. Following differential gene expression (DGE) analysis, we
found that only 10 genes were significantly DE between these two conditions (FDR < 0.1,
Supplementary Table 2), 9 of which were upregulated in CAT *vs* FMC63 CAR T-cells. Among those, we
found genes involved in cytotoxicity (*GNLY*, *GZMK*) and markers of T-cell activation such as MHC class
II molecules (MHCII) (*HLA-DRA* and *HLA-DPA*) (Fig. 3a)*.* This supports our previous observation of gene
expression in unstimulated CAT resembling more that of antigen activated CAR T-cells than gene
expression in unstimulated FMC63 (Fig. 2c). Mass cytometry analyses using a panel of antibodies
against markers of T-cell activation (Supplementary Table 3), confirmed that while unstimulated
FMC63 and CAT CAR T-cells showed similar EMD scores for the majority of the proteins investigated
(Supplementary Fig. 4a), unstimulated CAT CAR T-cells have significantly stronger activation priming
with higher expression of Granzyme B and CD25 ($p < 0.05$) (Fig. 3b). Antigen-independent CAR
activation, also known as tonic signalling, has been associated with CAR T-cell accelerated
differentiation and exhaustion, resulting in impaired anti-tumour efficacy *in vivo*[21-23]. To determine if
the activation priming observed in unstimulated CAT CAR T-cells was a consequence of CAR signalling
due to residual CD19 expressing cells during manufacture or tonic antigen-independent signalling, we
tested whether residual CD19+ B-lymphocytes were detectable in the CAR T-cell product and a
potential source of antigen specific activation. We monitored the proportion of CD19+ B-lymphocytes
in culture at different timepoints. While B-cells were detectable at day 0 in all samples (5.97% of cells
on average), at day 8 (5 day stimulation with CD3/CD28 beads + 3 day rest) they could only be detected
in the UNTR condition (1.43% of cells on average), and had been completely depleted from both CAR

constructs, as shown by FlowSOM analysis of mass cytometry data, clustering single-cells by cell types, and the relative frequencies (Fig. 3c and Supplementary Fig. 4b)[24].

As both CARs were equally capable of clearing CD19-expressing B-cells found at low-ratio in the manufacture product, we investigated whether the stronger activation in low- *vs* high-affinity CARs at both RNA and protein levels resulted from distinct T-cell differentiation responses of these CAR T-cells to CD19-priming. For instance, low receptor affinity and avidity in TCRs have been suggested to preferentially promote differentiation into memory subsets and to favour T-cell persistence *in vivo*[15,25]. Interestingly, CAT CAR T-cells exhibited a substantial increase in the fraction of CD8 central memory T-cells (T_{CM}, CD62+CD45RA-) by FACS as compared to both UNTR and FMC63 conditions (Fig. 3d top). In addition, when using the gene signature-based method xCell to computationally infere immune cell type composition from our gene expression profiles, we were able to confirm the enrichment in CD8 T_{CM} in unstimulated CAT CAR T-cells compared to FMC63 (Fig. 3d bottom)[26]. This suggests that CAR T-cells activation in response to residual B-lymphocytes present during the manufacture may preferentially induce a T_{CM} phenotype in low-affinity CAT CAR T-cells.

CD19 stimulation of low-affinity CAT CAR T-cells results in a distinct transcriptomic and protein profile than high-affinity FMC63 CAR T-cells with enhanced activation/proliferation

We next assessed the impact of "activation priming" and the increased proportion of T_{CM} on the molecular response to antigen stimulation of unstimulated CAT CAR T-cells. Importantly, the stronger basal activation observed in these cells did not prevent an even stronger response when exposed to CD19-expressing NALM6, as shown by a greater increase than FMC63 in the expression of activation, proliferation and cytotoxic markers, relatively to the unstimulated constructs (Supplementary Fig. 5 and 6). In agreement, after 24 h stimulation with NALM6, we identified 51 DE genes, 35 of which were upregulated in CAT compared to FMC63 (Fig. 4a and Supplementary Table 4). CD19 stimulation in CAT CAR T-cells also led to significantly augmented expression of immune stimulatory/proliferation cytokines (*IFNG, CSF2, CXCL8*), and IFN-γ responsive genes (*CIITA*) (Fig. 4a and Supplementary Fig. 7a). Conversely, CAT CAR T-cells displayed significantly decreased expression of the genes encoding for CRIF1 (*GADD45GIP1*), an inhibitor of cell cycle progression, and for FOXP3, the Treg-associated transcription factor known to be only transiently expressed in the initial stages of Th1 response and rapidly downregulated afterwards[27,28] (Fig. 4a). These results suggest that following antigenic stimulation CAT CAR T-cells show a different transcriptomic profile resulting in stronger activation and proliferation than stimulated FMC63. These observations were further substantiated at the protein level. Indeed, among the markers investigated (Supplementary Fig. 7b), CAT CAR T-cells exhibited a

marked increase in the expression (as measured by EMD score) of CD25 and NFAT1, indicators of T-
cell activation.

In addition to a significant increment of pRB and CD8 in CAT CAR T-cells, indicating an increased
fraction of proliferating CD8+ CAR T-cells, we noted enhanced expression of the IL7 receptor (CD127),
commonly associated to memory T-cell phenotypes[29,30] (Fig. 4b). Moreover, the augmented gene
expression of the MHCII trans-activator CIITA observed in CAT, resulted in a corresponding trend
toward the increase of HLA-DR protein[31,32] (Fig. 4b). Interestingly, in addition to being known as late
markers of T-cell activation, MHCII molecules have also been suggested to play a role in T-cell-T-cell
contact, potentially leading to the reinforcement of the immune response[33]. Additionally, CAT CAR T-
cells showed increased expression of the *TNFSF4* gene (Fig. 4a), encoding for the ligand of the T-cell
co-stimulatory receptor OX40 (OX40L). While OX40L is mainly expressed by antigen presenting cells
to promote T-cell activation during antigenic stimulation, it is also expressed in activating T-cells,
where leads to a homotypic OX40L-OX40 signalling axis promoting T-cell longevity and memory
differentiation[34]. Further, we also noted the increased expression of the inflammatory chemokine
genes *CCL4* and *CCL3L1* and genes involved in cell migration (*FLT1*, *DOCK5*) and focal adhesion
(*COL6A3*) in stimulated CAT CAR T-cells (Supplementary Fig. 7a). Importantly, the secretion of CCL4
and CCL3 by activated T-cells has been shown to mediate T-cell attraction toward one another,
promoting T-cell clusters formation and synaptic cytokines exchange coordinating immune
responses[35-38]. These results suggest that homotypic interactions between CAT CAR T-cells may play a
role in the enhanced activation/proliferation observed following CD19 antigenic stimulation
compared with FMC63 CAR T-cells.

The enhanced functionality of low-affinity CD19 CAR T-cells is associated to a distinct cytokine
polyfunctional phenotype upon antigen stimulation

To identify the mechanism underlying the stronger activation of CAT CAR T-cells, we next analysed
their cytokine production at the single-cell level by mass cytometry. We focused our analysis on T1 (3
h stimulation) as it recapitulates the median expression peak of cytokines in stimulated T-cells[39]. The
overall protein intensities measured by EMD scores (Supplementary Fig. 8a) indicated an increased
expression of Granzyme B (also observed in unstimulated CAT *vs* FMC63, Fig. 3b) and a trend toward
the upregulation of additional effector cytokines (IFN-γ, TNF-α) and of immune stimulatory molecules
(GM-CSF and IL-2). No upregulation was observed for Th2/immune-modulatory cytokines (IL-4, IL-5
and TGF-ß) and for Perforin B (Supplementary Fig. 8b). Because most individual differences in cytokine

levels were not statistically significant, we investigated the overall pattern of cytokine co-expression in CAT and FMC63 CAR T-cell responses. The ability of a single T-cell to express simultaneously more than 1 cytokine (polyfunctionality) has been linked to productive immune responses[40,41] and more recently described as a distinctive feature of CAR T-cells correlating with their improved anti-tumour efficacy[42,43]. We measured the frequency at which the 12 cytokines included in our analysis were co-expressed in single CAR T-cells, thus providing a comprehensive profile of their cytokine polyfunctionality. Upon stimulation with NALM6, 30.10% of FMC63 and 45.26% of CAT CAR T-cells were polyfunctional. Furthermore, not only the frequency of polyfunctional CAR T-cells, but also the number of cytokines co-expressed was higher in CAT than in FMC63, with a marked increment in the proportion of cells expressing combinations of 3 or more cytokines (7.33% in FMC63 *vs* 17.50% in CAT) (Fig. 5a). Importantly, a robust increase in CAT CAR T-cell cytokine polyfunctionality was only observed upon CD19 stimulation, in support of the antigenic specificity of the response (Fig. 5a, b and Supplementary Fig. 9a, b). The polyfunctional profiles of stimulated CAT CAR products were dominated by combinations of cytokines involving the effector molecules TNF-α, IL-2 and Granzyme B (Fig. 5b). Among all the possible combinations analysed, the cytokine co-expression patterns significantly enriched in CAT *vs* FMC63 involved effector (IFN-γ, TNF-α, Granzyme B), stimulatory (IL-2, GM-CSF) and T-cell memory-associated (IL-23)[44] cytokines (Fig. 5c).

p38 MAPK signalling axis is preferentially activated in response to low-affinity CD19 CAT CARs

To dissect the signalling axes resulting from for FMC63 and CAT differential cytokine profiles, we analysed the intracellular signalling of CAR T-cells using mass cytometry at 24 h post-stimulation. In particular, we measured the activation of mammalian Target of Rapamycin Complex 1 (mTORC1) dependent cell proliferation and protein translation signals, effectors of the TCR/CAR CD3ζ chain (pZAP70, pERK, pp38), and their downstream transcription factors (NFAT1, pCREB, FOXP3). To ensure that we only captured CAR dependent signals, CD3/CD28 beads were removed from the culture 48 h before NALM6 stimulation. Upon antigen stimulation, CAT significantly upregulated the mTORC1 downstream effectors pS6 and pRB ($p < 0.01$) (Fig. 6a), consistent with the increased proliferation observed in pre-clinical studies[12]. In addition, CAT CAR T-cells showed enhanced phosphorylation of ZAP70 (pZAP70) ($p < 0.05$) (Fig. 6a), a proximal effector of the CD3ζ chain (present in the CAR constructs Supplementary Fig. 1a) whose phosphorylation initiates the TCR intracellular signalling. Downstream of the pZAP70 cascade, the phosphorylated form of p38 mitogen activated protein kinase (MAPK) was upregulated in CAT CAR T-cells compared with FMC63 CAR T-cells ($p < 0.05$), while no difference was observed in the induction of another pZAP70 effector, pERK (Fig. 6a). Among the

transcription factors analysed, CAT CAR T-cells exhibited higher levels of NFAT1 and FOXP3

(Supplementary Fig. 10a). Of note, FOXP3 transient expression in T effectors (as demonstrated by its

concurrent transcript decrease, Fig. 4a) correlates with sustained levels of CD127, proliferation and

pro-inflammatory cytokine secretion[27].

To model CAT CAR downstream signalling network, we performed a density resampled estimate of

mutual information (DREMI) analysis[45] on our mass cytometry results. While EMD scores measure

intensity in individual proteins, DREMI scores quantify the intensity of interaction (direct or indirect)

between pairs of proteins. We focused on signalling differences between CAT and FMC63 and built

interaction networks centred on those differing molecules and connections. Up (red) and down (blue)

regulation of individual proteins, relative to the average protein expression in each condition, was

represented by EMD scores in the network nodes (circles in Fig. 6b, Supplementary Fig. 10b), while

interaction between two proteins was represented by DREMI scores (Supplementary Fig. 10c) in the

network edges (arrows in Fig. 6b, Supplementary Fig. 10b). While many of the signalling modules

identified showed similar activation, as expected from CD19 CARs only differing for their antigen

affinity (Supplementary Fig. 8a right), our analysis identified a signalling axis around p38 MAPK, which

was selectively upregulated in CAT upon antigen stimulation (Fig. 6b and Supplementary Fig. 10c). Of

note, p38 MAPK can be activated both downstream of CD3ζ chain stimulation (through pZAP70,

antigen-dependent activation) or in response to cytokines signalling (antigen-independent activation).

It is however only essential for cytokine-mediated activation in T-cells and thus of relevance to this

activation route in CAT CAR T-cells.[46,47] Further, p38 MAPK pathway regulates the stability and

translation of several anti-tumoural cytokine transcripts, including IL-2, IFN-γ, TNF-α, GM-CSF,

CXCL8,[48,49] some of which were upregulated at the mRNA and/or protein levels in CAT expressing CAR

T-cells (Fig. 4, Supplementary Fig. 8b). Lastly, the activation of p38 induces NFAT expression,

promoting its interaction with the coactivator CREB-binding protein and increased transcription of T-

cell activation genes, including cytokines. These results, combined with the marked polyfunctional

phenotype observed in CAT (Fig. 5), point to a self-reinforcing circuit of cytokine mediated activation

in CAT CAR T-cells, as depicted in Fig. 6c.

Low-affinity CAT CAR T-cells display increased trogocytosis

To determine the role of different T-cell subsets in the induction of cytokine polyfunctionality in CAT

CAR T-cells, we correlated the expression of cytokines and T-cell phenotypic surface markers, using

mass cytometry DREMI scores. Interestingly, among all the surface markers tested (Supplementary

Fig. 11a), CD19 (the CARs cognate antigen) established the greatest number of significant correlations

(with TNF-α, GM-CSF, IL-2 and TGF-β) in CAT CAR T-cells (Fig. 7a). Indeed FlowSOM analysis revealed

that CD19 expression was consistently detected in both CD4+ and CD8+ CAR T-cells (Fig. 7b). We

reasoned that if CD19 signal was originating from cell doublets (CAR T-NALM6), we should detect its

expression at both transcript and protein level. Interestingly, RNA-seq performed at 24 h after

stimulation had negligible expression of *CD19* and *CD22* (another B-cell marker), in contrast to *CD3*

mRNAs, arguing against the CD3+CD19+ signal originating from cell doublets (Supplementary Fig. 11b).

This observation was further substantiated by the absence of increase in DNA staining intensity

(marker of doublets) in the CD19+ CAR T-clusters by mass cytometry (Supplementary Fig. 11c). CD19

CAR T-cells have been recently reported to transiently expose CD19 cognate antigen on their surface

membrane, following target cell engagement. This process, trogocytosis, involves the transfer of

membrane fragments from antigen expressing cells to lymphocytes. As a consequence of CAR T-cells

becoming targets themselves, trogocytosis has been linked to fratricide killing in FMC63 CAR T-cells[50].

Our results highlighted a transient kinetic of the *bona fide* trogocytosis observed, peaking at 3 h after

NALM6 stimulation in both CAT and FMC63 (Fig. 7b, c), and decreasing after 24 h, while persisting only

in CAT CAR T-cells for longer (Fig. 7b, c and Supplementary Fig. 12a). The increased levels of

trogocytosis in CAT CAR T-cells suggests that engagements between CAT CAR T-cells and their targets

may be unusually frequent, consistent with the serial triggering pattern observed in low-affinity

immune receptors[14,51]. The longer persistence of CAT trog+ CAR T-cells in culture may result from

reduced fratricide since tumour cells (NALM6) expressed significantly higher levels of CD19 protein

than trog+ CAT CAR T-cells (Fig. 7d). As a result, trog+ CAR T-cells spared from fratricide killing may

serve as an additional source of CD19-dependent CAR T stimulation after NALM6 cells have been

cleared. To corroborate this hypothesis, we analysed trog+ CAR T-cell functional phenotypes by mass

cytometry at different timepoints after stimulation (Supplementary Fig. 12b). At 3h post-stimulation,

when CAT and FMC63 have similar frequencies of trog+ CAR T-cells, trog+ CAT CAR T-cells showed

higher markers of activation and cytokine polyfunctionality (Supplementary Fig. 12c, d). Strikingly, at

24 h after stimulation (when we observe no residual NALM6 in the co-culture, Fig. 7e), we noticed

that trog+ CAT CAR T-cells preserved a robust activation phenotype, with enhanced expression of

proliferation markers (pRB, CD3), T-cell activation factors (CD25, NFAT1, FOXP3), lytic cytokines

(Granzyme B, Perforin B) and p38 MAPK signalling (Fig. 7f and Supplementary Fig. 12e). Furthermore,

trog+ CAT CAR T-cells at 24 h maintained the ability to co-express combinations of effector (IFN-γ,

Granzyme B) and stimulatory (IL-2) cytokines. In contrast, trog+ FMC63 shifted to higher expression

of TGF-β (alone or in combination with other cytokines), which is commonly associated to the

abrogation of T-cell proliferation and cytotoxicity (Fig. 7g and Supplementary Fig. 12e). Altogether

these results point to a self-reinforcing role of trogocytosis in low-affinity CD19 CAT CAR T-cells.

Discussion

Modulating CAR T-cell affinity may enable us to enhance anti-tumour response and long-term tumour

surveillance, while minimizing CAR T-cell related toxicity. To this aim, understanding the molecular

determinants of treatment efficacy in low-affinity CARs is crucial to inform their design and optimize

their manufacture. We thus investigated the molecular features of low-affinity CD19 CAT CAR T-cells,

compared them with those of a high-affinity FMC63-based one, to begin unravelling the molecular

mechanism behind CAT clinical benefits[12]. In this regard, low-affinity CAT CARs induce stronger anti-

tumour responses than high-affinity ones, an apparent paradox, which could be explained by the

faster target off-rate of low-affinity CARs. Faster dissociation requires fewer targets to serially trigger

a larger number of CARs and amplify anti-tumoural response[14]. This would align with the proposed

model of temporal and spatial summation of T-cell activation, in which signals from serially triggered

immunoreceptors can be accumulated and integrated overtime to reach the threshold required for T-

cell activation[51]. Indeed, we found that residual B-cells expressing CD19 in the CAR T-cell product

prime both CAT and FMC63 CARs, with CAT however having a more pronounced expansion of CD8

T_{CM}. This is consistent with the hypothesis that CAT serial triggering may amplify T-cell activation from

lower antigen levels[14] and points to a boosting role of CD19 antigen priming during CAR T-cell

manufacture.

Our transcriptomic and protein profiling revealed that unstimulated CAT CAR T-cells are functionally

closer to antigen activated CAR T-cells than FMC63 CAR T-cells. It is not clear whether this "activation

priming" results from stimulation by residual CD19+ cells during manufacture or antigen-independent

CAR tonic signalling to their molecular phenotype. Mass cytometry analyses of unstimulated CAR T-

cells did not reveal activation of CAR-downstream signalling in CAT nor in FMC63, arguing against CAR-

dependent tonic signalling being responsible for CAT activation priming. On the other hand, the

transcriptomic profile of unstimulated CAT CAR T-cells, identified a number of upregulated activation

genes (*HLA-DBP1*, *HLA-DRA* etc). Despite these genes being commonly associated to T-cell activation,

their basal expression is also increased in memory T-cells as compared to $T_{NAÏVE}$ cells. This is because

memory T-cells are characterized by an open chromatin conformation favouring the access of

transcription factors to immune response genes, thus ensuring a pool of readily available mRNAs that

can be rapidly translated following stimulation[52]. When stimulated with CD19-expressing NALM6 cell

line, CAT CAR T-cells showed a distinct transcriptomic and protein response to CD19 antigenic
stimulation than FMC63 CAR T-cells with increased expression of proliferation, activation and
cytotoxic markers at both RNA and protein levels. *In vivo* clonal kinetics analyses have shown that
single CD19 CAR T-cells that have higher expression of cytotoxic related genes in the manufacture
product, many of which in common with ours (including *IFNG*, *HLA-DRA*, *CCL4*), gave rise to superior
in vivo expansion and survival, significantly contributing to later timepoints after adoptive transfer in
patients[53]. While we did not observe major differences in single cytokines, upon stimulation CAT CAR
T-cells were uniquely characterized by a polyfunctional pattern of cytokine expression, with a marked
increase in the frequency of single CAR T-cells expressing >=3 cytokines when compared to FMC63.
CAT polyfunctional profile was dominated by combinations of effector cytokines, consistent with their
potent anti-tumour activity. It has been recently shown that the ability of CAR T-cells to produce
multiple cytokines (polyfunctionality) in response to antigen exposure may be associated with
improved anti-tumour responses *in vivo*[42].

Analyses of intracellular signalling by mass cytometry revealed the increased phosphorylation of p38
MAPK in stimulated CAT as compared to FMC63. P38 MAPK is central to promoting mRNA stability of
proinflammatory genes which, intriguingly, may contribute to increased cytokine co-expression in
CAT. Furthermore, the activation of p38 can induce NFAT and pCREB-dependent transcription of T-
cell activation genes, including cytokines[48]. Analysis of gene expression also revealed the upregulation
of chemoattractive cytokines CCL4 and CCL3L1 in CAT, which induce T-cell homotypic interactions and
promote the reciprocal exchange of self-reinforcing signals such as OX40L,[34,37] which are upregulated
in CAT as well. These data suggest enhanced p38 MAPK phosphorylation in CAT CAR T-cells may play
a key role in establishing a self-reinforcing circuit of cytokine-mediated activation as depicted in Fig.
6c. In support, stimulated CAT CAR T-cells consistently formed larger clusters than FMC63 CAR T-cells
in culture (data not shown). T-cell aggregates are also instrumental in mediating the synaptic exchange
of cytokines, thus further amplifying and coordinating immune responses. Notably, cytokine crosstalk
among T-cells can trigger antigenic-independent stimulation and this T-cell activation route is
preferentially taken by memory T-cells as opposed to other T-cell subsets, due to their intrinsically
higher cytokine responsiveness[46]. Of note, p38 MAPK can be activated downstream of both antigen-
and cytokine- mediated stimulation through different mechanisms, but it is only indispensable for the
latter[14], suggesting a cause and effect relationship between the polyfunctional phenotype observed
and the activation of p38 MAPK signalling axis.

Finally, we associated the cytokine polyfunctional phenotype in CAT to CD19 trogocytosis in CAR T-
cells. Trogocytosis requires the uptake of the target antigen by T-cells and can result in either immune
suppression (T-cell anergy and fratricide killing), as recently shown for FMC63- based CARs[50], or
stimulation (increasing cytokine secretion and cell proliferation)[54]. Interestingly, we noted that the
proportion of CAT CAR T-cells undergoing trogocytosis (trog+) was higher than in FMC63 the longer
after stimulation. Considering that one unique property of low-affinity CAR T-cells is the ability to
selectively kill target cells with high levels of antigen (tumour cells) while sparing low antigen-
expressing cells (trog+ CAR T), it is tempting to speculate that reduced fratricide killing is responsible
for the increased persistence of trog+ CAT CAR T-cells in culture. We speculate that trogocytosed CD19
in CAT CAR T-cells may provide sustained intracellular signalling to CAT CAR T-cells between successive
short rounds of antigen ligation typical of low-affinity CARs, consistent with a previously proposed
model for the stimulatory role of trogocytosis in CD4 T-cells[55], enhancing activation of low-affinity CAR
T-cells. In conclusion, we describe the molecular mechanisms underlying the low-affinity CAT CAR T-
cell functional phenotype. We provide evidence that the potent and long-term anti-tumour responses
observed with low-affinity CAT CAR T-cells reflect a distinct pattern of both activation priming and
downstream signalling compared to high-affinity CD19 CAR T-cells which is established and sustained
in CAR T-cells by a self-reinforcing circuit activated via cytokines polyfunctional crosstalk. This work
may inform the future design of versatile CAR T-cells, capable of balancing safety, efficacy and long-
term persistence.
456

457   **Methods**
458

459   **Human peripheral blood samples and cell lines**
460

Peripheral blood was collected from 13 HDs after having obtained informed consent. Fresh
mononuclear cells were isolated by Ficoll density gradient centrifugation (day 0) and used for
experiments. All analyses were performed on fresh T- or CAR T-cells except for 96 h memory subset
FACS analysis. HD details and relative experimental procedure are reported in Supplementary Table
1. The human ALL cell line NALM6 was purchased from DSMZ. Cells were cultured according to
manufacturer's instructions in complete RPMI 1640 medium, GlutaMAX™ Supplement (Gibco™,
Thermo Fisher Scientific, Waltham, MA, USA) supplemented with 10% of heat-inactivated Foetal
Bovine Serum (FBS) (Gibco™, Thermo Fisher Scientific), 100 IU penicillin and 100 μg/ml streptomycin
(Sigma-Aldrich, St. Louis, MO, USA).
470

Ex vivo **T-cell expansion**

Freshly isolated PBMCs from HD1-HD8 were cultured in TexMACS^{TM} medium (Miltenyi Biotec, Bergisch

Gladbach, Germany), an optimised T-cell medium. To induce T-cell expansion, CD3/CD28 beads (CTS^{TM}

Dynabeads^{TM} CD3/CD28, Thermo Fisher Scientific) were added to cells in MACS GMP Cell

Differentiation Bags (Miltenyi Biotec) at a 1:3 lymphocyte:bead ratio. CD3/CD28 beads were

magnetically removed from the culture on day 5 and CAR T-cells rested for 48 h before proceeding to

antigen stimulation.

CAR construct generation

CD19 FMC63 and CAT CAR transfer vectors were generated as previously described[12] and were

provided by P.J.A.. The vector structure is schematised in Supplementary Fig. 1a.

Lentiviral production by transient transfection

Lentiviral production was performed by transient transfection according to the following procedure.

The day before transfection, 5.5 x 10^6 293T cells were seeded per 175 cm^2 flasks in complete DMEM,

high glucose, GlutaMAX^{TM} Supplement (Gibco^{TM}, Thermo Fisher Scientific) supplemented with 10%

FBS.

To generate viral particles, 293T cells were co-transfected with second generation LV packaging

plasmids pMD2.G and pCMV-dR8.74 (plasmidFactory, Bielefeld, Germany) and CAR transfer vectors

using GeneJuice® Transfection Reagent (Merck-KGaA, Darmstadt, Germany) according to

manufacturer's recommendations. LV supernatants were collected at 48 and 72 h post-transfection,

pooled, filtered and stored at -80°C before use.

Lentiviral transduction of T-cells

Following overnight activation with CD3/CD28 beads, 0.5 x 10^6 beads-activated T-cells from HD1-HD8

were suspended in 0.5 ml of TexMACS, transduced to express CD19 CAR construct (FMC63 or CAT)

with 1 ml of LV supernatant in RetroNectin® (Takara Bio, Kusatsu, Shiga, Japan)-coated 24-well plates

and spinoculated at 1,000 g for 40 min at room temperature. Generally, 6-8 x 10^6 beads-activated T-

cells per donor per construct were seeded for transduction. Transduction efficiency was assessed by

FACS evaluating mCherry (LV fluorescent marker) expression, at day 8 (7 days after transduction).

UNTR beads-activated T-cells per each donor were used as a control throughout the experiment and

were subjected to the same experimental conditions as LV-transduced ones.

NALM6 and CAR T-cells co-cultures

To assess the anti-tumour activity of CD19 CAR T-cells, we set-up co-cultures with CD19-expressing

NALM6 for 3, 24 or 96 h (stimulated conditions) on day 7 (6 days after transduction and 2 days post-

CD3/CD28 beads removal). Briefly, beads-activated T-cells from HD1-HD8 were incubated for 48 h in

optimised T-cell medium before stimulation. 0.1×10^6 UNTR or FMC63 or CAT beads-activated T-cells

were seeded in 96-well plates in complete TexMACS medium in a 1:1 ratio with irradiated (40 Gy)

NALM6. Generally, $0.1-1.5 \times 10^6$ beads-activated T-cells per donor per condition (UNTR, FMC63 and

CAT) were stimulated.

Unstimulated, beads-activated T-cells per each donor, condition and time point were kept in culture

under the same experimental conditions as antigen-stimulated T-cells.

Flow cytometry activated cell sorting and antibody staining

All FACS experiments included Fluorescence minus one (FMO) and single-antibody stained BD^{TM}

CompBeads (BD Biosciences) controls to set expression threshold and to calculate compensation,

respectively. Live cells were selected based on their non-permeability and lack of fluorescence

associated with DAPI Solution (BD Biosciences, Franklin Lakes, NJ, USA). The combination of

monoclonal antibodies used to identify T-cell subsets is described in the main text. Following 24 or 96

h of stimulation, cells from HD1-HD8 in all the experimental conditions were labelled with BV510-anti

CD3 (clone OKT3), APC-CY7-anti CD4 (clone SK3), PerCP/CY5.5-anti CD8 (clone RPA-T8), AF647-anti

CD62L (clone DREG-56), BV785-anti CD45RA (clone HI100) (BioLegend, San Diego, CA, USA) in order

to detect CD4, CD8 and memory subsets within live T-/CAR T-lymphocytes. PBMCs from HD9-HD13

were labelled with PE-anti CD19 (clone HIB19; BioLegend) to detect B-cells within lymphocytes in the

samples soon after isolation. Experiments were performed on a cell sorter FACSAria^{TM} III (BD

Biosciences) and on a CytoFLEX analyzer (Beckman Coulter Inc., Brea, CA, USA) and analysed with

FlowJo^{TM} software v10.6.1 (BD Biosciences) and Cytobank platform (www.cytobank.org).

RNA sequencing libraries preparation and sequencing

RNA sequencing libraries were generated from 8 donors (HD1-HD8), across six experimental
conditions, UNTR T-cells or FMC63 or CAT CAR T-cells in either absence or presence of NALM6
(unstimulated or stimulated). 1,000 T-cells/CAR T-cells were FACS-sorted in single PCR tubes
containing 4 ul of QIAGEN lysis buffer from QIAseq FX Single Cell RNA Library Kit (QIAGEN, Hilden,
Germany) and stored at -80°C. The kit steps include reverse transcription, cDNA amplification and
PCR-free NGS library preparation. Libraries were purified using AMPure XP magnetic beads (Beckman
Coulter Inc.,) according to the manufacturer instructions. After beads purification the libraries were
resuspended in 50 µL of buffer EB (Qiagen) and stored at -20 °C. Size and concentration of the libraries
generated was assessed using High Sensitivity Bioanalyzer (Agilent, Santa Clara, CA, USA) and Qubit
High-Sensitivity DNA kit (Invitrogen™, Thermo Fisher Scientific), respectively. Libraries were
equimolarly pooled to a final concentration of 10 nM and were sequenced with Illumina HiSeq 3000
(150 bp pair-end reads) at UCL Genomics.

RNA sequencing analysis

Sequencing quality control and mapping was performed on Fedora 32. DGE analysis and downstream
analyses were performed in *R 4.0.2* on Windows 10 (build 18362). Additional details can be found in
the transcriptomic Supplementary Bioinformatics Material (SBM). An FDR smaller than 0.1 (when
multiple tests were performed) or a *P* value smaller than 0.05 (when only one test was performed)
was considered statistically significant. Quality control and adapters trimming was performed with
FastQC 0.11.9[56] and *cutadapt 2.10*[57]. Sequences were mapped with *STAR 2.7.3a*[58] to the release 31 of
the Human Genome (GRCh38.p12)[59] and summarised with *featureCounts* from the *subread 2.0.0*
software package[60]. Only genes annotated as protein-coding genes were included into the DE and
downstream analyses. These were further filtered by the *filterByExpr* function from the *edgeR*
package[61] in order to remove lowly expressed genes. DE was assessed with *DESeq2*[62]. FDR were
calculated using Benjamini-Hochberg correction[63].

Only the genes which passed the *filterByExpr* filtering were used for GSEA pathway analysis. GSEA was
performed with *fgsea* package[64] on a subset of pathways from the MSigDB's Hallmarks pathways[65]
(see SBM Table III). The log$_2$Fold Change of the DE conditions compared with their matched UNTR
condition was used as an input of the analysis. Relative cell type enrichment was performed with
xCell[26] on a subset of cell types (SBM Table IV).

Mass cytometry analysis

Following 3 (T1) or 24 h (T2) of stimulation, samples from HD4-HD8 in all experimental conditions

were treated with Brefeldin A (BioLegend) (1:1,000) at 37°C for 4 h to favour intracellular cytokine

accumulation and were fixed with 1.6% of formaldehyde for 10 min at room temperature. Fixed

samples were stained with 0.05 µM Cell-ID™ Cisplatin-194-Pt (Fluidigm, South San Francisco, CA,

USA), a live-dead marker, for 1 min on a rocker. Samples for all experimental conditions from a given

donor and stimulation time point were then barcoded using the Cell-ID™ 20-Plex Pd Barcoding Kit

(Fluidigm) according to manufacturer's instructions. 12 samples were labelled concurrently in the

same tube following a three-step staining protocol. Briefly, 2.6-3.7 x 10^6 cells were firstly stained with

antibodies against surface markers (CD3, CD4, CD8a, CD62L, CD45RA, CD19, CD25, CD127, HLA-DR).

Following permeabilization with eBioscience™ Foxp3/Transcription Factor Staining Buffer Set

(Thermo Fisher Scientific) for 30 min at 4°C, cells were subsequently stained for the detection of

intracellular cytokines (IL-4, IL-5, IL-6, IL-23, IL-17A, IL-2, TNF-α, IFN-γ, GM-CSF, TGF-β, Granzyme B

and Perforin B) and cytoplasmic protein (RFP, used to detect mCherry expression). Cells were then

fixed in ice-cold 50% Methanol for 10 min at 4°C and underwent antibody staining for nuclear targets

(NFAT1 and FOXP3) and phosphoproteins (pSTAT4, pSTAT5, pp38, pERK1/2, pZAP70, pS6, pCREB and

pRB). Mass cytometry antibody details are listed in Supplementary Table 3. Stained cells were fixed

with 1.6% of formaldehyde for 10 min at room temperature on a rocker and incubated with Cell-ID™

Intercalator-Ir (Fluidigm) overnight at 4°C. The following day, experiments were performed on a Helios

mass cytometer (Fluidigm). EQ™ Four Element Calibration Beads (Fluidigm) were added to cell

suspensions immediately before acquisition to guarantee inter-sample comparability.

Mass cytometry data pre-processing and analysis

The resulting FCS files were normalised against EQ™ Four Element Calibration Beads, debarcoded and

uploaded on Cytobank platform. Viable single-cells were cleaned based on Gaussian parameters

(Event length, Centre, Offset and Residual), DNA content (193-Ir DNA) and Cisplatin (194-Pt) and gated

on CD19, CD3 and mCherry markers to detect B-cells, T-/CAR T-lymphocytes and CD19+ CAR T-cells.

Cell populations were exported from Cytobank and computationally analysed in Python (Python

version 3.7.6) or in R (R version 3.6.1) as described below.

Specifically, (EMD[19] and DREMI[45] scores were computed separately for each given stimulation

condition and time point between T-, CAR T-lymphocytes or CD19+ CAR T-cells and the entire cell

population as previously described[66] using CyGNAL repository (https://github.com/TAPE-

Lab/CyGNAL).

For the comparison between unstimulated and stimulated conditions, EMD scores were computed

separately for each given donor, culture condition (FMC63, CAT) and time point between CD3+ CAR

T-cells and the entire cell population within conditions.

The PCA performed on EMD scores of all 32 markers and the event counts in CD3+ T-cells from HD4-

HD8 across all the experimental conditions (Supplementary Fig. 3a, c-d) highlighted the lack of

activation of HD6 CAR T-cells upon antigen stimulation. For this reason, HD6 was excluded from the

analyses. PCA in Fig. 2d was generated on EMD scores of all 32 markers calculated on CD3+ T-cells

across all experimental conditions at T2. Heatmaps were generated using *pheatmap* package in R on

selected EMD (30 markers) scores or DREMI scores statistically different between the 2 CARs

normalised by row across specified experimental conditions.

Network analyses were performed using *tidygraph* and *ggraph* packages in R to compare signalling

connections in the two CAR constructs in specified experimental conditions. Nodes and edges were

coloured according to HD-averaged EMD and DREMI scores, respectively, which were computed on

CAR T-lymphocytes.

Additionally, pre-gated FCS files were imported in R, arcsinh transformed (cofactor = 5) and cell

clustered using the R packages *FlowSOM*[24] and *ConsensusClusterPlus*[67] as computed in[68]. The code for

the dimensional reduction analysis provided[68] was modified using R package *umap* for UMAP analysis.

UMAP analyses were performed downsampling at 1,000 cells and using default parameters.

Cytokine co-expression analysis was computed in R on CAR T (CD3+mCherry+) or trog+ CAR T

(CD3+mCherry+CD19+) pre-gated FCS files concatenated per donor for each given experimental

condition. An expression threshold of 1 was set for all the arcsinh transformed cytokine markers

except for TGF-ß, for which a threshold of 4 was selected. Chord diagrams of cytokine co-expression

were created with the R package *circlize*[69].

DNA staining intensity was assessed on unstimulated CAR T and on stimulated CD19- (CD19 expression

< 1) and CD19+ (CD19 expression >= 1) CAR T pre-gated FCS files arcsinh transformed.

Statistical analyses

Data are shown as mean $\pm$ standard error (se) and statistical analyses were performed in R or

GraphPad Prism 7 (GraphPad Software, LA Jolla, CA, USA) calculating Paired samples t-tests across

experimental conditions unless otherwise stated.

EMD, DREMI and polyfunctionality-related statistical analyses between selected experimental

conditions are reported in Supplementary Table 5.

Data availability

The RNA sequencing data and analyses are available at NCBI's Gene Expression Omnibus (GEO) data repository with the accession code GSE157584 and in GitHub (https://github.com/EduardoGCCM/CATvsFMC63_Michelozzi).
Mass cytometry raw and processed data will be made publicly available at https://community.cytobank.org/.

References

1. Miliotou, A.N. & Papadopoulou, L.C. CAR T-cell Therapy: A New Era in Cancer Immunotherapy. *Curr Pharm Biotechnol* **19**, 5-18 (2018).

2. Pehlivan, K.C., Duncan, B.B. & Lee, D.W. CAR-T Cell Therapy for Acute Lymphoblastic Leukemia: Transforming the Treatment of Relapsed and Refractory Disease. *Curr Hematol Malig Rep* **13**, 396-406 (2018).

3. Lee, D.W.*, et al.* Current concepts in the diagnosis and management of cytokine release syndrome. *Blood* **124**, 188-195 (2014).

4. Gust, J.*, et al.* Endothelial Activation and Blood-Brain Barrier Disruption in Neurotoxicity after Adoptive Immunotherapy with CD19 CAR-T Cells. *Cancer Discov* **7**, 1404-1419 (2017).

5. Watanabe, K., Kuramitsu, S., Posey, A.D., Jr. & June, C.H. Expanding the Therapeutic Window for CAR T Cell Therapy in Solid Tumors: The Knowns and Unknowns of CAR T Cell Biology. *Front Immunol* **9**, 2486 (2018).

6. Kowolik, C.M.*, et al.* CD28 costimulation provided through a CD19-specific chimeric antigen receptor enhances in vivo persistence and antitumor efficacy of adoptively transferred T cells. *Cancer Res* **66**, 10995-11004 (2006).

7. Singh, A.P.*, et al.* Development of a quantitative relationship between CAR-affinity, antigen abundance, tumor cell depletion and CAR-T cell expansion using a multiscale systems PK-PD model. *MAbs* **12**, 1688616 (2020).

8. Chmielewski, M., Hombach, A., Heuser, C., Adams, G.P. & Abken, H. T cell activation by antibody-like immunoreceptors: increase in affinity of the single-chain fragment domain above threshold does not increase T cell activation against antigen-positive target cells but decreases selectivity. *J Immunol* **173**, 7647-7653 (2004).

9. Caruso, H.G.*, et al.* Tuning Sensitivity of CAR to EGFR Density Limits Recognition of Normal Tissue While Maintaining Potent Antitumor Activity. *Cancer Res* **75**, 3505-3518 (2015).

673   10.   Liu, X*., et al.* Affinity-Tuned ErbB2 or EGFR Chimeric Antigen Receptor T Cells Exhibit an
674   Increased Therapeutic Index against Tumors in Mice. *Cancer Res* **75**, 3596-3607 (2015).

675   11.   Imai, C*., et al.* Chimeric receptors with 4-1BB signaling capacity provoke potent cytotoxicity
676   against acute lymphoblastic leukemia. *Leukemia* **18**, 676-684 (2004).

677   12.   Ghorashian, S*., et al.* Enhanced CAR T cell expansion and prolonged persistence in pediatric
678   patients with ALL treated with a low-affinity CD19 CAR. *Nat Med* **25**, 1408-1414 (2019).

679   13.   Ying, Z*., et al.* A safe and potent anti-CD19 CAR T cell therapy. *Nat Med* **25**, 947-953 (2019).

680   14.   Valitutti, S., Muller, S., Cella, M., Padovan, E. & Lanzavecchia, A. Serial triggering of many T-
681   cell receptors by a few peptide-MHC complexes. *Nature* **375**, 148-151 (1995).

682   15.   Caserta, S., Kleczkowska, J., Mondino, A. & Zamoyska, R. Reduced functional avidity promotes
683   central and effector memory CD4 T cell responses to tumor-associated antigens. *J Immunol*
684   **185**, 6545-6554 (2010).

685   16.   Wang, X*., et al.* Phase 1 studies of central memory-derived CD19 CAR T-cell therapy following
686   autologous HSCT in patients with B-cell NHL. *Blood* **127**, 2980-2990 (2016).

687   17.   Wang, X*., et al.* Comparison of naive and central memory derived CD8(+) effector cell
688   engraftment fitness and function following adoptive transfer. *Oncoimmunology* **5**, e1072671
689   (2016).

690   18.   Turtle, C.J*., et al.* CD19 CAR-T cells of defined CD4+:CD8+ composition in adult B cell ALL
691   patients. *J Clin Invest* **126**, 2123-2138 (2016).

692   19.   Orlova, D.Y*., et al.* Earth Mover's Distance (EMD): A True Metric for Comparing Biomarker
693   Expression Levels in Cell Populations. *PLoS One* **11**, e0151859 (2016).

694   20.   Fuzzy Analysis (Program FANNY). in *Finding Groups in Data* 164-198 (1990).

695   21.   Gomes-Silva, D*., et al.* Tonic 4-1BB Costimulation in Chimeric Antigen Receptors Impedes T
696   Cell Survival and Is Vector-Dependent. *Cell Rep* **21**, 17-26 (2017).

697   22.   Frigault, M.J*., et al.* Identification of chimeric antigen receptors that mediate constitutive or
698   inducible proliferation of T cells. *Cancer Immunol Res* **3**, 356-367 (2015).

699   23.   Long, A.H*., et al.* 4-1BB costimulation ameliorates T cell exhaustion induced by tonic signaling
700   of chimeric antigen receptors. *Nat Med* **21**, 581-590 (2015).

701   24.   Van Gassen, S*., et al.* FlowSOM: Using self-organizing maps for visualization and interpretation
702   of cytometry data. *Cytometry A* **87**, 636-645 (2015).

703   25.   Krummey, S.M*., et al.* CD45RB Status of CD8(+) T Cell Memory Defines T Cell Receptor Affinity
704   and Persistence. *Cell Rep* **30**, 1282-1291 e1285 (2020).

705   26.   Aran, D., Hu, Z. & Butte, A.J. xCell: digitally portraying the tissue cellular heterogeneity
706   landscape. *Genome Biol* **18**, 220 (2017).

27.   Allan, S.E.*, et al.* Activation-induced FOXP3 in human T effector cells does not suppress
proliferation or cytokine production. *Int Immunol* **19**, 345-354 (2007).

28.   Ran, Q.*, et al.* CRIF1 interacting with CDK2 regulates bone marrow microenvironment-induced
G0/G1 arrest of leukemia cells. *PLoS One* **9**, e85328 (2014).

29.   Carrette, F. & Surh, C.D. IL-7 signaling and CD127 receptor regulation in the control of T cell
homeostasis. *Semin Immunol* **24**, 209-217 (2012).

30.   Kaech, S.M.*, et al.* Selective expression of the interleukin 7 receptor identifies effector CD8 T
cells that give rise to long-lived memory cells. *Nat Immunol* **4**, 1191-1198 (2003).

31.   Holling, T.M., van der Stoep, N., Quinten, E. & van den Elsen, P.J. Activated human T cells
accomplish MHC class II expression through T cell-specific occupation of class II transactivator
promoter III. *J Immunol* **168**, 763-770 (2002).

32.   Gourley, T.S. & Chang, C.H. Cutting edge: the class II transactivator prevents activation-
induced cell death by inhibiting Fas ligand gene expression. *J Immunol* **166**, 2917-2921 (2001).

33.   Di Rosa, F.*, et al.* HLA class II molecules transduce accessory signals affecting the CD3 but not
the interleukin-2 activation pathway in T blasts. *Hum Immunol* **38**, 251-260 (1993).

34.   Soroosh, P., Ine, S., Sugamura, K. & Ishii, N. OX40-OX40 ligand interaction through T cell-T cell
contact contributes to CD4 T cell longevity. *J Immunol* **176**, 5975-5987 (2006).

35.   Sabatos, C.A.*, et al.* A synaptic basis for paracrine interleukin-2 signaling during homotypic T
cell interaction. *Immunity* **29**, 238-248 (2008).

36.   Thurley, K., Gerecht, D., Friedmann, E. & Hofer, T. Three-Dimensional Gradients of Cytokine
Signaling between T Cells. *PLoS Comput Biol* **11**, e1004206 (2015).

37.   Castellino, F.*, et al.* Chemokines enhance immunity by guiding naive CD8+ T cells to sites of
CD4+ T cell-dendritic cell interaction. *Nature* **440**, 890-895 (2006).

38.   Hugues, S.*, et al.* Dynamic imaging of chemokine-dependent CD8+ T cell help for CD8+ T cell
responses. *Nat Immunol* **8**, 921-930 (2007).

39.   Yiu, H.H., Graham, A.L. & Stengel, R.F. Dynamics of a cytokine storm. *PLoS One* **7**, e45027
(2012).

40.   Betts, M.R.*, et al.* HIV nonprogressors preferentially maintain highly functional HIV-specific
CD8+ T cells. *Blood* **107**, 4781-4789 (2006).

41.   Caccamo, N.*, et al.* Multifunctional CD4(+) T cells correlate with active Mycobacterium
tuberculosis infection. *Eur J Immunol* **40**, 2211-2220 (2010).

42.   Rossi, J.*, et al.* Preinfusion polyfunctional anti-CD19 chimeric antigen receptor T cells are
associated with clinical outcomes in NHL. *Blood* **132**, 804-814 (2018).

43.   Xhangolli, I.*, et al.* Single-cell Analysis of CAR-T Cell Activation Reveals A Mixed TH1/TH2
Response Independent of Differentiation. *Genomics Proteomics Bioinformatics* **17**, 129-139
(2019).

44.   Frucht, D.M. IL-23: a cytokine that acts on memory T cells. *Sci STKE* **2002**, pe1 (2002).

45.   Krishnaswamy, S.*, et al.* Systems biology. Conditional density-based analysis of T cell signaling
in single-cell data. *Science* **346**, 1250689 (2014).

46.   Geginat, J., Sallusto, F. & Lanzavecchia, A. Cytokine-driven proliferation and differentiation of
human naive, central memory, and effector memory CD4(+) T cells. *J Exp Med* **194**, 1711-1719
(2001).

47.   Li, C.*, et al.* Activation of p38 mitogen-activated protein kinase is critical step for acquisition
of effector function in cytokine-activated T cells, but acts as a negative regulator in T cells
activated through the T-cell receptor. *Immunology* **132**, 104-110 (2011).

48.   Dodeller, F. & Schulze-Koops, H. The p38 mitogen-activated protein kinase signaling cascade
in CD4 T cells. *Arthritis Res Ther* **8**, 205 (2006).

49.   Clark, A.R., Dean, J.L. & Saklatvala, J. Post-transcriptional regulation of gene expression by
mitogen-activated protein kinase p38. *FEBS Lett* **546**, 37-44 (2003).

50.   Hamieh, M.*, et al.* CAR T cell trogocytosis and cooperative killing regulate tumour antigen
escape. *Nature* **568**, 112-116 (2019).

51.   Rachmilewitz, J. & Lanzavecchia, A. A temporal and spatial summation model for T-cell
activation: signal integration and antigen decoding. *Trends Immunol* **23**, 592-595 (2002).

52.   Weng, N.P., Araki, Y. & Subedi, K. The molecular basis of the memory T cell response:
differential gene expression and its epigenetic regulation. *Nat Rev Immunol* **12**, 306-315
(2012).

53.   Sheih, A.*, et al.* Clonal kinetics and single-cell transcriptional profiling of CAR-T cells in patients
undergoing CD19 CAR-T immunotherapy. *Nat Commun* **11**, 219 (2020).

54.   Ahmed, K.A., Munegowda, M.A., Xie, Y. & Xiang, J. Intercellular trogocytosis plays an
important role in modulation of immune responses. *Cell Mol Immunol* **5**, 261-269 (2008).

55.   Osborne, D.G. & Wetzel, S.A. Trogocytosis results in sustained intracellular signaling in CD4(+)
T cells. *J Immunol* **189**, 4728-4739 (2012).

56.   Andrews, S. FastQC: A quality control tool for high throughput sequence data. (2019).

57.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
2011 **17**, 3 (2011).

58.   Dobin, A.*, et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

59.   Frankish, A.*, et al.* GENCODE reference annotation for the human and mouse genomes.
Nucleic Acids Research **47**, D766-D773 (2018).

60.   Liao, Y., Smyth, G.K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).

61.   Chen, Y., Lun, A. & Smyth, G. From reads to genes to pathways: differential expression analysis
of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline [version 2;
peer review: 5 approved]. *F1000Research* **5**(2016).

62.   Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

63.   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B
(Methodological)* **57**, 289-300 (1995).

64.   Korotkevich, G., Sukhov, V. & Sergushichev, A. Fast gene set enrichment analysis. *bioRxiv*,
060012 (2019).

65.   Liberzon, A.*, et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection.
Cell Syst **1**, 417-425 (2015).

66.   Qin, X.*, et al.* Cell-type-specific signaling networks in heterocellular organoids. *Nat Methods*
17, 335-342 (2020).

67.   Wilkerson, M.D. & Hayes, D.N. ConsensusClusterPlus: a class discovery tool with confidence
assessments and item tracking. *Bioinformatics* **26**, 1572-1573 (2010).

68.   Nowicka, M.*, et al.* CyTOF workflow: differential discovery in high-throughput high-
dimensional cytometry datasets. *F1000Res* **6**, 748 (2017).

69.   Gu, Z., Gu, L., Eils, R., Schlesner, M. & Brors, B. circlize Implements and enhances circular
visualization in R. *Bioinformatics* **30**, 2811-2812 (2014).

Acknowledgements

We are grateful to Dr Ayad Eddaudi (UCL ICH Flow Cytometry facility) for support in cell-sorting. We
are grateful to Dr Thomas Adejumo (Fluidigm) for mass cytometry valuable suggestions and assistance
with protocols design. We thank Dr Anne Marijn Kramer (Amsterdam UMC) for generating CD19
FMC63 and CAT CAR transfer vector plasmids with S.G. and M.P. and P.J.A. We thank Mr Efstratios
Kirtsios (UCL) for useful discussion. The authors acknowledge the contribution of UCL Genomics
Facility. This work was supported by the NIHR GOSH BRC (NIHR GOSH BRC 17PA01), the views
expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, or the

Department of Health. Part of this work was supported by the Leukaemia UK John Goldman Fellowship

to A.G. (2018/JGF/003), the Rosetrees Trust fund to A.G. (M700) and the Academy of Medical Sciences

Springboard Award to A.G. (SBF004\1025).

Author contributions

I.M.M. designed, performed and analyzed experiments, performed bioinformatic analyses of mass

cytometry experiments and contributed to writing the manuscript. E.G-C performed bioinformatic

analyses of transcriptomic data and wrote the relative bioinformatic supplementary information.

R.V.C.P. performed experiments and bioinformatic analyses of mass cytometry and transcriptomic

data. F.C-R provided data analysis tools and contributed to bioinformatic analyses of mass cytometry

experiments. P.P-C. performed transcriptomic bioinformatic analyses data checks and contributed to

writing the bioinformatic supplementary information. J. S., M.S., S.W.W and A.Gu. performed

experiments. S.G. and M.P. provided CAR constructs. C.J.T provided expertise in mass cytometry and

reagents. P.J.A. provided reagents and expertise and contributed to writing the manuscript. S.C.

supervised the bioinformatic analyses and contributed to writing the manuscript. A.G. designed and

supervised the project, performed and analyzed experiments and wrote the manuscript. All authors

provided critical feedback and helped shape the research, analysis and manuscript.

Competing interests

The authors declare no competing interests.

Correspondence

Corrispondente to Dr Alice Giustacchini.

Fig. 1: Generation and phenotypic characterisation of CAR T-cells from HD-PBMCs.

a Experimental workflow. Peripheral Blood Mononuclear Cells (PBMCs) were isolated from HDs and LV transduced to express CD19 CAR construct (FMC63 or CAT) following overnight activation with CD3/CD28 beads.

Six days after transduction, CAR T-cells were cultured without (unstimulated) or with target cells (NALM6) at 1:1 ratio (stimulated). Unstimulated and stimulated cells were analysed by flow cytometry and sorted for RNA-sequencing 24 hours post-stimulation. Mass cytometry analysis was performed on unstimulated and stimulated cells at 3 h (T1) and 24 h (T2) post-stimulation.

Activated UNTR T-cells were used as a control throughout the experiment.

b (left) Representative FACS plots for mCherry expression in UNTR T-cells and FMC63 and CAT CAR T-cells at 7 days post-transduction. (middle) Spaghetti plots showing transduction levels of CAR T-cells as percentage of mCherry+ (in CD3+) and (right) as MFI of mCherry in transduced T-cells. Lines connect results from individual donors. Unstimulated (top) and stimulated conditions (bottom) are reported.

c (left) Representative FACS plots showing CD4 and CD8 composition of UNTR T-cells and FMC63 and CAT CAR T-cells. (middle) Variation ($\log_2$ fold change) of CD4 and CD8 proportion in UNTR T-cells and FMC63 and CAT CAR T-cells. The dotted horizontal line (0) represents the conditions in which CD4=CD8. Unstimulated (top) and stimulated conditions (bottom) are reported. (right) Percentage of CD8+ cells in unstimulated and stimulated UNTR T-cells and FMC63 and CAT CAR T-cells.

Data represent mean ± se.

b-c Statistical significance was calculated by Paired t-test. *$P < 0.05$, ***$P < 0.001$, ****$P < 0.0001$. n = 8 HDs, n = 2 independent experiments.

Each experimental condition is indicated by a specific colour code (Unstimulated conditions = UNTR light grey, FMC63 light blue, CAT orange; stimulated conditions = UNTR grey, FMC63 blue, CAT red).

FIG 2

Fig. 2: RNA-seq and mass cytometry analyses of unstimulated and stimulated untransduced and CAR-transduced T-cells.

a Correlation of mCherry MFI measured by FACS with normalised RNA-seq counts (CPM) aligning to the scFv region of each of the two CARs (n = 6 HDs, n = 2 independent experiments). Pearson correlation and *P* values are displayed.

b Correlation of the percentage of CD8+ T-cells detected by FACS with *CD8* mRNA levels (n=6 HDs, n = 1 independent experiment). Pearson correlation and *P* values are displayed.

c PCA of the top 500 variable genes from RNA-seq analysis across all the experimental conditions (n = 6 HDs, n = 2 independent experiments).

d PCA of mass cytometry EMD scores computed at T2 (24 h stimulation) in CD3+ cells across all the experimental conditions (n = 4 HDs, n = 1 independent experiment).

e Heatmap showing the hierarchical clustering of all the experimental conditions analysed displaying the top 20 variable genes from RNA-seq PC1 (n = 6 HDs, n = 2 independent experiments).

f Fuzzy clustering analysis of RNA-seq data across all experimental conditions (n = 6 HDs, n = 2 independent experiments).

g GSEA of HALLMARK gene sets for (1) FMC63 or (2) CAT CAR T unstimulated condition *vs* UNTR unstimulated cells, (3) FMC63 or (4) CAT CAR T stimulated condition *vs* UNTR stimulated cells (n = 6 HDs, n = 2 independent experiments). Non-significant enrichment scores were set to zero. Gene sets with no significant enrichment scores are not shown.

a-g Each experimental condition is indicated by a specific colour code (Unstimulated conditions = UNTR light grey, FMC63 light blue, CAT orange; stimulated conditions = UNTR grey, FMC63 blue, CAT red).

FIG 3

Fig. 3: Molecular characterisation of unstimulated CAR-transduced T-cells.

a (left) Volcano plot showing DE genes between unstimulated FMC63 and CAT CAR T-cells. The dashed horizontal line represents the statistical significance threshold (FDR < 0.1). (right) The barplots show the expression of selected DE genes (FDR < 0.1) in unstimulated untransduced and transduced T-cells (n = 6 HDs, n = 2 independent experiments).

b (left) UMAP showing the expression of selected markers (CD4, CD8a, CD25, Granzyme B) detected by mass cytometry at T2 (24 h stimulation) in unstimulated CD3+ cells across all the experimental conditions. The colour of the scale reflects the protein expression levels.

(right) The barplots show the expression of mass cytometry EMD scores for CD25 and Granzyme B in unstimulated CD3+ cells across all the experimental conditions at T2. The dotted horizontal line (0) represents the average expression of a specific marker across all the experimental conditions analysed. (n = 4 HDs, n = 1 independent experiment).

c (left) UMAP representation of the 4 cell populations (CD8, CD4, CD3 low/neg, CD19) identified by FlowSOM analysis in unstimulated samples analysed by mass cytometry at T2 (24 h stimulation). The cell types are indicated by different colours. (right) Percentage of residual B-cells detected by mass cytometry in unstimulated samples at T2. (n = 4 HDs, n = 1 independent experiment).

d (top) Barplot showing the percentage of T_{CM} (CD45RA-CD62L+) in unstimulated CD8+ UNTR T-cells and FMC63 and CAT CAR T-cells (n = 3 HDs, n = 1 independent experiment) measured by FACS. (bottom) xCell score inferring the enrichment of CD8 T_{CM} in bulk RNA-seq results from unstimulated UNTR (CD3+) T-cells and FMC63 and CAT (CD3+mCherry+) CAR T-cells (n = 6 HDs, n = 2 independent experiments).

a-d Barplots show mean ± se. Statistical significance was calculated by Paired t-test. *P < 0.05, **P < 0.01. Each experimental condition is indicated by a specific colour code (UNTR light grey, FMC63 light blue, CAT orange).

Fig. 4: Molecular characterisation of stimulated CAR-transduced T-cells.

a (left) Volcano plot showing DE genes between FMC63 and CAT CAR T-cells upon NALM6 co-culture. The dashed horizontal line represents the statistical significance threshold (FDR < 0.1). (right) The barplots show the expression of selected DE genes (FDR < 0.1) in stimulated UNTR T-cells and in CAR T-cells (n = 6 HDs, n = 2 independent experiments).

b (left) UMAP showing the expression of selected markers (CD4, CD8a, pRB, CD25, NFAT1, HLA-DR, CD127) detected by mass cytometry at T2 (24 h stimulation) in stimulated CD3+ cells across all the experimental conditions (UNTR, FMC63, CAT). The colour of the scale reflects the protein expression levels.

(right) The barplots show the expression of mass cytometry EMD scores for CD8a, pRB, CD25, NFAT1, HLA-DR and CD127 in stimulated CD3+ cells across all the experimental conditions at T2 (24 h stimulation). The dotted horizontal line (0) represents the average expression of a specific marker across all the experimental conditions analysed. (n = 4 HDs, n = 1 independent experiment). Statistical significance was calculated by Paired t-test. *P < 0.05, **P < 0.01.

a-b Barplots show mean ± se. Each experimental condition is indicated by a specific colour code (UNTR grey, FMC63 blue, CAT red).

FIG 5

Fig. 5: Cytokine polyfunctionality in CAR-transduced T-cells at 3 h post-stimulation.

a The stacked barplots show the percentage of stimulated (top) and unstimulated (bottom) CAR T-cells (CD3+mCherry+) expressing 1 to 4 or >=5 cytokines/cell at T1 (3 h stimulation) as measured by mass cytometry.

b The circos plots show all the combinations of the 12 cytokines analysed in stimulated FMC63 (left) and CAT (right) CAR T-cells at T1 (3 h stimulation) by mass cytometry. The numbers indicate patterns of cytokine co-expression (from 1 to 7 cytokines/cell). A specific colour code has been assigned to each cytokine.

c The barplots show the percentage of the statistically significant combinations of cytokines co-expressed in stimulated FMC63 and CAT CAR T-cells at T1 (3 h stimulation) by mass cytometry. Data represent mean ± se. Statistical significance was calculated by Paired t-test. *$P < 0.05$, **$P < 0.01$. Each experimental condition is indicated by a specific colour code (FMC63 = blue, CAT = red).

a-c n = 4 HDs, n = 1 independent experiment.

FIG 6

Fig. 6: Intracellular signalling analyses in CAR-transduced T-cells at 24 h post-stimulation

a The barplots show the expression of mass cytometry EMD scores for selected phosphoproteins (pS6, pRB, pZAP70, pp38, pERK) in stimulated FMC63 and CAT CAR T-cells at T2 (24 h stimulation). The dotted horizontal line (0) represents the average expression of a specific marker across all the experimental conditions analysed. Barplots show mean ± se (n = 4 HDs, n = 1 independent experiment). Statistical significance was calculated by Paired t-test. *$P < 0.05$, **$P < 0.01$. Each experimental condition is indicated by a specific colour code (FMC63 blue, CAT red).

b Differential signalling network in stimulated FMC63 (left) and CAT (right) CAR T-cells at T2 (24 h stimulation) post-stimulation. The nodes are coloured by HD-averaged EMD scores computed between stimulated CAR T-cells at T2 and arrows, whose transparency reflects HD-averaged DREMI scores, indicate differential markers connectivity between the two CAR products ($P <= 0.05$, n = 4 HDs, n = 1 independent experiment).

c Schematized model of the proposed intracellular signalling occurring preferentially in activated CAT as compared to FMC63 CAR T-cells upon CD19+ target cell engagement. Stimulated CAT CAR T-cells show sustained activation of mTORC1 pathway with consequent elevated protein synthesis and cellular proliferation and survival (pS6 and pRB). Moreover, p38 MAPK cascade, initiated by the CAR CD3ζ chain through pZAP70, results in cytokine genes transcription (probably mediated by NFAT1 and pCREB) and mRNA stabilisation. The expressed cytokines (including IL-2, IL-4, TNF-α, IFN-γ, GM-CSF) could in turn sustain CAT CAR T-cell activation in an antigen independent manner.

FIG 7

Fig. 7: Molecular characterisation of trogocytic CAR-transduced T-cells.

a Heatmap showing CD19 related-mass cytometry DREMI scores computed at T1 (3 h stimulation) in CD3+ CAR T-cells across all the stimulated experimental conditions.

b UMAP representation across experimental conditions of the changes in proportion of 4 cell populations (CD8, CD4, CD8 CD19, CD4 CD19) identified by FlowSOM analysis in CAR T-cells analysed by mass cytometry at T1 (3 h stimulation) and T2 (24 h stimulation). The CAR T-cell subtypes are indicated by different colours. (n = 4 HDs, n = 1 independent experiment).

c Percentage of trog+ CAR T-cells detected by mass cytometry in unstimulated (at T2, 24 h stimulation) and in stimulated transduced samples (at T1 and T2).

d Median intensity of CD19 expression in B-cells and in trog+ CAR T-cells from stimulated CAR-transduced samples at T1 (3 h stimulation) by mass cytometry. Statistical significance was calculated by Mann-Withney test. *$P < 0.05$.

e Percentage of residual NALM6 cells detected by mass cytometry in stimulated samples at T2 (24 h stimulation).

f The barplots show the expression of mass cytometry EMD scores for pRB, pp38, CD25, NFAT1, Granzyme B and Perforin B in stimulated trog+ CAR T-cells at T2 (24 h stimulation). The dotted horizontal line (0) represents the average expression of a specific marker across all the experimental conditions analysed.

g The barplots show the percentage of the statistically different combinations of cytokines co-expressed in stimulated trog+ FMC63 and CAT CAR T-cells at T2 (24 h stimulation) by mass cytometry.

a, c, e-g Statistical significance was calculated by Paired t-test. *$P < 0.05$, **$P<0.01$, ***$P<0.001$.

a, c-g Data represent mean ± se (n = 4 HDs, n = 1 independent experiment). Each experimental condition is indicated by a specific colour code (Unstimulated conditions = FMC63 light blue, CAT orange; stimulated conditions = UNTR grey, FMC63 blue, CAT red).

Suppl FIG 1

Supplementary Fig. 1: Generation and phenotypic characterisation of CAR T-cells from HD-PBMCs.

a (left) Schematic representation of the FMC63 and CAT bicistronic lentiviral vector constructs and (right) of FMC63 and CAT CAR structure. Both constructs are second generation CARs composed by an anti-CD19 scFv, derived from FMC63 or CAT13.1E10 hybridoma, a CD8α hinge, a CD8α transmembrane domain (TM), a 4-1BB costimulatory domain and a CD3ζ signalling domain linked to a reporter gene (mCherry), marker of transduction.

b The bar graph shows the percentage of CD4+ cells in unstimulated and stimulated UNTR T-cells and FMC63 and CAT CAR T-cells. Each experimental condition is indicated by a specific colour code (Unstimulated conditions = UNTR light grey, FMC63 light blue, CAT orange; stimulated conditions = UNTR grey, FMC63 blue, CAT red). Data represent mean ± se. Statistical significance was calculated by Paired t-test. **$P < 0.01$. n = 8 HDs, n = 2 independent experiments.

Suppl FIG 2

Supplementary Fig. 2: RNA-seq analyses of unstimulated and stimulated untransduced and CAR-transduced T-cells.

a Correlation of mCherry MFI measured by FACS with *mCherry* mRNA levels (n = 6 HDs, n = 2 independent experiments). Pearson correlation and *P* values are displayed.

b Correlation of the percentage of CD4+ T-cells detected by FACS with *CD4* mRNA levels (n = 6 HDs, n = 2 independent experiments). Pearson correlation and *P* values are displayed.

c Heatmap showing the hierarchical clustering of all the experimental conditions analysed displaying the top 100 variable genes from RNA-seq PC1 (n = 6 HDs, n = 2 independent experiments).

d 3D fuzzy clustering analysis of RNA-seq data across all experimental conditions (n = 6 HDs, n = 2 independent experiments).

a-d Each experimental condition is indicated by a specific colour code (Unstimulated conditions = UNTR light grey, FMC63 light blue, CAT orange; stimulated conditions = UNTR grey, FMC63 blue, CAT red).

Suppl FIG 3

Supplementary Fig. 3: Mass cytometry analysis of unstimulated and stimulated untransduced and CAR-transduced T-cells.

a PCA plot of EMD scores from unstimulated and stimulated untransduced and CAR-transduced CD3+ T-cells at 3 h (T1) and 24 h (T2) post-stimulation (n= 5 HDs, n = 1 independent experiment). The labels specify each individual donor and the timepoint of the analysis, the colours identify each experimental condition.

b Bar plots of the top contributing variables to PC1 and PC2 in Fig. 2d. The dashed horizontal red line represents the expected average contribution of the variables.

c Bar plots showing the event counts in CD3+ T-cells measured for each sample. Samples are grouped by donor and the number of cells is displayed on top of the bars (n= 5 HDs, n = 1 independent experiment.

d Bar plots of the top contributing variables to PC1 and PC2 in panel a. The dashed horizontal red line represents the expected average contribution of the variables.

a-d Each experimental condition is indicated by a specific colour code (Unstimulated conditions = UNTR light grey, FMC63 light blue, CAT orange; stimulated conditions = UNTR grey, FMC63 blue, CAT red).

a

b

	UNTR unstimulated	FMC63 unstimulated	CAT unstimulated
CD3			
CD19			

Expression intensity

High

Low

UMAP2

UMAP1

Supplementary Fig. 4: Molecular characterisation of unstimulated CAR-transduced T-cells.

a (left) Heatmap showing mass cytometry EMD scores computed at T1 (3 h stimulation) and (right) T2 (24 h stimulation) in CD3+ cells across all the unstimulated experimental conditions (n = 4 HDs, n = 1 independent experiment).

Each condition is indicated by a specific colour code (UNTR light grey, FMC63 light blue, CAT orange).

b UMAP showing the expression of selected markers (CD3, CD19) detected by mass cytometry at T2 (24 h stimulation) in unstimulated CD3+ cells across all the experimental conditions (n = 4 HDs, n = 1 independent experiment). The colour scale reflects the protein expression levels.

Suppl FIG 5

Supplementary Fig. 5: Mass cytometry EMD scores in unstimulated and stimulated CAT CAR-transduced T-cells.

(top) The barplots show mass cytometry EMD scores for activation/proliferation markers and cytokines differentially expressed between unstimulated and stimulated CAT CAR T-cells at T1 (3 h stimulation) and (bottom) at T2 (24 h stimulation). The dotted horizontal line (0) represents the average expression of a specific marker across all the experimental conditions analysed. (n = 4 HDs, n = 1 independent experiment).

Data represent mean ± se. Statistical significance was calculated by Paired t-test. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Each experimental condition is indicated by a specific colour code (Unstimulated condition orange; stimulated condition red).

Suppl FIG 6

Supplementary Fig. 6: EMD scores in unstimulated and stimulated FMC63 CAR-transduced T-cells.

(top) The barplots show mass cytometry EMD scores for activation/proliferation markers and cytokines differentially expressed between unstimulated and stimulated FMC63 CAR T-cells at T1 (3 h stimulation) and (bottom) at T2 (24 h stimulation). The dotted horizontal line (0) represents the average expression of a specific marker across all the experimental conditions analysed. (n = 4 HDs, n = 1 independent experiment).

Data represent mean ± se. Statistical significance was calculated by Paired t-test. *$P < 0.05$, **$P < 0.01$. Each experimental condition is indicated by a specific colour code (Unstimulated condition light blue; stimulated condition blue).

a

Suppl FIG 7

Condition
- UNTR stimulated
- FMC63 stimulated
- CAT stimulated

b

T1

T2

Supplementary Fig. 7: Molecular characterisation of stimulated CAR-transduced T-cells.

a The barplots show the expression of selected DE genes (FDR < 0.1) in stimulated untransduced and transduced T-cells (n = 6 HDs, n = 2 independent experiments).

b (left) Heatmap showing mass cytometry EMD scores computed at T1 (3 h stimulation) and (right) T2 (24 h stimulation) in CD3+ cells across all the stimulated experimental conditions (n = 4 HDs, n = 1 independent experiment).

Each experimental condition is indicated by a specific colour code (UNTR grey, FMC63 blue, CAT red).

Suppl FIG 8

Supplementary Fig. 8: Molecular characterisation and cytokine expression of stimulated CAR-transduced T-cells.

a (left) Heatmap showing mass cytometry EMD scores computed at T1 (3 h stimulation) and (right) T2 (24 h stimulation) in stimulated FMC63 and CAT CAR T-cells (CD3+mCherry+).

b The barplots show the expression of mass cytometry EMD scores for selected effector (Granzyme B, IFN-γ, TNF-α, Perforin B), stimulatory (GM-CSF, IL-2, IL-5) and regulatory (IL-4 and TGF-β) cytokines in stimulated CAR T-cells at T1 (3 h stimulation). The dotted horizontal line (0) represents the average expression of a specific marker across all the experimental conditions analysed.

a-b Data represent mean ± se. Statistical significance was calculated by Paired t-test. *$P < 0.05$.

Each experimental condition is indicated by a specific colour code (UNTR grey, FMC63 blue, CAT red).

Suppl FIG 9

a

FMC63   CAT

b

Supplementary Fig. 9: Cytokine polyfunctionality in stimulated CAR-transduced T-cells at 24 h post-stimulation.

a (left) The circos plots show all the combinations of the 12 cytokines analysed in stimulated FMC63 and (right) CAT CAR T-cells at T2 (24 h stimulation) by mass cytometry. A specific colour code has been assigned to each cytokine. The numbers indicate the frequencies of different patterns of cytokine co-expression (from 1 to 6 cytokines/cell).

b (left) The stacked barplots show the percentage of stimulated and unstimulated (right) CAR T-cells (CD3+mCherry+) expressing 1 to 4 or >=5 cytokines/cell at T2 (24 h stimulation) as measured by mass cytometry.

a-b n = 4 HDs, n = 1 independent experiment.

Suppl FIG. 10

Supplementary Fig. 10: Characterization of intracellular signalling in unstimulated and stimulated CAT and FMC63 CAR T-cells at 24 h post-stimulation.

a The barplots show the expression of mass cytometry EMD scores for NFAT1 and FOXP3 transcription factors in stimulated FMC63 and CAT CAR T-cells at T2 (24 h stimulation). The dotted horizontal line (0) represents the average expression of a specific marker across all the experimental conditions analysed. Barplots show mean ± se (n = 4 HDs, n = 1 independent experiment). Statistical significance was calculated by Paired t-test. *$P < 0.05$.

b Differential signalling network in unstimulated FMC63 and CAT CAR T-cells at T2 (24 h stimulation) post co-culture. The nodes are coloured by HD-averaged EMD scores computed between unstimulated CAR T-cells at T2 and arrows, whose transparency reflects HD-averaged DREMI scores, indicate differential markers connectivity between the 2 CAR products (P <=0.05, n = 4 HDs, n = 1 independent experiment).

c (left) Heatmap showing mass cytometry DREMI scores differentially expressed in unstimulated and (right) stimulated FMC63 and CAT CAR T-cells at T2 (24 h stimulation) (n = 4 HDs, n = 1 independent experiment).

a-b Each condition is indicated by a specific colour code (unstimulated conditions = FMC63 light blue, CAT orange; stimulated conditions = FMC63 blue, CAT red).

Suppl FIG 11

Supplementary Fig. 11: Characterization of CD19-expressing CAR T-cells.

a Heatmap showing mass cytometry DREMI scores computed at T1 (3 h stimulation) in CD3+ CAR T-cells across all the stimulated experimental conditions.

b The barplots show the expression of B-cell (*CD19* and *CD22*)- and T-cell (*CD3D, E, G*)-related genes in stimulated untransduced and transduced T-cells. (n = 6 HDs, n = 2 independent experiments). The significance threshold is FDR < 0.1.

c Histograms showing the DNA marker intensity expression detected by mass cytometry at T1 (3 h stimulation), T2 (24 h stimulation) in unstimulated CAR T-cells and in stimulated trog- and trog+ CAR T-products.

a-c Each condition is indicated by a specific colour code (FMC63 blue, CAT red).

Suppl FIG 12

Supplementary Fig. 12: Molecular characterisation of CAT and FMC63 CAR T-cells undergoing trogocytosis.

a The barplots show the expression of mass cytometry EMD scores for CD19 in stimulated FMC63 and CAT CAR T-cells at T2 (24 h stimulation). The dotted horizontal line (0) represents the average expression of a specific marker across all the experimental conditions analysed.

b (left) Heatmap showing mass cytometry EMD scores computed at T1 (3 h stimulation) and (right) at T2 (24 h stimulation) in trog+ CAR T-cells across all the stimulated experimental conditions.

c The barplots show the expression of mass cytometry EMD scores for CD25 and pS6 in stimulated FMC63 and CAT CAR T-cells at T1 (3 h stimulation). The dotted horizontal line (0) represents the average expression of a specific marker across all the experimental conditions analysed.

d The barplots show the percentage of the statistically different combinations of cytokines co-expressed in stimulated trog+ FMC63 and CAT CAR T-cells at T1 (3 h stimulation) by mass cytometry.

e The barplots show the expression of mass cytometry EMD scores for CD3, CD127, FOXP3 and TGF-ß in stimulated FMC63 and CAT CAR T-cells at T2 (24 h stimulation). The dotted horizontal line (0) represents the average expression of a specific marker across all the experimental conditions analysed.

a, c-e Barplots show mean ± se (n = 4 HDs, n = 1 independent experiment). Statistical significance was calculated by Paired t-test. *$P < 0.05$, ** $P<0.01$.

a-e Each condition is indicated by a specific colour code (FMC63 blue, CAT red).