Title:

EEG Cross-Frequency Phase Synchronization as an Index of

Memory Matching in Visual Search

Authors:

Anna Lena Biel[1], Tamas Minarik[1,2], Paul Sauseng[1]

Affiliations:

1 Department of Psychology, Ludwig-Maximilians-Universität München,

Leopoldstr. 13, 80802 Munich, Germany

2 School of Psychology, University of Birmingham,

Edgbaston, Birmingham, B15 2TT, UK

Contact:

Prof. Dr. Paul Sauseng

Department of Psychology

LMU Munich

Leopoldstr. 13

80802 Munich, Germany

Tel: +49 89 2180 6159

e-mail: paul.sauseng@lmu.de

Short title:

Memory matching in visual search

Abstract

Visual perception is influenced by our expectancies about incoming sensory information. It is assumed that mental templates of expected sensory input are created and compared to actual input, which can be matching or not. When such mental templates are held in working memory, cross-frequency phase synchronization (CFS) between theta and gamma band activity has been proposed to serve matching processes between prediction and sensation. We investigated how this is affected by the number of activated templates that could be matched by comparing conditions where participants had to keep either one or multiple templates in mind for successful visual search. We found that memory matching appeared as transient CFS between EEG theta and gamma activity in an early time window around 150ms after search display presentation, in right hemispheric parietal cortex. Our results suggest that for single template conditions, stronger transient theta-gamma CFS at posterior sites contralateral to target presentation can be observed than for multiple templates. This lends evidence to the idea of sequential attentional templates and is understood in line with previous theoretical accounts strongly arguing for transient synchronization between posterior theta and gamma phase as a neuronal correlate of matching incoming sensory information with contents from working memory.

Introduction

Working memory and selective attention interact in many situations of our everyday life, influencing how we perceive the world. Image yourself looking for your car keys that you must have left somewhere in the kitchen. During your search, you will scan a rich visual environment for something that matches the representation of keys that you have in mind. Such situations are commonly described as visual search. Brought to a cognitive psychology laboratory, participants in a visual search paradigm are usually asked to search for a target object among a number of distractor objects presented on a computer screen. Current theories of attention hold that when we are searching for a target, then keeping a template representation of the target in working memory – a so called attentional template – leads to a bias in the competition for neuronal resources in favour of template-matching stimuli (Bundesen, 1990; Bundesen et al., 2005; Desimone & Duncan, 1995; Duncan & Humphreys, 1989). Insight into the neuronal mechanisms underlying the activation of such mental templates and their comparison with sensory input comes from studies in healthy humans (Soto et al., 2007; Spaak et al., 2015) patients with frontal lesions (Soto et al., 2006; Yago et al., 2004), lesion studies in primates (Everling et al., 2006; Iba & Sawaguchi, 2002; Rossi et al., 2007), as well as from formal theoretical models (Friston, 2005). Thereof, especially prefrontal brain regions are known to be involved in visual search and the top-down control of visual perception from working memory by impacting on lower visual cortex (for review, see Soto et al., 2008).

Interactions between higher and lower brain areas, as assumed to be involved in visual search, can be well investigated by analysis of oscillatory brain activity. Interaction within or between brain areas is implemented by synchronous neuronal activity, as reflected by rhythmical oscillations of the field potential which can be recorded using scalp electroencephalography (EEG). Oscillatory EEG activity is commonly reported to play a functional role for perceptual

and cognitive processes (Buzsáki & Draguhn, 2004; Fell & Axmacher, 2011; Fries, 2005). Two

brain areas are assumed to be functionally coupled when their activity is more synchronous than

what would be expected from random fluctuations. It has been suggested that the complexity of

the neural network(s) involved will determine the frequency range of the dynamics in in a given

interaction (Buschman & Miller, 2007; Fell & Axmacher, 2011; Fries, 2005), such that long-

range interactions during top-down processes draw on lower frequencies in the theta band (~6

Hz) and alpha band (~10 Hz), whereas higher frequency, such as gamma band (>30 Hz),

interactions characterize more local, small-network interactions.

During tasks requiring a high level of top-down control, long-range interregional

synchronization between human prefrontal and parietal areas has repeatedly been found for

oscillatory brain activity in the theta or alpha frequency range, for example during highly

demanding working memory tasks (e.g. Sarnthein et al., 1998; Sauseng et al., 2005; von Stein et

al., 2000). Synchronous gamma band activity has been linked to bottom-up processes such as

feature-binding and awareness (for review, see Engel & Singer, 2001). In the context of attention

and working memory, however, gamma band activity has been associated with processing

demands related to object representations, directed attention and active maintenance or

manipulation of information (e.g. Axmacher et al., 2006; Friese et al., 2013; Jensen et al., 2007).

Thereby, the common underlying mechanism is a need for comparison of sensory input with

memory content as proposed in the Match and Utilization Model by Herrmann and colleagues

(2004). This is supposed to be realized through two computational steps that gamma-band

responses are involved in: Through bottom-up like matching of visual input to memory contents,

reflected by an early, evoked gamma band around 100-150ms after stimulus onset; and through

utilizing matched information for memory updating or the reallocation of attention, reflected by a

later, induced gamma band response around 300ms post-stimulus and jittered in phase. However,

it has been argued that this model would well account for the matching with long-term memory

information but less well for the matching with short-term memory information, i.e. mental

templates kept in working memory (Holz et al., 2010; Sauseng et al., 2015). The argument here is

that the neuronal signature of matching sensory input with working memory templates should not

be solely explicable by an evoked gamma response, but that a long-range fronto-parietal network

drawing on theta band oscillations is expected to be involved. Theta band activity has been

shown to generally have a strong influence on local cortical activity both in the human and

animal brain, namely by entraining neuronal spiking and fast oscillatory activity, such as gamma

band activity (Canolty et al., 2006; Fell & Axmacher, 2011; O'Keefe & Recce, 1993; Sirota et

al., 2008). From studies using EEG in humans, perceptual and working memory processes have

been associated with theta-gamma frequency interaction (Demiralp et al., 2007; Griesmayr et al.,

2010; Sauseng et al., 2008, 2009; Schack et al., 2002). Such cross-frequency coupling is

commonly taken as an indicator of an exchange of information between global and local neuronal

networks (see Sauseng & Klimesch, 2008), and recently, especially cross-frequency phase

synchronization has been proposed to integrate neuronal processing which is distributed into

neuronal assemblies and across frequency bands by connecting the most central network nodes

(Palva & Palva, 2018). This may be associated with consistent spike-time relationships between

the oscillating neuronal populations; and cross-frequency phase-synchronized input to pyramidal

layer 5 cells may facilitate neuronal bursting of these cells (Palva & Palva, 2018). So, concerning

the activation of mental templates from working memory and their comparison with sensory

input, cross-frequency phase synchronization between theta and gamma band oscillations can be

regarded as a candidate neuronal mechanism underlying this process (Sauseng et al., 2010, 2015).

Indeed, there is evidence in support of the involvement of a transient theta to gamma phase

synchronization in posterior parietal brain areas in integrating top-down controlled mental

templates with bottom-up visual processing (Holz et al., 2010; Sauseng et al., 2008). In cases

where our expectancies and the actual visual input match, a higher transient phase

synchronization than in case of a non-match has been found between posterior theta and gamma

oscillations. For example, in a delayed-match-to-sample working memory paradigm, Holz and

colleagues (2010) found stronger posterior EEG theta to gamma phase synchronization for

congruent in comparison to incongruent trials 150-200 ms post probe presentation. Additionally,

the authors reported a resetting of theta phase shortly before this, leading them to propose that a

posterior phase resetting of theta band oscillations could enable the transient cross-frequency

synchronization with high frequency activity in the gamma band range found shortly after. This is

supported by the idea that cross-frequency phase synchronization could be a candidate

mechanism for integrating cognitive functions, such as the representation of sensory information

and attentional or executive functions, by connecting the most central network nodes between

distributed neuronal networks that support these functions (Palva & Palva, 2018).

In the current study, we asked whether the number of activated mental templates that

could be matched with sensory input does influence memory matching in visual search, as

reflected by a transient cross-frequency interaction between theta and gamma frequencies.

Assuming that we sequentially keep in mind multiple templates (Lisman & Idiart, 1995; Olivers

et al., 2011; Van Vugt et al., 2014), memory matching should occur relatively early, a bit later or

even much later in a given trial, depending on whether the sequence's first, second, or a later

mental template could be matched to the current visual input. This means that in conditions

where multiple mental templates could be matched to one out of several possible targets

appearing on screen, the memory matching mechanism and likewise its resulting neuronal

correlate, is supposed to display more temporal variability across trials. Therefore, lower overall

theta to gamma phase synchronization values, which are measured through an index aggregated

over trials, are expected to be exhibited than when a single mental template enables a precise

matching and thus a temporally aligned theta to gamma phase synchronization.

We designed a visual search paradigm where displays with four abstract symbols were

shown to participants, each display containing one target among distractors. Participants had to

indicate in which quadrant of the display their target symbol had been presented. We varied the

number of mental templates that had to be kept in mind for successfully performing the visual

search. In separate experimental blocks, the target could be either one single symbol (i.e. one

item had to be held in memory) or one out of a set of three target symbols (i.e. three items would

have to be retained). In the single template condition, we expected that around 150-200 ms after

search array onset, a transient increase in theta-gamma phase synchronisation should arise over

right-hemispheric posterior brain areas for targets located in the contralateral hemifield, relative

to ipsilateral targets, because this would corroborate evidence from other task paradigms (Holz et

al., 2010; Sauseng et al., 2008, 2009). Conversely, such transient increase in phase

synchronisation should not arise in a condition where three mental templates were required for

successful search performance, because a larger number of mental templates that could

potentially be matched to visual input would modulate cross-frequency phase interactions,

resulting in a temporally more variable pattern of synchronization across trials.

Methods

Participants

Thirty-five typically developed volunteers participated in the experiment. All gave written informed consent prior to their participation and received financial compensation or course credits upon completion. Four participants had to be excluded from analysis because their percentages of correct responses were in the range of chance level, suggesting they were merely guessing, in at least one condition of interest. Two more participants were excluded based on too noisy EEG recordings. In the remaining sample that was included in the analyses (n=29), mean age was 24.7 years ($SD = 2.8$) and 7 participants were male. All but one were right-handed, as assessed by the Edinburgh Handedness Inventory (Oldfield, 1971) and all reported normal or corrected to normal vision. The study was approved by the local Ethics Review Board and conducted according to the Declaration of Helsinki.

Apparatus and Stimuli

Participants were seated in a comfortable chair in a dimly lit room and were wearing an EEG cap (Easycap®) for registering EEG signals. They had a standard computer keyboard placed on their lap. Their left and right index and middle fingers were placed on four buttons of the numbers block, namely buttons 1, 2, 4, and 5, which were marked by coloured stickers. Each button represented one of four quadrants of a visual search display. Stimuli were presented on a 22-inch Samsung S22C450 monitor with a resolution of 1280 x 1024 and a 75 Hz refresh rate, which was placed centrally and at a distance of 80 cm from an observer. Stimulus presentation

was controlled using Presentation 0.71 (Neurobehavioural Systems®), which was synchronized

with recording of the EEG signals in BrainVision Recorder 2.0.4 (BrainProducts®).

All stimuli were presented against black background, with a white fixation cross in the

centre of the screen (see figure 1A). As stimuli, 16 different abstract symbols were created. None

of these abstract symbols were known to the observers. Thus, rather than using semantic memory

contents, participants needed to retain a mental template of the visual target(s) in working

memory. Four stimuli were used as targets (1-4) and 12 other stimuli as distractors (A-L). To

counterbalance which target and distractor stimuli appeared in the single vs. triple template

conditions, four experimental versions were used (*Version I:*1 and A-F vs. 2, 3, 4 and G-L;

Version II: 2 and G-L vs. 1, 3, 4 and A-F; *Version III*: 3 and A-F vs. 1, 2, 4 and G-L ; *Version IV*:

4 and G-L vs. 1, 2, 3 and A-F). For each template condition, target and distractor stimuli were

composed into 48 different search displays. Thus, in the *single template condition*, 12 search

displays had the target symbol in the same quadrant, while three distractor symbols, randomly

drawn from a subset of 6, were placed into the remaining three quadrants. For the *triple template*

condition, a display could contain one out of three possible targets, thus each target was placed in

each quadrant four times, accompanied by three randomly drawn distractors. This resulted in a

total of 256 search displays being used. Mask screens displayed circular Gabor gratings at the

same four locations of the search items, consisting of 9 white and 10 black lines each and

oriented vertically (135°).

Figure 1. Illustration of the visual search task. A: Exemplary trial sequence of the visual search
task. Trials consisted of a fixation (600-1000ms), a search display (200ms), a mask screen
(1000ms), and a fixation (1500ms); Inter-trial intervals (ITI) showed a blank screen (800-
1200ms). Search displays contained one target among three distractors and participants indicated
in which quadrant of the search display the target had been presented by button press. B:
Experimental procedure. The experiment consisted of two parts with counterbalanced order
across the four experimental versions (shown here: version IV). Prior to each part, consisting of
four task blocks each, participants memorized and practiced the target(s).

Task and Procedure

We recorded EEG from participants while they completed a visual search task, where

they searched for a target stimulus among distractors. At the beginning of each trial (see figure

1A), a central fixation cross was presented for a random duration between 600 and 1000ms,

which participants were instructed to fixate during the whole trial. Next, the search display was

displayed for a duration of 200ms, and immediately masked for 1000ms. The fixation cross

remained on the screen for another 1500ms. The target stimulus was presented equally often in

each quadrant of the search display (25% of the trials). Participants indicated in which quadrant

of the search display the target stimulus had been presented by pressing the respective button on

the numbers block, for upper left (button 4) upper right (button 5), lower left (button 1) or lower

right (button 2). They were instructed to respond as accurately as possible, and as soon as

possible after presentation of the search display. So accuracy was emphasized over speed. As an

inter-trial interval, a blank screen was shown for a random duration of 800 to 1200ms, adding up

to a total trial duration of 4500ms, before the next trial started. Participants were instructed to

keep their eyes fixated at the central fixation cross during the whole task.

Our paradigm contained two types of task conditions in which the number of possible

targets, and thus the number of mental templates to be held in mind, could be either one or three

(*Template*; single vs. triple). These template conditions were presented in two separate

experimental parts (see figure 1B), each consisting of 192 trials, divided into four blocks with 48

trials. Across the four experimental versions the order in which template conditions were

presented was counterbalanced (Version I and III: triple, then single; Version II and IV: single,

then triple). Version was randomly assigned to a participant and determined which targets and

distractor sets were assigned to which condition (see apparatus and stimuli). Within a block,

search displays were presented in randomized order. In the triple template condition, targets

appeared equally often. Participants took breaks between blocks, resulting in a total of about 40

minutes to complete the experiment.

In order to familiarize the participants with the targets, a training was completed prior to

each condition (see figure 1B). During a memorization phase before each practice block, the

respective single target or three targets were shown to participants in a printed version, and

participants were asked to memorize them well. Practice blocks had fewer trials than the actual

experimental blocks and served to make participants familiar with their targets. The same

target(s) and distractors as in the respective template condition were displayed here, however,

only with one instead of four stimuli on screen. Upon target presentation, participants decided

whether the displayed stimulus was (one of) their target(s) or not. They did so with a button

press, where they were asked to press left (no target) and right (target) arrow buttons on the

keyboard. At least two memorization phases and practice blocks were completed per condition. If

necessary, more were administered, until participants were confident in discriminating between

target(s) and distractors and performed well above chance in doing so.

11   The whole experiment, including the preparation of the EEG cap, instructions, breaks,

12   training blocks and experimental blocks, took about 2 hours.

13   **EEG Data Acquisition and Preprocessing**

14   EEG was registered from 60 scalp locations with Ag-AgCl electrodes arranged according

to the extended 10-10-system in a TMS compatible electrode cap (Easycap®), using a BrainAmp

MRplus amplifier (BrainProducts®). Two electrodes were placed above and next to the left eye

for recording horizontal and vertical eye movements and blinks. An additional ring-electrode on

the tip of the nose was used as a recording reference and the ground electrode was placed at

electrode position FPz. Electrode impedances were kept below 15 kΩ. EEG data were digitized at

1000 Hz in a frequency range above 0.016 Hz. A notch filter was set at 50 Hz. Butterworth zero

phase filters were used.

22   EEG data were pre-processed using BrainVision Analyzer 2.0.4 (Brain Products®). Raw

data was re-referenced using an average reference of all EEG channels. After filtering with a low-

cutoff of 0.5 Hz (48 dB/oct) and a high-cutoff of 100 Hz (48 dB/oct), visual inspection was used

for excluding data sections with large artifacts during task breaks. Next, semiautomatic Ocular

Correction with Independent Component Analysis (Ocular Correction ICA) was applied to

correct for artifacts caused by eye blinks and eye movements. Only trials including a correct

response that was given within 3000 ms after search display onset were retained. Data were then

segmented into 2000ms epochs to avoid edge artifacts in later analysis steps, ranging from

1000ms before to 1000ms after onset of a search display. Finally, epochs that contained

remaining artifacts due to eye movements or muscle activity were rejected manually. On average,

the number of trials that remained after these procedures were 75.2 trials (78.3%) for targets on

the left side of the screen and 79.3 trials (82.6%) for targets on the right in the Single template

condition. In the Triple template condition, on average 54.0 trials (56.2%) remained for left

hemifield target positions and 55.2 trials (57.5%) for targets on the right.

Cross-frequency phase synchronization index

Source-space EEG signals obtained from the brain regions of interest (ROIs; see below

for details) were decomposed using continuous wavelet transformation using Morlet wavelets. In

order to extract one lower frequency band that is centred over the typical theta range and

comparable to the study by Holz and colleagues (2010), for several lower frequency bands,

wavelet coefficients were extracted with 5 frequency steps ranging from 1 Hz to 12 Hz, using a

5-cycle complex Morlet parameter. Thus, the frequency of interest for theta band activity had a

central frequency of 6.50 Hz (bandwith 5.2–7.80 Hz). Both the theta and gamma band activity

was derived using the same Morlet parameter, to be able to detect a transient modulation of phase

in the gamma frequency range. Activity in several higher frequency bands was extracted with 6

frequency steps ranging from 30 Hz to 80 Hz and with a 5-cycle complex Morlet parameter. The

purpose of this was to extract gamma bands that are comparable to the study by Holz and

colleagues (2010) and to cover the whole range from 30 Hz to 80 Hz, but with little overlap to

avoid redundancy and to reduce data for statistical analysis. Thus, three of these higher frequency

bands were extracted as frequencies of interest for gamma band activity which were centered

around 40 Hz (bandwith 32–48 Hz), 60 Hz (bandwith 48–72 Hz), and 70 Hz (bandwith 56–84

Hz).

Next, continuous phase values were extracted from the wavelet coefficients' complex

values, for lower and higher frequency bands. To quantify their phase consistency across trials,

we calculated the cross-frequency phase synchronization index (PSI), similar to Schack and

Weiss (2005) or Palva and colleagues (2005), through custom-made scripts in MATLAB

R2015b. So first, for each trial and sampling point, slow frequency band and high frequency band

phase values were multiplied with the central frequency of the other band, as described with the

equation $m \cdot f_n = n \cdot f_m$ (where f_n and f_m denote the sampling point-wise low and high frequency

phase values, respectively, and m and n are the central frequencies of the low and high frequency

bands). Next, the phase differences across these adjusted signals was calculated for each trial and

sampling point by subtracting sampling point-wise high frequency from low frequency adjusted

signals. PSIs across trials were calculated as the average vector length of these phase differences,

by taking the square root of the sum of the squared sine and cosine values of the phase

differences, averaged across trials, yielding an index ranging from 0 to 1, where 1 indicates

largest synchrony in phase. From these sampling point-wise PSIs, we then created averaged PSIs

for time windows of 50ms length, starting at stimulus onset up to 450 ms and an averaged PSI for

a pre-stimulus time window of 200 ms, starting 200ms pre-stimulus up to stimulus onset. These

were transformed using Rayleigh's Z ($rzPSI = n*PSI^2$). This was done to account for the

number of trials (n) that went into calculation of the index which were overall lower in the triple

template condition (only correct, artifact-free trials were entered into the index, see above), since

usually, measures of phase-synchronization are sensitive to the difference of trial numbers across

conditions used (Cohen, 2014). Note that this yields an index not ranging from 0 to 1, but ranging

from 0 to n, where larger values indicate larger synchrony in phase.

Regions of interest and time of interest

Brain regions of interest (ROIs) for analysis of posterior theta and gamma band activity

were identified in source space in order to attenuate effects of volume conduction and to reduce

multi-channel EEG data. We therefore transformed EEG data from scalp-level data into voxel-

based Low Resolution Electromagnetic Tomography (LORETA) data (Pascual-Marqui et al.,

2002) using BrainVision Analyzer 2.0.4 (Brain Products®). Here, a standard brain based on the

MNI-305 brain template and a 3-shell spherical head model is used, and the source space

comprises the cortical gray matter and hippocampus in the Talairach atlas with 2394 voxels at 7

mm spatial resolution. Based on the literature (Holz et al., 2010; as well as Sauseng et al., 2008),

we were interested to compare posterior theta and gamma activity in bilateral posterior ROIs.

While we do not assume that these are the only brain areas involved in this task, a source-specific

analysis in the study by Sauseng and colleagues (2008) showed strong effects of cross-frequency

phase synchronization in a similar task, with bilateral posterior sources located within extrastriate

areas, covering the left and right superior occipital gyrus for the majority of subjects. Thus, for all

further analyses in source space, we manually selected bilateral posterior ROIs in the left and

right superior occipital gyrus (see figure 2 for a visualization of the left and right superior

occipital gyrus, as implemented in the AAL atlas (Tzourio-Mazoyer et al., 2002)): a left superior

occipital ROI (centroid at MNI: -15, -95, 15) and the homologue right superior occipital ROI

(centroid at MNI: 15, -95, 15).

Within these ROIs, we analysed differences between our experimental conditions in the

time window of interest. The time window of interest (TOI) was 150-200 ms after visual search

display onset, a typical time window found in the previous studies (Holz et al., 2010; as well as

Sauseng et al., 2008).

6
7   **Figure 2.** Illustration of the selected regions of interest (ROIs) in source space. A left middle
8   occipital ROI (centroid at MNI -15, -95, 15) and the homologue right middle occipital ROI
9   (centroid at MNI 15, -95, 15) were selected for all further analyses in source space. Highlighted
10   in red are the left middle occipital gyrus and right middle occipital gyrus.

Event-related Potentials

To relate our data to the existing EEG research on visual search, in which the N2pc ERP

component has been described as an important neuronal signature (Eimer, 2014; Luck, 2012), we

computed scalp-level grand average ERP waveforms for left and right target locations in both

template conditions. These were averaged for posterior parietal electrodes contra- or ipsilateral

relative to target location (P2, P4, P6, P8, PO4, PO8, and O2; or P1, P3, P5, P7, PO3, PO7, and

O2) and baseline-corrected using a pre-stimulus time window of 200 ms, starting 200ms pre-

stimulus up to stimulus onset. Based on visual inspection (see figure 3), ERP waves began to

differ between contralateral and ipsilateral target presentations from 220 ms onwards, which is in

the N2pc latency range. The N2pc, which is consistently found in visual search tasks, is an an

ERP component exhibiting an enhanced negativity at posterior electrodes contralateral to target

presentation and is typically interpreted as an electrophysiological marker of attentional capture.

Figure 3. Event-related potentials (ERPs) averaged for posterior parietal electrodes, contra- or
ipsilateral relative to target location in both template conditions.

Behavioural data

As a measure of task performance, the percentage of correct responses was computed for

each subject and in both template conditions. Additionally, the median across reaction times from

trials with a correct response was calculated (note, however, that the task instructions had

emphasized accuracy over speed, so reaction times should be interpreted with this in mind).

Statistical methods

For both the behavioural and the EEG data, statistical analyses were carried out using statistical software R 3.6.1 (R Core Team, 2019) and for data visualization, plots were created using the package ggplot2 3.2.1 (Wickham, 2016). To compare behavioural data from the single template condition vs. triple template condition, two-tailed paired-samples t-tests were used on task accuracy and response times. For the analysis of EEG data, linear mixed effects models (LMMs) were implemented with the lme4 package 1.1.21 (Bates et al., 2007), contrasts matrices were derived using the hypr package 0.1.7 (Rabe et al., 2020; Schad et al., 2019) and model summary tables were produced using the lmerOut package 0.5 (Alday, 2018). As an advantage over traditional repeated-measures ANOVA, LMMs estimate the difference between conditions instead of only the significance of a difference between conditions (see Payne et al. (2015) and Alday et al. (2017) for an overview on LMMs, parameter estimation and model fitting, examples of their use for EEG data analyses and further literature recommendations on LMMs). The consistency of theta-gamma phase difference, as measured by the Rayleigh's z-transformed cross-frequency phase synchronization indices (rzPSIs), was analysed for condition differences separately for the left and right posterior ROI (see above for details on the ROIs). This was done because data stemmed from sources located separate hemispheres and because other studies from our group have previously found lateralized effects of cross-frequency synchronization (see Sauseng et al. (2009) or Holz et al. (2010)). While the whole time series of rzPSIs was inspected descriptively, data were analysed for condition differences exclusively in the time window of interest found in previous studies, 150-200 ms after target onset (see above for details on the TOI) to reduce data for statistical analysis. Here, based on our hypothesis, we were mainly interested in an interaction between template condition and target location or any higher-order interaction involving these two factors. Adding rzPSIs for theta-to-40 Hz, theta-to-60 Hz, theta-

to-70 Hz into the model enabled us to assess whether a broadband or rather frequency specific

theta-to-gamma band effect were involved. In separate analyses for each ROI, we used a LMM

where rzPSIs from that ROI were predicted by the fixed effects COND (Template condition:

single, triple), TARG (Target location: contralateral, ipsilateral), CFS (Cross-frequency

synchronization: theta-to-40 Hz, theta-to-60 Hz, theta-to-70 Hz), and their interactions. The

model included a single random-effects term for the intercept of the individual subjects SUBJ.

For LMM modeling, the categorical variables were encoded with sequential difference contrasts

(for 2-level predictors COND and TARG: (1/2, -1/2); for 3-level predictor CFS: (-2/3, 1/3, 1/3)

and (-1/3, -1/3, 2/3)). Thus, the intercept is estimated as the grand average across all conditions

and resulting fixed effect estimates can be interpreted as main effects. For the model summaries

we regarded contrast coefficients with absolute t values larger than 1.96 as indicative of a precise

estimate. T-values above 1.96 can be treated as approximating the two-tailed 5% significance

level since a t-distribution with a high degree of freedom approaches the z distribution (Baayen et

al., 2008). The reported models were fit based on restricted maximum likelihood estimation.

Data availability

Data needed to reproduce all reported findings are available from the authors upon

reasonable request trough access to our data repository (https://osf.io/h2j6d/).

Results

Behavioural analyses

Task accuracy (measured as the percentage of correct responses) was higher in the SINGLE template condition (M = 86.08%, SD = 14.51%) than in the TRIPLE template condition (M = 63.031%, SD = 11.56%) as indicated by a significant paired samples t-test (t(28) = 5.94, p <0.001, d = 1.76). Similarly, reaction times (computed as the median across correct trial's reaction times) in the SINGLE template condition (M = 698.03ms, SD = 276.14ms) were significantly faster than in the TRIPLE template condition (M = 888.07ms, SD = 322.22ms; t(28) = -3.87, p < 0.001, d = -0.63). Both these results indicate that behavioural task performance was better when participants had to search for one target among distractors than for one out of three possible targets.

EEG analyses

Theta-gamma phase synchronization in the right hemispheric ROI

Figure 5 shows single-subject rzPSIs and their group average from the right hemispheric ROI in the time window 150-200 ms after visual search display onset. A summary of model fit for rzPSIs from the right hemispheric ROI in the time window 150-200 ms after visual search display onset and the fixed effects COND (single, triple), TARG (contralateral, ipsilateral), CFS (theta-to-40 Hz, theta-to-60 Hz, theta-to-70 Hz), their interactions as well as the random effect for SUBJ can be seen in table S0 in the supplemental materials. A visualization of the fixed effects is provided in figure 4. The grand mean rzPSIs have an estimate of 0.8 as represented by the intercept. TARG has an effect (0.062, t = 2.5) indicating that targets located in the contralateral

hemifield elicited larger rzPSIs than targets at ipsilateral locations, but there also is an interaction

between COND and TARG (0.18, t = 3.6), indicating that this target-related difference is larger in

the single than in the triple template condition. Importantly, no other contrast involving the

interaction effect between COND and TARG exceeded the threshold of absolute t values larger

than 1.96. So this critical effect does not interact with gamma frequency dependent differences,

although three contrasts involving the factor CFS yield precise estimates with absolute t values

larger than 1.96: One contrast shows that rzPSIS for Theta-to-70Hz are smaller than for Theta-to-

60Hz (-0.068, t = -2.2), however, there also is an interaction with COND, reflecting that this

gamma frequency dependent difference in rzPSIs is smaller for single than triple template

conditions (0.15, t = 2.5). The interaction between the other CFS contrast and COND indicates

that the difference between rzPSIs for Theta-to-60Hz and for Theta-to-40Hz is larger for single

than triple template conditions (-0.14, t = -2.3).

For an illustration of the whole time-series for rzPSIs from the right hemispheric ROI,

figure 6 shows the descriptives of group average rzPSIs (averaged across theta-to-40 Hz, theta-to-

60 Hz or theta-to-70 Hz cross-frequency synchronization) from the right hemispheric ROI for

post-stimulus time windows of 50ms length and for a pre-stimulus baseline.

Figure 4. Visualization of fixed-effect estimates for model fit of the cross-frequency phase
synchronization indices (Rayleigh's z-transformed; rzPSIs), measuring the consistency of theta-
gamma phase difference, from the right hemispheric ROI in the time window 150-200 ms after
visual search display onset. The model includes a random-effects term for the intercept of
individual subjects and the fixed effects COND (single, triple), TARG (contralateral, ipsilateral),
CFS (Theta-to-40 Hz, Theta-to-60 Hz, Theta-to-70 Hz), and interactions between them. Note:
Dots represent values of the estimated coefficients and lines show their standard deviations.

Figure 5. Cross-frequency phase synchronization indices (Rayleigh's z-transformed; rzPSIs),

measuring the consistency of theta-gamma phase difference, from the right hemispheric ROI in

the time window 150-200 ms after visual search display onset. RzPSIs are displayed separately

for single or triple template conditions (in color), for contralateral or ipsilateral target locations

(on the x axis) and for theta-to-40 Hz, theta-to-60 Hz or theta-to-70 Hz cross-frequency

synchronization (in separate panels). Single-subject indices (as thin lines) are overlayed by group

averages (as thick lines).

Figure 6. Cross-frequency phase synchronization indices (Rayleigh's z-transformed; rzPSIs),

measuring the consistency of theta-gamma phase difference, from the right hemispheric posterior

ROI in windows of 50ms length, starting at stimulus onset 0ms up to 450 ms, and in a 200ms pre-

stimulus baseline time. Group averaged rzPSIs are shown separately for single or triple template

conditions (in color and in separate panels) and for contralateral or ipsilateral target locations (as

line-type). Indices are averaged across theta-to-40 Hz, theta-to-60 Hz or theta-to-70 Hz cross-

frequency synchronization.

Theta-gamma phase synchronization for the left hemispheric ROI

For rzPSIs from the left hemispheric ROI in the time window 150-200 ms after visual search display onset (see supplemental materials figure S1), a summary of model fit for the fixed effects COND (single, triple), TARG (contralateral, ipsilateral), CFS (theta-to-40 Hz, theta-to-60 Hz, theta-to-70 Hz), interactions between them, and a single random-effects term for the intercept of the individual subjects can be seen in the supplemental materials in table S1.

Unlike the results from the right hemispheric ROI, for rzPSIs from the left hemispheric ROI, no contrast exceeded the threshold of absolute t values larger than 1.96. The grand mean rzPSIs have an estimate of 0.81 as represented by the intercept. Note that the by subject variation beyond the variability induced by the residual error was estimated as zero, i.e. the random effects matrix was singular for this model. However, since dropping a by-subject random effect of zero will have no effect on the fixed effect estimates, it was kept in the model.

For an illustration of the whole time-series for rzPSIs from the left hemispheric ROI, supplemental materials S2 shows the group average rzPSIs (averaged across theta-to-40 Hz, theta-to-60 Hz or theta-to-70 Hz cross-frequency synchronization) from the left hemispheric ROI for post-stimulus time windows of 50ms length and for a pre-stimulus baseline.

Control analyses

We conducted several control analyses in order to investigate whether the critical interaction between COND and TARG that we found for theta-gamma cross-frequency synchronization in the right-hemispheric ROI is frequency-specific and to exclude a spurious effect relying on evoked responses. These were conducted only for the right-hemispheric ROI because the critical effect was exclusively found for this ROI.

First, to control for the possibility that cross-frequency synchronization is rather between

gamma and broadband lower frequencies in general than specifically between gamma and the

theta frequency range, the same analyses as in the main analysis were carried out, but for cross-

frequency phase synchronization indices between gamma frequencies and the alpha frequency

range. If alpha-gamma phase synchronization showed the same pattern of results as theta-gamma

cross-frequency synchronization, specifically the critical interaction from the main analysis, the

effect would not be frequency specific. Next, to control for the possibility of spurious effects of

theta-gamma phase synchronization due to simultaneous but unrelated evoked activity in

response to probe presentation in both theta and gamma frequency bands, we analysed spectral

amplitudes as well as the phase locking factor (PLF) for theta and gamma frequencies. If a

simultaneous increase in spectral amplitudes or a simultaneous phase resetting in response to

stimulus onset can be found at both theta and gamma frequencies, this could lead to artificial

cross-frequency phase synchronization despite the two frequencies not interacting with each

other. This would be the case if spectral amplitudes or rzPLFs for theta and gamma frequencies

showed the same pattern of results as the main analysis, specifically the critical interaction.

Alpha-gamma phase synchronization for the right hemispheric ROI

To investigate the frequency specificity of the observed interaction between COND and

TARG that we found for theta-gamma cross-frequency synchronization in the right-hemisheric

ROI in the main analysis, the same analyses were carried out for cross-frequency phase

synchronization between gamma frequencies and the alpha frequency range. Thus, after wavelet

coefficients were extracted with 5 frequency steps ranging from 1 Hz to 12 Hz, using a 5-cycle

complex Morlet parameter, the central frequency of interest for this control analysis was at 9.25

Hz (7.40-11.10 Hz) in order to obtain phase estimates from the alpha frequency range. All

following analysis steps were identical to the previously described steps for the main analysis of

theta-gamma phase synchronization (see methods section for details).

Contrary to the effects observed for the main analysis, in the control analysis for alpha-

gamma rzPSIs from the right hemispheric ROI, no contrast exceeded the threshold of absolute t

values larger than 1.96. The grand mean rzPSIs have an estimate of 0.8 as represented by the

intercept (see supplemental materials figure S3 & S4 for data visualisation and supplemental

materials table S2 for a summary of model fit).

Amplitudes and PLF for the right hemispheric ROI

To control for the possibility of spurious effects of theta-gamma phase synchronization

due to evoked activity in response to probe presentation, we further analysed spectral amplitudes

for theta and gamma frequencies. Spectral amplitudes were calculated as the wavelet coefficients'

real values which were then averaged across trials for the same frequencies of interest as in the

main analysis: For theta band activity, spectral amplitudes with a central frequency of 6.50 Hz

(bandwith 5.2–7.80 Hz) were extracted through a continuous wavelet transform (5 frequency

steps from 1 Hz to 12 Hz, 5-cycle complex Morlet parameter). For gamma band activity, three

bands which were centered around 40 Hz (bandwith 32–48 Hz), 60 Hz (bandwith 48–72 Hz), and

70 Hz (bandwith 56–84 Hz) were extracted from a continuous wavelet transform (6 frequency

steps from 30 Hz to 80 Hz, 5-cycle complex Morlet parameter. As for the main analysis, we

computed averages of amplitudes for time windows of 50ms length and for a pre-stimulus time

window of 200 ms. We then also calculated the phase-locking factor (PLF; transformed using

Rayleigh's Z; $rzPLF = n*PLF^2$) for theta and gamma frequencies separately. This was done to

analyse the inter-trial consistency of phase-locking relative to stimulus onset within both

frequency bins. For this, their phase values were extracted as in the main analysis (see methods

section for details). PLFs were then calculated as the average vector length of these phase values

(Bonnefond & Jensen, 2012; Tallon-Baudry et al., 1996), which were also averaged into time

windows and transformed using Rayleigh's Z as in the main analysis.

Thus, for demonstrating that the observed interaction between COND and TARG for

theta-gamma cross-frequency synchronization in the right-hemisheric ROI in the main analysis is

not an artefact from filtering an evoked response, the same analyses were carried out for theta

amplitudes and for gamma amplitudes as well as theta and gamma phase-locking factors from the

right hemispheric ROI. While the three gamma bands' spectral amplitudes as well as their phase-

locking factors were calculated separately, they were averaged before entering them into the

model because the critical effect in the main analysis did not interact with gamma frequency

dependent differences. So for gamma frequencies, both these control analyses were conducted for

the average of all three gamma bands.

In the control analysis for theta amplitudes, no contrast exceeded the threshold of absolute

t values larger than 1.96. The grand mean theta amplitudes have an estimate of 2.4 (see

supplemental materials figure S9 & S10 for data visualisation and supplemental materials table

S5 for a summary of model fit). Similarly, in the control analysis for gamma amplitudes, all

contrasts remained below the threshold of absolute t values of 1.96. Here, the intercept indicated

that the grand mean gamma amplitudes have an estimate of 3.2 (see supplemental materials

figure S11 & S12 for data visualisation and supplemental materials table S6 for a summary of

model fit). The visualisation of the whole time-series from the theta and gamma band amplitudes

from the right hemispheric ROI illustrates that there is no simultaneous increase in both bands in

response to stimulus onset. In contrast to theta amplitudes, gamma amplitudes did not show any

stimulus-locked increase.

In the control analysis for theta rzPLFs, the intercept indicated that the grand mean theta

rzPLFs have an estimate of 38. The effect of COND (12, t value = 5.8) indicated that theta

rzPLFs are larger in single than triple template conditions (see supplemental materials figure S5

& S6 for data visualisation and supplemental materials table S3 for a summary of model fit).

However, in the control analysis for gamma rzPLFs, no contrast exceeded the threshold of

absolute t values larger than 1.96. Here, the grand mean gamma rzPLFs have an estimate of 0.95,

as indicated by the intercept (see supplemental materials figure S7 & S8 for data visualisation and

supplemental materials table S4 for a summary of model fit). When comparing the illustration of

the whole time-series from the theta rzPLFs and for gamma rzPLFs from the right hemispheric

ROI, it can be seen that gamma rzPLFs did not show a stimulus-locked resetting of phase.

Similar to the amplitudes control analysis, there is no indication for a simultaneous increase of

phase-locking of both bands in response to stimulus onset.

Discussion

Cross-frequency synchronization between theta and gamma band EEG activity has been

proposed to serve matching processes between prediction and sensation in visual perception

(Sauseng et al., 2010, 2015). In this study, we investigated how these electrophysiological

correlates of memory matching are affected by the number of activated internal templates which

can be compared to incoming sensory information. To perform the visual search task of this

experiment, one has to hold in mind a template of a single or of multiple targets' visual properties

so that it can be matched with the incoming stimulus. We expected to find stronger transient theta

phase to gamma phase synchronization at posterior sites that are contralateral relative to target

location compared to ipsilateral targets around 150-200ms after search display presentation in the

single template condition, but less so in the triple template condition.

In line with this, we found stronger theta-to-gamma phase synchronization in this early

time window at a ROI in the right middle occipital gyrus, elicited by targets presented in the

contralateral hemifield than by ipsilateral targets. An increase in theta to gamma phase

synchronization contralateral relative to ipsilateral to target locations is well in line with what can

be expected due to the lateralized organization of the visual system. This difference between

contra- and ipsilateral target locations was larger in the single template condition than in the triple

template condition. Specifically, in the single template condition, theta to gamma phase

synchronization was higher for contralateral targets than for ipsilateral targets. This was not the

case in the triple template condition. Thus, our data lend support to our hypothesis that memory

matching was less precise in conditions where one out of three mental templates had to be

matched with a target, whereas single template conditions enabled efficient memory matching.

This evidence from the single template condition in the current visual search paradigm

corroborates previous findings showing a higher transient phase synchronization between

posterior theta and gamma activity in cases where our expectancies match the actual visual input

than in case of a non-match (Holz et al., 2010; Sauseng et al., 2008). Thus, our data from the

single template condition fit well into the proposed framework (Sauseng et al., 2010, 2015),

which proposes that cross-frequency phase synchronization between theta and gamma

frequencies early after target presentation can be regarded as a candidate neuronal mechanism

underlying the matching of mental templates from working memory with sensory input.

Conversely, the triple template condition does not reveal such dynamics early after target

presentation; possibly indicating that multiple templates are held sequentially in working

memory, and that in a given trial depending on whether the sequence's first, second, or third

mental template could be matched to the current visual input, memory matching occurred

relatively early, a bit later or even much later, leading to overall more temporal variability across

trials. This is then reflected in the cross-frequency phase synchronization mechanism investigated

here. Based on proposals for a limited capacity of human visual working memory, supposedly

around three to four items (Luck & Vogel, 2013), one might expect that likewise, we could

maintain up to three or four simultaneous search templates for visual search as well. However,

there is evidence that not all working memory items influence the guidance of selective attention,

but that only active memory items function as an attentional template and directly affect

perception, whereas accessory memory items have relatively little influence on visual selection

(Olivers et al., 2011). The conclusion here is that working memory items generally compete for

the status of 'attentional template', which can only be achieved by one item at a time. So, although

working memory could store multiple objects, observers could only actively look for one at a

time. Thus, multiple-template search should require switching between mental templates or

sequentially looking for them. Interestingly, though, Beck and colleagues (2012) propose that

observers can concurrently keep two templates active in simultaneous search because when they

explicitly asked participants to simultaneously search for two templates, their gaze frequently

switched between them without switch costs. However, evidence showing clear switch costs for

selection has been reported when only one out of two potential targets was available, suggesting

that observers cannot actively search for multiple objects if they are not able to freely choose the

target category (Ort et al., 2017, 2018). This is again well in line with the idea that only one

search template at a time has priority and will guide visual attention (Olivers et al., 2011) and Ort

and colleagues argue that both lines of evidence can be explained from a reactive versus

proactive cognitive control framework (Braver, 2012). In this framework, when multiple targets

are all available for search, participants can proactively prepare for any target, resulting in a lack

of switch costs. Conversely, when only one of multiple possible targets is present for search, the

currently displayed target might not match the target that the participant anticipated. Reactive

control would follow this conflict, leading to increased processing times.

This latter case is similar to the current study's triple template condition, where only one

of three possible targets was presented for search. While the design of our task does not allow for

the analysis of inter-trial switch costs, we do see significantly fewer correct responses as well as

behavioural slowing in response times to targets in the triple template blocks compared to targets

in single template blocks. Slower and less accurate search performance has also been reported

during simultaneous search for two targets compared to search for either target alone, indicating

that subjects can probably not perform two simultaneous matching processes (Huang & Pashler,

2007; Menneer et al., 2007). So most likely, in a given trial in the current study's triple template

blocks, a currently displayed target possibly did not match the target that the participant

anticipated. This means that across trials, the match between memory templates and visual input

would occur at varying points in time, which should also be reflected in the underlying neuronal

mechanism of memory matching. Conversely, in the single template condition, certainty about

the mental template that has to be matched with sensory input was high in each trial and enabled

a temporally precise matching process across trials. Our data support this interpretation well,

showing that theta to gamma phase synchronization, the proposed underlying mechanism of

memory matching, was higher for contralateral targets than for ipsilateral targets in the single

template condition, whereas this was not the case in the triple template condition. Note that as a

measure for theta-gamma cross-frequency phase synchrony, we analysed the consistency of

phase difference between the two frequencies over trials (Holz et al., 2010; Palva, 2005; Sauseng

et al., 2008). This measure does not require the two frequencies to be coupled continuously. High

estimates of phase synchrony will be achieved when there is a fixed relation between low and

high frequencies in each trial, independent of absolute phase difference between them and of

phase-locking to stimulus of either of them, whereas low estimates will be achieved when phase

relations vary over trials . To ensure that differences between conditions were not based on

merely different trial counts in the single and triple template condition, PSI values were

transformed using Rayleigh's z transform (Cohen, 2014). Naturally, this does not eliminate the

difference in signal-to-noise ratio between conditions, which was most likely lower in the triple

template condition. Note, however, that this pattern of results was found even though in both

conditions, only trials with a correct response for which we assume that successful memory

matching must have taken place at some point were used to calculate theta-gamma phase

synchronization indices.

To discuss the pattern of results in the triple template condition, one might want to

speculate about how the memory matching mechanism investigated here might rely on pre-

stimulus working memory retention mechanisms. Generally, cross-frequency interactions

between gamma band activity and slower brain waves have frequently been suggested to be

involved in multi-item working memory, such as for multi-item working memory retention. A

prominent computational model assumes that separate memory items are represented by single

gamma waves which are nested into a theta wave (Jensen & Lisman, 1998; Lisman & Idiart,

1995). It is hypothesized that with this mechanism, multiple memory items (gamma waves nested

into a theta cycle) can be actively held in parallel in working memory. In the light of this

framework, one would assume that before search display presentation, in the triple template

condition, the three mental templates would each be represented by separate gamma cycles

nested into a theta wave one after another. Thus, upon search display presentation, it would be

crucial as to whether the first, second or third item (gamma wave) incidentally matches with the

one on the search display, leading to a temporal variability in the range of two gamma cycles.

Another theoretical framework which entails cross-frequency synchronization between theta and

gamma as the neural basis for multi-item WM retention argues that during retention, each item is

coded by an entire gamma burst, i.e. multiple cycles, nested into a theta wave (Herman et al.,

2013; Van Vugt et al., 2014). Following these ideas, there would be a temporal variability of

memory matching in the triple template condition in the range of two theta cycles, depending on

whether visual input matches with the first, second or third item (gamma burst). In this study,

grand mean reaction time differences in the triple template condition were longer than reaction

times in the single template condition by about 190 ms, which is in the range of a theta cycle.

Speculatively, this would fit rather well with the predictions derived from the latter framework,

where due to the expected temporal variability of two theta cycles, average reaction times would

be expected to be around the length of one theta cycle longer when one out of three potential

targets can be matched with visual input. Note, however that our task instructions had not

emphasized a speeded but rather an accurate response, so we cannot draw strong conclusions

here. To examine these predictions more closely, studies with a more precise measurement of

response times would be required.

While we find a clearly right-lateralized effect, there are some previous studies reporting

bilateral effects of stronger theta-gamma phase synchronization; e.g. in a cued visual attention

task (Sauseng et al., 2008). However, in another study, Holz and colleagues (2010) found a right-

lateralized effect of CFS in a visual delayed-match-to-sample task. Similar effects showing a

lateralized theta-locked gamma phase synchronization for memory loads 3-4 in visual working

memory are also reported by Sauseng and colleagues (2009). Other studies have also found

primarily right-hemispheric brain areas to be relevant for visual search. For example, activity

enhancements were reported in bilateral superior parietal cortex, but only extending into the

intraparietal sulcus of the right hemisphere during visual search compared to overt orienting,

(Nobre et al., 2003) and activation was completely right-lateralized for monitoring functions in

visual search (Vallesi, 2014). More evidence for clear right-hemispheric dominance for search

organization comes from lesion-studies, for example that lesions in the right parietal, temporal

and occipital cortex were related to disorganized search (Ten Brink et al., 2016).

The right-hemispheric effects from our data seem to be specific for theta-gamma phase

synchronization since a control analysis for cross-frequency phase synchronization between alpha

and gamma did not show similar results. Yet, for alpha-gamma phase synchronization there was

an interaction between template condition and target location for the left hemispheric posterior

source. However, the direction of this effect (stronger ipsilateral PSI) was contrary to what would

be expected due to the lateralized organization of the visual system.

Previous studies have reported similar cross-frequency interaction either between theta

and gamma frequencies around 30-50Hz in a cued visual attention task (Sauseng et al., 2008) or

between theta and higher gamma activity (Holz et al., 2010; Sauseng et al., 2009). In the current

data from our visual search paradigm, it seems that the difference between right posterior rzPSIs

for Theta-to-70Hz and for Theta-to-60Hz is smaller for single than triple template conditions;

whereas the difference for Theta-to-60Hz and for Theta-to-40Hz seems to be larger for single

than triple template conditions. However, the critical effect between template condition and target

location in the main analysis did not interact with gamma frequency dependent differences. This

speaks rather for a broadband gamma effect than a selective effect of theta and a narrow gamma

sub-band in the current visual search paradigm. The critical interaction from the main analysis

seems to be rather frequency-specific to theta-gamma phase synchronization, however, because

contrary to the effects observed for the main analysis, all contrasts remained below threshold in a

control analysis with alpha-gamma phase synchronization.

Because spurious effects of theta-gamma phase synchronization might arise due to evoked

activity in response to probe presentation, we analysed amplitudes and the phase locking factor

for theta and gamma frequencies to control for this. If both frequencies showed a simultaneous

increase in amplitudes or a simultaneous phase resetting in response to stimulus onset, the data

might indicate artificial cross-frequency phase synchronization in the absence of true interactions

between the two frequencies. However, none of these control analyses showed a simultaneous

increase in amplitudes or a simultaneous phase resetting in response to stimulus onset nor a

similar pattern of results as the main analysis. Thus, the results from these control analyses rather

indicate it being implausible that the observed interaction between COND and TARG for theta-

gamma cross-frequency synchronization in the right-hemisheric ROI in the main analysis, is due

to an artefact from simultaneous evoked activity in response to probe presentation in both theta

and gamma frequency bands.

Taken together, our data lend support to the hypothesis that neuronal networks operating

at theta and gamma frequency do become more synchronized in phase during an early time

window following visual search display onset, when a single template has to be retained

compared to triple template conditions. This adds to previous theoretical accounts that have

strongly argued for a transient synchronization between theta and gamma phase over posterior

electrode sites as a neuronal correlate of matching of incoming sensory information with memory

contents from working memory (Sauseng et al., 2010, 2015). It also lends evidence to the idea of

sequential attentional templates (Lisman & Idiart, 1995; Olivers et al., 2011; Van Vugt et al.,

2014). For future studies, it would be interesting to investigate the temporal dynamics of such

matching processes during the acquisition and consolidation phase of attentional templates.

Studying more naturalistic contexts of template to input matching where, for example, templates

are acquired via learning, could further illuminate the involvement of cross frequency interactions

in template to input matching.

References

Alday, P. M. (2018). *lmerOut: LaTeX Output for Mixed Effects Models with lme4*.

    https://bitbucket.org/palday/lmerout

Alday, P. M., Schlesewsky, M., & Bornkessel-Schlesewsky, I. (2017). Electrophysiology Reveals

    the Neural Dynamics of Naturalistic Auditory Language Processing: Event-Related

    Potentials Reflect Continuous Model Updates. *ENeuro*, *4*(6).

    https://doi.org/10.1523/ENEURO.0311-16.2017

Axmacher, N., Mormann, F., Fernández, G., Elger, C. E., & Fell, J. (2006). Memory formation

    by neuronal synchronization. *Brain Research Reviews*, *52*(1), 170–182.

    https://doi.org/10.1016/j.brainresrev.2006.01.007

Baayen, R. H., Davidson, D. J., & Bates, D. M. (2008). Mixed-effects modeling with crossed

    random effects for subjects and items. *Journal of Memory and Language*, *59*(4), 390–412.

    https://doi.org/10.1016/j.jml.2007.12.005

Bates, D., Sarkar, D., Bates, M. D., & Matrix, L. (2007). The lme4 package. *R Package Version*,

    2(1), 74.

Beck, V. M., Hollingworth, A., & Luck, S. J. (2012). Simultaneous Control of Attention by

    Multiple Working Memory Representations. *Psychological Science*, *23*(8), 887–898.

    https://doi.org/10.1177/0956797612439068

Bonnefond, M., & Jensen, O. (2012). Alpha oscillations serve to protect working memory

    maintenance against anticipated distracters. *Current Biology*, *22*(20), 1969–1974.

    https://doi.org/10.1016/j.cub.2012.08.029

Braver, T. S. (2012). The variable nature of cognitive control: A dual mechanisms framework.

    Trends in Cognitive Sciences, *16*(2), 106–113. https://doi.org/10.1016/j.tics.2011.12.010

Bundesen, C. (1990). A theory of visual attention. *Psychological Review*, *97*(4), 523–547.

   https://doi.org/10.1037/0033-295X.97.4.523

Bundesen, C., Habekost, T., & Kyllingsbæk, S. (2005). A Neural Theory of Visual Attention:

   Bridging Cognition and Neurophysiology. *Psychological Review*, *112*(2), 291–328.

   https://doi.org/10.1037/0033-295X.112.2.291

Buschman, T. J., & Miller, E. K. (2007). Top-down versus bottom-up control of attention in the

   prefrontal and posterior parietal cortices. *Science (New York, N.Y.)*, *315*(5820), 1860–

   1862. https://doi.org/10.1126/science.1138071

Buzsáki, G., & Draguhn, A. (2004). Neuronal oscillations in cortical networks. *Science (New*

   York, N.Y.), *304*(5679), 1926–1929. https://doi.org/10.1126/science.1099745

Canolty, R. T., Edwards, E., Dalal, S. S., Soltani, M., Nagarajan, S. S., Kirsch, H. E., Berger, M.

   S., Barbaro, N. M., & Knight, R. T. (2006). High Gamma Power Is Phase-Locked to

   Theta Oscillations in Human Neocortex. *Science*, *313*(5793), 1626–1628.

   https://doi.org/10.1126/science.1128115

Cohen, M. X. (2014). *Analyzing Neural Time Series Data: Theory and Practice*. MIT Press.

Demiralp, T., Bayraktaroglu, Z., Lenz, D., Junge, S., Busch, N. A., Maess, B., Ergen, M., &

   Herrmann, C. S. (2007). Gamma amplitudes are coupled to theta phase in human EEG

   during visual perception. *International Journal of Psychophysiology*, *64*(1), 24–30.

   https://doi.org/10.1016/j.ijpsycho.2006.07.005

Desimone, R., & Duncan, J. (1995). Neural Mechanisms of Selective Visual Attention. *Annual*

   Review of Neuroscience, *18*(1), 193–222.

   https://doi.org/10.1146/annurev.ne.18.030195.001205

Duncan, J., & Humphreys, G. W. (1989). Visual search and stimulus similarity. *Psychological*

   Review, *96*(3), 433–458. https://doi.org/10.1037/0033-295X.96.3.433

Eimer, M. (2014). The neural basis of attentional control in visual search. *Trends in Cognitive*

Sciences, *18*(10), 526–535. https://doi.org/10.1016/j.tics.2014.05.005

Engel, A. K., & Singer, W. (2001). Temporal binding and the neural correlates of sensory

awareness. *Trends in Cognitive Sciences*, *5*(1), 16–25. https://doi.org/10.1016/S1364-

6613(00)01568-0

Everling, S., Tinsley, C. J., Gaffan, D., & Duncan, J. (2006). Selective representation of task-

relevant objects and locations in the monkey prefrontal cortex. *European Journal of*

Neuroscience, *23*(8), 2197–2214. https://doi.org/10.1111/j.1460-9568.2006.04736.x

Fell, J., & Axmacher, N. (2011). The role of phase synchronization in memory processes. *Nature*

Reviews Neuroscience, *12*(2), 105–118. https://doi.org/10.1038/nrn2979

Fries, P. (2005). A mechanism for cognitive dynamics: Neuronal communication through

neuronal coherence. *Trends in Cognitive Sciences*, *9*(10), 474–480.

https://doi.org/10.1016/j.tics.2005.08.011

Friese, U., Köster, M., Hassler, U., Martens, U., Trujillo-Barreto, N., & Gruber, T. (2013).

Successful memory encoding is associated with increased cross-frequency coupling

between frontal theta and posterior gamma oscillations in human scalp-recorded EEG.

NeuroImage, *66*, 642–647. https://doi.org/10.1016/j.neuroimage.2012.11.002

Friston, K. (2005). A theory of cortical responses. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, *360*(1456), 815–836.

https://doi.org/10.1098/rstb.2005.1622

Griesmayr, B., Gruber, W. R., Klimesch, W., & Sauseng, P. (2010). Human frontal midline theta

and its synchronization to gamma during a verbal delayed match to sample task.

Neurobiology of Learning and Memory, *93*(2), 208–215.

https://doi.org/10.1016/j.nlm.2009.09.013

Herman, P. A., Lundqvist, M., & Lansner, A. (2013). Nested theta to gamma oscillations and precise spatiotemporal firing during memory retrieval in a simulated attractor network. *Brain Research*, *1536*, 68–87. https://doi.org/10.1016/j.brainres.2013.08.002

Herrmann, C. S., Munk, M. H. J., & Engel, A. K. (2004). Cognitive functions of gamma-band activity: Memory match and utilization. *Trends in Cognitive Sciences*, *8*(8), 347–355. https://doi.org/10.1016/j.tics.2004.06.006

Holz, E. M., Glennon, M., Prendergast, K., & Sauseng, P. (2010). Theta-gamma phase synchronization during memory matching in visual working memory. *NeuroImage*, *52*(1), 326–335. https://doi.org/10.1016/j.neuroimage.2010.04.003

Huang, L., & Pashler, H. (2007). A Boolean map theory of visual attention. *Psychological Review*, *114*(3), 599–631. https://doi.org/10.1037/0033-295X.114.3.599

Iba, M., & Sawaguchi, T. (2002). Involvement of the Dorsolateral Prefrontal Cortex of Monkeys in Visuospatial Target Selection. *Journal of Neurophysiology*, *89*(1), 587–599. https://doi.org/10.1152/jn.00148.2002

Jensen, O., Kaiser, J., & Lachaux, J. P. (2007). Human gamma-frequency oscillations associated with attention and memory. *Trends in Neurosciences*, *30*(7), 317–324. https://doi.org/10.1016/j.tins.2007.05.001

Jensen, O., & Lisman, J. E. (1998). An oscillatory short-term memory buffer model can account for data on the Sternberg task. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *18*(24), 10688–10699.

Lisman, J. E., & Idiart, M. A. (1995). Storage of 7 +/- 2 short-term memories in oscillatory subcycles. *Science*, *267*(5203), 1512–1515. https://doi.org/10.1126/science.7878473

Luck, S. J. (2012). Electrophysiological correlates of the focusing of attention within complex visual scenes: N2pc and related ERP components. In S. J. Luck & E. S. Kappenmann

(Eds.), *The Oxford handbook of event-related potential components* (pp. 329–360).

Oxford University Press. https://doi.org/10.1093/oxfordhb/9780195374148.013.0161

Luck, S. J., & Vogel, E. K. (2013). Visual working memory capacity: From psychophysics and

neurobiology to individual differences. *Trends in Cognitive Sciences*, *17*(8), 391–400.

https://doi.org/10.1016/j.tics.2013.06.006

Menneer, T., Barrett, D. J. K., Phillips, L., Donnelly, N., & Cave, K. R. (2007). Costs in

searching for two targets: Dividing search across target types could improve airport

security screening. *Applied Cognitive Psychology*, *21*(7), 915–932.

https://doi.org/10.1002/acp.1305

Nobre, A. C., Coull, J. T., Walsh, V., & Frith, C. D. (2003). Brain activations during visual

search: Contributions of search efficiency versus feature binding. *NeuroImage*, *18*(1), 91–

103. https://doi.org/10.1006/nimg.2002.1329

O'Keefe, J., & Recce, M. L. (1993). Phase relationship between hippocampal place units and the

EEG theta rhythm. *Hippocampus*, *3*(3), 317–330. https://doi.org/10.1002/hipo.450030307

Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory.

Neuropsychologia, *9*(1), 97–113. https://doi.org/10.1016/0028-3932(71)90067-4

Olivers, C. N. L., Peters, J., Houtkamp, R., & Roelfsema, P. R. (2011). Different states in visual

working memory: When it guides attention and when it does not. *Trends in Cognitive

Sciences*, *15*(7), 327–334. https://doi.org/10.1016/j.tics.2011.05.004

Ort, E., Fahrenfort, J. J., & Olivers, C. N. L. (2017). Lack of Free Choice Reveals the Cost of

Having to Search for More Than One Object. *Psychological Science*, *28*(8), 1137–1147.

https://doi.org/10.1177/0956797617705667

Ort, E., Fahrenfort, J. J., & Olivers, C. N. L. (2018). Lack of free choice reveals the cost of
multiple-target search within and across feature dimensions. *Attention, Perception &
Psychophysics*, *80*(8), 1904–1917. https://doi.org/10.3758/s13414-018-1579-7

Palva, J. M. (2005). Phase Synchrony among Neuronal Oscillations in the Human Cortex.
Journal of Neuroscience, *25*(15), 3962–3972. https://doi.org/10.1523/JNEUROSCI.4250-
04.2005

Palva, J. M., & Palva, S. (2018). Functional integration across oscillation frequencies by cross-
frequency phase synchronization. *The European Journal of Neuroscience*, *48*(7), 2399–
2406. https://doi.org/10.1111/ejn.13767

Palva, J. Matias, Palva, S., & Kaila, K. (2005). Phase synchrony among neuronal oscillations in
the human cortex. *The Journal of Neuroscience: The Official Journal of the Society for
Neuroscience*, *25*(15), 3962–3972. https://doi.org/10.1523/JNEUROSCI.4250-04.2005

Pascual-Marqui, R. D., Esslen, M., Kochi, K., & Lehmann, D. (2002). Functional imaging with
low-resolution brain electromagnetic tomography (LORETA): A review. *Methods and
Findings in Experimental and Clinical Pharmacology*, *24 Suppl C*, 91–95.

Payne, B. R., Lee, C.-L., & Federmeier, K. D. (2015). Revisiting the incremental effects of
context on word processing: Evidence from single-word event-related brain potentials.
Psychophysiology, *52*(11), 1456–1469. https://doi.org/10.1111/psyp.12515

R Core Team. (2019). A language and environment for statistical computing. *R Foundation for
Statistical Computing, Vienna, Austria.* URL https://www.R-project.org/.

Rabe, M. M., Vasishth, S., Hohenstein, S., Kliegl, R., & Schad, D. J. (2020). hypr: An R package
for hypothesis-driven contrast coding. *The Journal of Open Source Software*.
https://doi.org/10.21105/joss.02134

Rossi, A. F., Bichot, N. P., Desimone, R., & Ungerleider, L. G. (2007). Top Down Attentional

Deficits in Macaques with Lesions of Lateral Prefrontal Cortex. *Journal of Neuroscience*,

27(42), 11306–11314. https://doi.org/10.1523/JNEUROSCI.2939-07.2007

Sarnthein, J., Petsche, H., Rappelsberger, P., Shaw, G. L., & von Stein, A. (1998).

Synchronization between prefrontal and posterior association cortex during human

working memory. *Proceedings of the National Academy of Sciences of the United States*

of America, *95*(12), 7092–7096. https://doi.org/10.1073/pnas.95.12.7092

Sauseng, P., Conci, M., Wild, B., & Geyer, T. (2015). Predictive coding in visual search as

revealed by cross-frequency EEG phase synchronization. *Frontiers in Psychology*, *6*,

1655. https://doi.org/10.3389/fpsyg.2015.01655

Sauseng, P., Griesmayr, B., Freunberger, R., & Klimesch, W. (2010). Control mechanisms in

working memory: A possible function of EEG theta oscillations. *Neuroscience and*

Biobehavioral Reviews, *34*(7), 1015–1022.

https://doi.org/10.1016/j.neubiorev.2009.12.006

Sauseng, P., & Klimesch, W. (2008). What does phase information of oscillatory brain activity

tell us about cognitive processes? *Neuroscience and Biobehavioral Reviews*, *32*(5), 1001–

1013. https://doi.org/10.1016/j.neubiorev.2008.03.014

Sauseng, P., Klimesch, W., Gruber, W. R., & Birbaumer, N. (2008). Cross-frequency phase

synchronization: A brain mechanism of memory matching and attention. *NeuroImage*,

40(1), 308–317. https://doi.org/10.1016/j.neuroimage.2007.11.032

Sauseng, P., Klimesch, W., Heise, K. F., Gruber, W. R., Holz, E., Karim, A. A., Glennon, M.,

Gerloff, C., Birbaumer, N., & Hummel, F. C. (2009). Brain oscillatory substrates of visual

short-term memory capacity. *Current Biology: CB*, *19*(21), 1846–1852.

https://doi.org/10.1016/j.cub.2009.08.062

Sauseng, P., Klimesch, W., Schabus, M., & Doppelmayr, M. (2005). Fronto-parietal EEG coherence in theta and upper alpha reflect central executive functions of working memory. *International Journal of Psychophysiology: Official Journal of the International Organization of Psychophysiology*, *57*(2), 97–103. https://doi.org/10.1016/j.ijpsycho.2005.03.018

Schack, B., Vath, N., Petsche, H., Geissler, H. G., & Möller, E. (2002). Phase-coupling of theta-gamma EEG rhythms during short-term memory processing. *International Journal of Psychophysiology*, *44*(2), 143–163. https://doi.org/10.1016/S0167-8760(01)00199-4

Schack, Baerbel, & Weiss, S. (2005). Quantification of phase synchronization phenomena and their importance for verbal memory processes. *Biological Cybernetics*, *92*(4), 275–287. https://doi.org/10.1007/s00422-005-0555-1

Schad, D. J., Vasishth, S., Hohenstein, S., & Kliegl, R. (2019). How to capitalize on a priori contrasts in linear (mixed) models: A tutorial. *Journal of Memory and Language*, *110*. https://doi.org/10.1016/j.jml.2019.104038

Sirota, A., Montgomery, S., Fujisawa, S., Isomura, Y., Zugaro, M., & Buzsáki, G. (2008). Entrainment of Neocortical Neurons and Gamma Oscillations by the Hippocampal Theta Rhythm. *Neuron*, *60*(4), 683–697. https://doi.org/10.1016/j.neuron.2008.09.014

Soto, D., Hodsoll, J., Rotshtein, P., & Humphreys, G. W. (2008). Automatic guidance of attention from working memory. *Trends in Cognitive Sciences*, *12*(9), 342–348. https://doi.org/10.1016/j.tics.2008.05.007

Soto, D., Humphreys, G. W., & Heinke, D. (2006). Dividing the mind: The necessary role of the frontal lobes in separating memory from search. *Neuropsychologia*, *44*(8), 1282–1289. https://doi.org/10.1016/j.neuropsychologia.2006.01.029

Soto, D., Humphreys, G. W., & Rotshtein, P. (2007). Dissociating the neural mechanisms of memory-based guidance of visual selection. *Proceedings of the National Academy of Sciences*, *104*(43), 17186–17191. https://doi.org/10.1073/pnas.0703706104

Spaak, E., Fonken, Y., Jensen, O., & de Lange, F. P. (2015). The Neural Mechanisms of Prediction in Visual Search. *Cerebral Cortex*, bhv210. https://doi.org/10.1093/cercor/bhv210

Tallon-Baudry, C., Bertrand, O., Delpuech, C., & Pernier, J. (1996). Stimulus Specificity of Phase-Locked and Non-Phase-Locked 40 Hz Visual Responses in Human. *Journal of Neuroscience*, *16*(13), 4240–4249. https://doi.org/10.1523/JNEUROSCI.16-13-04240.1996

Ten Brink, A. F., Biesbroek, J. M., Kuijf, H. J., Van der Stigchel, S., Oort, Q., Visser-Meily, J. M. A., & Nijboer, T. C. W. (2016). The right hemisphere is dominant in organization of visual search-A study in stroke patients. *Behavioural Brain Research*, *304*, 71–79. https://doi.org/10.1016/j.bbr.2016.02.004

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., Mazoyer, B., & Joliot, M. (2002). Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *NeuroImage*, *15*(1), 273–289. https://doi.org/10.1006/nimg.2001.0978

Vallesi, A. (2014). Monitoring mechanisms in visual search: An fMRI study. *Brain Research*, *1579*, 65–73. https://doi.org/10.1016/j.brainres.2014.07.018

Van Vugt, M. K., Chakravarthi, R., & Lachaux, J.-P. (2014). For whom the bell tolls: Periodic reactivation of sensory cortex in the gamma band as a substrate of visual working memory maintenance. *Frontiers in Human Neuroscience*, *8*. https://doi.org/10.3389/fnhum.2014.00696

von Stein, A., Chiang, C., & König, P. (2000). Top-down processing mediated by interareal

synchronization. *Proceedings of the National Academy of Sciences of the United States of*

America, *97*(26), 14748–14753. https://doi.org/10.1073/pnas.97.26.14748

Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer.

Yago, E., Duarte, A., Wong, T., Barcelo, F., & Knight, R. T. (2004). Temporal kinetics of

prefrontal modulation of the extrastriate cortex during visual attention. *Cognitive,*

Affective, & Behavioral Neuroscience, *4*(4), 609–617.

https://doi.org/10.3758/CABN.4.4.609

Contributions:

A.L.B., T.M. and P.S. conceived the experiment. A.L.B. programmed the experimental paradigm, recruited participants, conducted the experiment, and analyzed the data. A.L.B. wrote the manuscript, supervised by P.S.. T.M. and P.S. critically revised the manuscript.

Conflict of interest:

The authors declare no competing interests.

Achnowledgements:

This work was funded by Deutsche Forschungsgemeinschaft (DFG) SA 1872/2-1. We thank Eva Margraf and Patricia Christian for their support with data collection.