Population Genomic Analyses Separate Historic Translocations from Contemporary Gene Flow in Arkansas White-tailed Deer (*Odocoileus virginianus*)

Tyler K. Chafin[1] | Zachery D. Zbinden[1] | Marlis R. Douglas[1] | Bradley T. Martin[1] | Christopher R. Middaugh[2] | M. Cory Gray[2] | Jennifer R. Ballard[2] | Michael E. Douglas[1]

[1]Department of Biological Sciences, University of Arkansas, Fayetteville, Arkansas 72701, USA; Email: (TKC) tkchafin@uark.edu; (BTM) btm002@uark.edu; (ZDZ) zdzbinde@uark.edu; (MED) med1@uark.edu; (MRD) mrd1@uark.edu

[2]Arkansas Game and Fish Commission, Research, Evaluation, and Compliance Division, Little Rock, Arkansas 72205, USA; Email: (CRM) christopher.middaugh@agfc.ar.gov; (JRB) jennifer.ballard@agfc.ar.gov; (MCG) cory.gray@agfc.ar.gov

Running Title: Population genomics of white-tailed deer

Keywords: Gene flow; Management; Population connectivity; ddRADseq; SNPs; Population structure; Historical population dynamics; Genetic drift; Wildlife

24 **ABSTRACT**

Approximately 100 years ago, hunter-harvest eliminated white-tailed deer (WTD; *Odocoileus*

26 *virginianus*) in eastern North America, which subsequently served as a catalyst for wildlife

management as a national priority. An extensive stock-replenishment effort soon followed, with

28 WTD broadly translocated among states as a means of reestablishment. Now, contemporary

issues focus on reverberations from a global (and fatal) epizootic disease in Cervidae (chronic

30 wasting disease, CWD). These cumulative impacts have effectively obscured the traditional

signals of post-translocation gene flow and dispersal in North American WTD. To develop

32 baseline data for its adjudication, we applied cutting-edge molecular and biogeographic tools to

process 1,143 WTD sampled state-wide in AR, with 54,102 single nucleotide polymorphisms

34 (SNPs) derived via reduced-representation genomic sequencing. We then employed Simpson's

diversity index to visualize landscape genetic patterns previously obscured by extensive

36 translocations, and this allowed us to summarize multidimensional ancestry assignments and

identify spatio-genetic transitions. We then sub-sampled transects and tested clinal patterns

38 across loci for concordance and/or coincidence. Two salient results emerged: (A) Genetic echoes

from historic translocations are widely apparent; and (B) Geographic filters (major rivers; urban

40 centers; highways) now act as inflection points for the distribution of more contemporary

ancestry. These results, in synergy, yielded a state-wide assessment of how historic

42 translocations, as well as ongoing processes, have acted to dictate contemporary population

structure of Arkansas WTD. In addition, the analytical framework employed herein effectively

44 deciphered the extant/historic drivers of WTD distribution in AR. It is also applicable for other

biodiversity elements with demographic histories equally as complex.

1 | INTRODUCTION

48   Distribution and abundance are key natural history attributes that provide a foundation for

organismal ecology (Andrewartha & Birch, 1956; Kissling et al., 2018). Each is geographically

50   and hierarchically structured, often across several spatial scales, and in response to a non-random

distribution of resources and refugia (Turner, 1989). Species distributions and abundances are

52   increasingly impacted by habitat fragmentation, over-exploitation, and translocation in the

Anthropocene (Baker et al., 2017; Corlett, 2015), with the functional integrity of these attributes

54   being severely challenged. Hunter-harvest, an additional anthropogenic impact, also exerts a

relatively consistent global impact on biodiversity (Darimont et al., 2009). Given the manner by

56   which abundances and distributions have been so impacted, other attributes such as dispersal and

population connectivity have subsequently emerged as focal points for contemporary wildlife

58   management, (Crooks & Sanjayan, 2006; Cushman, Elliot, Macdonald, & Loveridge, 2016).

Here we focus on these latter attributes as we evaluate historic and contemporary population

60   structure in a wild Arkansas (AR) ungulate (white-tailed deer, WTD; *Odocoileus virginianus*).

62   **1.1 | Dispersal, connectivity, and population structure**

Dispersal and connectivity have been traditionally assayed for management purposes using direct

64   methods such as radio- and satellite-tracking (Kays, Crofoot, Jetz, & Wikelski, 2015), yet both

have deficiencies such as low sample-sizes and a perspective that must extrapolate from

66   individuals to populations (Katzner & Arlettaz, 2020). An indirect approach, on the other hand,

relies instead upon gene flow (i.e., movements, demographics) and genetic drift (i.e.,

68   heterozygosity, effective population size) to quantify population-level movement ecology

captured over many generations (Bossart & Prowell, 1998; Comte & Olden, 2018). Landscape

70	genetics, for example (Richardson, Brady, Wang, & Spear, 2016), quantifies movement patterns

by evaluating the spatial structure inherent to genetic diversity (as mediated by dispersal and

72	gene flow), and does so for greater numbers at a fraction of the per-individual cost (Picard et al.,

2017; van Rees, Reed, Wilson, Underwood, & Sonsthagen, 2018; Wang & Shaffer, 2017).

74	We now recognize that genotypes within a species (i.e. gene pool) vary across

landscapes, often coalescing as discretely recognizable sub-populations (i.e., intra-specific

76	population structure driven by mutation, selection, gene flow, genetic drift), and visualized/

interpreted at varying degrees both temporally and spatially (Barton & Clark, 1990; Slatkin,

78	1989). They span from the afore-mentioned sub-populations through continuously decreasing

gradients of genetic similarity that extend over geographic or environmental distance (i.e.,

80	isolation by distance: Avise 1992; Bohonak 1999; Bradburd et al. 2018). Structure can be

ascertained by sampling individuals across appropriate spatial scales, relative to the dispersal

82	kernel of the organism (Nathan, Klein, Robledo-Arnuncio, & Revilla, 2012). It can then be

employed to test hypotheses about the likely causes of observed genetic patterns (Bradburd,

84	Ralph, & Coop, 2016; Miles, Rivkin, Johnson, Munshi-South, & Verrelli, 2019). However, gene

flow and genetic drift are often conflated, and their deconstruction is an important aspect when

86	attempting to clarify how population genetic structure is driven.

Patterns of genetic diversity are also impacted by multiple extrinsic and intrinsic factors.

88	For WTD, those extrinsic may include: Rivers, interstate highways, large tracks of forested

habitat (Robinson, Samuel, Lopez, & Shelton, 2012); agricultural land use, climate-related

90	factors (Brinkman, Deperno, Jenks, Haroldson, & Osborn, 2005); and landscape connectivity

(Koen, Tosa, Nielsen, & Schauber, 2017). Intrinsic factors, on the other hand, may reflect:

92	Population density (Lutz, Diefenbach, & Rosenberry, 2015); age structure/ sex ratio (Long,

Diefenbach, Rosenberry, & Wallingford, 2008); and social hierarchy (Nixon & Mankin, 2016).

Factors that traditionally shape dispersal and genetic diversity within/ across regions can difficult to quantify in that anthropogenic actions such as translocation or trafficking can obscure

existing patterns (Brown, Hull, Updike, Fain, & Ernest, 2009; Shephard, Ogden, Tryjanowski, Olsson, & Galbusera, 2013). This is particularly true for WTD, as it is one of the most

recreationally important animal species in North America (Knoche & Lupi, 2012), with populations intensely impacted by both hunter harvest and game management (Waller &

Alverson, 1997; Wolverton, Kennedy, & Cornelius, 2007). For example, at the beginning of the 20^{th} century, WTD and other heavily hunted species (e.g., wild turkey) saw heavy declines and

widespread extirpations. In a census of the north-central United States, Leopold (1931) noted "… some deer" were still extant in southern Missouri at this time, and this dovetails with WTD

numbers estimated in Arkansas (~500 scattered across isolated refugia; Holder, 1951). Subsequent AR efforts to bolster resident numbers and re-populate depleted areas involved

translocations within-state, as well as importations from out-of-state. Although successful, a result of these efforts is that apparent genetic patterns are likely to be strongly reflective of

artificial, rather than natural, movements.

A first step in designating factors that have shaped WTD population structure and

110   dispersal is to estimate that genetic component that remains within AR WTD genomes as a residual of historic translocations and subsequent genetic drift. Perhaps the most pressing need to

112   understand WTD dispersal is the impact these data have with regard to the potential containment and mitigation of a widespread and fatal neurodegenerative disease of cervids (chronic wasting

114   disease; CWD), that has now become a panzootic (Williams and Young 1980; Escobar et al. 2019; Mawdsley 2020). Thus, efforts to contain and mitigate its spread are paramount for

116 wildlife management, not only in North America but also globally (Leiss et al., 2017).

118 **1.2 | Dispersal, connectivity, and management**

An understanding of the manner by which WTD disperse across the landscape has clear

120 management implications. Its landscape movements are many and varied, as driven by habitat

quality, climate, and anthropogenic benefits. Region-specific patterns are needed so as to

122 develop broader generalizations regarding species-specific movement ecology (Brinkman et al.,

2005).

124   The goal of our study was to ascertain if signatures of historic translocations are apparent

in, and have contributed to, the genetic diversity and structure of Arkansas WTD, and

126 furthermore, if they can indeed be parsed from ongoing gene flow and genetic drift. To do so, we

surveyed a broad array of nuclear genomic markers using ddRAD sequencing (Peterson et al.,

128 2012). Our first objective was to characterize respective patterns of anthropogenic translocation

and natural re-colonization via refugial populations, and secondarily to seek evidence for

130 geographic barriers throughout the state that seemingly prevent or actively filter WTD dispersal.

However, we hypothesized that the former situation would obscure the latter, in that long-range

132 anthropogenically-mediated displacement of individuals violates methodological assumptions

that stipulate gene flow as occurring in a spatially consistent manner (Bradburd & Ralph, 2019).

134   To address this, we repurposed existing formulas that summarized ancestry assignments

by reducing their dimensionality, and mitigated methodological artefacts due to translocation

136 (Simpson, 1949). Our result was an interpolated 'surface' representing putative intraspecific

suture zones. We also wished to define the respective roles played by stochastic and

138 deterministic processes in generating these zones, particularly given historic population

fluctuations and the internal/ external translocations that conflated the demography of AR WTD.

140   To do so, we borrowed components of clinal variation theory so as to hypothesize the manner by

which individual loci should transition across these spaces, relative to a genome-wide average

142   (Barton, 1983; Barton & Hewitt, 1985; Endler, 1973; Hewitt, 2001; Polechová & Barton, 2011;

Slatkin, 1973).

144   Although clinal variation is often taken as evidence for selection, genetic drift in concert

with spatially variable gene flow can generate patterns in individual loci that mimic those seen in

146   adaptive clines (Vasemägi, 2006). With drift operating alone, allele frequencies at each locus

may wander 'upward' or 'downward' across space as a function of the initial allele frequency.

148   Range expansion or dissemination from refugia can also yield a type of 'rolling' founder effect

that can mirror a clinal pattern (Excoffier, Foll, & Petit, 2009; Hallatschek & Nelson, 2008;

150   Hewitt, 2000; Keller et al., 2013). Given this, we might predict that multi-locus patterns at

intraspecific suture zones generated solely as a result of genetic drift in expanding populations

152   will be dominated by stochastic directional change, with each drifting either up or down with

respect to the genome-wide average (Santangelo, Johnson, & Ness, 2018). On the other hand, a

154   strong barrier to movement would yield a discontinuity in the rate of this change across space

(Barton, 2008; Nagylaki, 1976; Slatkin, 1973). The existence of a barrier (either impacting

156   dispersal or as a strong fitness differential) thus implies the presence of an inflection point that

would induce a sigmoidal change in locus-specific ancestries (the "width" of this cline is the

158   inverse of the slope at the inflection point; Slatkin 1973; Endler 1977; Fitzpatrick 2013).

        Given these predictions, we summarized locus-wise patterns using two clinal parameters:

160   α, which describes the direction of genetic change (i.e. directionally), and β, which indicates the

rate of change, or 'width' of a cline. We examined the manner by which these two parameters

162   varied, then developed hypotheses regarding how those processes impacted our observed

population structure. We then considered the evolutionary implications, and limitations, of these

164   results in the context of WTD management.

166   **2 | METHODS**

2.1 | Sampling and data collection

168   During 2016-2019, 1,720 tissues were collected by the Arkansas Game and Fish Commission

(AGFC), representing all 75 Arkansas counties. A combination of targeted sampling, road-kill

170   surveys, and a voluntary state-wide CWD testing program were deployed (see Douglas et al.,

2020). Age and sex were collected where possible, with the former estimated by tooth

172   development and wear (Severinghaus, 1949). Data for an additional 30 samples were also

attained from Wisconsin, as a means to test for signals of historically recorded translocation

174   efforts involving Wisconsin stock (Holder, 1951). From these, a subset of 1,208 samples were

chosen for sequencing.

176   We homogenized tongue or ear tissue (stored at -20°C) and extracted genomic DNA

using QIAamp Fast Tissue kits (Qiagen, Inc), with verification via gel electrophoresis (2%

178   agarose). Samples with sufficient yields of high molecular weight DNA (>200ng minimum)

were then enzymatically fragmented via incubation at 37°C, using high-fidelity *NsiI* and *MspI*

180   restriction enzymes (New England Biolabs, Inc.), following enzyme and size selection

optimization using *in silico* digests (Chafin, Martin, Mussmann, Douglas, & Douglas, 2018) of

182   several available reference genomes hosted by NCBI: *Odocoileus virginianus*

(GCA_002102435.1), *Capreolus capreolus* (GCA_000751575.1), and *Capra hircus*

184   (GCF_001704415.1).

Digests were purified using Ampure XP beads (Beckman-Coulter, Inc.) and standardized

186   to 100ng per sample. Unique inline barcodes (Peterson et al., 2012) were then ligated using T4

DNA Ligase (following manufacturer protocols; New England Biolabs, Inc.). Samples were then

188   multiplexed (N=48) prior to automated size selection at 300-450bp using a Pippin Prep (Sage

Sciences). Adapter extension was performed over 12 PCR cycles using TruSeq-compatible

190   indexed primers (Illumina, Inc.) and Phusion high-fidelity *taq* polymerase (New England

Biolabs, Inc.) Additional quality controls (e.g., qPCR and fragment analysis) were performed on

192   final libraries prior to 1x100 single-end sequencing on the Illumina HiSeq 4000 (Genomics and

Cell Characterization Facility, University of Oregon/Eugene), with a total of N=96 samples

194   pooled per lane.

Raw reads were demultiplexed using the pyRAD pipeline (Eaton, 2014), and those with

196   barcode mismatches were discarded. Demultiplexed reads were further filtered by removing

those having >4 nucleotides below a quality threshold of 99% accuracy. Reads were then

198   clustered into putative loci within-individuals, allowing for a maximum distance threshold of

15%. This was done using the VSEARCH algorithm (Rognes, Flouri, Nichols, Quince, & Mahé,

200   2016) as implemented in pyRAD, so as to remove read clusters with 3+ indels, >5 ambiguous

consensus nucleotides, or a coverage <20X or >500X. Putative homologs were identified using

202   among-individual clustering with the same parameters, and additionally removing any locus

having >2 alleles per individual, >70% heterozygosity for any polymorphic site, >10

204   heterozygous sites, or <50% individual recovery (see *github.com/tkchafin/scripts* for post-

processing and file formatting scripts).

2.2 | Derivation of population structure

208   We first sub-sampled the dataset to one SNP per locus and excluded those with a minor allele

count <2. We then inferred population structure (ADMIXTURE: Alexander et al., 2009) with

210   parallel processing (ADMIXPIPE: Mussmann et al., 2020). Model selection (i.e. for K, the number

of populations) followed a cross-validation approach with results aggregated from 20

212   independent replicates (CLUMPAK: Kopelman et al., 2015).

Individual-level ADMIXTURE results were summarized as a 'surface' with spatial

214   discontinuities represented as interpolated assignment probabilities. Here, we constructed state-

wide rasters, as representing per-pixel probabilities or 'ancestry proportions,' using Empirical

216   Bayesian Kriging (EBK: ARCMAP 10.7.1, Esri, Inc.). Probability surfaces were then summarized

as evenness and diversity of ancestries in a given cell using Simpson's index (Simpson, 1949)

218   (where K=number of statewide sub-populations). Our use of the diversity index was based on a

straightforward prediction: Areas representing spatial transitions between populations will have a

220   correspondingly low certainty of assignment to a given sub-population ancestry. Likewise, those

with low inter-population exchange will be comprised of genetically similar individuals assigned

222   with high probability to the endemic ancestry. Thus, intraspecific suture zones represent a

marked transition from one population to another, as identified by site-wise diversity in

224   assignment probabilities (i.e., "ancestry diversity").

We expect spatial variation in ancestry diversity to be inversely proportional to true rates

226   of gene flow, in that this quantity (as well as the ancestry proportions from which it is computed)

are a product of gene flow averaged over many generations. Of note is the fact that the method

228   only examines local ancestry probabilities and will not be dominated by translocation-related

artefacts (as are methods based on interpolated pairwise genetic distances or F_{ST}). Patterns based

230   on Simpson's diversity were contrasted with those inferred using a form of 2D-stepping stone

model (EEMS; Petkova et al. 2015), which was run with 2 million MCMC iterations (1 million

232   as burn-in), and sampled every 1,000 iterations [following parameter sweeps that tuned MCMC

acceptance rates to fall between 20% and 50% (Roberts, Gelman, & Gilks, 1997)].

2.3 | Estimating clinal parameters

236   We examined the nature of intraspecific suture zones by examining how individual loci

transitioned across these regions, relative to the genome-wide average. To do so, we defined

238   eight transects across putative suture zones, sampling 32-73 individuals per transect ($\bar{x}$ =55.38).

Individuals were chosen to represent each sub-population using a probability threshold applied to

240   ADMIXTURE results. Loci were filtered to remove all SNPs with missing data in >50% of

individuals, and further restricted with regard to computational time by retaining only those loci

242   with a sufficient allele frequency differential [computed as $\delta > 0.50$; (Gregorius & Roberds,

1986)]. Locus-wise clinal patterns were then inferred using a Bayesian method developed

244   originally for hybrid zones (Gompert & Buerkle, 2011, 2012). Open source Python code for

filtering SNP matrices and generating necessary input are available at:

246   (*github.com/tkchafin/scripts/phylip2introgress.pl* and *phylip2bgc.pl*).

Analyses were performed for each transect across 4 replicates, each using 1 million

248   MCMC iterations, discarding the first 500,000 as burn-in, with output thinned to every 500

iterations. Results were summarized by visualizing a 2-D density of cline shape parameters.

250   These are: α (=cline center) that describes an increase ($\alpha > 0$) or decrease ($\alpha < 0$) in the

probability of locus-specific ancestry from a parental population; and β (=cline rate) that defines

252   the rate of transition in probabilities of locus-specific ancestries having either steep ($\beta > 0$) or

wide ($\beta < 0$) shapes (Gompert, Parchman, & Buerkle, 2012). In this context, a locus which does

254   not deviate from the genome-wide pattern would have $\alpha=\beta=0$. Deviation in a directional manner

(i.e., an increase or decrease of one ancestry over another) is described by α, whereas deviation

256   in the rate of ancestry change around an inflection point (i.e., sigmoidal) is described by β.

Statistical outliers were designated using the method of Gompert and Buerkle (2011). BGC

258   results were parsed and visualized using the ClinePlotR R package (Martin et al. 2020a,b)

260   **2.4 | Relative dispersal by age and sex**

            Of particular interest in wildlife management is the backwards inference of geographic

262   positioning from genotypes—that is, the geolocation of 'origination' points for sampled animals.

This could be used, for example, to ascertain the geographic origin of poached individuals, or to

264   estimate post-natal dispersal. To this end, we used the novel, deep-learning method LOCATOR

(Battey, Ralph, & Kern, 2020) to predict the geographic origin of samples without relying upon

266   explicit assumptions about population genetic processes underlying spatial genetic differentiation

(Bradburd & Ralph, 2019). This analysis was performed iteratively across each individual, using

268   the remaining samples to train the LOCATOR classifier, with 100 bootstrap pseudo-replicates to

assess variance in geolocation. Given computational constraints, we performed the analysis using

270   a subset of 5000 SNPs having a minor allele frequency >10%.

            We estimated relative dispersal distances as the Euclidean distance between sampled

272   localities and the centroid of predicted coordinates, under the assumption that the distance

between predicted and collection locations is the result of lifetime dispersal, at least for samples

274   for which geolocation variance is low among pseudo-replicates. These results were then

partitioned by age, sex and CWD-status.

276   Our second approach examined the decay in genetic relatedness as a function of distance

from each individual, measured as a Prevosti distance (R-package poppr: Kamvar et al., 2014).

278 Here, the assumption is that recently dispersed individuals will be, on average, more genetically

dissimilar from resident individuals, whereas resident individuals having an appreciable

280 reproductive output will be less so. These calculations were limited to individuals that had

neighboring samples within a 5km radius, thus implicitly restricting the analysis to high-density

282 sampling regions. We note, however, that the traditional aging method employed for AR WTD

(Severinghaus 1949) seemingly has reduced accuracy in older deer, potentially suggesting

284 caution in the interpretation of results (Cook & Hart, 1979; Gee, Holman, Causey, Rossi, &

Armstrong, 2002; Mitchell & Smith, 1991).

3 | RESULTS

288 ### 3.1 | Data processing

Our raw data represented N=1,143 samples, including N=29 from Wisconsin (Table S1). We

290 removed N=83 that had missing metadata, discrepancies with coordinates, or <50% of loci

present. Assembly in pyRAD yielded an average of 25,584 loci per sample (σ=8,639). After

292 removing loci present in <50% of samples and excluding those containing potential paralogs

(e.g. excessive heterozygosity or >2 alleles per locus), our final dataset contained 35,420 loci,

294 from which 2,655,584 SNPs were catalogued. Of these, 54,102 were excluded as singletons. To

limit signal redundancy, we then condensed the data to one SNP per locus, yielded a final matrix

296 of 35,099 SNPs for analyses of population structure.

298 ### 3.2 | Population structure and 'ancestry surfaces'

Cross-validation, performed on N=20 replicates each for subpopulation model (K=1-20),

300   revealed the optimal number of clusters as K=8. Spatial orientation of these samples (Fig. 1)

provided a geographic definition, with some subpopulations qualitatively defined by apparent

302   landscape features, such as the Arkansas River Valley as the southern extent of subpopulations

k=3 and k=6.

304   Two ancestries ($k1$ and $k8$) largely dominated south of the Arkansas River, bounded by

Interstate 30 to the north and the Ouachita River to the south (Figs. 1-2), each of which supports

306   the argument that genetic structure is defined by large geographic barriers. The southwestern

portion of the state has two ancestral assignments ($k1$ and $k3$), with the latter having mixed

308   representation in the north-central section (potentially an artefact of weak differentiation rather

than true shared ancestry). The southeastern section is dominated by a single gene pool ($k8$),

310   which coincidentally subsumed all Wisconsin samples, a strong signature of genetic variability

as a residual of historic translocations.

312   A greater amount of endemic structure occurred in the Ozark Mountains, north of the

Arkansas River, where six sub-populations were evident. The most broadly distributed ($k5$) was

314   to the east in the Mississippi alluvial plains, extending westward across the mainstem of the

White River then northward towards the confluence of the Black and White rivers, where it

316   grades into several distinct yet loosely defined sub-populations (Figs. 1-2). The northwestern

corner of the state was the most heterogeneous, with four primarily endemic gene pools (k=2,3,4

318   and 7; Fig. 2). The northern-most of these was approximately bounded by the White River (Fig.

2), and graded westward into an area of high apparent admixture (Fig. 1). The remaining

320   northwestern region was defined by several gene pools showing considerable admixture and

spatially weak transitions, suggesting reduced gene flow but with geographic and/ or

322   environmental boundaries reasonably porous.

Effective migration surfaces (EEMS) failed to capture any ascertainable pattern relating

324   to spatially defined population structure (Fig. S1). Geographic breaks separating sub-populations

(=suture zones) were captured instead by reducing the dimensionality of interpolated assignment

326   probabilities as a continuous Simpson's diversity index (Fig. 3). This, in turn, reflects a

dependence on homogeneity of local assignments, rather than global patterns compounded by

328   long-distance transplants.

330   **3.3 | Intraspecific genomic clines**

Genomic clines varied substantially among transects (Figs. 4, S2). Most inter-population

332   comparisons within northwest Arkansas, to include $k2$ x $k3$ (both eastern and southern transition

zones), $k2$ x $k6$, and $k4$ x $k7$, indicated variation primarily restricted to cline centers (α), with

334   cline steepness (β) at a minimum. The variation in locus-wise pattern for these cases (hereafter

termed "α-dominant"), indicated a directional change in the representation of reference

336   populations across the transect, but without a noticeable 'inflection' point, as implied by non-

zero cline rates (β).

338   Remaining comparisons, including one additional transect from northwest Arkansas ($k3$ x

$k4$), primarily reflected variation in cline rate (β), in that loci varied most prominently with

340   regard to steepness/width of transition around an inflection point respective to genome-wide

ancestry. Although two transects showed minor exception to this pattern (i.e., $k3$ x $k6$ varied

342   along both parameters; and $k4$ x $k7$ showed minimal variation in either), the contrasting variation

between α-dominant versus β-dominant transects suggests that different processes underly

344   ancestry transitions.

346   **3.4 | Estimating dispersal using geolocation analysis**

Individuals from densely sampled regions could be assigned to geographic origin using the

348   'deep-learning' approach, with CWD-positive individuals assigned with a mean bootstrap

distance from centroid prediction generally <15km (Fig. S3). However, we did find assignment

350   error (e.g., among bootstraps) was elevated in low-density sampling regions (Figs. 5, S4), which

resulted in higher estimated individual dispersal distances (Fig. S4A, B). Given that variance in

352   dispersal estimates dropped considerably below ~25km (Fig. S4C), a conservative threshold of

10km was chosen and all individuals having a mean bootstrap-centroid distance above that were

354   removed for the purposes of dispersal estimates. After filtering, N=110 samples remained (Fig.

5A). A higher error threshold (i.e., 20km) allowed a greater number of total samples to be

356   incorporated (N=264; Fig. 5B), with several collected as 'roadkill' appearing as long-distance

transfers. Although these results represent low-precision assignments, they do underscore the

358   capacity of the method with regard to the identification of transported individuals (e.g., as

poached or illegally dumped deer, or carcasses transported across state lines or regional

360   management zones).

Geo-located results for those individuals passing a strict error filter demonstrated a

362   dispersal distance for males approximately doubled that of females across all age classes

(statistically significant only for the Y2-2.5 class due to low sample sizes; Fig. 6). This pattern

364   was established as early as the Y1-1.5 group, indicating apparent male dispersal by that age.

Smaller dispersal distances were found for fawns across both sexes (Fig. 6), again corroborating

366   *a priori* biological expectations.

Patterns of genetic dissimilarity also showed an age x sex effect, with greater genetic

368   dissimilarity for neighboring Y0-1 males than females, and with a shift towards reduced

dissimilarity in males >5 (Fig. S5). This again supports that male deer in Arkansas have

370   dispersed by the Y1-1.5 age class. These results potentially reflect age-biased reproduction as

well. Males contributed disproportionately to their local gene pools by age 5 (i.e., producing

372   offspring with resident females), thereby creating a pattern of lower genetic dissimilarity among

neighboring individuals, regardless of distance.

4 | DISCUSSION

376   **4.1 | Genetic footprints of historic management**

The prolonged history of WTD hunter-harvest, and subsequent long-range translocations into

378   Arkansas and the surrounding region (Ellsworth et al., 1994; Holder, 1951), provide a sideboard

to our study in that artificial long-distance movements such as these violate the assumptions

380   inherent with many spatially-explicit methods. However, we found estimates of ancestry

diversity and probability surfaces to be qualitatively robust in that they firmly recapitulated the

382   record of historic translocations (Holder, 1951). This, in turn, necessitates an explanation of

historic context.

384   In the early 20th century, following many decades of over-hunting, AR WTD declined to

<500 individuals (Holder, 1951). In response, Arkansas Game and Fish Commission (AGFC)

386   implemented an extensive restocking program (1941-1951) by establishing three primary sources

as its basis. The first was Howard County WTD farm (southwestern Arkansas; Fig. 7),

388   established from locally transplanted central Arkansas individuals (Wynn, 1943). It is now

located within the epicenter of population $k1$ (Figs. 2, 7), with ancestry shared elsewhere in the

390   state, thus supporting it as a local translocation epicenter.

A second major source was the Sylamore District (Ozark Mountains; Wood, 1944;

392   Holder, 1951), where individuals were naturally abundant (Fig. S7). Individuals from this region

have a consistently higher probability of assignment to $k3$ (Figs. 2, 7), where records indicate

394   ~81% of repopulated individuals in the Gulf Coastal Plain originated from the Sylamore cluster

(Holder, 1951; Karlin, Heidt, & Sugg, 1989; Wood, 1944). Our analyses are in agreement, in that

396   many individuals in southwestern Arkansas (Figs. 1, 6) reflect a mixed assignment to the

"Sylamore" cluster. They also comprised ~36% of stocking efforts in the Mississippi Delta

398   region (Karlin et al., 1989).

Our results also indicate some mixed assignment of individuals to $k3$, although a more

400   widespread representation is found in $k5$. AGFC surveys (~1942-1945) indicate WTD as

relatively abundant in the southeastern Delta region (Figs. 7, S7), allegedly linked with re-

402   population efforts following the disastrous 1927 Mississippi flood. This event, coupled with

over-hunting, nearly extirpated WTD in the region, save small numbers sustained by local

404   sportsmen (Holder, 1951; Fig. S7). We hypothesize WTD in the Delta region largely descended

from those efforts, given the contemporary homogeneity of ancestry assignment for this region.

406   A third (extraneous) WTD source was the Sandhill Game Farm (Babcock, WI; Wood

1944). Records indicate that ~64% of releases into the Mississippi Delta region originated out-

408   of-state, the majority from Wisconsin (Holder, 1951; Karlin et al., 1989). Our Wisconsin samples

were unanimously assigned to a gene pool prominently represented in the southern Delta region

410   ($k8$; Fig. 2), firmly establishing its genetic legacy as extending from imported deer.

Mitochondrial haplotypes putatively originating from Wisconsin were also uncovered in

412   Missouri, Kentucky, and Mississippi (Budd, Berkman, Anderson, Koppelman, & Eggert, 2018;

DeYoung et al., 2003; Doerner et al., 2005).

414   All genetic 'clusters' that lacked spatial cohesion in our analyses can be connected to the

three major stocking sources involved in earlier restoration efforts (per reference to historic

416   records). The remaining sub-populations, primarily in the Ozarks (k2, 4, 6, and 7) may represent

natural re-colonization from refugial populations (Figs. 2, 7), an hypothesis supported by early

418   census data (Figs. S7, S8).

420   **4.2 | Landscape-drivers of contemporary population structure**

A primary focus in our study was the contemporary genetic structure of AR WTD, especially

422   with regard to the manner by which these data promote landscape resistance and potentially

constrain CWD infection (Hemming-Schroeder, Lo, Salazar, Puente, & Yan, 2018; Kelly et al.,

424   2014). Yet several prerequisites are apparent in this regard. For example, one question which is

apparent but previously constrained by technological limitations is the degree to which potential

426   patterns of genetic variability in WTD have been conflated by anthropogenically-mediated

translocations.

428   We addressed this issue by utilizing next-generation molecular techniques to derive

highly variable SNP markers. We also implemented advanced geospatial procedures that

430   visualized the spatial transitions inherent within WTD ancestry in such a way that translocation

histories were not a limiting factor. We did so by interpolating our assignment probabilities from

432   ADMIXTURE, then applying Simpson's diversity index as a means of reducing the

dimensionality of these probabilities. We also examined locus-specific patterns within our

434   transitions so as to polarize stochastic versus deterministic processes.

        Previous studies concluded that landscape genetic inferences were limited at best, due to

436   complex interactions between historic translocations and subsequent population growth (Leberg

        et al. 1994; Leberg and Ellsworth 1999; Budd et al. 2018). However, many of these studies relied

438   upon (now) obsolescent molecular markers (e.g., mtDNA or reduced panels of microsatellite

        DNA markers) that captured substantially less polymorphism than do next-generation methods

440   (Hodel et al., 2017; Jeffries et al., 2016; Lemopoulos et al., 2019). Subsequent studies at finer

        spatial scales have supported the role of large-scale geomorphic configurations (e.g., rivers,

442   highways) as semipermeable barriers to WTD gene flow (Kelly et al., 2014; Locher et al., 2015;

        Miller et al., 2020; Robinson et al., 2012), despite the potential complication caused by re-

444   stocking efforts, and in accordance with radio-telemetry data (Peterson et al., 2017).

        Spatio-genetic patterns can still be informative with regards to landscape-level dispersal,

446   despite rapid transitions between historical and contemporary conditions (Epps & Keyghobadi,

        2015) which might be expected to obfuscate relationships. For example, spatial genetic patterns

448   in invasive species are also attributable to large-scale environmental features, despite oft-reduced

        levels of genetic diversity (Lopez, Hurwood, Dryden, & Fuller, 2014; Sacks, Brazeal, & Lewis,

450   2016; Zalewski, Piertney, Zalewska, & Lambin, 2009). The same can be said for populations

        occupying landscapes defined by recent anthropogenic developments, such as large urban

452   centers, suggesting the potential for such patterns to rapidly emerge (Beninde et al., 2016;

        Combs, Puckett, Richardson, Mims, & Munshi-South, 2018; Kimmig et al., 2020). In the case of

454   translocation, the magnitude of analytical artefacts are expected to be both scale-dependent, and

        a function of the specific analytical assumptions. For example, classic tests of 'isolation-by-

456   distance' implicitly assume a negative relationship between geographic proximity and pairwise

patterns of genetic distance (Meirmans, 2012; Rousset, 1997). Here, translocations are

458   demonstrably inconsistent with this expectation.

The assignment method used in our study yielded patterns of genetic similarity that

460   reflected translocations as depicted in historic records (Fig. 7). We sought to effectively

'remove' this artificial signal so as to more appropriately expose landscape features that could

462   potentially modulate deer dispersal. Our approach (i.e., reducing dimensionality of assignment

probabilities) produces relatively straightforward predictions about the ensuing metric: Areas

464   with an elevated flux of individuals should have elevated homogeneity with regard to

interpolated ancestry assignment (i.e., ADMIXTURE ancestry proportions). By the same logic,

466   regions that reflect transitions in ancestry proportions would also demonstrate reduced

homogeneity. Although spatial assignments can also be vulnerable to artefacts, such as the over-

468   fitting of discrete clusters within otherwise continuous populations (Bradburd et al., 2017), they

effectively identify linear barriers to gene flow (Blair et al., 2012). Coincidentally, our approach

470   to dimensionality-reduction also revealed numerous linear subdivisions that aligned with major

landscape barriers, such as Arkansas rivers (Fig. 3). Other putative transitions approximately

472   corresponded with large urban centers.

474   **4.3 | Separating stochastic versus deterministic processes in clinal patterns**

It is difficult to disentangle contemporary demographic processes (versus analytical

476   idiosyncrasies) in the formation of these patterns. As a means of clarification, we generated

multi-locus genomic patterns (Epps & Keyghobadi, 2015) under the expectation that observed

478   deviations of locus-wise clines from the genome-wide average would dictate the relative

importance of stochastic versus deterministic processes (Barton, 1983; Barton & Hewitt, 1985;

Slatkin, 1973; Vasemägi, 2006). We found eight transects that varied in either the width of locus-wise transitions (β), or the direction of change in allele representation (α). Interestingly, 'α-

dominant' transects were found within the Ozark region, previously hypothesized as being naturally re-colonized from local refugia (i.e., $k2$ x $k3$E/S, $k2$ x $k6$, and $k4$ x $k7$; Figs. 4, S2). This

hypothesis is also supported by state-wide surveys (1940's) that found occupied WTD habitat roughly corresponding with each Ozark sub-population (Figs. 7, S8).

All transects displayed near-zero β values across loci as well, suggesting either stochastic processes (genetic drift, bottlenecks) or sampling artefacts as drivers of ancestry transitions,

given our expectation that a 'true' hard barrier between discrete populations would yield an under-representation of inter-population heterozygotes (as described by β).

One interpretation of this result is that genetic drift at the edge of an expanding refugium creates a 'rolling' founder effect, with alleles being over-represented ($\alpha>0$) or under-represented

($\alpha<0$) across the re-colonized territory (Excoffier & Ray, 2008; Hallatschek & Nelson, 2008). This phenomenon—termed "gene surfing"—can generate wholly neutral patterns that appear to

be adaptive (Peischl, Dupanloup, Bosshard, & Excoffier, 2016; Travis et al., 2007), and may explain the presence of numerous α-outlier loci in those transects (Fig. 4).

Interestingly, all of our β-dominant transects crossed major rivers: Transect $k3$ x $k6$ spanned the Arkansas River; $k3$ x $k5$ the White River; and $k1$ x $k8$ the Ouachita River. Transect

$k3$ x $k4$ crosses the smaller Buffalo River, which may explain the depressed variability in cline steepness (Fig. 4). Non-zero β values implicate variations on a sigmoidal cline shape, with values

describing rates of change in allele frequencies that are either steep ($\beta>0$) or wide ($\beta<0$).

         Classically, clinal patterns are established by selection, or the existence of hard

502   boundaries to gene flow (Endler, 1973; Slatkin, 1973). However, spatially variable gene flow

coupled with drift can generate clinal patterns without necessitating selection across a

504 considerable proportion of the genome (Vasemägi, 2006). Selection could feasibly be involved,

at least for regions involving translocation of WTD subspecies, in that crosses between

506 subspecies have elicited fitness impacts such as dystocia (abnomal maternal labor due to shape,

size, or position of the fetus; Galindo-Leal and Weber 1994). Previous studies have also

508 underscored the genetic cost of inter-subspecific stocking (Hopken, Lum, Meyers, & Piaggio,

2015), such as anomalous variability in breeding time among other southern-recovered WTD

510 populations (Sumners et al., 2015). Some limited evidence also exists for a pheromonal basis for

reproductive isolation among mule deer subspecies [*Odocoileus hemionus*; (Müller-Schwarze &

512 Müller-Schwarze, 1975)].

514 **4.3 | Hunter-harvest and its management implications**

Our results identify within AR WTD a diagnosable genetic signature of historic translocations

516 within the genomes of extant populations. This simultaneously underscores the success of early

restocking efforts, while also reiterating long-standing concerns about the genetic and/ or

518 phenotypic impacts of anthropogenic translocations (Meffe & Vrijenhoek, 1988). It also

reinforces the need to formalize as an explicit management consideration the intraspecific

520 taxonomy of WTD, both to better understand the (yet to be seen) evolutionary consequences of

past translocations, and to provide a more proactive baseline going forward (Cronin, 2003;

522 Gippoliti, Cotterill, Groves, & Zinner, 2018).

Resource agencies, in particular, should prioritize those populations that retain endemic

524 genetic diversity, and by so doing, tacitly acknowledge the importance of preserving this

evolutionary legacy. Such recognition would capitalize on the many years of natural selection

526   that have operated on those populations prior to anthropogenic interference. It would also open

the door for potential variation in hunter-harvest, such as by having a restricted season strictly for

528   'legacy' AR deer.

Our patterns of age- and sex-biased dispersal distances (Fig. 5, 6) largely recapitulate

530   expectations based on WTD life-history: Male dispersal established by the yearling cohort (Long

et al., 2008; McCoy, Hewitt, & Bryant, 2005; Shaw, Lancia, Conner, & Rosenberry, 2006);

532   females lack age-dependent dispersal (Lutz et al., 2015). The potential utility of the approach lies

within the context of a multi-state genetic monitoring program for WTD. While this would

534   necessitate inter-state investment and coordination, potential benefits of a 'next-generation'

approach would be in preserving the evolutionary heritage of WTD, supporting the adaptive

536   management of a biodiversity resource of national importance, as well as promoting the strong

tradition of hunting in North America (Geist, Mahoney, & Organ, 2000).

4.4 | CWD expansion and its management implications

540   Our results also suggest the existence of several discontinuities in gene flow across the state.

Given that CWD is a paramount issue in the global management of WTD (Rivera, Brandt,

542   Novakofski, & Mateus-Pinilla, 2019; Uehlinger, Johnston, Bollinger, & Waldner, 2016), a

valuable application of our results would be towards disease containment. Given that the spread

544   of CWD will be more rapid within rather than among populations, monitoring efforts should

focus on detecting 'breaches' between genetically distinct regions (which are most often divided

546   by major rivers and urban centers, as herein; Fig. 3). Interestingly, the current boundaries for the

AGFC 'Deer Management Units' (DMUs) (Meeker et al., 2019), are remarkably consistent with

548   many of the intraspecific suture zones identified in our study. Thus, potential mitigation efforts

should work to constrain the transportation of potentially infectious material (such as carcasses)

550 from CWD-detected DMUs. Management actions should reduce the outward dispersal of

individuals from DMUs, and one approach is to reduce the density of yearling males, which

552 often comprise >50% of emigrating individuals (McCoy et al., 2005; Nelson, 1993; Nixon,

Hansen, Brewer, & Chelsvig, 1991). A similar action could be management to reduce local

554 population densities. The latter, in turn, acts to drive dispersal in that robust population numbers

are a catalyst for increasing social pressure as well as resource competition (Long et al., 2008;

556 Lutz et al., 2015; Shaw et al., 2006).

One harvest strategy ['Quality Deer Management,' QDM; (Brothers and Ray, 1975;

558 Hamilton et al., 1995)] has several management goals operating in parallel to improve WTD

demography. These are: Actively decreasing overall population density; elevating the proportion

560 of mature males in the population; and increasing the female component of the sex ratio. The

success of this approach was assayed by evaluating WTD emigration rates pre- and post-

562 application (Shaw et al., 2006). Two important aspects were noted: Emigration probabilities

decreased, as did mating behavior of yearling males. The implications of a management strategy

564 focusing on these explicit goals would be a serious reduction in CWD dissemination. It is unclear

how this paradigm could impact long-term disease trajectories within CWD-prevalent DMUs,

566 although lower emigration probabilities and decreased densities would concomitantly depress the

probability of disease contraction for a given individual.

4.5 | Selection for CWD genetic variants and its management implications

570 Previous CWD research in WTD populations has focused on the prevalence of a polymorphism

in the *PRNP* gene [responsible for the malformed protein product that constitutes the prion agent

572 of CWD (Johnson et al., 2011; Race, Meade-White, Miller, Fox, & Chesebro, 2011; Rivera et al.,

2019)]. Those WTD exhibiting a non-synonymous transition at amino acid residue 96 (from G to

574 S) had decreased susceptibility to CWD. This variant had a significantly greater presence among

older individuals in CWD-detected areas, suggesting increased survivability for those individuals

576 manifesting that variant (Chafin et al., 2020; Douglas et al., 2020). Results herein support this

result, in that older males (>4 years) are genetically more similar to neighboring individuals than

578 to younger males (Fig. S6). This in turn suggests a non-linear increase in reproductive success

with age [or perhaps with other collinear factors as well, such as body weight or antler size

580 (Newbolt et al., 2017)].

Directing management efforts towards an older age-structure could therefore have an

582 ancillary benefit in that it would promote the relative reproduction of individuals having the 96S

variant, given its presence at a significantly higher probability among more mature individuals

584 (Douglas et al. 2020b). Our results could also be used to explore those environmental variables

potentially responsible for resistance to movement at the landscape-level. This aspect, in

586 conjunction with ongoing work on spatially variable genetic susceptibility, could promote the

development of risk assessment models that assay disease spread.

DATA AVAILABILITY STATEMENT

590 All raw sequence files will be accessioned in the NCBI Sequence Read Archive (SRA) under

BioProject XXXXXX following acceptance; relevant curated (i.e. assembled and filtered) data

592 will be archived on Dryad (doi: XXXXXX); codes and custom scripts developed in support of

this work are also available as open-source via GitHub under the GNU Public License:

594 github.com/tkchafin (and as cited in-text).

REFERENCES

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, *19*(9), 1655–1664.

Andrewartha, H., & Birch, L. (1956). *The distribution and abundance of animals*. Chicago: University of Chicago Press.

Avise, J. C. (1992). Molecular population structure and the biogeographic history of a regional fauna: A case history with lessons for conservation biology. *Oikos*, *63*(1), 62.

Baker, K. H., Gray, H. W. I., Ramovs, V., Mertzanidou, D., Akin Pekşen, Bilgin, C. C., … Hoelzel, A. R. (2017). Strong population structure in a species manipulated by humans
since the Neolithic: The European fallow deer (*Dama dama dama*). *Heredity*, *119*(1), 16–26.

Barton, N., & Clark, A. (1990). Population structure and processes in evolution. In *Population Biology* (pp. 115–173). Springer Berlin.

Barton, N. H. (2008). The effect of a barrier to gene flow on patterns of geographic variation. *Genetics Research*, *90*(1), 139–149.

Barton, Nicholas H. (1983). Multilocus Clines. *Evolution*, *37*(3), 454–471.

Barton, Nicholas H., & Hewitt, G. M. (1985). Analysis of hybrid zones. *Annual Review of*
612   *Ecology and Systematics*, *16*(1), 113–148.

Battey, C. J., Ralph, P. L., & Kern, A. D. (2020). Predicting geographic location from genetic
614   variation with deep neural networks. *ELife*, *9*, e54507.

Beninde, J., Feldmeier, S., Werner, M., Peroverde, D., Schulte, U., Hochkirch, A., & Veith, M.
616   (2016). Cityscape genetics: structural vs. functional connectivity of an urban lizard population. *Molecular Ecology*, *25*(20), 4984–5000.

618   Blair, C., Weigel, D. E., Balazik, M., Keeley, A. T. H., Walker, F. M., Landguth, E., … Balkenhol, N. (2012). A simulation-based evaluation of methods for inferring linear barriers
620   to gene flow. *Molecular Ecology Resources*, *12*(5), 822–833.

Bohonak, A. J. (1999). Dispersal, gene flow, and population structure. *The Quarterly Review of*
622   *Biology*, *74*(1), 21–45.

Bossart, J. L., & Prowell, D. P. (1998). Genetic estimates of population structure and gene flow:
624   Limitations, lessons and new directions. *Trends in Ecology and Evolution*, *13*(5), 202–206.

Bradburd, G. S., Coop, G. M., & Ralph, P. L. (2018). Inferring continuous and discrete
626   population genetic structure across space. *Genetics*, 210, 33–52.

Bradburd, G. S., & Ralph, P. L. (2019). Spatial population genetics: It's about time. *Annual*

628   *Review of Ecology, Evolution, and Systematics*, *50*(1), 427–449.

Bradburd, G. S., Ralph, P. L., & Coop, G. M. (2016). A spatial framework for understanding
630   population structure and admixture. *PLoS Genetics*, *12*(1), 1–38.

Brinkman, T. J., Deperno, C. S., Jenks, J. A., Haroldson, B. S., & Osborn, R. G. (2005).
632   Movement of femal white-tailed deer: effects of climate and intensive row-crop agriculture.
       Journal of Wildlife Management, *69*(3), 1099–1111.

634   Brothers, A., & Ray, M. E. (1975). *Producing quality whitetails*. Laredo, TX: Wildlife Services.

Brown, S. K., Hull, J. M., Updike, D. R., Fain, S. R., & Ernest, H. B. (2009). Black bear
636   population genetics in California: Signatures of population structure, competitive release,
       and historical translocation. *Journal of Mammalogy*, *90*(5), 1066–1074.
638   https://doi.org/10.1644/08-mamm-a-193.1

Budd, K., Berkman, L. K., Anderson, M., Koppelman, J., & Eggert, L. S. (2018). Genetic
640   structure and recovery of white-tailed deer in Missouri. *Journal of Wildlife Management*,
       82(8), 1598–1607.

642   Chafin, T. K., Douglas, M. R., Martin, B. T., Zbinden, Z. D., Middaugh, C. M., Ballard, J., …
       Douglas, M. E. (2020). Age structuring and spatial heterogeneity in prion protein gene
644   (PRNP) polymorphism in Arkansas white-tailed deer. *BioRxiv*, doi:
       10.1101/2020.07.15.205039.

646   Chafin, T. K., Martin, B. T., Mussmann, S. M., Douglas, M. R. M. E., & Douglas, M. R. M. E.
       (2018). FRAGMATIC: in silico locus prediction and its utility in optimizing ddRADseq
648   projects. *Conservation Genetics Resources*, *10*(3), 325–328.

Combs, M., Puckett, E. E., Richardson, J., Mims, D., & Munshi-South, J. (2018). Spatial
650   population genomics of the brown rat (*Rattus norvegicus*) in New York City. *Molecular
       Ecology*, *27*(1), 83–98.

652   Comte, L., & Olden, J. D. (2018). Fish dispersal in flowing waters: A synthesis of movement-
       and genetic-based studies. *Fish and Fisheries*, 19(6), 1063-1077.

654   Cook, R. L., & Hart, R. V. (1979). Ages assigned known-age Texas white-tailed deer: Tooth
       wear versus cementum analysis. In *Proceedings of the Annual Conference of the
656   Southeastern Association of Fish and Wildlife Agencies*.

Corlett, R. T. (2015). The Anthropocene concept in ecology and conservation. *Trends in Ecology
658   and Evolution*. https://doi.org/10.1016/j.tree.2014.10.007

Cronin, M. A. (2003). Research on deer taxonomy and its relevance to management. *Ecoscience*,
660   *10*(4), 432–442. https://doi.org/10.1080/11956860.2003.11682791

Crooks, K. R., & Sanjayan, M. (2006). *Connectivity Conservation*. (K. R. Crooks & M.
662   Sanjayan, Eds.), *Connectivity Conservation*. Cambridge: Cambridge University Press.

664 Cushman, S. A., Elliot, N. B., Macdonald, D. W., & Loveridge, A. J. (2016). A multi-scale assessment of population connectivity in African lions (Panthera leo) in response to landscape change. *Landscape Ecology*, *31*(6), 1337–1353.

666 Darimont, C. T., Carlson, S. M., Kinnison, M. T., Paquet, P. C., Reimchen, T. E., & Wilmers, C. C. (2009). Human predators outpace other agents of trait change in the wild. *Proceedings of* 668 *the National Academy of Sciences of the United States of America*, *106*(3), 952–954.

DeYoung, R. W., Demarais, S., Honeycutt, R. L., Rooney, A. P., Gonzales, R. A., & Gee, K. L. 670 (2003). Genetic consequences of white-tailed deer (*Odocoileus virginianus*) restoration in Mississippi. *Molecular Ecology*, *12*(12), 3237–3252.

672 Doerner, K. C., Braden, W., Cork, J., Cunningham, T., Rice, A., Furman, B. J., & McElray, D. (2005). Population Genetics of Resurgence : White-Tailed Deer in Kentucky. *Journal of* 674 *Wildlife Management*, *69*(1), 345–355.

Douglas, M. R., Chafin, T. K., Zbinden, Z. D., Martin, B. T., & Douglas, M. E. (2020). White- 676 tailed deer in Arkansas: Genetic connectivity and chronic wasting disease susceptibility. *Final Report to the Arkansas Game and Fish Commission*.

678 Eaton, D. A. R. (2014). PyRAD: assembly of de novo RADseq loci for phylogenetic analyses. *Bioinformatics*, *30*(13), 1844–1849.

680 Ellsworth, D. L., Honeycutt, R. L., Silvy, N. J., Smith, M. H., Bickham, J. W., & Klimstra, W. D. (1994). White-Tailed Deer Restoration to the Southeastern United States: Evaluating 682 Genetic Variation. *The Journal of Wildlife Management*, *58*(4), 686.

Endler, J. A. (1973). Gene flow and population differentiation. *Science*, *179*(4070), 243–250.

684 Endler, J. A. (1977). *Geographic variation, speciation, and clines*. Princeton, NJ: Princeton University Press.

686 Epps, C. W., & Keyghobadi, N. (2015). Landscape genetics in a changing world: Disentangling historical and contemporary influences and inferring change. *Molecular Ecology*, *24*(24), 688 6021–6040.

Excoffier, L., Foll, M., & Petit, R. J. (2009). Genetic Consequences of Range Expansions. 690 *Annual Review of Ecology, Evolution, and Systematics*, *40*(1), 481–501.

Excoffier, L., & Ray, N. (2008). Surfing during population expansions promotes genetic 692 revolutions and structuration. *Trends in Ecology and Evolution*, *23*(7), 347–351.

Fitzpatrick, B. M. (2013). Alternative forms for genomic clines. *Ecology and Evolution*, *3*(7), 694 1951–1966.

Galindo-Leal, C., & Weber, M. (1994). Translocation of deer subspecies: Reproductive 696 implications. *Wildlife Society Bulletin*, *22*(1), 117–120.

698 Gee, K. L., Holman, J. H., Causey, M. K., Rossi, A. N., & Armstrong, J. B. (2002). Aging white-tailed deer by tooth replacement and wear: A critical evaluation of a time-honored technique. *Wildlife Society Bulletin*, *30*(2), 387–393.

700 Geist, V., Mahoney, S. P., & Organ, J. F. (2000). Why hunting has defined the North American model of wildlife conservation. *Transactions of the 66th North American Wildlife and*
702 *Natural Resources Conference*, *10*, 3977–3990.

Gippoliti, S., Cotterill, F. P. D., Groves, C. P., & Zinner, D. (2018). Poor taxonomy and genetic
704 rescue are possible co-agents of silent extinction and biogeographic homogenization among ungulate mammals. *Biogeographia*, *33*(1), 41–54.

706 Gompert, Z., & Buerkle, A. C. (2011). Bayesian estimation of genomic clines. *Molecular Ecology*, *20*, 2111–2127.

708 Gompert, Z., & Buerkle, C. A. (2012). bgc□: Software for Bayesian estimation of genomic clines. *Molecular Ecology Resources*, *12*(6), 1168–1176.

710 Gompert, Z., Parchman, T. L., & Buerkle, C. a. (2012). Genomics of isolation in hybrids. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *367*(1587), 439–
712 450.

Gregorius, H. R., & Roberds, J. H. (1986). Measurement of genetical differentiation among
714 subpopulations. *Theoretical and Applied Genetics*, *71*, 826–834.

Hallatschek, O., & Nelson, D. R. (2008). Gene surfing in expanding populations. *Theoretical*
716 *Population Biology*, *73*(1), 158–170.

Hamilton, R. J., Know, W. M., & Guynn, D. C. (1995). How quality deer management works. In
718 K. V. Miller & R. L. Marchinton (Eds.), *Quality whitetails: the why and how of Quality Deer Management* (pp. 7–18). Mechanicsburg, PA: Stackpole.

720 Hemming-Schroeder, E., Lo, E., Salazar, C., Puente, S., & Yan, G. (2018). Landscape genetics: A toolbox for studying vector-borne diseases. *Frontiers in Ecology and Evolution*,
722 *6*(March), 1–11.

Hewitt, G. M. (2000). The genetic legacy of the Quaternary ice ages. *Nature*, *405*(6789), 907–
724 913.

Hewitt, G. M. (2001). Speciation, hybrid zones and phylogeography—or seeing genes in space
726 and time. *Molecular Ecology*, *10*(3), 537–549.

Hodel, R. G. J., Chen, S., Payton, A. C., McDaniel, S. F., Soltis, P., & Soltis, D. E. (2017).
728 Adding loci improves phylogeographic resolution in red mangroves despite increased missing data: Comparing microsatellites and RAD-Seq and investigating loci filtering.
730 *Scientific Reports*, *7*(1), 1–14.

Holder, T. H. (1951). A survey of Arkansas game. *Arkansas Game and Fish Commission*

732   *Federal Aid Publication Project II-R.*, 57–79.

Hopken, M. W., Lum, T. M., Meyers, P. M., & Piaggio, A. J. (2015). Molecular assessment of
734   translocation and management of an endangered subspecies of white-tailed deer
(*Odocoileus virginianus*). *Conservation Genetics*, *16*(3), 635–647.

736 Jeffries, D. L., Copp, G. H., Handley, L. L., Håkan Olsén, K., Sayer, C. D., Hänfling, B., …
Hanfling, B. (2016). Comparing RADseq and microsatellites to infer complex
738   phylogeographic patterns, an empirical perspective in the Crucian carp, *Carassius
carassius*, L. *Molecular Ecology*, *25*(13), 2997–3018.

740 Johnson, C. J., Herbst, A., Duque-Velasquez, C., Vanderloo, J. P., Bochsler, P., Chappell, R., &
McKenzie, D. (2011). Prion protein polymorphisms affect chronic wasting disease
742   progression. *PLoS ONE*, *6*(3).

Kamvar, Z. N., Tabima, J. F., & Grünwald, N. J. (2014). Poppr: an R package for genetic
744   analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ*, *2*,
e281.

746 Karlin, A. A., Heidt, G. A., & Sugg, D. W. (1989). Genetic variation and heterozygosity in
white-tailed deer in Southern Arkansas. *The American Midland Naturalist*, *121*(2), 273–
748   284.

Katzner, T. E., & Arlettaz, R. (2020). Evaluating contributions of recent tracking-based animal
750   movement ecology to conservation management. *Frontiers in Ecology and Evolution*,
7(519), 1-10.

752 Kays, R., Crofoot, M. C., Jetz, W., & Wikelski, M. (2015). Terrestrial animal tracking as an eye
on life and planet. *Science*, *348*(6240), aaa2478.

754 Keller, I., Wagner, C. E., Greuter, L., Mwaiko, S., Selz, O. M., Sivasundar, a., … Seehausen, O.
(2013). Population genomic signatures of divergent adaptation, gene flow and hybrid
756   speciation in the rapid radiation of Lake Victoria cichlid fishes. *Molecular Ecology*, *22*(11),
2848–2863.

758 Kelly, A. C., Mateus-Pinilla, N. E., Brown, W., Ruiz, M. O., Douglas, M. R., Douglas, M. E., …
Novakofski, J. (2014). Genetic assessment of environmental features that influence deer
760   dispersal: Implications for prion-infected populations. *Population Ecology*, *56*(2), 327–340.

Kimmig, S. E., Beninde, J., Brandt, M., Schleimer, A., Kramer-Schadt, S., Hofer, H., … Frantz,
762   A. C. (2020). Beyond the landscape: Resistance modelling infers physical and behavioural
gene flow barriers to a mobile carnivore across a metropolitan area. *Molecular Ecology*,
764   *29*(3), 466–484.

Kissling, W. D., Ahumada, J. A., Bowser, A., Fernandez, M., Fernández, N., García, E. A., …
766   Hardisty, A. R. (2018). Building essential biodiversity variables (EBVs) of species
distribution and abundance at a global scale. *Biological Reviews*, *93*(1), 600–625.

768   Knoche, S., & Lupi, F. (2012). The economic value of publicly accessible deer hunting land. *Journal of Wildlife Management*, *76*(3), 462–470.

770   Koen, E. L., Tosa, M. I., Nielsen, C. K., & Schauber, E. M. (2017). Does landscape connectivity shape local and global social network structure in white-tailed deer? *PLOS ONE*, *12*(3),
772   e0173570.

        Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A., & Mayrose, I. (2015). Clumpak:
774   a program for identifying clustering modes and packaging population structure inferences across K. *Molecular Ecology Resources*, *15*(5), 1179–1191.

776   Leberg, P. L., & Ellsworth, D. L. (1999). Further evaluation of the genetic consequences of translocations on southeastern white-tailed deer populations. *The Journal of Wildlife*
778   *Management*, *63*(1), 327–334.

        Leiss, W., Westphal, M., Tyshenko, M. G., Croteau, M. C., Oraby, T., Adamowicz, W., …
780   Krewski, D. (2017). Challenges in managing the risks of chronic wasting disease. *International Journal of Global Environmental Issues*, *16*(4), 277–302.
782   https://doi.org/10.1504/IJGENVI.2017.086716

        Lemopoulos, A., Prokkola, J. M., Uusi-Heikkilä, S., Vasemägi, A., Huusko, A., Hyvärinen, P.,
784   … Vainikka, A. (2019). Comparing RADseq and microsatellites for estimating genetic diversity and relatedness — Implications for brown trout conservation. *Ecology and*
786   *Evolution*, *9*(4), 2106–2120.

        Locher, A., Scribner, K. T., Moore, J. A., Murphy, B., & Kanefsky, J. (2015). Influence of
788   landscape features on spatial genetic structure of white-tailed deer in human-altered landscapes. *Journal of Wildlife Management*, *79*(2), 180–194.
790   https://doi.org/10.1002/jwmg.826

        Long, E. S., Diefenbach, D. R., Rosenberry, C. S., & Wallingford, B. D. (2008). Multiple
792   proximate and ultimate causes of natal dispersal in white-tailed deer. *Behavioral Ecology*, *19*(6), 1235–1242.

794   Lopez, J., Hurwood, D., Dryden, B., & Fuller, S. (2014). Feral pig populations are structured at fine spatial scales in tropical Queensland, Australia. *PLoS ONE*, *9*(3).

796   Lutz, C. L., Diefenbach, D. R., & Rosenberry, C. S. (2015). Population density influences dispersal in female white-tailed deer. *Journal of Mammalogy*, *96*(3), 494–501.

798   Martin, B. T., Douglas, M. R., Chafin, T. K., Placyk, J. S., Birkhead, R. D., Phillips, C. A., & Douglas, M. E. (2020a). Contrasting signatures of introgression in North American box
800   turtle (*Terrapene* spp.) contact zones. *Molecular Ecology*. Early View.

        Martin, B. T., Chafin, T. K., Douglas, M. R., & Douglas, M. E. (2020b). ClinePlotR: Visualizing
802   genomic clines and detecting outliers in R. *bioRxiv*. doi:10.1101/2020.09.05.284109

        Mawdsley, J. R. (2020). Phylogenetic patterns suggest broad susceptibility to chronic wasting

804   disease across Cervidae. *Wildlife Society Bulletin*, *44*(1), 152–155.

McCoy, J. E., Hewitt, D. G., & Bryant, F. C. (2005). Dispersal by yearling male white-tailed
806   deer and implications for management. *Journal of Wildlife Management*, *69*(1), 366–376.

Meeker, R., Brown, J., Carner, B., Stephens, K., Dugger, G., Groves, B., … White, D. J. (2019).
808   Arkansas Game and Fish Strategic Deer Management Plan. *Arkansas Game and Fish
        Commission. Little Rock, AR.*

810 Meffe, G. K., & Vrijenhoek, R. C. (1988). Conservation genetics in the management of desert
        fishes. *Conservation Biology*, *2*(2), 157–169.

812 Meirmans, P. G. (2012). The trouble with isolation by distance. *Molecular Ecology*, *21*(12),
        2839–2846.

814 Miles, L. S., Rivkin, L. R., Johnson, M. T. J., Munshi-South, J., & Verrelli, B. C. (2019,
        September). Gene flow and genetic drift in urban environments. *Molecular Ecology*.
816   Blackwell Publishing Ltd.

Miller, W. L., Miller-Butterworth, C. M., Diefenbach, D. R., & Walter, W. D. (2020).
818   Assessment of spatial genetic structure to identify populations at risk for infection of an
        emerging epizootic disease. *Ecology and Evolution*, *10*(9), 3977–3990.

820 Mitchell, C. J., & Smith, W. P. (1991). Reliability of techniques for determining age in southern
        white-tailed deer. *Journal of the Tennessee Adacemy of Science*, *66*(3), 117–120.

822 Müller-Schwarze, D., & Müller-Schwarze, C. (1975). Subspecies specificity of response to a
        mammalian social odor. *Journal of Chemical Ecology*, *1*(1), 125–131.

824 Mussmann, S. M., Douglas, M. R., Chafin, T. K., & Douglas, M. E. (2020). ADMIXPIPE:
        Population analyses in ADMIXTURE for non-model organisms. *BMC Bioinformatics*, 21,
826   337.

Nagylaki, T. (1976). Clines with variable migration. *Genetics*, *83*(4), 867–886.

828 Nathan, R., Klein, E., Robledo-Arnuncio, J. J., & Revilla, E. (2012). Dispersal kernels: review.
        In J. Clobert, M. Baguette, T. G. Benton, & J. M. Bullock (Eds.), *Dispersal ecology and
830   evolution* (pp. 187–210). Oxford, UK: Oxford University Press.

Nelson, M. E. (1993). Natal dispersal and gene flow in white-tailed deer in northeastern
832   Minnesota. *Journal of Mammalogy*, *74*(2), 316–322.

Newbolt, C. H., Acker, P. K., Neuman, T. J., Hoffman, S. I., Ditchkoff, S. S., & Steury, T. D.
834   (2017). Factors influencing reproductive success in male white-tailed deer. *Journal of
        Wildlife Management*, *81*(2), 206–217.

836 Nixon, C. M., Hansen, L. P., Brewer, P. A., & Chelsvig, J. E. (1991). Ecology of white-tailed
        deer in an intensively farmed region of Illinois. *Wildlife Monographs*, *118*, 3–77.

838   Nixon, C. M., & Mankin, P. C. (2016). Social and Environmental Influences on Female White-Tailed Deer Dispersal Behavior. *Great Plains Research*, *26*(2), 85–92.
840   https://doi.org/10.1353/gpr.2016.0017

Peischl, S., Dupanloup, I., Bosshard, L., & Excoffier, L. (2016). Genetic surfing in human
842   populations: from genes to genomes. *Current Opinion in Genetics and Development*, *41*, 53–61.

844   Peterson, B. E., Storm, D. J., Norton, A. S., & Van Deelen, T. R. (2017). Landscape influence on dispersal of yearling male white-tailed deer. *Journal of Wildlife Management*, *81*(8), 1449–
846   1456.

Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double digest
848   RADseq: An inexpensive method for *de novo* SNP discovery and genotyping in model and non-model species. *PLoS ONE*, *7*(5), e37135.

850   Petkova, D., Novembre, J., & Stephens, M. (2015). Visualizing spatial population structure with estimated effective migration surfaces. *Nature Genetics*, *48*(1), 94–100.

852   Picard, C., Dallot, S., Brunker, K., Berthier, K., Roumagnac, P., Soubeyrand, S., … Thébaud, G. (2017). Exploiting Genetic Information to Trace Plant Virus Dispersal in Landscapes.
854   *Annual Review of Phytopathology*, *55*(1), 139–160.

Polechová, J., & Barton, N. (2011). Genetic drift widens the expected cline but narrows the
856   expected cline width. *Genetics*, *189*(1), 227–235.

Race, B., Meade-White, K., Miller, M. W., Fox, K. A., & Chesebro, B. (2011). *In vivo*
858   comparison of chronic wasting disease infectivity from deer with variation at prion protein residue 96. *Journal of Virology*, *85*(17), 9235–9238.

860   Richardson, J. L., Brady, S. P., Wang, I. J., & Spear, S. F. (2016). Navigating the pitfalls and promise of landscape genetics. *Molecular Ecology*, *25*(4), 849–863.

862   Rivera, N. A., Brandt, A. L., Novakofski, J. E., & Mateus-Pinilla, N. E. (2019). Chronic wasting disease in cervids: Prevalence, impact and management strategies. *Veterinary Medicine:*
864   *Research and Reports*, *10*, 123–139. https://doi.org/10.2147/vmrr.s197404

Roberts, G. O., Gelman, A., & Gilks, W. R. (1997). Weak convergence and optimal scaling of
866   random walk Metropolis algorithms. *Annals of Applied Probability*, *7*(1), 110–120.

Robinson, S. J., Samuel, M. D., Lopez, D. L., & Shelton, P. (2012). The walk is never random:
868   Subtle landscape effects shape gene flow in a continuous white-tailed deer population in the Midwestern United States. *Molecular Ecology*, *21*(17), 4190–4205.

870   Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: a versatile open source tool for metagenomics. *PeerJ*, *4*, e2409v1.

872   Rousset, F. (1997). Genetic differentiation and estimation of gene flow from F-statistics under

isolation by distance. *Genetics*, *145*, 1219–1228.

874   Sacks, B. N., Brazeal, J. L., & Lewis, J. C. (2016). Landscape genetics of the nonnative red fox
         of California. *Ecology and Evolution*, *6*(14), 4775–4791.

876   Santangelo, J. S., Johnson, M. T. J., & Ness, R. W. (2018). Modern spandrels: The roles of
         genetic drift, gene flow and natural selection in the evolution of parallel clines. *Proceedings*
878   *of the Royal Society B: Biological Sciences*, *285*(1878).

         Severinghaus, C. W. (1949). Tooth development and wear as criteria of age in white-tailed deer.
880   *The Journal of Wildlife Management*, *13*, 195–216. https://doi.org/10.2307/3796089

         Shaw, J. C., Lancia, R. A., Conner, M. S., & Rosenberry, C. S. (2006). Effect of population
882   demographics and social pressures on white-tailed deer dispersal ecology. *Journal of*
         Wildlife Management, *70*(5), 1293–1301.

884   Shephard, J. M., Ogden, R., Tryjanowski, P., Olsson, O., & Galbusera, P. (2013). Is population
         structure in the European white stork determined by flyway permeability rather than
886   translocation history? *Ecology and Evolution*, *3*(15), 4881–4895.

         Simpson, E. H. (1949). Measurement of diversity. *Nature*, *163*(1943), 688.

888   Slatkin, M. (1973). Gene flow and selection in a cline. *Genetics*, *75*(4), 733–756.

         Slatkin, Montgomery. (1989). Population structure and evolutionary progress. *Genome*, *31*(1),
890   196–202. https://doi.org/10.1139/g89-034

         Sumners, J. A., Demarais, S., DeYoung, R. W., Honeycutt, R. L., Rooney, A. P., Gonzales, R.
892   A., & Gee, K. L. (2015). Variable breeding dates among populations of white-tailed deer in
         the southern United States: The legacy of restocking? *Journal of Wildlife Management*,
894   *79*(8), 1213–1225.

         Travis, J. M. J., Münkemüller, T., Burton, O. J., Best, A., Dytham, C., & Johst, K. (2007).
896   Deleterious mutations can surf to high densities on the wave front of an expanding
         population. *Molecular Biology and Evolution*, *24*(10), 2334–2343.

898   Turner, M. G. (1989). Landscape ecology: the effect of pattern on process. *Annual Review of*
         Ecology and Systematics. Vol. 20. https://doi.org/10.1146/annurev.es.20.110189.001131

900   Uehlinger, F. D., Johnston, A. C., Bollinger, T. K., & Waldner, C. L. (2016). Systematic review
         of management strategies to control chronic wasting disease in wild deer populations in
902   North America. *BMC Veterinary Research*, *12*(1).

         van Rees, C. B., Reed, J. M., Wilson, R. E., Underwood, J. G., & Sonsthagen, S. A. (2018).
904   Landscape genetics identifies streams and drainage infrastructure as dispersal corridors for
         an endangered wetland bird. *Ecology and Evolution*, *8*(16), 8328–8343.

906   Vasemägi, A. (2006). The adaptive hypothesis of clinal variation revisited: Single-locus clines as

a result of spatially restricted gene flow. *Genetics*, *173*(4), 2411–2414.

908   Waller, D. M., & Alverson, W. S. (1997). The white-tailed deer: A keystone herbivore. *Wildlife Society Bulletin*, *25*(2), 217–226.

910   Wang, I. J., & Shaffer, H. B. (2017). Population genetic and field-ecological analyses return similar estimates of dispersal over space and time in an endangered amphibian. 912   *Evolutionary Applications*, *10*(6), 630–639.

Williams, E. S., & Young, S. (1980). Chronic wasting disease of captive mule deer: a 914   spongiform encephalopathy. *Journal of Wildlife Diseases*, *16*(1), 89–98.

Wolverton, S., Kennedy, J. H., & Cornelius, J. D. (2007). A paleozoological perspective on 916   white-tailed deer (Odocoileus virginianus texana) population density and body size in central Texas. *Environmental Management*, *39*(4), 545–552.

918   Wood, R. (1944). Arkansas' Deer Transplanting Program. *Transactions of the North American Wildlife Conference*, *9*, 162–167.

920   Wynn, S. (1943). Howard County Game Refuge and Deer Farm. *The Arkansas Historical Quarterly*, *2*(4), 340–345.

922   Zalewski, A., Piertney, S. B., Zalewska, H., & Lambin, X. (2009). Landscape barriers reduce gene flow in an invasive carnivore: Geographical and local genetic structure of American 924   mink in Scotland. *Molecular Ecology*, *18*(8), 1601–1615.

926   **Figures and Tables**

928   **Figure 1**: **Ancestry proportions for 1,143 Arkansas white-tailed deer**, as inferred using the

program ADMIXTURE. Samples are represented as pie charts plotted at absolute collection

930   coordinates, with colors of assignment probabilities proportional to a particular subpopulation.

932   **Figure 2**: **Assignment probabilities for eight populations (k=1 through k=8)**, interpolated

using Empirical Bayesian Kriging. P(k)=1.0 corresponds to 100% probability of ancestry per

934   raster cell, and P(k)=0.0 corresponds to 0%. Individual samples are represented as black dots.

936   **Figure 3: Simpson's diversity of interpolated ancestry proportions.** Each raster cell was

assigned 8 values equal to the expected proportion of ancestry corresponding to the 8 genetic

938   clusters (Fig. 1), then summarized by calculating Simpson's diversity index.

940   **Figure 4: Relationships between genomic cline parameters** contrasted among eight sub-

sampled transects =of intraspecific suture zones in white-tailed deer (Fig. 3). Contour plots show

942   relative densities of SNPs varying in cline rate (β), representing the steepness of clines, and cline

center (α), representing bias in SNP ancestry. Outlier loci are highlighted in blue.

Figure 5: Summary of geo-location predictions in LOCATOR as a random subset of N=490

946   individuals, at two error thresholds: 10km (=A) and 20km (=B), thus constraining results to

N=110 (=A) and N=264 (=B). Error calculated as mean distance of bootstrap predictions from

948   the centroid predicted for each individual (see Fig. S4). A black dot denotes the predicted

location of an individual and a colored dot indicates the 'true' location (colors proportional to

950   measurement error in km).

952   **Figure 6: Inferred dispersal distances, partitioned by sex and age class**. Dispersal distances

were calculated as the difference (in km) between the predicted and true locations, excluding all

954   individuals with a mean prediction error of 10km (see Figs. 14-15). Sample sizes for each group

are given in black, with the mean in red below each box plot. P-values are reported for within-

956   age two-sample t-tests comparing means from males and females.

958 **Figure 7: Proposed translocation pathways and refugial populations in Arkansas**, showing

(left) translocation events supported by genetics along (dashed line), as well as those with

960 anecdotal support given limited historical records (solid), from three primary stocking sources:

Sandhill Game Farm in Wisconsin (W); 'Sylamore District' state refuge sites (S); and the

962 Howard County Game Farm (H). Also shown are major inhabited regions of white-tailed deer

(right) estimated from 1942-47 game surveys, excluding small regions (see full version in Online

964 Supplementary Material). Putative inhabited regions are also annotated according to their

hypothesized associations with contemporary genetic clusters (Fig. 1-2).

Assignment probability

0.00 - 0.20 0.21 - 0.40 0.41 - 0.60 0.61 - 0.80 0.81 - 1.00

979 **Supplementary Figures and Tables**

980 **Table S1:** Numbers (=N) of white-tailed deer genotyped across ~35,000 SNP loci. Samples were
981 collected from 75 counties (=County) in Arkansas (2016-2019). Code indicates standard 2-letter
982 county abbreviation. Samples removed due to missing data are not included.
983

County	Code	N	County	Code	N
Arkansas	AR	11	Lee	LE	9
Ashley	AS	1	Lincoln	LI	8
Baxter	BA	19	Little River	LR	10
Benton	BE	19	Logan	LO	32
Boone	BO	34	Lonoke	LN	11
Bradley	BR	2	Madison	MA	39
Calhoun	CA	14	Marion	MR	39
Carroll	CR	50	Miller	MI	8
Chicot	CH	3	Mississippi	MS	2
Clark	CL	7	Monroe	MO	10
Clay	CY	15	Montgomery	MN	2
Cleburne	CE	8	Nevada	NE	13
Cleveland	CV	5	Newton	NW	140
Columbia	CO	4	Ouachita	OU	12
Conway	CN	11	Perry	PE	8
Craighead	CG	7	Phillips	PH	9
Crawford	CW	6	Pike	PI	8
Crittenden	CT	8	Poinsett	PO	10
Cross	CS	10	Polk	PL	3
Dallas	DA	7	Pope	PP	62
Desha	DE	11	Prairie	PR	10
Drew	DR	14	Pulaski	PU	11
Faulkner	FA	14	Randolph	RA	8
Franklin	FR	20	Saline	SA	11
Fulton	FU	9	Scott	SC	2
Garland	GA	14	Searcy	SE	32
Grant	GR	5	Sebastian	SB	17
Greene	GE	16	Sevier	SV	12
Hempstead	HE	9	Sharp	SH	16
Hot Spring	HS	3	St. Francis	SF	7
Howard	HO	11	Stone	ST	15
Independence	IN	7	Union	UN	6
Izard	IZ	6	Van Buren	VB	23
Jackson	JA	2	Washington	WA	17
Jefferson	JE	13	White	WH	10
Johnson	JO	45	Woodruff	WO	9
Lafayette	LA	14	Yell	YE	31
Lawrence	LW	17	**Total**		**1,143**

Figure S1: **Effective migration rates and intra-population diversity** ($\log_{10}$ scale), calculated
986   from effective migration surfaces (EEMS). Rates are plotted according to colored bin, with
   divisions calculated as natural breaks using the Jenks algorithm in ArcMAP.
988

Figure S2: Sampled transects employed for genomic cline analysis with selected groups within each coalesced within colored hulls: Reference population 1 (p1; red); Reference population 2 (p2; green); and putative admixed individuals (popA; blue).

Figure S3: LOCATOR Geo-located results for CWD-positive individuals derived via

LOCATOR. Individuals are represented by dots, pairs of which are connected by a single line.
A black dot denotes predicted location, while a colored dot indicates 'true' (observed = sampled)

location (color proportional to distances separating each). Inserted figures offer full prediction
results for three selected samples, with bootstrap estimates demarked by a 95% contour.

1002 **Figure S4: Effects of error thresholds on inferred dispersal distances as predicted by LOCATOR analysis.** Inferred dispersal
kernels were computed at several error thresholds: (A) Error computed as mean distance of predicted coordinates for each individual
1004 at 100 bootstrap estimates from predicted centroid (=interpretation of localized vs. dispersed predictions). Dispersal distances
calculated as difference (in km) between predicted centroid versus 'true' (observed) location. Inferred dispersal distances generally
1006 increase for individuals with larger prediction error; (B) Larger stochastic variation in centroid location for individuals with lower
predicted precision among bootstraps; (C) Standard deviation of dispersal distances for the latter computed in a sliding window of
1008 10km along the x-axis (prediction error).

Figure S5: Spatial patterns of genetic dissimilarity among white-tailed deer partitioned by age and sex. Genetic distances between individuals and neighbors derived from 5,000 randomly

sampled SNPs depicted across physical distance (x-axis). Results depict different cohorts of males (A) and females (B) by age.

1016

Figure S6: **Census estimates for white-tailed deer from surveys in 1942-1946**

1018

1020

1022

1024

1026

1028

1030

1032

Figure S7: Subpopulation probabilities overlain with 'deer occupied territories' circa 1942-
1034 **1947** (in green). Polygons representing deer occupation were manually compiled from the
Wildlife and Cover Map of Arkansas (1942-47), prepared by Arkansas Game and Fish
1036 Commission in cooperation with U. S. Fish and Wildlife Service as one aspect of a Federal Aid
Project (drawn by Flaun M. Tolar). Map accessed from the University of Arkansas Library
1038 Arkansas Collection (Special Collections). A higher resolution geo-referenced version available
at a later date (reproduction/ copy permissions granted by University of Arkansas Libraries).
1040

1042