Dengue activates mTORC2 signaling to counteract apoptosis and maximize viral replication

Christoph C. Carter[1,2,#,*], Jean Paul Olivier[1,*], Alexis Kaushansky[1,3], Fred D. Mast[1], John D. Aitchison[1,3,4]

[1]Center for Global Infectious Disease Research, Seattle Children's Research Institute, Seattle WA 98109; [2]Division of Allergy and Infectious Diseases, Department of Medicine, University of Washington, Seattle WA 98109; [3] Department of Pediatrics, University of Washington, Seattle WA 98195; Department of Biochemistry, University of Washington, Seattle WA 98195

Running title: *Dengue activates mTORC2 for viral replication*

#Present address: Gilead Sciences, Inc., 333 Lakeside Drive, Foster City CA 94404
* contributed equally

To whom correspondence should be addressed:
  John D. Aitchison, PhD
  Center for Global Infectious Disease Research,
  Seattle Children's Research Institute, Seattle WA 98019;
  john.aitchison@seattlechildrens.org;
  Tel.(206) 884-3125; Fax. (206) 884-3104.

Keywords: Dengue virus (DENV), mammalian target of rapamycin (mTOR), mammalian target of rapamycin complex 2 (mTORC2), apoptosis, non-structural protein 5, flavivirus, pathogenesis, viral replication

ABSTRACT

The mechanistic target of rapamycin (mTOR) functions in at least two distinct complexes: mTORC1, which regulates cellular anabolic-catabolic homeostasis, and mTORC2, which is an important regulator of cell survival and cytoskeletal maintenance. mTORC1 has been implicated in the pathogenesis of flaviviruses including dengue, where it contributes to the establishment of a pro-viral autophagic state. In contrast, the role of mTORC2 in viral pathogenesis is unknown. In this study, we explore the consequences of a physical protein-protein interaction between dengue non-structural protein 5 (NS5) and host cell mTOR proteins during infection. Using shRNA to differentially target mTORC1 and mTORC2 complexes, we show that mTORC2 is required for optimal dengue replication. Furthermore, we show that mTORC2 is activated during viral replication, and that mTORC2 counteracts virus-induced apoptosis, promoting the survival of infected cells. This work reveals a novel mechanism by which the dengue flavivirus can promote cell survival to maximize viral replication.

INTRODUCTION

Mechanistic target of rapamycin (mTOR) is an essential serine/threonine kinase that functions in several key aspects of mammalian cell biology (reviewed in (1)). mTOR exerts its actions as a component of at least two distinct complexes, mTORC1 and mTORC2. mTORC1 is composed of five proteins: mTOR, Raptor, mLST8, PRAS40 and Deptor. mTORC1 functions as a master regulator of anabolic/catabolic homeostasis. In conditions of nutritional abundance, mTOR phosphorylates p70 ribosomal protein S6K (S6K) and eukaryotic initiation factor 4E binding protein 1 (4E-BP1), leading to increased protein synthesis. In conditions of nutrient scarcity mTORC1 is inactivated, stimulating autophagy, which allows for the recycling and turnover of cellular organelles and protein complexes (1).

mTORC2 is composed of six proteins: mTOR, Rictor, mLST8, SIN1, PRRL5L and Deptor. mTORC2 has distinct roles from mTORC1 in cellular physiology, but these roles are less well understood than those of mTORC1. mTORC2 promotes cell survival and proliferation through phosphorylation of AKT at ser473 (2). It is also known to play a role in the maintenance of the actin cytoskeleton, and when inactivated results in morphological abnormalities in some cell lines (3, 4). It has been suggested that mTORC2 may modulate translational machinery due to its association with ribosomes; however, the ramifications of this interaction are not well understood (5). Although mTORC1 is the canonical regulator of autophagy, mTORC2 has been implicated in the regulation of specific autophagic processes such as chaperone-mediated autophagy and mTORC1-independent autophagy (6, 7).

Given the important role of mTORC1 in regulating cellular metabolism, it is not surprising that several viruses have evolved mechanisms to modulate mTORC1 signaling (8, 9). Numerous viruses have been shown to manipulate mTORC1 activity during infection; some viruses activate mTORC1 to maintain cellular anabolic machinery, whereas others suppress mTORC1 activity to favor cap-independent viral protein synthesis (8, 9). In the case of dengue, virus-induced modulation of mTORC1 has been suspected due to the importance of autophagy in dengue infection (10-13), although this interaction of the virus with mTORC1 has not been comprehensively investigated. The importance of mTORC1 in dengue replication is supported by studies demonstrating increased viral replication in the response to pharmacologic mTOR inhibition and a recent study implicating mTORC1 in the dengue-induced activation of lipophagy (14, 15).

In contrast to mTORC1, the role of mTORC2 in virus-host interaction is poorly understood. Activation of mTORC2 has been documented in human cytomegalovirus, West Nile and influenza infection (16-18), but the functional role of mTORC2 in these infections remains unknown. Furthermore, to our knowledge no role for mTORC2 has been described in dengue infection.

Here, we describe a role for mTORC2 in promoting cell survival during dengue infection. We find that the dengue non-structural protein 5 (NS5) interacts with mTORC1 and mTORC2 complexes, and that dengue infection leads to the activation of mTORC2 signaling. We report that inactivation of mTORC2 signaling leads to a decrease in viral replication and an increase in virus-induced apoptosis and cell death. These findings suggest a mechanism by which dengue counteracts apoptosis to maintain cell survival and maximize viral replication.

91
RESULTS
Dengue NS5 protein interacts with mTORC1 and mTORC2
A previous quantitative proteomics approach (I-DIRT; Isotopic Differentiation of
Interactions as Random or Targeted(19)), designed to identify *bona fide* dengue-host
protein-protein interactions, defined a high-confidence protein interaction network
including a predicted interaction between the dengue NS5 protein and mTOR (20). To
validate and further study this interaction, we performed co-immunoprecipitation
experiments using exogenously expressed NS5. GFP-tagged NS5 or GFP alone was
transfected into 293FT cells, leading to modest expression of the fusion protein in the
cytosol with nuclear accumulation, as has been previously reported (21). Protein
complexes were then affinity purified using GFP-specific nanobodies (22). Subsequent
western blotting of affinity purified NS5 complexes revealed mTOR, Raptor, and Rictor
proteins, demonstrating that NS5 interacts with both mTORC1 and mTORC2 (Fig. 1).
105
mTORC2 is activated during dengue replication and is required for efficient viral
replication
To investigate what impact NS5 interactions may have on dengue viral infection, and to
explore the differential effect of NS5 interaction with mTORC1 and mTORC2, the
expression of each complex was silenced using lentivirus-expressed shRNA targeting
mTOR protein, Raptor (a component of mTORC1) or Rictor (a component of mTORC2)
or, as a control, a nonspecific scrambled oligo-sequence (23). We used HepG2
hepatoma cells for these experiments as they have been used extensively in studies of
dengue replication, apoptosis, autophagy and ER stress (15, 20, 24, 25). shRNA
transduction resulted in substantially decreased protein abundance of mTOR, Raptor
and Rictor (Fig. 2*A*). To assess whether Raptor and Rictor knockdown inhibited the
signaling activity of mTORC1 and mTORC2, we examined the phosphorylation status of
well-characterized downstream targets (p70 S6K thr389 for mTORC1 and AKT ser473
for mTORC2) (23, 26). Knockdown of Raptor led to diminished mTORC1 signaling as
evidenced by decreased S6K thr389 phosphorylation, and knockdown of Rictor led to
diminished mTORC2 signaling as evidenced by decreased AKT ser473 phosphorylation
(Fig. 2*A*). We observed that depletion of mTORC1 led to reciprocal activation of
mTORC2 activity, consistent with prior reports suggesting that mTORC1 represses
mTORC2 signaling (27) (Fig. 2*A*). The converse did not appear to be the case, as
depletion of mTORC2 did not lead to increased phosphorylation of S6K by mTORC1
(Fig. 2*A*). Furthermore, mTOR knockdown diminished levels of Raptor and Rictor, but
knockdown of neither Raptor or Rictor influenced levels of the other (Fig. 2*A*).
We next assessed the effect of reduced mTOR protein, Raptor (mTORC1), and
Rictor (mTORC2) on dengue serotype 2 replication by infecting respective knockdown
cells and quantifying the amount of infectious virus released. Interestingly, Rictor
knockdown led to a substantial decrease in the amount of viral replication, while Raptor
knockdown led to a small but significant increase in replication. No significant effect on
viral replication was observed when mTOR protein was knocked down (Fig. 2*B*).
Having observed a decrease in viral replication with mTORC2 inactivation, we
next asked whether dengue infection affected the activity of mTORC2. To do this, we
infected HepG2 cells with dengue and assessed the phosphorylation status of the

downstream targets AKT and S6K by western blot. In these experiments, mTORC2-
specific phosphorylation at AKT ser473 was increased in infected cells compared to
mock treated cells, and the ratio of ser473 phosphorylated AKT to total AKT was
consistently increased in infected cells (Fig. 2*C*, p < 0.001). Using mTOR, Raptor and
Rictor knockdown cells, we again observed an increase in AKT ser473 phosphorylation
in infected cells, which was abrogated in Rictor knockdown cells, demonstrating that
mTORC2 is required for dengue-induced AKT phosphorylation. As in Fig. 2*A*,
knockdown of mTOR protein or Raptor led to increased mTORC2 activity, which was
not further increased by dengue infection (Fig. 2*D*).
146
mTORC2 inhibition does not affect cellular morphology, growth rates or dengue-
induced LC3-II accumulation
The increase in mTORC2 activity during dengue infection and the requirement for
mTORC2 activity for maximal viral replication led us to investigate potential physiologic
functions of mTORC2 that are exploited by the virus. Because mTORC2 has been
shown to play a role in regulating cell proliferation (28), we considered the possibility
that our findings could be explained by altered cell growth rates. We assessed cell
growth rates in knockdown cells by measuring cell counts and CFSE dilution. No
significant differences in growth rates were observed upon comparison of mTORC2
knockdown cells with control cells (Fig. 3*A* and *B*). Because mTOR has also been
shown to play a role in the maintenance of cell morphology by modulating ion channel
activity and altering dynamics of the actin cytoskeleton (3, 4), we examined the
morphology of the mTORC2-depleted cells and their actin cytoskeleton. Cells treated
with shRNA targeting Rictor showed normal cellular morphology, normal cell spreading
and a morphologically normal actin cytoskeleton (Fig. 3*C*), suggesting that mTORC2
inactivation was not causing derangements in cell architecture in this setting.
Although autophagy is canonically regulated by mTORC1, we also considered the
possibility that inactivation of mTORC2 suppresses dengue replication by altering
autophagy through indirect regulatory effects on mTORC1. While the effect that dengue
has on the movement of cargo through the canonical autophagy pathway is unclear, it is
well known that dengue induces the accumulation of autophagosomes, characterized by
accumulation of lipidated LC3 protein (LC3-II) (10, 12, 13, 15). To assess effects of
mTORC2 inhibition on the pro-autophagic activity of dengue, we measured LC3-II
isoform levels in dengue-infected knockdown cells and observed similar degrees of
dengue-induced LC3-II accumulation (Fig. 3*D*), suggesting that mTORC2 inactivation
does not block the effects of dengue on autophagy.
173
mTORC2 inactivation causes increased dengue-mediated apoptosis and cell
death
We next asked whether mTORC2 could function as a pro-survival factor in infected
cells, given the known role for mTORC2 in regulating cell survival (28-30). We infected
Rictor knockdown or control cells with dengue for 24 or 36 h and, as a positive control
for apoptosis, treated the cells with staurosporine. We then asked whether apoptosis or
cell death was altered in the context of mTORC2 inhibition by measuring activated
caspase 3 expression and cell viability within infected and uninfected cells using flow
cytometry. At 24 hours post-infection (hpi), we observed a similar frequency of infected

cells when comparing control cells to Rictor knockdown cells, arguing against an early
block in viral replication in mTORC2-deficient cells. However, at 36 hpi, Rictor
knockdown was associated with a significant decrease in the proportion of infected cells
(Fig. 4*A* and *B*). Assessment of activated caspase 3 expression and cell viability at 24 h
revealed a small increase in apoptosis and cell death in mTORC2-deficient cells, which
were not statistically significant (Fig. 4*A*, *C*, and *D*). However, by 36 hpi, a marked
increase in apoptosis and cell death was observed in infected mTORC2 knockdown
cells, both of which were highly significant (4*A*, *C*, and *D*, $p < 0.001$ for activated
caspase 3 level, $p < 0.01$ for cell viability). Interestingly, while staurosporine treatment
increased cell death and apoptosis at similar levels in both scramble and Rictor
knockdown cells, suggesting that the increased apoptosis seen in dengue-infected cells
was specific for virus-induced apoptosis. We also measured the release of dengue from
infected cells and found that less infectious virus was released from Rictor knockdown
cells, and that the difference increased from 24 to 36 hpi, indicating that virus production
is diminished as apoptotic cell death increases.
198
DISCUSSION
In this study, we describe a role for host mTORC2 in dengue replication, demonstrating
that dengue can interact with the cellular mTOR signaling system, activating mTORC2
and thus promoting cell survival and permitting efficient viral replication. Specifically, the
data we present here suggest that mTORC2 plays a critical role in supporting viral
production by countering virus-induced apoptosis of the host cell. The induction of
apoptosis in dengue infection has been well-described, occurring in several cell types
both *in vitro* and *in vivo*, including endothelial cells, dendritic cells, and hepatocytes (31-
34). The finding of apoptotic cells in human autopsy specimens from severe dengue
cases has led to speculation that apoptosis contributes to pathogenesis in these cases
(32). While apoptosis may contribute to tissue damage and pathogenesis from the host
perspective, it is also an important mechanism for host control of viral infection,
triggering cell death before infectious progeny can be released (35) and shaping the
subsequent immune response. Since uncontrolled apoptosis would be detrimental to
viral replication, it is not surprising that dengue has evolved a mechanism to attenuate
the induction of apoptosis in infected cells. In the present study, we found that
decreasing mTORC2 signaling in infected cells leads to an increased frequency of
apoptosis in cell death, and the induction of apoptosis corresponds to a decrease in the
release of viral progeny from infected cells. Interestingly, the susceptibility of mTORC2-
deficient cells to apoptosis was specific for dengue infection in our experiments, as
neither baseline apoptosis nor apoptosis in response to staurosporine treatment
increased over control upon depletion of mTORC2. Furthermore, we found that dengue
infection triggers the activation of mTORC2, which likely represents a viral adaptation to
maintain cell survival during infection. While this strategy has not been described in
other viral infections, it has been reported as mechanism for cancer cell survival and
metastasis in several malignancies (36).
Dengue NS5 protein binds to mTOR, suggesting that the viral protein modulates
mTOR signaling during infection. NS5 appears to bind both mTORC1 and mTORC2,
evidenced by the co-immunoprecipitation of Raptor and Rictor with NS5. The molecular
consequences of NS5-mTOR interaction remain to be investigated. One possibility is

that NS5 binding to mTORC2 directly facilitates activation of the complex or acts as an
adaptor protein to stabilize interactions with downstream targets of the complex. Given
that there is crosstalk between mTORC1 and mTORC2 signaling (37), it is also possible
that interaction of NS5 with mTORC1 could activate mTORC2, either via de-repression
of mTORC2 or through alterations in mTOR protein stoichiometry.
The findings we present here also highlight the role of proteomic approaches in
understanding virus-host interactions. Much of the recent focus in dengue research has
utilized high-throughput genetic screens employing approaches such as RNAi and
CRISPR (38-40). While these genetic screens have identified host dependency factors,
because they rely on measuring viral gene expression or cell survival as the readout for
infection resistance, they are likely biased toward host factors involved in the early
stages of infection such as viral entry and gene expression as opposed to host factors
needed for late infection events such as viral release, maturation, and host cell survival.
For this reason, it is perhaps not surprising that components of the autophagy
machinery and mTOR signaling have not been consistently identified in these screens,
despite the roles of these host factors reported here and by others (14, 15). Moreover,
genetic screens stop short of identifying specific molecular interactions between virus
and host that mediate regulatory processes, which can be elucidated when interactions
between viral and host proteins are interrogated via proteomics. Nonetheless, it is
notable that Rictor inhibition was associated with decreased relative infection in 2 RNAi
libraries previously reported, although that finding did not meet the authors' criteria for
further validation (40). In contrast to genetic screens, proteomics methods can identify
host interactions at all stage of the viral life cycle, and subsequent validation can identify
important host interactions. When combined with quantitative approaches to distinguish
high-probability interactors from non-specific binding, proteomics approaches can
identify host factors with high validity (20).
The observation that mTORC2 is required for efficient dengue replication raises
the possibility of mTOR as a target for host-directed antiviral therapeutic development.
mTOR is a highly "druggable" target, and intense interest in the mTOR pathways in
oncology and neuropathology fields has spurred the development of multiple new small
molecule inhibitors with high specificity for mTOR (29, 41). In the solid organ transplant
infectious disease field, data have begun to emerge that certain mTOR inhibitors (used
as anti-organ rejection medications) may have anti-viral properties, reducing the risk of
some viral reactivation syndromes (42). In the case of dengue *per se*, candidate
compounds would likely need to have mTORC2 specificity, since mTORC1 inhibition
appears to enhance viral replication. While most newer generation mTOR inhibitors
have dual mTORC1/C2 specificity, a recently developed compound CID613034 has
been described that specifically inhibits mTORC2, demonstrating the feasibility of
specific mTORC2 inhibition (43). Since multiple biochemical steps occur between
mTORC2 assembly and the resultant anti-apoptotic outcome, components of the
mTORC2 signaling cascade might also provide useful targets for host-based
interventions.
The role of mTORC2 signaling in the pathogenesis of other viral infections
remains to be determined; however, there is evidence that other viruses including West
Nile, influenza and human cytomegalovirus may also stimulate mTORC2 signaling (16-
18). Whether mTORC2 is important for viral replication and serves as an anti-apoptotic

mechanism in those viruses remains to be determined, but it is possible that mTORC2
modulation is a common mechanism used by several viruses to counteract the host's
programmed cell death response. If that is the case, host-directed therapeutic
interventions targeting mTORC2 could be active against multiple viral pathogens.
279
MATERIALS AND METHODS
Antibodies and reagents – The following antibodies were obtained from Cell Signaling
Technology (Danvers, MA) and used at 1:1000 for western blotting: Raptor clone
24C10, Rictor clone D16H9, phospho-Rictor thr1135 clone D30A6, mTOR 7C10, AKT
clone 11E7, phospho-AKT ser473 clone D9E (used at 1:2000), p70 S6K clone 49D7,
phospho-p70 S6K thr389 clone 108D2, LC3A/B clone D3U4C, and B-actin clone 13E5.
The dengue 2 E protein antibody PA5-32246 was used at 1:10,000 for western blotting
(Thermo-Fisher Scientific, Waltham, MA). Secondary detection for western blotting used
anti-rabbit HRP antibody diluted 1:10,000 (Amersham ECL, GE Healthcare, Chicago,
IL). Polyclonal affinity purified NS3 and NS5 proteins were produced as previously
described (20). The pan-flavivirus E antibody 4G2 was prepared from hybridoma
supernatants and purified by protein A/G chromatography. For flow cytometry
experiments, 4G2 was conjugated to FITC according to the manufacturer's instructions
(FluoroTag kit, Sigma-Aldrich, St. Louis, MO). The dimerized GFP nanobody construct
LaG-16–G4S–LaG-2 (green lobster) was a gift from Michael Rout, and was prepared as
previously described (22). Phalloidin-Alexafluor 488 was obtained from Thermo-Fisher
scientific and used according to their instructions. Staurosporine was obtained from
Millipore Sigma (Burlington MA) and was used at 5 µM concentration.
298
Plasmids – The lentiviral shRNA plasmids pLKO.1 scramble, Raptor_2, mTOR_2, and
Rictor_2 were gifts from David Sabatini (Addgene plasmids 1864, 1858, 1856, and
1854). pMD2.G and psPAX2 were gifts from Didier Trono (Addgene plasmids 12259
and 12260). Dengue NS5-GFP fusion construct was generated by inserting the dengue
New Guinea C NS5 coding sequence from pDVW601 (44) into pACGFP1-N1 (Clontech)
as previously described (20).
305
Cell culture – HepG2 and Vero cells were cultured at 37 °C in 5% CO_2, in medium
composed of MEM supplemented with 10% FBS, 1× non-essential amino acids, 50
units/mL of penicillin and 50 µg/mL of streptomycin. Medium was replenished frequently
during experiments to avoid signaling changes caused by nutrient or growth factor
depletion. The *Aedes albopictus* derived cell line C6/36 was propagated at 28 °C in 5%
CO_2 in medium composed of MEM supplemented with 10% FBS, 1× non-essential
amino acids, 50 units/mL of penicillin and 50 µg/mL of streptomycin. The B lymphocyte
hybridoma cell line D1-4G2-4-15 was obtained from the ATCC (Manassas, VA), and
was maintained in ATCC Hybri-Care Medium supplemented with 10% FBS and 1.5 g/L
sodium bicarbonate.
316
317   *Transfections* – 293FT cells were plated to 70-80% density in antibiotic free medium.
318   Transfection complexes were prepared by mixing plasmid DNA with polyethyleneimine
319   (PEI Max 40k, Polysciences Inc., Warrington, PA) in a 1:4 mass ratio. After 15 min

320 incubation at room temperature DNA complexes were added dropwise to the cell
321 cultures.
322
Affinity capture – To prepare GFP affinity-capture beads, the dimerized GFP nanobody
LaG-16–G4S–LaG-2 (green lobster) was covalently linked to magnetic beads
(Dynabeads M-270 epoxy Thermo Fisher Scientific) as previously described (45).
Briefly, 5 µg of nanobody were used per 1 mg of Dynabeads, with conjugations carried
out in 0.1 M sodium phosphate buffer and 1 M ammonium sulfate, with an 18- to 20-h
incubation at 30 °C. Beads were then washed sequentially with 0.1 M sodium
phosphate buffer, 100 mM glycine pH 2.5, 10 mM Tris-HCl pH 8.8, 100 mM
triethylamine, 1× PBS (4 times), PBS + 0.5% Triton X-100, and 1× PBS. For affinity
capture experiments, cells were harvested 48 h after transfection. Cells were washed
with ice cold PBS and then lysed with 1% Triton X-100, 0.5% sodium deoxycholate, 110
mM potassium acetate pH 7.5, 20 mM HEPES, 2 mM $MgCl_2$, 25 mM NaCl, and 1×
protease/phosphatase inhibitor cocktail (Cell Signaling Technology). Lysates were
clarified by centrifugation for 10 min at 13,000 × g at 4 °C. Affinity capture beads were
immediately added to the clarified lysate and incubated for 10 min at room temperature
with rotation. Beads were then washed 3× with lysis buffer, and bound protein
complexes eluted with 1.1× LDS sample buffer for 10 min at 70 °C. For SDS-PAGE and
western blot analysis, 10× reducing agent (Thermo-Fisher Scientific) was added and
samples were heated for an additional 10 min at 70 °C. SDS-PAGE and western blot
analysis were performed as described below. Gel staining was performed using Sypro
Ruby fluorescent gel stain (Thermo-Fisher Scientific) according to the manufacturer's
instructions.
344
shRNA-mediated gene silencing – To generate lentiviral vector stocks, shRNA
constructs were cotransfected with pMD2.G and psPAX2 into HEK 293T cells.
Supernatants were harvested, passed through 0.45 µm filters, layered on 20% sucrose
cushions, and centrifuged at 100,000 × g for 4 h at 4 °C. Lentiviral pellets were
resuspended in OptiMEM and stored at -80 °C until use. For lentiviral transductions,
viral stocks were diluted to the desired concentration with OptiMEM and 0.8 µg/mL
polybrene and added to cells. At 48 h post-transduction, cells containing stably
integrated constructs were selected using 2 µg/mL puromycin. Experiments were
performed on cell lines that were maintained and passaged for no more than 3 weeks
before discarding and establishing fresh cell lines.
355
Virus and infections – DENV-2 MON601, a molecular clone of DENV-2 New Guinea
strain C[46], was generated by transfection of *in vitro*-transcribed RNA into Vero cells,
followed by no more than 5 passages in C6/36 cells. Virus was propagated by infecting
80% confluent C6/36 monolayers with low-passage stock virus at a MOI of 0.01, and
harvesting infectious supernatants 5-7 days post-infection. Infectious supernatants were
cleared of cellular debris by centrifugation and stored at -80 °C until use. Virus stocks
and experimental infectious supernatants were titrated using a flow cytometry approach
which has been described elsewhere (46). Briefly, serially diluted virus stocks were
used to infected Vero cells in a multi-well plate. Cells were harvested 20-24 h post-
infection, fixed and permeabilized, and stained with 4G2-FITC. The percentage of

infected cells was then used to calculate the number of fluorescence forming units
(FFU) per milliliter of inoculum. For experimental infections, virus was diluted in
OptiMEM to the desired MOI and incubated on cells for 90 min at 37 °C. Virus was then
removed, the cells washed, and complete growth medium added.
370
Western blot analysis – Cells were placed on ice and washed with ice-cold PBS. Cells
were then collected and lysed with Triton X-100 lysis buffer (1% Triton X-100, 120 mM
NaCl, 1 mM EDTA, 40 mM HEPES pH 7.4, and 1× protease and phosphatase inhibitor
cocktail (Cell Signaling Technology)). To prepare whole cell extracts, SDS lysis buffer
(2% SDS, 50 mM Tris pH 7.4, 5% glycerol, 5 mM EDTA, 1 mM NaF, 1 mM DTT, and 1×
phosphatase & protease inhibitor cocktail) was used instead of Triton X-100 lysis buffer.
Protein concentration was determined using BCA assay and a BSA standard curve, and
equivalent amounts of protein were mixed with 4× LDS sample buffer and 10× reducing
agent (Thermo-Fisher Scientific), followed by denaturation at 70 °C for 10 min. Proteins
were then resolved on 4-12% Bis-Tris gels (for lower molecular weight proteins) or 3-8%
Tris-Acetate gels (for higher molecular weight proteins) (NuPage, Thermo-Fisher
Scientific) and run in MOPS or Tris-Acetate running buffer respectively. Proteins were
transferred to PVDF membrane and blocked in 5% milk/TBS-T for 1-2 h. Primary
antibodies were diluted in 5% BSA/TBS-T, and incubated overnight at 4 °C. Membranes
were washed and incubated with anti-rabbit HRP antibody for 1-2 h at room
temperature. Membranes were then washed with TBS-T, exposed to chemiluminescent
substrate, and imaged using a digital CCD platform (Fluorchem E, Protein Simple, San
Jose, CA). Band densitometry was performed using ImageJ software.
389
390   *Fluorescence microscopy* – Cells were fixed with 4% paraformaldehyde/PBS for 15 min
391   at room temperature, permeabilized with 0.1% Triton X-100/PBS for 10 min at room
392   temperature and blocked with 5% normal goat serum in 0.05% Tween-20/PBS. Cells
393   were then stained with phalloidin-Alexafluor 488 (Thermo Fisher Scientific) for 1 h at
394   room temperature and mounted using medium containing DAPI.
395
Flow cytometry – For viability analysis, cells were trypsinized, washed with PBS, and
stained with a cell impermeable amine reactive dye (LIVE/DEAD Violet A.K.A. LD405,
ThermoFisher Scientific) according to the manufacturer's instructions. Cells were fixed
and permeabilized using the BD Cytofix/Cytoperm kit according to the manufacturer's
instructions (BD Biosciences). Permeabilized cells were stained with fluorophore-
conjugated antibodies as indicated in the text. Cells were analyzed on a BD LSR-II
cytometer, and data were analyzed using FlowJo software.
403

ACKNOWLEDGMENTS

F.D.M was supported by a postdoctoral fellowship of the Canadian Institutes for Health Research. This study was supported by grant R01GM101183 to A.K. and grants R21AI124266 and P41GM109824 to J.D.A. from the National Institutes of Health.

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest with the contents of this article.

REFERENCES

1.	Laplante, M., and Sabatini, D. M. (2009) mTOR signaling at a glance. *J Cell Sci* **122**, 3589-3594
2.	Laplante, M., and Sabatini, D. M. (2012) mTOR signaling in growth control and disease. *Cell* **149**, 274-293
3.	Jacinto, E., Loewith, R., Schmidt, A., Lin, S., Ruegg, M. A., Hall, A., and Hall, M. N. (2004) Mammalian TOR complex 2 controls the actin cytoskeleton and is rapamycin insensitive. *Nat Cell Biol* **6**, 1122-1128
4.	Sarbassov, D. D., Ali, S. M., Kim, D. H., Guertin, D. A., Latek, R. R., Erdjument-Bromage, H., Tempst, P., and Sabatini, D. M. (2004) Rictor, a novel binding partner of mTOR, defines a rapamycin-insensitive and raptor-independent pathway that regulates the cytoskeleton. *Curr Biol* **14**, 1296-1302
5.	Zinzalla, V., Stracka, D., Oppliger, W., and Hall, M. N. (2011) Activation of mTORC2 by association with the ribosome. *Cell* **144**, 757-768
6.	Arias, E., Koga, H., Diaz, A., Mocholi, E., Patel, B., and Cuervo, A. M. (2015) Lysosomal mTORC2/PHLPP1/Akt Regulate Chaperone-Mediated Autophagy. *Mol Cell* **59**, 270-284
7.	Lampada, A., O'Prey, J., Szabadkai, G., Ryan, K. M., Hochhauser, D., and Salomoni, P. (2017) mTORC1-independent autophagy regulates receptor tyrosine kinase phosphorylation in colorectal cancer cells via an mTORC2-mediated mechanism. *Cell Death Differ* **24**, 1045-1062
8.	Buchkovich, N. J., Yu, Y., Zampieri, C. A., and Alwine, J. C. (2008) The TORrid affairs of viruses: effects of mammalian DNA viruses on the PI3K-Akt-mTOR signalling pathway. *Nat Rev Microbiol* **6**, 266-275
9.	Le Sage, V., Cinti, A., Amorim, R., and Mouland, A. J. (2016) Adapting the Stress Response: Viral Subversion of the mTOR Signaling Pathway. *Viruses* **8**
10.	Lee, Y. R., Lei, H. Y., Liu, M. T., Wang, J. R., Chen, S. H., Jiang-Shieh, Y. F., Lin, Y. S., Yeh, T. M., Liu, C. C., and Liu, H. S. (2008) Autophagic machinery activated by dengue virus enhances virus replication. *Virology* **374**, 240-248
11.	Lee, Y. R., Hu, H. Y., Kuo, S. H., Lei, H. Y., Lin, Y. S., Yeh, T. M., Liu, C. C., and Liu, H. S. (2013) Dengue virus infection induces autophagy: an in vivo study. *J Biomed Sci* **20**, 65
12.	Heaton, N. S., and Randall, G. (2010) Dengue virus-induced autophagy regulates lipid metabolism. *Cell Host Microbe* **8**, 422-432
13.	Metz, P., Chiramel, A., Chatel-Chaix, L., Alvisi, G., Bankhead, P., Mora-Rodriguez, R., Long, G., Hamacher-Brady, A., Brady, N. R., and Bartenschlager,

R. (2015) Dengue Virus Inhibition of Autophagic Flux and Dependency of Viral Replication on Proteasomal Degradation of the Autophagy Receptor p62. *J Virol* **89**, 8026-8041

14. Mateo, R., Nagamine, C. M., Spagnolo, J., Mendez, E., Rahe, M., Gale, M., Jr., Yuan, J., and Kirkegaard, K. (2013) Inhibition of cellular autophagy deranges dengue virion maturation. *J Virol* **87**, 1312-1321

15. Jordan, T. X., and Randall, G. (2017) Dengue Virus Activates the AMP Kinase-mTOR Axis To Stimulate a Proviral Lipophagy. *J Virol* **91**

16. Kuss-Duerkop, S. K., Wang, J., Mena, I., White, K., Metreveli, G., Sakthivel, R., Mata, M. A., Munoz-Moreno, R., Chen, X., Krammer, F., Diamond, M. S., Chen, Z. J., Garcia-Sastre, A., and Fontoura, B. M. A. (2017) Influenza virus differentially activates mTORC1 and mTORC2 signaling to maximize late stage replication. *PLoS Pathog* **13**, e1006635

17. Shives, K. D., Beatman, E. L., Chamanian, M., O'Brien, C., Hobson-Peters, J., and Beckham, J. D. (2014) West nile virus-induced activation of mammalian target of rapamycin complex 1 supports viral growth and viral protein expression. *J Virol* **88**, 9458-9471

18. Kudchodkar, S. B., Yu, Y., Maguire, T. G., and Alwine, J. C. (2006) Human cytomegalovirus infection alters the substrate specificities and rapamycin sensitivities of raptor- and rictor-containing complexes. *Proc Natl Acad Sci U S A* **103**, 14182-14187

19. Tackett, A. J., DeGrasse, J. A., Sekedat, M. D., Oeffinger, M., Rout, M. P., and Chait, B. T. (2005) I-DIRT, a general method for distinguishing between specific and nonspecific protein interactions. *J Proteome Res* **4**, 1752-1756

20. Carpp, L. N., Rogers, R. S., Moritz, R. L., and Aitchison, J. D. (2014) Quantitative proteomic analysis of host-virus interactions reveals a role for Golgi brefeldin A resistance factor 1 (GBF1) in dengue infection. *Mol Cell Proteomics* **13**, 2836-2854

21. Kumar, A., Buhler, S., Selisko, B., Davidson, A., Mulder, K., Canard, B., Miller, S., and Bartenschlager, R. (2013) Nuclear localization of dengue virus nonstructural protein 5 does not strictly correlate with efficient viral RNA replication and inhibition of type I interferon signaling. *J Virol* **87**, 4545-4557

22. Fridy, P. C., Li, Y., Keegan, S., Thompson, M. K., Nudelman, I., Scheid, J. F., Oeffinger, M., Nussenzweig, M. C., Fenyo, D., Chait, B. T., and Rout, M. P. (2014) A robust pipeline for rapid production of versatile nanobody repertoires. *Nat Methods* **11**, 1253-1260

23. Sarbassov, D. D., Guertin, D. A., Ali, S. M., and Sabatini, D. M. (2005) Phosphorylation and regulation of Akt/PKB by the rictor-mTOR complex. *Science* **307**, 1098-1101

24. Thepparit, C., Khakpoor, A., Khongwichit, S., Wikan, N., Fongsaran, C., Chingsuwanrote, P., Panraksa, P., and Smith, D. R. (2013) Dengue 2 infection of HepG2 liver cells results in endoplasmic reticulum stress and induction of multiple pathways of cell death. *BMC Res Notes* **6**, 372

25. Suksanpaisan, L., Cabrera-Hernandez, A., and Smith, D. R. (2007) Infection of human primary hepatocytes with dengue virus serotype 2. *J Med Virol* **79**, 300-307

26. Burnett, P. E., Barrow, R. K., Cohen, N. A., Snyder, S. H., and Sabatini, D. M. (1998) RAFT1 phosphorylation of the translational regulators p70 S6 kinase and 4E-BP1. *Proc Natl Acad Sci U S A* **95**, 1432-1437

27. Julien, L. A., Carriere, A., Moreau, J., and Roux, P. P. (2010) mTORC1-activated S6K1 phosphorylates Rictor on threonine 1135 and regulates mTORC2 signaling. *Mol Cell Biol* **30**, 908-921

28. Oh, W. J., and Jacinto, E. (2011) mTOR complex 2 signaling and functions. *Cell Cycle* **10**, 2305-2316

29. Zou, Z., Chen, J., Yang, J., and Bai, X. (2016) Targeted Inhibition of Rictor/mTORC2 in Cancer Treatment: A New Era after Rapamycin. *Curr Cancer Drug Targets* **16**, 288-304

30. Lamming, D. W., Demirkan, G., Boylan, J. M., Mihaylova, M. M., Peng, T., Ferreira, J., Neretti, N., Salomon, A., Sabatini, D. M., and Gruppuso, P. A. (2014) Hepatic signaling by the mechanistic target of rapamycin complex 2 (mTORC2). *FASEB J* **28**, 300-315

31. Torrentes-Carvalho, A., Azeredo, E. L., Reis, S. R., Miranda, A. S., Gandini, M., Barbosa, L. S., and Kubelka, C. F. (2009) Dengue-2 infection and the induction of apoptosis in human primary monocytes. *Mem Inst Oswaldo Cruz* **104**, 1091-1099

32. Limonta, D., Capo, V., Torres, G., Perez, A. B., and Guzman, M. G. (2007) Apoptosis in tissues from fatal dengue shock syndrome. *J Clin Virol* **40**, 50-54

33. Lin, J. C., Lin, S. C., Chen, W. Y., Yen, Y. T., Lai, C. W., Tao, M. H., Lin, Y. L., Miaw, S. C., and Wu-Hsieh, B. A. (2014) Dengue viral protease interaction with NF-kappaB inhibitor alpha/beta results in endothelial cell apoptosis and hemorrhage development. *J Immunol* **193**, 1258-1267

34. Martins Sde, T., Silveira, G. F., Alves, L. R., Duarte dos Santos, C. N., and Bordignon, J. (2012) Dendritic cell apoptosis and the pathogenesis of dengue. *Viruses* **4**, 2736-2753

35. Orzalli, M. H., and Kagan, J. C. (2017) Apoptosis and Necroptosis as Host Defense Strategies to Prevent Viral Infection. *Trends Cell Biol* **27**, 800-809

36. Kim, L. C., Cook, R. S., and Chen, J. (2017) mTORC1 and mTORC2 in cancer and the tumor microenvironment. *Oncogene* **36**, 2191-2201

37. Tyakht, A. V., Ilina, E. N., Alexeev, D. G., Ischenko, D. S., Gorbachev, A. Y., Semashko, T. A., Larin, A. K., Selezneva, O. V., Kostryukova, E. S., Karalkin, P. A., Vakhrushev, I. V., Kurbatov, L. K., Archakov, A. I., and Govorun, V. M. (2014) RNA-Seq gene expression profiling of HepG2 cells: the influence of experimental factors and comparison with liver tissue. *BMC Genomics* **15**, 1108

38. Sessions, O. M., Barrows, N. J., Souza-Neto, J. A., Robinson, T. J., Hershey, C. L., Rodgers, M. A., Ramirez, J. L., Dimopoulos, G., Yang, P. L., Pearson, J. L., and Garcia-Blanco, M. A. (2009) Discovery of insect and human dengue virus host factors. *Nature* **458**, 1047-1050

39. Marceau, C. D., Puschnik, A. S., Majzoub, K., Ooi, Y. S., Brewer, S. M., Fuchs, G., Swaminathan, K., Mata, M. A., Elias, J. E., Sarnow, P., and Carette, J. E. (2016) Genetic dissection of Flaviviridae host factors through genome-scale CRISPR screens. *Nature* **535**, 159-163

40. Savidis, G., McDougall, W. M., Meraner, P., Perreira, J. M., Portmann, J. M., Trincucci, G., John, S. P., Aker, A. M., Renzette, N., Robbins, D. R., Guo, Z.,

Green, S., Kowalik, T. F., and Brass, A. L. (2016) Identification of Zika Virus and
Dengue Virus Dependency Factors using Functional Genomics. *Cell Rep* **16**, 232-
246
41. Xie, J., Wang, X., and Proud, C. G. (2016) mTOR inhibitors in cancer therapy.
F1000Res **5**
42. Pascual, J., Royuela, A., Fernandez, A. M., Herrero, I., Delgado, J. F., Sole, A.,
Guirado, L., Serrano, T., de la Torre-Cisneros, J., Moreno, A., Cordero, E.,
Gallego, R., Lumbreras, C., Aguado, J. M., Spanish Society of Transplantation,
V., and Immune Response Investigation Study, G. (2016) Role of mTOR inhibitors
for the control of viral infection in solid organ transplant recipients. *Transpl Infect*
Dis **18**, 819-831
43. Benavides-Serrato, A., Lee, J., Holmes, B., Landon, K. A., Bashir, T., Jung, M. E.,
Lichtenstein, A., and Gera, J. (2017) Specific blockade of Rictor-mTOR
association inhibits mTORC2 activity and is cytotoxic in glioblastoma. *PLoS One*
12, e0176599
44. Pryor, M. J., Carr, J. M., Hocking, H., Davidson, A. D., Li, P., and Wright, P. J.
(2001) Replication of dengue virus type 2 in human monocyte-derived
macrophages: comparisons of isolates and recombinant viruses with substitutions
at amino acid 390 in the envelope glycoprotein. *Am J Trop Med Hyg* **65**, 427-434
45. Cristea, I. M., and Chait, B. T. (2011) Conjugation of magnetic beads for
immunopurification of protein complexes. *Cold Spring Harb Protoc* **2011**, pdb
prot5610
46. Lambeth, C. R., White, L. J., Johnston, R. E., and de Silva, A. M. (2005) Flow
cytometry-based assay for titrating dengue virus. *J Clin Microbiol* **43**, 3267-3272

FIGURE LEGENDS

FIGURE 1**. Dengue NS5 interacts with mTORC1 and mTORC2**. *A,* Expression of
NS5-GFP fusion protein. 293FT cells were transfected with expression plasmids
encoding GFP or NS5-GFP fusion protein, and live cells were imaged by fluorescence
microscopy. Bar = 100 μm. *B*, Cells were transfected with GFP or NS5-GFP plasmids
as in (*A*), and lysates were affinity captured using anti-GFP nanobodies. 50% of the
eluates or 1% of the input lysate were then analyzed by western blot analysis with the
indicated antibodies.

FIGURE 2. **Dengue infection activates mTORC2 signaling and is required for**
maximal viral replication. *A*, Selective inactivation of mTORC1 and mTORC2 in
HepG2 cells. Cells were transduced with lentivirus encoding shRNA directed against
mTOR protein, Raptor, or Rictor, or with a non-specific scramble sequence, and then
selected with puromycin. Lysates were analyzed by western blot with the indicated
antibodies. *B*, mTORC2 inactivation diminishes dengue replication. Cells were
transduced with lentiviral shRNA vectors as in (*A*) and were then infected with dengue
MON601 at a MOI of 1. 24 h later cell culture supernatants were collected and titrated
on Vero cells. The data represent scramble-normalized values from 4 independent
experiments. *C*, mTORC2 is activated during dengue infection. HepG2 cells were
infected with dengue at MOI of 4. Cells were collected at 36 h post-infection and
analyzed by western blot with the indicated antibodies. The bar graph shows the ratio of
phospho-AKT to total AKT band intensity averaged from 7 independent experiments. *D*,
Rictor knockdown abrogates mTORC2 activation by dengue. Cells were transduced
with shRNA encoding lentivirus as in (*A*) and were infected with dengue at MOI of 4.
Cells were collected at 36 hpi and were analyzed by western blot with the indicated
antibodies. Error bars are one standard deviation. p values are derived from 2-tailed
Student's t test; n.s. denotes $p > 0.05$, * denotes $p < 0.05$, ** denotes $p < 0.01$, and ***
denotes $p < 0.001$.

FIGURE 3. **mTORC2 inactivation does not alter HepG2 cell morphology,**
cytoskeletal architecture, growth rate, or LC3 lipidation in response to dengue
infection. *A*, HepG2 cells were transduced with lentiviral vectors encoding shRNA
directed against mTOR protein, Raptor or Rictor, or with a non-specific scramble
sequence and were selected with puromycin. Cells were imaged by light microscopy
(top panels) or stained with fluorophore-labeled phalloidin and imaged with fluorescence
microscopy (bottom panels). Bar = 100 μm *B*, Cells were transduced with shRNA
lentiviral vectors as in (*A*) and were plated at equal densities. Cells were then counted
using an automated cell counter 4 days later. Data represent cell counts from 4
separate cultures. *C*, Cells were transduced with shRNA lentiviral vectors as in (*A*) and
were plated at equal densities. Cells were then loaded with CFSE and analyzed by flow
cytometry 24 h (open histograms) and 96 h (shaded histograms) later. Bar graph
represents the ratio of CFSE mean fluorescence intensity between the 24 and 96 h
timepoint, averaged from 4 separate cultures. *D*, mTORC2 inhibition does not alter
dengue-mediate LC3-II lipidation. Cells were transduced with shRNA lentiviral vectors
as in (*A*) and were infected with dengue at MOI 4. Cells were harvested at 36 h post-
infection and lysates were analyzed by western blot with the indicated antibodies. The

bar graph shows the ratio of LC3-II band intensity between dengue infected and mock
treated cells from 3 independent experiments. Error bars are one standard deviation. p
values are derived from 2-tailed Student's t test; n.s. denotes $p > 0.05$, * denotes $p <$
0.05, ** denotes $p < 0.01$, and *** denotes $p < 0.001$.

FIGURE 4. **mTORC2 inhibition leads to increased dengue-induced apoptosis and**
cell death. *A*, HepG2 cells were infected with dengue at MOI of 4 for 24 or 36 h, mock
treated, or treated with 5 µM staurosporine overnight. Cells were then collected and
stained with a cell-impermeable amine reactive dye (LD405), followed by fixation,
permeabilization, and staining with anti-activated caspase 3 and anti-flavivirus E protein
antibodies. Infected (E-positive) cells were gated and activated caspase 3 (CC3) and
LD405 expression were analyzed within that population. Numbers represent population
frequencies in each gate/quadrant. *B*, Cells were treated as in (*A*), and the percent of
infected (E-positive) cells was calculated. The bar graph shows the average values for 3
independent experiments, each normalized to the scramble condition. *C*, Cells were
treated as in (*A*), and the percent of infected cells expressing activated caspase 3 was
calculated. The bar graph represents Z-normalized average values from 3 independent
experiments. *D*, Cells were treated as in (*A*), and the percent of infected cells positive
for LD405 was calculated. The bar graph represents *Z*-normalized values from 3
independent experiments. p values are derived from 2-tailed Student's t test; n.s.
denotes $p > 0.05$, * denotes $p < 0.05$, ** denotes $p < 0.01$, and *** denotes $p < 0.001$.

REFERENCES

1. Laplante M, Sabatini DM. 2009. mTOR signaling at a glance. J Cell Sci 122:3589-94.
2. Laplante M, Sabatini DM. 2012. mTOR signaling in growth control and disease. Cell 149:274-93.
3. Jacinto E, Loewith R, Schmidt A, Lin S, Ruegg MA, Hall A, Hall MN. 2004. Mammalian TOR complex 2 controls the actin cytoskeleton and is rapamycin insensitive. Nat Cell Biol 6:1122-8.
4. Sarbassov DD, Ali SM, Kim DH, Guertin DA, Latek RR, Erdjument-Bromage H, Tempst P, Sabatini DM. 2004. Rictor, a novel binding partner of mTOR, defines a rapamycin-insensitive and raptor-independent pathway that regulates the cytoskeleton. Curr Biol 14:1296-302.
5. Zinzalla V, Stracka D, Oppliger W, Hall MN. 2011. Activation of mTORC2 by association with the ribosome. Cell 144:757-68.
6. Arias E, Koga H, Diaz A, Mocholi E, Patel B, Cuervo AM. 2015. Lysosomal mTORC2/PHLPP1/Akt Regulate Chaperone-Mediated Autophagy. Mol Cell 59:270-84.
7. Lampada A, O'Prey J, Szabadkai G, Ryan KM, Hochhauser D, Salomoni P. 2017. mTORC1-independent autophagy regulates receptor tyrosine kinase phosphorylation in colorectal cancer cells via an mTORC2-mediated mechanism. Cell Death Differ 24:1045-1062.
8. Buchkovich NJ, Yu Y, Zampieri CA, Alwine JC. 2008. The TORrid affairs of viruses: effects of mammalian DNA viruses on the PI3K-Akt-mTOR signalling pathway. Nat Rev Microbiol 6:266-75.
9. Le Sage V, Cinti A, Amorim R, Mouland AJ. 2016. Adapting the Stress Response: Viral Subversion of the mTOR Signaling Pathway. Viruses 8.
10. Lee YR, Lei HY, Liu MT, Wang JR, Chen SH, Jiang-Shieh YF, Lin YS, Yeh TM, Liu CC, Liu HS. 2008. Autophagic machinery activated by dengue virus enhances virus replication. Virology 374:240-8.
11. Lee YR, Hu HY, Kuo SH, Lei HY, Lin YS, Yeh TM, Liu CC, Liu HS. 2013. Dengue virus infection induces autophagy: an in vivo study. J Biomed Sci 20:65.
12. Heaton NS, Randall G. 2010. Dengue virus-induced autophagy regulates lipid metabolism. Cell Host Microbe 8:422-32.
13. Metz P, Chiramel A, Chatel-Chaix L, Alvisi G, Bankhead P, Mora-Rodriguez R, Long G, Hamacher-Brady A, Brady NR, Bartenschlager R. 2015. Dengue Virus Inhibition of Autophagic Flux and Dependency of Viral Replication on Proteasomal Degradation of the Autophagy Receptor p62. J Virol 89:8026-41.
14. Mateo R, Nagamine CM, Spagnolo J, Mendez E, Rahe M, Gale M, Jr., Yuan J, Kirkegaard K. 2013. Inhibition of cellular autophagy deranges dengue virion maturation. J Virol 87:1312-21.
15. Jordan TX, Randall G. 2017. Dengue Virus Activates the AMP Kinase-mTOR Axis To Stimulate a Proviral Lipophagy. J Virol 91.
16. Kuss-Duerkop SK, Wang J, Mena I, White K, Metreveli G, Sakthivel R, Mata MA, Munoz-Moreno R, Chen X, Krammer F, Diamond MS, Chen ZJ, Garcia-Sastre A, Fontoura BMA. 2017. Influenza virus differentially activates mTORC1 and

mTORC2 signaling to maximize late stage replication. PLoS Pathog
13:e1006635.
17.   Shives KD, Beatman EL, Chamanian M, O'Brien C, Hobson-Peters J, Beckham
JD. 2014. West nile virus-induced activation of mammalian target of rapamycin
complex 1 supports viral growth and viral protein expression. J Virol 88:9458-71.
18.   Kudchodkar SB, Yu Y, Maguire TG, Alwine JC. 2006. Human cytomegalovirus
infection alters the substrate specificities and rapamycin sensitivities of raptor-
and rictor-containing complexes. Proc Natl Acad Sci U S A 103:14182-7.
19.   Tackett AJ, DeGrasse JA, Sekedat MD, Oeffinger M, Rout MP, Chait BT. 2005. I-
DIRT, a general method for distinguishing between specific and nonspecific
protein interactions. J Proteome Res 4:1752-6.
20.   Carpp LN, Rogers RS, Moritz RL, Aitchison JD. 2014. Quantitative proteomic
analysis of host-virus interactions reveals a role for Golgi brefeldin A resistance
factor 1 (GBF1) in dengue infection. Mol Cell Proteomics 13:2836-54.
21.   Kumar A, Buhler S, Selisko B, Davidson A, Mulder K, Canard B, Miller S,
Bartenschlager R. 2013. Nuclear localization of dengue virus nonstructural
protein 5 does not strictly correlate with efficient viral RNA replication and
inhibition of type I interferon signaling. J Virol 87:4545-57.
22.   Fridy PC, Li Y, Keegan S, Thompson MK, Nudelman I, Scheid JF, Oeffinger M,
Nussenzweig MC, Fenyo D, Chait BT, Rout MP. 2014. A robust pipeline for rapid
production of versatile nanobody repertoires. Nat Methods 11:1253-60.
23.   Sarbassov DD, Guertin DA, Ali SM, Sabatini DM. 2005. Phosphorylation and
regulation of Akt/PKB by the rictor-mTOR complex. Science 307:1098-101.
24.   Thepparit C, Khakpoor A, Khongwichit S, Wikan N, Fongsaran C,
Chingsuwanrote P, Panraksa P, Smith DR. 2013. Dengue 2 infection of HepG2
liver cells results in endoplasmic reticulum stress and induction of multiple
pathways of cell death. BMC Res Notes 6:372.
25.   Suksanpaisan L, Cabrera-Hernandez A, Smith DR. 2007. Infection of human
primary hepatocytes with dengue virus serotype 2. J Med Virol 79:300-7.
26.   Burnett PE, Barrow RK, Cohen NA, Snyder SH, Sabatini DM. 1998. RAFT1
phosphorylation of the translational regulators p70 S6 kinase and 4E-BP1. Proc
Natl Acad Sci U S A 95:1432-7.
27.   Julien LA, Carriere A, Moreau J, Roux PP. 2010. mTORC1-activated S6K1
phosphorylates Rictor on threonine 1135 and regulates mTORC2 signaling. Mol
Cell Biol 30:908-21.
28.   Oh WJ, Jacinto E. 2011. mTOR complex 2 signaling and functions. Cell Cycle
10:2305-16.
29.   Zou Z, Chen J, Yang J, Bai X. 2016. Targeted Inhibition of Rictor/mTORC2 in
Cancer Treatment: A New Era after Rapamycin. Curr Cancer Drug Targets
16:288-304.
30.   Lamming DW, Demirkan G, Boylan JM, Mihaylova MM, Peng T, Ferreira J,
Neretti N, Salomon A, Sabatini DM, Gruppuso PA. 2014. Hepatic signaling by the
mechanistic target of rapamycin complex 2 (mTORC2). FASEB J 28:300-15.
31.   Torrentes-Carvalho A, Azeredo EL, Reis SR, Miranda AS, Gandini M, Barbosa
LS, Kubelka CF. 2009. Dengue-2 infection and the induction of apoptosis in
human primary monocytes. Mem Inst Oswaldo Cruz 104:1091-9.

32.   Limonta D, Capo V, Torres G, Perez AB, Guzman MG. 2007. Apoptosis in
721   tissues from fatal dengue shock syndrome. J Clin Virol 40:50-4.
722   33.   Lin JC, Lin SC, Chen WY, Yen YT, Lai CW, Tao MH, Lin YL, Miaw SC, Wu-Hsieh
723   BA. 2014. Dengue viral protease interaction with NF-kappaB inhibitor alpha/beta
724   results in endothelial cell apoptosis and hemorrhage development. J Immunol
725   193:1258-67.
726   34.   Martins Sde T, Silveira GF, Alves LR, Duarte dos Santos CN, Bordignon J. 2012.
727   Dendritic cell apoptosis and the pathogenesis of dengue. Viruses 4:2736-53.
728   35.   Orzalli MH, Kagan JC. 2017. Apoptosis and Necroptosis as Host Defense
729   Strategies to Prevent Viral Infection. Trends Cell Biol 27:800-809.
730   36.   Kim LC, Cook RS, Chen J. 2017. mTORC1 and mTORC2 in cancer and the
731   tumor microenvironment. Oncogene 36:2191-2201.
732   37.   Tyakht AV, Ilina EN, Alexeev DG, Ischenko DS, Gorbachev AY, Semashko TA,
733   Larin AK, Selezneva OV, Kostryukova ES, Karalkin PA, Vakhrushev IV, Kurbatov
734   LK, Archakov AI, Govorun VM. 2014. RNA-Seq gene expression profiling of
735   HepG2 cells: the influence of experimental factors and comparison with liver
736   tissue. BMC Genomics 15:1108.
737   38.   Sessions OM, Barrows NJ, Souza-Neto JA, Robinson TJ, Hershey CL, Rodgers
738   MA, Ramirez JL, Dimopoulos G, Yang PL, Pearson JL, Garcia-Blanco MA. 2009.
739   Discovery of insect and human dengue virus host factors. Nature 458:1047-50.
740   39.   Marceau CD, Puschnik AS, Majzoub K, Ooi YS, Brewer SM, Fuchs G,
741   Swaminathan K, Mata MA, Elias JE, Sarnow P, Carette JE. 2016. Genetic
742   dissection of Flaviviridae host factors through genome-scale CRISPR screens.
743   Nature 535:159-63.
744   40.   Savidis G, McDougall WM, Meraner P, Perreira JM, Portmann JM, Trincucci G,
745   John SP, Aker AM, Renzette N, Robbins DR, Guo Z, Green S, Kowalik TF, Brass
746   AL. 2016. Identification of Zika Virus and Dengue Virus Dependency Factors
747   using Functional Genomics. Cell Rep 16:232-246.
748   41.   Xie J, Wang X, Proud CG. 2016. mTOR inhibitors in cancer therapy. F1000Res
749   5.
750   42.   Pascual J, Royuela A, Fernandez AM, Herrero I, Delgado JF, Sole A, Guirado L,
751   Serrano T, de la Torre-Cisneros J, Moreno A, Cordero E, Gallego R, Lumbreras
752   C, Aguado JM, Spanish Society of Transplantation V, Immune Response
753   Investigation Study G. 2016. Role of mTOR inhibitors for the control of viral
754   infection in solid organ transplant recipients. Transpl Infect Dis 18:819-831.
755   43.   Benavides-Serrato A, Lee J, Holmes B, Landon KA, Bashir T, Jung ME,
756   Lichtenstein A, Gera J. 2017. Specific blockade of Rictor-mTOR association
757   inhibits mTORC2 activity and is cytotoxic in glioblastoma. PLoS One
758   12:e0176599.
759   44.   Pryor MJ, Carr JM, Hocking H, Davidson AD, Li P, Wright PJ. 2001. Replication
760   of dengue virus type 2 in human monocyte-derived macrophages: comparisons
761   of isolates and recombinant viruses with substitutions at amino acid 390 in the
762   envelope glycoprotein. Am J Trop Med Hyg 65:427-34.
763   45.   Cristea IM, Chait BT. 2011. Conjugation of magnetic beads for
764   immunopurification of protein complexes. Cold Spring Harb Protoc 2011:pdb
765   prot5610.

766   46.   Lambeth CR, White LJ, Johnston RE, de Silva AM. 2005. Flow cytometry-based
767   assay for titrating dengue virus. J Clin Microbiol 43:3267-72.

A

| Negative | GFP | NS5-GFP |

B

IP: GFP   1% input

GFP   NS5-GFP   GFP   NS5-GFP

Immunoblot

mTOR

Raptor

Rictor

NS5