1 **Title:** Host phylogeny and host ecology structure the mammalian gut microbiota at different
2 taxonomic scales
3
4 **Running title:** Gut microbiota of African herbivores
5

6 **Author names:** Connie A. Rojas[1,2,3], Santiago A. Ramírez-Barahona[4], Kay E.
7 Holekamp[1,2,3], Kevin. R. Theis[3,5]
8
[1]Department of Integrative Biology, Michigan State University, Lansing, MI, USA
[1]Ecology, Evolution and Behavior Program, Michigan State University, Lansing, MI, USA
[3]BEACON Center for the Study of Evolution in Action, Michigan State University, Lansing,
MI, USA
[4]Departament of Botany, Institute of Biology, Universidad Nacional Autónoma de México,
Mexico City, MX
[5]Department of Biochemistry, Microbiology and Immunology, Wayne State University
School of Medicine, Detroit, MI, USA
17
18 **Name and email address of all authors:**
19 Connie Rojas, rojascon@msu.edu (Corresponding author)
20 Santiago A. Ramírez-Barahona, santiago.ramirez@ib.unam.mx
21 Kay E. Holekamp, holekamp@msu.edu
22 Kevin R. Theis, ktheis@med.wayne.edu
23
24 **Full postal address of corresponding author:**
25 Natural Science Building
26 288 Farm Lane, Rm. 321
27 East Lansing, MI 48824
28

29 **Keywords (3-10):** herbivore gut microbiome, African mammals, phylosymbiosis, diet, 16S
30 rRNA gene sequencing, host ecology, host geography
31
32 **word count for the abstract (max 350 words):** 228
33 **combined number of main tables and figures:** 7
34 **References:** 109

Abstract
The gut microbiota is critical for host function. Among mammals, host phylogenetic relatedness and diet are strong drivers of gut microbiota structure, but one factor may be more influential than the other. Here, we used 16S rRNA gene sequencing to determine the relative contributions of host phylogeny and host dietary guild in structuring the gut microbiotas of 11 herbivore species from 5 families living sympatrically in southwest Kenya. Herbivore species were classified as grazers, browsers, or mixed-feeders. We found that gut microbiotas were highly species-specific, and that host taxonomy accounted for more variation in the gut microbiota (30%) than did host dietary guild (10%) or sample month (8%). Overall, similarity among gut microbiotas increased with host phylogenetic relatedness (r=0.73), yet this relationship was not consistent among seven closely related Bovid host species (r=0.21 NS). Within bovids, host dietary guild explained more of the variation in the gut microbiota than did host species. Lastly, while we found that the gut microbiotas of herbivores residing in southwest Kenya converge with those of distinct populations of conspecifics from central Kenya, fine-scale differences in the abundances of bacterial amplicon sequence variants (ASVs) between individuals from the two regions were also observed. Overall, our findings suggest that host phylogeny and taxonomy strongly structure the gut microbiota, but these gut microbial communities could be furthered modified by host ecology, especially among closely related host species.

Background
The gut microbiota performs critical roles for its host and is essential to host functioning. In mammals, resident gut microbes promote the digestive efficiency of their hosts by synthesizing vitamins, breaking down fiber, and supplementing the host with energy released from fermentation [1–4]. The gut microbiota also interacts with the host immune system, and may also modulate behavior [5–7]. Due to the critical importance of the gut microbiota for host performance, research has focused on determining the forces that shape its assembly and composition. Decades of research show that across vertebrate hosts, the gut microbiota is predominantly shaped by host phylogeny and ecology. Closely related host species tend to have more similar gut microbiotas than more distantly related host species [8–12] and this congruence between host phylogenetic relatedness and gut microbiota similarity is termed "phylosymbiosis" [13–15]. However, gut microbiotas can also be further shaped by their host's ecology, including their host's diet, habitat, and geographic location [16–19]. Thus, although both of these host factors may shape the gut microbiota, their relative contributions might be influenced by a variety of variables including the taxonomic breadth of the host species surveyed, and the diversity of host habitats, diets, and physiologies represented.

Similarity in the gut microbiota among individuals could be due to hosts providing potential colonizing microbes with similar ecological niches [20, 21]. This could arise from overlap in diet, habitat, physiology, and behavior among hosts, as well as from shared evolutionary history. Thus, if phylosymbiosis is observed, both host phylogenetic relatedness and ecology could be contributing to the pattern. A diversity of studies have disentangled the effects of these two factors and have shown that phylosymbiosis can be observed among hosts that share habitats or diets, and among hosts that reside in different habitats and consume different diets. For example, in mice, voles, and shrews, gut microbiotas tend to be more similar among closely related host species, despite these animals occupying different habitats [22]. In populations of American pikas (*Ochotona princeps*) from different mountain ranges, a cladogram of gut microbiota similarity was congruent with a phylogeny

of host genetic similarity [23]. Among hosts with overlapping diets, gut communities still
exhibit patterns consistent with phylosymbiosis, as has been documented for folivorous
primates [9]. Furthermore, in 33 species of sympatric herbivores from the Laikipia region in
central Kenya, host phylogenetic relatedness strongly predicted gut microbiota composition
(r = 0.91) and was weakly correlated with host diet (r = 0.28) [24], suggesting that
convergence of gut microbiotas among closely related hosts was not due to similarities in
their diet.
106
Here, we build upon this work and use 16S rRNA gene sequencing to determine the relative
influences of host phylogenetic relatedness and host ecology in structuring the gut
microbiota of 11 species of herbivores living sympatrically in the Masai Mara National
Reserve (henceforth the Masai Mara) in southwestern Kenya. We survey the gut
microbiotas of African buffalo, domestic cattle, common eland, impala, Kirk's dik-dik,
Thompson's gazelle, topi, Masai giraffe, common warthog, plains zebra, and African
elephant (Table 1). These species represent 5 mammalian families (Bovidae, Elephantidae,
Equidae, Giraffidae, and Suidae) and three dietary guilds: grazers, browsers, and mixed-
feeders. Furthermore, we compare the gut microbiotas of conspecific herbivores from the
Masai Mara and Laikipia to determine the extent to which host geography and/or local
habitat influence gut microbiota composition and patterns of phylosymbiosis. The two
regions differ in their altitude, vegetation, mammal densities, and degree of human
disturbance [25–29], any of which could potentially affect the gut microbiota compositions of
their resident herbivores.
121
Specifically, we surveyed the gut microbiotas of 11 species of herbivores to 1)
determine the relative contributions and amount of variance in the gut microbiota that is
explained by host taxonomy (family and species) and host diet, 2) ascertain whether
phylosymbiosis is observed across broad host taxonomic scales (i.e. among all study
species) and lesser host taxonomic scales (i.e. among 7 closely related Bovid species), and
3) examine the influences of host phylogenetic relatedness and host ecology (diet and
geography) on the gut microbiota of conspecific Masai Mara (this study) and Laikipia
herbivore hosts [24]. Collectively, our findings elucidate the factors shaping the gut
microbiota of hosts at greater and lesser taxonomic scales.
131
Results
Aim 1: Determine the amount of variance in the gut microbiota explained by host
taxonomy and host diet
Microbiota composition. Our analyses showed that some bacterial taxa were
widely shared among host species and dietary guilds, whereas others were abundant only
in particular host species. All herbivore gut microbiotas were dominated by two bacterial
phyla, *Firmicutes* (51% average relative abundance across samples), and *Bacteroidetes*
(32%) (Figure S1). The most abundant bacterial families were *Ruminococcaceae* (30.8%),
Rikenellaceae (11.4%), *Lachnospiraceae* (10.9%), and *Prevotellaceae* (8%) (Figure 1A).
Prevalent bacterial genera included *Alistipes*, *Bacteroides*, *Ruminococcus*, and *Treponema*
(Figure S2). A total of 10 out of 11, 930 (0.08%) Amplicon Sequence Variants (ASVs) were
present in 90% of samples pooled across all host species; 7 were assigned to the family
Ruminococcaceae, 1 to *Peptococcaceae*, and 2 to *Lachnospiraceae* (*Agathobacter*).
According to a blast search against the NCBI nucleotide database, sequences from the 7
Ruminococcaceae ASVs were highly similar to sequences from uncultured
Ruminococcaceae strains, uncultured rumen bacteria, and uncultured anaerobic bacteria.

Of these 10 ASVs, only two were abundant across samples (e.g. ASV10543
Ruminococcaceae), five were modestly abundant in specific host species (e.g. ASV71
Agathobacter in elephants), and 3 were present at very low abundances in all samples (e.g.
ASV7824 *Peptococcaceae*) (Figure S3). The latter 3 ASVs do not appear to be
contamination introduced during DNA extraction and sequencing, as these sequences are
highly similar to those found in rumen and fecal samples.
154
Nonetheless, variation in gut microbiota compositions among host species and
dietary guilds was clearly evident. Indicator species analysis showed that the gut
microbiotas of elephants were significantly associated with *Endomicrobiaceae* and
Desulfobulbaceae, those of Zebras with *Helicobacteraceae* and *Deltaproteobacteria*, and
those of warthogs with *Myxococcales* and *Coxiellaceae* (Figure 1B). Giraffe gut microbiotas
were highly associated with *Enterobacteriaceae*, *Bifidobacteriaceae*, and *Bacillaceae*.
Similarly, there were bacterial types that were indicative of specific dietary guilds. Grazer
gut microbiotas were characterized by *Sphingobacteriaceae*, *Flavobacteriaceae*,
Neisseriaceae, and *Lentisphaeria* (Figure 1C). Browser gut microbiotas had 11 indicator
bacterial taxa, including *Bacillaceae*, *Coriobacteriales, Methanomicrobia* and
Rubrobacteriaceae. Lastly, the gut microbiotas of mixed feeders were highly associated
with *Synergistaceae, Succinivibrionaceae*, and *Bacteroidales*, among other bacteria.
167
More fine-scale analysis of the presence and absence of bacterial ASVs also
revealed that the gut microbiotas of our studied herbivores contained microbes that were
biased towards particular host species. These were bacterial ASVs that were present in
75% samples for that host species, and absent in 97% of other samples. Buffalo, cattle,
topi, and impala gut microbiotas mostly contained ASVs that were present in other
herbivores, as <3% of their ASVs were biased to any of these host species. Between 4%
and 8% of ASVs comprising the gut microbiota of dik-diks, eland, elephant, Thompson's
gazelle, and giraffe were biased towards these host species. Warthogs and zebras
however, harbored more unusual microbiotas, as 70-77% of their ASVs were rarely
detected in the guts of the other African mammals.
178
Microbiota alpha-diversity : Gut microbiota richness, evenness, and phylogenetic
diversity also varied with host taxonomy (family and species) and dietary guild (Table 2,
Figure 2). Post-hoc comparisons revealed that hosts from the Suidae and Elephantidae
families generally harbored less rich and less even gut communities than the other
surveyed host families (Figure 2A; Table S1). Moreover, equids harbored more
phylogenetically diverse gut communities than all other herbivores (Table S1). Across the
three alpha-diversity metrics, browsers had less diverse gut microbiotas than grazers or
mixed-feeders (Figure 2A, Table S2). Within a single host family (i.e. Bovidae), the gut
microbiotas of buffalo, dikdiks, and gazelles were less rich, even, and phylogenetically
diverse than those of all other sampled bovids (Figure 2B, Table S3). In bovid hosts,
browsers also had less diverse gut microbiotas than grazers or mixed-feeders (Figure 2B,
Table S3).
191
Additionally, we determined whether *similarity* in gut microbiota alpha-diversity was
correlated with a *quantitative measure* of host diet, while accounting for variation in the gut
microbiota associated with host phylogenetic relatedness. Specifically, we used previously
published dietary C4 (%) data [30–33] from these 11 herbivore species as our numerical
measure of diet (Table S4). C4 (%) values quantify the degree of grazing by the host and

reflect the proportion of an animal's diet composed of C4 monocotyledon grasses relative to
C3 trees, shrubs, and forbs. Similarity in gut microbiota *evenness* (Shannon diversity)
increased with host overlap in dietary C4, but this was not true of gut microbiota richness or
phylogenetic diversity (partial Mantel Chao1 r=0.03, p=0.34; Shannon r=0.35, p=0.028, PD
r=0.097, p=0.25).
202
Microbiota beta-diversity: Across the surveyed herbivores, host family explained
on average ~23% of the variation in gut microbiota structure, followed by host dietary guild
(10%), sample month (8%), and host species (7.5%) (PERMANOVA analyses, Table 3).
Regardless of whether distance matrices took into account the proportions of bacterial taxa,
their presence/absence, or their phylogenetic relatedness, the percent variation explained
by each host factor was consistent. Therefore, for brevity, we only present PCoA ordination
plots using the Bray-Curtis index. These plots show that gut microbiotas primarily partition
by host family, and also secondarily by host dietary guild (Figure 3A).
211
Within Bovidae, host dietary guild was a slightly stronger predictor of the gut
microbiota than host species or sample month. On average, host dietary guild accounted for
17.3% of the variation, whereas host species explained 12.2% of the variation, and sample
month contributed to 7.3% of the variation (PERMANOVA analyses, Table 3). PCoA
ordinations showed that the gut microbiotas of bovids clustered by host dietary guild and
host species (Figure 3B). When controlling for host dietary guild and sample month, the
microbiota is highly-species specific and host species accounts for 40% of the observed
variation across samples (PERMANOVA: Bray-Curtis R^2=0.42; Jaccard R^2=0.39; Weighted
Unifrac R^2=0.40, Unweighted Unifrac R^2=0.41; all p=0.001) (Figure S4).
221
Aim 2: Ascertain whether phylosymbiosis is observed among herbivore hosts at
broad and lesser taxonomic scales
Although a dendrogram of gut microbiota similarity did not closely reflect host phylogeny
(Figure 3C), gut microbiota similarity did increase overall with host phylogenetic relatedness
among the 11 species of herbivores (mantel test Bray-Curtis r=0.76, p=0.006; Jaccard
r=0.72, p=0.006; Weighted Unifrac r=0.75, p=0.007; Unweighted Unifrac r=0.72, p=0.007)
(Figure 3D). The gut microbiotas tended to be more similar among closely related host taxa
(e.g. buffalo and cattle) than among distantly related host taxa (e.g. impala and elephant).
Despite warthogs being more closely related to bovids than to elephants and zebras, they
had highly dissimilar communities from all of the herbivore species examined, and pairwise
comparisons that include this host species deviate from the trendline (~1.0 Bray-Curtis
dissimilarity, Figure 3D).
234
Importantly, at a lesser host taxonomic scale, among closely related host species in
the Bovid family, we did not find a consistent relationship between gut microbiota similarity
and host phylogenetic relatedness (mantel test Bray-Curtis r=0.27, p=0.09; Jaccard r=0.20,
p=0.08; Weighted Unifrac r=0.22, p=0.059; Unweighted Unifrac r=0.16, p=0.08) (Figure 3D).
However, when taking into account fine-scale variation associated with host diet (%C4)
(Table S4), gut microbiota similarity is modestly correlated with host phylogenetic
relatedness (partial Mantel Bray-Curtis r=0.43, p=0.013; Jaccard r=0.35, p=0.016; Unifrac
r=0.33, p=0.017; Unweighted Unifrac r=0.29, p=0.017). When repeating this analysis on the
dataset that includes all herbivore hosts, the original finding remained unchanged;
accounting for fine-scale variation in host %C4 does not affect the phylosymbiosis pattern
observed at a broad host taxonomic scale (partial Mantel Bray-Curtis r=0.77, p=0.002,

Jaccard r=0.72, p=0.007, Unifrac r=0.75, p=0.004, Unweighted Unifrac r=0.72, p=0.01). In
summary, we observed phylosymbiosis among hosts spanning a broad taxonomic scale
involving multiple families, but not consistently among hosts spanning a lesser taxonomic
scale involving only members of the Bovid family.
250
Aim 3: Examine influences of host phylogenetic relatedness and host ecology on the
gut microbiota of conspecific hosts
Lastly, we further analyzed the influences of host phylogenetic relatedness and host
ecology (diet and geography) on the gut microbiota of 8 herbivore species from two distinct
populations in the Masai Mara (this study) and the Laikipia region, respectively (Kartzinel *et*
al. [24]). The eight herbivore species overlapping both studies were African buffalo,
domestic cattle, common eland, impala, giraffe, plains zebra, common warthog, and African
elephant.
259
We found that gut microbiota structure varied little among hosts from the two
geographic regions, as this factor accounted for <3% of the variation in the gut microbiota
(PERMANOVA analyses, Table S5). The gut microbiotas were primarily structured by host
species and host dietary guild, which explained on average, 38% and 11% of the variation,
respectively (PERMANOVA analyses, Table S5); sample month explained an additional 7%
of the variation. Ordination plots confirm these findings and demonstrate that samples
primarily cluster by host species (Figure 4A), although some separation of samples based
on host geographic region is also observed, particularly among cattle, impala, and giraffe.
268
For this dataset, a total of 3 ASVs out of 18, 039 (0.01%) were present across 90%
of samples; two were classified as *Ruminococcaceae*, and 1 as *Lachnospiraceae*. All 3
ASVs were among the 10 ASVs present across 90% of Masai Mara samples. To further
compare the gut microbiotas of conspecific hosts, we analyzed the relative abundances of
the 32 most abundant ASVs in the dataset. A heatmap of these 32 ASVs demonstrate that
there were ASVs that were found at comparable proportions in conspecific hosts, as well as
ASVs that were differentially abundant among conspecific hosts (Figure 4B, Table S6). For
example, ASV4033 *Prevotellaceae* is similarly abundant between Masai Mara and Laikipia
hosts, but ASV15828 *Kiritimatiellae* appears to be enriched in cattle from Laikipia compared
to cattle from the Masai Mara. Linear mixed models that statistically compared the relative
abundances of these ASVs between the two groups showed that 13 of the 32 ASVs were
actually differentially abundant between conspecifics (Table S6).
281
To extend these analyses and identify additional ASVs that may be differentially
abundant among conspecifics, we also conducted Linear discriminant analysis Effect Size
(LEfSe) for each host species. Roughly 30% of the ASVs in the gut microbiotas of each host
species were enriched in hosts from one population relative to the other (Table S7). ASVs
that were typically enriched were classified as *Ruminococcaceae*,
Lachnospiraceae, *Rikenellaceae*, *Clostridiales*, *Bacteroidales*, and *Kiritimatiellae* (Figure
4C). Herbivores from Laikipia tended to be enriched in *Methanocorpusculum, Clostridiales,*
and *Kiritimatiellae* ASVs relative to Masai Mara herbivores (Figure 4C). Masai Mara gut
microbiotas were overrepresented by *Lachnospiraceae* and *Treponema* ASVs. Interestingly,
hosts from both geographic regions could be enriched in taxonomically similar ASVs. For
example, eland in Laikipia were enriched in 3 ASVs classified as *Prevotellaceae*,
*Rikenellacea*e, and *Ruminococcaceae,* respectively, and eland from the Masai Mara were
enriched in 3 different ASVs that were also classified as *Prevotellaceae*, *Rikenellacea*e, and

Ruminococcaceae (Figure 4C). These findings suggest that variation in the gut microbiotas
of these herbivore conspecifics is observable at the level of specificity of bacterial ASVs.
297
Patterns consistent with phylosymbiosis were also observed in this combined
dataset, despite herbivore hosts occupying different geographic regions in Kenya, and
representing distinct populations. Gut microbiota similarity increased with host phylogenetic
relatedness (mantel test Bray-Curtis r=0.60, p=0.03; Jaccard r=0.64, p=0.04; Weighted
Unifrac r=0.54, p=0.003; Unweighted Unifrac r=0.66, p=0.04), although the strength of the
correlation was less than that of Masai Mara herbivores only (r=0.76), or what was
previously reported for Laikipia herbivores (r=0.91) [24]. Consistent with our earlier findings,
phylosymbiosis was not evident among the four species of bovids that overlapped between
the two studies (mantel test Bray-Curtis r=0.30, p=0.25; Jaccard r=0, p=0.5; Weighted
Unifrac r=-0.52, p=0.91; Unweighted Unifrac r=-0.49, p=0.83). When taking into account gut
microbiota variation attributable to host diet (% dietary C4), both findings remained
unchanged (<u>partial Mantel test all herbivores</u>: Bray-Curtis r=0.60, p=0.03, Jaccard r=0.64,
p=0.04, Weighted Unifrac r=0.53, p=0.03, Unweighted Unifrac r=0.65, p=0.05 0.05; <u>bovids</u>
<u>only</u>: Bray-Curtis r=0.36, p=0.16, Jaccard r=0, p=0.5, Weighted Unifrac r=-0.52, p=0.87,
Unweighted Unifrac r=-0.61, p=0.83).
313
Discussion
Principal findings of study. The primary purpose of this study was to determine the
relative contributions of host phylogenetic relatedness and dietary guild in structuring the
gut microbiotas of 11 species of sympatric African herbivores. We also compared the gut
microbiotas of herbivores from the Masai Mara to herbivores from Laikipia, Kenya to
determine the extent to which two distinct populations of identical herbivore species varied
in their gut microbiotas. We found that gut microbiotas were highly species-specific, but also
varied with host ecology, including host dietary guild and sample month, particularly in
closely related Bovid species. Furthermore, gut microbiota similarity increased with host
phylogenetic relatedness at broad host taxonomic scales, but at lesser taxonomic scales,
the phylosymbiosis signal was only apparent after accounting for fine-scale variation
attributable to host diet (%C4). Lastly, although the gut microbiotas of conspecific herbivore
hosts converged and primarily clustered by host species, differences between conspecifics
in the relative abundances of their bacterial ASVs were also observed. Collectively, our
findings suggest that mammalian gut microbiotas are strongly shaped by host phylogenetic
relatedness and taxonomy, but they can be further modified by host ecology, including host
diet and geography.
331
Aim 1: Determine the amount of variance in the gut microbiota explained by host
taxonomy and host diet
Across the surveyed herbivores, gut microbiota composition, diversity, and
structured varied with host taxonomy. Species-specificity of the gut microbiota is
widespread, and is commonly reported in the majority of comparative gut microbiome
studies [11, 34–36]. Host species may vary in their body size, behavior, neuroendocrine
system, immune function, and metabolism, any of which could potentially shape the
structure of their gut microbiotas [37–40]. When comparing gut microbiota alpha-diversity,
results showed that warthogs and elephants harbored less diverse gut communities than
the other sampled herbivores. Due to their omnivory, warthogs have a greater dietary
breadth than the other studied herbivores, yet they harbored less diverse microbiotas. This
is in accordance with prior findings, which report that the most diverse diets do not always

correlate with the most diverse gut microbiotas [24, 41, 42]. Furthermore, analyses showed
that browsers had the least diverse gut microbiotas, potentially because they consume
vegetation that has a higher lignin content and a lower fiber digestibility than grass [43].
Specialized bacterial metabolisms may be required to digest this tougher plant material.
Additionally, group size has been shown to correlate with gut microbiota diversity [44, 45],
and the browsers in our study (giraffes, dikdiks) forage in smaller groups than grazers
(buffalo, zebras) and mixed-feeders (gazelles, impala), which forage in herds. Frequent
social interactions and interactions with a greater number of individuals is known to promote
species richness in individual gut microbiotas [45, 46].
353
Similar to findings from a plethora of microbiome studies, the gut microbiota
structure of the studied herbivores also varied with host diet. Hosts from each dietary guild
consume food sources that vary in their structure, chemistry, and nutrition quality; which
require morphological, physiological and behavioral adaptations [47, 48]. For example,
grazers mostly feed on grasses, which have thicker cell walls, a lower protein content, and
use the C_4 photosynthetic pathway compared to browse (e.g. leaves, shrubs, and woody
vegetation), which have a higher protein content, and use C_3 photosynthesis [47, 48]. To
efficiently extract energy from these different food sources, browsers and grazers evolved
adaptations in their salivary chemistry, tooth morphology, gut structure, and speed of
digestion [49, 50]. These adaptations, along with the actual nutrients hosts are providing to
their microbes, potentially contribute to gut microbiota divergence among hosts from
different dietary guilds.
366
Despite differences in the gut microbiota among host species and dietary guilds,
there were some features of the gut microbiota that were shared across individuals from
multiple species. Across our surveyed herbivores, the most abundant bacterial taxa in the
gut microbiota were *Ruminococcaceae*, *Rikenellaceae, Lachnospiraceae,* and
Prevotellaceae which represent core taxa previously found in the gut microbiotas of many
ruminants and herbivores in general, including cervids and bovids [8, 51], equids [52],
elephants [53], and giraffes [54]. *Ruminococcaceae* and *Lachnospiraceae* have also been
found in the guts of folivorous primates [3] and in domestic pigs [55, 56]. Members of these
bacterial families are responsible for digesting the cellulose, hemicellulose, lignin, and
protein found in vegetation, and fermenting these into short-chain fatty acids (SCFAs) [57].
SCFAs represent usable forms of energy for the hosts [58] and contribute to host
colonocyte growth, immune defense, and anti-inflammatory responses [1]. These bacterial
taxa also possess fiber-degrading capabilities and can provide their hosts with protection
against ingested toxic plant secondary metabolites [59].
381
382   Interestingly, 7 of the 10 ASVs that were present in 90% of Masai Mara herbivores
383 were classified as *Ruminococcaceae* and were sequences highly similar to uncultured
384 *Ruminococcaceae* strains extracted from bovine, goat, and sheep rumens [60], suggesting
385 that these "core" microbes could be functionally important for the host.
386
Aim 2: Ascertain whether phylosymbiosis is observed among herbivore hosts at
broad and lesser taxonomic scales
Our results showed that phylosymbiosis was observed across a broad host taxonomic
scale, among 11 species of herbivores living sympatrically in the Masai Mara. Patterns of
phylosymbiosis have been documented extensively in many vertebrate groups, including
primates, rodents, ruminants, carnivores, reptiles, and insects [9–12, 24, 61, 62]. Evidence

of phylosymbiosis among host species living in *sympatry* specifically, has been previously
documented in seven species of deer mice [63], six species of Malagasy mammals [64],
twelve species of lemurs [65], and nine species of diurnal, non-human primates [66].
396
The mechanisms and processes that yield patterns of phylosymbiosis have not yet
been elucidated, but host ecological and phenotypic traits are likely acting as filters and thus
shaping microbial community assembly. Closely related hosts are potentially colonized by
taxonomically similar microbes due to similarities in their morphology, anatomy, digestive
physiologies, and immune system components [37, 38, 67]. Specifically, related hosts may
possess similar antimicrobial peptides and toll-like receptors that serve to filter the same
bacterial clades from the environment [68, 69]. Closely related hosts may further develop
immune tolerance via adaptive immunity to the same symbiotic, commensal, and transient
microbes [68, 69]. Lastly, some phylogenetically related hosts may also possess similar
social group structures and pathways for transmitting microbes among group-mates,
thereby contributing to patterns of phylosymbiosis. Overall, accumulation of differences in
traits as hosts diverged from one another could potentially provide enough niche
differentiation in the gut to promote the divergence of their symbiotic bacterial communities.
410
At a lesser host taxonomic scale, within our sampled group of closely related Bovid
species in the Masai Mara, variation in the gut microbiota was more strongly associated
with host diet than host species, and a pattern consistent with phylosymbiosis was only
apparent if fine-scale variation in diet (%C4) was accounted for. This phylosymbiosis signal
was weaker (r=0.35) than that observed for all herbivores (r=0.73). Similarly, other studies
report that among closely related hosts, host ecology more strongly predicts the structure of
the gut microbiota than host relatedness. For example, in lemurs (*Eulemur* spp.,
Propithecus spp.), phylosymbiosis was observed across but not within two host lineages,
and within host lineages, host habitat (dry forest vs. rainforest) was significantly correlated
with gut microbiota diversity [65]. In populations of yellow (*Papio cynecephalus*) and anubis
baboons (*Papio anubis*), gut microbiota dissimilarity did not increase with host genetic
distance, but did vary with their habitat's soil chemistry [70]. Because the bovids surveyed
here are closely related, their gut microbiotas are already very similar, and variation can
result from fine-scale differences in diet (proportions of grass vs. shrubs vs. trees
consumed) [43, 71–73]. Nonetheless, some of the variation in the gut microbiota of bovids
is not attributable to host diet and can be explained by host phylogenetic relatedness. Thus,
even among closely related host species, both ecological and evolutionary forces shape gut
microbial communities.
429
Aim 3: Examine influences of host phylogenetic relatedness and host ecology on the
gut microbiota of conspecific hosts
While gut microbiota structure was primarily associated with host species in the combined
Masai Mara and Laikipia dataset, differences in gut microbiota composition between
conspecific hosts from the two populations were also evident. Herbivores of the same
species may possess similar evolutionary trajectories, physiologies, and behaviors, and
thus may be providing microbes with similar niches for colonization, which is why at a broad
level their gut microbiotas converge. However, the two geographic regions do vary in their
climate, soil geochemistry, plant communities, and resident herbivore species [25–29, 74],
and potentially in their bacterial species pools, which could lead to the fine-scale microbiota
differences among conspecifics. This finding was supported by our data; over 30% of ASVs
were differentially enriched between Masai Mara and Laikipia hosts. Laikipia herbivores for

example, tended to be enriched in ASVs classified as *Methanocorpusculum, Clostridiales,*
and *Kiritimatiellae*, while Masai Mara herbivores had an overrepresentation of ASVs
belonging to *Lachnospiraceae* and *Treponema*. Abundances of the methanogenic
Methanocorpusculum are related to forage type and geographic location in cattle [75], and
in the mammalian gut, *Lachnospiraceae* are associated with a high-fat diet [76].
Furthermore, in the bovine rumen, *Treponema* degrade hemicellulose and their growth
increases in the presence of pectin [77, 78], a carbohydrate abundant in non-woody plants.
This suggests that differences in the ASV abundances between conspecific hosts likely
reflect fine-scale differences in their diets and habitats. Interestingly, both *Clostridiales* and
Lachnospiraceae are major microbial taxa of the mammalian gut and comprise fermentative
bacteria that synthesize SCFAs from the hydrolysis of starches and sugars [57]; thus,
conspecific hosts can be enriched in taxonomically distinct microbes that perform similar
functions.
455
456 Notably, 3 "core" bacterial ASVs were present in > 90% of samples from the 8 species of
457 herbivores sampled from two allopatric populations. The 3 ASVs were highly similar to
458 sequences from uncultured *Ruminococcaceae* strains and other anaerobic, rumen bacteria,
459 illustrating that these ASVs a) may be easily acquired from the environment, and b) could
460 be essential for host digestion.
461
Lastly, we found that phylosymbiosis was also evident among conspecific African
herbivores living in allopatry, although the strength of the phylosymbiotic signal was slightly
reduced compared to that observed for either sympatric population considered in isolation.
Overlap in gut microbiota structure is thought to be lower in allopatric animal populations
than in sympatric animals due to variation introduced by habitat, dietary differences, and the
spatial limits of bacterial dispersal [12]. It is important to note that differences between the
gut microbiotas of Laikipia and Masai Mara conspecifics could also be potentially
attributable to differences in sampling, DNA extraction, and sequencing protocols between
the two studies [79–81]. Collectively, our findings show that mammalian gut microbiotas
converge among closely related host species and among conspecifics, but can be
differentiated with variation introduced by the host's ecology.
473
Conclusions
Our study showed that among 11 species of African herbivores living in sympatry, gut
microbiotas are highly species-specific and exhibit patterns congruent with phylosymbiosis.
However, these gut microbiotas are also shaped by their host's ecology, and we observed
differences between hosts from different dietary guilds . Furthermore, among eight species
of herbivores residing in two geographic regions in Kenya, gut microbiotas were similar
among hosts of the same species, but also exhibited fine scale differences in the
abundances of their bacterial ASVs. Overall, these findings suggest that related hosts are
providing microbes with similar niches for colonization, but these microbial niches are
further shaped by host diet, geography, and local environmental conditions. Future studies
should examine whether phylosymbiosis is evident at the functional level in the gut
metagenomes of African herbivores and in metazoan taxa in general. The gut microbiotas
between distantly related host species could be taxonomically dissimilar, but functionally
similar due to the widespread functional redundancy observed in microbial metabolisms.
These studies will be necessary to further our understanding of the processes and
mechanisms potentially underlying patterns of phylosymbiosis.
490

Methods
Study location and sampling.
Fecal samples (N=181) were collected opportunistically from 11 species of herbivores
permanently residing in the Talek and Mara Triangle regions of the Masai Mara (1°22′19″S,
34° 56′17″E) from March-June 2018 (Table 1). This Reserve is covered by open rolling
grassland interspersed with seasonal watercourses and riparian vegetation. It has two rainy
seasons (March-May and November-December, with annual rainfall >1000mm) [82], and
most of our sampling took place during the rainy months, particularly during the month of
March (Figure S5). Although the Masai Mara is home to small resident populations of zebra
and wildebeest, millions of these individuals migrate into the Reserve from July-October
every year. Because our sampling occurred before July, samples from wildebeest and
zebras were limited.

For fecal sample collection, we either observed animals defecating or identified
species-of-origin based on the size, shape, and consistency of fresh dung, following
Kartzinel *et al.* [24]. Samples were then placed in sterile cryogenic vials and stored in liquid
nitrogen until they were transported to Michigan State University, where they remained
frozen at -80°C until nucleic acid extraction. For a list of samples and their associated
metadata, see the Github repository for this project
(https://github.com/rojascon/Rojas_et_al_2020_African_herbivores_gut_microbiome).

While we did not directly collect diet data from the surveyed herbivores, we used
Kingdon's *East African Mammals* [83–86] to classify our study species into grazers,
browsers, and mixed-feeders. To obtain more fine-scale data on host diet, we also compiled
dietary C4 (%) data for these herbivores from previously published studies (Table S4) [30–
33]. Percent C4 values reflect the proportion of monocotyledon grasses consumed relative
to trees, shrubs, and forbs.

DNA extraction and 16S rRNA gene sequencing.
Fecal samples were sent to the University of Chicago at Illinois (UIC) Sequencing Core for
automated DNA extractions using QIAGEN DNeasy PowerSoil kits (Valencia, CA, USA).
DNA concentrations of the fecal sample extracts were quantified using Qubit. The V4 region
of the 16S rRNA gene was targeted for sequencing on the Illumina MiSeq platform at the
Michigan State University Genomics Core, using published protocols by Caporaso *et al.*
2012 [87] and Kozich *et al.* 2013 [88].

Sequence processing and bioinformatics.
Sequences from Masai Mara herbivore gut microbiotas were processed in R (v.3.6.2) [89]
using the Divisive Amplicon Denoising Algorithm (DADA2) pipeline (v.1.14.1) [90] to infer
amplicon sequence variants (ASVs). Briefly, reads were filtered for quality, allowing for 2
and 3 errors per forward and reverse read, respectively (trimLeft = c(10, 10), maxN = 0,
maxEE = 2, truncQ = 2). Forward reads were trimmed to 240bp and reverse reads to
200bp; these paired-end reads were merged. Sequences were then dereplicated to remove
redundancy and ASVs were inferred by pooling reads from all samples. Prior to creating the
ASV abundance table, chimeras were removed and ASVs were taxonomically classified
using the SILVA rRNA gene reference database (v.132) [91] with an 80% confidence
threshold. ASVs taxonomically assigned as Eukarya, Chloroplasts, or Mitochondria were
removed from the dataset, as were those of unknown Kingdom origin; 12,938 total ASVs
remained. The resulting ASV table and the taxonomic designations of the ASVs are

available on GitHub. On average, samples retained over 70% (± 11%) of their total
sequences after processing in DADA2. Nineteen samples did not amplify well (<400
sequences after processing) and were removed from the dataset. Most of these samples
belonged to browser species (giraffes and dik-diks) suggesting that there may have been
PCR inhibitors in their fecal samples (e.g. humic acid, tannins) that prevented successful
extraction of DNA or library preparation. Table 1 has the sample sizes (N) for each study
species before and after this filtering.
547
Microbiota composition analyses
Statistical analyses and data visualization were completed in R unless otherwise stated. To
visualize microbiota composition, stacked barplots were constructed in ggplot2 (v.3.3.2)
[92]. These plots showed the bacterial phyla, families, and genera with average relative
abundances greater than 1% across samples. We also identified the ASVs that were
present in >90% of samples across all host species, and the relative abundances of these
ASVs were displayed as heatmaps using the R pheatmap package (v.1.0.12) [93].
Sequences from the 10 ASVs were blasted against the National Center for Biotechnology
Information (NCBI) Nucleotide database [60] to find similar biological sequences from
known bacterial taxa. Furthermore, we also identified ASVs that were biased towards
particular host species; these were ASVs that were present in >75% of the samples for a
particular host species (e.g. giraffes) and absent in 97% of samples from the other host
species.
561
To detect the bacterial taxa strongly associated with particular host families or
dietary guilds, we used the R indicspecies package (v.1.7.9) [94], which calculates an
indicator value for each bacterial taxon based on its prevalence in a given group and
absence in others. A table of bacterial family relative abundances was used as input, and
significance was assessed via permutation tests using 999 permutations (α=0.05). Bacterial
families with indicator values >0.4 were plotted in ggplot2.
568
Microbiota α-diversity statistical analyses
Prior to alpha-diversity analyses, we controlled for the potential influences of sequencing
depth by subsampling all samples to 17,000 sequences using the mothur (v.1.42.3) [95]
sub.sample command. Four fecal samples did not meet this sequence cutoff criterion, and
were excluded from all alpha-diversity analyses. Mothur was used to construct rarefaction
curves of ASV richness vs. sequencing depth (Figure S6) and Good's coverage values
averaged 97.78 ± 0.91 across all samples, indicating that sample coverage was high and
appropriate for characterizing fecal microbiota profiles. These values are comparable to
those typically reported in other mammalian gut microbiota studies [8, 96, 97].
578
Microbiota alpha-diversity was estimated using Chao1 Richness, Shannon diversity,
and Faith's phylogenetic diversity in R. Chao1 Richness and Shannon indices were
calculated using the phyloseq package (v.1.33.0) [98]. To obtain measures of Faith's
Phylogenetic Diversity (PD), we constructed a phylogenetic tree of ASV sequences using
phangorn (v. 2.5.5) [99] and calculated PD using the picante package (v.1.8.1) [100]. The
effects of predictor variables on each measure of alpha-diversity across all samples were
evaluated via linear mixed models (LMMs) using the lme4 package (v.1.1.23) [101],
specifying host dietary guild and host family as fixed variables and sample month as a
random effect. A similar model was also built for bovid samples, which included host
species as a predictor in lieu of host family. The significance of each predictor variable was

determined by calculating likelihood ratio Chi-2 test statistics (α=0.05) on the full models
using the car package (v.3.0.7) [102]. These tests were followed by TukeyHSD post-hoc
tests with Benjamini-Hochberg adjustments to control for multiple comparisons. Boxplots of
microbiota alpha-diversity were generated in ggplot2.
593
To further quantify the influence of host diet on the three metrics of gut microbiota
alpha-diversity, we conducted partial Mantel tests in the R vegan package (v.2.5.7) [103].
Specifically, we evaluated whether similarity in gut microbiota alpha-diversity was
associated with similarity in dietary C4(%) after taking into account variation due to host
phylogenetic relatedness. The 3 matrices used as input were i) a dissimilarity matrix of gut
microbiota alpha-diversity, ii) a dissimilarity matrix of host %C4, and iii) a matrix of host
divergence times.
601

Microbiota β-diversity analyses and testing for phylosymbiosis
In order to determine the relative contributions and amount of variance explained by host
predictor variables, permutational multivariate analyses of variance (PERMANOVA) tests
based on Bray-Curtis, Jaccard, and Unifrac distance matrices were run using vegan. Bray-
Curtis/Jaccard distances were estimated using vegan, whereas weighted and unweighted
Unifrac distances were estimated using phyloseq. Bray-Curtis and weighted Unifrac
distances take into account the abundances of bacterial taxa while Jaccard and unweighted
Unifrac metrics only consider their presence or absence. Both UniFrac metrics utilize
information on the phylogenetic diversity of bacterial members when calculating microbiota
similarity. PERMANOVA model #1 included sample month, host dietary guild, host family,
and host species as predictors (in this order). Model 2 was identical, except it was restricted
to the Bovid dataset and excluded the host family term. Model 3 quantified the effect of the
host species predictor while controlling for dietary guild (strata=dietary guild). Microbiota
similarity and groupings across samples were visualized via Principal Coordinates Analysis
(PCoA) plots.
617

To test for phylosymbiosis, i.e. the congruence between host phylogenetic
relatedness and gut microbiota similarity, mean divergence times (mya) were calculated
between every pair of host species in R. First, we retrieved 1000 phylogenetic trees that
included all species of Artiodactyla and African elephants (*Loxodonta Africana*) from
Upham's et al. (2019) Mammalian supertree [104]. The trees were randomly sampled from
the posterior distribution of Upham's supertrees (Mammals birth-death tip-dated DNA-only
trees) using the VertLife online resource (http://vertlife.org/). Each tree was pruned to
include only the species in this study, and branch lengths (i.e. divergence times between
each pair of host species) were extracted using the R ape package (v.5.4.1) [105]. All 1000
trees showed the same phylogenetic relationships among the study species and matrices of
mean divergence times were estimated from those trees. To determine the strength of the
phylosymbiosis signal, we conducted mantel tests on gut microbiota dissimilarity matrices
and matrices of host divergence times, setting 999 permutations and Spearman
correlations. We also ran partial Mantel tests to evaluate the strength of the phylosymbiosis
signal while taking into account gut microbiota variation attributable to diet. The 3 matrices
used as input were i) a gut microbiota dissimilarity matrix, ii) a matrix of host divergence
times, and iii) a dissimilarity matrix of host %C4 (Table S4).
635

636   We visualized the phylosymbiosis findings by plotting gut microbiota similarity (0-1)
637   against host phylogenetic divergence time (mya) in ggplot2. We also constructed a

638   consensus phylogeny of our host species and compared it against a dendrogram of gut
639   microbiota dissimilarity, which was calculated using hierarchical clustering with the R stats
640   package [89] and plotted using the ape package.
641
Comparisons of Masai Mara and Laikipia herbivores.
In order to compare the gut microbiotas of Masai Mara (1°22′19″S, 34° 56′17″E) herbivores
to the gut microbiotas of their conspecifics in Laikipia (0°17′33″N, 36° 53′55″E) (~ 290 km
from the Masai Mara), we downloaded all publicly available sequences from Kartzinel *et al.*
[24], and combined them with the raw 16S rRNA gene sequences from this study (Masai
Mara herbivores). These were then processed together in DADA2. A total of eight herbivore
species overlapped between the two studies: African buffalo, domestic cattle, common
eland, impala, giraffe, warthog, plains zebra, and African elephant. The majority of samples
from Kartizinel *et al.* [24] and from our study were collected during the wet seasons in their
respective regions (Table S8), although, in general, Laikipia is more arid than the Masai
Mara, with only 300-600mm precipitation annually [28, 106, 107]. For a list of all samples
(N=305), and their associated meta data, see the *Availability of data and materials* section.
654
The bioinformatics processing and statistical analyses were performed as described
above, with a few exceptions. In DADA2, forward and reverse reads were trimmed to 240bp
and 150bp, respectively, to better account for sequence quality. Up to 2 errors were allowed
per forward read and up to 4 errors per reverse read. To identify the strongest predictors of
gut microbiota structure, we constructed a PERMANOVA model that included sample
month, host geographic region, host dietary guild, and host species as variables (in this
order). PCoA ordinations and testing for phylosymbiosis were conducted as described
earlier. To visualize gut microbiota compositions between Masai Mara and Laikipia
herbivores, a heatmap of the 32 most abundant bacterial ASVs was constructed using R
pheatmap. To statistically evaluate whether conspecific hosts differed in their abundances
of these 32 ASVs, we ran 32 distinct LMMs in R, each specifying an ASV relative
abundance as the dependent variable, host region as a fixed effect and host species as a
random effect (to restrict analyses to within a host species). p-values were adjusted for
multiple comparisons using the Bonferroni-Hochberg method. To extend this analysis to
include more ASVs, we conducted Linear discriminant analysis Effect Size (LEfSe) [108] in
the Galaxy platform [109] using default parameters. Only ASVs >0.01% average relative
abundance across samples were included in the dataframe uploaded to Galaxy. ASVs that
were enriched in hosts from one geographic region relative to the other were visualized via
diverging dot plots in R with the ggplot2 package.
674
List of abbreviations used in this manuscript
Amplicon Sequence Variant (ASV)
Linear Mixed Model (LMM)
Permutational Multivariate Analysis of Variance (PERMANOVA)
Principal Coordinates Analysis (PCoA)
Linear discriminant analysis Effect Size (LEfSe)
National Center for Biotechnology Information (NCBI)
Short Chain Fatty Acids (SCFAs)
University of Chicago at Illinois (UIC)
Divisive Amplicon Denoising Algorithm (DADA2)
Faith's Phylogenetic Diversity (PD)
Millions of years (mya)

Institutional Animal Care and Use Committee (IACUC)

DECLARATIONS
Ethical Approval and Consent to participate
Our research and procedures were most recently approved on April 16, 2019 (IACUC
approval no. PROTO201900126) and comply with the ethical standards of Michigan State
University and Kenya.

Consent for publication
Not applicable

Availability of data and materials
The 16S rRNA gene sequence data from this study were deposited in NCBI's Sequence
Read Archive under BioProject PRJNA656793 and accession numbers SAMN15803511-
SAMN15803691. Sample metadata, data output by DADA2 (ASV table & ASV taxonomic
classifications), data obtained from LEfSe, and R scripts for all analyses and figures
included in this manuscript are available on Github
(https://github.com/rojascon/Rojas_et_al_2020_African_herbivores_gut_microbiome).

Supplementary Materials
Supplementary tables and figures can be found online at bioRxiv, alongside this manuscript
submission.

Competing Interests
The authors declare that there are no competing interests.

Funding
This work was supported by funds from NSF grants OISE1853934, IOS 1755089, and
OIA0939454 to K.E.H. and colleagues, the latter via the BEACON Center for the Study of
Evolution in Action. The corresponding author (C.A.R.) was also supported by a Graduate
Research Fellowship from NSF, a Predoctoral Fellowship from the Ford Foundation, and a
summer fellowship awarded by the MSU Ecology, Evolution, and Behavior program.

Author Contributions
K.R.T., K.E.H., and C.A.R. designed the study, C.A.R. and Mara Hyena Project field
assistants collected the samples. C.A.R., S.R.B., and K.R.T. analyzed the data. C.A.R.,
K.R.T, and K.E.H. wrote the manuscript and all authors approved the final version of the
manuscript.

Acknowledgements
We thank the field assistants and graduate students from the Mara Hyena Project for
assisting with fecal sample collection in the field. We also thank the Kenya Wildlife Service,
the Kenyan National Commission on Science, Technology and Innovation, the Kenyan
National Environmental Management Authority, the Narok County Government, the
Naboisho Conservancy, the Mara Conservancy, and Brian Heath for allowing us to conduct
this research.

Author's information
Not applicable

Tables
Table 1. List of host study species and their associated metadata.

Order	Family	Species	Dietary Guild	Total Samples (N)	Analyzed samples (N)
Cetartiodactyla	*Bovidae*	African buffalo	grazer	18	17
Cetartiodactyla	*Bovidae*	Domestic cattle	grazer	14	13
Cetartiodactyla	*Bovidae*	Common eland	mixed feeder	8	8
Cetartiodactyla	*Bovidae*	Impala	mixed feeder	20	20
Cetartiodactyla	*Bovidae*	Kirk's dik dik	browser	37	31
Cetartiodactyla	*Bovidae*	Thomson's gazelle	mixed feeder	14	14
Cetartiodactyla	*Bovidae*	Topi	grazer	19	19
Cetartiodactyla	*Giraffidae*	Masai giraffe	browser	25	18
Cetartiodactyla	*Suidae*	Warthog	mixed feeder	9	8
Perissodactyla	*Equidae*	Plains zebra	grazer	5	5
Proboscidea	*Elephantidae*	African elephant	mixed feeder	12	12

Table 2. Microbiota richness, evenness, and phylogenetic diversity vary with host taxonomy and dietary guild.

	Factor	Chao 1 Richness	Shannon diversity	Phylogenetic diversity
Across study species (N=161)	Host family	$\chi^2=53.58$, **p<0.001**	$\chi^2=33.45$ **p<0.001**	$\chi^2=18.31$ **p=0.001**
	Host dietary guild	$\chi^2=79.027$ **p<0.001**	$\chi^2=73.72$ **p<0.001**	$\chi^2=52.61$ **P<0.001**
Within bovids (N=118)	Host species	$\chi^2=57.03$ **p<0.001**	$\chi^2=10.81$ **p=0.02**	$\chi^2=50.53$ **p<0.001**
	Host dietary guild	$\chi^2=19.55$ **p<0.0001**	$\chi^2=33.91$ **p<0.0001**	$\chi^2=10.67$ **p<0.01**

Shown are the likelihood ratio χ^2 test statistics obtained for linear mixed effects models specifying host dietary guild and host family as predictor variables, sample date as a random effect, and an alpha-diversity metric as a dependent variable. A similar model restricted to bovids was also constructed; it specified host species instead of host family. Significant p-values (α=0.05) are bolded.

Table 3. Host taxonomy and dietary guild shape the gut microbiotas of African herbivores.

Analysis	Host factors	Bray-Curtis (% variance explained)	Jaccard (% variance explained)	Weighted Unifrac (% variance explained)	Unweighted Unifrac (% variance explained)
Across all host study species (N=165)	Host family	22.33, **p=0.001**	20.59, **p=0.001**	26.38, **p=0.001**	24.23, **p=0.001**
	Host dietary guild	11.25, **p=0.001**	10.21, **p=0.001**	9.71, **p=0.001**	10.30, **p=0.001**
	Host species	9.40, **p=0.001**	8.6, **p=0.001**	4.45, **p=0.001**	7.23, **p=0.001**
	sample month	7.42, **p=0.001**	6.80, **p=0.001**	10.21, **p=0.001**	7.66, **p=0.001**
Across bovids (N=122)	Host dietary guild	18.26, **p=0.001**	15.91, **p=0.001**	18.35, **p=0.001**	16.77, **p=0.001**
	Host species	15.16, **p=0.001**	13.38, **p=0.001**	8.23, **p=0.001**	12.16, **p=0.001**
	sample month	7.56, **p=0.001**	7.03, **p=0.001**	7.59, **p=0.001**	7.17, **p=0.001**

Shown are the R^2 values (% variance explained) and p-values for PERMANOVA tests (y ~ sample month + host dietary guild + host taxonomy) based on 4 types of distance matrices. Bray-Curtis and Weighted Unifrac distance matrices take into consideration the proportions of bacterial taxa, while Jaccard and unweighted Unifrac take into account only their presence or absence. Both Unifrac distances account for phylogenetic relatedness among bacterial types. Significant p-values (α=0.05) are bolded.

Figure legends

Figure 1. Gut microbiota composition of African herbivores. A) Stacked bar plots
showing the relative frequency of 16S rRNA gene sequences assigned to each bacterial
family (or order, if a family-level classification could not be assigned) across samples.
Samples are grouped by host species, and each color represents a bacterial family. **B)**
Bacterial families significantly associated with particular herbivore families as determined by
indicator species analysis. Differences in these taxa abundances can explain differences in
the microbiota among the different groups. **C)** Bacterial families significantly associated
with herbivores from different dietary guilds as determined by indicator species analysis.

Figure 2. Host taxonomy and dietary guild are associated with gut microbiota
diversity in African herbivores. A) Boxplots of microbiota evenness (Shannon diversity)
among host families and dietary guilds across all studied herbivores, and **B)** among host
species and dietary guilds within the family *Bovidae*. Boxes that do not share any letters
represent statistically significant comparisons; see Tables S1-S3 for all post-hoc
comparisons. Thicker dots represent outlier values.

811
Figure 3. African herbivore gut microbiotas exhibit patterns consistent with
phylosymbiosis. A) PCoA plots constructed from Bray-Curtis dissimilarity matrices. Each
point represents a sample and is color-coded by host family (left) or host dietary guild
(right). Closeness of points indicates high community similarity. The percentage of variance
accounted for by each principal-coordinate axis is shown in the axis labels. **B)** PCoA plots
constructed from Bray-Curtis dissimilarity matrices of bovid species only. Each point is
color-coded by host species (left) or host dietary guild (right). **C)** Phylogenetic tree of host
species (left) obtained from pruning Upham's *et al.* 2019 Mammalian supertree, compared
against a dendogram (right) of gut microbiota similarity using hierarchical clustering. **D)**
Scatterplot of pairwise host divergence times (in millions of years) vs. gut microbiota
similarity (Bray-Curtis distances) across all sampled herbivores (left) and within the single
host family *Bovidae* (right).

Figure 4. The gut microbiotas of conspecific African herbivores broadly converge, but also exhibit differences in their ASV abundances. We compared the gut microbiotas of eight species of herbivores residing in both the Masai Mara (this study) and Laikipia (Kartzinel *et al.* 2019) regions in Kenya. **A)** PCoA plot constructed from Bray-Curtis dissimilarity matrices. Each point represents a sample and is color-coded by host species; shape shading indicates geographic region. **B)** Heatmap of the 32 most abundant bacterial ASVs residing in the gut microbiotas of Masai Mara and Laikipia herbivores. Samples are grouped by host species, and are color-coded by host geographic region. **C)** ASVs enriched in Masai Mara vs. Laikipia herbivores as determined by LEfSe. Each dot represents a unique ASV and is color-coded by host geographic region. A total of 210 ASVs are displayed (those with LDA >3.2) and their family or genus level classification are on the x-axis.

A)

C)

B)

847
848
Additional files
Additional file 1. Tables S1-S8. Table S1-S3: Multiple-comparison testing of gut microbiota alpha-diversity among host families, species, and dietary guilds. **Table S4:** Previously published dietary %C4 data for the surveyed herbivores. **Table S5:** PERMANOVA tests that included sample month, geographic region, host dietary guild and host species for the combined Masai Mara and Laikipia dataset. **Table S6:** Linear models testing whether conspecific hosts differ in the relative abundances of 32 dominant ASVs. **Table S7:** Proportion of ASVs that were differentially enriched in herbivores from one geographic region compared to the other according to LEfSe. **Table S8:** Sample sizes for each month for the combined Masai Mara and Laikipia dataset (.xlxs 29 KB).

Additional file 2. Figures S1-S6. Figure S1-S2: Stacked bar plots showing relative abundances of top bacterial phyla and genera. **Figure S3:** Heatmap of the relative abundances of 10 widely-shared bacterial ASVs. **Figure S4:** PCoA ordinations showing sample clustering within a dietary guild (grazers, browsers, mixed-feeders). **Figure S5:** Stacked barplot of the proportion of samples from each host species that were collected each month (for the Masai Mara dataset). **Figure S6:** Rarefaction curves of ASV richness for the study samples (.pdf 1.9 MB).

References
1. Jovel J, Dieleman LA, Kao D, Mason AL, Wine E. The Human Gut Microbiome in Health and Disease. Metagenomics. 2018;:197–213. doi:10.1016/B978-0-08-102268-9.00010-0.
2. Ungerfeld E, Leigh M, Forster R, Barboza P. Influence of Season and Diet on Fiber Digestion and Bacterial Community Structure in the Rumen of Muskoxen (Ovibos moschatus). Microorganisms. 2018;6:89. doi:10.3390/microorganisms6030089.
3. Greene LK, McKenney EA, O'Connell TM, Drea CM. The critical role of dietary foliage in maintaining the gut microbiome and metabolome of folivorous sifakas. Sci Rep. 2018;8:1–13.
4. den Besten G, van Eunen K, Groen A, Venema K, Reijngoud D, Bakker B. The role of short-chain fatty acids in the between diet, gut microbiota, and host energy metabolism. J Lipid Res. 2013;54:2325–40. doi:10.1194/jlr.R036012.
5. Park W. Gut microbiomes and their metabolites shape human and animal health. J Microbiol. 2018;56:151–3. doi:10.1007/s12275-018-0577-8.
6. Münger E, Montiel-Castro AJ, Langhans W, Pacheco-López G. Reciprocal Interactions Between Gut Microbiota and Host Social Behavior. Front Integr Neurosci. 2018;12:1–14. doi:10.3389/fnint.2018.00021.
7. Pennisi E. Gut bacteria linked to mental well-being and depression. Science (80-). 2019;363:569–569. doi:10.1126/science.363.6427.569.
8. Li J, Zhan S, Liu X, Lin Q, Jiang J, Li X. Divergence of Fecal Microbiota and Their Associations With Host Phylogeny in Cervinae. Front Microbiol. 2018;9:1–11. doi:10.3389/fmicb.2018.01823.
9. Amato KR, Sanders J, Song SJ, Nute M, Metcalf JL, Thompson LR, et al. Evolutionary trends in host physiology outweigh dietary niche in structuring primate gut microbiomes. ISME J. 2019;13:576–587. doi:10.1038/s41396-018-0175-0.

10. Nishida AH, Ochman H. Rates of gut microbiome divergence in mammals. Mol Ecol.
2018;27:1884–97.

11. Youngblut ND, Reischer GH, Walters W, Schuster N, Walzer C, Stalder G, et al.
Host diet and evolutionary history explain different aspects of gut microbiome diversity
among vertebrate clades. Nat Commun. 2019;10:1–15. doi:10.1038/s41467-019-10191-
3.

12. Moeller AH, Suzuki TA, Lin D, Lacey EA, Wasser SK, Nachman MW. Dispersal
limitation promotes the diversification of the mammalian gut microbiota. PNAS.
2017;114:13768–73. doi:10.1073/pnas.1700122114.

13. Kohl KD. Ecological and evolutionary mechanisms underlying patterns of
phylosymbiosis in host-associated microbial communities. Philos Trans R Soc B Biol
Sci. 2020;375.

14. Lim SJ, Bordenstein SR. An introduction to phylosymbiosis. Proc R Soc B Biol Sci.
2020;287.

15. Brooks AW, Kohl KD, Brucker RM, Van Opstal EJ, Bordenstein SR. Phylosymbiosis:
Relationships and Functional Effects of Microbial Communities across Host
Evolutionary History. PLoS Biol. 2016;14:1–19. doi:10.1371/journal.pbio.2000225.

16. Watson SE, Hauffe HC, Bull M, McKinney MA, Atwood TA, Perkins SE. Global
change-driven use of onshore habitat impacts polar bear faecal microbiota. ISME J.
2019. doi:10.1038/s41396-019-0480-2.

17. Angelakis E, Yasir M, Bachar D, Azhar EI, Lagier JC, Bibi F, et al. Gut microbiome
and dietary patterns in different Saudi populations and monkeys. Sci Rep. 2016;6:1–9.
doi:10.1038/srep32191.

18. Goertz S, Menezes AB De, Birtles RJ, Id JF, Lowe E, Maccoll ADC, et al.
Geographical location influences the composition of the gut microbiota in wild house
mice (Mus musculus domesticus) at a fine spatial scale. PLoS One. 2019;14:1–16.

19. Orkin JD, Campos FA, Guadamuz A, Melin AD, Myers MS, Hernandez SEC.
Seasonality of the gut microbiota of free-ranging white-faced capuchins in a tropical dry
forest. ISME J. 2019;13:183–196. doi:10.1038/s41396-018-0256-0.

20. Hillerislambers J, Adler PB, Harpole WS, Levine JM, Mayfield MM. Rethinking
Community Assembly through the Lens of Coexistence Theory. Annu Rev Ecol Evol
Syst. 2012;43:227–48. doi:10.1146/annurev-ecolsys-110411-160411.

21. Smith B, Wilson JB. Community convergence: Ecological and evolutionary. Folia
Geobot. 2002;37:171–83.

22. Knowles SCL, Eccles RM, Baltrūnaitė L. Species identity dominates over
environment in shaping the microbiota of small mammals. Ecol Lett. 2019;22:826–37.

23. Kohl KD, Varner J, Wilkening JL, Dearing MD. Gut microbial communities of
American pikas (Ochotona princeps): Evidence for phylosymbiosis and adaptations to
novel diets. J Anim Ecol. 2018;87:323–30.

24. Kartzinel TR, Hsing JC, Musili PM, Brown BRP, Pringle RM. Covariation of diet and
gut microbiome in African megafauna. PNAS. 2019;116:1–6.

25. Ottichilo WK, De Leeuw J, Skidmore AK, Prins HHT, Said MY. Population trends of
large non-migratory wild herbivores and livestock in the Masai Mara ecosystem, Kenya,
between 1977 and 1997. Afr J Ecol. 2000;38:202–16.

26. Goheen JR, Augustine DJ, Veblen KE, Kimuyu DM, Palmer TM, Porensky LM, et al.
Conservation lessons from large-mammal manipulations in East African savannas: The

KLEE, UHURU, and GLADE experiments. Ann N Y Acad Sci. 2018;1429:31–49.

27. Veldhuis MP, Ritchie ME, Ogutu JO, Morrison TA, Beale CM, Estes AB, et al. Cross-boundary human impacts compromise the Serengeti-Mara ecosystem. Science (80-). 2019;363:1424–8. doi:10.1126/science.aav0564.

28. Augustine DJ. Response of native ungulates to drought in semi-arid Kenyan rangeland. Afr J Ecol. 2010;48:1009–20.

29. Broten MD, Said M. Population Trends of Ungulates in and around Kenya's Masai Mara Reserve. In: Serengeti II: Dynamics, Management, and Conservation of an Ecosystem. 1995. p. 169–93.

30. Cerling TE, Harris JM, Passey BH. Diets of East African Bovidae based on stable isotope analysis. J Mammal. 2003;84:456–70.

31. Kartzinel TR, Chen PA, Coverdale TC, Erickson DL, Kress WJ, Kuzmina ML, et al. DNA metabarcoding illuminates dietary niche partitioning by African large herbivores. PNAS. 2015;112:8019–24.

32. Codron D, Codron J, Lee-Thorp JA, Sponheimer M, De Ruiter D, Sealy J, et al. Diets of savanna ungulates from stable carbon isotope composition of faeces. J Zool. 2007;273:21–9.

33. Codron J, Lee-Thorp JA, Sponheimer M, Codron D, Grant R, De Ruiter D. Elephant (Loxodonta Africana) diets in Kruger National Park, South Africa: Spatial and landscape differences. J Mammal. 2006;87:27–34.

34. Ley RE, Hamady M, Lozupone C, Turnbaugh PJ, Ramey RR, Bircher JS, et al. Evolution of Mammals and Their Gut Microbes. Science (80-). 2008;320:1647.

35. McKenzie VJ, Song SJ, Delsuc F, Prest TL, Oliverio AM, Korpita TM, et al. The effects of captivity on the mammalian gut microbiome. Integr Comp Biol. 2017;57:690–704.

36. Phillips CD, Phelan G, Dowd SE, McDonough MM, Ferguson AW, Delton Hanson J, et al. Microbiome analysis among bats describes influences of host phylogeny, life history, physiology and geography. Mol Ecol. 2012;21:2617–27.

37. Groussin M, Mazel F, Sanders JG, Smillie CS, Lavergne S, Thuiller W, et al. Unraveling the processes shaping mammalian gut microbiomes over evolutionary time. Nat Commun. 2017;8:14319. doi:10.1038/ncomms14319.

38. Groussin M, Mazel F, Alm EJ. Co-evolution and Co-speciation of Host-Gut Bacteria Systems. Cell Host Microbe. 2020;28:12–22. doi:10.1016/j.chom.2020.06.013.

39. Godon JJ, Arulazhagan P, Steyer JP, Hamelin J. Vertebrate bacterial gut diversity: Size also matters. BMC Ecol. 2016;16:1–9.

40. Liu P, Cheng A, Huang S, Lu H, Oshida T. Body-size Scaling is Related to Gut Microbial Diversity, Metabolism and Dietary Niche of Arboreal Folivorous Flying Squirrels. Sci Rep. 2020;:1–12. doi:10.1038/s41598-020-64801-y.

41. Budd K, Gunn JC, Eggert LS, Finch T, Klymus K, Sitati N. Effects of diet, habitat, and phylogeny on the fecal microbiome of wild African savanna (Loxodonta africana) and forest elephants (L . cyclotis). Ecol Evol. 2020;00:1–14.

42. Metcalf JL, Song SJ, Morton JT, Weiss S, Seguin-Orlando A, Joly F, et al. Evaluating the impact of domestication and captivity on the horse gut microbiome. Sci Rep. 2017;7.

43. Illius AW, Gordon IJ. The functional significance of the browser-grazer dichotomy in African ruminants. Oecologia. 1994;98:167–75.

44. Miller EA, Livermore J, Alberts SC, Tung J, Archie EA. Ovarian cycling and
reproductive state shape the vaginal microbiota in wild baboons. Microbiome. 2017;5:1–
14.
45. Moeller AHA, Foerster S, Wilson ML, Pusey AE, Hahn BH, Ochman H. Social
behavior shapes the chimpanzee pan-microbiome. Sci Adv. 2016;2:e1500997–
e1500997. doi:10.1126/sciadv.1500997.
46. Johnson KVA. Gut microbiome composition and diversity are related to human
personality traits. Hum Microbiome J. 2020;15 December 2019:100069.
doi:10.1016/j.humic.2019.100069.
47. Shipley L. Grazers and browsers: how digestive morphology affects diet selection.
Grazing Behav Livest Wildl. 1999;:20–7.
http://www.cnr.uidaho.edu/range456/readings/shipley.pdf.
48. Venter JA, Vermeulen MM, Brooke CF. Feeding Ecology of Large Browsing and
Grazing Herbivores. In: Gordon IJ, Prins HHT, editors. The Ecology of Browsing and
Grazing II. Cham: Springer International Publishing; 2019. p. 127–53. doi:10.1007/978-
3-030-25865-8_5.
49. Hofmann RR. Evolutionary steps of ecophysiological adaptation and diversification
of ruminants: a comparative view of their digestive system. Oecologia. 1989;78:443–57.
doi:10.1007/BF00378733.
50. Spencer LM. Morphological Correlates of Dietary Resource Partitioning in the
African Bovidae. J Mammal. 1995;76:448–71. doi:10.2307/1382355.
51. Henderson G, Cox F, Ganesh S, Jonker A, Young W, Janssen PH, et al. Rumen
microbial community composition varies with diet and host, but a core microbiome is
found across a wide geographical range. Sci Rep. 2015;5:14567.
52. Edwards JE, Shetty SA, Berg P Van Den, Burden F, Doorn DA Van, Pellikaan WF,
et al. Multi-kingdom characterization of the core equine fecal microbiota based on
multiple equine (sub) species. Anim Microbiome. 2020;2:1–16.
53. Ilmberger N, Güllert S, Dannenberg J, Rabausch U, Torres J, Wemheuer B, et al. A
comparative metagenome survey of the fecal microbiota of a breast-and a plant-fed
asian elephant reveals an unexpectedly high diversity of glycoside hydrolase family
enzymes. PLoS One. 2014;9:1–12.
54. Roggenbuck M, Sauer C, Poulsen M, Bertelsen MF, Sørensen SJ. The giraffe
(Giraffa camelopardalis) rumen microbiome. FEMS Microbiol Ecol. 2014;90:237–46.
55. Tan SC, Chong CW, Yap IKS, Thong KL, Teh CSJ. Comparative assessment of
faecal microbial composition and metabonome of swine, farmers and human control.
Sci Rep. 2020;10:1–13.
56. Zeng B, Zhang S, Xu H, Kong F, Yu X, Wang P, et al. Gut microbiota of Tibetans
and Tibetan pigs varies between high and low altitude environments. Microbiol Res.
2020;235 November 2019:126447. doi:10.1016/j.micres.2020.126447.
57. Biddle A, Stewart L, Blanchard J, Leschine S. Untangling the genetic basis of
fibrolytic specialization by Lachnospiraceae and Ruminococcaceae in diverse gut
communities. Diversity. 2013;5.
58. Flint HJ, Scott KP, Duncan SH, Louis P, Forano E. Microbial degradation of complex
carbohydrates in the gut. Gut Microbes. 2012; August:289–306.
59. Kohl KD, Denise Dearing M. The woodrat gut microbiota as an experimental system
for understanding microbial metabolism of dietary toxins. Front Microbiol. 2016;7:1–9.

60. Coordinators NR. Database resources of the National Center for Biotechnology
Information. Nucleic Acids Res. 2018;46. doi:10.1093/nar/gkx1095.
61. Scheelings TF, Moore RJ, Van TTH, Klaassen M, Reina RD. Microbial symbiosis
and coevolution of an entire clade of ancient vertebrates: the gut microbiota of sea
turtles and its relationship to their phylogenetic history. Anim Microbiome. 2020;2:2–12.
62. Tinker KA, Ottesen EA. Phylosymbiosis across deeply diverging lineages in
omnivorous cockroaches. Appl Environ Microbiol. 2020;86:e02513-19.
doi:10.1128/AEM.02513-19.
63. Schmidt E, Mykytczuk N, Schulte-Hostedde AI. Effects of the captive and wild
environment on diversity of the gut microbiome of deer mice (Peromyscus maniculatus).
ISME J. 2019;13:1293–305. doi:10.1038/s41396-019-0345-8.
64. Perofsky AC, Lewis RJ, Meyers LA. Terrestriality and bacterial transfer: a
comparative study of gut microbiomes in sympatric Malagasy mammals. ISME J.
2019;13:50–63. doi:10.1038/s41396-018-0251-5.
65. Greene LK, Clayton JB, Rothman RS, Semel BP, Semel MA, Gillespie TR, et al.
Local habitat , not phylogenetic relatedness , predicts gut microbiota better within
folivorous than frugivorous lemur lineages. Biol Lett. 2019;15:5–11.
66. Gogarten JF, Davies TJ, Benjamino J, Gogarten JP, Graf J, Mielke A, et al. Factors
influencing bacterial microbiome composition in a wild non-human primate community in
Taï National Park, Côte d'Ivoire. ISME J. 2018;12:2559–74. doi:10.1038/s41396-018-
0166-1.
67. Amato KR, Mallott EK, Mcdonald D, Dominy NJ, Goldberg T, Lambert JE, et al.
Convergence of human and Old World monkey gut microbiomes demonstrates the
importance of human ecology over phylogeny. Genome Biol. 2019;20:201.
68. Gerardo NM, Hoang KL, Stoy KS. Evolution of animal Immunity in the light of
beneficial symbioses. Philos Trans R Soc B2. 2020;375:20190601.
69. Popkes M, Valenzano DR. Microbiota-host interactions shape ageing dynamics.
Philos Trans R Soc B2. 2020;375:20190596.
70. Grieneisen LE, Charpentier MJE, Alberts SC, Blekhman R, Bradbur G, Tung J, et
al. Genes, geology and germs: Gut microbiota across a primate hybrid zone are
explained by site soil properties, not host species. Proc R Soc B Biol Sci. 2019;286.
71. Muegge BD, Kuczynski J, Kinghts D, Clemente JC, Fontana L, Henrissat B, et al.
Diet drives Convergence in Gut Microbiome Functions Across Mammalian Phylogeny
and Within Humans. Science (80-). 2017;332:970–4.
72. Clemens ET, Maloiy GMO. Digestive physiology of East African wild ruminants.
Comp Biochem Physiol -- Part A Physiol. 1983;76:319–33.
73. Pérez-Barbería JF, Gordon IJ, Illius AW. Phylogenetic analysis of stomach
adaptation in digestive strategies in African ruminants. Oecologia. 2001;129:498–508.
74. Shorrocks B. The Biology of African Savannas. New York: Oxford University Press;
2007.
75. Wright ADG, Auckland CH, Lynn DH. Molecular diversity of methanogens in feedlot
cattle from Ontario and Prince Edward Island, Canada. Appl Environ Microbiol.
2007;73:4206–10. doi:10.1128/AEM.00103-07.
76. Daniel H, Gholami AM, Berry D, Desmarchelier C, Hahne H, Loh G, et al. High-fat
diet alters gut microbiota physiology in mice. ISME J. 2014;8:295–308.
doi:10.1038/ismej.2013.155.

77. Liu J, Pu YY, Xie Q, Wang JK, Liu JX. Pectin Induces an In Vitro Rumen Microbial
Population Shift Attributed to the Pectinolytic Treponema Group. Curr Microbiol.
2014;70:67–74. doi:10.1007/s00284-014-0672-y.
78. Xie X, Yang C, Guan LL, Wang J, Xue M, Liu JX. Persistence of Cellulolytic Bacteria
Fibrobacter and Treponema After Short-Term Corn Stover-Based Dietary Intervention
Reveals the Potential to Improve Rumen Fibrolytic Function. Front Microbiol. 2018;9
JUN:1363. doi:10.3389/fmicb.2018.01363.
79. Kennedy K, Hall MW, Lynch MDJ, Moreno-Hagelsieb G, Neufeld JD. Evaluating
bias of Illumina-based bacterial 16S rRNA gene profiles. Appl Environ Microbiol.
2014;80:5717–22.
80. De La Cuesta-Zuluaga J, Escobar JS. Considerations For Optimizing Microbiome
Analysis Using a Marker Gene. Front Nutr. 2016;3:26. doi:10.3389/fnut.2016.00026.
81. Mallott EK, Malhi RS, Amato KR. Assessing the comparability of different DNA
extraction and amplification methods in gut microbial community profiling. Access
Microbiol. 2019.
82. Green DS, Zipkin EF, Incorvaia DC, Holekamp KE. Long-term ecological changes
influence herbivore diversity and abundance inside a protected area in the Mara-
Serengeti ecosystem. Glob Ecol Conserv. 2019;20:e00697.
doi:10.1016/j.gecco.2019.e00697.
83. Kingdon J. East African Mammals: Volume I. Chicago: The University of Chicago
Press; 1984.
84. Kingdon J. East African Mammals: Volume IIIC. Chicago: The University of Chicago
Press; 1982.
85. Kingdon J. East African Mammals: Volume IIID. Chicago: The University of Chicago
Press; 1982.
86. Kingdon J. East African Mammals: Volume IIIB. Chicago: The University of Chicago
Press; 1979.
87. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al.
Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq
platforms. ISME J. 2012;6:1621–4. doi:10.1038/ismej.2012.8.
88. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. Development of a
Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon
Sequence Data on the MiSeq Illumina Sequencing Platform. Appl Environ Microbiol.
2013;79:5112–20. doi:10.1128/AEM.01043-13.
89. R Core Team. R: a language and environment for statistical computing. In: R
Foundation for Statistical Computing. Vienna, Austria; 2019.
https://www.gbif.org/tool/81287/r-a-language-and-environment-for-statistical-computing.
Accessed 28 Aug 2019.
90. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:
High-resolution sample inference from Illumina amplicon data. Nat Methods.
2016;13:581–3.
91. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA
ribosomal RNA gene database project: improved data processing and web-based tools.
Nucleic Acids Res. 2013;41:D590-6. doi:10.1093/nar/gks1219.
92. Wickham H. ggplot2: Elegant Graphics for Dat Analysis. New York, NY: Springer-
Verlag New York; 2009. doi:10.1007/978-0-387-98141-3.

93. Kolde R. Pheatmap: Pretty Heatmaps. R package version 1.0.12. 2019;:1–8.
https://cran.r-project.org/package=pheatmap.
94. De Cáceres M, Legendre P. Associations between species and groups of sites:
Indices and statistical inference. Ecology. 2009;90:3566–74.
95. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al.
Introducing mothur: open-source, platform-independent, community-supported software
for describing and comparing microbial communities. Appl Environ Microbiol.
2009;75:7537–41. doi:10.1128/AEM.01541-09.
96. Jin L, Wu D, Li C, Zhang A, Xiong Y, Wei R, et al. Bamboo nutrients and
microbiome affect gut microbiome of giant panda. Symbio. 2020;80:293–304.
97. Bo T-B, Zhang X-Y, Wen J, Deng K, Qin X-W, Wang D-H. The microbiota-gut-brain
interaction in regulating host metabolic adaptation to cold in male Brandt's voles
(Lasiopodomys brandtii). ISME J. 2019. doi:10.1038/s41396-019-0492-y.
98. McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. PLoS One. 2013;8:1–11.
99. Schliep KP. phangorn: Phylogenetic analysis in R. Bioinformatics. 2011;27:592–3.
100. Kembel SW, Cowan PD, Helmus MR, Cornwell WK, Morlon H, Ackerly DD, et al.
Picante: R tools for integrating phylogenies and ecology. Bioinformatics. 2010;26:1463–
4.
101. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using
lme4. J Stat Softw. 2015;67:1–48. doi:10.18637/jss.v067.i01.
102. Fox J, Weisberg S, Fox J. An R companion to applied regression. Thousand Oaks,
CA: Sage; 2011.
103. Oksanen J, Blanchet F, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan:
Community Ecology Package. R Packag version 24-6. 2018. https://cran.r-
project.org/web/packages/vegan/index.html. Accessed 28 Aug 2019.
104. Upham N, Esselstyn J, Jetz W. Inferring the mammal tree: species-level sets of
phylogenies for questions in ecology, evolution, and conservation. PLoS Biol.
2019;17:e3000494.
105. Paradis E, Claude J, Strimmer K. APE: Analyses of phylogenetics and evolution in
R language. Bioinformatics. 2004;20:289–90.
106. Ochieng EO. Characterizing the spatial distributions of elephants in Mpala, Kenya.
2015.
107. Kartzinel TR, Pringle RM. Multiple dimensions of dietary diversity in large
mammalian herbivores. J Anim Ecol. 2020;0:1–15.
108. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, et al.
Metagenomic biomarker discovery and explanation. Genome Biol. 2011;12:R60.
doi:10.1186/gb-2011-12-6-r60.
109. Afgan E, Baker D, Batut B, van den Beek M, Bouvier D, Čech M, et al. The Galaxy
platform for accessible, reproducible and collaborative biomedical analyses: 2018
update. Nucleic Acids Res. 2018;46:W537–44. doi:10.1093/nar/gky379.
1165