**Outer membrane lipoprotein DolP interacts with the BAM complex and**

**promotes fitness during envelope stress response**

David Ranava[1,§], Yiying Yang[1,§], Luis Orenday-Tapia[1,§], François Rousset[2],

Catherine Turlan[1], Lun Cui[2], Violette Morales[1], Cyril Moulin[1], Gladys Munoz[1],

Jérôme Rech[1], Anne Caumont-Sarcos[1], Cécile Albenne[1], David Bikard[2], Raffaele

Ieva[1*]

[1]Laboratoire de Microbiologie et Génétique Moléculaires (LMGM), Centre de

Biologie Intégrative (CBI), Université de Toulouse, CNRS, UPS, Toulouse 31062,

France.

[2]Synthetic Biology Group, Microbiology Department, Institut Pasteur, Paris 75015,

France.

[§]These authors contributed equally to this paper.

*Correspondance to :

Raffaele Ieva, raffaele.ieva@univ-tlse3.fr

**Abstract**

In Gram-negative bacteria, coordinated remodelling of the outer membrane (OM) and the peptidoglycan is crucial for envelope integrity. Envelope stress caused by unfolded OM proteins (OMPs) activates sigmaE($\sigma^E$) in Enterobacteria. Whereas $\sigma^E$ upregulates biogenesis factors, including the $\beta$-barrel assembly machinery (BAM) that catalyzes OMP-folding, elevated $\sigma^E$ activity can be detrimental for OM integrity. We report that DolP (YraP), a $\sigma^E$-upregulated OM lipoprotein needed for envelope integrity, is a novel interactor of BAM and that BamA is a critical determinant of the BAM-DolP complex. Mid-cell recruitment of DolP had been reported to promote septal peptidoglycan remodelling during cell division, but the role of DolP during envelope stress was unknown. Using a synthetic-defect CRISPRi screen, we found that *dolP* is critical in cells with elevated or constitutive $\sigma^E$ activation. Differently from other $\sigma^E$-upregulated proteins, DolP is not directly implicated in OMP assembly. Envelope stress or OM-overaccumulating BamA, however, interferes with DolP septal recruitment. Taken together, our results uncover DolP as a fitness factor during $\sigma^E$ activation, and support a model where differential septal recruitment of DolP contributes to link envelope stress to a late step of cell division.

## Introduction

The outer membrane (OM) of Gram-negative bacteria forms a protective barrier against harmful compounds, including several antimicrobials. This envelope structure surrounds the inner membrane and the periplasm that contains the peptidoglycan, a net-like structure made of glycan chains and interconnecting peptides. During cell division, the multi-layered envelope structure is remodelled by the divisome machinery [1]. At a late step of division, septal peptidoglycan synthesized by the divisome undergoes splitting, initiating the formation of the new poles of adjacent daughter cells. Finally, remodelling of the OM barrier completes formation of the new poles in the cell offspring. The mechanisms by which cells coordinate OM remodelling with peptidoglycan splitting, preserving the permeability barrier of this protective membrane, are ill-defined [2].

Integral outer membrane proteins (OMPs) are crucial to maintain the OM permeability barrier. OMPs fold into amphipathic β-barrel structures that span the OM and carry out a variety of tasks. Porins are OMPs that facilitate the diffusion of small metabolites. Other OMPs function as cofactor transporters, secretory channels, or machineries for the assembly of proteins and lipopolysaccharide (LPS) [3,4], a structural component of the external OM leaflet that prevents the diffusion of noxious chemicals [4]. The β-barrel assembly machinery (BAM) is a multi-subunit complex that mediates the folding and membrane insertion of OMPs transiting through the periplasm [5,6]. The essential and evolutionarily conserved BamA insertase subunit is an OMP consisting of an amino (N)-terminal periplasmic domain made of polypeptide transport-associated (POTRA or P) motifs and a carboxy (C)-terminal 16-stranded β-barrel membrane domain that catalyzes OMP biogenesis [7,8]. The flexible pairing of β-strands 1 and 16 of the BamA β-barrel controls a lateral gate connecting the interior of the barrel towards the surrounding lipid bilayer [9-12]. Conformational dynamics of the BamA β-barrel region proximal to the lateral gate is

thought to locally increase the entropy of the surrounding lipid bilayer [9,13,14] and

assist the insertion of nascent OMPs into the OM [11,15,16]. The N-terminal periplasmic

portion of BamA from the enterobacterium *Escherichia coli* contains five POTRA

motifs that serve as a scaffold for four lipoproteins, BamBCDE, which assist BamA

during OMP biogenesis [17-19]. The N-terminal POTRA motif is also the docking site of

the periplasmic chaperone SurA [20]. Together with the chaperones Skp and DegP,

SurA contributes to monitor unfolded OMPs transported into the periplasm by the

inner membrane general secretory (Sec) pathway [21,22].

Defective OMP assembly causes periplasmic accumulation of unfolded

protein transport intermediates. This envelope stress is signalled across the inner

membrane to induce a sigmaE ($\sigma^E$)-mediated transcriptional response [23]. In the

absence of stress, $\sigma^E$ is sequestered by the inner membrane-spanning RseA factor.

By-products of misfolded OMP turnover activate degradation of RseA, liberating $\sigma^E$

[24]. The $\sigma^E$ response copes with stress i) by upregulating genes involved in OMP

biogenesis, such as the *bam* genes [25], and ii) by lowering the OMP biogenesis

burden via a post-transcriptional mechanism [26]. Whereas $\sigma^E$ is essential [27], a tight

control of cytosolic $\sigma^E$ availability is necessary for optimal cell fitness and to prevent

a possible detrimental effect on the envelope [28-31]. Remarkably, the functions of a

number of genes upregulated by $\sigma^E$ remains unknown. Among those, *yraP* encodes

a ~20 kDa OM-anchored lipoprotein largely conserved in $\gamma$ and $\beta$ proteobacteria [32-

36]. YraP is crucial for OM integrity [34] and pathogenicity [32], and it was proposed to be

important for OMP biogenesis [34]. A recent study showed that YraP localizes at the

mid-cell during a late step of cell division, where it contributes to the regulation of

septal peptidoglycan splitting by an unknown mechanism [33]. These observations do

not explain why YraP is upregulated in response to $\sigma^E$ activation and how YraP

helps coping with envelope stress.

During cell division, envelope stress caused by defective OMP biogenesis

would place an additional burden on the already-complicated envelope reshaping

process faced by cells. It remains unknown whether stress caused by unfolded

OMPs influences envelope remodelling at the forming poles of two adjacent

daughter cells. Here we present a functional investigation of *yraP* prompted by a

genome-wide synthetic-defect screen. We demonstrate genetic interactions of *yraP*

with *bam* genes and we provide evidence that YraP can associate with the BAM

complex via an interaction with the BamA subunit. YraP is not critical for OMP

assembly but promotes fitness upon activation of the $\sigma^E$ response and under

conditions that impair secretion or assembly of OMPs in the OM. Upon envelope

stress or when BAM overaccumulates in the OM, YraP loses its association with the

mid-cell, thus suggesting that YraP contributes to link envelope stress response and

septal peptidoglycan hydrolysis during a late step of cell division. Hence, we

propose to rename YraP as DolP (underline{d}ivision- and underline{O}M stress-associated underline{l}ipoprotein).

**Results**

**A genome-wide synthetic-defect screen identifies *dolP* genetic interactions**

The mutant allele Δ*dolP::kan* [37] was introduced into *E. coli* BW25113 by P1

transduction. The resulting Δ*dolP* strain grew normally on LB medium but was

highly susceptible to vancomycin, an antibiotic normally excluded from the OM of

wild-type (*dolP*⁺) cells (Fig. 1A). The expression of C-terminally fused DolP variants

in place of the wild-type protein restored growth in the presence of vancomycin (Fig.

1A and Fig. S1B). This result supports the notion that DolP is important for envelope

integrity [33-36]. However, the role of DolP during envelope stress remains poorly

understood.

To gain insights into the function of DolP, we subjected Δ*dolP* cells to a

genome-wide synthetic-defect screen exploiting a Clustered Regularly Interspersed

Short Palindromic Repeat interference (CRISPRi) approach. Targeting of the

catalytically inactive dCas9 nuclease by gene-specific single guides RNAs

(sgRNAs) enables gene repression (Fig. 1B) [38]. The EcoWG1 sgRNA library

targeting the entire genome of *E. coli* MG1655 [39] was introduced into isogenic Δ*dolP*

or *dolP*⁺ MG1655-derivative strains. The fitness of each knockdown was then

compared in these backgrounds by deep-sequencing of the sgRNA library after ~17

growth generations. The outputs obtained from two independent tests were highly

reproducible (Fig. S2A). A strong fitness defect in the Δ*dolP* strain was caused by

the targeting of *envC* (Fig. 1C, Fig. S2B and Tables S1 and S2), followed by the

targeting of *ftsX* and *ftsE* (Fig. 1C, Tables S1 and S2). A validation growth test

showed that the synthetic fitness defect observed for Δ*dolP* cells was caused by

dCas9-dependent silencing of *ftsX* and *envC* (Fig. 1D, panels 6 and 7). The ABC

transporter-like complex FtsE/FtsX has multiple roles in organizing the cell

divisome, including the recruitment of periplasmic EnvC, a LytM domain-containing

factor required for the activation of amidases that hydrolase septal peptidoglycan [40].

This peptidoglycan remodelling reaction is mediated by two sets of highly controlled

and partially redundant amidases, AmiA/AmiB and AmiC [41,42]. Whereas AmiA and

AmiB are activated at the inner membrane/peptidoglycan interface by the

coordinated action of FtsE/FtsX and EnvC, activation of AmiC requires the OM-

anchored LytM domain-containing lipoprotein NlpD [43,44]. Under laboratory

conditions, the activity of only one of these two pathways is sufficient for septal

peptidoglycan splitting, whereas inhibition of both pathways leads to formation of

chains of partially divided cells, *i.e.* cells that have begun to divide but are blocked

at the step of septal peptidoglycan splitting [44]. A recent report showed that *dolP* is

necessary to complete septal peptidoglycan splitting and to promote cell separation

when the AmiA/AmiB pathway is inactive, somehow linking DolP to AmiC activation

[33]. Thus, the reduced fitness caused by silencing of *envC, ftsE or ftsX* in ∆*dolP* cells

(Fig. 1C) may be explained by impaired cell separation when both the AmiA/AmiB

and the AmiC pathways are not active. In keeping with this notion, *amiA* itself was

found among the negative fitness hits of the CRISPRi screen (Fig. 1C). *amiB* was

not a hit (Table S1), probably because AmiA is sufficient to split septal

peptidoglycan in the absence of other amidases [45].

Most importantly, the CRISPRi approach identified novel *dolP*-genetic

interactions that had a score similar to that obtained for *amiA* (Fig. 1C). These

included an interaction with *rseA*, encoding the inner membrane $\sigma^E$-sequestering

factor, as well as with *bamD*, encoding an essential subunit of the BAM complex

[34,46]. In accordance with the screen output, a serial dilution assay confirmed that

silencing of *bamD* (as well as *bamE*, encoding a stoichiometric interactor of BamD)

impairs growth of cells lacking DolP (Fig. 1D, panels 8 and 9). In addition, the

interaction of *dolP* with *rseA* was also confirmed in the genetic background of the

BW25113 strain (Fig. 1E). Further genes, mainly involved in OMP biogenesis and

homeostasis, or more generally in protein secretion, had a lower interaction score

(Table S1 and Fig. S2C) and are highlighted in Fig. 1C. These include genes

encoding the Sec ancillary complex SecDF-YajC, which contributes to the efficient secretion of proteins including OMPs [22] and was shown to interact with the BAM complex [47,48]. Collectively, the results of the CRISPRi screen indicate that the function of DolP is particularly critical for cell fitness upon inactivation of septal peptidoglycan hydrolysis by AmiA, as well as when OMP biogenesis is impaired.

**DolP does not play a crucial role in OMP biogenesis but improves fitness when the OM undergoes stress**

At a first glance, the newly identified genetic interaction between *dolP* and *bamD* (Figs. 1C and 1D) points to a possible role of DolP in OMP biogenesis. However, we found that the protein composition of the crude envelope fraction (Fig. S3A) and more specifically the assembly of two trimeric porins, LamB and OmpC (Figs. S3B and S3C) were not affected by the deletion of *dolP*. Similarly, the assembly kinetics of an autotransporter OMP was not impaired in Δ*dolP* cells (Fig. S3D). Importantly, cells lacking DolP had a normal $\sigma^E$ activity as the expression of $\sigma^E$-regulated genes, such as *bamA* or *bamE*, was not upregulated (Fig. 2A). Taken together, these observations suggest that DolP does not play a major function in OMP biogenesis. Yet, DolP promotes optimal cell fitness under envelope stress conditions. The genetic interaction of *dolP* with *bamD* or *surA* (Fig. 1C) may also suggest that DolP is required to optimally survive the envelope modifications consequent to activation of the $\sigma^E$ response in the corresponding deletion strains [34,46,49]. This hypothesis is supported by the observation that $\sigma^E$ activation caused by *rseA* silencing worsens the fitness of Δ*dolP* cells. We also noted that depletion of the $\sigma^E$-dependent *micA* gene, which encodes a small antisense RNA that reduces the expression of some OMPs during envelope stress, caused a negative fitness in wild-type but not in Δ*dolP* cells (Tables S1 and S2). Although this CRISPRi hit resulted from two independent experimental repeats, it is not highlighted in Figure 1C as the short nucleotide sequence of *micA* was targeted by only two sgRNAs (in

contrast to most genes that were targeted by 5 sgRNAs), and thus it was

considered as potentially non-significant by our statistical analysis. The fact that

$\Delta dolP$ tolerates lack of σE-dependent OMP downregulation would further support

the notion that the OMP assembly burden does not impair growth of this strain.

To further test the importance of DolP under envelope stress conditions, we

deleted *dolP* in a strain lacking *bamB,* which was identified with a lower genetic

score by the CRISPRi approach. In cells lacking BamB, the $\sigma^E$ response is partially

activated, causing the upregulation of *bam* genes (Fig. 2B). A strain carrying the

simultaneous deletion of *dolP* and *bamB* was viable but growth-defective (Fig. 2C).

Normal growth was restored by ectopic expression of a C-terminally, polyhistidine

tagged DolP protein variant (Fig. 2C). Phase-contrast analysis of the $\Delta dolP \Delta bamB$

strain revealed a number of cells with altered morphology or releasing vesicles as

well as several dead cells (Fig. 2D), indicating of a high degree of envelope stress

and possibly OM instability [50]. We conclude that although DolP is not crucial for

OMP biogenesis, its expression promotes optimal fitness probably by strengthening

OM integrity under the burden caused by envelope stress conditions and the

consequent $\sigma^E$-dependent response.

**DolP interacts with BamA assembled in the OM**

The observed interaction between *dolP* and *bam* genes prompted us to

investigate whether DolP physically interacts with the BAM complex. To this end, a

construct encoding C-terminally protein A-tagged DolP was ectopically

overproduced via the isopropylthiogalactoside (IPTG)-inducible *trc* promoter (P$_{trc}$).

The envelope of cells expressing DolP$^{ProtA}$ was solubilized using digitonin as main

mild-detergent component prior to IgG-affinity chromatography (Fig. 3A, coomassie

staining). Site-specific enzymatic cleavage of a protease-specific amino acid linker

between DolP and the protein A tag was used for protein elution. Notably, BamA,

BamC, BamD and BamE were immunodetected in the elution fraction of protein A-

tagged DolP (Fig. 3A, lane 3). In contrast, the inner membrane protein CyoA and cytosolic RpoB were not detected. We next adapted a previously established methodology and a $P_{trc}$-derivative construct [51] to ectopically overproduce BAM$^{ProtA}$ (consisting of wild-type BamABCD and a C-terminally protein A-tagged BamE protein variant) and isolate the BAM complex via IgG-affinity purification. Importantly, in addition to the BamE bait and other subunits of the BAM complex, DolP was also immunodetected in the elution fraction (Fig. 3B, lane 3). Other proteins of the bacterial envelope (Skp, OmpA, and $F_1\beta$ of the $F_1F_O$ ATP synthase) or cytosolic RpoB were not detected (Fig. 3B). Taken together, our native pull-down analysis indicates that DolP and BAM have affinity for each other.

To explore whether the central BAM subunit, BamA, is a critical determinant of the BAM-DolP interaction, we performed Ni-affinity purification using the solubilized envelope fraction obtained from cells overproducing BamA and C-terminally polyhistidine-tagged DolP. Under these conditions, BamA was efficiently co-eluted together with DolP$^{His}$, demonstrating that BamA and DolP can interact even in the absence of stoichiometric amounts of the BAM lipoproteins (Fig. 3C, coomassie staining). To assess if the interaction of DolP and BAM takes place at the OM, DolP$^{His}$ was overproduced together with an assembly-defective form of BamA, BamA$^{\Delta P1}$ [20]. When overproduced together with the BAM lipoproteins, BamA$^{\Delta P1}$ largely accumulated in the periplasm, failing to efficiently assemble into the BAM complex (Figs. S4A and S4B). When expressed together with DolP$^{His}$, assembly-defective BamA$^{\Delta P1}$ was highly depleted in the corresponding eluate (Fig. 3C, lane 7), even though BamA$^{\Delta P1}$ was only marginally reduced in the crude envelope fraction with respect to wild-type BamA (Fig. 3C, lane 3). In contrast to BamA$^{\Delta P1}$, the BamA$^{\Delta P2}$ variant, which is efficiently assembled into the OM (Fig. S4), was co-eluted with DolP$^{His}$ to a similar extent as wild-type BamA (Fig. 3C, lane 8). This result indicates that DolP has affinity for OM-assembled BamA.

In seeking a detergent that would interfere with the interaction of BAM and

DolP, and allow their purification as separate components, we solubilized the

envelope fraction with increasing amounts of n-dodecyl β-D-maltoside (DDM), a

detergent previously used to isolate the native BAM complex [51]. At concentrations of

DDM between 0.3% (w/v) and 1% (w/v), we observed a drastic reduction in the

amounts of BAM subunits that were co-eluted with DolP$^{His}$ (Fig. 3D), indicating that

the BAM-DolP interaction is sensitive to DDM. We thus used 1% (w/v) DDM to

extract and purify His-tagged DolP or His-tagged BAM as separate components

(Fig. 3E). When analyzed by blue native-PAGE and immunoblotting, purified DolP

gave rise to a diffused signal at around 450 kDa (Fig. 3F, lane 1) suggesting a

dynamic multimeric organization of this protein. Purified BAM migrated as expected

at 250 kDa (Fig. 3F, lane 7). When purified BAM and DolP were pre-incubated in

the presence of a low DDM concentration and subsequently resolved by blue

native-PAGE, a complex with an apparent molecular weight higher than that of the

BAM complex was detected with both anti-BamA and anti-DolP specific antibodies

(Fig. 3F, lanes 3 to 6). Taken together our results demonstrate that the BAM

complex and DolP interact to form a larger complex and that OM-assembled BamA

is a critical determinant of this interaction. Only a portion of BAM and DolP

associate to form the BAM-DolP complex, indicating that the interaction is

substoichiometric.

**269    BamA overaccumulation in the OM reduces DolP mid-cell localization**

In light of our observation that DolP interacts with BAM, we asked whether

the envelope localization patterns of DolP and BAM are reciprocally linked. First, we

monitored the effect of DolP expression on the localization of the chromosomally

encoded BamD$^{mCherry}$ subunit of the BAM complex. This protein generated a

fluorescence signal throughout the envelope that was not affected by the lack or the

overproduction of DolP (Fig. S5). Next, we checked the effect of BAM

overaccumulation in the OM on DolP localization. DolP associates with the OM and

accumulates at mid-cell during a late step of cell division [33]. To monitor the

localization of DolP, we used a strain harbouring a chromosomal *dolP-gfp* fusion

(Fig. 1A). The localization of the DolP$^{GFP}$ fusion protein (Fig. S6A) was analyzed

concomitantly to other two chromosomally-encoded markers of the division septum,

ZipA$^{mCherry}$ or NlpD$^{mCherry}$. ZipA is involved in an early step of divisome assembly

and accumulates at division sites before, as well as, during envelope constriction

(Fig. S6B) [52]. Instead, NlpD is a late marker of cell division involved in the activation

of AmiC and accumulates at septa that are already undergoing constriction (Fig.

S6C)[42,44]. DolP$^{GFP}$ accumulated at mid-cell sites where the envelope appeared

invaginated, showing a localization pattern similar to that of NlpD$^{mCherry}$ (Fig. S6C)

[33]. Strikingly, we found that ectopic BAM overproduction (Fig. S7A) depleted

DolP$^{GFP}$ from mid-cell sites (Fig. 4A, center). In contrast, no obvious effects on cell

division nor on mid-cell recruitment of ZipA$^{mCherry}$ and NlpD$^{mCherry}$ was observed (Fig.

S7B). The depletion of DolP at mid-cell sites was dependent on BamA

overaccumulation, as the overproduction of only the four BAM lipoproteins did not

prevent DolP$^{GFP}$ mid-cell localization (Fig. 4A, right). In another control experiment,

we observed that the overproduction of a different OM-spanning protein, OmpA, did

not prevent DolP mid-cell localization (Fig. S7C). To check whether the OMP-

biogenesis activity of BamA impairs DolP mid-cell localization, we made use of an

inactive BamA mutant form (Fig. S8A) harbouring a polyhistidine peptide extension

at its C-terminal β-strand[53]. Similar to the overaccumulation of the full BAM

complex, the overaccumulation of BamA$^{His}$ interfered with DolP mid-cell localization

(Figs. S8B and S8C), without affecting ZipA$^{mCherry}$ and NlpD$^{mCherry}$ (Fig. S7D),

indicating that the cellular localization of DolP does not depend on the OMP-

assembly activity of BamA. In contrast, the ability of BamA to assemble into the OM

was a critical determinant of the observed septal depletion of DolP. In fact, the

periplasm-accumulating BamA$^{\Delta P1/His}$ variant (Fig. S8D) did not impair DolP$^{GFP}$ mid-

cell localization (Fig. S8B, center), whereas the OM-overaccumulating BamA$^{\Delta P2/His}$

did (Figs. S8B, right, S8C, and S8D). Taken together, these results suggest that the

overaccumulation of BamA in the OM interferes with the recruitment of DolP at mid-

cell sites.

**DolP mid-cell localization is impaired under envelope stress conditions**

Because we observed that DolP is critical for fitness under envelope stress

conditions, we asked whether the localization of DolP$^{GFP}$ would be affected in

mutants than manifest envelope stress. To this end, we analyzed the localization of

DolP$^{GFP}$ in strains lacking either the OMP chaperone SurA or the lipoprotein BamB.

Both Δ*surA* and Δ*bamB* strains have an enhanced OM permeability and produce

higher levels of BAM complex due to activation of the $\sigma^E$ response [49,54] (Fig. 2B).

Importantly, the frequency of mid-cell labelling by DolP$^{GFP}$ was reduced by more

than half in Δ*surA* cells (Fig. 4B). As observed for the Δ*surA* strain, DolP$^{GFP}$

accumulated at the mid-cell with a lower frequency also when *bamB* was deleted

(Fig. S8E). Together, our results indicate that the depletion of DolP at mid-cell sites

occurs during the physiological response to conditions that cause envelope stress.

**Discussion**

Despite its importance in maintaining OM integrity, the reason why DolP is

upregulated by $\sigma^E$ has remained unknown [32-34]. Here we demonstrate an

unprecedented interaction of DolP with the BAM complex. Differently from other

factors upregulated upon $\sigma^E$ activation, DolP does not directly promote OMP

biogenesis. Instead, DolP is crucial for fitness in cells with elevated $\sigma^E$ activity and

its cell localization is sensitive to envelope stress as well as to the overaccumulation

of BAM in the OM. Our results have important implications for the understanding of

the mechanisms that coordinate OM and peptidoglycan remodelling and that help

preserving envelope integrity.

We have obtained several pieces of evidence suggesting that DolP is not

strictly required for OMP assembly into the OM under our experimental conditions.

Lack of DolP, in fact, does not reduce the steady state levels of OMPs, nor it

impairs the kinetics of assembly of an autotransporter virulence factor. Furthermore,

we could not detect activation of the $\sigma^E$ response in cells lacking DolP. Activation of

$\sigma^E$ has been reported to alleviate defects caused by lack or inactivation of OMP

biogenesis factors [55-58]. On the contrary, activation of $\sigma^E$ caused a fitness defect in a

$\Delta dolP$ strain. This result suggests that, under envelope stress conditions, DolP

plays a role different from OMP assembly and probably only indirectly related to

OMP biogenesis. In this respect, it is notable that the CRISPRi screen highlighted a

synthetic defect caused by lack of DolP and depletion of the Sec ancillary complex

SecDF-YajC, which is important for secretion of proteins across the inner

membrane, including OMPs [22]. Trans-envelope interactions of this complex with

BAM have been proposed to optimize the flux of unfolded OMPs across the

periplasm [47,48]. It is thus likely that conditions causing a depletion of OMPs in the

OM are particularly detrimental in cells lacking DolP.

Although the details of how DolP contributes to maintain OM integrity and

how it supports growth under envelope stress conditions will warrant further studies,

it is worth noting that DolP consists of two BON (bacterial OsmY and nodulation)

domains, folding motifs predicted to interact with hydrophobic ligands such as lipids

[59]. Phospholipid-binding by DolP has been reported in a recent study [60]. Two

aspects of our results hint at a plausible role of the lipid-binding function of DolP in

promoting fitness during envelope stress. The first is related to the biochemical

interaction of DolP with the BamA subunit of the BAM complex, which is predicted

to locally interfere with the organization of the OM lipid bilayer and to generate an

energetically favourable environment for the insertion of nascent OMPs [61,62]. BamA-

mediated membrane destabilization was shown by molecular dynamics simulations,

as well as by reconstituting BamA into proteoliposomes [9,63]. Thus, it is possible that

during envelope stress the increased amount of BamA has an impact on the

organization of the OM and contributes to some degree to the described detrimental

effect of elevated $\sigma^E$ activity [30,31]. It remains to be seen how DolP docks to BamA,

and whether it influences the organization of BAM sites and of the surrounding lipid

bilayer. The second aspect consists in *dolP* being a fitness factor in cells with an

active $\sigma^E$ response and reduced OMP biogenesis. OMPs represent a major

component of the OM and are likely to occupy nearly half of its surface[62]. OMPs are

largely arranged in clusters [64-66] embedded by highly organized LPS molecules in

the external leaflet of the OM [4] and the rigidity of their $\beta$-barrel structures contributes

to the mechanical stiffness of the OM [67]. The importance of OMPs for OM integrity

is highlighted by the fact that mutant strains lacking some of the major OMP

components (OmpA, OmpC and OmpF) present an enhanced OM permeability to

several antibiotics and detergents [30,68]. We propose that DolP plays a role in

preserving the structure of the OM, when BAM and OMP levels are inversely altered

by the activation of the $\sigma^E$ transcriptional response [25,54].

Another key result that emphasizes the importance of DolP during envelope

stress is the observation that septal localization of DolP is impaired under this

condition. Distinct biogenesis and surveillance pathways are required to maintain

the protective function of the multi-layered envelope of Gram-negative bacteria.

Emerging evidence is uncovering an intricate trans-envelope network of protein-

protein interactions that connect biogenesis machineries, stress-sensing factors and

their associated transducing pathways across the envelope layers [2,47,69-71]. DolP

septal localization was shown to be critical for efficient NlpD-mediated activation of

AmiC during septal peptidoglycan splitting [33]. Cells lacking both NlpD and AmiC

have reduced OM integrity [33], which may contribute to explain the vancomycin

sensitivity of $\Delta dolP$ cells. It is however unknown whether envelope stress caused by

the accumulation of unfolded OMPs influences peptidoglycan remodelling.

Intriguingly, our observation that mid-cell localization of DolP is reduced in *surA* or

*bamB* deletion mutant strains points to a role of DolP in linking envelope stress to

septal peptidoglycan hydrolysis. Whereas we have demonstrated that the

overaccumulation of BamA in the OM impairs the recruitment of DolP to the mid-

cell, it is possible that also other envelope features that are modified under

envelope stress conditions, for instance as a consequence of OMP depletion,

contribute to influence DolP localization. We suggest that reduced levels of DolP at

mid-cell sites, and thus impaired AmiC activation [33], can play an important role in

coping with envelope stress, for instance by regulating the window of time available

to restore normal OMP biogenesis prior to completing the formation of the new

poles in the cell offspring.

## Materials and Methods

### Bacterial strains and growth conditions

All *E. coli* strains used in this study are listed in Table S3. Strains newly generated for this study derive from BW25113 [Δ(*araD-araB*)*567* Δ(*rhaD-rhaB*)*568* Δ*lacZ4787* (::rrnB-3) *hsdR514 rph-1*] [72] or MG1655 (F⁻ λ⁻ *ilvG*⁻ *rfb-50 rph-1*) [73]. Deletions of *dolP, envC, rseA, surA,* or *bamB* in a BW25113 strain was achieved by P1 transduction of the Δ*dolP::kan^R^*, Δ*envC::kan^R^*, Δ*rseA::kan^R^*, Δ*surA::kan^R^* and Δ*bamB::kan^R^* alleles, respectively, obtained from the corresponding Keio collection strains [37]. BW25113 derivative strains harbouring chromosomal fusions of constructs encoding superfolder GFP downstream of *dolP* or mCherry downstream of *nlpD*, *zipA* and *bamD* were obtained by λ-red recombination as previously described [74]. Briefly, a kanamycin-resistance cassette was amplified from plasmid pKD4 using oligonucleotides carrying extensions of approximately 50 nucleotides homologous to regions immediately upstream or downstream the stop codon of the interested gene. *Dpn*I-digested and purified PCR products were electroporated into the BW25113 or derivative strains. Recombinant clones were selected at 37°C on LB agar plates containing kanamycin. When necessary, the *kan^R^* cassette inserted into a mutated locus (gene deletion or fusion) was removed upon transformation with the heat-curable plasmid pCP20 [74]. The MG1655 derivative strain, LC-E75, harbouring a dCas9-encoding construct under the control of the P*tet* promoter has been described [38]. Cells were cultured in home-made lysogeny broth (LB) medium (1% (w/v) tryptone, 0.5% (w/v) yeast extract, 5 mg/ml (NaCl), commercially available Miller LB Broth (Sigma) or M9 minimal medium containing M9 salts (33.7 mM $Na_2HPO_4$, 22 mM $KH_2PO_4$, 8.55 mM NaCl, 9.35 mM $NH_4Cl$) and supplemented with 0.2% w/v glycerol and all the amino acids, or all amino acids except methionine and cysteine in the case of $^{35}S$ pulse-chase labeling. Antibiotics were used at the following concentrations: ampicillin 100 μg/ml, kanamycin 50 μg/ml, vancomycin 60

μg/ml. For spot tests, cells were cultured to mid-log phase, washed with M9 salts

and serially diluted in ice-cold M9 salts prior to spotting on agar plates.

**Plasmid construction**

All plasmids used in data figures are listed in Table S4. Plasmids for the

ectopic expression of BAM subunits, DolP, or OmpA are derived from a pTRC99a

vector. The plasmid pBAM$^{His}$ (pJH114), which harbours a P$_{trc}$ promoter followed by

the *bamABCDE* open reading frames and an octahistidine tag fused downstream of

*bamE* was described [51]. The region of pBAM$^{His}$ comprising the segment that spans

from the *bamA* start codon to the *bamE* stop codon was deleted by site-directed

mutagenesis, generating pCtrl. Plasmids pBamA$^{His}$ was generated by restriction-

free cloning inserting the *bamA* ORF without its stop codon downstream of the P$_{trc}$

promoter and upstream of the octahistidine encoding region in pCtrl. The *dolP* ORF

amplified from the BW25113 genomic DNA was used to replace the *bamABCDE*

ORFs in pJH114 by restriction-free cloning, generating pDolP$^{His}$. The plasmid

pBamA-DolP$^{His}$ was generated by restriction-free cloning of the *bamA* ORF between

the P$_{trc}$ promoter and *dolP* in pDolP$^{His}$. Site-directed mutagenesis was conducted on

pBAM$^{His}$ (pJH114) to obtain pBamBCDE$^{His}$. A sequence encoding the tobacco etch

virus protease cleavage site (TEV site) followed by a tandem Protein-A tag was

amplified from pYM10 [75] and fused by restriction-free cloning with the last codon of

the *bamE* gene in pBAM$^{His}$ to generate pBAM$^{ProtA}$. A stop codon was introduced

downstream of the *bamE* last codon to generate pBAM. Plasmids encoding the ΔP1

and ΔP2 BamA variant were obtained by site-directed mutagenesis deleting the

portion of *bamA* ORFs corresponding to residues E22-K89 or P92-G172,

respectively. The TEV site and the tandem Protein-A construct amplified from

pYM10 was inserted by restriction-free cloning downstream of the *dolP* last codon in

pDolP$^{His}$, generating pDolP$^{ProtA}$. pDolP was derived from pDolP$^{ProtA}$ using site-

directed mutagenesis to introduce a stop codon immediately downstream of the

*dolP* ORF. The *ompA* ORF was amplified from the BW25113 genomic DNA and

inserted by restriction-free cloning between Ptrc and the His-tag encoding construct

of pCtrl to generate pOmpA[His]. The sgRNAs plasmids are derived from psgRNAcos

[38]. To generate sgRNA-encoding plasmids the DNA sequences

AGCTGCACCTGCTGCGAATA (*bamD* sgRNA, plasmid pCAT187),

GTAAACCACTCGCTCCAGAG (*bamE* sgRNA, plasmid pCAT189),

CTCATCCGCGTGGGCGGAAA (*envC* sgRNA, plasmid pCAT191), and

CTGAGCCGCCGACCGATTTA (*ftsX* sgRNA, pCAT193) were inserted into a *Bsa*I

site of the psgRNAcos.

**CRISPRi screen and data analysis**

Strain LC-E75 (*dolP*[+]) and its Δ*dolP* derivative were transformed with the

EcoWG1 library which contains 5 guides per gene as previously described [39]. After

culturing pooled transformant cells in LB at 37°C to early exponential phase (optical

density at 600 nm [$OD_{600}$] = 0.2), a sample was withdrawn for plasmid isolation

($t_{start}$). Subsequently, cultures were supplemented with 1 μM anhydrotetracycline

(aTc) to induce dCas9 expression and further incubated at 37°C. When cultures

reached an $OD_{600}$ of 2 they were diluted 1:100 into LB supplemented with 1 μM aTc

and incubated at the same temperature until an $OD_{600}$ of 2. This step was repeated

one more time prior to withdrawing a sample for isolation of plasmid DNA ($t_{end}$).

Sequencing indexes were used to assign reads to each sample. Illumina

sequencing samples were prepared and analyzed as previously described [38].

Briefly, a two-step PCR was performed with Phusion polymerase (Thermo

Scientific) using indexed primers. The first PCR adds the first index and the second

PCR adds the second index and flow-cell attachment sequences. Pooled PCR

products were gel-purified. Sequencing was performed on a NextSeq550 machine

(Illumina). The total number of reads obtained for each sample was used to

normalize raw reads by sample size. Replicates were pooled to increase depth

before another normalization by sample size. Guides with less than 100 normalized

read counts in initial time points were discarded. For each screen, sgRNA fitness

was calculated as the log2-transformed ratio of normalized reads counts between

the final and the initial time point:

$$log2FC = log2\left(\frac{Normalized\ reads_{final} + 1}{Normalized\ reads_{initial} + 1}\right)$$

For each sample, log2FC values were centered by subtracting the median log2FC

of non-targeting control guides. We then calculated for each sgRNA the difference

of log2FC value between the $\Delta dolP$ screen and the $dolP^+$ screen. Guides were

ranked from the lowest negative values (negative fitness effect in $\Delta dolP$ compared

to $dolP^+$) to the highest positive values (positive fitness effect in $\Delta dolP$ compared to

$dolP^+$) and the significance of the interaction between $dolP$ and each gene was

evaluated by performing a minimum hypergeometric (mHG) test on the ranked list

for each gene using the mHG R package (v. 1.1) [76]. False-discovery rate (FDR) was

used to correct p-values for multiple testing. For each gene, the median difference

of log2FC between $\Delta dolP$ and $dolP^+$ screens was used as a measure of the genetic

interaction.

## Cell fractionation

To prepare whole-cell lysates, cells were cultured to early exponential phase

($OD_{600}$ = 0.2-0.3) in LB medium at 37°C and collected. Where indicated, IPTG was

added 30 min prior to cell collection. Cells were pelleted by centrifugation, washed

once with M9 salt and lysed with Laemmli Sample Buffer (BioRad) (69 mM Tris-HCl,

pH 6.8, 11.1% [v/v] glycerol, 1.1% [w/v] lithium dodecyl sulfate [LDS], 0.005% [w/v]

bromophenol blue, supplemented with 357 mM β-mercaptoethanol and 2 mM

phenylmethylsulfonyl fluoride [PMSF]). The total protein content was denatured at

98°C for 5 min prior SDS-PAGE analysis.

To perform lysozyme/EDTA lysis, cells were cultured to early exponential phase,

collected by centrifugation, resuspended in 33 mM Tris-HCl pH 8 to an $OD_{600}$ of 1.

The cell suspension was then supplemented with 0.1 mg/ml lysozyme (Sigma), 2

mM EDTA and incubated on ice for 20 minutes to induce lysis. After addition of 10

mM $MgSO_4$, the membrane fraction was collected by centrifugation at 16000x $g$. The

supernatant was further centrifuged at 100000x $g$ to remove any residual

membrane fraction, which was discarded. The obtained soluble fraction was

subjected to protein precipitation by adding 10% (w/v) trichloroacetic acid (TCA).

TCA precipitates were solubilized in Laemmli Sample Buffer prior to SDS-PAGE

analysis.

The crude envelope fraction used for native affinity purification of affinity

tagged BAM complex or DolP was prepared from cells that were cultured in LB until

an $OD_{600}$ of approximately 0.25 and then supplemented with 400 $\mu$M IPTG for 1

hour to induce ectopic expression. Cells were collected by centrifugation at 6000x $g$

at 4°C, resuspended in 20 mM Tris-HCl pH 8, and mechanically disrupted using a

Cell Disruptor (Constant Systems LTD) set to 0.82 kPa. The obtained cell lysate

fractions were clarified by centrifugation at 6000x $g$ and 4°C. The supernatant was

then subjected to ultracentrifugation at 100000x $g$ at 4°C to collect the envelope

fraction.

**Isolation of native protein complexes by IgG- or nickel-affinity**

**chromatography**

The envelope fraction was resuspended at a concentration of approximately

10 mg/ml in solubilization buffer (20 mM Tris-HCl pH 7.4, 100 mM NaCl, 0.1 mM

EDTA, 2 mM PMSF) supplemented with EDTA-free protease inhibitor cocktail

(Roche), and 1.1% (w/v) of a mild detergent component corresponding to digitonin

(Merck) and n-dodecyl $\beta$-D-maltoside (DDM, Merck) as indicated in Figure Legends.

To facilitate extraction of membrane proteins, samples were subjected to mild

agitation for 1 hour at 4°C. Insoluble material was removed by centrifugation at

16000x $g$ at 4°C. To perform IgG affinity purification, membrane-extracted proteins

were incubated for 1.5 hours at 4°C with purified human IgG (Sigma) that had been

previously coupled to CNBr-activated Sepharose beads (GE Healthcare). After

extensive washes of the resin with solubilization buffer containing 0.3% (w/v)

digitonin and 0.03% (w/v) DDM, bound proteins were eluted by incubation with

AcTEV protease (ThermoFisher) overnight at 4°C under mild agitation. To perform

nickel (Ni)-affinity purification, membrane-extracted proteins were supplemented

with 20 mM imidazole and incubated with Protino Ni-NTA agarose beads

(Macherey-Nagel) for 1 hour at 4°C. After extensive washes of the resin with

solubilization buffer supplemented with 50 mM imidazole, 0.3% (w/v) digitonin,

0.03% (w/v) DDM, and the EDTA-free protease inhibitor cocktail (Roche), bound

proteins were eluted using the same buffer supplemented with 500 mM imidazole.

***In vitro* reconstitution of the BAM-DolP interaction and BN-PAGE analysis**

Envelope fractions were obtained from cells carrying pBAM[His] or pDolP[His]

and cultured until early exponential phase in LB medium at 37°C and subsequently

supplemented with 400 μM IPTG for 1.5 hours to induce the expression of the BAM

complex genes or *dolP*. The envelope fractions were solubilized and purified by Ni-

affinity and size exclusion chromatography, adapting a previously published

protocol. Briefly, after membrane solubilization with 50 mM Tris-HCl pH 8.0, 150

mM NaCl, and 1% (w/v) DDM, and removal of insoluble material by

ultracentrifugation at 100000x $g$, 4°C, soluble proteins were loaded onto a Ni-

column (HisTrap FF Crude, GE Healthcare) pre-equilibrated with 50 mM Tris-HCl

pH 8.0, 150 mM NaCl, and 0.03% (w/v) DDM (equilibration buffer), using an ÄKTA

Purifier 10 (GE Healthcare) at 4°C. The column containing bound proteins was

washed with equilibration buffer supplemented with 50 mM imidazole. Proteins were

eluted in equilibration buffer, applying a gradient of imidazole from 50 mM to 500

mM and further separated by gel filtration using an HiLoad 16/600 Superdex 200

(GE Healthcare) in equilibration buffer. Eluted proteins were concentrated using an

ultrafiltration membrane with a 10 kDa molecular weight cutoff (Vivaspin 6,

Sartorius). To reconstitute the BAM-DolP complex *in vitro,* equimolar concentrations

of purified BAM and DolP were used. Purified proteins were mixed in equilibration

buffer for 1 hour at 4°C or for 30 min at 25°C as indicated in Figure Legends. The

reaction was further diluted 1:4 times in ice-cold blue native buffer (20 mM Tris-HCl

pH 7.4, 50 mM NaCl, 0.1 mM EDTA, 1% [w/v] digitonin, 10% w/v glycerol) and ice

cold blue native loading buffer (5% coomassie brilliant blue G-250, 100 mM Bis-

Tris-HCl, pH 7.0, 500 mM 6-aminocaproic acid) prior to loading onto home-made 5-

13% blue native polyacrylamide gradient gels. Resolved protein complexes were

blotted onto a PVDF membrane and immunodecorated. Where non-relevant gel

lanes were removed, a white space was used to separate contiguous parts of the

same gel.

**Pulse-chase *in vivo* protein labeling**

Pulse-chase labeling of cellular proteins with radioactive $^{35}$S methionine and

cysteine was conducted as previously described [77]. Briefly, overnight cultures were

washed and freshly diluted into M9 medium to $OD_{600}$ = 0.03 and incubated at 37 °C.

When the cultures reached $OD_{600}$ = 0.3, ectopic expression of EspP was induced by

adding 200 μM IPTG for 30 min. $^{35}$S labelling was performed by supplementing

cultures with a mixture of [$^{35}$S]-methionine and [$^{35}$S]-cysteine followed after 30

seconds by the addition of an excess of cold methionine and cysteine. Samples

were withdrawn at different chase times, immediately mixed with ice and subjected

to TCA precipitation, prior to immunoprecipitation.

**Antibodies**

Proteins separated by SDS-PAGE or blue native-PAGE and blotted onto

PVDF membranes were immunodecorated using epitope-specific rabbit polyclonal

antisera with the following exceptions. The $F_1\beta$ subunit of the ATP $F_1F_O$ synthase

was detected using a rabbit polyclonal antiserum raised against an epitope of the

homologous protein of Saccharomyces cerevisiae (Atp2). RpoB was detected using

a mouse monoclonal antibody (NeoClone Biotechnology). The secondary

immunodecoration was conducted using anti-rabbit or anti-mouse peroxidase-

conjugated antibodies produced in goat (Sigma). EspP immunoprecipitation was

conducted using a rabbit polyclonal antiserum specific for a C-terminal epitope of

EspP.

**Epifluorescence microscopy and analysis**

Overnight cultures of *E. coli* BW25113 and its derivative strains were diluted

into fresh M9 medium containing 0.2% glycerol and grown at 30°C to $OD_{600}$ = 0.2-

0.3. When indicated, cultures were supplemented with 400 μM of IPTG to induce

ectopic expression of plasmid-borne genes for at least 1 hour prior to collecting

samples for microscopy analysis. Culture volumes of 0.6 μl were deposited directly

onto slides coated with 1% (w/v) agarose in a phosphate-buffered saline solution

and visualized by epifluorescence microscopy. Cells were imaged at 30°C using an

Eclipse TI-E/B Nikon wide field epifluorescence inverted microscope with a phase

contrast objective (Plan APO LBDA 100X oil NA1.4 Phase) and a Semrock filter

mCherry (Ex: 562BP24; DM: 593; Em: 641BP75) or FITC (Ex: 482BP35; DM: 506;

Em: 536BP40). Images were acquired using a CDD OrcaR2 (Hamamatsu) camera

with illumination at 100% from a HG Intensilight source and with an exposure time

of 1-3 seconds. Nis-Elements AR software (Nikon) was used for image capture.

Image analysis was conducted using the Fiji software. Counting of cells with

DolP[GPF] signals at mid-cell sites was performed using the Fiji Cell Counter plugin.

**Acknowledgments**

We thank Harris Bernstein (NIDDK/NIH, Bethesda, MD) for providing key reagents and for comments on the manuscript. We thank Tanneke Den Blaauwen (University of Amsterdam) and Nathalie Dautin (IBPC/CNRS, Paris) for discussion. We thank the Light Imaging Toulouse CBI (LITC) platform for assistance with and maintenance of the microscopy instrumentation. Financial support was provided by: the Fondation pour la Recherche Médicale to D.R.; the Chinese Scholarship Council to Y.Y.; the Ecole Normale Supérieure to F.R.; the European Research Council (Europe Union's Horizon 2020 research and innovation program, grant agreement No 677823), the French governmental Investissement d'Avenir program and the Laboratoire d'Excellence "Integrative Biology of Emerging Infectious Diseases" (ANR-10-LABX-62-IBEID) to D.B.; the CNRS ATIP-Avenir program to R.I.

**Author contribution**

R.I. conceived the project and wrote the manuscript; D.B. conceived and developed the CRISPRi approach; R.I., D.B., D.R. L.C. and C.A. supervised the experiments; D.R., Y.Y., L.O., F.R., L.C., V.M., C.M., G.M., A.C.-S, C.T., J.R. and C.A. performed the experiments and analyzed the data together with R.I. and D.B.; A.C.-S, D.R., Y.Y., L.O. and F.R. prepared the figures; all authors discussed the project and the experimental results, and commented on the manuscript.

642 **Figures and Figure Legends**

**Figure 1.**

**Fig. 1. Genome-wide screen of *dolP* interactions.**

(**A**) The deletion of *dolP* impairs OM integrity. The indicated strains were serially

diluted and spotted onto LB agar plates lacking or supplemented with 60 μg/ml

vancomycin as indicated.

(**B**) Schematic representation of the CRISPR-based gene silencing approach. LC-

E75 (*dolP*$^+$) or its Δ*dolP* derivative strain, both carrying d*cas9* under the control of

an anhydrotetracycline (aTc)-inducible promoter in their chromosome were

transformed with a library of plasmids encoding gene-specific sgRNAs. The library

covers any *E. coli* MG1655 genetic features with an average of 5 sgRNAs per gene.

Pooled transformed cells were cultured to early exponential phase prior to plasmid

extraction and quantitative Illumina sequencing to assess the initial distribution of

sgRNA constructs in each culture (t$_{start}$). Upon addition of 1 μM aTc to induce

sgRNA-mediated targeting of d*cas9* for approximately 17 generations, samples of

cells from each culture were newly subjected to plasmid extraction and Illumina

sequencing to determine the final distribution of sgRNA constructs (t$_{end}$).

(**C**) Left: Comparison of gene scores obtained in *dolP*$^+$ and Δ*dolP* screens. The log2

fold-change (log2FC) between t$_{end}$ and t$_{start}$ calculated for each sgRNAs (Fig. S2B)

was grouped by gene target, and their median (Table S1) was used to derive fitness

gene scores. Right: Volcano plot of the *dolP* genetic interaction (GI) scores. The x-

axis shows a genetic interaction score calculated for each gene based on the

minimum hypergeometric (mHG) test conducted on the ranked difference of

sgRNA-specific log2FC values between the Δ*dolP* and the *dolP*$^+$ screens. The y-

axis shows the log10 of the false discovery rate FDR of the test. The dashed line

shows FDR = 0.05. In both panels, genes highlighted in orange have FDR < 0.05

and GI>1 whereas genes highlighted in red have FDR < 0.05 and GI>2.

(**D** and **E**) Validation of the genetic interactions determined in (C). (D) LC-E75

(*dolP*$^+$) or its Δ*dolP* derivative strain expressing sgRNAs that target the indicated

genes were serially diluted and spotted on LB agar lacking or supplemented with

aTc to induce expression of d*cas9*, as indicated. (E) BW25113 derivative cells

deleted of *rseA* or both *rseA* and *dolP* were transformed with an empty vector

(pCtrl) or a plasmid encoding DolP (pDolP$^{His}$). Ectopic expression of DolP$^{His}$ was

driven by the leaky transcriptional activity of P$_{trc}$ in the absence of IPTG. (D and E)

10-fold serial dilutions of the indicated transformants were spotted on LB agar.

**Figure 2.**

**Fig. 2. DolP promotes fitness under envelope stress conditions.**

(A) One and three folds of a sample containing the total cell lysate fraction from a BW25113 (*dolP*[+]) strain and a derivative Δ*dolP* strain were analyzed by SDS-PAGE and immunoblotting using the indicated antisera.

(B) One, two and three folds of a sample containing the total cell lysate fraction from a BW25113 (*bamB*[+]) strain and a derivative Δ*bamB* strain were analyzed by SDS-PAGE and immunoblotting using the indicated antisera.

(C) BW25113 and derivative cells deleted of *dolP, bamB* or both genes were cultured, serially diluted and spotted on LB agar. A BW25113 derivative strain deleted of both *dolP* and *bamB* was cultured, serially diluted and spotted on LB agar supplemented with ampicillin.

(D) Overnight cultures of BW25113 (control), Δ*dolP*, Δ*bamB and* Δ*dolP* Δ*bamB,* were freshly diluted in LB medium and re-incubated at 30°C until $OD_{600}$ = 0.3. Cell samples were visualized on 1% (w/v) agarose pads by phase contrast microscopy. Bar = 5 μm.

**Figure 3.**

**Fig. 3. DolP associates with the BAM complex via an interaction with BamA.**

(**A**, **B**). The envelope fraction of BW25113 cells carrying the indicated plasmids were solubilized with 1% (w/v) digitonin and 0.1% (w/v) DDM and subjected to IgG affinity purification of either protein A-tagged DolP (A) or BamE (B). The load and elution fractions were analyzed by SDS-PAGE. The coomassie staining of the elution of protein A-tagged DolP (A) or protein A-tagged BamE (B) are shown below the diagrams representing overproduced proteins. Blotted protein from load and elution fractions were detected by immunodecoration using the indicated antisera. Load 0.5% (A) 1% (B); Elution 100%. The asterisk indicates the products of TEV digestion obtained with samples containing DolP$^{ProtA}$ (A) or BamE$^{ProtA}$ (B).

(**C**) The envelope fraction of BW25113 cells carrying the plasmids overproducing His-tagged DolP and the indicated BamA protein variants (deleted of POTRA1 or of POTRA2) were solubilized with 1% (w/v) digitonin and 0.1% (w/v) DDM and subjected to Ni-affinity purification. The load and elution fractions were analyzed by SDS-PAGE. The coomassie staining of the elution of His-tagged DolP overproduced together with wild-type BamA is shown below the diagram representing overproduced proteins. Blotted protein from load and elution fractions were detected by immunodecoration using the indicated antisera. Load 2%; Elution 100%.

(**D**) Equal aliquots of the envelope fraction of BW25113 cells expressing His-tagged DolP were solubilized with the indicated concentrations of digitonin and DDM (digitonin [w/v], DDM [w/v]): 1%, 0.1% (lane 1), 0.8%, 0.3% (lane 2), 0.3%, 0.8% (lane 3), 0.1%, 1% (lane 4). His-tagged DolP was purified by Ni-affinity chromatography. In all cases, proteins were eluted in the presence of 0.3% (w/v) digitonin and 0.03% (w/v) DDM. Load 0.2%; Elution 100%.

(**E**) The envelope fraction of BW25113 cells overproducing DolP$^{His}$ or the BAM complex containing C-terminally His-tagged BamE was subjected to protein extraction with 1% (w/v) DDM and Ni-affinity purification and gel filtration

chromatographies. The elution fractions were analyzed by SDS-PAGE and

coomassie staining. The double asterisk indicates a contaminant protein in the

elution of DolP.

(**F**) Purified His-tagged BAM complex and DolP were incubated alone for 1 hour at

4°C (lanes 1, 2 and 7), or together for 1 hour at 4°C (lanes 3 and 6) or for 30 min at

25°C (lanes 4 and 5), prior to blue native-PAGE and immunoblotting using the

indicated antisera.

**Figure 4.**

**Fig. 4. BamA overaccumulation in the OM and envelope stress interfere with**

**mid-cell localization of DolP.**

(**A**) Overnight cultures of BW25113 cells harboring the chromosomal fusion *dolP-*

*gfp* and transformed with either pCtrl (empty vector) or pBAM$^{His}$, or pBamBCDE$^{His}$

were freshly diluted in minimal M9 medium, incubated at 30°C until $OD_{600}$ = 0.1 and

supplemented with 400 $\mu$M IPTG for 1 hour. Cell samples were visualized on 1%

(w/v) agarose pads by phase contrast and fluorescence microscopy. Arrowheads

indicate envelope constriction sites between forming daughter cells. Bar = 5 $\mu$m.

(**B**) Overnight cultures of BW25113 (control) or $\Delta surA$ derivative cells carrying the

*dolP-gfp* chromosomal fusion were freshly diluted in LB medium and incubated at

30°C until $OD_{600}$ = 0.3. Cell samples were visualized as in (A). Bar = 5 $\mu$m. The

percentage of cells presenting fluorescent DolP$^{GFP}$ signals at division septa in $\Delta surA$

cells was normalized to the same fraction obtained for the control cells and reported

below the micrograph. More than 1000 cells were counted in each sample.

**References**

1 den Blaauwen, T., Hamoen, L. W. & Levin, P. A. The divisome at 25: the
road ahead. *Current opinion in microbiology* **36**, 85-94,
doi:10.1016/j.mib.2017.01.007 (2017).
2 Egan, A. J. F., Errington, J. & Vollmer, W. Regulation of peptidoglycan
synthesis and remodelling. *Nature reviews. Microbiology*,
doi:10.1038/s41579-020-0366-3 (2020).
3 Calmettes, C., Judd, A. & Moraes, T. F. Structural Aspects of Bacterial Outer
Membrane Protein Assembly. *Advances in experimental medicine and*
*biology* **883**, 255-270, doi:10.1007/978-3-319-23603-2_14 (2015).
4 Nikaido, H. Molecular basis of bacterial outer membrane permeability
revisited. *Microbiology and molecular biology reviews : MMBR* **67**, 593-656,
doi:10.1128/mmbr.67.4.593-656.2003 (2003).
5 Ranava, D., Caumont-Sarcos, A., Albenne, C. & Ieva, R. Bacterial
machineries for the assembly of membrane-embedded beta-barrel proteins.
*FEMS microbiology letters* **365**, doi:10.1093/femsle/fny087 (2018).
6 Schiffrin, B., Brockwell, D. J. & Radford, S. E. Outer membrane protein
folding from an energy landscape perspective. *BMC biology* **15**, 123,
doi:10.1186/s12915-017-0464-5 (2017).
7 Webb, C. T., Heinz, E. & Lithgow, T. Evolution of the beta-barrel assembly
machinery. *Trends in microbiology* **20**, 612-620,
doi:10.1016/j.tim.2012.08.006 (2012).
8 Voulhoux, R., Bos, M. P., Geurtsen, J., Mols, M. & Tommassen, J. Role of a
highly conserved bacterial protein in outer membrane protein assembly.
*Science (New York, N.Y.)* **299**, 262-265, doi:10.1126/science.1078973
(2003).
9 Noinaj, N., Kuszak, A. J., Gumbart, J. C., Lukacik, P., Chang, H., Easley, N.
C., Lithgow, T. & Buchanan, S. K. Structural insight into the biogenesis of
beta-barrel membrane proteins. *Nature* **501**, 385-390,
doi:10.1038/nature12521 (2013).
10 Iadanza, M. G., Higgins, A. J., Schiffrin, B., Calabrese, A. N., Brockwell, D.
J., Ashcroft, A. E., Radford, S. E. & Ranson, N. A. Lateral opening in the
intact beta-barrel assembly machinery captured by cryo-EM. *Nature*
*communications* **7**, 12865, doi:10.1038/ncomms12865 (2016).
11 Gu, Y., Li, H., Dong, H., Zeng, Y., Zhang, Z., Paterson, N. G., Stansfeld, P.
J., Wang, Z., Zhang, Y., Wang, W. & Dong, C. Structural basis of outer
membrane protein insertion by the BAM complex. *Nature* **531**, 64-69,
doi:10.1038/nature17199 (2016).
12 Bakelar, J., Buchanan, S. K. & Noinaj, N. The structure of the beta-barrel
assembly machinery complex. *Science (New York, N.Y.)* **351**, 180-186,
doi:10.1126/science.aad3460 (2016).
13 Doerner, P. A. & Sousa, M. C. Extreme Dynamics in the BamA beta-Barrel
Seam. *Biochemistry* **56**, 3142-3149, doi:10.1021/acs.biochem.7b00281
(2017).
14 Gessmann, D., Chung, Y. H., Danoff, E. J., Plummer, A. M., Sandlin, C. W.,
Zaccai, N. R. & Fleming, K. G. Outer membrane beta-barrel protein folding is
physically controlled by periplasmic lipid head groups and BamA.
*Proceedings of the National Academy of Sciences of the United States of*
*America* **111**, 5878-5883, doi:10.1073/pnas.1322473111 (2014).

15  Doyle, M. T. & Bernstein, H. D. Bacterial outer membrane proteins assemble
via asymmetric interactions with the BamA beta-barrel. *Nature*
*communications* **10**, 3358, doi:10.1038/s41467-019-11230-9 (2019).
16  Tomasek, D., Rawson, S., Lee, J., Wzorek, J. S., Harrison, S. C., Li, Z. &
Kahne, D. Structure of a nascent membrane protein as it folds on the BAM
complex. *Nature*, doi:10.1038/s41586-020-2370-1 (2020).
17  Sklar, J. G., Wu, T., Gronenberg, L. S., Malinverni, J. C., Kahne, D. &
Silhavy, T. J. Lipoprotein SmpA is a component of the YaeT complex that
assembles outer membrane proteins in Escherichia coli. *Proceedings of the*
*National Academy of Sciences of the United States of America* **104**, 6400-
6405, doi:10.1073/pnas.0701579104 (2007).
18  Kim, S., Malinverni, J. C., Sliz, P., Silhavy, T. J., Harrison, S. C. & Kahne, D.
Structure and function of an essential component of the outer membrane
protein assembly machine. *Science (New York, N.Y.)* **317**, 961-964,
doi:10.1126/science.1143993 (2007).
19  Wu, T., Malinverni, J., Ruiz, N., Kim, S., Silhavy, T. J. & Kahne, D.
Identification of a multicomponent complex required for outer membrane
biogenesis in Escherichia coli. *Cell* **121**, 235-245,
doi:10.1016/j.cell.2005.02.015 (2005).
20  Bennion, D., Charlson, E. S., Coon, E. & Misra, R. Dissection of beta-barrel
outer membrane protein assembly pathways through characterizing BamA
POTRA 1 mutants of Escherichia coli. *Molecular microbiology* **77**, 1153-
1171, doi:10.1111/j.1365-2958.2010.07280.x (2010).
21  Rizzitello, A. E., Harper, J. R. & Silhavy, T. J. Genetic evidence for parallel
pathways of chaperone activity in the periplasm of Escherichia coli. *Journal*
*of bacteriology* **183**, 6794-6800, doi:10.1128/jb.183.23.6794-6800.2001
(2001).
22  Crane, J. M. & Randall, L. L. The Sec System: Protein Export in Escherichia
coli. *EcoSal Plus* **7**, doi:10.1128/ecosalplus.ESP-0002-2017 (2017).
23  Walsh, N. P., Alba, B. M., Bose, B., Gross, C. A. & Sauer, R. T. OMP peptide
signals initiate the envelope-stress response by activating DegS protease via
relief of inhibition mediated by its PDZ domain. *Cell* **113**, 61-71,
doi:10.1016/s0092-8674(03)00203-4 (2003).
24  Ades, S. E. Regulation by destruction: design of the sigmaE envelope stress
response. *Current opinion in microbiology* **11**, 535-540,
doi:10.1016/j.mib.2008.10.004 (2008).
25  Rhodius, V. A., Suh, W. C., Nonaka, G., West, J. & Gross, C. A. Conserved
and variable functions of the sigmaE stress response in related genomes.
*PLoS biology* **4**, e2, doi:10.1371/journal.pbio.0040002 (2006).
26  Guillier, M., Gottesman, S. & Storz, G. Modulating the outer membrane with
small RNAs. *Genes & development* **20**, 2338-2348,
doi:10.1101/gad.1457506 (2006).
27  De Las Penas, A., Connolly, L. & Gross, C. A. SigmaE is an essential sigma
factor in Escherichia coli. *Journal of bacteriology* **179**, 6862-6864,
doi:10.1128/jb.179.21.6862-6864.1997 (1997).
28  Missiakas, D., Mayer, M. P., Lemaire, M., Georgopoulos, C. & Raina, S.
Modulation of the Escherichia coli sigmaE (RpoE) heat-shock transcription-
factor activity by the RseA, RseB and RseC proteins. *Molecular microbiology*
**24**, 355-371, doi:10.1046/j.1365-2958.1997.3601713.x (1997).

29  De Las Penas, A., Connolly, L. & Gross, C. A. The sigmaE-mediated response to extracytoplasmic stress in Escherichia coli is transduced by RseA and RseB, two negative regulators of sigmaE. *Molecular microbiology* **24**, 373-385, doi:10.1046/j.1365-2958.1997.3611718.x (1997).

30  Nicoloff, H., Gopalkrishnan, S. & Ades, S. E. Appropriate Regulation of the sigma(E)-Dependent Envelope Stress Response Is Necessary To Maintain Cell Envelope Integrity and Stationary-Phase Survival in Escherichia coli. *Journal of bacteriology* **199**, doi:10.1128/jb.00089-17 (2017).

31  Mitchell, A. M. & Silhavy, T. J. Envelope stress responses: balancing damage repair and toxicity. *Nature reviews. Microbiology* **17**, 417-428, doi:10.1038/s41579-019-0199-0 (2019).

32  Morris, F. C., Wells, T. J., Bryant, J. A., Schager, A. E., Sevastsyanovich, Y. R., Squire, D. J. P., Marshall, J., Isom, G. L., Rooke, J., Maderbocus, R., Knowles, T. J., Overduin, M., Rossiter, A. E., Cunningham, A. F. & Henderson, I. R. YraP Contributes to Cell Envelope Integrity and Virulence of Salmonella enterica Serovar Typhimurium. *Infection and immunity* **86**, doi:10.1128/iai.00829-17 (2018).

871  33  Tsang, M. J., Yakhnina, A. A. & Bernhardt, T. G. NlpD links cell wall remodeling and outer membrane invagination during cytokinesis in Escherichia coli. *PLoS genetics* **13**, e1006888, doi:10.1371/journal.pgen.1006888 (2017).

875  34  Onufryk, C., Crouch, M. L., Fang, F. C. & Gross, C. A. Characterization of six lipoproteins in the sigmaE regulon. *Journal of bacteriology* **187**, 4552-4561, doi:10.1128/jb.187.13.4552-4561.2005 (2005).

878  35  Seib, K. L., Haag, A. F., Oriente, F., Fantappie, L., Borghi, S., Semchenko, E. A., Schulz, B. L., Ferlicca, F., Taddei, A. R., Giuliani, M. M., Pizza, M. & Delany, I. The meningococcal vaccine antigen GNA2091 is an analogue of YraP and plays key roles in outer membrane stability and virulence. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology* **33**, 12324-12335, doi:10.1096/fj.201900669R (2019).

884  36  Bos, M. P., Grijpstra, J., Tommassen-van Boxtel, R. & Tommassen, J. Involvement of Neisseria meningitidis lipoprotein GNA2091 in the assembly of a subset of outer membrane proteins. *The Journal of biological chemistry* **289**, 15602-15610, doi:10.1074/jbc.M113.539510 (2014).

888  37  Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K. A., Tomita, M., Wanner, B. L. & Mori, H. Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. *Molecular systems biology* **2**, 2006.0008, doi:10.1038/msb4100050 (2006).

38  Cui, L., Vigouroux, A., Rousset, F., Varet, H., Khanna, V. & Bikard, D. A CRISPRi screen in E. coli reveals sequence-specific toxicity of dCas9. *Nature communications* **9**, 1912, doi:10.1038/s41467-018-04209-5 (2018).

39  Calvo-Villamanan, A., Ng, J. W., Planel, R., Menager, H., Chen, A., Cui, L. & Bikard, D. On-target activity predictions enable improved CRISPR-dCas9 screens in bacteria. *Nucleic acids research*, doi:10.1093/nar/gkaa294 (2020).

40  Pichoff, S., Du, S. & Lutkenhaus, J. Roles of FtsEX in cell division. *Research in microbiology* **170**, 374-380, doi:10.1016/j.resmic.2019.07.003 (2019).

41  Heidrich, C., Templin, M. F., Ursinus, A., Merdanovic, M., Berger, J., Schwarz, H., de Pedro, M. A. & Holtje, J. V. Involvement of N-acetylmuramyl-L-alanine amidases in cell separation and antibiotic-induced autolysis of

Escherichia coli. *Molecular microbiology* **41**, 167-178, doi:10.1046/j.1365-
2958.2001.02499.x (2001).

42   Uehara, T., Dinh, T. & Bernhardt, T. G. LytM-domain factors are required for
daughter cell separation and rapid ampicillin-induced lysis in Escherichia coli.
*Journal of bacteriology* **191**, 5094-5107, doi:10.1128/jb.00505-09 (2009).

43   Yang, D. C., Peters, N. T., Parzych, K. R., Uehara, T., Markovski, M. &
Bernhardt, T. G. An ATP-binding cassette transporter-like complex governs
cell-wall hydrolysis at the bacterial cytokinetic ring. *Proceedings of the*
*National Academy of Sciences of the United States of America* **108**, E1052-
1060, doi:10.1073/pnas.1107780108 (2011).

44   Uehara, T., Parzych, K. R., Dinh, T. & Bernhardt, T. G. Daughter cell
separation is controlled by cytokinetic ring-activated cell wall hydrolysis. *The*
*EMBO journal* **29**, 1412-1422, doi:10.1038/emboj.2010.36 (2010).

45   Chung, H. S., Yao, Z., Goehring, N. W., Kishony, R., Beckwith, J. & Kahne,
D. Rapid beta-lactam-induced lysis requires successful assembly of the cell
division machinery. *Proceedings of the National Academy of Sciences of the*
*United States of America* **106**, 21872-21877, doi:10.1073/pnas.0911674106
(2009).

46   Malinverni, J. C., Werner, J., Kim, S., Sklar, J. G., Kahne, D., Misra, R. &
Silhavy, T. J. YfiO stabilizes the YaeT complex and is essential for outer
membrane protein assembly in Escherichia coli. *Molecular microbiology* **61**,
151-164, doi:10.1111/j.1365-2958.2006.05211.x (2006).

47   Carlson, M. L., Stacey, R. G., Young, J. W., Wason, I. S., Zhao, Z., Rattray,
D. G., Scott, N., Kerr, C. H., Babu, M., Foster, L. J. & Duong Van Hoa, F.
Profiling the Escherichia coli membrane protein interactome captured in
Peptidisc libraries. *eLife* **8**, doi:10.7554/eLife.46615 (2019).

48   Alvira, S., Watkins, D. W., Troman, L., Allen, W. J., Lorriman, J.,
Degliesposti, G., Cohen, E. J., Beeby, M., Daum, B., Gold, V. A. M., Skehel,
J. M. & Collinson, I. Inter-membrane association of the Sec and BAM
translocons for bacterial outer-membrane biogenesis. *bioRxiv*, 589077,
doi:10.1101/589077 (2020).

49   Rouviere, P. E. & Gross, C. A. SurA, a periplasmic protein with peptidyl-prolyl
isomerase activity, participates in the assembly of outer membrane porins.
*Genes & development* **10**, 3170-3182, doi:10.1101/gad.10.24.3170 (1996).

50   McBroom, A. J. & Kuehn, M. J. Release of outer membrane vesicles by
Gram-negative bacteria is a novel envelope stress response. *Molecular*
*microbiology* **63**, 545-558, doi:10.1111/j.1365-2958.2006.05522.x (2007).

51   Roman-Hernandez, G., Peterson, J. H. & Bernstein, H. D. Reconstitution of
bacterial autotransporter assembly using purified components. *eLife* **3**,
e04234, doi:10.7554/eLife.04234 (2014).

52   Hale, C. A. & de Boer, P. A. Direct binding of FtsZ to ZipA, an essential
component of the septal ring structure that mediates cell division in E. coli.
*Cell* **88**, 175-185, doi:10.1016/s0092-8674(00)81838-3 (1997).

53   Hartmann, J. B., Zahn, M., Burmann, I. M., Bibow, S. & Hiller, S. Sequence-
Specific Solution NMR Assignments of the beta-Barrel Insertase BamA to
Monitor Its Conformational Ensemble at the Atomic Level. *Journal of the*
*American Chemical Society* **140**, 11252-11260, doi:10.1021/jacs.8b03220
(2018).

54   Vertommen, D., Ruiz, N., Leverrier, P., Silhavy, T. J. & Collet, J. F.
Characterization of the role of the Escherichia coli periplasmic chaperone

SurA using differential proteomics. *Proteomics* **9**, 2432-2443,
doi:10.1002/pmic.200800794 (2009).

55  Konovalova, A., Schwalm, J. A. & Silhavy, T. J. A Suppressor Mutation That
Creates a Faster and More Robust sigmaE Envelope Stress Response.
*Journal of bacteriology* **198**, 2345-2351, doi:10.1128/jb.00340-16 (2016).

56  Kern, B., Leiser, O. P. & Misra, R. Suppressor Mutations in degS Overcome
the Acute Temperature-Sensitive Phenotype of DeltadegP and DeltadegP
Deltatol-pal Mutants of Escherichia coli. *Journal of bacteriology* **201**,
doi:10.1128/jb.00742-18 (2019).

57  Hart, E. M., O'Connell, A., Tang, K., Wzorek, J. S., Grabowicz, M., Kahne, D.
& Silhavy, T. J. Fine-Tuning of sigma(E) Activation Suppresses Multiple
Assembly-Defective Mutations in Escherichia coli. *Journal of bacteriology*
**201**, doi:10.1128/jb.00745-18 (2019).

58  Leiser, O. P., Charlson, E. S., Gerken, H. & Misra, R. Reversal of the ΔdegP
phenotypes by a novel rpoE allele of Escherichia coli. *PloS one* **7**, e33979,
doi:10.1371/journal.pone.0033979 (2012).

59  Yeats, C. & Bateman, A. The BON domain: a putative membrane-binding
domain. *Trends in biochemical sciences* **28**, 352-355, doi:10.1016/s0968-
0004(03)00115-4 (2003).

60  Bryant, J. A., Morris, F. C., Knowles, T. J., Maderbocus, R., Heinz, E.,
Boelter, G., Alodaini, D., Colyer, A., Wotherspoon, P. J., Staunton, K. A.,
Jeeves, M., Browning, D. F., Sevastsyanovich, Y. R., Wells, T. J., Rossiter,
A. E., Bavro, V. N., Sridhar, P., Ward, D. G., Chong, Z. S., Icke, C., Teo, A.,
Chng, S. S., Roper, D. I., Lithgow, T., Cunningham, A. F., Banzhaf, M.,
Overduin, M. & Henderson, I. R. Structure-function analyses of dual-BON
domain protein DolP identifies phospholipid binding as a new mechanism for
protein localisation. *bioRxiv*, 2020.2008.2010.244616,
doi:10.1101/2020.08.10.244616 (2020).

61  Fleming, K. G. A combined kinetic push and thermodynamic pull as driving
forces for outer membrane protein sorting and folding in bacteria.
*Philosophical transactions of the Royal Society of London. Series B,
Biological sciences* **370**, doi:10.1098/rstb.2015.0026 (2015).

62  Horne, J. E., Brockwell, D. J. & Radford, S. E. Role of the lipid bilayer in
outer membrane protein folding in Gram-negative bacteria. *The Journal of
biological chemistry* **295**, 10340-10367, doi:10.1074/jbc.REV120.011473
(2020).

63  Sinnige, T., Weingarth, M., Renault, M., Baker, L., Tommassen, J. & Baldus,
M. Solid-state NMR studies of full-length BamA in lipid bilayers suggest
limited overall POTRA mobility. *Journal of molecular biology* **426**, 2009-2021,
doi:10.1016/j.jmb.2014.02.007 (2014).

64  Rassam, P., Copeland, N. A., Birkholz, O., Tóth, C., Chavent, M., Duncan, A.
L., Cross, S. J., Housden, N. G., Kaminska, R., Seger, U., Quinn, D. M.,
Garrod, T. J., Sansom, M. S., Piehler, J., Baumann, C. G. & Kleanthous, C.
Supramolecular assemblies underpin turnover of outer membrane proteins in
bacteria. *Nature* **523**, 333-336, doi:10.1038/nature14461 (2015).

65  Gunasinghe, S. D., Shiota, T., Stubenrauch, C. J., Schulze, K. E., Webb, C.
T., Fulcher, A. J., Dunstan, R. A., Hay, I. D., Naderer, T., Whelan, D. R., Bell,
T. D. M., Elgass, K. D., Strugnell, R. A. & Lithgow, T. The WD40 Protein
BamB Mediates Coupling of BAM Complexes into Assembly Precincts in the

Bacterial Outer Membrane. *Cell reports* **23**, 2782-2794, doi:10.1016/j.celrep.2018.04.093 (2018).

Jarosławski, S., Duquesne, K., Sturgis, J. N. & Scheuring, S. High-resolution architecture of the outer membrane of the Gram-negative bacteria Roseobacter denitrificans. *Molecular microbiology* **74**, 1211-1222, doi:10.1111/j.1365-2958.2009.06926.x (2009).

Lessen, H. J., Fleming, P. J., Fleming, K. G. & Sodt, A. J. Building Blocks of the Outer Membrane: Calculating a General Elastic Energy Model for β-Barrel Membrane Proteins. *Journal of chemical theory and computation* **14**, 4487-4497, doi:10.1021/acs.jctc.8b00377 (2018).

Choi, U. & Lee, C. R. Distinct Roles of Outer Membrane Porins in Antibiotic Resistance and Membrane Integrity in Escherichia coli. *Frontiers in microbiology* **10**, 953, doi:10.3389/fmicb.2019.00953 (2019).

Delhaye, A., Collet, J. F. & Laloux, G. A Fly on the Wall: How Stress Response Systems Can Sense and Respond to Damage to Peptidoglycan. *Frontiers in cellular and infection microbiology* **9**, 380, doi:10.3389/fcimb.2019.00380 (2019).

Macritchie, D. M. & Raivio, T. L. Envelope Stress Responses. *EcoSal Plus* **3**, doi:10.1128/ecosalplus.5.4.7 (2009).

Grabowicz, M. & Silhavy, T. J. Envelope Stress Responses: An Interconnected Safety Net. *Trends in biochemical sciences* **42**, 232-242, doi:10.1016/j.tibs.2016.10.002 (2017).

Grenier, F., Matteau, D., Baby, V. & Rodrigue, S. Complete Genome Sequence of Escherichia coli BW25113. *Genome announcements* **2**, doi:10.1128/genomeA.01038-14 (2014).

Blattner, F. R., Plunkett, G., 3rd, Bloch, C. A., Perna, N. T., Burland, V., Riley, M., Collado-Vides, J., Glasner, J. D., Rode, C. K., Mayhew, G. F., Gregor, J., Davis, N. W., Kirkpatrick, H. A., Goeden, M. A., Rose, D. J., Mau, B. & Shao, Y. The complete genome sequence of Escherichia coli K-12. *Science (New York, N.Y.)* **277**, 1453-1462, doi:10.1126/science.277.5331.1453 (1997).

Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 6640-6645, doi:10.1073/pnas.120163297 (2000).

Knop, M., Siegers, K., Pereira, G., Zachariae, W., Winsor, B., Nasmyth, K. & Schiebel, E. Epitope tagging of yeast genes using a PCR-based strategy: more tags and improved practical routines. *Yeast (Chichester, England)* **15**, 963-972, doi:10.1002/(sici)1097-0061(199907)15:10b<963::Aid-yea399>3.0.Co;2-w (1999).

McLeay, R. C. & Bailey, T. L. Motif Enrichment Analysis: a unified framework and an evaluation on ChIP data. *BMC bioinformatics* **11**, 165, doi:10.1186/1471-2105-11-165 (2010).

Ieva, R. & Bernstein, H. D. Interaction of an autotransporter passenger domain with BamA during its translocation across the bacterial outer membrane. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 19120-19125, doi:10.1073/pnas.0907912106 (2009).

**Supplementary Materials**

**Outer membrane lipoprotein DolP interacts with the BAM complex and**

**promotes fitness during envelope stress response**

David Ranava, Yiying Yang, Luis Orenday-Tapia, François Rousset, Catherine

Turlan, Lun Cui, Violette Morales, Cyril Moulin, Gladys Munoz, Jérôme Rech, Anne

Caumont-Sarcos, Cécile Albenne, David Bikard, Raffaele Ieva

# Figure S1.

12
**Fig. S1. The deletion of *dolP* severely impairs growth in the presence of**
**vancomycin.**

(**A**) BW25113 and Δ*dolP* cells were cultured in LB. The cell densities of both cultures

were monitored by measuring the $OD_{600}$ at regular time intervals. The graph reports

mean values of independent cultures ± standard deviation (SD, *n* = 3).

(**B**) BW25113 and Δ*dolP* cells carrying the indicated plasmids were serially diluted

and spotted on LB containing 60 μg/ml vancomycin where indicated. Ectopic

expression was driven by the leaky transcriptional activity of $P_{trc}$ in the absence of

IPTG*.

22
23

**Figure S2.**

**Fig. S2. Reproducibility of the CRISPRi screens and ontology analysis of gene hits.**

(**A**) The raw read counts of experimental replicates are very well correlated (Pearson's r > 0.97) in both $dolP^+$ (left) and $\Delta dolP$ (right) screens.

(**B**) Log2FC values of each sgRNA are compared between the WT and $\Delta dolP$ screens. Only guides targeting *envC*, *bamD*, and *rseA* are highlighted in orange, red and cyan, respectively.

(**C**) Gene Ontology analysis of the genes with a significant synthetic interaction with *dolP*. Using FDR<0.05 as a threshold, 27 genes were selected for Gene Ontology (GO) analysis for cellular component (upper panel) or biological process (bottom panel)[78]. Both analyses show an overrepresentation of genes associated with cell envelope and membrane function.

**Figure S3.**

**Fig. S3. DolP is not crucial for OMP biogenesis.**

(**A**) The protein content of the envelope fraction of BW25113 (*dolP*[+]) or $\Delta dolP$ cells was mixed with SDS-PAGE loading buffer and analyzed by SDS-PAGE and coomassie staining.

(**B** and **C**) The protein content of the envelope fraction of BW25113 (*dolP*[+]) and $\Delta dolP$ cells was mixed with SDS-PAGE loading buffer, heated at the indicated temperatures, separated by SDS-PAGE and immunoblotted using anti-LamB (B) or anti-OmpC (C) antisera. LamB and OmpC form complexes (trimers), which are resistant to SDS and require significant heating to be denatured. At progressively higher temperatures, LamB (B) and OmpC (C) complexes are denatured and the two proteins migrate as monomers. The amounts of LamB and OmpC complexes or denatured monomers and the temperature-dependent patterns of complex-to-monomer conversion do not appear to depend on *dolP* expression.

(**D**) The biogenesis of EspP, an autotransporter OMP of enterohemorrhagic *E. coli* heterologously expressed in BW25113 cells, was monitored in time. The plasmid pRI22[79] harbouring the EspP encoding gene under the control of an IPTG-inducible promoter was transformed into wild-type BW25113 (*dolP*[+]) and $\Delta dolP$ cells. EspP is a serine protease autotransporter of *Enterobacteriaceae*, and consists of an N-terminal passenger domain secreted on the cell surface and a C-terminal domain which is ultimately folded into an integral OM β-barrel structure[80]. EspP biogenesis requires both periplasmic chaperones, such as SurA, and the BAM complex[79]. Once EspP assembly and secretion are completed, the passenger domain is cleaved by an autocatalytic reaction that is catalyzed by residues of the folded β-barrel domain facing the interior of the barrel. Thus, the proteolytic reaction can be exploited to infer about completion of β-barrel folding and assembly[80,81]. To monitor the kinetics of EspP autoproteolysis, cells carrying pRI22 were cultured in M9 minimal medium until early exponential phase, supplemented with 200 μM IPTG and subjected to [35]S

pulse-chase labelling. The total protein contents of samples collected after the

indicated chase times were subjected to immunoprecipitation using an antiserum

specific for an EspP C-terminal peptide. The asterisk indicates a degradation product

of EspP, likely generated by the OM protease OmpT, which cleaves the N-terminal

region of the passenger domain as it is secreted across the OM[82]. A band

corresponding to the EspP β-barrel domain is generated with a similar time

dependence in both *dolP*[+] and Δ*dolP* cells.

77
78

**Figure S4.**

79
80

 **Fig. S4. Overproduced BAM accumulates in the OM**

(**A**) The crude envelope fractions of cells overproducing BAM$^{ProtA}$, BAM$^{\Delta P1/ProtA}$, or

BAM$^{\Delta P2/ProtA}$ were solubilized with 1% (w/v) digitonin and 0.1% (w/v) DDM and

subjected to native IgG-affinity chromatography. Elutions were analyzed by blue

native-PAGE and coomassie staining (lanes 1-3) or SDS-PAGE and immunoblotting

(lanes 4-6). Blue native-PAGE of the elution fraction containing wild-type BamA

resolved a coomassie stainable complex migrating with an apparent mass of 250

kDa, which corresponds to the BAM complex (lane 1). A roughly similar amount of

BAM complex was detected in the elution obtained with BamA$^{\Delta P2/ProtA}$ samples,

although in this case the BAM$^{\Delta P2}$ complex migrated slightly faster, accounting for the

BamA$^{\Delta P2}$ mass difference (lane 3). In contrast, the amount of BAM$^{\Delta P1/ProtA}$ variant was

considerably lower (lane 2). The asterisk indicates that BamE is obtained from TEV

digestion of BamE$^{ProtA}$.

(**B**) Cells overproducing the indicated variants of the BAM complex were fractionated.

The membrane and soluble fractions, obtained upon treatment of collected cells with

lysozyme and EDTA were analyzed by SDS-PAGE and immunoblotting. BamA$^{\Delta P1}$

was depleted from the total membrane fraction (lanes 1-3), and accumulated in the

soluble fraction containing the periplasmic protein DsbA (lanes 4-6).

Contrast          BamD<sup>mCherry</sup>

Control

ΔdolP

pDolP<sup>His</sup>

**Figure S5.**

**Fig. S5. Effect of lack or overproduction of DolP on BAM localization.**

BW25113 derivative cells harbouring a *bamD-mCherry* chromosomal fusion were

visualized on 1% (w/v) agarose pads by fluorescence and phase-contrast

microscopy. Top: *dolP*⁺ cells (control); Center: Δ*dolP* cells: Bottom: *dolP*⁺ cells

carrying pDolP$^{His}$. Cells were cultured at 30°C in minimal M9 medium to OD$_{600}$ = 0.5.

IPTG (400 μM) was supplemented for 1 hour to induce ectopic expression of DolP$^{His}$.

Cells were visualized on 1% (w/v) agarose pads by fluorescence and phase-contrast

microscopy. Bar = 5 μm.

**A**

**B**

**C**

# Figure S6.

**Fig. S6. DolP accumulates at mid-cell sites during a late step of cell division.**

(**A**) The total protein content of cells harbouring the *dolP-gfp* chromosomal fusion

(lanes 1) or wild-type *dolP* (lanes 2) were analyzed by SDS page followed by

coomassie staining or immunoblotting using a DolP specific antiserum.

(**B,C**) BW25113-derivative cells harbouring the chromosomal fusion *dolP-gfp* and

either *zipA-mCherry* (B) or *nlpD-mCherry* (C) were cultured at 30°C in minimal M9

medium to $OD_{600}$ = 0.2-0.3 and visualized on 1% (w/v) agarose pads by fluorescence

and phase-contrast microscopy. Bar = 5 $\mu$m.

120
121

**Figure S7.**

**Fig. S7. Overproduction of BAM does not influence septal recruitment of NlpD and ZipA.**

(**A**) BW25113 cells harbouring pBAM$^{His}$ cultured and supplemented with 400 $\mu$M IPTG for 1 hour prior to collecting cells. The protein content of the envelope fraction (lane 1) was analysed by SDS-PAGE and coomassie staining and compared to the protein content of the envelope fraction of BW25113 cells (lane 2).

(**B**) BW25113-derivative cells harbouring the chromosomal fusion *dolP-gfp* and *zipA-mCherry* or *nlpD-mCherry* were transformed with pBAM$^{His}$ (ectopic overproduction of all BAM subunits, with His-tagged BamE). Cells were then cultured at 30°C in minimal M9 medium supplemented with 400 $\mu$M IPTG for 1 hour until OD$_{600}$ = 0.2-0.3, and visualized on 1% (w/v) agarose pads by fluorescence microscopy. The arrowheads indicate the localization of DolP$^{GFP}$, ZipA$^{mCherry}$ or NlpD$^{mCherry}$ at division septa.

(**C**) Top: BW25113-derivative *dolP-gfp* cells transformed with pOmpA (ectopic overproduction of His-tagged OmpA) were culture in LB medium and supplemented with 400 mm IPTG or no IPTG for 1 hour prior to collecting cells. The total protein content was analyzed by SDS-PAGE and immunoblotting using the indicated antisera. Bottom: BW25113-derivative *dolP-gfp* cells transformed with pOmpA$^{His}$ were cultured in minimal M9 medium supplemented with 400 $\mu$M IPTG for 1 hour until OD$_{600}$ = 0.2-0.3, and visualized (B). Bar = 5 $\mu$m.

(**D**) BW25113-derivative cells harbouring the chromosomal fusion *dolP-gfp* and *zipA-mCherry* or *nlpD-mCherry* were transformed with pBamA$^{His}$ (ectopic overproduction of a partially inactive His-tagged form of BamA). Cells were then cultured at 30°C in minimal M9 medium supplemented with 400 $\mu$M IPTG for 1 hour until OD$_{600}$ = 0.2-0.3, and visualized as in (B).

**A**

**B**

pBamA$^{His}$  pBamA$^{\Delta P1/His}$  pBamA$^{\Delta P2/His}$

**C**

**D**

**E**

Figure S8.

**Fig. S8. BamA overaccumulation in the OM and envelope stress impair septal DolP localization.**

(**A**) The JCM166 BamA depletion strain or its transformants carrying plasmids encoding BamA$^{His}$ or wild-type BamA and DolP were cultured in the presence of 0.02% (w/v) arabinose (which induces transcription of a chromosomal copy of wild-type *bamA* engineered downstream of an arabinose-inducible P$_{bad}$ promoter). Serial dilutions were spotted on LB agar devoid of or supplemented with 0.02% (w/v) arabinose, as indicated. In the absence of arabinose, growth is supported by the non-induced, leaky P$_{trc}$ transcription of the wild-type BamA-encoding gene, and not of its variant encoding C-terminally His-tagged BamA.

(**B**) Overnight cultures of BW25113 cells harboring the chromosomal *dolP-gfp* fusion and carrying the indicated plasmids were diluted in minimal M9 medium supplemented with 400 $\mu$M IPTG. The cells were grown at 30°C to OD$_{600}$ = 0.2-0.3 and visualized on 1% (w/v) agarose pads by fluorescence microscopy. The arrows indicate envelope constriction sites between forming daughter cells. Bar = 5 $\mu$m.

(**C**) Counting of BW25113 cells presenting septal DolP$^{GFP}$ fluorescent signals. *dolP-gfp* cells carring pBAM$^{His}$, pBamA$^{His}$, pBamA$^{\Delta P1/His}$ or pBamA$^{\Delta P2/His}$ (analyzed by fluorescence microscopy and shown in Fig. 4A and in Fig. S8B) were counted. The percentage of cells presenting fluorescent DolP$^{GFP}$ signals at the mid-cell in samples overproducing all five BAM subunits or only one of the indicated BamA variants was normalized to the same fraction obtained for cells carrying the control empty vector (pCtrl). Bar charts display a mean value ± SD (*n* = 3). More than 300 cells were counted in each experiment.

(**D**) BW25113 cells carrying the indicated plasmids were cultured and supplemented with 400 $\mu$M IPTG to induce ectopic expression of BamA and its mutant forms. Collected cells were subjected to lysozyme/EDTA lysis to obtain the total membrane and soluble fractions. The protein contents of the indicated cell fractions were separated by SDS-PAGE and immunodecorated with the indicated antisera.

(**E**) Overnight cultures of ∆*bamB* cells harboring the chromosomal *dolP-gfp* fusion

were diluted in minimal M9 medium, cultured at 30°C to $OD_{600}$ = 0.3 and visualized

on 1% (w/v) agarose pads by fluorescence microscopy. The arrows indicate

envelope constriction sites between forming daughter cells. Bar = 5 μm. The

percentage of cells presenting fluorescent $DolP^{GFP}$ signals at division septa in ∆*bamB*

cells was normalized to the same fraction obtained for the control cells (BW25113

carrying the *dolP-gfp* chromosomal fusion) and reported below the micrograph. More

than 300 cells were counted.

**Table S1. Log2FC values of sgRNAs in the WT or ΔdolP screen.**

See source data file (Table S1)

**Table S2. Genetic interaction scores.**

See source data file (Table S2)

**Table S3: List of strains**

| Name | Genotype and relevant features | Source |
|---|---|---|
| BW25113 | Δ(araD-araB)567 Δ(rhaD-rhaB)568 ΔlacZ4787(::rrnB-3) hsdR514 rph-1 (wild-type reference) | Grenier et al., 2014 [83] |
| ΔenvC | BW25113 envC::kan | This study |
| ΔrseA | BW25113 rseA::kan | This study |
| ΔdolP | BW25113 dolP::kan | This study |
| ΔbamB | BW25113 bamB::kan | This study |
| ΔbamB ΔdolP | BW25113 ΔbamB dolP::kan | |
| ΔdolP ΔrseA | BW25113 ΔdolP rseA::kan | This study |
| dolP-gfp | BW25113 dolP-gfp | This study |
| ΔsurA dolP-gfp | BW25113 surA::kan dolP-gfp | This study |
| ΔbamB dolP-gfp | BW25113 bamB::kan dolP-gfp | This study |
| ΔdolP bamD-mCherry | BW25113 ΔdolP bamD-mCherry | This study |
| bamD-mCherry | BW25113 bamD-mCherry | This study |
| dolP-gfp nlpD-mCherry | BW25113 dolP-gfp nlpD-mCherry | This study |
| dolP-gfp zipA-mCherry | BW25113 dolP-gfp zipA-mCherry | This study |
| JCM166 | MC4100 ara^{r/-} Δ(λatt-lom)::bla P$_{BAD}$yaeT araC ΔyaeT (BamA depletion strain) | Wu et al., 2005 [84] |
| LC-E75 | F⁻ λ⁻ ilvG⁻ rfb-50 rph-1 186attB::P$_{tet}$-dcas9, λattB::mCherry | Cui et al., 2018 [85] |
| LC-E75 ΔdolP | LC-E75 dolP::kan | This study |

 **Table S4: List of plasmids**

| Name | Relevant features | Source |
|---|---|---|
| pCtrl | Reference empty vector for ectopic protein expression | This study |
| pBAM | Ectopic expression of all 5 BAM subunits | This study |
| pBAM$^{His}$ (pJH114) | Ectopic expression of all 5 BAM subunits; BamE is C-terminally His-tagged | Roman Hernandez *et al*., 2014 [86] |
| pBAM$^{ProtA}$ | Ectopic expression of all 5 BAM subunits; BamE is C-terminally Protein A-tagged; a TEV site amino acid linker is positioned immediately upstream of the tag | This study |
| pBAM$^{\Delta P1/ProtA}$ | Ectopic expression of all 5 BAM subunits; BamA harbours the deletion of POTRA1 and BamE is C-terminally Protein A-tagged; a TEV site amino acid linker is positioned immediately upstream of the tag | This study |
| pBAM$^{\Delta P2/ProtA}$ | Ectopic expression of all 5 BAM subunits; BamA harbours the deletion of POTRA2 and BamE is C-terminally Protein A-tagged; a TEV site amino acid linker is positioned immediately upstream of the tag | This study |
| pBamBCDE$^{His}$ | Ectopic expression of BamB, BamC, BamD and BamE subunits; BamE is C-terminally His-tagged | This study |
| pBamA$^{His}$ | Ectopic expression of C-terminally His-tagged BamA | This study |
| pBamA$^{\Delta P1/His}$ | Ectopic expression of a C-terminally His-tagged BamA variant lacking POTRA1 | This study |
| pBamA$^{\Delta P2/His}$ | Ectopic expression of a C-terminally His- | This study |

| | | |
|---|---|---|
| | tagged BamA variant lacking POTRA2 | |
| pBamA-DolP$^{His}$ | Ectopic expression of BamA and C-terminally His-tagged DolP | This study |
| pBamA$^{\Delta P1}$-DolP$^{His}$ | Ectopic expression of a BamA variant lacking POTRA1 and C-terminally His-tagged DolP | This study |
| pBamA$^{\Delta P2}$-DolP$^{His}$ | Ectopic expression of a BamA variant lacking POTRA2 and C-terminally His-tagged DolP | This study |
| pDolP | Ectopic expression of wild-type DolP | This study |
| pDolP$^{ProtA}$ | Ectopic expression of C-terminally protein A-tagged DolP; a TEV site amino acid linker is positioned immediately upstream of the tag | This study |
| pDolP$^{His}$ | Ectopic expression of C-terminally His-tagged DolP | This study |
| pOmpA$^{His}$ | Ectopic expression of C-terminally His-tagged OmpA | This study |
| pRI22 | Ectopic expression of EspP | Ieva *et al.*, 2009 [79] |
| psgRNAcos | Reference empty vector for expression of sgRNAs | Cui *et al.*, 2018 [85] |
| psgRNA*bamD* (pCAT187) | Expression of a *bamD*-specific sgRNA | This study |
| psgRNA*bamE* (pCAT189) | Expression of a *bamE*-specific sgRNA | This study |
| psgRNA*envC* (pCAT191) | Expression of a *envC*-specific sgRNA | This study |
| psgRNA*ftsX* (pCAT193) | Expression of a *ftsX*-specific sgRNA | This study |

**Supplementary References**

Ashburner, M., Ball, C. A., Blake, J. A., Botstein, D., Butler, H., Cherry, J. M., Davis, A. P., Dolinski, K., Dwight, S. S., Eppig, J. T., Harris, M. A., Hill, D. P., Issel-Tarver, L., Kasarskis, A., Lewis, S., Matese, J. C., Richardson, J. E., Ringwald, M., Rubin, G. M. & Sherlock, G. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nature genetics* **25**, 25-29, doi:10.1038/75556 (2000).

Ieva, R. & Bernstein, H. D. Interaction of an autotransporter passenger domain with BamA during its translocation across the bacterial outer membrane. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 19120-19125, doi:10.1073/pnas.0907912106 (2009).

Albenne, C. & Ieva, R. Job contenders: roles of the beta-barrel assembly machinery and the translocation and assembly module in autotransporter secretion. *Molecular microbiology* **106**, 505-517, doi:10.1111/mmi.13832 (2017).

Dautin, N., Barnard, T. J., Anderson, D. E. & Bernstein, H. D. Cleavage of a bacterial autotransporter by an evolutionarily convergent autocatalytic mechanism. *The EMBO journal* **26**, 1942-1952, doi:10.1038/sj.emboj.7601638 (2007).

Szabady, R. L., Peterson, J. H., Skillman, K. M. & Bernstein, H. D. An unusual signal peptide facilitates late steps in the biogenesis of a bacterial autotransporter. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 221-226, doi:10.1073/pnas.0406055102 (2005).

Grenier, F., Matteau, D., Baby, V. & Rodrigue, S. Complete Genome Sequence of Escherichia coli BW25113. *Genome announcements* **2**, doi:10.1128/genomeA.01038-14 (2014).

Wu, T., Malinverni, J., Ruiz, N., Kim, S., Silhavy, T. J. & Kahne, D. Identification of a multicomponent complex required for outer membrane biogenesis in Escherichia coli. *Cell* **121**, 235-245, doi:10.1016/j.cell.2005.02.015 (2005).

Cui, L., Vigouroux, A., Rousset, F., Varet, H., Khanna, V. & Bikard, D. A CRISPRi screen in E. coli reveals sequence-specific toxicity of dCas9. *Nature communications* **9**, 1912, doi:10.1038/s41467-018-04209-5 (2018).

Roman-Hernandez, G., Peterson, J. H. & Bernstein, H. D. Reconstitution of bacterial autotransporter assembly using purified components. *eLife* **3**, e04234, doi:10.7554/eLife.04234 (2014).