Title

Extraintestinal pathogenic *Escherichia coli* (ExPEC) are associated with prolonged carriage of

extended-spectrum β-lactamase-producing *E. coli* acquired during travel

Author list

Boas C.L. van der Putten[1,2]*, Jarne M. van Hattem[1], John Penders[3,4], COMBAT Consortium[#],

Daniel R. Mende[1], Constance Schultsz[1,2]

[1]Department of Medical Microbiology, Amsterdam UMC, University of Amsterdam, the

Netherlands

[2]Department of Global Health, Amsterdam Institute for Global Health and Development,

Amsterdam UMC, University of Amsterdam, the Netherlands

[3]School for Public Health and Primary Care (Caphri), Department of Medical Microbiology,

Maastricht University Medical Centre, Maastricht, Netherlands

[4]School for Nutrition and Translational Research in Metabolism (NUTRIM), Maastricht University

Medical Centre, Maastricht, Netherlands

*Corresponding author. Email: boas.vanderputten@amsterdamumc.nl

Telephone: +31205664868.

[#]Members are listed in the appendixes

Article type: Original article

Abstract word count: 246

Main text word count: 2521

Abstract

Objectives Extended-spectrum β-lactamase-producing *Escherichia coli* (ESBL-Ec) are frequently

acquired during international travel, contributing to the global spread of antimicrobial

resistance. Human-adapted ESBL-Ec are predicted to exhibit increased intestinal carriage

duration, resulting in a higher likelihood of onward human-to-human transmission. Yet,

bacterial determinants of increased carriage duration are unknown. Previous studies analysed

small traveler cohorts, with short follow-up times, or did not employ high-resolution molecular

typing, and were thus unable to identify bacterial traits associated with long-term carriage.

Methods In a prospective cohort study of 2001 international travelers, we analysed 160 faecal

ESBL-Ec isolates from all 38 travelers who acquired ESBL-Ec during travel and subsequently

carried ESBL-Ec for at least 12 months after return, by whole-genome sequencing. For 17

travelers, we confirmed the persistence of ESBL-Ec strains through single nucleotide variant

typing. To identify determinants of increased carriage duration, we compared the 17 long-term

carriers (≥12 months carriage) with 33 age-, sex- and destination-matched short-term carriers

(<1 month carriage). Long-read sequencing was employed to investigate ESBL plasmid

persistence.

Results We show that in healthy travelers with very low antibiotic usage, extraintestinal

pathogenic lineages of *E. coli* (ExPEC) are significantly more likely to persist than other *E. coli*

lineages. The long-term carriage of *E. coli* from ExPEC lineages is mainly driven by sequence type

131 and phylogroup D *E. coli*.

Conclusions Although ExPEC frequently cause extra-intestinal infections such as bloodstream

infections, our results imply that ExPEC are also efficient intestinal colonizers, which potentially

contributes to their onward transmission.

Introduction

International travel contributes significantly to the spread of extended-spectrum β-lactamase

(ESBL) gene positive *Escherichia coli* (ESBL-Ec).[1,2] Genetically diverse ESBL-Ec are frequently

acquired during international travel.[3–5] Whilst travel-acquired ESBL-Ec typically are lost during

travel or within the first month after return (Figure 1A),[6] ESBL-Ec and ESBL genes have been

detected for more than 12 months after return in a proportion of travelers.[1,3–5,7] ESBL genes can

persist through at least two different mechanisms.[8] Bacterial strains that carry ESBL-encoding

genes in their chromosome or on a stable plasmid can persist over time as part of the local

microbiome (strain persistence, Figure 1B). In addition, ESBL genes can be located on mobile

genetic elements (MGEs) which persist in the microbiome by their ability to transfer between

different bacterial hosts (MGE persistence, Figure 1C).

ESBL-Ec lineages which are capable of long-term colonization and are adapted to the human

intestinal tract are likely to contribute to onward transmission of ESBL-Ec and ESBL genes.

However, it is unknown which ESBL-Ec lineages are capable of persistence after return from

travel, due to a lack of studies with a sufficiently large sample size and a prospective

longitudinal study design. Additionally, high-resolution typing methods such as whole-genome

sequencing (WGS) are needed to reliably determine whether the strains have been carried over

a long period. One study employed WGS to investigate persistence in 16 travelers who acquired

ESBL-Ec abroad and showed that only one traveler carried a travel-acquired ESBL-Ec strain for at

least 7 months.[7] A very recent study analysed data from 11 travellers which were colonised for

>3 months.[9] Due to the low sample sizes in both studies, few ESBL-Ec attributes could be

identified that were significantly associated with long-term carriage. Armand-Lefèvre et al.

reported an association between phylogroups B2/D/F with prolonged carriage duration (p =

0.02).[9] Other studies either did not employ WGS or focused on short-term carriage only.

We studied a cohort of 2001 Dutch international travelers (COMBAT cohort).[1] Six hundred and

thirty-three travelers acquired ESBL gene-positive Enterobacterales during travel abroad, of

whom 38 travelers (6.0%) were colonized for ≥12 months after acquisition of ESBL gene-positive

Enterobacterales, all of which were *E. coli*. Here we report on host and bacterial characteristics

associated with ESBL genes persistence and the mechanism through which the ESBL genes

persisted. We identified an association between extraintestinal pathogenic *E. coli* and long-term

carriage.

Materials and methods
77

Travelers and isolates selection

We included all 38 travelers from the COMBAT cohort (N=2001) who were colonized for ≥12

months with *Escherichia coli* which possessed ESBL genes belonging to a single ESBL gene group

at return from travel and at all subsequent time points (1, 3, 6, and 12 months after return), and

who tested ESBL-negative before travel (see reference 1 for a detailed description of sampling

and microbiological methods).[1] In short, faecal samples were inoculated in tryptic soy broth

containing 50 mg/L vancomycin to select for Enterobacterales. After overnight incubation, the

broth was subcultured on chromID ESBL agar plates (bioMérieux). Morphologically different

colonies were isolated, with a maximum of five isolates per faecal sample. Eighty-five ESBL-Ec

isolates sampled at return from travel (T0) and 75 ESBL-Ec isolates sampled 12 months after

return from travel (T12) were included in the current study (Figure 2).

***In silico* typing**

DNA extraction and library preparation were performed on all available isolates using the

Qiagen Blood and Tissue DNA extraction kit (Cat No./ID: 69506) and Kapa HTP library prep kit

(Kit Code KK8234), respectively. Whole-genome sequencing was performed on Illumina HiSeq

2500 by the Amsterdam UMC Core Facility Genomics.

Sequencing data were analysed using a Snakemake v5.7.1[10] pipeline, available at https://

github.com/boasvdp/COMBAT (v1.1.0 archived at https://doi.org/10.5281/zenodo.4582689). In

short, Illumina sequencing data were trimmed using fastp v0.20.0,[11] assembled using the Shovill

wrapper v1.0.9 (https://github.com/tseemann/shovill) for SPAdes,[12] and resistance genes were

identified using AMRfinderplus v3.2.3.[13] *E. coli* phylogroups were predicted using EzClermont

v0.4.3.[14] Multi-locus sequence typing (MLST) was performed with the mlst script

(https://github.com/tseemann/mlst), using the Achtman scheme for *Escherichia coli*.[15]

Analysis of strain persistence

We defined strain persistence as the presence of one or more ESBL-Ec isolates in faecal samples

from a single traveler obtained at return from travel (T0), as well as 12 months thereafter (T12),

with a maximum of 25 SNPs per 5 Mbp difference in the core genome.[16]

To assess strain persistence, we calculated single nucleotide polymorphism differences between

isolates to identify which isolates belonged to the same strain. We mapped sequencing reads

using Snippy v4.4.5 (https://github.com/tseemann/snippy) on MLST-specific reference genomes

selected with ReferenceSeeker v1.6.3[17] to obtain core genome alignments. Reference genomes

used for all MLST sequence types, together with core genome alignment lengths, can be found

in table S1. For each MLST, IQtree v1.6.12[18] was used to infer a phylogeny from the core

genome alignment under the model advised by Modelfinder.[19] Recombination events in the

core genome alignment were identified using ClonalFrameML v1.12[20] and masked using maskrc-

svg v0.5 (https://github.com/kwongj/maskrc-svg). SNP differences were counted using snp-dists

v0.7.0 (https://github.com/tseemann/snp-dists) and alignment lengths were calculated using a

modified version of snp-dists v0.7.0 (https://github.com/boasvdp/snp-dists). SNP counts were

scaled to 5 Mbp, to approximate the number of genome-wide SNPs.[16]

To determine whether ESBL plasmids had persisted independent of bacterial host, we employed

long read sequencing (Oxford Nanopore Technologies). We generated Oxford Nanopore

Technologies sequencing data according to Van der Putten et al. (2020).[21] In short, strains were

grown overnight at 37 ˚C in liquid LB. DNA extraction and library preparation were performed

using the Qiagen MagAttract HMW DNA Kit (Cat. No. 67563) and ONT native barcoding kit (Cat.

No. EXP-NBD114), respectively. The library was subsequently sequenced on an ONT MinION

flowcell. Raw read data was filtered using Filtlong v0.2.0 (https://github.com/rrwick/Filtlong)

and assembled with corresponding Illumina data using Unicycler v0.4.8.[22] Quality control was

implemented at several steps in the pipeline using FastQC v0.11.8,[23] Quast v4.6.3,[24] and

MultiQC v1.6.[25] Plasmid comparison was performed using ANICalculator.[26]

Comparison with short-term carriers

Seventeen long-term carriers were matched by age (range +/- 7 years), sex, and travel

destination (United Nations subregions) to thirty-three travelers who were colonized for less

than a month using SPSS 26 (Figure 2). Illumina WGS was performed as described before on all

ESBL-Ec isolated at return from travel from these short-term carriers (total 41 isolates).

Plotting and statistical analysis

Data were plotted using ggplot2 v3.1.1,[27] ggthemes v2.4.0,[28] and patchwork[29] in R v3.5.1.

Tabular data were analysed using Pandas v0.24.2[30] in Python v3.6.7. Statistical analysis was

performed using Fisher's exact test as implemented in the Python library SciPy.[31]

Results
Data characteristics

Out of 2001 Dutch international travelers, we included 34 travelers with samples available, out

of all 38 travelers whose faecal samples were positive for *E. coli* harbouring the same ESBL

group gene at return (T0), one month after return from travel (T1) and 12 months after return

from travel (T12), but were ESBL-negative before travel (Figure 2).[1] We included a median of 1

ESBL-Ec isolate per traveler per timepoint (range: 1-5 isolates). SNP typing showed that acquired

ESBL-Ec isolates were genetically diverse, with some travelers acquiring up to four distinct

strains.

ESBL-Ec strain persistence

SNP analysis

Twenty-two out of 34 travelers harboured persistent strains based on SNP typing (Figure 2). Of

the 12 travelers who did not harbour persistent strains, ten carried strains with differing MLST

profiles between T0 and T12, and two carried strains from a single ST but with the number of

core genome-wide SNP differences exceeding the threshold (25 SNPs for a 5 Mbp genome).[16]

Clonal ST38 lineages

Non-persistent, yet highly conserved strains that are widely disseminated within the human

population could potentially be misidentified as persistent when using SNV analyses. Hence, we

compared SNV distances between isolates obtained from unrelated travelers (i.e., belonging to

different households) to identify strains shared across our study population. Conserved strains

belonging to two ST38 lineages were identified in five unrelated travelers. Lineage ST38-$bla_{CTX-M-27}$

$_{27}$ was identified in three unrelated travelers and lineage ST38-$bla_{CTX-M-14}$ in two other, unrelated

travelers. All five unrelated travelers returned from different countries and continents,

suggesting these two lineages have disseminated widely. Whole genomes from lineages ST38-

$bla_{CTX-M-27}$ and ST38-$bla_{CTX-M-14}$ were also abundantly present in public data and a phylogenetic

analysis of their core genomes confirmed their high similarity (Figure S1). Hence, we could not

conclusively determine whether these highly clonal strains had persisted in the five travelers

concerned, or whether these strains were re-acquired from an unknown source. Therefore, we

excluded the travelers harbouring these clonal ST38 strains from further analyses to identify

lineages potentially associated with persistence. ST38 strains colonizing four additional travelers

were not related to the ST38-$bla_{CTX-M-27}$ and ST38-$bla_{CTX-M-14}$ lineages and were thus not excluded

from further analyses.

We identified two related travelers who harboured highly similar ST69 isolates (9 genome-wide

SNPs). The most likely explanation of this observation is that the travellers acquired this strain

from the same source or one from the other, and that it has persisted in both travellers since.

Plasmid analysis

For six travelers, we detected identical ESBL genes from isolates sampled at return from travel

and twelve months thereafter, but we could exclude ESBL gene persistence in persistent isolates

based on SNP typing (Table S1). To determine whether the plasmid carrying these ESBL genes

had persisted independent of bacterial host (Figure 1B), we employed long read sequencing. For

one traveler, we detected a pair of ESBL-Ec isolates belonging to different MLST types in which

an almost identical plasmid (99.8% nucleotide identity) was identified at T0 and T12, suggesting

ESBL gene persistence by plasmid transfer between different *E. coli* hosts.

In summary, we identified persistent ESBL-Ec strains in 17 out of 34 travelers with prolonged

ESBL-Ec carriage and persistent ESBL-plasmid in one traveler (Table S1). Although we included

multiple ESBL-Ec isolates for some travelers, we did not observe multiple persistent strains

within any single traveler.

Comparison of ESBL-Ec from long-term and short-term carriers

For each long-term carrier (≥12 months carriage) harbouring a persistent strain, two short-term

carriers (<1 month carriage) were matched by age, sex and travel destination. For one long-term

carrier, only a single matching short-term carrier could be identified, resulting in a comparison

of 17 isolates from 17 long-term carriers and 42 isolates from 33 matched short-term carriers,

which were sequenced.

Antibiotic usage was low before, during, and after travel and similar between long-term and

short-term carriers (Table 1). None of the travelers were admitted to the hospital during or after

travel in either group. One single traveler returned to the same country as visited during index

travel, within 12 months after return from index travel.

Following the recent definition of common extraintestinal pathogenic *E. coli* (ExPEC) lineages

according to Manges *et al.* (2019),[32] persistent strains belonged significantly more often to

ExPEC lineages than non-persistent strains. In 15 out of 17 long-term carriers, ExPEC were

identified as the persistent strain while in only 7 out of 33 short-term carriers ExPEC were

detected (odds ratio 27.86, 95% confidence interval: 5.11-151.74). This difference appears to be

driven mostly by ST131 and phylogroup D strains.

Discussion

We demonstrated long-term carriage of travel-acquired ESBL-positive *Escherichia coli* in 17

travelers out of a cohort of 2001 travelers, which was driven by persistence of ESBL-Ec

belonging to ExPEC lineages. In a single traveler, the persistence of ESBL-Ec seemed to be due to

an ESBL plasmid which shifted between bacterial hosts after colonising the traveler's gut.

Our study strengthens the finding of Armand-Lefèvre et al.[9] that ESBL-Ec lineage is associated

with persistent carriage after acquisition during travel. Interestingly, we come to similar

conclusions although we have focused on long-term carriage (≥12 months) as compared to the

previous study (>3 month carriage). Earlier work could not make strong assertions about the

persistence of strains, either due to the limited number of included travelers, because the

typing methods employed were insufficiently discriminating between isolates, because of a

limited duration of follow-up, or because the study population was not representative of

community exposure.[3–5,7] In a recent study in Laos, a group of travelers attending a course at

local hospitals acquired a very high diversity of ESBL-Ec *E. coli* immediately upon arrival and it

was shown that this acquisition of resistant *E. coli* during travel is highly dynamic.[6] However, the

short follow-up in this study did not allow analysis of long-term outcomes of the acquisition of

ESBL-Ec.

Previous studies have shown that a limited number of *E. coli* lineages contribute to a large

fraction of *E. coli*-mediated extraintestinal disease, including urinary tract infections and

bloodstream infections.[32] These extraintestinal pathogenic *E. coli* are commonly referred to as

ExPEC. While ExPEC display pathogenic potential, epidemiological studies often find ExPEC

colonizing the human gut in the absence of symptoms.[33,34] In fact, it has been proposed that

ExPEC have evolved towards particularly efficient intestinal colonization and their

extraintestinal virulence is an evolutionary "byproduct".[35] The long-term persistence of ExPEC

lineages, acquired after relatively short travel duration as observed in our study (median 20

days, interquartile range: 17 – 29 days) suggests that ExPEC lineages have spread globally

successfully due to their adaptation to the human intestinal tract. It should however be noted

that we restricted our analysis to ESBL-producing *E. coli* and duration of persistence of

susceptible ExPEC in this cohort is unknown.

Whole-genome sequencing allowed us to detect two clonal ST38 lineages which were shared

between unrelated travelers. By additional analysis of publicly available WGS data, these ST38

lineages were shown to have spread globally. The extremely high degree of similarity within

these ST38 lineages interfered with reliable strain identification based on core genome DNA

sequence analysis. Future studies should explore the application of SNP typing using the

pangenome rather than only the core genome, for example through software like Pandora[36].

These novel methods however currently display a higher error rate than the core genome

analysis approach used in the present study[36]. Awareness of the circulation of ST38 or other

rapidly expanding lineages that are indistinguishable by their core genome is important, for

example for management of suspected outbreaks of ST38 in hospital settings.

Our current study focused solely on ESBL-producing *E. coli* and found an association between

ExPEC carriage and increased carriage duration. Our approach could also be applied to non-

ESBL-producing ExPEC, to determine whether acquisition of ExPEC in general, independent of

their antibiotic susceptibility profile, is more likely to results in increased carriage duration after

travel. The frequency of sampling can be considered a limitation of this study. However, when

comparing between two timepoints, twelve months apart, we could still identify very closely

related isolates in samples from the same traveler, indicating strain persistence. Additionally,

we only had isolates available for household members of four long-term carriers. Including

more household members in future studies might allow us to estimate the likelihood of onward

transmission following long-term carriage.

Applying genomic epidemiology to a large traveler cohort, we have shown that ESBL-positive *E.*

coli acquired during travel are able to persist for more than a year. The strains that showed the

longest carriage duration belonged predominantly to pathogenic ExPEC lineages. Our data imply

that long-term carriage of resistant *E. coli* is governed by bacterial characteristics which are

associated with lineage. This finding possibly allows a more precise risk assessment for

international travelers returning with travel-acquired resistant *E. coli*.

Acknowledgements

The authors would like to thank Arie van der Ende for the helpful discussions. We thank

SURFsara (https://www.surfsara.nl) for the support in using the Lisa Compute Cluster.

Funding information

The COMBAT study was funded by Netherlands Organization for Health, Research and

Development (ZonMw; 50-51700-98-120) and EU-H2020 programme (COMPARE, 643476). BP

was funded through an internal grant of the Amsterdam UMC ("flexibele OiO beurs").

Declaration of interests

We declare no competing interests.

Data and code availability

All Illumina and Oxford Nanopore Technologies sequencing data used in this study are currently

available free of restrictions at NCBI under project accession number PRJEB40103. Metadata

linking isolates to travelers, required to reproduce our analyses, are currently available free of

restrictions at the GitHub repository of this project

(https://www.github.com/boasvdp/COMBAT, v1.1.0 archived through Zenodo at

https://doi.org/10.5281/zenodo.4582689).

All code is available free of restrictions under the MIT license at

https://www.github.com/boasvdp/COMBAT (v1.1.0 archived through Zenodo at

https://doi.org/10.5281/zenodo.4582689).

Ethical statement

The COMBAT study was approved by the Medical Research Ethics Committee, Maastricht

University Medical Centre (METC 12-4-093). All participants provided written informed consent.

Appendices

281   Figure S1: Phylogeny of clonal complex 38 genomes with continent of origin.

282   Table S1: All SNP comparisons between isolates with relevant metadata (169 rows).

283   Supplemental information: Members of the COMBAT consortium.

References

1.   Arcilla MS, van Hattem JM, Haverkate MR, et al. Import and spread of extended-spectrum β-
lactamase-producing Enterobacteriaceae by international travellers (COMBAT study): a
prospective, multicentre cohort study. *Lancet Infect Dis*. 2017;17(1):78-85. doi:10.1016/S1473-
3099(16)30319-X

2.   Ruppé E, Andremont A, Armand-Lefèvre L. Digestive tract colonization by multidrug-resistant
Enterobacteriaceae in travellers: An update. *Travel Med Infect Dis*. 2018;21:28-35.
doi:10.1016/j.tmaid.2017.11.007

3.   Ruppé E, Armand-Lefèvre L, Estellat C, et al. High Rate of Acquisition but Short Duration of Carriage
of Multidrug-Resistant Enterobacteriaceae After Travel to the Tropics. *Clin Infect Dis Off Publ Infect
Dis Soc Am*. 2015;61(4):593-600. doi:10.1093/cid/civ333

4.   Paltansing S, Vlot JA, Kraakman MEM, et al. Extended-spectrum β-lactamase-producing
enterobacteriaceae among travelers from the Netherlands. *Emerg Infect Dis*. 2013;19(8):1206-
1213. doi:10.3201/eid.1908.130257

5.   Pires J, Kuenzli E, Kasraian S, et al. Polyclonal Intestinal Colonization with Extended-Spectrum
Cephalosporin-Resistant Enterobacteriaceae upon Traveling to India. *Front Microbiol*. 2016;7:1069.
doi:10.3389/fmicb.2016.01069

6.   Kantele A, Kuenzli E, Dunn SJ, et al. Dynamics of intestinal multidrug-resistant bacteria colonisation
contracted by visitors to a high-endemic setting: a prospective, daily, real-time sampling study.
Lancet Microbe. 2021;0(0). doi:10.1016/S2666-5247(20)30224-X

7.   Bevan ER, McNally A, Thomas CM, Piddock LJV, Hawkey PM. Acquisition and Loss of CTX-M-
Producing and Non-Producing Escherichia coli in the Fecal Microbiome of Travelers to South Asia.
mBio. 2018;9(6). doi:10.1128/mBio.02408-18

8.   Sheppard AE, Stoesser N, Wilson DJ, et al. Nested Russian Doll-Like Genetic Mobility Drives Rapid
Dissemination of the Carbapenem Resistance Gene blaKPC. *Antimicrob Agents Chemother*.
2016;60(6):3767-3778. doi:10.1128/AAC.00464-16

9.   Armand-Lefèvre L, Rondinaud E, Desvillechabrol D, et al. Dynamics of extended-spectrum beta-
lactamase-producing Enterobacterales colonization in long-term carriers following travel abroad.
Microb Genomics. 2021;7(7). doi:10.1099/mgen.0.000576

10.   Köster J, Rahmann S. Snakemake—a scalable bioinformatics workflow engine. *Bioinformatics*.
2012;28(19):2520-2522. doi:10.1093/bioinformatics/bts480

11.   Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*.
2018;34(17):i884-i890. doi:10.1093/bioinformatics/bty560

12.   Bankevich A, Nurk S, Antipov D, et al. SPAdes: A New Genome Assembly Algorithm and Its
Applications to Single-Cell Sequencing. *J Comput Biol*. 2012;19(5):455-477.
doi:10.1089/cmb.2012.0021

13.   Feldgarden M, Brover V, Haft DH, et al. Validating the AMRFinder Tool and Resistance Gene
Database by Using Antimicrobial Resistance Genotype-Phenotype Correlations in a Collection of
Isolates. *Antimicrob Agents Chemother*. 2019;63(11). doi:10.1128/AAC.00483-19

14.   Waters NR, Abram F, Brennan F, Holmes A, Pritchard L. Easy phylotyping of Escherichia coli via the
EzClermont web app and command-line tool. *Access Microbiol*. Published online 2020.
doi:10.1099/acmi.0.000143

15.   Wirth T, Falush D, Lan R, et al. Sex and virulence in Escherichia coli: an evolutionary perspective.
Mol Microbiol. 2006;60(5):1136-1151. doi:10.1111/j.1365-2958.2006.05172.x

16.   Gorrie CL, Silva AGD, Ingle DJ, et al. Systematic analysis of key parameters for genomics-based real-
time detection and tracking of multidrug-resistant bacteria. *bioRxiv*. Published online September
25, 2020:2020.09.24.310821. doi:10.1101/2020.09.24.310821

17.   Schwengers O, Hain T, Chakraborty T, Goesmann A. ReferenceSeeker: rapid determination of
333   appropriate reference genomes. *J Open Source Softw*. 2020;5(46):1994. doi:10.21105/joss.01994

334   18.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective Stochastic
335   Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol*. 2015;32(1):268-274.
336   doi:10.1093/molbev/msu300

337   19.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: fast model
338   selection for accurate phylogenetic estimates. *Nat Methods*. 2017;14(6):587-589.
339   doi:10.1038/nmeth.4285

340   20.   Didelot X, Wilson DJ. ClonalFrameML: Efficient Inference of Recombination in Whole Bacterial
341   Genomes. *PLOS Comput Biol*. 2015;11(2):e1004041. doi:10.1371/journal.pcbi.1004041

342   21.   van der Putten BCL van der, Roodsant TJ, Haagmans MA, Schultsz C, Ark KCH van der. Five
343   Complete Genome Sequences Spanning the Dutch Streptococcus suis Serotype 2 and Serotype 9
344   Populations. *Microbiol Resour Announc*. 2020;9(6). doi:10.1128/MRA.01439-19

345   22.   Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome assemblies from short
346   and long sequencing reads. *PLOS Comput Biol*. 2017;13(6):e1005595.
347   doi:10.1371/journal.pcbi.1005595

348   23.   Andrews S, others. *FastQC: A Quality Control Tool for High Throughput Sequence Data*. Babraham
349   Bioinformatics, Babraham Institute, Cambridge, United Kingdom; 2010.

350   24.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome assemblies.
351   *Bioinformatics*. 2013;29(8):1072-1075. doi:10.1093/bioinformatics/btt086

352   25.   Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for multiple tools
353   and samples in a single report. *Bioinformatics*. 2016;32(19):3047-3048.
354   doi:10.1093/bioinformatics/btw354

355   26.   Varghese NJ, Mukherjee S, Ivanova N, et al. Microbial species delineation using whole genome
356   sequences. *Nucleic Acids Res*. 2015;43(14):6761-6771. doi:10.1093/nar/gkv657

357   27.   Wickham H. *Ggplot2: Elegant Graphics for Data Analysis*. Springer; 2016.

358   28.   Arnold JB. ggthemes: Extra Themes, Scales and Geoms for "ggplot2." *R Package Version*. 2017;3(0).

359   29.   Pedersen TL. patchwork: The Composer of ggplots. *R Package Version 00*. 2017;1.

360   30.   team T pandas development. *Pandas-Dev/Pandas: Pandas*. Zenodo; 2020.
361   doi:10.5281/zenodo.3509134

362   31.   Jones E, Oliphant T, Peterson P, others. *SciPy: Open Source Scientific Tools for Python*.; 2001.
363   http://www.scipy.org/

364   32.   Manges AR, Geum HM, Guo A, Edens TJ, Fibke CD, Pitout JDD. Global Extraintestinal Pathogenic
365   Escherichia coli (ExPEC) Lineages. *Clin Microbiol Rev*. 2019;32(3). doi:10.1128/CMR.00135-18

33. Denamur E, Clermont O, Bonacorsi S, Gordon D. The population genetics of pathogenic Escherichia
coli. *Nat Rev Microbiol*. 2021;19(1):37-54. doi:10.1038/s41579-020-0416-x

34. Verschuuren TD, van Hout D, Arredondo-Alonso S, et al. Comparative genomics of ESBL-producing
Escherichia coli (ESBL-Ec) reveals a similar distribution of the 10 most prevalent ESBL-Ec clones and
ESBL genes among human community faecal and extra-intestinal infection isolates in the
Netherlands (2014–17). *J Antimicrob Chemother*. 2021;76(4):901-908. doi:10.1093/jac/dkaa534

35. Le Gall T, Clermont O, Gouriou S, et al. Extraintestinal Virulence Is a Coincidental By-Product of
Commensalism in B2 Phylogenetic Group Escherichia coli Strains. *Mol Biol Evol*. 2007;24(11):2373-
2384. doi:10.1093/molbev/msm172

36. Colquhoun RM, Hall MB, Lima L, et al. Nucleotide-resolution bacterial pan-genomics with reference
graphs. *bioRxiv*. Published online November 24, 2020:2020.11.12.380378.
doi:10.1101/2020.11.12.380378

Table 1. Characteristics of long-term and short-term carriers. Matching was performed on sex,

age and travel destination (United Nations subregions). These characteristics are depicted in

italic. Data are presented as number (%) for all characteristics except age and travel duration,

which is presented as median (IQR).

		Long-term carriers (n = 17)	Short-term carriers (n = 33)
Male		4 (23.5%)	8 (24.2%)
Age (years)		50.2 (41.7-59.4)	51.4 (36.5-58.1)
Continent visited during index travel	*Asia*	16 (94.1%)	31 (93.9%)
	North and South America	1 (5.9%)	2 (6.1%)
Travel duration (days)		20 (17-28)	19 (14-21)
Admitted to hospital during index travel		0 (0%)	0 (0%)
Travel to the same country as index travel, within 12 months after return from index travel		1 (5.9%)	0 (0%)
Antibiotic usage	Within 3 months before index travel	0 (0%)	0 (0%)
	During index travel	3 (17.6%)	2 (6.1%)
	Within 1 month after return from index travel	0 (0%)	0 (0%)
	Within 1 to 3 months after return from index travel	1 (5.9%)	2 (6.7%)
	Within 3 to 6 months after return from index travel	0 (0%)	2 (7.1%)
	Within 6 to 12 months after return from index travel	3 (17.6%)	2 (7.4%)
Traveler's diarrhoea		6 (35.3%)	9 (27.3%)

Figure 1. Schematic overview of main persistence mechanism of ESBL genes in the gut of

returning travelers. Gray arrows indicate carriage duration over time. A) No persistence, where

the colonizing strain is lost during the follow-up period. B) Strain persistence, where a bacterial

strain harboring an ESBL gene is continuously present. C) Persistence through mobile genetic

elements, where the original strain harboring the ESBL gene has been lost, but the ESBL gene

(located on a mobile genetic element) has been passed on to another strain.

Figure 2. Study flowchart. Strain persistence was defined as isolates from a single traveler,

sample twelve months apart with identical ESBL alleles, identical MLST profiles and 25 or fewer

genome-wide SNPs difference. Clonal ST38 isolates were excluded. Seventeen long-term

carriers (colonized ≥12 months with a single strain) were matched on sex, age and travel

destination to thirty-three short-term carriers (colonized <1 month).

Figure S1. Core genome phylogeny of 1805 clonal complex 38 *E. coli* strains. Outer ring indicates

continent on which the strain was isolated. The two clonal ST38 lineages present in the COMBAT

collection are marked in red and blue. Tree and metadata available through iTOL:

https://itol.embl.de/tree/2131278347268071589210657.