A phospho-regulated ensemble signal motif of α-TAT1 drives dynamic microtubule acetylation

Abhijit Deb Roy[1*], Evan G. Gross[2], Gayatri S. Pillai[2], Shailaja Seetharaman[3,4], Sandrine Etienne-Manneville[3], Takanari Inoue[1*]

[1] Department of Cell Biology and Center for Cell Dynamics, Johns Hopkins University School of Medicine, 855 North Wolfe Street, Baltimore, MD 21205, USA

[2] The Johns Hopkins University, Baltimore, MD 21218, USA

[3]Cell Polarity, Migration and Cancer Unit, Institut Pasteur, UMR3691 CNRS, Equipe Labellisée Ligue Contre le Cancer, F-75015, Paris, France.

[4]Université Paris Descartes, Sorbonne Paris Cité, 12 Rue de l'École de Médecine, 75006 Paris, France.

[*] Correspondence:
Abhijit Deb Roy abhijit.debroy@gmail.com
Takanari Inoue jctinoue@jhmi.edu

Abstract:

Spatiotemporal patterns of microtubule modifications such as acetylation underlie diverse cel-

lular functions. While the molecular identity of the acetylating agent, α-tubulin N-

acetyltransferase 1 (α-TAT1), as well as the functional consequences of microtubule acetyla-

tion have been revealed, the molecular mechanisms that regulate multi-tasking α-TAT1 action

for dynamic acetylation remain obscure. Here we identified a signal motif in the intrinsically

disordered C-terminus of α-TAT1, which comprises three functional elements - nuclear export,

nuclear import and cytosolic retention. Their balance is tuned via phosphorylation by serine-

threonine kinases to determine subcellular localization of α-TAT1. While the phosphorylated

form binds to 14-3-3 adapters and accumulates in the cytosol for maximal substrate access,

the non-phosphorylated form is sequestered inside the nucleus, thus keeping microtubule

acetylation minimal. As cancer mutations have been reported to this motif, the unique ensem-

ble regulation of α-TAT1 localization may hint at a role of microtubule acetylation in aberrant

physiological conditions.

Introduction:

Acetylation of Lysine-40 of α-tubulin is an evolutionarily conserved post-translational modification observed across eukaryotic species[1–3], which is involved in diverse physiological and pathological states[4–6]. Acetylation is mainly observed in polymerized microtubules[7–9] and may provide structural flexibility to stabilize microtubules against bending forces[10–13]. In cultured cells, microtubule acetylation mediates focal adhesion dynamics, adaptation to extracellular matrix rigidity as well as regulation of tissue stiffness[14–17]. Additionally, acetylated microtubules regulate touch sensation in *M. musculus*, *C. elegans* and *D. melanogaster,* suggesting a role in mechano-response[18–21]. Microtubule acetylation has been implicated in axonal transport in neurons[22–24], migration in cancer cells[5,25–27], autophagy[6,28,29], podosome stabilization in osteoclasts[30] and viral infections[31–34]. α-TAT1 is the only known acetyltransferase for α-tubulin[35,36] in mammals. α-TAT1 predominantly catalyzes α-tubulin in stable polymerized microtubules[19,37], and may have additional effects on microtubules independent of its catalytic activity[38].

Although microtubule acetylation is spatially and temporally regulated downstream of many molecular signaling pathways, little is known about how these pathways converge on α-TAT1 to achieve such dynamic patterns. In the present study, we used computational sequence analyses and live cell microscopy to identify a conserved motif in the intrinsically disordered C-terminus of α-TAT1, consisting of an NES and an NLS, that mediates its spatial distribution. We show that cytosolic localization of α-TAT1 is critical for microtubule acetylation. We further demonstrate that nuclear localization of α-TAT1 is inhibited by the action of serine-threonine kinases, specifically cyclin dependent kinases (CDKs), Protein Kinase A (PKA) and Casein kinase 2 (CK2) and identify 14-3-3 proteins as binding partners of α-TAT1. Our findings establish a novel role of the intrinsically disordered C-terminus in

62   controlling α-TAT1 function by regulating its intracellular localization downstream of kinase

63   and phosphatase activities.

 Results:

 α-TAT1 localization mediates microtubule acetylation: α-TAT1 has an N-terminal catalytic

domain that shows homology to other acetyltransferases, while its C-terminus was not

resolved in crystal structures[39] (Fig. 1a). Based on its amino acid sequence, α-TAT1 C-

terminus was predicted to be intrinsically disordered by both IUPred2A[40] and PrDOS[41]

prediction servers (Fig. 1a, Supplementary Fig. S1a, b, c). To explore if the intra-cellular

localization of α-TAT1 is regulated, we sought to identify any localization signals present in the

α-TAT1 amino acid sequence (Supplementary Fig. S1a). The prediction program NetNES[42]

identified a putative NES in α-TAT1 C-terminus (Fig. 1a, Supplementary Fig. S2). On the other

hand, PSORT-II[43] subcellular localization program predicted that α-TAT1 should be

predominantly localized to the nucleus due to the presence of a putative class 4 NLS[44] in its

C-terminus (Fig. 1a). The region encompassing the putative NES and NLS is conserved

across the human α-TAT1 isoforms (Supplementary Fig. S3a), as well as across mammalian

α-TAT1 proteins (Supplementary Fig. S3b). To test whether α-TAT1 indeed showed any

intracellular distribution pattern, we expressed mVenus-α-TAT1 in HeLa cells, and observed

distinct nuclear exclusion in most cells, although a subset of cells showed lack of exclusion

(Fig. 1b). Based on our categorization of the distribution patterns (see Methods, Fig. 1c), we

determined that approximately 77% of cells showed cytosolic distribution, 22% showed

diffused pattern (i.e., both cytosolic and nuclear distribution) and 1% of cells showed nuclear

enrichment of mVenus-α-TAT1 (Fig. 1d). These observations were consistent with ratiometric

analysis (see Methods, Fig 1c) of mVenus-α-TAT1 distribution (Fig. 1e). Time lapse

microscopy showed temporal changes in fluorescence intensity of mVenus-α-TAT1 in cell

nuclei and cytosol (Supplementary Fig. S4a, b), suggesting that α-TAT1 localization is

dynamic by nature.

α-TAT1 preferentially acetylates polymerized microtubules, which are typically

cytosolic. Based on this, we hypothesized that spatial regulation of α-TAT1 may control its

function. Exogenous expression of mVenus-α-TAT1 or its catalytic domain (residues 1-

236)[19,39] (Fig. 1f) was sufficient to significantly increase α-tubulin acetylation in HeLa cells

compared to non-transfected cells (Fig. 1g, h). To test whether nuclear localization may

sufficiently sequester α-TAT1 from microtubules, we tethered the NLS from cMyc to α-TAT1

catalytic domain and thus localized it to the nucleus (Fig. 1f, g top panel). Exogenous

expression of NLS-mVenus-α-TAT1(1-236) did not increase α-tubulin acetylation levels

compared to non-transfected cells (Fig. 1 g, h), suggesting that nuclear sequestration of α-

TAT1 inhibits its function.

α-TAT1 undergoes Exportin 1 dependent nuclear export: To dissect the molecular

mechanisms of α-TAT1 localization in the cytosol, we speculated that α-TAT1 is actively

exported out of the nucleus, and/or that α-TAT1 binds to a protein that keeps the complex out

of the nucleus. We began by testing the first possibility by assessing involvement of the

nuclear export machinery. Exportin 1 (Exp1), also called Chromosome region maintenance 1

protein homolog (CRM1), mediates nuclear export of many proteins[45,46]. GFP-α-TAT1, but not

GFP, co-immunoprecipitated with endogenous Exp1 (Fig. 2a), indicating an interaction

between these two proteins. Treatment with 100 nM Leptomycin-B (LMB), an inhibitor of Exp1

mediated nuclear export[47], significantly decreased the number of cells displaying nuclear

exclusion of mVenus-α-TAT1 compared to vehicle (Fig 2b, c, d). Inhibition of nuclear export

was initiated within an hour of LMB treatment, although some cells were refractory to the

treatment (Supplementary Fig. S5a, b). Decreased LMB concentrations (1 nM and 10 nM)

had a comparable impact as 100 nM dosage (Supplementary Fig. S5c, d). Furthermore, LMB

treatment induced significant reduction in α-tubulin acetylation levels in HeLa cells within 4

hours, compared to vehicle (Fig. 2e, f). Our data suggest that α-TAT1 is actively exported

from the nucleus to the cytosol in an Exp1 dependent manner and that this export facilitates

α-TAT1 function. The residual exclusion of α-TAT1 could be due to Exp1-independent nuclear

export pathways or association with other cytosolic proteins, which was tested later (see

below).

Nuclear export of α-TAT1 is mediated by a C-terminal NES: Our data demonstrate that α-

TAT1 function is linked to Exp1 mediated nuclear export. To examine whether nuclear export

of α-TAT1 is regulated by its catalytic activity, we expressed a catalytic dead mutant, mVenus-

α-TAT1(D157N)[39], in HeLa cells. mVenus-α-TAT1(D157N) did not display any loss of nuclear

exclusion (Supplementary Fig. S6a, b, c). The catalytic domain, mVenus-α-TAT1(1-236),

displayed a complete loss of nuclear exclusion; whereas the C-terminus, mVenus-α-

TAT1(236-323), displayed a distribution pattern comparable to that of WT (Fig. 3a, b, c, d). In

addition, inhibition of Exp1 by 100 nM LMB significantly reduced the nuclear exclusion of

mVenus-α-TAT1 C-terminus (Fig. 3e). Exclusion of α-TAT1 C-terminus (size $\approx$ 38 kDa) further

indicates that nuclear exclusion of mVenus-α-TAT1 (size $\approx$ 63 kDa) is not simply due to size

exclusion of passive diffusion into nuclei, and demonstrates that nuclear exclusion of α-TAT1

is a transferable property mediated by its C-terminus.

Exp1 dependent nuclear export is typically mediated by binding with short stretches of

hydrophobic, often Leucine rich, NES[42,45] that are often found in disordered regions of the

cargo proteins. As previously mentioned, NetNES suggested the presence of a conserved

NES between V286 and L297 in α-TAT1 C-terminus (Supplementary Fig S2). Interestingly,

this region is also predicted to be a site of protein-protein interactions by ANCHOR2

prediction software[40] (Supplementary Fig. S1b). Truncation of this putative NES, α-TAT1-

delNES, abrogated its nuclear exclusion (Fig. 3a, b, c, d). Although alanine substitution of the

hydrophobic residues in the V286-L297 region, α-TAT1(VL/A), decreased its nuclear

exclusion, the loss of exclusion was less than expected from our observations with LMB

treatment (Supplementary Fig. S6a, b, c). On further examination of the NetNES prediction,

we observed that the NES predicted by the Hidden Markov Model (Supplementary Fig. S2)

included the residues between L282 and L297. Additional Alanine substitutions of L282 and

F285, α-TAT1(NES/A) (Supplementary Fig. S6d), further reduced nuclear exclusion of α-TAT1

(Fig. 3a, b, c, d, Supplementary Fig. S6a, b, c), suggesting that these residues contribute to

the nuclear export of α-TAT1. Taken together, our data suggest that α-TAT1 has a

hydrophobic NES in its C-terminus.

α-TAT1 interacts with 14-3-3 proteins. Our observations thus far with nuclear exclusion of

mVenus-α-TAT1 indicate that the putative NLS identified by the PSORT prediction server is

either non-functional or is basally inhibited with occasional activation. 14-3-3 protein binding

has been reported to negatively regulate nuclear import by inhibiting binding of importins to

NLS[48–50]. Part of the putative NLS sequence "PAQRRRTR" bears similarity to 14-3-3 binding

motif RXX(pS/pT)XP[51], and 14-3-3-Pred[52], a 14-3-3 interaction prediction server identified

T322 as a potential 14-3-3 binding site. To examine whether α-TAT1 bound to 14-3-3 proteins,

we co-expressed GFP-α-TAT1 with HA-tagged 14-3-3-β or 14-3-3-ζ in HEK-293T cells and

performed a co-immunoprecipitation assay. GFP-α-TAT1, but not GFP alone, co-precipitated

with HA-14-3-3-β and HA-14-3-3-ζ (Fig. 4a). These observations are consistent with our

observations in mass spectrometry analysis of α-TAT1 in HEK cells that identified 14-3-3-β

and 14-3-3-ζ as potential interactors[53].

Serine/threonine kinase activities mediate cytosolic localization of α-TAT1. 14-3-3

binding to proteins is mediated by phosphorylated serine and threonine residues[51]. NetPhos

prediction server identified over 30 putative phosphosites in α-TAT1, of which nine residues

have been reported to be phosphorylated in phospho-proteomic studies (Supplementary Fig.

S7a, b). Treatment with 100 nM Staurosporin, a pan-kinase inhibitor, significantly increased

nuclear localization of α-TAT1 (Fig. 4b, c, Supplementary Fig. S8), suggesting that

phosphorylation negatively regulated nuclear localization of α-TAT1. To identify the specific

kinases that regulate α-TAT1 localization, we treated cells expressing mVenus-α-TAT1 with

several serine-threonine kinase inhibitors (summarized in Supplementary Table T1). Both RO-

3306 (inhibitor for CDK1 and CDK2[54,55]) and PD0332991 (also called Pablociclib, inhibitor for

CDK4 and CDK6[56,57]), but not Purvalanol-B (inhibits CDK1, CDK2 and CDK5[55]), increased

nuclear localization of mVenus-α-TAT1 in HeLa cells (Fig. 4b, c, Supplementary Fig. S8).

Silmitasertib (also called CX-4945, inhibitor for CK2[58]) but not D-4476 (inhibitor for CK1[59])

also had a similar impact (Fig. 4b, c, Supplementary Fig. S8). Although H-89 (inhibitor for

PKA) only had a moderate effect on mVenus-α-TAT1 localization, co-expression of mCFP-

PKI, a more potent inhibitor for PKA[60], considerably increased nuclear localization of mVenus-

α-TAT1 (Fig 4c, Supplementary Fig. S8, Supplementary Table T1). Finally, treatment with

LJI308 (inhibitor for Ribosomal S9 kinase) appeared to further increase nuclear exclusion of

mVenus-α-TAT1 (Fig. 4c, Supplementary Table T1). These data demonstrate that nuclear

localization of α-TAT1 is negatively regulated by CDKs, CK2 and PKA.

Nuclear localization of α-TAT1 is phospho-inhibited: Since α-TAT1(1-284) did not display

nuclear exclusion (Fig. 3b, c, d), we reasoned that the phosphosites which inhibit nuclear

localization of α-TAT1 might be located between F285 and R323. NetPhos prediction server

identified S294, T303, S315 and T322 as potential phosphosites in this region

(Supplementary Fig. S7a, indicated in red box), wherein only S315 and T322 have been

reported to be phosphorylated in phosphoproteomic studies (Supplementary Fig. S7b) and

they also flank the putative NLS (Fig. 5a). Importin-α binds with NLS enriched in basic

residues through a charge-based interaction[44]. Phosphorylation of amino acids adjacent to

such an NLS may inhibit the association of Importin-α binding through a disruption of the

charge balance in the NLS region[61]. Alanine substitution of T322, but not of S315, significantly

increased nuclear localization of mVenus-α-TAT1 (Fig. 5a, b, c, d, Supplementary Fig. S9a, b,

c). Alanine substitution of both S315 and T322, α-TAT1(ST/A), showed considerably more

nuclear localization of mVenus-α-TAT1 than T322 alone (Fig 5a, b, c, d). These data suggest

that T322 phosphorylation inhibits nuclear localization of α-TAT1, while S315 may play a co-

operative role in such inhibition. Substitution of S315 with acidic residues (S315D) appeared

to boost nuclear exclusion of α-TAT1, whereas substitution of T322 with acidic residues

(T322E) or both (ST/DE) displayed increased diffused pattern, but not increased nuclear

accumulation (Supplementary Fig. S9a, b, c). This may be because these acidic residues,

unlike phosphate moieties, do not sufficiently counter the basic residues in the NLS; or that

phospho-T322, and to a lesser extent, phospho-S315 phosphorylation may be involved in

protein-protein interactions, possibly with 14-3-3 isoforms, that inhibit nuclear localization of α-

TAT1. These observations suggest that the phosphate moiety in phosphorylated T322 and

S315 is critical for inhibition of α-TAT1 nuclear import.

One possible explanation of increased nuclear localization of T322A mutant is that

phospho-T322 mediates nuclear export of α-TAT1. Truncation of the putative NLS including

T322, α-TAT1-delNLS, did not increase nuclear localization (Fig. 5a, b, c, d), indicating that

T322 did not mediate nuclear export of α-TAT1. Furthermore, alanine substitution of the

hydrophobic residues in α-TAT1 NES as well as S315 and T322, α-TAT1(NES/A, ST/A) not

only abrogated nuclear exclusion, but considerably increased nuclear accumulation of α-TAT1

(Fig. 5a, b, c, d), suggesting additive effects of NES inhibition and NLS activation. α-

TAT1(NES/A, ST/DE) mutant also showed diffused pattern but not nuclear accumulation

(Supplementary Fig. S9a, b, c). Taken together, our observations suggest that phospho-T322

inhibits α-TAT1 NLS and that the α-TAT1 NES and NLS act independently of one another.

Discussion:

One of the bottlenecks in elucidating the role of microtubule acetylation in biological phenomena is the knowledge gap of how upstream molecular signaling pathways control α-TAT1 function to modulate microtubule acetylation. Our study demonstrates that intracellular α-TAT1 localization is a dynamically regulated process, orchestrated by a balance of nuclear export and import, which modulates microtubule acetylation levels (Fig. 5e). To our knowledge, this is the first study to identify the molecular mechanisms that spatially regulate α-TAT1. We have demonstrated a hitherto unknown role of the inherently disordered α-TAT1 C-terminus and identified novel interactions with 14-3-3 proteins and several kinases. TAK1 dependent phosphorylation of α-TAT1 Serine-237 has been reported to stimulate its catalytic property[62]. In neurons, p27[kip1] directly binds to α-TAT1 and stabilizes it against proteasomal degradation[24], thus enhancing α-tubulin acetylation. Our observation that spatial sequestration of α-TAT1 from microtubules modulates acetylation dynamics suggests a role of the nucleus as a reservoir or sequestration chamber to control protein access of substrates. Regulated spatial sequestration of biomolecules can control their action[63–67] and aberrant localization of proteins have been reported in many diseases[68–71]. Our study further highlights the role of spatial signaling processes in controlling protein function.

We have demonstrated active nuclear export of α-TAT1 by Exp1 through an NES rich in hydrophobic residues, which was critical for efficient microtubule acetylation. In addition, we have identified an NLS consistent with non-canonical class IV NLS[44]. Interestingly, position 7 of this NLS, which should not be an acidic residue, is occupied by Threonine-322. Since phosphorylation of threonine can significantly increase its net negative charges, it is ideally situated to act as an ON/OFF switch for the NLS. Although we have identified Threonine-322 to be the critical phospho-residue that inhibits nuclear import, Serine-315 appears to provide

additional inhibition. The increased nuclear localization of ST/A mutant over T322A mutant raises the possibility that S315 and T322 may aggregate signals from different signaling pathways to fine-tune α-TAT1 localization.

Our data demonstrate that nuclear localization of α-TAT1 is inhibited by kinase action, possibly on Threonine-322 and Serine-315. Specifically, our study shows a role of CDKs, PKA and CK2 in coordinating spatial distribution of α-TAT1. Such phospho-regulation of α-TAT1 provides a possible mechanism for the changes in α-TAT1 localization and microtubule acetylation observed at different stages of the cell cycle[72]. We identified Threonine-322 to be a putative binding site for 14-3-3 proteins and demonstrated that α-TAT1 binds to 14-3-3-β and 14-3-3-ζ proteins. 14-3-3s typically interact with phospho-serines or phospho-threonines in intrinsically disordered regions and may mediate nuclear transport of proteins by masking NES or NLS[73]. Furthermore, 14-3-3 proteins may significantly alter the structure of their binding partners to align along their rigid α-helical backbone, to expose or hide critical binding sites[74]. Comparable kinase-mediated regulation of nuclear export and nuclear import has previously been reported in a few transcription regulators[75–78]. In particular, regulation of Cdc25 localization by Checkpoint kinase1 (Chk1) mediated phosphorylation of and subsequent recruitment of 14-3-3-β to an NLS-proximal phosphosite is virtually the same as our proposed model (Fig. 5e) of α-TAT1 localization[78,79], suggesting that such kinase-mediated balancing of nuclear export and import is a general strategy for protein localization. α-TAT1 is unique in this aspect in that unlike the other proteins that are spatially regulated in this manner, α-TAT1 has no known substrates in the nucleus and that its nuclear localization appears to be primarily to sequester it from microtubules. Of course, it is possible that nuclear import of α-TAT1 facilitates interactions with presently unidentified substrates located in the nucleus. In a similar vein, 14-3-3 proteins and Exp1 are also acetylated[80,81], and it is intriguing to speculate that these might be substrates of α-TAT1.

It is worthwhile to consider that a significant number of post-translational modifications

of α-TAT1 appear on its intrinsically disordered C-terminus (Supplementary Fig. 7b).

Disordered regions may act as a signaling hub by interacting with multiple proteins, thus

facilitating complex formation and acting as integrators of signaling pathways[82]. We have

demonstrated the presence of an NES, NLS, phosphorylation sites and putative 14-3-3

binding sites within the α-TAT1 C-terminus. That it is well conserved across mammalian

species as well as in all the human isoforms suggests a critical role of the α-TAT1 C-terminus

in its function. Indeed, numerous cancer-associated mutations curated in COSMIC[83]

(www.cancer.sanger.ac.uk) and TCGA Research Network (www.cancer.gov/tcga) databases,

in the ATAT1 gene are located in the intrinsically disordered C-terminal region. More

specifically, there are a considerable number of deletions, frame shifts and missense

mutations encompassing the NES and the NLS regions, which may be expected to affect the

spatial distribution of α-TAT1. Whether these mutations underlie the pathogenesis in these

cancers remain to be examined. Considering the role of microtubule acetylation in a wide

array of cellular activities, it may be conjectured that loss of spatial regulation of α-TAT1 may

be present in other diseases as well.

In conclusion, we propose a new model for regulation of microtubule acetylation

through spatial sequestration of α-TAT1 (Fig. 5e), which include three key aspects: presence

of an NES that facilitates Exp1 mediated nuclear export, presence of an NLS to mediate

nuclear import and finally, modulation of this nuclear import by kinases. Further investigation

into the role of specific kinases on α-TAT1 localization may yield a better understanding of its

function in cellular processes and pathologies and help identify new therapeutic targets.

Materials and Methods

Cell culture and transfection: HeLa and HEK-293T cells were cultured in DMEM basal media and passaged every third day of culture. For optimal growth, the media were supplemented with 10% (v/v) fetal bovine serum and Penicillin/Streptomycin, and the cells were maintained under standard cell culture conditions (37 °C and 5% CO_2). The cell lines were regularly checked for mycoplasma contamination. FuGENE 6 reagent (Promega, Madison, WI) was used for transient transfection of HeLa cells according to the manufacturer's instructions. For immunoprecipitation assays, HEK cells were transfected using calcium phosphate method.

DNA plasmids: H2B-mCherry construct was a generous gift from Dr. Sergi Regot. α-TAT1 construct was a generous gift from Dr. Antonina Roll-Mecak. The α-TAT1 construct was subcloned into the pTriEx-4 vector (Novagen) using PCR and restriction digestion with mVenus at the N terminus and α-TAT1 at the C terminus. H2B-mCherry and CFP-PKI constructs were respectively subcloned into mCherry-C1 and mCer3-C1 vectors (Clontech). GFP-αTAT1 construct was a gift from from Dr. Philippe Chavrier and Dr. Guillaume Montagnac. HA-14-3-3 plasmids were a generous gift from Dr. Michael Yaffe. As indicated in the results and figure legends, tags of compatible fluorescent proteins including Cerulean, mVenus and mCherry were appended to facilitate detection. Unless specified otherwise, the termini of tagging were positioned as in the orders they were written. Truncations of α-TAT1 were generated by PCR. Point mutations of α-TAT1 were generated using overlapping PCR. The open reading frames of all DNA plasmids were verified by Sanger sequencing.

Sequence alignment: Protein sequence alignment was performed using Clustal-W[84] (https://www.ebi.ac.uk/Tools/msa/clustalo/).

Nuclear transport and kinase inhibitors: LMB was purchased from LC Laboratories (catalog # L6100). SB203580 (Sigma Aldrich, catalog # S8307), Doramapimod (BIRB 796, Selleck Chemicals, catalog # S1574), CHIR99021 (Sigma Aldrich, catalog # SML1046), Sostrastaurin (Selleck Chemicals, catalog # S2791), RO-3306 (Selleck Chemicals, catalog # S7747), Ipatasertib (RG7440, Selleck Chemicals, catalog # S2808), Capivasertib (AZD5363, Selleck Chemicals, catalog # S8019), Silmitasertib (CX 4945, Selleck Chemicals, catalog # S2248), KU-55933 (Sigma Aldrich, catalog # SML1109) were generous gifts from Dr. Sergi Regot. SB239063 (Sigma Aldrich, catalog # S0569) was a generous gift from Dr. Jun Liu. The rest of the kinase inhibitors were purchased as indicated: Staurosporine (Sigma Aldrich, catalog # 569397), LJI308 (Sigma Aldrich, catalog # SML1788), Y-27632 (LC Laboratories, catalog # Y-5301), Gö 6976 (Sigma Aldrich, catalog # 365250), Gö 6983 (Sigma Aldrich, catalog # G1918), H-89 (Sigma Aldrich, catalog # B1427), D4476 (BioVision, catalog # 1770), KN-62 (Selleck Chemicals, catalog # S7422), KU-57788 (MedChemExpress, catalog # HY-11006), Purvalanol B (AdipoGen Life Sciences, catalog # SYN-1070), KT-5823 (Cayman Chemicals, catalog # 10010965), PD0332991 (Sigma Aldrich, catalog # PZ0199).

Immunofluorescence assays: HeLa cells were transiently transfected with mVenus-α-TAT1, mVenus-α-TAT1 catalytic domain and NLS-mVenus-α-TAT1 catalytic domain. 24 hours post-transfection, cells were fixed using ice-cold methanol for 10 minutes, washed thrice with cold PBS, blocked with 1% BSA in PBS for one hour and then incubated overnight at 4°C with monoclonal antibodies against tubulin (Millipore, catalog # MAB1864) and acetylated α-Tubulin (Sigma Aldrich, catalog # T7451). Next day, the samples were washed thrice with cold PBS and incubated with secondary antibodies (Invitrogen) for one hour at room temperature, after which they were washed thrice with PBS and images were captured by microscopy. For LMB treatment, HeLa cells were dosed with 100 nM LMB or equal volume of vehicle (EtOH),

incubated for 4 hours, after which methanol fixation and immunostaining was performed as described above.

Immunoprecipitation assays: HEK293T cells were transiently transfected with pEGFP-c1 (GFP-Ctl) or GFP-αTAT1 with HA-14-3-3β or HA-14-3-3ζ using the calcium phosphate method. Cell lysates were prepared by scraping cells using 1X lysis buffer (10X recipe- 50 mM Tris pH 7.5, triton 20%, NP40 10%, 2 M NaCl, mixed with cOmplete protease inhibitor tablet - Roche, Product number 11873580001). Cell lysates rotated on a wheel at 4°C for 15 min and centrifuged for 10 min at 13,000 rpm 4°C to pellet the cell debris. A small volume of the supernatant was used as the soluble input. Soluble detergent extracts were incubated with GFP nanobody (NanoTag, N0310) for 1 h at 4°C. Samples were then centrifuged and washed thrice with wash buffer (250 mM NaCl, 0.1% Triton X-100 in PBS). The resin and the soluble input were then mixed with Laemmli buffer (composed of 60 mM Tris-HCl pH 6.8, 10% glycerol, 2% SDS and 50 mM DTT with the addition of protease and phosphatase inhibitors). Samples were boiled 5 min at 95°C before loading in polyacrylamide gels. Gels were transferred for western blot and membranes were blocked with TBST (0.1% Tween) and 5% milk and incubated 1 h with the primary antibody and 1 h with HRP-conjugated secondary antibody. Bands were revealed with ECL chemiluminescent substrate (Biorad). Two different western blots were used to visualize GFP-Input and HA-14-3-3 proteins due to similar molecular weights. Antibodies used: GFP-HRP (NB600-313, Novus Biologicals), anti-HA (rat; Merck; 11867423001), anti-exportin-1 (mouse; BD Transduction Laboratories™; 611832). Secondary HRP antibodies were all purchased from Jackson ImmunoResearch.

Microscopy and image analyses: All imaging was performed with an Eclipse Ti microscope (Nikon) with a 100X objective (1.0X zoom and 4X4 binning) and Zyla 4.2 sCMOS camera (Andor), driven by NIS Elements software (Nikon). Time lapse imaging was performed at 15 min intervals for 10-15 hours. All live cell imaging was conducted at 37°C, 5% CO_2 and 90%

humidity with a stage top incubation system (Tokai Hit). Vitamin and phenol red-free media (US Biological) supplemented with 2% fetal bovine serum were used in imaging to reduce background and photobleaching. Inhibitors and vehicles were present in the imaging media during imaging. All image processing and analyses were performed using Metamorph (Molecular Devices, Sunnyvale, CA, USA) and FIJI software (NIH, Bethesda, MD, USA).

For categorical analysis of mVenus-α-TAT1 localization, images were visually inspected and classified as displaying either cytosolic, diffused, or nuclear localization of mVenus fluorescence signal. For ratiometric analysis, the ratio of the fluorescence intensity from region of interest ($\approx$ 10 µm diameter) in the nucleus to that in a perinuclear area was used to minimize any volumetric artifacts (Fig. 1c). To determine the baseline Nuc/cyto ratio for cytosolic (<0.8) and nuclear (>1.2), we visually identified WT mVenus-α-TAT1 cells, with or without LMB treatment, that showed distinctly cytosolic or nuclear localization and used the rounded average ratio values from these cells (n > 100 cells). All cells showing nuc/cyto ratio in between were classified as diffused. For both categorical and ratiometric analyses, H2B-mCherry signal was used to identify nuclei; in a few cases where H2B-mCherry signal was absent, phase images were used to identify the nuclei. Cells displaying too much (near saturation in 16 bit) or too little (approximately less than 1.5-fold signal over background) of mVenus or mCherry fluorescence signal, or those which appeared to be dying on visual inspection were not included during image acquisition.

For immunofluorescence assays with exogenous expression of mVenus-α-TAT1 plasmids, transfected cells were identified by the presence of mVenus fluorescence signal. The ratio of acetylated α-Tubulin over α-Tubulin (Ac. α-Tub/α-Tub) for transfected cells was normalized against that for non-transfected cells averaged over 20 untransfected cells from the same dish. For LMB and vehicle treatment, Ac. α-Tub/α-Tub ratios are shown.

Statistical analyses: Microsoft Excel (Microsoft, Redmond, WA, USA) and R (R Foundation

for Statistical Computing, Vienna, Austria) were used for statistical analyses. Exact number of

samples for each data set are specified in the respective figure legends. Data was pooled

from at least three independent experiments (technical replicates) and within each experiment

typically data from at least 40 cells were obtained (biological replicates). In some cases with

kinase inhibitors that induced cell death, we collected data from smaller number of cells per

experiment but increased the number of experiments to ensure sufficient data. Sample sizes

were chosen based on the commonly used range in the field without performing any statistical

power analysis. Normal probability plot (Supplementary Fig. S10) was utilized to confirm nor-

mal distribution of the Nuc/Cyto ratio of mVenus-αTAT1. Extreme outliers ($<Q1 - 3 \times IQR$ or

$>Q3+3 \times IQR$) were excluded from plots and statistical analyses. *P*-values were obtained from

two-tailed Students *t*-test assuming unequal variance. Exact *P*-values for kinase inhibitor as-

says are available in Supplementary Table T1.

Data availability: All relevant data and source codes are included. Plasmid constructs will be

available through Addgene.

Acknowledgements

We thank Allen Kim for discussions that led to initiation of this project and Amy F. Peterson for help with the kinase inhibitor assays. We thank Yuta Nihongaki and Helen D. Wu for constructive discussions, as well as Robert DeRose for proofreading. This project was supported by American Heart Association fellowship 20POST35220046 (ADR), discretionary funds (TI), the La Ligue contre le cancer (S-CR17017) and Centre National de la Recherche Scientifique and Institut Pasteur. SS is funded by the ITN PolarNet Marie Curie grant and Fondation pour la Recherche Médicale and is enrolled at the Ecole Doctorale Frontières du Vivant (FdV) – Programme Bettencourt.

Contributions

ADR initiated the project and designed and performed most of the experiments and data analyses. GSP and EGG performed experiments and data analyses under the guidance of ADR and TI. SS performed the immunoprecipitation assays under the guidance of SEM. ADR and TI wrote the final version of the manuscript based on contributions from all the authors.

Competing interests

The authors declare no competing interests.

Figure legends

Figure 1. Intracellular distribution of α-TAT1 mediates its function.

a) Cartoon showing predicted NES and NLS in intrinsically disordered C-terminus of α-TAT1, adapted from PDB: 4GS4, b) intracellular distribution of mVenus-α-TAT1, red dotted lines outline nuclei as identified by H2B-mCherry in lower panel, c) cartoon showing criteria for categorical and ratiometric analyses of α-TAT1 distribution, d) categorical analysis (n = 1032 cells) and e) ratiometric analysis (n = 304 cells) of mVenus-α-TAT1 localization, f) cartoon showing α-TAT1 mutants used in g) immunofluorescence assays showing levels of acetylated α-tubulin and total α-tubulin, transfected cells are indicated with red arrowheads, h) ratio of acetylated α-tubulin to total tubulin intensities with exogenous expression of α-TAT1 and its mutants, normalized against that of non-transfected cells, (WT: 50, catalytic domain: 44, NLS-catalytic domain: 48 cells). Scale bar = 10μm. ***: $P<0.001$ and NS: not significant, Student's t-test.

Figure 2. α-TAT1 undergoes Exp1 mediated nuclear export.

a) Co-immunoprecipitation of endogenous Exp1 with GFP-α-TAT1, b) intracellular distribution of mVenus-α-TAT1 with vehicle (EtOH) and 100 nM LMB treatment, nuclei are indicated in red dotted lines, c) categorical analysis (WT: 1032, vehicle: 450, LMB: 495 cells) and d) ratiometric analysis (WT: 304, vehicle: 210, LMB: 240 cells) of mVenus-α-TAT1 localization with vehicle and LMB treatment, e) immunofluorescence images showing acetylated and total α-tubulin in HeLa cells with vehicle or LMB treatment, f) ratio of acetylated to total α-tubulin with vehicle or LMB treatment (vehicle:120, LMB: 130 cells). Scale bar = 10μm. ***: $P<0.001$ and NS: not significant, Student's t-test.

Figure 3. Intracellular distribution of α-TAT1 is mediated by its C-terminus.

a) Cartoon showing α-TAT1 mutant design, b) categorical analysis (WT: 1032, Cat. Dom: 371,

C-term: 290, delNES: 343 and NES/A: 501 cells) and c) ratiometric analysis (WT: 304, Cat.

Dom: 213, C-term: 220, delNES: 221 and NES/A: 212 cells) of mVenus-α-TAT1 mutant locali-

zation as indicated, d) representative images showing intracellular distribution of mVenus-α-

TAT1 mutants as listed, red dotted lines outline nuclei, e) ratiometric analysis of intracellular

distribution of mVenus-α-TAT1 C-term with 100 nM LMB (C-term: 220, LMB: 209 cells. Scale

bar = 10µm. ***: $P<0.001$ and NS: not significant, Student's t-test.

Figure 4. Nuclear localization of α-TAT1 is phospho-inhibited.

a) Co-immunoprecipitation of GFP-α-TAT1 with HA-14-3-3β and HA-14-3-3ζ proteins, b) intra-

cellular distribution of mVenus-α-TAT1 with Staurosporine, RO-3306, PD0332991 and Sil-

mitasertib treatment, nuclei are indicated in red dotted lines, c) ratiometric analysis of mVe-

nus-α-TAT1 localization with kinase inhibitors (WT:304, STS: 183, RO-3306: 221,

PD0332991: 234, Purvalanol-B: 243, D4476:253, Silmitasertib: 253, H89: 219, PKI: 208, Ipa-

tasertib: 259, Capivsertib: 218, Go6976: 297, Go6983: 241, Sostrastaurin: 401, KT-5823: 269,

SB203580: 259, Birb796: 262, KU55933: 317, LJI308: 229, CHIR99021: 227, KN-62:210, KU-

57788: 249, Y-27632: 161 cells). Scale bar = 10µm. ***: $P<0.001$ and NS: not significant, Stu-

dent's t-test.

Figure 5. α-TAT1 has a C-terminal phospho-inhibited NLS.

a) Cartoon showing α-TAT1 putative NLS flanked by potential phosphosites and correspond-

ing mutants, b) intracellular distribution of mVenus-α-TAT1 mutants as indicated, nuclei are

outlined in red dotted lines, c) ratiometric analysis (WT: 304, T322A: 228, ST/A: 194, delNLS:

208, NES/A, ST/A: 225 cells) and d) categorical analysis (WT: 1032, T322A: 346, ST/A: 290,

delNLS: 563, NES/A, ST/A: 476 cells) of intracellular localization of mVenus-α-TAT1 mutants,

e) proposed model of α-TAT1 localization and its impact on microtubule acetylation. Scale bar

= 10µm. ***P<0.001 and NS, not significant, Student's *t*-test.

References:

1. L'Hernault, S. W. & Rosenbaum, J. L. Chlamydomonas alpha-tubulin is posttranslationally modified by acetylation on the epsilon-amino group of a lysine. *Biochemistry* **24**, 473–478 (1985).

2. Diggins, M. A. & Dove, W. F. Distribution of acetylated alpha-tubulin in Physarum polycephalum. *J. Cell Biol.* **104**, 303–309 (1987).

3. Janke, C. & Montagnac, G. Causes and Consequences of Microtubule Acetylation. *Curr. Biol.* **27**, R1287–R1292 (2017).

4. Schatten, G. *et al.* Acetylated alpha-tubulin in microtubules during mouse fertilization and early development. *Dev. Biol.* **130**, 74–86 (1988).

5. Boggs, A. E. *et al.* α-Tubulin Acetylation Elevated in Metastatic and Basal-like Breast Cancer Cells Promotes Microtentacle Formation, Adhesion, and Invasive Migration. *Cancer Res.* **75**, 203–215 (2015).

6. Esteves, A. R. *et al.* Acetylation as a major determinant to microtubule-dependent autophagy: Relevance to Alzheimer's and Parkinson disease pathology. *Biochim. Biophys. Acta BBA - Mol. Basis Dis.* **1865**, 2008–2023 (2019).

7. L'Hernault, S. W. & Rosenbaum, J. L. Chlamydomonas alpha-tubulin is posttranslationally modified in the flagella during flagellar assembly. *J. Cell Biol.* **97**, 258–263 (1983).

8. Maruta, H., Greer, K. & Rosenbaum, J. L. The acetylation of alpha-tubulin and its relationship to the assembly and disassembly of microtubules. *J. Cell Biol.* **103**, 571–579 (1986).

9. Piperno, G., LeDizet, M. & Chang, X. J. Microtubules containing acetylated alpha-tubulin in mammalian cells in culture. *J. Cell Biol.* **104**, 289–302 (1987).

10. Sudo, H. & Baas, P. W. Acetylation of Microtubules Influences Their Sensitivity to Severing

by Katanin in Neurons and Fibroblasts. *J. Neurosci.* **30**, 7215–7226 (2010).

11. Portran, D., Schaedel, L., Xu, Z., Théry, M. & Nachury, M. V. Tubulin acetylation protects

long-lived microtubules against mechanical ageing. *Nat. Cell Biol.* **19**, 391–398 (2017).

12. Xu, Z. *et al.* Microtubules acquire resistance from mechanical breakage through

intralumenal acetylation. *Science* **356**, 328–332 (2017).

13. Eshun-Wilson, L. *et al.* Effects of α-tubulin acetylation on microtubule structure and

stability. *Proc. Natl. Acad. Sci.* **116**, 10366–10371 (2019).

14. Joo, E. E. & Yamada, K. M. MYPT1 regulates contractility and microtubule acetylation to

modulate integrin adhesions and matrix assembly. *Nat. Commun.* **5**, 3510 (2014).

15. Bance, B., Seetharaman, S., Leduc, C., Boëda, B. & Etienne-Manneville, S. Microtubule

acetylation but not detyrosination promotes focal adhesion dynamics and astrocyte

migration. *J. Cell Sci.* **132**, (2019).

16. Swiatlowska, P., Sanchez-Alonso, J. L., Wright, P. T., Novak, P. & Gorelik, J. Microtubules

regulate cardiomyocyte transversal Young's modulus. *Proc. Natl. Acad. Sci.* **117**, 2764–

2766 (2020).

17. Coleman, A. K., Joca, H. C., Shi, G., Lederer, W. J. & Ward, C. W. Tubulin acetylation

increases cytoskeletal stiffness to regulate mechanotransduction in striated muscle.

bioRxiv 2020.06.10.144931 (2020) doi:10.1101/2020.06.10.144931.

18. Zhang, Y. *et al.* Identification of genes expressed in C. elegans touch receptor neurons.

Nature **418**, 331–335 (2002).

19. Shida, T., Cueva, J. G., Xu, Z., Goodman, M. B. & Nachury, M. V. The major α-tubulin K40

acetyltransferase αTAT1 promotes rapid ciliogenesis and efficient mechanosensation.

Proc. Natl. Acad. Sci. U. S. A. **107**, 21517–21522 (2010).

20. Morley, S. J. *et al.* Acetylated tubulin is essential for touch sensation in mice. *eLife* **5**,

e20813 (2016).

21. Yan, C. *et al.* Microtubule Acetylation Is Required for Mechanosensation in Drosophila.

Cell Rep. **25**, 1051-1065.e6 (2018).

22. Lin, S., Sterling, N. A., Junker, I. P., Helm, C. T. & Smith, G. M. Effects of αTAT1 and

HDAC5 on axonal regeneration in adult neurons. *PloS One* **12**, e0177496 (2017).

23. Fourcade, S. *et al.* Loss of SIRT2 leads to axonal degeneration and locomotor disability

associated with redox and energy imbalance. *Aging Cell* **16**, 1404–1413 (2017).

24. Morelli, G. *et al.* p27Kip1 Modulates Axonal Transport by Regulating α-Tubulin

Acetyltransferase 1 Stability. *Cell Rep.* **23**, 2429–2442 (2018).

25. Lee, C.-C., Cheng, Y.-C., Chang, C.-Y., Lin, C.-M. & Chang, J.-Y. Alpha-tubulin

acetyltransferase/MEC-17 regulates cancer cell migration and invasion through epithelial–

mesenchymal transition suppression and cell polarity disruption. *Sci. Rep.* **8**, 17477

(2018).

26. Castro-Castro, A., Janke, C., Montagnac, G., Paul-Gilloteaux, P. & Chavrier, P.

ATAT1/MEC-17 acetyltransferase and HDAC6 deacetylase control a balance of

acetylation of alpha-tubulin and cortactin and regulate MT1-MMP trafficking and breast

tumor cell invasion. *Eur. J. Cell Biol.* **91**, 950–960 (2012).

27. Oh, S. *et al.* Genetic disruption of tubulin acetyltransferase, αTAT1, inhibits proliferation

and invasion of colon cancer cells through decreases in Wnt1/β-catenin signaling.

Biochem. Biophys. Res. Commun. **482**, 8–14 (2017).

28. Geeraert, C. *et al.* Starvation-induced Hyperacetylation of Tubulin Is Required for the

Stimulation of Autophagy by Nutrient Deprivation. *J. Biol. Chem.* **285**, 24184–24194

(2010).

29. McLendon, P. M. *et al.* Tubulin hyperacetylation is adaptive in cardiac proteotoxicity by

promoting autophagy. *Proc. Natl. Acad. Sci.* **111**, E5178–E5186 (2014).

30. Destaing, O. *et al.* A novel Rho-mDia2-HDAC6 pathway controls podosome patterning

through microtubule acetylation in osteoclasts. *J. Cell Sci.* **118**, 2901–2911 (2005).

31. Elliott, G. & O'Hare, P. Herpes Simplex Virus Type 1 Tegument Protein VP22 Induces the

Stabilization and Hyperacetylation of Microtubules. *J. Virol.* **72**, 6448–6455 (1998).

32. Husain, M. & Harrod, K. S. Enhanced acetylation of alpha-tubulin in influenza A virus

infected epithelial cells. *FEBS Lett.* **585**, 128–132 (2011).

33. Sabo, Y. *et al.* HIV-1 induces the formation of stable microtubules to enhance early

infection. *Cell Host Microbe* **14**, (2013).

34. Zan, J. *et al.* Rabies Virus Infection Induces Microtubule Depolymerization to Facilitate

Viral RNA Synthesis by Upregulating HDAC6. *Front. Cell. Infect. Microbiol.* **7**, (2017).

35. Kalebic, N. *et al.* αTAT1 is the major α-tubulin acetyltransferase in mice. *Nat. Commun.* **4**,

1–10 (2013).

36. Kim, G.-W., Li, L., Ghorbani, M., You, L. & Yang, X.-J. Mice lacking α-tubulin

acetyltransferase 1 are viable but display α-tubulin acetylation deficiency and dentate

gyrus distortion. *J. Biol. Chem.* **288**, 20334–20350 (2013).

37. Szyk, A. *et al.* Molecular Basis for Age-Dependent Microtubule Acetylation by Tubulin

Acetyltransferase. *Cell* **157**, 1405–1415 (2014).

38. Kalebic, N. *et al.* Tubulin Acetyltransferase αTAT1 Destabilizes Microtubules

Independently of Its Acetylation Activity. *Mol. Cell. Biol.* **33**, 1114–1123 (2013).

39. Friedmann, D. R., Aguilar, A., Fan, J., Nachury, M. V. & Marmorstein, R. Structure of the α-

tubulin acetyltransferase, αTAT1, and implications for tubulin-specific acetylation. *Proc.*

Natl. Acad. Sci. U. S. A. **109**, 19655–19660 (2012).

40. Mészáros, B., Erdős, G. & Dosztányi, Z. IUPred2A: context-dependent prediction of

protein disorder as a function of redox state and protein binding. *Nucleic Acids Res.* **46**,

W329–W337 (2018).

41. Ishida, T. & Kinoshita, K. PrDOS: prediction of disordered protein regions from amino acid

sequence. *Nucleic Acids Res.* **35**, W460-464 (2007).

42. la Cour, T. *et al.* Analysis and prediction of leucine-rich nuclear export signals. *Protein*

Eng. Des. Sel. **17**, 527–536 (2004).

43. Horton, P. *et al.* WoLF PSORT: protein localization predictor. *Nucleic Acids Res.* **35**,

W585-587 (2007).

44. Kosugi, S. *et al.* Six Classes of Nuclear Localization Signals Specific to Different Binding

Grooves of Importin α. *J. Biol. Chem.* **284**, 478–485 (2009).

45. Kutay, U. & Güttinger, S. Leucine-rich nuclear-export signals: born to be weak. *Trends Cell*

Biol. **15**, 121–124 (2005).

46. Hutten, S. & Kehlenbach, R. H. CRM1-mediated nuclear export: to the pore and beyond.

Trends Cell Biol. **17**, 193–201 (2007).

47. Sun, Q. *et al.* Nuclear export inhibition through covalent conjugation and hydrolysis of

Leptomycin B by CRM1. *Proc. Natl. Acad. Sci.* **110**, 1303–1308 (2013).

48. Sekimoto, T., Fukumoto, M. & Yoneda, Y. 14-3-3 suppresses the nuclear localization of

threonine 157-phosphorylated p27Kip1. *EMBO J.* **23**, 1934–1942 (2004).

49. Mancini, M. *et al.* 14-3-3 ligand prevents nuclear import of c-ABL protein in chronic

myeloid leukemia. *Traffic Cph. Den.* **10**, 637–647 (2009).

50. Huang, X. *et al.* Shade-induced nuclear localization of PIF7 is regulated by

phosphorylation and 14-3-3 proteins in Arabidopsis. *eLife* **7**, (2018).

51. Yaffe, M. B. *et al.* The Structural Basis for 14-3-3:Phosphopeptide Binding Specificity. *Cell*

91, 961–971 (1997).

52. Madeira, F. *et al.* 14-3-3-Pred: improved methods to predict 14-3-3-binding

phosphopeptides. *Bioinforma. Oxf. Engl.* **31**, 2276–2283 (2015).

53. Seetharaman, S. *et al.* Microtubules tune mechanosensitive cell responses. *bioRxiv*

2020.07.22.205203 (2020) doi:10.1101/2020.07.22.205203.

54. Vassilev, L. T. *et al.* Selective small-molecule inhibitor reveals critical mitotic functions of

human CDK1. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 10660–10665 (2006).

55. Jorda, R. *et al.* How Selective Are Pharmacological Inhibitors of Cell-Cycle-Regulating

Cyclin-Dependent Kinases? *J. Med. Chem.* **61**, 9105–9120 (2018).

56. Fry, D. W. *et al.* Specific inhibition of cyclin-dependent kinase 4/6 by PD 0332991 and

associated antitumor activity in human tumor xenografts. *Mol. Cancer Ther.* **3**, 1427–1438

(2004).

57. Toogood, P. L. *et al.* Discovery of a Potent and Selective Inhibitor of Cyclin-Dependent

Kinase 4/6. *J. Med. Chem.* **48**, 2388–2406 (2005).

58. Ferguson, A. D. *et al.* Structural basis of CX-4945 binding to human protein kinase CK2.

FEBS Lett. **585**, 104–110 (2011).

59. Rena, G., Bain, J., Elliott, M. & Cohen, P. D4476, a cell-permeant inhibitor of CK1,

suppresses the site-specific phosphorylation and nuclear exclusion of FOXO1a. *EMBO*

Rep. **5**, 60–65 (2004).

60. Kim, A. K., Wu, H. D. & Inoue, T. Rational Design of a Protein Kinase A Nuclear-cytosol

Translocation Reporter. *Sci. Rep.* **10**, 9365 (2020).

61. Harreman, M. T. *et al.* Regulation of nuclear import by phosphorylation adjacent to nuclear

localization signals. *J. Biol. Chem.* **279**, 20613–20621 (2004).

62. Shah, N. *et al.* TAK1 activation of alpha-TAT1 and microtubule hyperacetylation control

AKT signaling and cell growth. *Nat. Commun.* **9**, 1–12 (2018).

63. Shimada, Y., Gulli, M. P. & Peter, M. Nuclear sequestration of the exchange factor Cdc24

by Far1 regulates cell polarity during yeast mating. *Nat. Cell Biol.* **2**, 117–124 (2000).

64. Volmat, V., Camps, M., Arkinstall, S., Pouysségur, J. & Lenormand, P. The nucleus, a site

for signal termination by sequestration and inactivation of p42/p44 MAP kinases. *J. Cell*

Sci. **114**, 3433–3443 (2001).

65. Formstecher, E. *et al.* PEA-15 mediates cytoplasmic sequestration of ERK MAP kinase.

Dev. Cell **1**, 239–250 (2001).

66. Yang, W., Wightman, R. & Meyerowitz, E. M. Cell Cycle Control by Nuclear Sequestration

of CDC20 and CDH1 mRNA in Plant Stem Cells. *Mol. Cell* **68**, 1108-1119.e3 (2017).

67. Lasker, K. *et al.* Selective sequestration of signalling proteins in a membraneless

organelle reinforces the spatial regulation of asymmetry in Caulobacter crescentus. *Nat.*

Microbiol. **5**, 418–429 (2020).

68. Chen, Y. *et al.* Aberrant subcellular localization of BRCA1 in breast cancer. *Science* **270**,

789–791 (1995).

69. Jiao, W. *et al.* Aberrant nucleocytoplasmic localization of the retinoblastoma tumor

suppressor protein in human cancer correlates with moderate/poor tumor differentiation.

Oncogene **27**, 3156–3164 (2008).

70. Hung, M.-C. & Link, W. Protein localization in disease and therapy. *J. Cell Sci.* **124**, 3381–

3392 (2011).

71. Wang, X. & Li, S. Protein mislocalization: mechanisms, functions and clinical applications

in cancer. *Biochim. Biophys. Acta* **1846**, 13–25 (2014).

72. Nekooki-Machida, Y. *et al.* Dynamic localization of α-tubulin acetyltransferase ATAT1

through the cell cycle in human fibroblastic KD cells. *Med. Mol. Morphol.* **51**, 217–226

(2018).

73. Pennington, K. L., Chan, T. Y., Torres, M. P. & Andersen, J. L. The dynamic and stress-

adaptive signaling hub of 14-3-3: emerging mechanisms of regulation and context-

dependent protein–protein interactions. *Oncogene* **37**, 5587–5604 (2018).

74. Yaffe, M. B. How do 14-3-3 proteins work?-- Gatekeeper phosphorylation and the

molecular anvil hypothesis. *FEBS Lett.* **513**, 53–57 (2002).

75. Oeckinghaus, A. & Ghosh, S. The NF-κB Family of Transcription Factors and Its

Regulation. *Cold Spring Harb. Perspect. Biol.* **1**, (2009).

76. Spencer, S. L. *et al.* The Proliferation-Quiescence Decision Is Controlled by a Bifurcation

in CDK2 Activity at Mitotic Exit. *Cell* **155**, 369–383 (2013).

77. Regot, S., Hughey, J. J., Bajar, B. T., Carrasco, S. & Covert, M. W. High-Sensitivity

Measurements of Multiple Kinase Activities in Live Single Cells. *Cell* **157**, 1724–1734

(2014).

78. Deneke, V. E., Melbinger, A., Vergassola, M. & Di Talia, S. Waves of Cdk1 activity in S-

phase synchronize the cell cycle in Drosophila embryos. *Dev. Cell* **38**, 399–412 (2016).

79. Uchida, S. *et al.* Binding of 14-3-3β but not 14-3-3σ controls the cytoplasmic localization of

CDC25B: binding site preferences of 14-3-3 subtypes and the subcellular localization of

CDC25B. *J. Cell Sci.* **117**, 3011–3020 (2004).

80. Choudhary, C. *et al.* Lysine acetylation targets protein complexes and co-regulates major

cellular functions. *Science* **325**, 834–840 (2009).

81. Andersen, J. L. *et al.* A biotin switch-based proteomics approach identifies 14-3-3ζ as a

target of Sirt1 in the metabolic regulation of caspase-2. *Mol. Cell* **43**, 834–842 (2011).

82. Wright, P. E. & Dyson, H. J. Intrinsically Disordered Proteins in Cellular Signaling and

Regulation. *Nat. Rev. Mol. Cell Biol.* **16**, 18–29 (2015).

83. Tate, J. G. *et al.* COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids*

Res. **47**, D941–D947 (2019).

84. F, M. *et al.* The EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic*

Acids Res. **47**, W636–W641 (2019).

Figure 1

a Catalytic domain (1-236)   Intrinsically disordered C-terminus (199-323)

282   297   315   322

Putative NES: LRPFVPEQELLRSLRL

Putative NLS: PAQRRRT^{322}

b

mVenus-α-TAT1   Cytosolic   Diffused   Nuclear

H2B-mCherry

c   cyto   diff   nuc

$\dfrac{Nuc}{Cyto}$ <0.8   0.8-1.2   >1.2

d Fraction of cells — Nuc, Diff, Cyto

e Nuc/Cyto — WT — nuc / 1.2 / diff / 0.8 / cyto

f

WT — Cat. domain — NES — NLS

Cat. domain

NLS-Cat. domain — NLS — Cat. domain

g   WT   Cat. domain   NLS-Cat. domain

mVenus-α-TAT1

anti-Ac. α-Tubulin

anti-Tubulin

h Ac. α-Tub/ α-Tub (a.u.) — ***   ***   NS — WT, Cat. domain, NLS-Cat.

Figure 2

Figure 3

a

WT — Cat. domain | NES | NLS

Cat. dom — Cat. domain

C-term — NES | NLS

delNES — Cat. domain

NES/A — Cat. domain | NES/A | NLS

NES: **L**RP**FV**PEQE**LL**RS**L**R**L**
NES/A: **A**RP**AA**PEQE**AA**RS**A**R**A**

b

Fraction of cells

Cytosolic ■ Diffused ■ Nuclear

c

Nuc/Cyto

*** NS *** ***

WT Cat. Dom C-term delNES NES/A

d

Cat. domain | C-terminus | delNES | NES/A

e

Nuc/Cyto

***

C-term | C-term + LMB

Figure 4

Figure 5

A phospho-regulated signal motif of α-TAT1 drives dynamic microtubule acetylation

Abhijit Deb Roy[1*], Evan G. Gross[2], Gayatri S. Pillai[2], Shailaja Seetharaman[3,4], Sandrine

Etienne-Manneville[3], Takanari Inoue[1*]

[1] Department of Cell Biology and Center for Cell Dynamics, Johns Hopkins University School

of Medicine, Baltimore, 855 North Wolfe Street, MD 21205, USA

[2] The Johns Hopkins University, Baltimore, MD 21218, USA

[3]Cell Polarity, Migration and Cancer Unit, Institut Pasteur, UMR3691 CNRS, Equipe

Labellisée Ligue Contre le Cancer, F-75015, Paris, France.

[4]Université Paris Descartes, Sorbonne Paris Cité, 12 Rue de l'École de Médecine, 75006

Paris, France.

[*] Correspondence:

Abhijit Deb Roy abhijit.debroy@gmail.com

Takanari Inoue jctinoue@jhmi.edu

20
21

a) Query: α-TAT1 Isoform 7

MEFPFDVDALFPERITVLDQHLRPPARRPGTTTPARVDLQQQIMTIIDELGKASAKAQNLSAPITSA
SRMQSNRHVVYILKDSSARPAGKGAIIGFIKVGYKKLFVLDDREAHNEVEPLCILDFYIHESVQRH
GHGRELFQYMLQKERVEPHQLAIDRPSQKLLKFLNKHYNLETTVPQVNNFVIFEGFFAHQHRPPA
PSLRATRHSRAAAVDPTPAAPARKLPPKRAEGDIKPYSSSDREFLKVAVEPPWPLNRAPRRATP
PAHPPPRSSSLGNSPERGP**LRPFVPEQELLRSLRL**CPPHPTARLLLAADPGGS**PAQRRRT**^{322}R

Supplementary Figure S1. αTAT1 C-terminus is disordered. a) amino acid sequence of αTAT1
isoform 7 used as the query, the putative NES and NLS are in bold, b) IUPred2 (and ANCHOR2)
and c) PrDOS results suggesting that αTAT1 C-terminus is disordered. The threshold to be
considered disordered is 0.5 for both.

NetNES optimized prediction: LRPF**V**PEQE**LL**RS**L**R**L**

Hidden Makov Model (HMM) algorithm prediction: **L**RP**FV**PEQE**LL**RS**L**R**L**

Supplementary Figure S2. NetNES prediction for NES in αTAT1 identifying the putative NES, the predicted NES from NetNES optimized algorithm (pink) and that from Hidden Markov Model (blue) are shown with the hydrophobic residues indicated in bold.

Supplementary Figure S3. The putative NES and NLS in α-TAT1 are evolutionarily conserved. a) Alignment of human α-TAT1 isoforms, b) alignment of different mammalian α-TAT1 showing the C-terminal region including the putative NES and NLS (enclosed in boxes), genus of the mammalian α-TAT1 are shown. Alignments were performed using ClustalW (https://www.ebi.ac.uk/Tools/msa/clustalo/)

a)

α-TAT1

b)

Supplementary Figure S4. α-TAT1 intracellular localization is dynamic in nature. a) Spontaneous temporal changes in spatial distribution of mVenus-α-TAT1, scale bar = 10 μm b) temporal changes in nuclear fluorescence intensity of mVenus-α-TAT1 in HeLa cells, each line indicates normalized nuclear intensity of mVenus- α-TAT1 in a single cell, n = 20 cells

Supplementary Figure S5. α-TAT1 undergoes CRM1 dependent nuclear export. a), b) Temporal changes in mVenus-α-TAT1 localization on 100 nM LMB treatment after LMB addition at time 0, scale bar = 10 μm, for b), each line indicates normalized nuclear intensity of mVenus- α-TAT1 in a single cell, black line shows the mean and gray line indicate the 95% C.I., n = 26 cells , c) categorical analysis (WT: 1032, vehicle: 450, 1 nM: 329, 10 nM: 291, 100 nM: 495 cells) and d) ratiometric analysis (WT: 304, vehicle: 210, 1 nM: 203, 10 nM: 200, 100 nM: 240 cells) of mVenus-α-TAT1 localization on vehicle (EtOH) or LMB treatment at 1, 10 and 100 nM. ***$P<0.001$ and NS, not significant, Student's *t*-test.

a
NES: **L**RP**FV**PEQE**LL**RS**L**R**L**
VL/A: LRPF**A**PEQE**AA**RS**A**R**A**
NES/A: **A**RP**AA**PEQE**AA**RS**A**R**A**

Supplementary Figure S6. α-TAT1 has a C-terminal NES. a) α-TAT1 NES predicted by NetNES and mutants using alanine substitution, b) categorical (WT: 1032, D157N: 429, LMB: 495, VL/A: 436, NES/A: 501 cells), and c) ratiometric (WT: 304, D157N: 217, LMB: 240, VL/A: 201, NES/A: 212 cells) analyses of intracellular localization of mVenus-α-TAT1 and mutants as indicated. ***$P<0.001$ and NS, not significant, Student's t-test.

Supplementary Figure S7. a) Putative phosphosites in α-TAT1 predicted by NetPhos, those in F285-R323 are highlighted, b) currently reported post-translational modifications in α-TAT1 curated by PhosphoSitePlus®, www.phosphosite.org.

Supplementary Figure S8. Categorical analysis of α-TAT1 localization on treatment with kinase inhibitors. WT: 1032, STS: 157, R0-3306: 796, PD0332991: 543, Purvalanol B: 992, D4476: 689, 418, H89: 251, PKI: 354, Ipatasertib: 546, Capivasertib: 385, Go6976: 712, Go6983: 606, Sostrastaurin: 605, KT5823: 570, SB203580: 505, Birb796: 991, KU55933: 815, LJI308: 522, CHIR99021: 484, KN62: 531, KU57788: 949, Y27632: 340 cells.

Supplementary Figure S9. α-TAT1 has a C-terminal phospho-inhibited NLS. a) Cartoon showing α-TAT1 NES and NLS flanked by potential phosphosites and mutants, b) categorical (WT:1032, S315A: 442, S315D: 412, T322E: 310, ST/DE: 258 cells) and c) ratiometric (WT: 304, S315A: 212, S315D: 223, T322E: 232, ST/DE: 195 cells) analyses of intracellular localization of mVenus-α-TAT1 mutants. ***$P<0.001$ and NS, not significant, Student's *t*-test.

Supplementary Figure S10. Normal probability plot of nuclear/cytosolic ratio of mVenus-α-TAT1 expressed in HeLa cells.

Drug	Conc. Used	target kinase	Incubation time	p value (t-test)
STS	100 nM	Pan kinase	4 hours	1.20187E-11
RO3306	10 µM	Cdk1, Cdk2	4 hours	1.87845E-10
PD0332991	1 µM	Cdk4, Cdk6	4 hours	1.29985E-10
Purvalanol B	1 µM	Cdk1, Cdk2, Cdk5	4 hours	0.086698844
D4476	1 µM	CK1	4 hours	0.244864719
Silmitasertib	10 µM	CK2	4 hours	7.34743E-12
H89	10 µM	PKA	4 hours	0.001868139
PKI	NA	PKA	Overnight	8.76993E-08
Ipatasertib	10 µM	PKB (Akt)	4 hours	0.026614985
Capivasertib	1 µM	PKB (Akt)	4 hours	0.928873535
Go6976	250 nM	PKC	4 hours	0.900843024
Go6983	250 nM	PKC	4 hours	0.068144954
Sostrastaurin	10 µM	PKC	4 hours	0.008916116
KT5823	5 µM	PKG	4 hours	0.581314971
SB203580	10 µM	p38 MAPK	4 hours	0.007945499
Birb796	10 µM	p38 MAPK	4 hours	0.768174955
KU55933	10 µM	ATM	4 hours	0.26994365
LJI308	10 µM	RSK	4 hours	9.62544E-07
CHIR99021	1 µM	GSK3	4 hours	0.309440886
KN62	10 µM	CaMK-II, P2RX7	4 hours	0.021582087
KU57788	1 µM	DNA-PK	4 hours	0.385979985
Y27632	10 µM	ROCK	4 hours	0.375473038

Supplementary Table T1. Details of kinase inhibitors used.