COVID-19 CG: Tracking SARS-CoV-2 mutations by locations and dates of interest

Albert Tian Chen[1], Kevin Altschuler, Shing Hei Zhan[2]*, Yujia Alina Chan[1]*, Benjamin E.

Deverman[1]*

[1]Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA,

United States of America.

[2]Department of Zoology & Biodiversity Research Centre, the University of British Columbia,

Vancouver BC, Canada.

*Correspondence to: bdeverma@broadinstitute.org, alinac@broadinstitute.org,

zhan@zoology.ubc.ca

Lead Contact

Benjamin E. Deverman

Stanley Center for Psychiatric Research

Broad Institute of MIT and Harvard

75 Ames St

Cambridge, MA 02142

Phone (626) 755-7523

bdeverma@broadinstitute.org

Abstract

COVID-19 CG is an open resource for tracking SARS-CoV-2 single-nucleotide variations

(SNVs) and lineages while filtering by location, date, gene, and mutation of interest. COVID-19

CG provides significant time, labor, and cost-saving utility to diverse projects on SARS-CoV-2

transmission, evolution, emergence, immune interactions, diagnostics, therapeutics, vaccines,

and intervention tracking. Here, we describe case studies in which users can interrogate (1)

SNVs in the SARS-CoV-2 Spike receptor binding domain (RBD) across different geographic

regions to inform the design and testing of therapeutics, (2) SNVs that may impact the

sensitivity of commonly used diagnostic primers, and (3) the recent emergence of a dominant

lineage harboring an S477N RBD mutation in Australia. To accelerate COVID-19 research and

public health efforts, COVID-19 CG will be continually upgraded with new features for users to

quickly and reliably pinpoint mutations as the virus evolves throughout the pandemic and in

response to therapeutic and public health interventions.

Introduction

Since the beginning of the pandemic, SARS-CoV-2 genomic data has been accumulating at an unprecedented rate (90,000+ virus genomes as of September, 2020 on the GISAID database) (Elbe and Buckland-Merrett, 2017; Shu and McCauley, 2017). Numerous countries have mobilized to sequence thousands of SARS-CoV-2 genomes upon the occurrence of local outbreaks, collectively and consistently contributing more than 10,000 genomes per month (**Figure S1A, B**). It is important to note that, despite the slow accumulation of potentially functional (nonsynonymous) mutations, there has been a steady increase in the number of variants with more than 6 nonsynonymous mutations compared to the WIV04 reference, an early isolate of SARS-CoV-2 that was collected in Wuhan in December, 2019 (**Figure S1C**). To evaluate the outcomes of anti-COVID-19 measures and detect keystone events of virus evolution, it is important to track changes in SARS-CoV-2 mutation and population dynamics in a location and date-specific manner. Indeed, several countries and the National Institutes of Health (NIH) have recognized how critical it is to collect SARS-CoV-2 genomic data to support contact tracing efforts and to inform public health decisions – these are paramount to the re-opening of countries and inter-regional travel (Collins 2020; Rockett et al. 2020; Oude Munnink, et al. 2020; Gudbjartsson et al. 2020; Pybus et al. 2020). Yet, the quantity and complexity of SARS-CoV-2 genomic data (and metadata) make it challenging and costly for the majority of scientists to stay abreast of SARS-CoV-2 mutations in a way that is meaningful to their specific research goals. Currently, each group or organization has to independently expend labor, computing costs, and, most importantly, time to curate and analyze the genomic data from GISAID before they can generate specific hypotheses about SARS-CoV-2 lineages and mutations in their population(s) of interest.

Results

To address this challenge, we built COVID-19 CoV Genetics (COVID-19 CG, covidcg.org), a

performant, interactive, and fully-scalable web application that tracks SARS-CoV-2 single-

nucleotide variants (SNVs) and lineages without sub-sampling. COVID-19 CG is a free, open

access interface that allows users to adapt analyses according to their dates and locations of

interest (**Figure 1A,B**; data processing workflow in **Figure S2**). Users can also select and

compare trends in SARS-CoV-2 lineage or SNV frequency across multiple locations (**Figure**

1C) as we will demonstrate using case studies. COVID-19 CG provides functionalities that, to

the best of our knowledge, cannot be found in other existing public browsers, and was designed

to empower these specific user groups:

Vaccine and therapeutics developers can inform the design and testing of their vaccine,

antibody, or small molecule by using COVID-19 CG to rapidly identify all of the variants in their

targeted SARS-CoV-2 protein or antigen, alongside the frequency of each variant in their

geographic location(s) of interest. Scientists can use COVID-19 CG to generate hypotheses and

experimentally determine whether the variants present in the location of vaccine/therapeutic

implementation may impact their product-specific interaction interface or antigen.

Case study of SNVs in the receptor binding domain (RBD) of the SARS-CoV-2 Spike:

Analyzing SNVs by geography and time is critical as the frequency of each SNV may vary

significantly across different regions over time. For instance, an S477N mutation in the RBD has

become dominant in Oceania (57.5% of Oceanian genotypes, all time) although it constitutes

only 4.36% of SARS-CoV-2 genotypes globally and has not been detected in Africa, Asia, or

South America (**Figure 2A**). SNV frequency in a given region can also shift over time, e.g., an

RBD N439K mutation not found in Ireland prior to July is now present in 79.5% of the genomes

collected mid-July through August (**Figure 2B**). Another rare RBD S477N mutation, which was

found in only 1.05% of the Australian SARS-CoV-2 sequences before June, now constitutes

more than 90% of the sequenced June through August isolates (**Figure 2C**). This geographical

and temporal variation is important to incorporate into the design and testing of therapeutic

antibodies (such as those under development as therapeutics by Regeneron that specifically

target the SARS-CoV-2 Spike RBD), as well as mRNA or recombinant protein-based vaccines.

This will help to assure developers of the efficacy of their therapeutics and vaccines against the

SARS-CoV-2 variants that are present in the intended location of implementation.

In addition, COVID-19 CG can be harnessed to track changes in SARS-CoV-2 evolution post-

implementation of therapeutics and vaccines. It will be crucial to watch for rare escape variants

that could resist drug- or immune-based interventions to eventually become the dominant

SARS-CoV-2 variant in the community. This need was particularly emphasized by a Regeneron

study that demonstrated that single amino acid variants could evolve rapidly in the SARS-CoV-2

Spike to ablate binding to antibodies that had been previously selected for their ability to

neutralize all known RBD variants; these amino acid variations were found either inside or

outside of the targeted RBD region, and some are already present at low frequency among

human isolates globally (Baum et al., 2020). The authors, Baum et al., suggested that these

rare escape variants could be selected under the pressure of single antibody treatment, and,

therefore, advocated for the application of cocktails of antibodies that bind to different epitopes

to minimize SARS-CoV-2 mutational escape. A recent study by Greaney et al. generated high-

resolution 'escape maps' delineating RBD mutations that could potentially result in virus escape

from neutralization by ten different human antibodies (Greaney et al., 2020). Based on lessons

learnt from the rise of multidrug resistant bacteria and cancer cells, it will be of the utmost

importance to continue tracking SARS-CoV-2 evolution even when multiple vaccines and

therapeutics are implemented in a given human population.

Diagnostics developers can evaluate their probe, primer, or point-of-care diagnostic according

to user-defined regional and temporal SARS-CoV-2 genomic variation. More than 665

established primers/probes are built into COVID-19 CG, and new diagnostics will be continually

incorporated into the browser. Users can also input custom coordinates or sequences to

evaluate their own target sequences and design new diagnostics.

Case study of SNVs that could impact the sensitivity of diagnostic primers: A recent

preprint alerted us to the finding that a common G29140T SNV, found in 22.3% of the study's

samples from Madera County, California, was adversely affecting SARS-CoV-2 detection by the

NIID_2019-nCoV_N_F2 diagnostic primer used at their sequencing center; the single SNV

caused a ~30-fold drop in the quantity of amplicon produced by the NIID_2019-nCov_N_F2/R2

primer pair (Vanaerschot et al., 2020). We used COVID-19 CG to detect other SNVs that could

impact the use of this primer pair, discovering that there are SARS-CoV-2 variants in several

countries with a different C29144T mutation at the very 3' end of the same NIID_2019-

nCoV_N_F2 primer (**Figure 3A**). As the authors of the preprint, Vanaerschot et al., noted, SNVs

could impact assay accuracy if diagnostic primers and probes are also being used to quantify

viral loads in patients. We found that at least ten other primer pairs could potentially be at risk in

different geographical regions due to SNVs that appear proximal to the 3' ends of primers

(**Figure 3B-K**): China-CDC-N-F and R; NIH, Thailand, WH-NIC N-F; US CDC 2019-nCoV-N1-R;

US CDC 2019-nCoV-N2-F; ARTIC-V3_nCoV-2019_11_RIGHT; ARTIC-V3_nCoV-

2019_13_LEFT; ARTIC-V3_nCoV-2019_34_LEFT; ARTIC-V3_nCoV-2019_39_LEFT (note that

the ARTIC primers are used for nanopore sequencing) (Tyson et al., 2020); WHO

N_Sarbarco_R1; and Institut Pasteur, Paris 12759Rv. We advocate that labs and clinics use

COVID-19 CG (https://covidcg.org) to check their most commonly used primers and probes

against the SARS-CoV-2 sequences that are prevalent in their geographic regions.

Researchers and public health professionals can use COVID-19 CG to gain insights as to

how the virus is evolving in a given population over time (e.g., in which genes are mutations

occurring, and do these lead to structural or phenotypic changes?). For example, users can

track D614G distributions across any region of interest over time. **Figure 4** shows a variety of

different D614G population dynamics in different areas. Nonetheless, we strongly caution

against inferring (i) chains or directionality of transmission and (ii) changes in the transmissibility

of any SARS-CoV-2 SNV based on population dynamics alone. Inconsistent sampling, sampling

biases, differences in founder host population traits (even median patient age), superspreading

events, regionally and temporally differential travel restrictions, and numerous other factors

instead of virus biological differences can influence the global distribution of SNVs (Grubaugh et

al., 2020).

Case study of Australia's new dominant SARS-CoV-2 variant: We discovered that the

SARS-CoV-2 Spike S477N mutation has become more prevalent in Australia (**Figure 5A**).

Globally, the S477N mutation was first detected in a single sample of lineage B.1.1.25 that was

collected on January 26, 2020 in Victoria, Australia; this is now the dominant SARS-CoV-2

variant in the region (**Figure 5B, C**). In particular, the set of SNVs that co-occur with the S477N

mutation in Australia (all time, as well as prior to May, 2020 before the most recent outbreak)

are different from the set of co-occurring SNVs in the United Kingdom (**Figure 5C**) —

suggesting that the S477N mutation occurred separately in the Australian and the UK lineages.

However, COVID-19 CG only reflects data contributed to GISAID. Variants of interest could be

present in other countries, but not yet known to the public because the sequencing centers in

those countries have not collected or deposited their data in GISAID. Furthermore, in instances

where only a singular, sporadic variant is detected (no sustained transmission), there is also the

possibility of sequencing error resulting in incorrect lineage assignment. Due to these caveats,

the genetic data must be used in combination with other types of data, such as from contact

tracing efforts, before it is possible to draw conclusions about the international transmission of

SARS-CoV-2 variants. In the case of the S477N variant that is now dominating in Australia, the

sequencing data alone indicate that the local transmission of this variant since January, 2020 in

Australia cannot be ruled out.

Discussion

COVID-19 CG (https://covidcg.org) was designed to be modular in order to continually integrate

datasets from other COVID-19 initiatives. We anticipate building modules for users to **(1)** map

emerging mutations onto structural interaction interfaces of interest (e.g., between virus protein

and therapeutic antibodies or host proteins) using existing and future structures on the Protein

Data Bank (PDB), **(2)** visualize mutations in isolates of interest in the context of different virus

protein phenotypes or mutational constraints of antibody epitopes according to emerging

genotype-to-phenotype maps (Greaney et al., 2020; Starr et al., 2020), **(3)** compare SARS-CoV-

2 mutations in different host species (e.g., humans versus minks) (Oude Munnink et al., 2020b),

(4) rapidly determine when and where each lineage or SNV has been detected around the

world, and **(5)** overlay important policy events or travel restrictions over time on the lineage or

SNV tracker to help guide user date range selection. In addition, as more detailed metadata is

generated by COVID-19 studies and initiatives, we will update the application to enable filtering

according to patient traits such as gender, age, ethnicity, and medical condition (e.g.,

symptoms, hospitalization).

COVID-19 CG (https://covidcg.org) was built to help scientists and professionals worldwide, with

varying levels of bioinformatics expertise, in their real-time analysis of SARS-CoV-2 genetic

data. We hope that COVID-19 CG will also motivate decision makers to sustain or accelerate

their sequencing of virus isolates in their geographical area for the purposes of informing

vaccine, therapeutics, and policy development. Collecting virus genomic data is particularly

relevant to regions that are experiencing increases in COVID-19 cases. If only sparse genomic

data are sampled, we risk the late detection of SARS-CoV-2 variants that exhibit enhanced

virulence or resistance against therapeutics or vaccination programs in these pandemic

hotspots. Furthermore, the widespread implementation of vaccines and antibody therapies

could stimulate the emergence and selection of new escape variants (Baum et al., 2020). To

compound these risks, SARS-CoV-2 transmission from humans to minks (and back into

humans) has already been detected at farms across the Netherlands, Denmark, Spain, and the

United States (Oude Munnink et al., 2020b). This process of species crossing, if left unchecked,

can result in the emergence of diverse SARS-CoV-2 variants.

Coordinated sequencing and contact tracing efforts (e.g., in the UK, Singapore, the

Netherlands, Italy, California, and Australia) emphasize the urgency of establishing open access

platforms to evaluate trends in virus introduction into each country or region in real time. Local

policymakers, public health researchers, and scientists can use **global** SARS-CoV-2 genetic

data, in complementation with contact tracing data, to better understand which lineages were

imported into their region (from which potential international locations), whether these were

introduced multiple times, and if particular lineages are dying out or persisting. Labs in

numerous countries are already making incredible efforts to sequence the SARS-CoV-2 variants

circulating in their local populations (**Figure 6**). When each country actively contributes to the

database of SARS-CoV-2 genomes, this protects against sampling biases that can impact the

ability to perform phylogenetic analysis and interpret global SARS-CoV-2 data. Towards this

goal that affects all of humanity, we advocate for the increased sequencing of SARS-CoV-2

isolates from patients (and infected animals) around the world, and for these data to be shared

in as timely a manner as possible.

Experimental procedures

Data Pipeline

Our data processing pipeline is written with the Snakemake scalable bioinformatics workflow engine (Koster and Rahmann, 2012), which modularizes our workflow and enables reproducibility and compatibility with cloud-computing. All code and relevant documentation are hosted on an open-source, publicly available GitHub repository (https://github.com/vector-engineering/COVID19-CG), providing example data for users to validate our pipeline.

Data Acquisition

SARS-CoV-2 sequences and metadata are downloaded daily from the GISAID EpiCov™ database (https://epicov.org/epi3/start), by navigating to the "Browse" tab and selecting sequences by "Submission Date". As of 2020-09-04, only 10,000 sequences can be downloaded from this selection at a time, so the selection is adjusted to include no more than 10,000 sequences. "Sequences", "Patient status metadata", and "Sequencing technology metadata" are downloaded separately, stored in separate folders, and renamed for ingestion into the data processing pipeline (see below).

Sequence Preprocessing

Based on best practices, we filter out sequences meeting any of the following criteria: (1) Present on the NextStrain's exclusion list (https://github.com/nextstrain/ncov/blob/master/defaults/exclude.txt), (2) Isolates from non-humans (animals, environmental samples, etc), (3) genome length less than 29,700 nt, or (4) >5% ambiguous base calls. Sequences which pass all preprocessing filters are carried onto the next steps.

Metadata Cleaning

We clean metadata with the aim of preserving the original intent of the authors and data

submitters while presenting simpler and unified versions to end users. Sequencing metadata is

cleaned to remove obvious typos, and to unify labels with the same meaning, e.g., "MinION"

and "Nanopore MinION". Location metadata is cleaned with the goal of simplifying the location

selector in the sidebar. Locations with excessive children are collapsed to the nearest upper

hierarchical grouping. E.g., if a state has individual data for 200+ towns, these towns will be

collapsed to the county level in order to facilitate easier data browsing. Typos and clear

identities are also unified to prevent the display of duplicate locations in the application.

SNV Assignments

SNVs and insertions/deletions (indels) at the nucleotide and amino acid level are determined by

aligning each sequence to the WIV04 reference sequence (WIV04 is a high quality December,

2019 isolate that is 100% identical to the first publicly available SARS-CoV-2 genome reference

Wuhan-Hu-1/NC_045512.2, excepting the sequences at the end of the genomes) using

bowtie2. Spurious SNVs and probable sequencing errors, defined as less than 3 global

occurrences, are filtered out prior to downstream analysis. SNVs involving ambiguous base

calls ("N" in the original sequences) are ignored. Indels resulting in frameshifts are ignored, and

SNVs/indels occurring in non-protein-coding regions are ignored when determining SNVs/indels

on the AA level.

Lineage/Clade Analysis

Viral lineages, as defined by the *pangolin* tool (Rambaut et al., 2020), and clades (Tang et al.,

2020) are provided by GISAID. In accordance with *pangolin*, SNVs present in >90% of

sequences within each lineage/clade will be assigned as lineage/clade-defining SNVs.

Application Compilation

The web application is written in Javascript, and primarily uses the libraries React.js, MobX, and

Vega. The code is compiled into javascript bundles by webpack. All sequence data is

compressed and injected inline as JSON into the javascript bundle – no server is needed to

serve data to end users. The compiled application files can then be hosted on any static server.

Application Deployment

COVID CG (https://covidcg.org) is hosted by Google Cloud Run. The application code is

assembled into a Docker image (see Dockerfile), with a build environment (node.js) and

deployment environment (NGINX).

Acknowledgments

B.D. and A.C. are supported by awards from the National Institute of Neurological Disorders and

Stroke (UG3NS111689) and a Brain Initiative award funded through the National Institute of

Mental Health (UG3MH120096) and from the Stanley Center for Psychiatric Research. Y.A.C. is

supported by a Broad Institute SPARC award and the Stanley Center for Psychiatric Research.

We gratefully acknowledge all of the authors from the originating laboratories responsible for

obtaining the specimens and the submitting laboratories where genetic sequence data were

generated and shared via the GISAID Initiative, on which this resource is based.

Data Availability

All of the data shown in this manuscript and displayed on COVID CG (https://covidcg.org) are

downloaded from the GISAID EpiCov™ database (https://www.gisaid.org). All code and relevant

documentation are hosted on an open-source, publicly available GitHub repository

(https://github.com/vector-engineering/COVID19-CG).

Author Contributions

Y.A.C., S.H.Z, and A.T.C. conceived the project and browser. B.E.D. supervised the work.

A.T.C. and K.A. developed the COVID CG web browser with input from all of the authors. S.H.Z.

advised the implementation of lineage and clade analysis. Y.A.C., B.E.D., and A.T.C. prepared

the figures, analyzed the data, and wrote the manuscript with input from all authors.

Declaration of Interests

Shing Hei Zhan is a Co-founder and Director of Bioinformatics at Fusion Genomics Corporation,

which develops molecular diagnostic assays for infectious diseases. The other authors declare

no competing interests.

References

Baum A, Fulton BO, Wloga E, Copin R, Pascal KE, Russo V, Giordano S, Lanza K, Negron N,

Ni M, Wei Y, Atwal GS, Murphy AJ, Stahl N, Yancopoulos GD & Kyratsous CA (2020) Antibody

cocktail to SARS-CoV-2 spike protein prevents rapid mutational escape seen with individual

antibodies. Science. Available at: http://dx.doi.org/10.1126/science.abd0831

Collins F (2020) Genome Data Help Track Community Spread of COVID-19. NIH Director's

Blog. Available at: https://directorsblog.nih.gov/2020/07/21/genome-data-helps-track-

community-spread-of-covid-19/

Dong E, Du H & Gardner L (2020) An interactive web-based dashboard to track COVID-19 in

real time. Lancet Infect. Dis. 20, 533–534.

van Dorp L, Richard D, Tan CCS, Shaw LP, Acman M & Balloux F (2020) No evidence for

increased transmissibility from recurrent mutations in SARS-CoV-2. Genomics, 501.

Elbe S & Buckland-Merrett G (2017) Data, disease and diplomacy: GISAID's innovative

contribution to global health. Glob Chall 1, 33–46.

Greaney AJ, Starr TN, Gilchuk P, Zost SJ, Binshtein E, Loes AN, Hilton SK, Huddleston J,

Eguia R, Crawford KHD, Dingens AS, Nargi RS, Sutton RE, Suryadevara N, Rothlauf PW, Liu

Z, Whelan SPJ, Carnahan RH, Crowe JE & Bloom JD (2020) Complete mapping of mutations to

the SARS-CoV-2 spike receptor-binding domain that escape antibody recognition. bioRxiv,

2020.09.10.292078. Available at: https://www.biorxiv.org/content/10.1101/2020.09.10.292078v1

Grubaugh ND, Hanage WP & Rasmussen AL (2020) Making Sense of Mutation: What D614G

Means for the COVID-19 Pandemic Remains Unclear. Cell. Available at:

https://linkinghub.elsevier.com/retrieve/pii/S0092867420308175

Gudbjartsson DF, Helgason A, Jonsson H, Magnusson OT, Melsted P, Norddahl GL,

Saemundsdottir J, Sigurdsson A, Sulem P, Agustsdottir AB, Eiriksdottir B, Fridriksdottir R,

Gardarsdottir EE, Georgsson G, Gretarsdottir OS, Gudmundsson KR, Gunnarsdottir TR,

Gylfason A, Holm H, Jensson BO, Jonasdottir A, Jonsson F, Josefsdottir KS, Kristjansson T,

Magnusdottir DN, le Roux L, Sigmundsdottir G, Sveinbjornsson G, Sveinsdottir KE, Sveinsdottir

M, Thorarensen EA, Thorbjornsson B, Löve A, Masson G, Jonsdottir I, Möller AD, Gudnason T,

Kristinsson KG, Thorsteinsdottir U & Stefansson K (2020) Spread of SARS-CoV-2 in the

Icelandic Population. N. Engl. J. Med. 382, 2302–2315.

Koster J & Rahmann S (2012) Snakemake--a scalable bioinformatics workflow engine.

Bioinformatics 28, 2520–2522. Available at: http://dx.doi.org/10.1093/bioinformatics/bts480

Langmead B, Trapnell C, Pop M & Salzberg SL (2009) Ultrafast and memory-efficient alignment

of short DNA sequences to the human genome. Genome Biology 10, R25. Available at:

http://dx.doi.org/10.1186/gb-2009-10-3-r25

Oude Munnink BB, Nieuwenhuijse DF, Stein M, O'Toole Á, Haverkate M, Mollers M, Kamga SK,

Schapendonk C, Pronk M, Lexmond P, van der Linden A, Bestebroer T, Chestakova I,

Overmars RJ, van Nieuwkoop S, Molenkamp R, van der Eijk AA, GeurtsvanKessel C, Vennema

H, Meijer A, Rambaut A, van Dissel J, Sikkema RS, Timen A, Koopmans M & Dutch-Covid-19

response team (2020) Rapid SARS-CoV-2 whole-genome sequencing and analysis for informed

public health decision-making in the Netherlands. Nat. Med. Available at:

http://dx.doi.org/10.1038/s41591-020-0997-y

Oude Munnink BB, Sikkema RS, Nieuwenhuijse DF, Molenaar RJ, Munger E, Molenkamp R,

van der Spek A, Tolsma P, Rietveld A, Brouwer M, Bouwmeester-Vincken N, Harders F, Honing

RH der, Wegdam-Blans MCA, Bouwstra R, GeurtsvanKessel C, van der Eijk A, Velkers F, Smit

L, Stegeman A, van der Poel WHM & Koopmans M (2020) Jumping back and forth:

anthropozoonotic and zoonotic transmission of SARS-CoV-2 on mink farms. bioRxiv,

2020.09.01.277152. Available at: https://www.biorxiv.org/content/10.1101/2020.09.01.277152v1

Pybus OG & Rambaut A with du Plessis L, Zarebski AE, Kraemer MUG, Raghwani J, Gutiérrez

B, Hill V, McCrone J, Colquhoun R, Jackson B, O'Toole A, Ashworth J, on behalf of the COG-

UK consortium (2020) Preliminary analysis of SARS-CoV-2 importation & establishment of UK

transmission lineages. Virological. Available at: https://virological.org/t/preliminary-analysis-of-

sars-cov-2-importation-establishment-of-uk-transmission-lineages/507

Rambaut A, Holmes EC, O'Toole Á, Hill V, McCrone JT, Ruis C, du Plessis L & Pybus OG

(2020) A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic

epidemiology. Nat Microbiol. Available at: http://dx.doi.org/10.1038/s41564-020-0770-5

Rockett RJ, Arnott A, Lam C, Sadsad R, Timms V, Gray K-A, Eden J-S, Chang S, Gall M,

Draper J, Sim EM, Bachmann NL, Carter I, Basile K, Byun R, O'Sullivan MV, Chen SC-A,

Maddocks S, Sorrell TC, Dwyer DE, Holmes EC, Kok J, Prokopenko M & Sintchenko V (2020)

360 Revealing COVID-19 transmission in Australia by SARS-CoV-2 genome sequencing and agent-

361 based modeling. Nat. Med. Available at: http://dx.doi.org/10.1038/s41591-020-1000-7

362 Shu Y & McCauley J (2017) GISAID: Global initiative on sharing all influenza data - from vision

363 to reality. Euro Surveill. 22. Available at: http://dx.doi.org/10.2807/1560-

364 7917.ES.2017.22.13.30494

365 Starr TN, Greaney AJ, Hilton SK, Crawford KHD, Navarro MJ, Bowen JE, Alejandra Tortorici M,

366 Walls AC, Veesler D & Bloom JD (2020) Deep mutational scanning of SARS-CoV-2 receptor

367 binding domain reveals constraints on folding and ACE2 binding. bioRxiv, 2020.06.17.157982.

368 Available at: https://www.biorxiv.org/content/10.1101/2020.06.17.157982v1

369 Tang X, Wu C, Li X, Song Y, Yao X, Wu X, Duan Y, Zhang H, Wang Y, Qian Z, Cui J & Lu J

370 (2020) On the origin and continuing evolution of SARS-CoV-2. Natl. Sci. Rev. 7, 1012–1023.

371 Tyson JR, James P, Stoddart D, Sparks N, Wickenhagen A, Hall G, Choi JH, Lapointe H,

372 Kamelian K, Smith AD, Prystajecky N, Goodfellow I, Wilson SJ, Harrigan R, Snutch TP, Loman

373 NJ & Quick J (2020) Improvements to the ARTIC multiplex PCR method for SARS-CoV-2

374 genome sequencing using nanopore. bioRxiv, 2020.09.04.283077. Available at:

375 https://www.biorxiv.org/content/10.1101/2020.09.04.283077v1

376 Vanaerschot M, Mann SA, Webber JT, Kamm J, Bell SM, Bell J, Hong SN, Nguyen MP, Chan

377 LY, Bhatt KD, Tan M, Detweiler AM, Espinosa A, Wu W, Batson J, Dynerman D, CLIAHUB

378 Consortium, Wadford DA, Puschnik AS, Neff N, Ahyong V, Miller S, Ayscue P, Tato CM, Paul S,

379 Kistler A, DeRisi JL & Crawford ED (2020) Identification of a polymorphism in the N gene of

380 SARS-CoV-2 that adversely impacts detection by a widely-used RT-PCR assay. bioRxiv,

381 2020.08.25.265074. Available at:

382 https://www.biorxiv.org/content/10.1101/2020.08.25.265074v1.full

Figure 1. The COVID-19 CG (https://covidcg.org) browser interface. (A) Users can select

SARS-CoV-2 genomes according to lineage, clade, or SNV, virus gene or protein, and

location(s). Genomes can also be filtered by metadata, and specifically analyzed at genomic

coordinates of interest, such as the target sites of 665 commonly used diagnostic primers and

probes. (B) In the "Compare lineages or SNVs" tab, users can visualize SARS-CoV-2 lineages

or SNVs by location, define their date range of interest, and see the corresponding SNVs at the

nucleotide or amino acid level. (C) In the "Compare locations" tab, users can compare the

frequencies of specific lineages or SNVs in multiple locations over time.

Figure 2. Mutational frequencies in the SARS-CoV-2 Spike receptor binding domain (RBD) across geographical location and time. Screen captures from the Compare AA SNVs tab are shown. **(A)** The top 10 RBD SNVs alongside the number of high quality sequences available on GISAID are shown for each region. **(B)** The top RBD SNVs for Ireland between mid July and August, 2020 are shown. The S439N mutant had not been previously detected in Ireland. **(C)** The top RBD SNVs for Australia between June and August, 2020 are shown. The S477N mutant constituted only 1.05% of the Australian SARS-CoV-2 genomes on GISAID prior to June.

Figure 3. Investigating diagnostic-targeted regions of the SARS-CoV-2 genome for SNVs that could impact primer/probe sensitivity. Images were downloaded from the Compare NT SNVs tab. Labels for specific mutations were added. Primer pairs that contain at least one primer with potentially impactful SNVs near the 3' end are shown. None of the 11 primer pairs shown here were designed with degenerate bases. (**A**) The G29140T has been demonstrated to impact the NIID_2019-nCOV_N_F2 primer sensitivity. (**B-K**) Primer pairs affected by SNVs with a global frequency of more than 80 instances are shown. (**B**) As an example, majors SNVs are colored accordingly in the China-CDC-N-F and R (forward and reverse) primers.

Figure 4. Population dynamics of Spike D614G in different regions. Images were

downloaded from the Compare Lineages tab of covidcg.org: The Spike D614 variants are

shown in grey, and the G614 variants are shown in green. Plots displaying different population

dynamics were deliberately selected. Time is shown on the horizontal axis and the number of

sequences is shown on the vertical axis; these differ per country depending on when and how

many samples were collected and whether the sequences were deposited onto GISAID by

August 31, 2020.

Figure 5. Frequency of the Spike S477N mutation in Australia over time. (A) Image

downloaded from the Compare NT SNVs tab of covidcg.org: SARS-CoV-2 variants bearing the

Spike S477N mutation (also known as the G22992A SNV; depicted in pink), the majority of

which lie in the B.1.1.25 lineage, have become the most prevalent form of SARS-CoV-2 in

Australia. **(B)** Image downloaded from the Compare Locations tab of covidcg.org: the

cumulative percent of sequences carrying the S477N mutation in Australia. **(C)** Images

downloaded from the Compare NT SNVs tab of covidcg.org: Co-occurring SNVs of G22992A

(Spike S477N) in Australia, all time versus prior to May, 2020, versus in the United Kingdom.

Figure 6. Global statistics of SARS-CoV-2 genomes contribution to GISAID. Interactive and

more comprehensive versions of the figure panels are displayed on the Global Sequencing

Coverage tab of covidcg.org. **(A)** A world map of countries labeled by the number of SARS-

CoV-2 sequences contributed per 1000 cases. **(B)** A bar graph showing the sequences per

1000 cases for the top five countries and the USA. Countries with less than 500 cases were

excluded from this plot. **(C)** A world map of countries labeled by median days between sample

collection and sequence deposition. **(D)** A bar graph showing the median days from collection to

deposition for the top five countries and the USA. These interactive displays are generated

using sequencing data from the GISAID EpiCov™ database (nextmeta file) and case data from

the JHU CSSE COVID-19 Data (Dong et al., 2020). Only samples that were collected between

March and May, 2020 were included to avoid biases from samples that have been collected in

the previous three months but not yet deposited onto GISAID.

Supplemental Figures

Figure S1. The number of global SARS-CoV-2 genome sequences and mutations is

accumulating. Data shown as of September 9, 2020. **(A)** Sequence deposition in GISAID

continues at a steady pace, albeit there is a lag between collection (red line) and submission

date (blue line). The rate of sequence submission is steady at >10,000 genomes per month. **(B)**

More than 100 countries have deposited SARS-CoV-2 genomes in GISAID. **(C)** The number of

SARS-CoV-2 variants with more than six nonsynonymous (NS) mutations continues to increase.

Figure S2. COVID-19 CG computational workflow. **(1)** Starting from the GISAID database, sequences are continuously updated, manually for now, but ultimately via automated data ingestion. **(2)** Based on best practices, we filter out sequences on NextStrain's exclusion list, non-human isolates, <29,700 nt, or with >5% ambiguous base calls (van Dorp et al., 2020). **(3)** SNVs at the nucleotide and amino acid level are determined by aligning (via *bowtie2*) each sequence to the WIV04 reference, a high quality December, 2019 isolate recommended by GISAID; NextStrain uses the 100% identical Wuhan-Hu-1 (Langmead et al., 2009). Importantly, spurious SNVs and probable sequencing errors are filtered out prior to downstream analysis. **(4)** Viral lineages, defined by the *pangolin* tool, are provided by GISAID. In accordance with *pangolin*, SNVs present in >90% of sequences within each lineage are assigned as lineage-defining SNVs. **(5)** The curated data and metadata, SNVs, and lineage-assigned SNVs are associated with their respective sequence identifier and compiled into a compact data set. **(6)** These data are uploaded onto the COVID-19 CG web application. **(7)** New analyses will be built into the COVID-19 CG application throughout the course of the pandemic. **(8-10)** Features and modules that integrate knowledge from other COVID-19 initiatives are continuously incorporated into COVID-19 CG.